Highly defended nudibranchs 'escape' to visually distinct background habitats

Cedric P. van den Berg[1,2], Matteo Santon[2], John A. Endler[3], Karen L. Cheney[1]

[1]School of the Environment, The University of Queensland, 4072 Brisbane, Australia

[2]School of Biological Sciences, University of Bristol, BS8 1QU Bristol, UK

[3]School of Life and Environmental Sciences, Deakin University, Geelong, Victoria 3216, Australia

Corresponding author: Cedric P. van den Berg

Email: c.vandenberg@bristol. ac.uk

CPvdB 0000-0001-6422-7237

MS 0000-0002-9397-4052

JAE 0000-0002-7557-7627

KLC 0000-0001-5622-949

Lay Summary

According to the 'escape and radiate' hypothesis, the ability to store potent chemical defences as
protection from predators enables animals to be minimally dependent on matching visual
backgrounds for camouflage. We found evidence supporting this hypothesis in 12 species of
nudibranch molluscs. Aposematic species were found on visual backgrounds that differed consistently
in appearance from the visual background of undefended cryptic species. However, we found no
difference in the variability of visual backgrounds between species with or without chemical defences.
This suggests that warning colouration in eastern Australian Dorid nudibranchs coincides with broadly
generalisable spatiochromatic properties of visual backgrounds.

Abstract

The 'escape and radiate' hypothesis predicts that once species have evolved aposematism, defended
species can utilise more visually diverse visual backgrounds as they 'escape' the need to be well

camouflaged. This enables species to explore new ecological niches, resulting in increased
diversification rates. To test this hypothesis' 'escape' component, we examined whether the
background habitats of 12 nudibranch mollusc species differed among species depending on the
presence and strength of chemical defences. We obtained a rich array of colour pattern statistics using
Quantitative Colour Pattern Analysis (QCPA) to analyse backgrounds viewed through the eyes of a
potential predator (triggerfish, *Rhinecanthus aculeatus*). Colour pattern analysis was done at viewing
distances simulating an escalating predation sequence. We identified four latent factors comprising
17 non-correlated colour pattern parameters, which captured the among-species variability
associated with differences in chemical defences. We found that chemically defended species, indeed,
were found on visually distinct backgrounds with increased colour and luminance contrast,
independent of viewing distance. However, we found no evidence for increased among-species
background diversity coinciding with the presence and strength of chemical defences. Our results
agree with the 'escape and radiate' hypothesis, suggesting that potent chemical defences in Dorid
nudibranchs coincide with spatiochromatic differences of visual background habitats perceived by
potential predators.

Keywords

Aposematism, camouflage, defensive animal colouration, escape and radiate hypothesis, signal
honesty, visual ecology, visual modelling, warning signals.

Introduction

Animals and plants use aposematic colour patterns to advertise defences to potential
predators (Poulton, 1890). One of the hypothesised benefits of an organism evolving aposematism is
the potential to 'escape' the costs of crypsis, including limited movement and thus access to resources
due to the need to match a specific background habitat or preventing detection through motion
contrast for primary defence (Endler, 1984; Merilaita & Tullberg, 2005; Regan & Beverley, 1984).
Therefore, Merilaita and Tullberg (2005) suggested that alternative visual defence strategies, such as
aposematism, are more likely to evolve in highly variable visual environments where crypsis is difficult
to achieve. The evolution of aposematic signalling provides increased access to resources across
ecological niches as individuals are not strongly restricted to particular habitats to avoid predator
detection. This reduced dependency on crypsis may drive increased speciation and diversification
rates (Arbuckle & Speed, 2015; Ehrlich & Raven, 1964). This process was coined 'escape and radiate'

by Thompson (1989) and has since been adopted as a core concept underpinning the role of chemical
defences in adaptive radiations (e.g. Arbuckle & Harris, 2021).

The evolution of potent chemical defences often coincides with aposematism (see (Ruxton et
al., 2018; Summers et al., 2015; White & Umbers, 2021) for reviews), and theoretical modelling
supports the conclusion that aposematic species are less constrained by the visual properties of their
habitat (habitat generalists) than undefended background-matching species (habitat specialists)
(Merilaita & Tullberg, 2005; Speed, Brockhurst, et al., 2010). The appearance of an animal against its
visual background fundamentally impacts its detectability by potential predators, a crucial factor
shaping the design and function of defensive colouration (see van den Berg, Endler, et al., 2023) for
discussion). Therefore, assuming relaxed selection for background matching, secondary defences in a
diverse prey community inhabiting visually complex habitats might facilitate chemically defended
species to inhabit distinct and possibly more diverse visual backgrounds than undefended species.
However, whether the presence of chemical defences qualitatively or quantitatively correlates with
general among-species differences in background habitats remains unknown.

Here, we tested the hypothesis that more defended species inhabit distinct and more variable
habitats using 12 species of co-occurring Eastern Australian nudibranch species (Fig. S1). Nudibranchs
are an intriguing system for the study of defensive animal colouration due to potent chemical defences
in many species (Avila, 1995; Winters et al., 2022; Winters, White, et al., 2018) and colour patterns
ranging and colour patterns that range from bold aposematic displays to near-perfect camouflage
(Debelius & Kuiter, 2007). Indeed, the evolution of specialised glands (mantle dermal formations) for
the storage and secretion of defensive chemicals (Carbone et al., 2013; Wägele et al., 2006), in
combination with the evolution of aposematic signalling, may have led to adaptive radiation in
nudibranchs (Gosliner, 2001).

Until recently, studies quantifying visual backgrounds in the context of defensive animal
colouration have relied on the independent consideration of spatial and spectral (colour and
luminance) properties (e.g. Cortesi & Cheney, 2010; Nokelainen et al., 2021). In this study, we
considered these pattern components in combination (van den Berg, Troscianko, et al., 2020) using
calibrated digital photography, the multispectral image calibration and analysis toolbox (MICA)
(Troscianko & Stevens, 2015) and its' integrated Quantitative Colour Pattern Analysis (QCPA)
framework (van den Berg, Troscianko, et al., 2020). This methodology enabled us to assess the
spatiochromatic information of natural backgrounds upon which nudibranch individuals (12 species,
n = 184) were found. QCPA provides a descriptive array of image statistics capturing the
spatiochromatic properties of each background according to the physiological limitations of an

ecologically relevant observer. Here, we used the visual system of a triggerfish (*Rhinecanthus*
aculeatus), a common, omnivorous reef fish. We analysed different viewing distances to model an
escalating predation sequence to account for changes in background appearance to a predator up
close (2cm) and at a distance (30cm).

We then compared these colour pattern parameters to measures of the strength of chemical
defence for each species, which ranged from highly unpalatable to palatable, using previously
published data from antifeedant assays with rockpool shrimp (*Palaemon serenus*) and brine shrimp
(Winters et al., 2022). We hypothesised that chemically defended (unpalatable) species would be
found on visually distinct, more variable visual backgrounds than undefended (palatable) species.
Furthermore, we predicted a strong correlation between the presence and strength of chemical
defences and differences in spatiochromatic properties of background habitats among species with
differing strengths of chemical defences at distances where prey detection and identification are most
likely (Endler, 1986, 1991; Ruxton et al., 2018).

Materials and Methods

Study species. We took calibrated digital photographs of 184 Dorid nudibranch individuals belonging
to 12 species from four locations on the east coast of Australia: Sunshine Coast (SE Queensland, QLD),
Gold Coast (SE QLD), Cook Island (New South Wales, NSW) and Nelson Bay (NSW) between March
2016 and February 2021 (Table S1, Fig. S1). Our study considers many of the more commonly found
Dorid nudibranchs in the study sites (e.g. (Larkin et al., 2018; Schubert & Smith, 2020; Smith & Davis,
2019)). Species were identified visually using various nudibranch ID guides (Coleman et al., 2015;
Debelius & Kuiter, 2007; Gosliner et al., 2018): *Aphelodoris varia* (n = 22), *Chromodoris elisabethina* (n
= 21), *Chromodoris kuiteri* (n = 17), *Chromodoris lochi* (n = 3), *Dendrodoris krusensterni* (n = 7),
Discodoris sp. (n = 13), *Doriprismatica atromarginata* (n = 27), *Glossodoris vespa* (n = 15); *Hypselodoris*
bennetti (n = 10); *Phyllidia ocellata* (n = 23); *Phyllidia varicosa* (n = 8); *Phyllidiella pustulosa* (n = 18);
numbers indicate sample sizes used in the analysis. *Discodoris sp*. in our study may have constituted
a mixture of closely related species, including *Sebadoris fragilis*, *Jorunna pantheris, Tayuva lilacina* and
undescribed species; however, these species are visually indiscriminable. Therefore, they are named
Discodoris sp. as they were found in the same locations, have no known chemical defences, are
predominantly nocturnal and are closely related (Larkin et al., 2018). Only one out of the 12 species
(*Doriprismatica atromarginata*) was found in comparable numbers across all sites in NSW and SE QLD,
with most species found either in NSW or SE QLD (Table S1). Nudibranchs were photographed
underwater against their natural habitat using a calibrated Olympus EPL-5 with a 60mm macro lens in
an Olympus PT-EP10 underwater housing using white LED illumination from a combination of VK6r

and PV62 Scubalamp video lights (van den Berg, Troscianko, et al., 2020). All pictures were taken at

roughly a 90-degree angle relative to each animal and its background (Fig. 1).

Figure 1. Representative nudibranch and background images for 4 out of the 12 species in this study. From top to bottom: *Discodoris sp., Phyllidia ocellata, Chromodoris kuiteri, Doriprismatica atromarginata.*

Image analysis. Using ImageJ (Schneider et al., 2012) and the MICA toolbox (Troscianko & Stevens,

2015), the images were manually segmented into regions of interest (ROI), selecting the animal from

its background and defining a size standard using a custom-made colour and grey standard placed in

each image (Fig. 1). All images were aligned, so the nudibranch was aligned so the majority of each

animal's body was oriented vertically (head up) before analysis using QCPA (van den Berg, Troscianko,

et al., 2020). To analyse the images, we used the visual system parameters of a trichromatic triggerfish,

Rhinecanthus aculeatus (with spectral sensitivities of 413 nm, 480 nm and 528 nm λ_{max}; Cheney et al.

2022). This species (adult total length, TL ~ 15 cm) is a typical shallow reef inhabitant found throughout

the Indo-Pacific and feeds on invertebrates, algae, and detritus (Randall et al., 1997) and is deemed a

potential predator of nudibranchs. We analysed for viewing distances of 2 cm and 30 cm, assuming a

triggerfish spatial acuity of 3 cycles per degree (Champ et al., 2014). The viewing distances reflect a

range covering the maximal distance at which the largest specimen and coarse background detail

would be detectable (30cm) to the closest possible distance, reflecting ultimate contact between

predator and prey or visual background 2cm). Following acuity modelling, the images were processed

with a Receptor Noise Limited (RNL) ranked filter (falloff: 3, radius: 5, repetition: 5) and clustered using

RNL clustering with a colour discrimination threshold of 2 ΔS (Cheney et al., 2019) and a luminance

contrast threshold of 4 ΔS (van den Berg, Hollenkamp, et al., 2020). We used Weber fractions based

on a relative photoreceptor abundance of 1:2:2:2 (sw:mw:lw:dbl) and photoreceptor noise of 0.05,

resulting in 0.07:0.05:0.05:0.05.

QCPA analysis was achieved using a custom batch script (van den Berg, Condon, et al., 2023),

resulting in a highly descriptive array of 157 colour pattern statistics for each animal's visual

background. These parameters were spread across the following subtypes of colour pattern analysis:

1) Colour Adjacency analysis (CAA). CAA uses a transition matrix tallying all the synonymous and non-

synonymous transitions along horizontal and vertical sampling transects across an image segmented

into colour pattern elements. This transition matrix is then used to describe the geometry of a colour

pattern (Endler, 2012; van den Berg, Troscianko, et al., 2020). 2) Visual contrast analysis (VCA). VCA

uses the relative abundance and spectral properties of colour pattern elements to describe visual

contrast (Endler & Mielke, 2005; van den Berg, Troscianko, et al., 2020). 3) Boundary strength analysis

(BSA). BSA uses the relative abundance of boundaries between colour pattern elements to describe

visual contrast (Endler et al., 2018; van den Berg, Troscianko, et al., 2020). 4) Local edge intensity

analysis (LEIA). LEIA quantifies the strength and abundance of edge contrast in an unsegmented image.

Statistics ending with 'hrz' or 'vrt' are the horizontal (across body axis) and vertical (along body axis)

versions (analysing the respective transition matrix only) of their respective statistic (analysing the full

transition matrix). For example, a background with algae or seagrass will likely have an elongated

aspect ratio, whereas a uniform sandy background will not. A detailed description of each pattern

statistic can be found in van den Berg, Troscianko, et al. (2020).

Level of chemical defence. As a measure of chemical defence for each species, we used previously

published data from antifeedant assays with rockpool shrimp (*Palaemon serenus*) and toxicity assays

with brine shrimp (Winters et al., 2022). Assay data for *Glossodoris vespa* is presented in Winters *et*

al. (2018). In summary, the data were obtained from extracting secondary metabolites from

nudibranchs and added to food pellets made from squid mantle at increasing concentrations. Effective

dose (ED_{50}) values were calculated based on the concentration that elicited a rejection response in at

least 50% of the shrimp. Lethal dose (LD_{50}) values were calculated based on the concentration that

killed at least 50% of brine shrimp. For this study, the resulting ED_{50} and LD_{50} values were normalised

and calculated as $1-ED_{50}$ / $1-LD_{50}$ to range from 0 (most palatable/toxic) to 1 (least palatable/toxic).

Where multiple estimates of ED_{50} existed for a species due to multiple extracts/assays being

performed, the average value was used. Only assays using whole-body extracts were considered to

allow for comparisons between species.

Due to the uneven spread of toxicity and palatability values among species, we classified each

species according to the shape of a sigmoidal response curve where species with values of 0 were

considered palatable, values up to 0.25 were considered weakly unpalatable, values between 0.25

and 0.74 medium unpalatable and values of 0.74 and higher as highly unpalatable. We chose 0.74

rather than 0.75 as the boundary between medium and high unpalatability, as this was the median

unpalatability of species with chemical defences (Table S1). To ensure at least three species in each

group of chemical defences and to allow for subsequent investigations of differences in backgrounds

between species with different levels of chemical fences, we separated the species into those who

had no known toxicity and unpalatability: those with some level of toxicity and moderate

unpalatability and those with some level of toxicity and high unpalatability (Fig S1).

Statistical analysis

To analyse the large dataset derived from the QCPA analysis, we only kept images that did not

produce any missing value for any background pattern metrics. VCA, CAA, and BSA metrics can

produce NaN or infinite values if a colour pattern has less than two colour pattern elements following

RNL clustering [60]. LEIA metrics do not suffer from this limitation. We then applied a Latent variable

Exploratory Factor Analysis (EFA) with the R package *psych* using the factoring method of Ordinary

Least Squares "ols" and the orthogonal rotation "varimax". To prepare the dataset for the EFA, we

first filtered the number of highly correlated QCPA metrics by keeping those that were less correlated

than 0.6 Pearson correlation, which reduced their number from 157 to 17. We then ran the factor

analysis based on four factors. The number of factors was selected by comparing the eigenvalues

calculated from the original dataset to the median eigenvalues extracted from 10,000 randomly

generated datasets with the same rows and columns as the original data. We selected factors with

eigenvalues greater than the median of the eigenvalues from the simulated data. We also computed

bootstrapped confidence intervals of the loadings by iterating the factor analysis 1000 times.

Due to data filtering for metrics less correlated than 0.6, the QCPA parameter listed for a given

loading is likely synonymous with various other parameters in our 157-dimension colour pattern

space. Therefore, the precise wording to describe each factor can vary substantially depending on

which colour pattern metrics are put into focus (for a complete list of parameter correlations, see

Table S2).

The scores of the factors extracted from the EFA were then used to implement four

phylogenetic, distributional linear mixed models to compare the natural backgrounds of nudibranchs

with different levels of chemical defences. Models were run in R v 4.1.2 (https://www.r-

project.org/) using the brms package (Bürkner, 2018), which fits Bayesian models using

Stan (https://mc-stan.org/). To account for the phylogenetic dependency of closely related species, all

models included the phylogenetic tree of the 12 species of nudibranchs (Fig. S2), with the tree from

(Cheney et al., 2014) pruned and missing species added according to their taxonomic classification in

the World Register of Marine Species (Appeltans et al., 2012). The phylogenetic model was

implemented   following   the   guidelines   of   the   online   brms   vignette

(https://cran.rproject.org/web/packages/brms/vignettes/brms_phylogenetics.html) based on de

Villemereuil & Nakagawa (de Villemereuil & Nakagawa, 2014).

The first model investigated differences in scores for latent factor 1 between nudibranchs with

different levels of chemical defences (see chemical defences section) using a Student distribution. The

model estimated the effect of the main categorial predictors level of chemical defence (low,

moderate, high) and observer distance (2 cm and 30 cm) and their interaction on the response

distribution's mean and the residual standard deviation. To account for repeated measurements of

each species, we also included species ID as a random intercept to the model. We further included

random slopes over distance because their relationship with the value of the response factor 1

changes among species.

The second, third and fourth models were identical to the first but used factor 2, factor 3 and

factor 4 as response variables. All models were fitted using weakly informative prior distributions

(normal with mean = 0 and sd = 5 for intercept and coefficients, exponential (1) for standard

deviations). Their performance was evaluated using posterior predictive model checking, which

compares model predictions with observed data to assess overall model fit. We ran four Markov-

Chain-Monte-Carlo (MCMC) chains for each model and obtained coefficient estimates from 16,000

post-warm-up samples. All model parameters reached reliable conversion indicators (Korner-

Nievergelt et al., 2015): A Monte Carlo standard error smaller than 5% of the posterior standard

deviation, an effective posterior sample size greater than 10% of the total sample size, and a $\hat{R}$ statistic

value smaller than 1.01.

For graphical displays of the results, we present - for each combination of chemical defences

and distance - the medians of latent factors values and their 95% Credible Intervals (CIs) of the

posterior distributions of fitted values for the population average obtained from the joint posterior

distributions of the model parameters (Korner-Nievergelt et al., 2015). The same posterior distribution

of fitted values was used to compute pairwise differences and their 95% CIs between all possible

combinations of the same two categorical predictors using the emmeans R package (Lenth, 2023). To

compare the variance of response values between all combinations of predictor levels, we also

compute the posterior distribution of all pairwise differences of the residual standard deviation on

the original scale (back-transformed from the log scale). The effect size of pairwise differences

increases with increasing deviation of such differences from zero, and the robustness of the result

increases with decreasing degree of overlap of the 95% Credible Intervals (CIs) with zero.

Results

Using the EFA, we identified four factors to describe how background pattern metrics differ between

multiple viewing distances and levels of chemical defences. While not intended to identify a maximal

amount of variability in colour pattern variation in our dataset, the four factors still explain 39% of

the total variation (factor 1: 11%; factor 2: 11%, factor 3: 11%, factor 4: 6%) (Fig 2). Looking at the

loadings of each factor, we can identify what latent variable they describe. While it would be possible

to discuss each factor in almost infinite depth, we keep their description to loadings of +/- 0.4 to

capture their main properties.

<u>Factor 1:</u> *Higher luminance and chromatic contrast coincides with smaller patch size*

Factor 1 (Fig. 2a) describes 11% of variation in visual backgrounds and captures a positive relationship

between increases in the mean RNL luminance edge contrast (e.g., *Lum.mean*) and RNL chromaticity

edge contrast (e.g. *Col.mean*) as measured by LEIA (unclustered image) and the average size of colour

pattern elements in an RNL clustered image (*CAA.PT*). The increase in average luminance and

chromatic edge contrast in visual backgrounds also coincides with an increase in variability of the

luminance contrast relative to the mean (e.g. *BSA.BCVL*).

The visual backgrounds of chemically defended species have noticeably higher levels of factor

1 than those of undefended species [difference (+- 95% CI)] (Fig 2a). This is true up close (2cm) as well

as from further away (30cm) for toxic species with moderate levels of unpalatability (2cm: -1.53 (-2.28

/ -0.78); 30cm: -1.05 (-1.83 / -0.23)) as well as toxic species with high levels of unpalatability (2cm: -

1.44 (-2.18 / -0.73); 30cm -1.17 (-1.92 / -0.37)). However, toxic species with moderate levels of

unpalatability have similar factor values to species with high levels of unpalatability up close (2cm:

0.08 (-0.38 / 0.54); 30cm: -0.13 (-0.63 / 0.36)). Therefore, chemically defended species were found on

visual backgrounds with more chromatic and achromatic contrast (Table S3). However, the variability

of backgrounds on which species were found did not differ between groups and remained similar at

2cm and 30cm (Table S4).

Factor 2: *Higher colour and luminance contrast between patches coincide with lower background homogeneity*

Factor 2 (Fig. 2b) describes 11% of the variation in visual backgrounds and mostly captures the correlation between simultaneous increases in the strength of luminance contrast (e.g. *BSA.BML, VCA.MSL*) and saturation contrast (e.g. *BSA.BMS*) between colour pattern elements and decreases in background evenness (e.g. *CAA.Qt* & *CAA.Qc*).

We found no differences in factor values between species with different levels of chemical defences [difference (+- 95% CI)] (Fig. 2b,Table S5) nor any differences between groups in the variability of factor values (Table S6) at 2cm or 30cm viewing distance. All groups showed higher factor values at 2cm compared to 30cm (Fig. 2b, undefended: 1.01 (0.57 / 1.48); toxic moderately unpalatable: 1.01 (0.67 / 1.31); toxic highly unpalatable: 0.80 (0.47 / 1.11)). The variability of backgrounds on which species were found did not differ between groups and remained similar at 2cm and 30cm (Table S6).

Factor 3: *Increased luminance and colour contrast variability correlate with reduced average colour pattern regularity*

Factor 3 captures the relationship between increases in the variability of luminance contrast in visual backgrounds (e.g. *Lum.kurtosis, Lum.CoV*) as well as the variability of colour contrast (e.g. *Col.kurtosis, VCA.CVS*) and a decrease of the average colour contrast (e.g. *Col.mean*) as well as colour pattern regularity (e.g. *CAA.Qc*).

We found no differences in factor values of backgrounds between groups with different levels of chemical defences [difference (+- 95% CI)] (Fig. 2c, Table S7) at either 2cm or 30cm. However, all groups showed increased factor values at 2cm compared to 30cm (Fig 2c; undefended: (0.50 (0.07 / 0.91)); toxic moderate unpalatable: toxic highly unpalatable species (0.34 (0.08 / 0.59)). No difference in the variability of factor values was detected between the background of groups at either viewing distance, and background variability was the same at 2cm and 30cm for all groups (Table S8).

Factor 4: *Increases in achromatic patch contrast correlate with decreases in achromatic boundary contrast, but increases in chromatic boundary contrast variability*

Factor 4 (Fig. 3d) describes a correlation between achromatic average patch contrast (e.g. *VCA.ML*) and measures of achromatic boundary contrast (e.g. *Lum.mean, BSA.BML*). Specifically, increases in mean luminance contrast relative to the size of colour pattern elements coincide with a decrease in the average luminance contrast of boundaries between colour pattern elements. Furthermore,

decreases in the mean luminance contrast of pattern boundaries correlate with an increase in

chromatic boundary contrast variability relative to the mean (e.g. *BSA.BCVSsat*).

We did not find any differences in factor values between the backgrounds of species with

different levels of chemical defences at either 2cm, or 30cm [difference (+- 95% CI)] (Fig. 2d, Table S9).

However, while undefended species showed no difference in factor values of backgrounds between

viewing distances (0.21 (-0.25 / 0.69)), both toxic and medium unpalatable species (0.53 (0.25 / 0.81)),

as well as toxic and highly unpalatable species (0.42 (0.16 / 0.70)), had higher factor values for their

visual backgrounds at 2cm compared to 30cm. No differences in background variability between

groups were detected at either viewing distance (Table S10), except for undefended species having

more variable backgrounds at 30cm compared to toxic and moderately defended species (0.45 (0.07

/ 0.90). The variability of visual backgrounds did not differ between viewing distances for either group

(Table S10).

Figure 2 Detailed visual representation of the loadings of factor 1 (A), factor 2 (C), factor 3 (E), and factor 4 (G). Greyed-out factor loadings indicate colour pattern descriptors with minor contributions (<0.4) to each factor. Differences between group estimates of factor values are given for factor 1 (B), factor 2(D), factor 3 (F), and factor 4 (H). Estimates are given for 2cm viewing distance (left panel half, white) and 30cm (right panel half, grey).

Discussion

Our analysis captured four latent variables describing correlations between the spatiochromatic properties of visual backgrounds on which animals from 12 species of nudibranch molluscs were found (Fig. 2). In agreement with one of our predictions derived from the 'escape and radiate' hypothesis, these latent variables showed that nudibranch species with chemical defences, irrespective of their relative strength, were found on visual backgrounds distinct in their appearance from backgrounds of undefended species according to the physiological limitations of a potential predator, a triggerfish (*Rhinecanthus aculeatus*) (Fig. 2a). However, while some visual properties of backgrounds varied significantly more up close (2cm) than from further away (30cm), we did not find any differences in the variability of visual backgrounds between species with different levels of chemical defences. This lack of among-species variability indicates that chemical defences and their relative strength do not correspond to increased among-species background variability in our dataset, contrary to the predictions derived from the 'escape and radiate' hypothesis (Merilaita & Tullberg, 2005; Thompson, 1989). Instead, we suggest that chemical defences in Dorid nudibranchs coincide with broad, yet equally variable, differences in visual background habitats.

As shown by the composition of each factor (Fig. 2), multicomponent descriptors of complex phenotypes can be difficult to reduce to a low-dimensional, intuitive, spatiochromatic property that adequately captures the underlying complexity and links it to perceptual and functional properties of colouration. Unlike numerical classifiers from artificial neuronal networks or other machine learning approaches used in high-dimensional image analyses for computer vision (e.g. Bileschi et al., 2007; Talas et al., 2020)), each parameter in our colour pattern space describes a specific spatiochromatic property (van den Berg, Troscianko, et al., 2020). Therefore, despite a range of complexity in associations with response variables (see, e.g. van den Berg et al., 2022) for discussion), we can make assumptions about the perceptual processes associated with our latent predictors.

Visual backgrounds of chemically defended species in our study can be broadly characterised by the presence of increased colour and luminance contrast between objects and surfaces when compared to visual backgrounds of undefended species (Fig. 2a). This difference in the visual appearance of background habitats to a potential predator was equally as strong when viewed from 2cm than at 30cm. Therefore, backgrounds of chemically defended species remained more spatiochromatically contrasting than those of undefended species (Fig. 2a), even at distances where substantial amounts of spatial information would be lost to a potential predator. This persistence of spatiochromatic variability in visual backgrounds across viewing distances is of interest in the context of distance-dependent selection pressures shaping the ecology and evolution of colour pattern

diversity in prey communities (e.g. Barnett et al., 2016; Endler, 1978; van den Berg et al., 2022; van

den Berg, Endler, et al., 2023). For example, imperfect mimicry among- and colour pattern

polymorphism within aposematic species could be shaped by their adaptive value in the context of

predator perception at multiple viewing distances. Persisting variability of visual backgrounds across

viewing distances could be selecting for perceptual similarities between animals and backgrounds in

a more general way than previously considered.

Aposematism is assumed to be widespread in Dorid nudibranchs (Rudman, 1991), with bold

colours and patterns coinciding with the presence of chemical defences (Cortesi & Cheney, 2010;

Winters et al., 2022; Winters, Wilson, et al., 2018). However, the coincidence of secondary defences

and boldly contrasting animal colouration (e.g. van den Berg, Endler, et al., 2023) does not mean that

the colours and patterns displayed by a species necessarily serve a warning function (see Summers et

al., 2015; White & Umbers, 2021) for review). Instead, the function of bold colouration in an

environment with increased colour and luminance contrast is likely complex and might even assist in

camouflage (Endler, 1978; Marshall & Stevens, 2014; van den Berg, Endler, et al., 2023). The

detectability of an aposematic animal is determined by its appearance against its visual background

and is likely fine-tuned to the cost-benefit trade-offs of increased predator encounters (e.g. Barnett

et al., 2016; van den Berg, Endler, et al., 2023). Therefore, assuming substantial (yet variable) degrees

of signalling honesty in Dorid nudibranchs (e.g. Cortesi & Cheney, 2010), our study suggests broadly

generalisable correlations between the presence of chemical defences in aposematic species and the

spatiochromatic properties of their visual habitat. However, these constraints can be variably

explained by the need for efficient camouflage (e.g. Barnett et al., 2016; Speed, Ruxton, et al., 2010)

or signalling efficacy (e.g. Endler & Mappes, 2004; Speed, Brockhurst, et al., 2010) at variable viewing

distances. How specific spatiochromatic properties of visual backgrounds highlighted by our study

impact the signalling function of aposematic colouration and camouflage in the considered species

would be of great interest in future behavioural experimentation studies.

Nudibranchs mainly use chemotaxis to move around their environment to find food and

mates, with visual input only relevant for basic phototaxis, such as determining daytime or detecting

shelter (Eakin et al., 1967). Therefore, unlike other aposematic species, such as insects or frogs

(Higginson et al., 2012; Rojas, 2017), nudibranchs will unlikely choose resting and foraging

microhabitats based on visual cues. Therefore, it is likely that the backgrounds on which species are

found are the indirect rather than direct consequence of correlations between their habitat's visual

appearance and other sensory modes and selective pressures shaping the efficacy of visual defences.

Thus, finding chemically defended species on distinct visual backgrounds fits well with assumed

radiation in feeding ecology, enabling the acquisition of secondary defences (e.g. Winters, White, et
al., 2018).

To our knowledge, no study has quantified the mobility and the diets of many nudibranch
species (but see Rudman & Bergquist, 2007), and the spatiotemporal distribution of food sources
remains poorly known. Some of the most unpalatable nudibranchs in this study are known to forage
on sponges containing highly potent secondary metabolites (e.g. latrunculin a) at least at some stage
during their lifetime (e.g. *Chromodoris elisabethina*) (Cheney et al., 2016). In contrast, others can
synthesise compounds *de novo* (Cimino & Ghiselin, 1999). Nudibranchs with more specialist diets may
need to increase mobility to find uncommon food sources. The sponge diet consequently allows the
animals to maintain functional levels of secondary metabolites or pigments required to maintain
salient aposematic colour patterns. Background specialisation might, therefore, be increased in
species dependent on such supplemental food sources and could influence within and among-species
background variability. However, whether the presence of secondary defences in the context of visual
background habitats also coincides with changes in defensive animal colouration remains an exciting
and crucial avenue for future research.

The mechanisms underlying the 'escape and radiate' hypothesis are often unclear, with little
empirical evidence of the process (e.g. Suchan & Alvarez, 2015). For example, evidence of increased
background variability could only be present during certain stages of (rapid) speciation in aposematic
animals. More extensive sampling of primary and secondary defences and corresponding visual
backgrounds within and among species, considering each species' geographic distribution, would be
of great interest for future research. Combined with an increased resolution of existing heterobranch
phylogenies (e.g. Layton et al., 2018, 2020), this would enable a more detailed investigation into the
presence and scale of animal background and colour pattern diversification and its correlation with
secondary defences.

The direction, speed and extent by which natural selection would constrain correlations
between the strength of secondary defences and background habitat is likely determined by multiple,
currently poorly understood factors such as the fitness benefits of specific colour patterns across
different visual backgrounds, the diversity of visual backgrounds inhabited by a given species, the
protective value of secondary compounds across complex predator communities and the heritability
of visual phenotypes, to name a few. Nevertheless, our study provides empirical support to the
possibility of secondary defences coinciding with generalisable differences in visual background
habitats in a complex community of nudibranch molluscs. We demonstrate this by employing
sophisticated methodology aimed at reflecting visual processing of ecologically relevant observers and

utilising a high-dimensional, highly differentiated approach to quantifying spatiochromatic properties of visual backgrounds.

Acknowledgements

We thank various volunteers for assistance with fieldwork and image analysis. We are grateful for the High-Performance Computing (HPC) infrastructure at The University of Queensland (Wiener & Awoonga) and Dr. Simone Blomberg, who provided infrastructure that contributed to the computing of image statistics.

Conflict of Interest

The authors have no conflict of interest to declare.

Data availability

The data is available on UQ's digital repository eSpace: https://doi.org/10.48610/8f7b70e

Author contributions

CPvdB conceived the original study with KLC & JAE providing critical input; CPvdB collected and analysed the data; CPvdB and KLC led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Funding

This work was funded by the Australian Research Council (FT190199313 and DP180102363 awarded to KLC and JAE), two Holsworth Wildlife Research Endowment grants awarded to CPvdB, a Swiss National Foundation Postdoc.Mobility Fellowship (P500PB_211070) awarded to CPvdB, a research grant from the Society of Conchologists awarded to CPvdB, and a student research grant from the Australasian Society for the Study of Animal Behaviour awarded to CPvdB. MS was supported by a MSCA 2021 postdoctoral fellowship (101066328) funded via the Engineering and Physical Sciences Research Council [grant number EP/X020819/1].

References

Appeltans, W., Bouchet, P., Boxshall, G. A., De Broyer, C., de Voogd, N. J., Gordon, D. P., Hoeksema,

B. W., Horton, T., Kennedy, M., Mees, J., Poore, G. C. B., Read, G., Stöhr, S., Walter, T. C., &

Costello, M. J. (eds). (2012). *World Register of Marine Species*. http://www.marinespecies.org

Arbuckle, K., & Harris, R. J. (2021). Radiating pain: venom has contributed to the diversification of

the largest radiations of vertebrate and invertebrate animals. *BMC Ecology and Evolution*,

21(1), 1–12. https://doi.org/10.1186/s12862-021-01880-z

Arbuckle, K., & Speed, M. P. (2015). Antipredator defenses predict diversification rates. *Proceedings*

of the National Academy of Sciences of the United States of America, *112*(44), 13597–13602.

https://doi.org/10.1073/pnas.1509811112

Avila, C. (1995). Natural products of opisthobranch molluscs: a biological review. *Oceanography and*

Marine Biology: An Annual Review, *33*, 487–559.

Barnett, J. B., Scott-Samuel, N. E., & Cuthill, I. C. (2016). Aposematism: balancing salience and

camouflage. *Biology Letters*, *12*(8), 20160335. https://doi.org/10.1098/rsbl.2016.0335

Bürkner, P.-C. (2018). Advanced Bayesian Multilevel Modeling with the R Package brms. *The R*

Journal, *10*(1), 395. https://doi.org/10.32614/RJ-2018-017

Carbone, M., Gavagnin, M., Haber, M., Guo, Y.-W., Fontana, A., Manzo, E., Genta-Jouve, G.,

Tsoukatou, M., Rudman, W. B., Cimino, G., Ghiselin, M. T., & Mollo, E. (2013). Packaging and

Delivery of Chemical Weapons: A Defensive Trojan Horse Stratagem in Chromodorid

Nudibranchs. *PLoS ONE*, *8*(4), e62075. https://doi.org/10.1371/journal.pone.0062075

Champ, C. M., Wallis, G., Vorobyev, M., Siebeck, U., & Marshall, J. (2014). Visual acuity in a species of

coral reef fish: *Rhinecanthus aculeatus*. *Brain, Behavior and Evolution*, *83*(1), 31–42.

https://doi.org/10.1159/000356977

Cheney, K. L., Cortesi, F., How, M. J., Wilson, N. G., Blomberg, S. P., Winters, A. E., Umanzör, S., &

Marshall, N. J. (2014). Conspicuous visual signals do not coevolve with increased body size in

marine sea slugs. *Journal of Evolutionary Biology*, *27*(4), 676–687.

https://doi.org/10.1111/jeb.12348

Cheney, K. L., Green, N. F., Vibert, A. P., Vorobyev, M., Marshall, N. J., Osorio, D. C., & Endler, J. A.

(2019). An Ishihara-style test of animal colour vision. *Journal of Experimental Biology*, *222*(1),

jeb189787. https://doi.org/10.1242/jeb.189787

Cheney, K. L., White, A., Mudianta, I. W., Winters, A. E., Quezada, M., Capon, R. J., Mollo, E., &

Garson, M. J. (2016). Choose your weaponry: Selective storage of a single toxic compound,

latrunculin a, by closely related nudibranch molluscs. *PLoS ONE*, *11*(1), 1–16.

https://doi.org/10.1371/journal.pone.0145134

Cimino, G., & Ghiselin, M. T. (1999). Chemical defense and evolutionary trends in biosynthetic

capacity among dorid nudibranchs (Mollusca: Gastropoda: Opisthobranchia). *Chemoecology*,

9(4), 187–207. https://doi.org/10.1007/s000490050052

Coleman, N., Cobb, G., & Mullins, D. (2015). *Nudibranchs Encyclopedia*. Thomas Slone, Masalai Press

and Tim Hochgrebe.

Cortesi, F., & Cheney, K. L. (2010). Conspicuousness is correlated with toxicity in marine

opisthobranchs. *Journal of Evolutionary Biology*, *23*(7), 1509–1518.

https://doi.org/10.1111/j.1420-9101.2010.02018.x

de Villemereuil, P., & Nakagawa, S. (2014). General Quantitative Genetic Methods for Comparative

Biology. In L. Z. Garamszegi (Ed.), *Modern Phylogenetic Comparative Methods and Their*

Application in Evolutionary Biology: Concepts and Practice (pp. 287–303). Springer Berlin

Heidelberg. https://doi.org/10.1007/978-3-662-43550-2_11

Debelius, H., & Kuiter, R. H. (2007). *Nudibranchs of the world*. IKAN-Unterwasserarchiv.

Eakin, R. M., Westfall, J. A., & Dennis, M. J. (1967). Fine Structure of the Eye of a Nudibranch

Mollusc, *Hermissenda Crassicornis*. *Journal of Cell Science*, *2*(3), 349–358.

https://doi.org/10.1242/jcs.2.3.349

Ehrlich, P. R., & Raven, P. H. (1964). Butterflies and Plants: A Study in Coevolution. *Evolution*, *18*(4),

586. https://doi.org/10.2307/2406212

Endler, J. A. (1978). A predator's view of animal color patterns. *Evolutionary Biology*, *11*(5), 320–364.

https://doi.org/10.1007/978-1-4615-6956-5_5

Endler, J. A. (1984). Progressive background in moths, and a quantitative measure of crypsis.

Biological Journal of the Linnean Society, *22*(3), 187–231. https://doi.org/10.1111/j.1095-

8312.1984.tb01677.x

Endler, J. A. (1986). Defense Against Predators. In *Predator-prey relationships, Perspectives and*

Approaches from the Study of Lower Vertebrates (pp. 109–134). University of Chicago Press.

Endler, J. A. (1991). Interactions between predators and prey. In J. R. Krebs & N. B. Davies (Eds.),

Behavioural Ecology (pp. 169–196). Blackwell Scientific.

Endler, J. A. (2012). A framework for analysing colour pattern geometry: Adjacent colours. *Biological*

Journal of the Linnean Society, *107*(2), 233–253. https://doi.org/10.1111/j.1095-

8312.2012.01937.x

Endler, J. A., Cole, G. L., & Kranz, A. M. (2018). Boundary strength analysis: Combining colour pattern

geometry and coloured patch visual properties for use in predicting behaviour and fitness.

Methods in Ecology and Evolution, *9*(12), 2334–2348. https://doi.org/10.1111/2041-

210X.13073

Endler, J. A., & Mappes, J. (2004). Predator Mixes and the Conspicuousness of Aposematic Signals.

The American Naturalist, *163*(4), 532–547. https://doi.org/10.1086/382662

Endler, J. A., & Mielke, P. W. (2005). Comparing entire colour patterns as birds see them. *Biological*

Journal of the Linnean Society, *86*(4), 405–431. https://doi.org/10.1111/j.1095-

8312.2005.00540.x

Gosliner, T. M. (2001). Aposematic coloration and mimicry in opisthobranch mollusks: new

phylogenetic and experimental data. *Bollettino Malacologico*, *37*(37), 163–170.

Gosliner, T. M., Valdes, A., & Behrens, D. W. (2018). *Nudibranch & Sea Slug Identification - Indo-*

Pacific (2nd ed.). New World Publications.

Higginson, A. D., Speed, M. P., & Ruxton, G. D. (2012). Effects of anti-predator defence through toxin

sequestration on use of alternative food microhabitats by small herbivores. *Journal of*

Theoretical Biology, *300*, 368–375. https://doi.org/10.1016/j.jtbi.2012.01.020

Korner-Nievergelt, F., Roth, T., von Felten, S., Guélat, J., Almasi, B., & Korner-Nievergelt, P. (2015).

Bayesian Data Analysis in Ecology Using Linear Models with R, BUGS, and STAN. Elsevier.

https://doi.org/10.1016/C2013-0-23227-X

Larkin, M. F., Smith, S. D. A., Willan, R. C., & Davis, T. R. (2018). Diel and seasonal variation in

heterobranch sea slug assemblages within an embayment in temperate eastern Australia.

Marine Biodiversity, *48*(3), 1541–1550. https://doi.org/10.1007/s12526-017-0700-9

Layton, K. K. S., Carvajal, J. I., & Wilson, N. G. (2020). Mimicry and mitonuclear discordance in

nudibranchs: New insights from exon capture phylogenomics. *Ecology and Evolution*, *10*(21),

11966–11982. https://doi.org/10.1002/ece3.6727

Layton, K. K. S., Gosliner, T. M., & Wilson, N. G. (2018). Flexible colour patterns obscure identification

and mimicry in Indo-Pacific Chromodoris nudibranchs (Gastropoda: Chromodorididae).

Molecular Phylogenetics and Evolution, *124*(February), 27–36.

https://doi.org/10.1016/j.ympev.2018.02.008

Lenth, R. V. (2023). *emmeans: Estimated Marginal Means, aka Least-Squares Means* (1.8.8).

https://cran.r-project.org/package=emmeans

Marshall, K. L. A., & Stevens, M. (2014). Wall lizards display conspicuous signals to conspecifics and

reduce detection by avian predators. *Behavioral Ecology*, *25*(6), 1325–1337.

https://doi.org/10.1093/beheco/aru126

Merilaita, S., & Tullberg, B. S. (2005). Constrained camouflage facilitates the evolution of

conspicuous warning coloration. *Evolution*, *59*(1), 38–45. https://doi.org/10.1111/j.0014-

3820.2005.tb00892.x

Nokelainen, O., Scott-Samuel, N. E., Nie, Y., Wei, F., & Caro, T. (2021). The giant panda is cryptic.

Scientific Reports, *11*(1), 1–10. https://doi.org/10.1038/s41598-021-00742-4

Poulton, E. B. (1890). *The Colours of Animals*. Kegan Paul, Trench & Trubner.

Randall, J. E., Allen, G. R., & Steene, R. C. (1997). *Fishes of the Great Barrier Reef and Coral Sea*.

Crawford House Publishing.

Regan, D., & Beverley, K. I. (1984). Figure–ground segregation by motion contrast and by luminance

contrast. *Journal of the Optical Society of America A*, *1*(5), 433.

https://doi.org/10.1364/josaa.1.000433

Rojas, B. (2017). Behavioural, ecological, and evolutionary aspects of diversity in frog colour

patterns. *Biological Reviews*, *92*(2), 1059–1080. https://doi.org/10.1111/brv.12269

Rudman, W. B. (1991). Purpose in pattern - the evolution of color in chromodorid Nudibranchs.

Journal of Molluskan Studies, *57*(1991), 5–21.

Rudman, W. B., & Bergquist, P. R. (2007). A review of feeding specificity in the sponge-feeding

Chromodorididae (Nudibranchia: Mollusca). *Molluscan Research*, *27*(2), 60–88.

Ruxton, G. D., Allen, W. L., Sherratt, T. N., & Speed, M. P. (2018). *Avoiding Attack* (Vol. 1). Oxford

University Press. https://doi.org/10.1093/oso/9780199688678.001.0001

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image

analysis. *Nature Methods*, *9*(7), 671–675. https://doi.org/10.1038/nmeth.2089

Schubert, J., & Smith, S. D. A. (2020). Sea slugs—"rare in space and time"—but not always. *Diversity*,
12(11), 1–14. https://doi.org/10.3390/d12110423

Smith, S. D. A., & Davis, T. R. (2019). Slugging it out for science: volunteers provide valuable data on
the diversity and distribution of heterobranch sea slugs. *Molluscan Research*, *39*(3), 214–223.
https://doi.org/10.1080/13235818.2019.1594600

Speed, M. P., Brockhurst, M. A., & Ruxton, G. D. (2010). The dual benefits of aposematism: Predator
avoidance and enhanced resource collection. *Evolution*, *64*(6), 1622–1633.
https://doi.org/10.1111/j.1558-5646.2009.00931.x

Speed, M. P., Ruxton, G. D., Blount, J. D., & Stephens, P. A. (2010). Diversification of honest signals in
a predator-prey system. *Ecology Letters*, *13*(6), 744–753. https://doi.org/10.1111/j.1461-
0248.2010.01469.x

Suchan, T., & Alvarez, N. (2015). Fifty years after Ehrlich and Raven, is there support for plant-insect
coevolution as a major driver of species diversification? *Entomologia Experimentalis et*
Applicata, *157*(1), 98–112. https://doi.org/10.1111/eea.12348

Summers, K., Speed, M. P., Blount, J. D., & Stuckert, A. M. M. (2015). Are aposematic signals honest?
A review. *Journal of Evolutionary Biology*, *28*(9), 1583–1599. https://doi.org/10.1111/jeb.12676

Thompson, J. N. (1989). Concepts of coevolution. *Trends in Ecology and Evolution*, *4*(6), 179–183.
https://doi.org/10.1016/0169-5347(89)90125-0

Troscianko, J., & Stevens, M. (2015). Image calibration and analysis toolbox – a free software suite
for objectively measuring reflectance, colour and pattern. *Methods in Ecology and Evolution*,
6(11), 1320–1331. https://doi.org/10.1111/2041-210X.12439

van den Berg, C. P., Condon, N. D., Conradsen, C. I., White, T. E., & Cheney, K. L. (2023). Automated
workflows using Quantitative Colour Pattern Analysis (QCPA): A guide to batch processing and
downstream data analysis. *BioRxiv*.
https://doi.org/https://doi.org/10.1101/2023.02.02.526788

van den Berg, C. P., Endler, J. A., & Cheney, K. L. (2023). Signal detectability and boldness are not the
same: the function of defensive coloration in nudibranchs is distance-dependent. *Proceedings*
of the Royal Society B: Biological Sciences, *290*(2003), 2022.12.20.521213.
https://doi.org/10.1098/rspb.2023.1160

van den Berg, C. P., Endler, J. A., Papinczak, D. E. J., & Cheney, K. L. (2022). Using colour pattern edge
contrast statistics to predict detection speed and success in triggerfish (*Rhinecanthus*

aculeatus). *Journal of Experimental Biology*. https://doi.org/10.1242/jeb.244677

van den Berg, C. P., Hollenkamp, M., Mitchell, L. J., Watson, E. J., Green, N. F., Marshall, N. J., &

Cheney, K. L. (2020). More than noise: Context-dependant luminance contrast discrimination in

a coral reef fish (*Rhinecanthus aculeatus*). *Journal of Experimental Biology*, *223*(21),

jeb.232090. https://doi.org/10.1242/jeb.232090

van den Berg, C. P., Troscianko, J., Endler, J. A., Marshall, N. J., & Cheney, K. L. (2020). Quantitative

Colour Pattern Analysis (QCPA): A comprehensive framework for the analysis of colour patterns

in nature. *Methods in Ecology and Evolution*, *11*(2), 316–332. https://doi.org/10.1111/2041-

210X.13328

Wägele, H., Ballesteros, M., & Avila, C. (2006). Defensive Glandular Structures In Opisthobranch

Molluscs — From Histology To Ecology. In *Oceanography and Marine Biology* (Vol. 44, Issue

June, pp. 197–276). https://doi.org/10.1201/9781420006391.ch5

White, T. E., & Umbers, K. D. L. (2021). Meta-analytic evidence for quantitative honesty in

aposematic signals. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1949).

https://doi.org/10.1098/rspb.2021.0679

Winters, A. E., Chan, W., White, A. M., Berg, C. P., Garson, M. J., & Cheney, K. L. (2022). Weapons or

deterrents? Nudibranch molluscs use distinct ecological modes of chemical defence against

predators. *Journal of Animal Ecology*, *91*(4), 831–844. https://doi.org/10.1111/1365-

2656.13643

Winters, A. E., White, A. M., Dewi, A. S., Mudianta, I. W., Wilson, N. G., Forster, L. C., Garson, M. J., &

Cheney, K. L. (2018). Distribution of Defensive Metabolites in Nudibranch Molluscs. *Journal of*

Chemical Ecology, *44*(4), 384–396. https://doi.org/10.1007/s10886-018-0941-5

Winters, A. E., Wilson, N. G., van den Berg, C. P., How, M. J., Endler, J. A., Marshall, N. J., White, A.

M., Garson, M. J., & Cheney, K. L. (2018). Toxicity and taste: unequal chemical defences in a

mimicry ring. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1880), 20180457.

https://doi.org/10.1098/rspb.2018.0457

Table S1

Summary table of all individuals in the dataset used for this study

Species	Unpalatability (1-ED_{50})	Toxicity (1-LD_{50})	Sampling site			
			Sunshine Coast, QLD	Gold Coast, QLD	Nelson Bay, NSW	Grand Total
Aphelodoris varia	0	0	0	0	22	22
Chromodoris elisabethina	0.55	1	21	0	0	21
Chromodoris kuiteri	0.74	1	10	7	0	17
Chromodoris lochi	0.61	1	3	0	0	3
Dendrodoris krusensterni	0	0	0	2	5	7
Discodoris sp.	0	0	0	0	13	13
Doriprismatica atromarginata	0.74	0.93	16	0	11	27
Glossodoris vespa	0.87	0.29	15	0	0	15
Hypselodoris bennetti	0.73	0.88	0	0	10	10
Phyllidia ocellata	0.63	0.4	23	0	0	23
Phyllidia varicosa	0.92	0.53	8	0	0	8
Phyllidiella pustulosa	0.86	0.14	18	0	0	18

Figure S1

Figure S1. Representative photographs of the 12 species used in this study grouped into categories of chemical defences based on whole-body extract assays with palaemon shrimp to assess unpalatability (1-Effective Dose, ED_{50}) and brine shrimp to assess toxicity (1-Lethal Dose, LD_{50}) values: A) *Aphelodoris varia;* B) *Dendrodoris krusensterni*; C) *Discodoris sp*; D) *Chromodoris elisabethina*; E) *Chromodoris lochi*; F) *Hypselodoris bennetti;* G) *Phyllidia ocellata*; H) *Chromodoris kuiteri*; I) *Doriprismatica atromarginata;* J) *Glossodoris vespa*; K) *Phyllidia varicosa;* L) *Phyllidiella pustulosa.*

Figure S2

Phylogenetic tree used in this study, modified from Cheney et al. 2014 as described in 'Methods'.

Table S2

Summary table of all >.6 Pearson correlation values between parameters with >0.4 loadings for each factor.

FACTOR 1			FACTOR 2			FACTOR 3			FACTOR 4		
Factor loading >0.4	Correlated Param. (>.6)	Cor	Factor loading >0.4	Correlated Param. (>.6)	Cor	Factor loading of >0.4	Correlated Param. (>.6)	Cor	Factor loading of >0.4	Correlated Param. (>.6)	Cor
BSA.BCVSsat.Hrz	BSA.BCVDmax	0.80	BSA.BML.Vrt	BSA.BML	1.00	BSA.BCVL.Hrz	CAA.St	0.61	BSA.BCVL.Hrz	CAA.St	0.61
	BSA.BCVSsat	1.00		BSA.BMSL	1.00		CAA.Ht	0.68		CAA.Ht	0.68
	BSA.BCVDmax.Hrz	0.80		BSA.BML.Hrz	1.00		CAA.St.Hrz	0.62		CAA.St.Hrz	0.62
	BSA.BCVDmax.Vrt	0.79		BSA.BMSL.Hrz	0.99		CAA.St.Vrt	0.61		CAA.St.Vrt	0.61
	BSA.BCVSsat.Vrt	1.00		BSA.BMSL.Vrt	1.00		CAA.Ht.Hrz	0.68		CAA.Ht.Hrz	0.68
VCA.ML.Vrt	VCA.ML	1.00	BSA.BMS.Hrz	BSA.BMS	1.00		CAA.Ht.Vrt	0.68		CAA.Ht.Vrt	0.68
	VCA.sL	0.67		BSA.BMS.Vrt	1.00		CAA.Qt.Hrz	0.63		CAA.Qt.Hrz	0.63
	VCA.MDmax	0.71	CAA.Qc.Vrt	CAA.Jc	0.82		CAA.Qt.Vrt	0.62		CAA.Qt.Vrt	0.62
	VCA.MSsat	0.65		CAA.Qc	0.77		BSA.BsL	0.87		BSA.BsL	0.87
	VCA.ML.Hrz	1.00		CAA.Scpl	0.75		BSA.BCVL	1.00		BSA.BCVL	1.00
	VCA.sL.Hrz	0.67		CAA.Qcpl	0.68		BSA.BsSL	0.82		BSA.BsSL	0.82
	VCA.sL.Vrt	0.67		CAA.Jc.Hrz	0.82		BSA.BCVSL	0.99		BSA.BCVSL	0.99
	VCA.MDmax.Hrz	0.71		CAA.Jc.Vrt	0.82		BSA.BsL.Hrz	0.87		BSA.BsL.Hrz	0.87
	VCA.MDmax.Vrt	0.71		CAA.Qc.Hrz	0.97		BSA.BsL.Vrt	0.87		BSA.BsL.Vrt	0.87
	VCA.MSsat.Hrz	0.65		CAA.Scpl.Hrz	0.75		BSA.BCVL.Vrt	0.99		BSA.BCVL.Vrt	0.99
	VCA.MSsat.Vrt	0.65		CAA.Scpl.Vrt	0.76		BSA.BsSL.Hrz	0.82		BSA.BsSL.Hrz	0.82
				CAA.Qcpl.Hrz	0.69		BSA.BCVSL.Hrz	1.00		BSA.BCVSL.Hrz	1.00
				CAA.Qcpl.Vrt	0.71		BSA.BsSL.Vrt	0.82		BSA.BsSL.Vrt	0.82
			CAA.Qt	CAA.Qcpl	0.86		BSA.BCVSL.Vrt	0.98		BSA.BCVSL.Vrt	0.98
				CAA.Qt.Hrz	0.85	Col.kurtosis	Col.CoV	0.67	BSA.BCVSsat.Hrz	BSA.BCVDmax	0.80
				CAA.Qt.Vrt	0.84		Col.skew	0.93		BSA.BCVSsat	1.00

VCA.MSL.Vrt

Group	Feature	Value
	CAA.Qcpl.Hrz	0.84
	CAA.Qcpl.Vrt	0.83
VCA.MSL.Vrt	VCA.MSL	1.00
	VCA.sSL	0.83
	VCA.MSL.Hrz	1.00
	VCA.sSL.Hrz	0.83
	VCA.sSL.Vrt	0.83

Group	Feature	Value
	Col.CoV.hrz	0.70
	Col.skew.hrz	0.94
	Col.kurtosis.hrz	0.99
	Col.CoV.vrt	0.69
	Col.skew.vrt	0.93
	Col.kurtosis.vrt	0.99
	BSA.BCVS	0.67
	BSA.BCVS.Hrz	0.67
	BSA.BCVS.Vrt	0.66
Lum.CoV	Lum.CoV.hrz	0.93
	Lum.CoV.vrt	0.93
Lum.kurtosis.vrt	Lum.skew	0.84
	Lum.kurtosis	0.98
	Lum.skew.hrz	0.71
	Lum.kurtosis.hrz	0.85
	Lum.skew.vrt	0.92
VCA.CVS.Hrz	VCA.sS	0.62
	VCA.CVS	1.00
	VCA.sS.Hrz	0.62
	VCA.sS.Vrt	0.62
	VCA.CVS.Vrt	1.00
	BSA.BCVS	0.73
	BSA.BCVS.Hrz	0.73
	BSA.BCVS.Vrt	0.73
VCA.MSL.Vrt	VCA.MSL	1.00
	VCA.sSL	0.83
	VCA.MSL.Hrz	1.00
	VCA.sSL.Hrz	0.83
	VCA.sSL.Vrt	0.83

Group	Feature	Value
	BSA.BCVDmax.Hrz	0.80
	BSA.BCVDmax.Vrt	0.79
	BSA.BCVSsat.Vrt	1.00
Lum.mean.vrt	Lum.mean	0.99
	Lum.sd	0.94
	Lum.mean.hrz	0.96
	Lum.sd.hrz	0.90
	Lum.sd.vrt	0.97
	CAA.C	0.68
	CAA.C.Hrz	0.67
	CAA.C.Vrt	0.69
VCA.ML.Vrt	VCA.ML	1.00
	VCA.sL	0.67
	VCA.MDmax	0.71
	VCA.MSsat	0.65
	VCA.ML.Hrz	1.00
	VCA.sL.Hrz	0.67
	VCA.sL.Vrt	0.67
	VCA.MDmax.Hrz	0.71
	VCA.MDmax.Vrt	0.71
	VCA.MSsat.Hrz	0.65
	VCA.MSsat.Vrt	0.65

Table S3

Pairwise contrasts for the model investigating latent factor 1 expressed as the median differences between groups with different strengths of chemical defences (see 'Methods' for details). The effect size of pairwise differences increases with increasing deviation of such differences from zero, and the robustness of the result increases with decreasing degree of overlap of the 95% Credible Intervals (CIs) with zero.

Factor 1: Contrasts	Difference	Lower CI	Upper CI
Undefended at 2cm - Toxic and moderately unpalatable at 2cm	-1.53	-2.28	-0.78
Undefended at 2cm - Toxic and highly unpalatable at 2cm	-1.44	-2.18	-0.73
Undefended at 2cm - Undefended at 30cm	-0.18	-0.61	0.27
Undefended at 2cm - Toxic and moderately unpalatable at 30cm	-1.23	-1.99	-0.46
Undefended at 2cm - Toxic and highly unpalatable at 30cm	-1.35	-2.10	-0.65
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 2cm	0.08	-0.38	0.54
Toxic and moderately unpalatable at 2cm - Undefended at 30cm	1.35	0.58	2.15
Toxic and moderately unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	0.30	-0.02	0.63
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 30cm	0.17	-0.28	0.65
Toxic and highly unpalatable at 2cm - Undefended at 30cm	1.26	0.47	2.04
Toxic and highly unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	0.22	-0.29	0.70
Toxic and highly unpalatable at 2cm - Toxic and highly unpalatable at 30cm	0.09	-0.19	0.38
Undefended at 30cm - Toxic and moderately unpalatable at 30cm	-1.05	-1.83	-0.23
Undefended at 30cm - Toxic and highly unpalatable at 30cm	-1.17	-1.92	-0.37
Toxic and moderately unpalatable at 30cm - Toxic and highly unpalatable at 30cm	-0.13	-0.63	0.36

Table S4

Pairwise variance contrasts for the model investigating latent factor 1 expressed as the median differences of the residual standard deviation on the original scale (back-transformed from the log scale) between groups with different strengths of chemical defences (see 'Methods' for details). The effect size of pairwise differences increases with increasing deviation of such differences from zero, and the robustness of the result increases with decreasing degree of overlap of the 95% Credible Intervals (CIs) with zero.

Factor 1: variance contrasts	Diference	Lower CI	Upper CI
Undefended at 2cm - Toxic and moderately unpalatable at 2cm	0.21	-0.07	0.55
Undefended at 2cm - Toxic and highly unpalatable at 2cm	-0.03	-0.30	0.31
Undefended at 2cm - Undefended at 30cm	-0.12	-0.53	0.22
Undefended at 2cm - Toxic and moderately unpalatable at 30cm	-0.21	-0.67	0.12
Undefended at 2cm - Toxic and highly unpalatable at 30cm	0.13	-0.09	0.33
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 2cm	-0.23	-0.49	0.01
Toxic and moderately unpalatable at 2cm - Undefended at 30cm	-0.33	-0.76	0.01
Toxic and moderately unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	-0.42	-1.09	0.08
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 30cm	-0.08	-0.45	0.22
Toxic and highly unpalatable at 2cm - Undefended at 30cm	-0.09	-0.52	0.25
Toxic and highly unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	-0.19	-0.81	0.26
Toxic and highly unpalatable at 2cm - Toxic and highly unpalatable at 30cm	0.16	-0.26	0.51
Undefended at 30cm - Toxic and moderately unpalatable at 30cm	-0.09	-0.71	0.44
Undefended at 30cm - Toxic and highly unpalatable at 30cm	0.25	-0.15	0.69
Toxic and moderately unpalatable at 30cm - Toxic and highly unpalatable at 30cm	0.34	-0.06	0.85

Table S5

Pairwise contrasts for the model investigating latent factor 2 expressed as the median differences between groups with different strengths of chemical defences (see 'Methods' for details). The effect size of pairwise differences increases with increasing deviation of such differences from zero, and the robustness of the result increases with decreasing degree of overlap of the 95% Credible Intervals (CIs) with zero.

Factor 2: Contrasts	Difference	Lower CI	Upper CI
Undefended at 2cm - Toxic and moderately unpalatable at 2cm	-0.1226	-0.614	0.402
Undefended at 2cm - Toxic and highly unpalatable at 2cm	-0.1056	-0.617	0.394
Undefended at 2cm - Undefended at 30cm	1.0072	0.568	1.482
Undefended at 2cm - Toxic and moderately unpalatable at 30cm	0.8871	0.352	1.417
Undefended at 2cm - Toxic and highly unpalatable at 30cm	0.6930	0.125	1.194
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 2cm	0.0175	-0.333	0.347
Toxic and moderately unpalatable at 2cm - Undefended at 30cm	1.1337	0.552	1.705
Toxic and moderately unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	1.0052	0.668	1.312
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 30cm	0.8158	0.410	1.164
Toxic and highly unpalatable at 2cm - Undefended at 30cm	1.1151	0.552	1.686
Toxic and highly unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	0.9896	0.615	1.363
Toxic and highly unpalatable at 2cm - Toxic and highly unpalatable at 30cm	0.7997	0.473	1.105
Undefended at 30cm - Toxic and moderately unpalatable at 30cm	-0.1239	-0.709	0.462
Undefended at 30cm - Toxic and highly unpalatable at 30cm	-0.3172	-0.971	0.230
Toxic and moderately unpalatable at 30cm - Toxic and highly unpalatable at 30cm	-0.1898	-0.615	0.215

Table S6

Pairwise variance contrasts for the model investigating latent factor 2 expressed as the median differences of the residual standard deviation on the original scale (back-transformed from the log scale) between groups with different strengths of chemical defences (see 'Methods' for details). The effect size of pairwise differences increases with increasing deviation of such differences from zero, and the robustness of the result increases with decreasing degree of overlap of the 95% Credible Intervals (CIs) with zero.

Factor 2: Variance Contrast	Difference	Lower CI	Upper CI
Undefended at 2cm - Toxic and moderately unpalatable at 2cm	0.16	-0.08	0.45
Undefended at 2cm - Toxic and highly unpalatable at 2cm	0.06	-0.19	0.33
Undefended at 2cm - Undefended at 30cm	-0.12	-0.48	0.21
Undefended at 2cm - Toxic and moderately unpalatable at 30cm	0.01	-0.31	0.26
Undefended at 2cm - Toxic and highly unpalatable at 30cm	-0.13	-0.43	0.11
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 2cm	-0.11	-0.32	0.11
Toxic and moderately unpalatable at 2cm - Undefended at 30cm	-0.28	-0.67	0.04
Toxic and moderately unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	-0.15	-0.65	0.24
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 30cm	-0.29	-0.73	0.05
Toxic and highly unpalatable at 2cm - Undefended at 30cm	-0.18	-0.55	0.14
Toxic and highly unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	-0.05	-0.47	0.28
Toxic and highly unpalatable at 2cm - Toxic and highly unpalatable at 30cm	-0.18	-0.67	0.21
Undefended at 30cm - Toxic and moderately unpalatable at 30cm	0.13	-0.34	0.55
Undefended at 30cm - Toxic and highly unpalatable at 30cm	0.00	-0.47	0.43
Toxic and moderately unpalatable at 30cm - Toxic and highly unpalatable at 30cm	-0.13	-0.53	0.27

Table S7

Pairwise contrasts for the model investigating latent factor 3 expressed as the median differences between groups with different strengths of chemical defences (see 'Methods' for details). The effect size of pairwise differences increases with increasing deviation of such differences from zero, and the robustness of the result increases with decreasing degree of overlap of the 95% Credible Intervals (CIs) with zero.

Factor 3: Contrasts	Difference	Lower CI	Upper CI
Undefended at 2cm - Toxic and moderately unpalatable at 2cm	0.23	-0.27	0.69
Undefended at 2cm - Toxic and highly unpalatable at 2cm	0.18	-0.26	0.66
Undefended at 2cm - Undefended at 30cm	0.50	0.07	0.91
Undefended at 2cm - Toxic and moderately unpalatable at 30cm	0.50	0.00	1.01
Undefended at 2cm - Toxic and highly unpalatable at 30cm	0.51	0.02	0.97
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 2cm	-0.05	-0.37	0.24
Toxic and moderately unpalatable at 2cm - Undefended at 30cm	0.26	-0.31	0.82
Toxic and moderately unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	0.27	-0.03	0.56
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 30cm	0.28	-0.08	0.60
Toxic and highly unpalatable at 2cm - Undefended at 30cm	0.32	-0.24	0.84
Toxic and highly unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	0.32	-0.02	0.69
Toxic and highly unpalatable at 2cm - Toxic and highly unpalatable at 30cm	0.33	0.07	0.56
Undefended at 30cm - Toxic and moderately unpalatable at 30cm	0.01	-0.59	0.60
Undefended at 30cm - Toxic and highly unpalatable at 30cm	0.01	-0.53	0.60
Toxic and moderately unpalatable at 30cm - Toxic and highly unpalatable at 30cm	0.01	-0.35	0.40

Table S8

Pairwise variance contrasts for the model investigating latent factor 3 expressed as the median differences of the residual standard deviation on the original scale (back-transformed from the log scale) between groups with different strengths of chemical defences (see 'Methods' for details). The effect size of pairwise differences increases with increasing deviation of such differences from zero, and the robustness of the result increases with decreasing degree of overlap of the 95% Credible Intervals (CIs) with zero.

Factor 3: Variance Contrast	Estimate	Lower CI	Upper CI
Undefended at 2cm - Toxic and moderately unpalatable at 2cm	0.15	-0.10	0.44
Undefended at 2cm - Toxic and highly unpalatable at 2cm	0.09	-0.15	0.38
Undefended at 2cm - Undefended at 30cm	-0.19	-0.62	0.15
Undefended at 2cm - Toxic and moderately unpalatable at 30cm	-0.12	-0.46	0.14
Undefended at 2cm - Toxic and highly unpalatable at 30cm	0.00	-0.24	0.19
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 2cm	-0.06	-0.26	0.14
Toxic and moderately unpalatable at 2cm - Undefended at 30cm	-0.34	-0.78	-0.01
Toxic and moderately unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	-0.27	-0.78	0.15
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 30cm	-0.15	-0.55	0.15
Toxic and highly unpalatable at 2cm - Undefended at 30cm	-0.28	-0.72	0.05
Toxic and highly unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	-0.21	-0.67	0.15
Toxic and highly unpalatable at 2cm - Toxic and highly unpalatable at 30cm	-0.09	-0.53	0.25
Undefended at 30cm - Toxic and moderately unpalatable at 30cm	0.08	-0.43	0.58
Undefended at 30cm - Toxic and highly unpalatable at 30cm	0.19	-0.23	0.66
Toxic and moderately unpalatable at 30cm - Toxic and highly unpalatable at 30cm	0.11	-0.23	0.51

Table S9

Pairwise contrasts for the model investigating latent factor 4 expressed as the median differences between groups with different strengths of chemical defences (see 'Methods' for details). The effect size of pairwise differences increases with increasing deviation of such differences from zero, and the robustness of the result increases with decreasing degree of overlap of the 95% Credible Intervals (CIs) with zero.

Factor 4: Contrasts	Estimate	Lower CI	Upper CI
Undefended at 2cm - Toxic and moderately unpalatable at 2cm	0.21	-0.51	0.96
Undefended at 2cm - Toxic and highly unpalatable at 2cm	0.31	-0.38	1.04
Undefended at 2cm - Undefended at 30cm	0.21	-0.25	0.69
Undefended at 2cm - Toxic and moderately unpalatable at 30cm	0.75	0.02	1.48
Undefended at 2cm - Toxic and highly unpalatable at 30cm	0.73	0.05	1.51
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 2cm	0.09	-0.31	0.53
Toxic and moderately unpalatable at 2cm - Undefended at 30cm	-0.01	-0.81	0.78
Toxic and moderately unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	0.53	0.25	0.81
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 30cm	0.52	0.08	0.98
Toxic and highly unpalatable at 2cm - Undefended at 30cm	-0.10	-0.92	0.65
Toxic and highly unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	0.44	0.02	0.84
Toxic and highly unpalatable at 2cm - Toxic and highly unpalatable at 30cm	0.42	0.16	0.70
Undefended at 30cm - Toxic and moderately unpalatable at 30cm	0.54	-0.26	1.35
Undefended at 30cm - Toxic and highly unpalatable at 30cm	0.53	-0.27	1.32
Toxic and moderately unpalatable at 30cm - Toxic and highly unpalatable at 30cm	-0.02	-0.42	0.44

Table S10

Pairwise variance contrasts for the model investigating latent factor 4 expressed as the median differences of the residual standard deviation on the original scale (back-transformed from the log scale) between groups with different strengths of chemical defences (see 'Methods' for details). The effect size of pairwise differences increases with increasing deviation of such differences from zero, and the robustness of the result increases with decreasing degree of overlap of the 95% Credible Intervals (CIs) with zero.

Factor 4: Variance Contrasts	Estimate	Lower CI	Upper CI
Undefended at 2cm - Toxic and moderately unpalatable at 2cm	0.08	-0.19	0.40
Undefended at 2cm - Toxic and highly unpalatable at 2cm	0.07	-0.19	0.39
Undefended at 2cm - Undefended at 30cm	-0.24	-0.66	0.11
Undefended at 2cm - Toxic and moderately unpalatable at 30cm	0.21	0.00	0.40
Undefended at 2cm - Toxic and highly unpalatable at 30cm	-0.05	-0.31	0.14
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 2cm	-0.01	-0.24	0.23
Toxic and moderately unpalatable at 2cm - Undefended at 30cm	-0.32	-0.80	0.04
Toxic and moderately unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	0.13	-0.27	0.46
Toxic and moderately unpalatable at 2cm - Toxic and highly unpalatable at 30cm	-0.13	-0.56	0.21
Toxic and highly unpalatable at 2cm - Undefended at 30cm	-0.31	-0.78	0.04
Toxic and highly unpalatable at 2cm - Toxic and moderately unpalatable at 30cm	0.14	-0.20	0.40
Toxic and highly unpalatable at 2cm - Toxic and highly unpalatable at 30cm	-0.12	-0.60	0.25
Undefended at 30cm - Toxic and moderately unpalatable at 30cm	0.45	0.07	0.90
Undefended at 30cm - Toxic and highly unpalatable at 30cm	0.19	-0.25	0.65
Toxic and moderately unpalatable at 30cm - Toxic and highly unpalatable at 30cm	-0.26	-0.59	0.03

Table S11

Coefficient estimates of the model investigating the scores for latent factor 1 between species of nudibranchs with different levels of chemical defences (N = 12, R^2 = 0.36). Estimates are based on a Student distribution with an identity link for the mean of the response distribution and a log link for its residual standard deviation (Sigma). The estimate is more likely to be non-zero when the credible intervals do not overlap with zero.

Coefficient	Mean	M. Error	95% CIs Low	High
Group-level effects				
Phylogenesis				
Sd [Intercept]	0.24	0.18	0.01	0.66
Species of nudibranch				
Sd [Intercept: Distance 2 cm]	0.21	0.12	0.02	0.48
Sd [Distance 30 cm]	0.12	0.10	0.00	0.36
Sigma: Sd [Intercept: Distance 2 cm]	0.12	0.09	0.00	0.35
Sigma: Sd [Distance 30 cm]	0.14	0.11	0.01	0.41
Population-level effects				
Intercept [Undefended, 2 cm]	-1.08	0.28	-1.63	-0.51
Sigma_Intercept [Undefended, 2 cm]	-0.24	0.16	-0.53	0.08
Chemical defence [Toxic and moderately unpalatable]	1.52	0.37	0.73	2.24
Chemical defence [Toxic and highly unpalatable]	1.44	0.36	0.73	2.18
Distance [30 cm]	0.18	0.22	-0.25	0.63
Chemical defence [Toxic and moderately unpalatable] x Distance [30 cm]	-0.48	0.28	-1.02	0.07
Chemical defence [Toxic and highly unpalatable] x Distance [30 cm]	-0.27	0.27	-0.80	0.25
Sigma: Chemical defence [Toxic and moderately unpalatable]	-0.30	0.21	-0.76	0.1
Sigma: Chemical defence [Toxic and highly unpalatable]	0.04	0.19	-0.36	0.38
Sigma: Distance [30cm]	0.13	0.21	-0.27	0.55
Sigma: Chemical defence [Toxic and moderately unpalatable] x Distance [30 cm]	0.09	0.28	-0.47	0.65
Sigma: Chemical defence [Toxic and highly unpalatable] x Distance [30 cm]	-0.32	0.26	-0.85	0.18

Table S12

Coefficient estimates of the model investigating the scores for latent factor 2 between species of nudibranchs with different levels of chemical defences (N = 12, R^2 = 0.28). Estimates are based on a Student distribution with an identity link for the mean of the response distribution and a log link for its residual standard deviation (Sigma). The estimate is more likely to be non-zero when the credible intervals do not overlap with zero.

Coefficient	Mean	M. Error	95% CIs	
			Low	High
Group-level effects				
Phylogenesis				
Sd [Intercept]	0.13	0.11	0	0.42
Species of nudibranch				
Sd [Intercept: Distance 2 cm]	0.14	0.09	0.01	0.35
Sd [Distance 30 cm]	0.16	0.12	0.01	0.44
Sigma: Sd [Intercept: Distance 2 cm]	0.09	0.08	0.00	0.28
Sigma: Sd [Distance 30 cm]	0.12	0.10	0.00	0.38
Population-level effects				
Intercept [Undefended, 2 cm]	0.31	0.20	-0.09	0.7
Sigma_Intercept [Undefended, 2 cm]	-0.29	0.14	-0.57	0
Chemical defence [Toxic and moderately unpalatable]	0.12	0.26	-0.39	0.63
Chemical defence [Toxic and highly unpalatable]	0.11	0.26	-0.38	0.63
Distance [30 cm]	-1.01	0.23	-1.48	-0.6
Chemical defence [Toxic and moderately unpalatable] x Distance [30 cm]	0.00	0.28	-0.56	0.56
Chemical defence [Toxic and highly unpalatable] x Distance [30 cm]	0.21	0.28	-0.33	0.78
Sigma: Chemical defence [Toxic and moderately unpalatable]	-0.24	0.19	-0.63	0.13
Sigma: Chemical defence [Toxic and highly unpalatable]	-0.07	0.17	-0.42	0.26
Sigma: Distance [30cm]	0.14	0.21	-0.28	0.54
Sigma: Chemical defence [Toxic and moderately unpalatable] x Distance [30 cm]	-0.17	0.27	-0.70	0.38
Sigma: Chemical defence [Toxic and highly unpalatable] x Distance [30 cm]	0.00	0.25	-0.48	0.5

Table S13

Coefficient estimates of the model investigating the scores for latent factor 3 between species of nudibranchs with different levels of chemical defences (N = 12, R^2 = 0.07). Estimates are based on a Student distribution with an identity link for the mean of the response distribution and a log link for its residual standard deviation (Sigma). The estimate is more likely to be non-zero when the credible intervals do not overlap with zero.

Coefficient	Mean	M. Error	95% CIs	
			Low	High
Group-level effects				
Phylogenesis				
Sd [Intercept]	0.11	0.11	0	0.39
Species of nudibranch				
Sd [Intercept: Distance 2 cm]	0.13	0.08	0.01	0.32
Sd [Distance 30 cm]	0.12	0.10	0.00	0.36
Sigma: Sd [Intercept: Distance 2 cm]	0.13	0.09	0.01	0.35
Sigma: Sd [Distance 30 cm]	0.12	0.10	0.00	0.36
Population-level effects				
Intercept [Undefended, 2 cm]	0.22	0.18	-0.14	0.58
Sigma_Intercept [Undefended, 2 cm]	-0.49	0.19	-0.87	-0.1
Chemical defence [Toxic and moderately unpalatable]	-0.23	0.24	-0.69	0.27
Chemical defence [Toxic and highly unpalatable]	-0.18	0.23	-0.63	0.3
Distance [30 cm]	-0.49	0.22	-0.91	-0.07
Chemical defence [Toxic and moderately unpalatable] x Distance [30 cm]	0.23	0.27	-0.30	0.75
Chemical defence [Toxic and highly unpalatable] x Distance [30 cm]	0.17	0.25	-0.32	0.66
Sigma: Chemical defence [Toxic and moderately unpalatable]	-0.27	0.24	-0.74	0.19
Sigma: Chemical defence [Toxic and highly unpalatable]	-0.14	0.22	-0.60	0.29
Sigma: Distance [30cm]	0.26	0.25	-0.22	0.75
Sigma: Chemical defence [Toxic and moderately unpalatable] x Distance [30 cm]	-0.11	0.32	-0.75	0.51
Sigma: Chemical defence [Toxic and highly unpalatable] x Distance [30 cm]	-0.27	0.30	-0.87	0.32

Table S14

Coefficient estimates of the model investigating the scores for latent factor 4 between species of nudibranchs with different levels of chemical defences (N = 12, R^2 = 0.15). Estimates are based on a Student distribution with an identity link for the mean of the response distribution and a log link for its residual standard deviation (Sigma). The estimate is more likely to be non-zero when the credible intervals do not overlap with zero.

Coefficient	Mean	M. Error	95% CIs Low	High
Group-level effects				
Phylogenesis				
Sd [Intercept]	0.23	0.18	0.01	0.66
Species of nudibranch				
Sd [Intercept: Distance 2 cm]	0.19	0.11	0.02	0.44
Sd [Distance 30 cm]	0.11	0.09	0.00	0.33
Sigma: Sd [Intercept: Distance 2 cm]	0.12	0.09	0.01	0.33
Sigma: Sd [Distance 30 cm]	0.10	0.08	0.00	0.3
Population-level effects				
Intercept [Undefended, 2 cm]	0.43	0.27	-0.09	0.98
Sigma_Intercept [Undefended, 2 cm]	-0.31	0.17	-0.64	0.04
Chemical defence [Toxic and moderately unpalatable]	-0.22	0.36	-0.99	0.49
Chemical defence [Toxic and highly unpalatable]	-0.32	0.36	-1.07	0.36
Distance [30 cm]	-0.21	0.24	-0.67	0.27
Chemical defence [Toxic and moderately unpalatable] x Distance [30 cm]	-0.33	0.27	-0.87	0.21
Chemical defence [Toxic and highly unpalatable] x Distance [30 cm]	-0.22	0.27	-0.75	0.31
Sigma: Chemical defence [Toxic and moderately unpalatable]	-0.11	0.21	-0.54	0.29
Sigma: Chemical defence [Toxic and highly unpalatable]	-0.09	0.20	-0.50	0.29
Sigma: Distance [30cm]	0.28	0.21	-0.13	0.69
Sigma: Chemical defence [Toxic and moderately unpalatable] x Distance [30 cm]	-0.62	0.27	-1.15	-0.09
Sigma: Chemical defence [Toxic and highly unpalatable] x Distance [30 cm]	-0.22	0.25	-0.72	0.27