Original article

DNA metabarcoding reveals the dietary profiles of a benthic marine crustacean, *Nephrops norvegicus*

Peter Shum[1,2], Janine Wäge-Recchioni[,2,3], Graham S Sellers[2], Magnus L Johnson[4], Domino A. Joyce[2]

[1] *Faculty of Science, Liverpool John Moores University, Liverpool, L3 3AF, UK*

[2] *School of Natural Sciences, University of Hull, Hull HU6 7RX, UK*

[3] *Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Seestrasse 15, 18119, Rostock, Germany*

[4] *School of Environmental Sciences, University of Hull, Hull HU6 7RX, UK*

Keywords: COI; DNA metabarcoding; gut contents; natural sampler; scampi

Corresponding Author: P.Shum[at]ljmu.ac.uk

Abstract

Norwegian lobster, *Nephrops norvegicus*, are a generalist scavenger and predator capable of short foraging excursions but can also suspension feed. Existing knowledge about their diet relies on a combination of methods including morphology-based stomach content analysis and stable isotopes, which are time and experience costly and often lack the resolution to distinguish prey items to species level. DNA metabarcoding overcomes many of the challenges associated with traditional methods and it is an attractive approach to study the dietary profiles of animals. Here, we present the diet of the commercially valuable *Nephrops norvegicus* using DNA metabarcoding of gut contents. Despite difficulties associated with host amplification, our cytochrome oxidase I (COI) molecular assay successfully achieves higher resolution information than previously used traditional approaches. We detected taxa that were likely consumed during different feeding strategies. Dinoflagellata, Chlorophyta and Bacillariophyta accounted for almost 50% of the prey items consumed, and are associated with suspension feeding, while fish with high discard rates were detected which are linked to active foraging. In addition, we were able to characterise biodiversity patterns by considering *Nephrops* as natural samplers, as well as detecting parasitic dinoflagellates (e.g., *Hematodinium* sp.) in over 50% of the samples, which are known to influence burrow related behaviour in infected individuals. The metabarcoding data presented here greatly enhances a better understanding of a species' ecological role that can be applied as a routine procedure in future studies for proper consideration in the management and decision-making of fisheries.

Introduction

Marine animals possess a diverse repertoire of feeding strategies linked to individual and species-specific foraging behaviour in complex, multifood environments. They engage in food acquisition at individual, population and community levels that shape ecosystem functioning across trophic levels. This influences biotic interactions and behaviour among species which drives phenotypic selection and eco-evolutionary feedbacks (Post and Palkovacs, 2009; Clinchy et al., 2013; Majdi et al., 2018). At an individual level, feeding is linked to nutrition and ecophysiology as the quantity and quality of food resources regulate individual survival, growth and fecundity. The ability to monitor these dietary profiles of animals, particularly those of commercial importance, across spatial scales is fundamental to understanding how environmental and anthropogenic activities influence nutrition. An accurate and comprehensive dietary matrix of species in marine ecosystems is an essential foundation for ecological fisheries models that are necessary for developing a better understanding of likely impacts of climate change on marine systems (Mackinson and Daskalov 2007).

In the marine environment, species are rarely observed to forage directly and most studies depend on the identification of prey remains in stomach contents or faeces to determine the prey items being consumed (Pompanon et al., 2012). The methodology for studying the diet of animals is varied and considers the type of sample collection, sample processing, and the identification and quantification of prey consumption. However, obtaining detailed diet information is challenging for many species because of the demanding effort required to directly observe and physically identify food items from stomach contents (Pompanon et al., 2012). This is particularly problematic in understudied species as traditional stomach content analyses rely on extensive experience to identify species specific characteristics of hard parts (e.g., otoliths, scales, cleithra, carapace) and soft contents with the aid of good references to identify items from digested, broken and finely comminuted material (Zacharia, 2017). But diet items that have been recently consumed can be rapidly digested and become quickly indiscernible which can underestimate dietary composition. This lack of consistent information can negatively impact food web models in estimating annual catch and consumption by predators (Aydin et al., 2007) and could lead to unexpected and undesirable management outcomes (Gaichas et al., 2010). DNA barcoding using the universal cytochrome oxidase I (COI, ~650bp) gene can improve detection of otherwise unidentifiable individual prey items, however there is reduced success for highly digested material

with time and resources needed to process separate items (Moran et al., 2016). Another popular

application is the use of carbon and nitrogen isotopic signatures (i.e., stable isotopes) in tissues to

measure the differential assimilation of dietary components which can overcome some of the

shortcomings of traditional dietary studies (Dalerum & Angerbjörn, 2005). For example,

Wieczorek et al. (2018) studied the diet of the lesser spotted dogfish using the stomach content

analysis and stable isotopes. They found that stomach content analysis presents a snapshot of the

diet that overestimate hard-bodied prey species while stable isotopes revealed that soft-bodied

filter feeders were by far the most important diet items, accounting for approximately 76% of the

energy assimilated. This highlights the advantage of complementary approaches to provide a better

overview of a species trophic position.

While stable isotope analysis is a powerful method to complement stomach content data in

estimating dietary profiles, it lacks the resolution to accurately recover species level information

due the deficiency in suitable isotopic reference data (Elsner et al., 2012; Galloway et al., 2015).

An alternative method that has quickly gained momentum in trophic ecology studies is DNA

metabarcoding. DNA metabarcoding embraces DNA barcoding of short DNA fragments and next

generation sequencing to identify species information from a variety of sample types, and

facilitates the detection of small, soft-bodied or cryptic species which might be overlooked during

traditional diet analysis (de Sousa et al., 2019). The power and utility of DNA metabarcoding has

become an attractive tool to identify food DNA consumed by animals to reveal dietary habits (van

Zinnicq Bergmann et al., 2021), parasite load (Cabodevilla et al., 2022), trophic niches (Meyer et

al., 2020), and local natural biodiversity (Siegenthaler et al., 2019).

Nephrops norvegicus, (also referred to as the Norwegian lobster, Dublin Bay Prawn or

scampi, *Nephrops* hereafter), is a benthic decapod crustacean and is one of the most important

economically valued fisheries in Europe generating a value of 50 M€, making it the second most

valuable landed species in the North Sea and Eastern Arctic region in 2019 (STECF, 2021). They

are distributed over semi-isolated mud patches throughout the North-Eastern Atlantic Ocean in the

North Sea as far south as the Canary Islands and extending into the Eastern Mediterranean Sea

(Johnson et al., 2013). *Nephrops* are thought to be crepuscular opportunistic predators and

scavengers, found to feed on fish, crustaceans, molluscs and other taxa (Cristo & Cartes, 1998).

Nephrops also possess a complex mode of feeding by capturing and ingesting suspended

particulate organic matter (i.e., POM_{susp}) from the water column, i.e., suspension feeding - a

strategy thought to be used especially by females for surviving starvation during the long breeding
period when they are restricted to their burrows (Loo et al., 1993). Santana et al. (2020) revealed
the importance of suspension feeding through stable isotopes and found that half of their diet was
made up of suspended particulate organic matter alone. Fish were shown to be another important
food source, but it is unclear which species contribute to their diet or whether their diet is
supplemented by discards arising from inshore fisheries (Santana et al., 2020). The link between
diet and behaviour in *Nephrops* is particularly important to fisheries as when they are in their
burrows they cannot be captured by fishers. The fishing fleet are aware that *Nephrops*, as well as
having a crepuscular emergence habit, will appear and disappear en masse in different grounds at
different times of year and fishers will move among the discrete fishing grounds (functional units)
as they become available (Ungfors et al., 2013)

Here we applied DNA metabarcoding to characterise the diet of *Nephrops norvegicus* from
specimens collected in the North and Irish Seas. *Nephrops* are a generalist forager and highly
commercial benthic crustacean and combined with a molecular approach offers a unique
opportunity to consider them as natural biodiversity samplers. Therefore, we aim to i) characterise
their feeding strategy using DNA metabarcoding of digested material in the gut and ii) examine
the biodiversity of prey consumed.

120
Materials and methods
Sample collection and processing
A total of 207 *Nephrops norvegicus* specimens were collected on board commercial fishing vessels
in the East (n=77) and West (n=68) of the North Sea - Fladen Grounds and the Irish Sea (n=63) in
January 2016. Specimens were collected from the seafloor and were stored on ice at sea and stored
at -80°C in the lab prior to dissection. In the lab, each specimen was thawed on ice after which the
total contents of the gastro-intestinal tract was placed into a DNeasy PowerSoil tube using sterile
forceps and DNA extraction was performed following the manufacturer's protocol. Purified DNA
extracts were quantified using dsDNA HS Assay kit Qubit fluorometer
(www.lifetechnologies.com).

131
Data generation, library preparation and sequencing
Each sample was PCR amplified targeting the 313bp fragment of the cytochrome oxidase I gene
(COI) (mICOIintF: GGWACWGGWTGAACWGTWTAYCCYCC; Leray *et al.*, (2013), matched

to jgHCO2198: TAIACYTCIGGRTGICCRAARAAYCA; Geller *et al*., 2013). Each sample was

amplified in triplicate and subsequently pooled to reduce biases in individual PCRs. A single step

PCR protocol was used containing indexed primers with 8 bp oligo tags differing in at least 3

bases. A variable number (2, 3 or 4) of degenerate bases (N's) were added to the beginning of each

primer to increase nucleotide diversity for sequencing. PCR reactions were carried out in 25 μL

volumes containing 12.5 μL of MyFi mix (Bioline), 1 μL of each forward and reverse primer (0.5

μM), 2 μL of DNA template (0.5-10 ng/μL) and 8.5 μL of molecular grade H_2O. The thermocycle

condition for the PCR was 10 min at 94ºC; 35 cycles at 94 ºC for 1 min, 45 ºC for 1 min and 72

ºC for 1 min; and a final elongation at 72 ºC for five minutes. The quality of all amplifications was

assessed through electrophoresis, running the PCR products on a 1% Sodium borate (1X SB) gel

stained with gel red (Biotium). All PCR products were purified using magnetic beads (0.8x, Omega

Mag-Bind) before all samples were pooled in equimolar amounts and normalized to 45 μL

containing 3 μg of total purified PCR product. Along with the samples, one positive (*Astatotilapia*

burtoni) and one negative (purified water) control was amplified in each plate and sequenced.

The Illumina library was constructed from 3 μg of total DNA using the NextFlex PCR-free

library preparation kit (BIOO Scientific) following the manufacturer's instructions. The library

was quantified by qPCR using NEBNext Library Quant Kit for Illumina, adjusted to a final

molarity of 15 pM and with a 10% PhiX control, was sequenced on an Illumina MiSeq platform

using v2 chemistry (2 x 250 bp paired-end) at the University of Hull.

Bioinformatic pipeline

The sequence reads were analysed using the OBITools software (Boyer *et al*., 2016). FastQC was

used to assess the quality of the reads and trimmed accordingly based on a minimum quality

threshold of 28 using *obicut*. Pair-end reads were aligned using *illuminapairend* and alignments

with a quality score <40 were discarded. The aligned dataset was demultiplexed using *ngsfilter*.

The aligned reads were further filtered for length 300-320 bp (*obigrep*) and reads containing

ambiguous bases were removed. The reads were then dereplicated using obiuniq and a chimera

removal step was performed using the uchime-denovo algorithm implemented in vsearch (Rognes

et al., 2016). Molecular Operational Taxonomic Unit (MOTU) clustering was carried out using

Swarm (d value of 13) (Mahé *et al*., 2014). A reference COI database was generated by in silico

PCR against the R134 release of the EMBL-EBI database using ecoPCR (Ficetola *et al*., 2010),

and taxonomic assignment for each MOTU was performed using the *ecotag* algorithm, which

implements a conservative lowest common ancestor approach.

Statistical analysis

To determine adequate sampling of the diet profiles for each *Nephrops* collection site, we

examined species diversity (presence-absence) using sample-based rarefaction and extrapolation

sampling curves (iNEXT, Hsieh *et al*., 2016). The rarefaction curves were extrapolated for each

collection site to 200 samples and the total species richness (S_{est}) for each site was estimated (Chao

1987). Rarefaction curves were used to determine the percentage of S_{est} sampled for each site by

dividing the cumulative number of expected species (S_{est}) by the estimated total species richness

of each site (total *S_{est}*). We calculated indices of the relative frequency of occurrence as the

occurrence per gut (O/G) index (the number of occurrences of a diet item divided by the total

number of gut samples). Alpha diversity was performed using a pairwise ANOVA of diversity

measure (Shannon index) of MOTUs for each site (*vegan*, Oksanen 2013). Analyses were

performed in the statistical programming environment R v.4.0.2 (R Core Team, 2020). Dietary

overlap per site was measured using Horn's similarity index implemented in the function *vegdist*

(*vegan*). >0.6 are considered very high overlap, and conversely, dietary overlap <0.4 indicates that

there are substantial differences in patterns of prey use (Ross 1986).

Network analysis

We examined the dietary differences between *Nephrops* collections by generating a quantitative

bipartite network (Barrat *et al*., 2004) implemented in R (*geomnet*, Tyner & Hofmann 2016),

where individuals grouped by location were linked to prey groups. The network was weighted to

visualise the proportional contribution of each prey group to the diet of *Nephrops* at a given

location. Furthermore, a unipartite network was generated to illustrate the dietary preference for

MOTUs at lower taxonomic ranks (species, genus or family level) between specimens among

locations. This analysis consisted of one set of nodes whereby two species can be connected

through trophic interactions (Delmas *et al*., 2019). The network was directed from predator to prey

and interactions were weighted using presence-absence abundance for each MOTU. The final

visualisation of the unipartite network was performed using the Force Atlas algorithm in Gephi

v0.9.2 (Bastian *et al*., 2009).

Results

Overall, a total number of 207 *Nephrops norvegicus* specimens collected from three locations were screened for gut contents using DNA metabarcoding from the North Sea (East (n=77) and West (n=68)) and Irish Sea (n=63), including one positive and one negative control. Illumina sequencing produced a total of 18,646,302 paired-end reads. After quality filtering (paired-end assembly, quality, length filtering and dereplication) and removing 21,592 potential chimeras (0.9%), the final table consisted of 815 MOTUs (12,228,986 reads). However, the vast majority of reads (98.6%, 12,054,773) aligned to Nephropidae which we considered host contamination and was therefore removed from further analysis. Furthermore, taxa unlikely to form the basis of *Nephrops* diet were removed (37 MOTUs, 5.2% reads, e.g., terrestrial species: hominid, Canidae, Insecta). With a modest number of remaining reads, we considered data with reliable taxonomic assignments 85% and greater, with each MOTU having a minimum of three reads. This resulted in a final dataset consisting of 94 *N. norvegicus* specimens (North Sea, East: 44, West: 25; Irish Sea: 25) with 116,154 reads in 119 MOTUs.

Table 1 Number of *N. norvegicus* specimens needed to detect various percentages of estimated total MOTU richness at three sites in the North Sea (East and West) and the Irish Sea. *n* is the total number of samples analysed and Obs is the total MOTU richness observed for each site.

Site	*n*	Obs	% Estimated total species richness			
			80	90	95	99
East	44	74%	62	101	130	158
West	25	49%	77	119	147	175
Irish	25	46%	84	131	163	194
Combined	94	80%	93	148	186	224

Rarefaction extrapolation curves were used to assess sampling effort of *Nephrops* diet among collections using presence-absence data **(Fig. 1)**. Rarefaction curves failed to reach saturation for each group suggesting increased sampling effort and/or sequencing depth is desirable to adequately obtain sufficient MOTU coverage. The analysis of total species richness revealed 74%, 49%, 46% of the estimated total MOTU richness was observed in diet composition from North

223 Sea East (*n*=44), West (*n*=25) and Irish Sea (*n*=25) respectively. Rarefaction curves indicated that

224 an average of 74 and 117 individuals per site were needed to detect 80% and 90% of the estimated

225 total MOTU species richness respectively **(Table 1)**.

227
Figure 1 Rarefaction and extrapolation curves for MOTU richness for diet sample estimates from three collections,
with symmetric 95% confidence intervals based on Colwell *et al*. (2012). Solid lines: rarefaction curves (observed
data). Dashed lines: extrapolation curves. Shaded area for each solid line: 95% confidence interval for the expected
rarefied MOTU richness. Shaded area for each dashed line: 95% confidence interval for the expected extrapolated
class richness up to a sample size of 200. Points indicate observed rarefaction values for each collection.

For alpha diversity analysis, we calculated the mean Shannon diversity and observed MOTUs

found in the diet of *N. norvegicus* in each site **(Fig. 2)**. A one-way ANOVA analysis comparing

diet diversity showed considerable differences across sites ($F = 5.17$, $p = 0.0067$). A Holm-

Bonferroni corrected posthoc *t*-test showed the North Sea East collection was notably different to

both the North Sea West and Irish Sea collections (paired *t*-test, $p < 0.001$). However, the observed

number of MOTUs was higher in the North Sea East (86) than the West (50) and Irish Sea (43).

Figure 2 Box plots showing alpha diversity of *N. norvegicus* gut contents for each site, North Sea East, West and Irish
Sea. The Shannon index was computed for all 94 specimens compared across locations. Bars above plots indicate
significance *** p<0.001, NS non-significant.
245

Overall diet composition

The 119 MOTUs identified from the gastro-intestinal tract across *Nephrops* specimens consisted

of 16 identified Phyla. *Nephrops* show high dietary overlap across sites (**Table S1; Fig. S1**) at

82.2%, whereas 10% of individuals show differences in diet, likely as a result of low sample

resolution. The bipartite plot demonstrates the broad associations of 22 prey groups between

collections (**Fig. 3**). The variety of prey sources was distributed over a diverse range of groups

with Dinoflagellata as the most frequently occurring taxa in the gut content (average occurrence

per gut, O/G = 0.44), followed by Chlorophyta (aO/G = 0.26), Holothuroidea (aO/G = 0.204),

Malacostraca (aO/G = 0.203), Asteroidea (aO/G = 0.17), Bacillariophyta (aO/G = 0.16) and

Nemertea (aO/G = 0.13) among others. While these groups appear abundant in the *Nephrops* diet,

Fungi was found to be the most diverse with 28 MOTUs followed by Actinopterygii (12 MOTUs),

Bacillariophyta (8 MOTUs), Malacostraca (5 MOTUs) and Ochrophyta (5 MOTUs) with the

remaining groups showing between 1-4 MOTUs. We found instances of some groups exclusively

reported for the North Sea East (e.g., Aves, Mammalia), West (Discosea, Florideophyceae,

Oomycota) and Irish Sea (Cephalopoda, Chondrichthyes).

261
262
Figure 3 Bipartite network including 94 *N. norvegicus* specimens (North Sea, East: 44, West: 25 and Irish Sea: 25)
and their prey items (grouped into 22 categories). The relative proportion of each prey category consumed by each *N.*
norvegicus group corresponds with the width of each interaction bar. The pie charts show the relative proportion of
Phyla within each group.

Prey composition

In evaluating dietary profiles of *Nephrops*, we considered MOTUs identified at lower taxonomic

ranks with >90% identity (58 MOTUs) to construct an empirical food web composed of 17 Fungi,

15 vertebrates, 15 invertebrates, 6 algae and 5 protists, totalling 61 nodes and 94 weighted edges

(connecting predatory interactions) based on presence-absence abundance (**Fig. 4**). The network

is directed with links originating from the predator node to the prey node. Overall, alga MOTUs

(aO/G = 0.18) dominated node occurrence followed by metazoan (aO/G = 0.04) and Fungi (aO/G

= 0.02). The metrics to determine key species within the network identified similar species with

the most central nodes identified through the weighted degree centrality metric showing protists

(Suessiales *sp*., O/G = 0.61; *Hematodinium sp.*, O/G = 0.60; Dinophyceae *sp*., O/G = 0.11) as the

top MOTUs followed by algae (*Micromonas bravo,* O/G = 0.27; *Chloroparvula pacifica*, O/G =

0.15) and invertebrates (*Asterias rubens*, O/G = 0.22; *Crossaster papposus* O/G = 0.10).

281
Figure 4 Unipartite network illustrating species composition digested for *N. norvegicus* specimens collected at three
sites. Venn diagram shows the number of species level MOTUs per site.

The North Sea East collection comprised the highest number of unique MOTUs (18) with
seven vertebrates, four invertebrates, two algae and five Fungi detected. Vertebrate MOTUs
composed mostly of fish species such as the common dragonet (*Callionymus lyra*, O/G = 0.02),
European plaice (*Pleuronectes platessa*, O/G = 0.04), Atlantic herring (*Clupea harengus*, O/G =
0.02) and the common dab (*Limanda limanda*, O/G = 0.18). Other vertebrates included a seabird
species, razorbill (*Alca torda*, O/G = 0.02), and two mammals, white-beaked dolphin
(*Lagenorhynchus albirostris*, O/G = 0.02) and harbour porpoise (*Phocoena phocoena*, O/G =
0.04). Four invertebrates detected were the black brittle star (*Ophiocomina nigra*, O/G = 0.02),
two hydroids (*Nemertesia antennina* O/G = 0.02, *Leuckartiara octona*, O/G = 0.11) and a
polychaete (*Pholoe pallida*, O/G = 0.04). By contrast, the North Sea West was composed of eight
unique MOTUs with two algae, two invertebrates, three Fungi and one amoeba detected. We
detected an invertebrate hydroid MOTU in the Family Campanariidae (O/G = 0.04) and the
European lobster (*Homarus gammarus*, O/G = 0.04). Alga identified were the brown Forkweed
alga (*Dictyota dichotoma*, O/G = 0.04) and red alga (*Ahnfeltia plicata*, O/G = 0.08). The Irish Sea
showed the least unique MOTUs with two vertebrates (northern pike, *Esox lucius*, O/G = 0.04;
small-spotted catshark, *Scyliorhinus canicula*, O/G = 0.04), one invertebrate (curled octopus,
Eledone cirrhosa, O/G = 0.04) and two Fungi detected.

Discussion
We used gut content metabarcoding of the commercially valuable *Nephrops norvegicus* to provide
a detailed detection of specific food items to enhance traditional, broad-scale trophic assignments
of prey. Our results indicate an opportunistic strategy that allows these generalist crustaceans to
effectively utilise a wide range of food sources, including Algae, Diatoms, Fish, Crustaceans,
Molluscs, Echinoderms, Nemerteans, Polychaetes, Mammals, Fungi and other taxa. In addition, a
single snapshot of *Nephrops* diet revealed estimates of biodiversity and species distributions for a
variety of food taxa in their native range, suggesting *Nephrops* could be used as a sentinel organism
for monitoring local biodiversity. Last, we observed high rates of infection by the parasitic
dinoflagellate *Hematodinium*, known to alter burrow-related behaviour, for *Nephrops* populations
in the North and Irish Sea which may have a detrimental impact on predation and fishing. Our
sample collection and modest sequencing data lend further support for the promise of using DNA

metabarcoding as a tool for measuring dietary profiles, biodiversity and monitoring pathogens in

natural *Nephrops* populations that are important to commercial fisheries.

Methodological considerations

DNA metabarcoding offers an enormous opportunity to observe greater taxonomic resolution to

study the feeding ecology of organisms but a few drawbacks in our study require consideration to

understand the dietary repertoire of *Nephrops*. First, isolating DNA from highly digested remains

presents a methodological constraint for successful PCR amplification. Our initial objective was

to include DNA metabarcoding data of the stomach contents of all *Nephrops* specimens, but we

were unable to obtain sufficient PCR amplification and therefore we broadened our findings to gut

contents which achieved greater PCR success. However, the presence of organic compounds in

the stomach and gut of *Nephrops* (e.g., digestive enzymes, Gibson & Barker, 1979s) cause DNA

damage which is one of the main contributors of PCR inhibition (Bernstein et al., 2005; Prieto et

al., 2006; Harms-Tuohy et al., 2016). The variation in diet will affect the secretion of the gastric

juices which play a significant role in PCR inhibition at different concentrations in the gut (Rotllant

et al., 2014). This can be highly problematic as high levels of PCR inhibition will be present in

many genomic DNA extracts that negatively influence PCR amplification (van der Reis et al.,

2018). For more accurate identification and quantification of diet components, the following

strategies could be considered to overcome DNA degradation and PCR inhibition: 1) DNA

extraction protocols with tailored inhibition removal steps (Sellers et al., 2018); 2) incorporation

of alternative proteins to enhance PCR amplification such as bovine serum albumin (BSA) or T4

gene 32 protein (gp32) (Schrader et al., 2012; van der Reis et al., 2018); 3) performing a DNA

repair procedure on genomic DNA template to allow increased PCR amplification success

(Ambers et al., 2014; Davidov et al., 2020) and 4) targeting a shorter fragment than the COI

(313bp) such as the 18S rRNA V9 (~134 bp) to account for advanced digestion and DNA damage

(Albaina et al., 2016).

Second, the choice of primer is an important factor that influences the quality of PCR

amplification and the desired taxonomic resolution. We used a versatile primer set that is known

to be highly effective in amplifying COI across invertebrate phyla (Leray et al., 2013) with an

expectation that high sequencing depth would recover an adequate characterisation of diet items.

Consequently, we generated high sequencing depth with over 18 million sequencing reads but

found over 98% of the data was assigned to *Nephrops* which we considered host contamination,
despite reports of cannibalism between conspecifics (Sardà & Valladares, 1990). Thus, there is
considerable scope for improvement to identify species that are concealed by amplification bias.
One solution to overcome host amplification is simply to target specific taxonomic groups such as
vertebrates using a well-established 12S rRNA assay (Miya et al., 2015; Siegenthaler et al., 2019).
This would greatly enhance the resolution of fish species detected in order to distinguish the
proportion of discards that might contribute to the *Nephrops* diet. However, a multimarker assay
would be needed to obtain a holistic view of eukaryotic diversity in the diet. Therefore, an
alternative solution is the design of a blocking primer which can block amplification of host DNA
in a complex sample (Craig et al., 2014, Weber et al., 2022). This can help preferentially bind non-
host DNA through incorporating nucleotide mismatches for *Nephrops* in the primer with a C3
spacer on the 3′ end which blocks extension of the host PCR amplification (Su et al., 2018; Liu et
al., 2019). Development of a blocking primer assay for *Nephrops* will offer increased opportunity
for species detection but it will require systematic PCR testing to determine the efficiency of
blocking both non-target and target DNA. Nevertheless, our rarefaction estimates illustrate our
COI assay recovered between 49% and 79% of prey diversity across sites for *Nephrops*, and
additional improvements will reveal further fine-scale dietary information.

Dietary profile of **Nephrops norvegicus**

Traditional diet composition analysis relies on visual observation methods of undigested remains
and morphology-based stomach content analysis of *Nephrops* from the Mediterranean and Atlantic
waters off the coast of Portugal show crustaceans and fish to be the main prey-groups (Cristo &
Cartes, 1998). However, this approach is labour-intensive, requires considerable taxonomic
expertise, and the assessment of dietary items is often hampered by the variable rate of prey
digestion in the gut (Hara et al., 2020), and lack of diagnostic features of digested and soft prey
items, thereby underestimating the dietary assemblage. Stable carbon isotope composition of
organic matter can trace the assimilation of nutrients present in animal tissue over a long period
and Santana et al. (2020) used stable isotope analysis of *Nephrops* collected in the West of Ireland
and found suspended particulate organic matter and fish to be important diet components followed
by plankton and invertebrate sources. Yet, this approach does not allow high-resolution analysis
of species-specific diet composition.

Here we implemented DNA metabarcoding of the gastrointestinal tract using a COI primer set to characterise the diet of *Nephrops* from the North and Irish Sea. Overall, our molecular assessment of *Nephrops* gut content displays a broad omnivorous diet. This consisted of 16 Phyla and 35 Classes with 40 MOTUs (36%, ≥98% identity) identified to species level. We report the dominance of Dinoflagellata, Chlorophyta and Bacillariophyta which comprise an average of 49% of the dietary composition for *Nephrops* across sites and this may reflect a nutritional advantage consistent with their ability to suspension feed. Unclassified taxa alone made up an average of 8.7% of consumed prey and an average of 13% consisted of Echinodermata, Chordata and Arthropoda, while the remaining taxa (Ascomyata, Annelida, Basidiomycota, Cnidaria, Discosea, Mollusca, Nemertea, Ochrophyta, Oomycota, Rhodophyta) accounted for an average of 11%. Overall, we observed high dietary overlap among *Nephrops* across sites and this may reflect the common abundance of particulate organic matter and benthic organisms on which they feed. Similarly, morphological stomach content assessment of *Nephrops* from the Eastern and Western Mediterranean and adjacent Atlantic showed no significant differences between sites or seasons, which was explained by the great similarity of the bathyal fauna (Cristo & Cartes, 1998). Nonetheless, we found instances of unique species in the diet of *Nephrops*, with 18 species exclusively found in the East of the North Sea (e.g., Actinopterygii, Aves, Mammalia), eight species in the West of the North Sea (Discosea, Florideophyceae, Oomycota) and five species found in the Irish Sea (Cephalopoda, Chondrichthyes).

Nephrops are benthic animals that burrow in soft sediment and emerge from their burrows to forage. Fluctuations in food availability may present challenging conditions, particularly when their nutritional status is influenced by density-dependent factors. For example, in high density areas, competition for food may limit their scope for growth (Johnson et al., 2013) and increased aggressive social behaviour could drive up the metabolic rate and thereby exhaust energy resources (Parslow-Williams, 1998; Bell et al., 2006; Campbell et al., 2009). The significance of suspension feeding allows individuals to overcome challenging scenarios related to food availability, particularly for females during the breeding season (Watts et al., 2016), and avoiding aggressive encounters between male conspecifics (Sbragaglia et al., 2017). It is suggested that suspension feeding is energetically efficient and can be more profitable for growth compared to active feeding given that 65-68% of the daily energy intake is achieved from suspension feeding (Loo et al., 1996; Parslow-Williams et al., 2002). We found nearly 60% of individuals (55/94) consume 50% or more

taxa that are likely as a result of suspension feeding and this pattern mirrors stable isotope analysis
from Santana et al. (2020). On the other hand, it is reported that females remain berried in normal
environmental conditions to avoid predation during long breeding periods and depend on
suspension feeding as an important strategy to survive starvation when restricted to burrows (Loo
et al., 1993). However, we found females to have similar dietary abundance as their male
counterparts showing patterns of suspension and active feeding and this pattern is in line with
Santana et al. (2020), who revealed male and female *Nephrops* have remarkably similar dietary
profiles. Therefore, it appears female *Nephrops* do not exclusively rely on suspension feeding and
may experience occasional feeding excursions attracted by available food in close proximity to the
burrow opening (Aguzzi et al., 2007; Watts et al., 2016). Thus, the detection of active food foraging
of prey such as invertebrates and fish along with other prey taxa shows the potential of *Nephrops*
to utilise alternative, accessible, and highly nutritional prey to maximise their energy uptake.

More broadly, a combination of DNA metabarcoding and stable isotope analysis could
significantly improve general data gathering to feed ecological models. The last broad scale
description of diets in the North Sea ("Year of the Stomach") was carried out by a European
consortium between 1981-1991 but, because of the labour-intensive nature of dietary identification
at the time, was limited to identifying diets of a few commercial fish species and results were
biased towards hard-bodied species (ICES 1997). Difficulties in identifying stomach contents
often mean that ecological modellers use pooled taxa such as "plankton" or "heterotrophic
benthos" in their diet matrices. Given the likely complexity within these groupings in terms of
variations in numbers of component species in each classification, that are predated upon and their
own trophic levels, this low resolution may limit the validity and utility of some ecological models
(Christensen and Walters 2004; Fulton et al. 2003). Metabarcoding may contribute to a route in
developing much more robust diet matrices for ecological models.

Sentinels of biodiversity
Biodiversity baselines play an important role in understanding the impact of multiple stressors
such as climate change and anthropogenic pressures on biodiversity (Vernooy et al., 2010).
Monitoring programmes enable the observation of ecological temporal variability and enhance our
capacity to manage species and ecosystems (Pöysä & Linkola., 2021). The past decade has
experienced a surge in the development of DNA-based approaches to support rapid biodiversity

assessments (Siegenthaler et al., 2018; Shum & Palumbi, 2021), with recent work harnessing the 'natural sampler' ability of organisms to collect DNA from the environment. For example, Mariani et al. (2019) utilised the water filtering efficiency of sponges to capture environmental DNA (eDNA) to recover highly informative biodiversity assemblages of vertebrate fauna from the Mediterranean Sea and Antarctica ocean. The concept of natural sampling has been further implemented in a range of different organisms to assess biodiversity using DNA metabarcoding of stomach contents from predatory or scavenging crustaceans (Siegenthaler et al., 2019; Nørgaard et al., 2021) to filter-feeding bivalve molluscs (Weber et al., 2022). In the case of *Nephrops*, their potential as a natural sampler is enhanced by their wide distribution from Iceland to as far south of the Canary Islands with their range extending into the eastern Mediterranean Sea. *Nephrops* have a sustainable fishery and are relatively resilient to the effects of trawling in some areas of high fishing pressure with landings maintained at historically high levels for over 40 years (ICES, 2021a). This means that these generalist predators and scavengers could be uniquely resourceful natural samplers in capturing benthic biodiversity.

The variation in food availability in the diet of *Nephrops* enhances the description of local biodiversity in the foraging area and helps build an inventory of species co-occurrence. We observed eight echinoderm species in the diet of *Nephrops* with overall greater occurrences of these species in the East of the North Sea than the West and Irish Sea. Some of these echinoderms represent species with limited geographical distributions showing the centre of their abundances in the North and Irish Seas (e.g. the Spiny mudlark urchin *Brissopsis lyrifera*, brittlestar *Amphiura filiformis* and Black brittlestar *Ophiocomina nigra*). Other echinoderms detected have greater distribution ranges with abundances along the North Atlantic coasts of North America and Europe (Common starfish, *Asterias rubens*) to cosmopolitan species inhabiting cold temperate waters (Common sunstar, *Crossaster papposus*). Identification of these species is important to establish biodiversity baselines particularly for indicator species that help monitor ecosystem health. For example, Sea Star Wasting Disease (SSWD) is an ongoing disease epidemic that leads to behavioural changes, lesions, loss of turgor, limb autotomy, and death characterised by rapid degradation (Hewson et al., 2014). This caused mass mortality of major sea star populations along much of the west coast of North America resulting in the functional extinction of charismatic species (e.g., sunflower sea star, *Pycnopodia helianthoides*, Hamilton et al., 2021). Recently however, a SSWD-like outbreak has been documented and shown to be susceptible in the common

sunstar (*Crossaster papposus*) in European waters from the Irish Sea and further research is needed

to understand the geographical extent of the outbreak (Smith et al., 2022). Although we report no

detections of *S. papposus* in the Irish Sea, we observed an encouraging number of occurrences

(n=10) in the North Sea that illustrate *Nephrops* can be employed to monitor the presence of these

keystone species. However, it is unclear whether the ingestion of echinoderms is a result of active

predation or scavenging. For example, the Common starfish (*Asterias rubens*) is used as a

biological indicator to assess the physical disturbance of bottom-trawl activity causing arm damage

and leaving severed remains scattered along the seafloor (Rogers et al., 2001). While active

predation on sea stars may occur, foraging the severed limbs can explain the occurrences in the

diet of *Nephrops*, including the detection of other benthic species.

479   Fish comprise an important component of the *Nephrops* diet (Cristo & Cartes, 1998;

Santana et al., 2020). The capture of certain fish may present little effort for *Nephrops* but it is

argued that the consumption of fish is subsidised from discards of commercial fishing activity

(Saila et al., 2002; Santana et al., 2020). However, it is unclear what species contribute to their

consumption. We found twelve species of fish in the gut of *Nephrops* but with low occurrences

across our samples. Most fish species consumed are demersal or benthopelagic that are commonly

found in the Northeast Atlantic with several species also distributed throughout the Western

Atlantic. These include haddock (*Melanogrammus aeglefinus*), whiting (*Merlangius merlangus*),

Norway pout (*Trisopterus esmarkii*) and European sprat (*Sprattus sprattus*) which were detected

in the North and Irish Seas, while Atlantic herring, (*Clupea harengus*), common dab (*Limanda*

limanda), European plaice (*Pleuronectes platessa*) and American plaice (*Hippoglossoides*

platessoides) were only detected in the North Sea. Many of these fish are commercially valuable

and they constitute a substantial proportion of discards each year that negatively affects sustainable

exploitation. For example, the mean discard rate for the European plaice between 2013-2017

accounted for a staggering 71% of the total catches from the Celtic Sea and Bristol Channel alone

(ICES, 2018). This high rate of discard mortality is intensified by the adverse impact of bottom

trawling of non-target fish communities, and this is demonstrated from the whiting fishery as

Nephrops-directed otter trawls accounted for 98% (1,030 tonnes) of discards from the Irish Sea in

2020 (ICES, 2021b). Therefore, a considerable proportion of fisheries discards contribute an

important food source for these marine scavengers that offer favourable opportunities for growth.

However, biodiversity estimates will be biased by seasonal fisheries that alter the diversity that are

available to *Nephrops*. Nevertheless, it is reasonable to consider biodiversity estimates of other
naturally occurring species as a result of active foraging such as small demersal fish species with
negligible discards or commercial importance (e.g., red bandfish, *Cepola macrophthalma*;
dragonet, *Callionymus lyra*). Other species that likely involved scavenging remains include
jellyfish (*Leuckartiara octona*), octopus (*Eledone cirrhosa*), catshark (*Scyliorhinus canicular*) and
dolphins (*Phocoena phocoena*, *Lagenorhynchus albirostris*). These species illustrate their
generalist behaviour and highlight the potential of *Nephrops* as natural samplers.

Pathogen surveillance

Further to biodiversity monitoring, diet analyses performed using DNA metabarcoding with
universal primers has an advantage of also detecting parasites as non-target identifications,
demonstrating that parasites can be readily amplified from gut samples (Weber et al., 2022). Our
approach allowed a survey of parasitic dinoflagellates (e.g., *Hematodinium* sp.) that infect a
growing number of crustacean genera globally, many of which are exploited as commercial
fisheries (Stentiford & Shields, 2005). They are considered the most significant known pathogen
of *N. norvegicus* (Field *et al*., 1992; Briggs & McAliskey, 2002). We recovered the presence of
Hematodinium in the gut of *Nephrops* in over 50% of individuals collected from the North and
Irish Seas. The significance of these detections may indicate *Hematodinium*-infection which
presents a negative impact on their behaviour, but the difference was marginal (overall prey
observations: infected: 298, 52%; uninfected: 271, 48%). This implies infected individuals were
exposed to more food resources possibly due to longer foraging durations. While the detection and
presence of the parasitic dinoflagellate *Hematodinium* may represent a major stressor to *Nephrops*,
further research is required to determine both the abundance of prey consumed as well as the
progression and developmental stage of the parasite (Albalat et al., 2012). No attempt was made
to confirm positive infection from our *Nephrops* collection as this finding represents a
serendipitous result, but it is reasonable to expect positive detections are true infections based on
the high prevalence of *Hematodinium*-infected *Nephrops* (Stentiford et al., 2001; Albalat et al.,
2012). Nevertheless, it is important to monitor the prevalence of this pathogen as it negatively
affects swimming performance and burrowing behaviour for *Nephrops* individuals that experience
a greater number of burrow departures and increased foraging time during illuminated periods
compared to uninfected individuals (Stentiford *et al*., 2001). This parasite is known to affect the

hosts' behaviour indirectly through energy depletion (Taylor et al., 1996) and an increased
foraging time is essential to provide the nutritional requirements for the host and the parasite
(Barber et al., 2000; Stentiford et al., 2001). We found a higher number of prey observations in
Hematodinium-infected *Nephrops* compared to uninfected individuals, but infected individuals
have a higher risk of predation and fishing mortality, which in turn can have a pronounced impact
on productivity and commercial catch quantity and value (Stentiford & Neil 2011; Albalat et al.,
2012).
538
Implications for fisheries management
From a practical standpoint, the present results reveal important considerations for the assessment
and management of these commercially valuable stocks. First, our gut content metabarcoding data
clarifies complex food web structure by generating unparalleled resolution of trophic interactions,
and this helps to overcome fragmented data with low resolution that has blurred existing ecosystem
models of North Sea *Nephrops* to support an ecosystem approach to fisheries management
(Mackinson & Daskalov, 2007). Previous models were unable to acknowledge the significance of
Nephrops feeding strategy through both suspension and active feeding for survival, revealed here
with DNA based methods. Second, fisheries discards appear to play an important role in *Nephrops*
nutrition as the fish species consumed are also among the highest recorded in discards from
commercial fishing operations. The capture method used by commercial trawling (i.e., bottom
trawling) will enable the rapid exploitation of fish species and indiscriminately catch non-target
species which then become discarded and can eventually supply the *Nephrops* diet. However,
while reducing bycatch and discarding remain conservation priorities, it is also crucial to
understand and anticipate the potential consequences of reducing discards for species that have
become reliant on them which may play an important structuring role for *Nephrops* as well as other
taxa (Lejeune et al., 2022). Last, it is difficult to ignore the strong pattern of *Hematodinium* sp.
abundance among samples as this parasitic dinoflagellate is known to affect the burrow emergence
behaviour of *Nephrops* (Stentiford et al., 2001) and negatively impacts the quality of the meat
which can render them unmarketable and cause a significant economic loss to fisheries (estimated
between GBP 2-4 million, Field et al., 1992; Stentiford et al., 2001; Albalat et al., 2012; Davies et
al., 2019). Further study is required to investigate the spatial and temporal prevalence patterns of
Hematodinium-infected *Nephrops* as this will be a particularly important consideration for

transferability (Lauria et al., 2015) and the release of hatchery reared individuals in appropriate
locations (Powell et al., 2015). Given the added value of data from gut content metabarcoding we
are in a position to radically improve the biological inference of *Nephrops* which can be
implemented in ecosystem models to provide a greater understanding of trophic links with discards
and help develop tailored management of this unique component of marine biodiversity.
567

Conclusion
DNA metabarcoding offers enhanced taxonomic resolution in *Nephrops* dietary profiles beyond
traditional morphology-based approaches and stable isotopes. The development of DNA
metabarcoding-based inferences recently proposed using organisms to act as natural samplers for
biodiversity assessments. We utilised the gut contents of *Nephrops* to characterise local
biodiversity and estimate the prevalence of *Hematodinium* infected individuals. Our results show
a strong dietary overlap in invertebrates, fish, algae and other taxa from the North and Irish Seas
and strengthen recent work indicating the significance of suspension feeding observed in
Nephrops. Their generalist foraging behaviour allowed detection of indicator species used in
routine environmental impact assessments and revealed the consumption of fish species associated
with the high rate of discards. These patterns expand our understanding in *Nephrops* trophic
ecology and offer interesting perspectives in methodological applications that indicate further
avenues of research.
581

583 **Acknowledgements**
584 We thank the following for their help during the project; Edward Whittle of Whitby Seafoods for
585 providing *Nephrops* samples; Robert Donnelly for technical assistance in the lab.
586

Author contributions
D.A.J. and M.L.J. conceptualised the study with input from P.S.. P.S., J.W-R. and G.S.S.
performed all laboratory work. P.S. carried out statistical analysis and drafted the manuscript with
contribution from D.A.J. and M.L.J. All authors provided feedback on the manuscript.

Data availability statement

Raw Illumina sequences can be found on NCBI's SRA database BioProject ID: PRJNA911567.

The data set including MOTUs, taxonomic assignment, abundances as well as OBITools

bioinformatics   scripts,   R   scripts,   and   sample   barcodes   are   available   at:

https://github.com/shump2/Nephrops-diet.

References

Aguzzi, J., Company, J.B. and Sarda, F., 2007. The activity rhythm of berried and unberried females of Nephrops norvegicus (Decapoda, Nephropidae). Crustaceana, 80(9), pp.1121-1134.

Albalat, A., Gornik, S.G., Beevers, N., Atkinson, R.J.A., Miskin, D. and Neil, D.M., 2012. Hematodinium sp. infection in Norway lobster Nephrops norvegicus and its effects on meat quality. Diseases of Aquatic Organisms, 100(2), pp.105-112.

Albaina, A., Aguirre, M., Abad, D., Santos, M. and Estonba, A., 2016. 18S rRNA V9 metabarcoding for diet characterization: a critical evaluation with two sympatric zooplanktivorous fish species. Ecology and Evolution, 6(6), pp.1809-1824.

Ambers, A., Turnbough, M., Benjamin, R., King, J. and Budowle, B., 2014. Assessment of the role of DNA repair in damaged forensic samples. International journal of legal medicine, 128(6), pp.913-921.

Aydin, K.Y., Gaichas, S.K., Ortiz, I., Kinzey, D.H. and Friday, N., 2007. A comparison of the Bering Sea, Gulf of Alaska, and Aleutian Islands large marine ecosystems through food web modeling.

Barber, I., Hoare, D. and Krause, J., 2000. Effects of parasites on fish behaviour: a review and evolutionary perspective. Reviews in Fish Biology and Fisheries, 10(2), pp.131-165.

Barrat, A., Barthelemy, M., Pastor-Satorras, R. and Vespignani, A., 2004. The architecture of complex weighted networks. Proceedings of the national academy of sciences, 101: 3747-3752.

Bastian, M., Heymann, S. and Jacomy, M., 2009, March. Gephi: an open source software for exploring and manipulating networks. In Proceedings of the International AAAI Conference on Web and Social Media (Vol. 3, No. 1).

Bell M. C., Redant F., Tuck I. 2006. Nephrops species, pp. 412–462. In, and Phillips B. F. (ed.), Lobsters: Biology, Management, Aquaculture and Fisheries. Wiley-Blackwell, Oxford.

Bernstein, H., C. Bernstein, C. M. Payne, K. Dvorakova and H. Garewal, 2005. Bile acids as carcinogens in human gastrointestinal cancers. Mutat. Res. 589: 47–65.

Boyer F, Mercier C, Bonin A, Le Bras Y, Taberlet P and Coissac E 2016. obitools: a unix-inspired software package for DNA metabarcoding. Molecular Ecology Resources, 16:176-182.

Briggs, R.P. and McAliskey, M., 2002. The prevalence of Hematodinium in Nephrops norvegicus from the western Irish Sea. Journal of the Marine Biological Association of the United Kingdom, 82(3), pp.427-433.

Cabodevilla X, Gómez-moliner BJ, Abad N, Madeira MJ (2022). Simultaneous analysis of the intestinal parasitesand diet through eDNA metabarcoding.Integrative Zoology 00, 1–15.

Campbell, N., Allan, L., Weetman, A. and Dobby, H., 2009. Investigating the link between Nephrops norvegicus burrow density and sediment composition in Scottish waters. ICES Journal of Marine Science, 66(9), pp.2052-2059.

Chao, A., 1987. Estimating the population size for capture-recapture data with unequal catchability. Biometrics, pp.783-791.

Christensen V, Walters CJ (2004) Ecopath with Ecosim: methods, capabilities and limitations. Ecol Modell 172:109–139. https://doi.org/10.1016/j.ecolmodel.2003.09.003

Clinchy, M., Sheriff, M.J. and Zanette, L.Y., 2013. Predator-induced stress and the ecology of fear. Functional Ecology, 27(1), pp.56-65.

Craig, C., Kimmerer, W. J., and Cohen, C. S. (2014). A DNA-based method for investigating feeding by copepod nauplii. J. Plankton. Res. 36, 271–275. doi: 10.1093/plankt/fbt104

Cristo, M. and Cartes, J.E., 1998. A comparative study of the feeding ecology of Nephrops norvegicus (L.), (Decapoda: Nephropidae) in the bathyal Mediterranean and the adjacent Atlantic. Scientia Marina, 62, pp.81-90.

da Silva Santana, C.A., Wieczorek, A.M., Browne, P., Graham, C.T. and Power, A.M., 2020. Importance of suspended particulate organic matter in the diet of Nephrops norvegicus (Linnaeus, 1758). Scientific reports, 10(1), pp.1-11.

Dalerum, F. and Angerbjörn, A., 2005. Resolving temporal variation in vertebrate diets using naturally occurring stable isotopes. Oecologia, 144(4), pp.647-658.

Davidov, K., Iankelevich-Kounio, E., Yakovenko, I., Koucherov, Y., Rubin-Blum, M. and Oren, M., 2020. Identification of plastic-associated species in the Mediterranean Sea using DNA metabarcoding with Nanopore MinION. Scientific reports, 10(1), pp.1-11.

Davies, C.E., Batista, F.M., Malkin, S.H., Thomas, J.E., Bryan, C.C., Crocombe, P., Coates, C.J. and Rowley, A.F., 2019. Spatial and temporal disease dynamics of the parasite Hematodinium sp. in shore crabs, Carcinus maenas. Parasites & vectors, 12(1), pp.1-15.

de Sousa, L.L., Silva, S.M. and Xavier, R., 2019. DNA metabarcoding in diet studies: Unveiling ecological aspects in aquatic and terrestrial ecosystems. Environmental DNA, 1(3), pp.199-214.

Delmas, E., Besson, M., Brice, M.H., Burkle, L.A., Dalla Riva, G.V., Fortin, M.J., Gravel, D., Guimarães Jr, P.R., Hembry, D.H., Newman, E.A. and Olesen, J.M., 2019. Analysing ecological networks of species interactions. Biological Reviews, 94: 16-36.

Elsner, M., Jochmann, M.A., Hofstetter, T.B., Hunkeler, D., Bernstein, A., Schmidt, T.C. and Schimmelmann, A., 2012. Current challenges in compound-specific stable isotope analysis of environmental organic contaminants. Analytical and bioanalytical chemistry, 403(9), pp.2471-2491.

Field RH, Chapman CJ, Taylor AC, Neil DM, Vickerman K (1992) Infection of the Norway lobster Nephrops norvegicus by a Hematodinium-like species of dinoflagellate on the west coast of Scotland. Dis aquat Org 13:l-15

Ficetola GF, Coissac E, Zundel S, Riaz T, Shehzad W, Bessière J, Taberlet P and Pompanon F 2010. An in silico approach for the evaluation of DNA barcodes. BMC genomics, 11: 434.

Fulton EA, Smith ADM, Johnson CR (2003) Effect of complexity on marine ecosystem models. Mar Ecol Prog Ser 253:1–16.

Gaichas, S.K., Aydin, K.Y. and Francis, R.C., 2010. Using food web model results to inform stock assessment estimates of mortality and production for ecosystem-based fisheries management. Canadian Journal of Fisheries and Aquatic Sciences, 67(9), pp.1490-1506.

Galloway, A.W., Brett, M.T., Holtgrieve, G.W., Ward, E.J., Ballantyne, A.P., Burns, C.W., Kainz, M.J., Müller-Navarra, D.C., Persson, J., Ravet, J.L. and Strandberg, U., 2015. A fatty acid based Bayesian approach for inferring diet in aquatic consumers. PloS one, 10(6), p.e0129723.

Geller, J., Meyer, C., Parker, M. and Hawk, H., 2013. Redesign of PCR primers for mitochondrial cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic surveys. Molecular ecology resources, 13(5), pp.851-861.

R. Gibson and P. L. Barker, 1979. The decapod hepatopancreas. Oceanography Marine Biol, Vol. 17, pp. 285–346.

Hamilton, S.L., Saccomanno, V.R., Heady, W.N., Gehman, A.L., Lonhart, S.I., Beas-Luna, R., Francis, F.T., Lee, L., Rogers-Bennett, L., Salomon, A.K. and Gravem, S.A., 2021.

Disease-driven mass mortality event leads to widespread extirpation and variable recovery potential of a marine predator across the eastern Pacific. Proceedings of the Royal Society B, 288(1957), p.20211195.

Harms-Tuohy, C.A., Schizas, N.V. and Appeldoorn, R.S., 2016. Use of DNA metabarcoding for stomach content analysis in the invasive lionfish Pterois volitans in Puerto Rico. Marine Ecology Progress Series, 558, pp.181-191.

Hewson, I., Button, J.B., Gudenkauf, B.M., Miner, B., Newton, A.L., Gaydos, J.K., Wynne, J., Groves, C.L., Hendler, G., Murray, M. and Fradkin, S., 2014. Densovirus associated with sea-star wasting disease and mass mortality. Proceedings of the National Academy of Sciences, 111(48), pp.17278-17283.

Hsieh, T.C., Ma, K.H. and Chao, A., 2016. iNEXT: an R package for rarefaction and extrapolation of species diversity (H ill numbers). Methods in Ecology and Evolution, 7(12), pp.1451-1456.

ICES (1997) Database report of the stomach sampling project, 1991. ICES Cooperative Research Report Number 219, pp422

ICES. 2018. Plaice (Pleuronectes platessa) in divisions 7.f and 7.g (Bristol Channel, Celtic Sea). In Report of the ICES Advisory Committee, 2018. ICES Advice 2018, ple.27.7fg. https://doi.org/10.17895/ices.pub.4483

ICES. 2021a. Norway lobster (Nephrops norvegicus) in Division 6.a, Functional Unit 13 (West of Scotland, the Firth of Clyde and Sound of Jura). In Report of the ICES Advisory Committee, 2021. ICES Advice 2021, nep.fu.13, https://doi.org/10.17895/ices.advice.7796.

ICES. 2021b. Whiting (Merlangius merlangus) in Division 7.a (Irish Sea). In Report of the ICES Advisory Committee, 2021. ICES Advice 2021, whg.27.7a, https://doi.org/10.17895/ices.advice.7887.

Johnson, M.P., Lordan, C. and Power, A.M., 2013. Habitat and ecology of Nephrops norvegicus. Advances in marine biology, 64, pp.27-63.

Lauria V, Power AM, Lordan C, Weetman A, Johnson MP (2015) Spatial Transferability of Habitat Suitability Models of Nephrops norvegicus among Fished Areas in the Northeast Atlantic: Sufficiently Stable for Marine Resource Conservation? PLoS ONE 10(2): e0117006. doi:10.1371/journal.pone.0117006

Le Loc'h, F., Hily C., 2005 Stable carbon and nitrogen isotope analysis of Nephrops norvegicus / Merluccius merluccius fishing grounds in the Bay of Biscay (Northeast Atlantic) Can. J. Fish. Aquat. Sci., 62:123-132

Lejeune, B., Mouchet, M.A., Mehault, S. and Kopp, D., 2022. Gut content metabarcoding reveals potential importance of fisheries discards consumption in marine fauna. Canadian Journal of Fisheries and Aquatic Sciences, 99(999), pp.1-10.

Leray, M., Yang, J.Y., Meyer, C.P., Mills, S.C., Agudelo, N., Ranwez, V., Boehm, J.T. and Machida, R.J., 2013. A new versatile primer set targeting a short fragment of the mitochondrial COI region for metabarcoding metazoan diversity: application for characterizing coral reef fish gut contents. Frontiers in zoology, 10(1), p.34.

Liu, C., Qi, R.J., Jiang, J.Z., Zhang, M.Q. and Wang, J.Y., 2019. Development of a blocking primer to inhibit the PCR amplification of the 18S rDNA sequences of Litopenaeus vannamei and its efficacy in Crassostrea hongkongensis. Frontiers in microbiology, 10, p.830.

Mackinson, S. and Daskalov, G., 2007. An ecosystem model of the North Sea to support an ecosystem approach to fisheries management: description and parameterisation. Sci. Ser. Tech Rep., Cefas Lowestoft, 142: 196pp.

Mahé F, Rognes T, Quince C, de Vargas C and Dunthorn M 2014. Swarm: robust and fast clustering method for amplicon-based studies. PeerJ, 2, p.e593.

Majdi, N., Hette-Tronquart, N., Auclair, E., Bec, A., Chouvelon, T., Cognie, B., Danger, M., Decottignies, P., Dessier, A., Desvilettes, C. and Dubois, S., 2018. There's no harm in having too much: A comprehensive toolbox of methods in trophic ecology. Food webs, 17, p.e00100.

Mariani, S., Baillie, C., Colosimo, G. and Riesgo, A., 2019. Sponges as natural environmental DNA samplers. Current Biology, 29(11), pp.R401-R402.

Meyer, J.M., Leempoel, K., Losapio, G. and Hadly, E.A., 2020. Molecular ecological network analyses: an effective conservation tool for the assessment of biodiversity, trophic interactions, and community structure. Frontiers in Ecology and Evolution, 8, p.588430.

Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J.Y., Sato, K., Minamoto, T., Yamamoto, S., Yamanaka, H., Araki, H. and Kondoh, M., 2015. MiFish, a set of universal PCR primers for metabarcoding environmental DNA from fishes: detection of more than 230 subtropical marine species. Royal Society open science, 2(7), p.150088.

Moran, Z., Orth, D.J., Schmitt, J.D., Hallerman, E.M. and Aguilar, R., 2016. Effectiveness of DNA barcoding for identifying piscine prey items in stomach contents of piscivorous catfishes. Environmental Biology of Fishes, 99(1), pp.161-167.

Nørgaard, L., Olesen, C.R., Trøjelsgaard, K., Pertoldi, C., Nielsen, J.L., Taberlet, P., Ruiz-González, A., De Barba, M. and Iacolina, L., 2021. eDNA metabarcoding for biodiversity assessment, generalist predators as sampling assistants. Scientific reports, 11(1), pp.1-12.

Oksanen, J., 2013. Vegan: ecological diversity. R Project, 368.

Parslow-Williams, P., Goodheir, C., Atkinson, R. J. A. & Taylor, A. C. Feeding energetics of the Norway lobster, Nephrops norvegicus in the Firth of Clyde, Scotland. Ophelia 56, 101–120 (2002).

Pöysä, H. and Linkola, P., 2021. Extending temporal baseline increases understanding of biodiversity change in European boreal waterbird communities. Biological Conservation, 257, p.109139.

Prieto, A.I., Ramos-Morales, F. and Casadesus, J., 2006. Repair of DNA damage induced by bile salts in Salmonella enterica. Genetics, 174(2), pp.575-584.

Pringle, R.M. and Hutchinson, M.C., 2020. Resolving food-web structure. Annual Review of Ecology, Evolution, and Systematics, 51(1).

Rogers, S.I., Ellis, J.R. and Dann, J., 2001. The association between arm damage of the common starfish, Asterias rubens, and fishing intensity determined from aerial observation. Sarsia, 86(2), pp.107-112.

Rognes T, Flouri T, Nichols B, Quince C and Mahé F 2016. VSEARCH: a versatile open source tool for metagenomics. PeerJ, 4, p.e2584.

Pompanon, F., Deagle, B.E., Symondson, W.O., Brown, D.S., Jarman, S.N. and Taberlet, P., 2012. Who is eating what: diet assessment using next generation sequencing. Molecular ecology, 21(8), pp.1931-1950.

Post, D.M. and Palkovacs, E.P., 2009. Eco-evolutionary feedbacks in community and ecosystem ecology: interactions between the ecological theatre and the evolutionary play. Philosophical Transactions of the Royal Society B: Biological Sciences, 364(1523), pp.1629-1640./

Powell A, Cowing DM, Scolding JWS, Shepherd-Waring T, Eriksson SP, Lupatsch I, Johnson ML, Shields RJ, Gowland D. (2015). Nephrops norvegicus: Hatchery Handbook. Deliverable 4.1., NEPHROPS project, EU Framework 7 Project no. 286903

Ross, S.T., 1986. Resource partitioning in fish assemblages: a review of field studies. Copeia, pp.352-388.

Rotllant, G., Mente, E., Gisbert, E. and Karapanagiotidis, I.T., 2014. Effects of different diets on the digestive physiology of adult Norway lobster Nephrops norvegicus. Journal of Shellfish Research, 33(1), pp.1-9.

Saila, S.B., Nixon, S.W. and Oviatt, C.A., 2002. Does lobster trap bait influence the Maine inshore trap fishery?. North American Journal of Fisheries Management, 22(2), pp.602-605.

Sardà, F. and Valladares, F.J., 1990. Gastric evacuation of different foods byNephrops norvegicus (Crustacea: Decapoda) and estimation of soft tissue ingested, maximum food intake and cannibalism in captivity. Marine Biology, 104(1), pp.25-30.

Sbragaglia, V., Leiva, D., Arias, A., Antonio García, J., Aguzzi, J. and Breithaupt, T., 2017. Fighting over burrows: the emergence of dominance hierarchies in the Norway lobster (Nephrops norvegicus). Journal of Experimental Biology, 220(24), pp.4624-4633.

Schrader, C., Schielke, A., Ellerbroek, L. and Johne, R., 2012. PCR inhibitors–occurrence, properties and removal. Journal of applied microbiology, 113(5), pp.1014-1026.

Scientific, Technical and Economic Committee for Fisheries (STECF) - The 2021 Annual Economic Report on the EU Fishing Fleet (STECF 21-08), EUR 28359 EN, Publications Office of the European Union, Luxembourg, 2021, ISBN 978-92-76-40959-5, doi:10.2760/60996, JRC 126139.

Sellers, G.S., Di Muri, C., Gómez, A. and Hänfling, B., 2018. Mu-DNA: a modular universal DNA extraction method adaptable for a wide range of sample types. Metabarcoding and Metagenomics, 2, p.e24556.

Shum, P. and Palumbi, S.R., 2021. Testing small-scale ecological gradients and intraspecific differentiation for hundreds of kelp forest species using haplotypes from metabarcoding. Molecular Ecology, 30(13), pp.3355-3373.

Siegenthaler, A., Wangensteen, O.S., Soto, A.Z., Benvenuto, C., Corrigan, L. and Mariani, S., 2019. Metabarcoding of shrimp stomach content: Harnessing a natural sampler for fish biodiversity monitoring. Molecular Ecology Resources, 19(1), pp.206-220.

Smith, S., Hewson, I. and Collins, P., 2022. Not another Pandemic! The first records of sea star wasting disease in European Waters. Authorea. April 20, 2022.

Stentiford, G.D. and Neil, D.M., 2011. Diseases of Nephrops and Metanephrops: a review. Journal of Invertebrate Pathology, 106(1), pp.92-109.

Stentiford G, Neil DM, Atkinson RJA. The relationship of Hematodinium infection prevalence in a Scottish Nephrops norvegicus population to season, moulting and sex. ICES J Mar Sci. 2001;58:814–23.

Stentiford, G.D., Neil, D.M. and Atkinson, R.J.A., 2001. Alteration of burrow-related behaviour of the norway lobster, nephrops norvegicus during infection by the parasitic dinoflagellate hematodinium. Marine & Freshwater Behaviour & Phy, 34(3), pp.139-156.

Stentiford, G.D. and Shields, J.D., 2005. A review of the parasitic dinoflagellates Hematodinium species and Hematodinium-like infections in marine crustaceans. Diseases of aquatic organisms, 66(1), pp.47-70.

Su, M., Liu, H., Liang, X., Gui, L. and Zhang, J., 2018. Dietary analysis of marine fish species: enhancing the detection of prey-specific DNA sequences via high-throughput sequencing using blocking primers. Estuaries and coasts, 41(2), pp.560-571.

Taylor, A.C., Field, R.H. and Parslow-Williams, P.J., 1996. The effects of Hematodinium sp.-infection on aspects of the respiratory physiology of the Norway lobster, Nephrops norvegicus (L.). Journal of Experimental Marine Biology and Ecology, 207(1-2), pp.217-228.

Tyner, S. and Hofmann, H., 2016. geomnet: Network Visualization in the 'ggplot2' Framework. R package version 0.2. 0.

van der Reis, A.L., Laroche, O., Jeffs, A.G. and Lavery, S.D., 2018. Preliminary analysis of New Zealand scampi (Metanephrops challengeri) diet using metabarcoding. PeerJ, 6, p.e5641.

van Zinnicq Bergmann, M.P., Postaire, B.D., Gastrich, K., Heithaus, M.R., Hoopes, L.A., Lyons, K., Papastamatiou, Y.P., Schneider, E.V., Strickland, B.A., Talwar, B.S. and Chapman, D.D., 2021. Elucidating shark diets with DNA metabarcoding from cloacal swabs. Molecular Ecology Resources, 21(4), pp.1056-1067.

Vernooy, R., Haribabu, E., Muller, M. R., Vogel, J. H., Hebert, P. D., Schindel, D. E., Shimura, J., & Singer, G. A. (2010). Barcoding life to conserve biological diversity: Beyond the taxonomic imperative. PLoS Biology, 8(7), e1000417.

Watts, A. J. R., Albalat, A., Smith, I. P., Atkinson, R. J. A. & Neil, D. M. 2016. Seasonal nutritional status in Norway lobsters, Nephrops norvegicus (L.): are females nutritionally compromised over the winter? Mar. Biol. Res. 12, 563–572.

Weber, S., Junk, I., Brink, L., Künzel, S., Veith, M., Teubner, D., Klein, R., Paulus, M. and Krehenwinkel, H., Molecular diet analysis in mussels and other metazoan filter feeders and an assessment of their utility as natural eDNA samplers. Molecular Ecology Resources.

Wieczorek, A.M., Power, A.M., Browne, P. and Graham, C.T., 2018. Stable-isotope analysis reveals the importance of soft-bodied prey in the diet of lesser spotted dogfish Scyliorhinus canicula. Journal of Fish Biology, 93(4), pp.685-693.

Zacharia, P. U. (2017). Trophic levels and methods for stomach content analysis of fishes. In: Course Manual Summer School on Advanced Methods for Fish Stock Assessment and Fisheries Management. Lecture Note Series No. 2/2017. Kochi: CMFRI, pp. 278–288.