MERLIN-DEFICIENT iPSCs AS AN *IN VITRO* MODEL SYSTEM FOR

STUDIYNG *NF2* PATHOGENESIS

Núria Catasús[1], Miguel Torres-Martin[1,2], Inma Rosas[1,2], Bernd Kuebler[3], Gemma Casals-Sendra[1], Helena Mazuelas[4], Alex Negro[1,2], Francesc Roca-Ribas[5], Emilio Amilibia[5], Begoña Aran[3], Anna Veiga[3], Ángel Raya[3], Bernat Gel[4], Ignacio Blanco[1,2], Eduard Serra[4], Meritxell Carrió[4*] and Elisabeth Castellanos[1,2*]

*Equally contributed

1) Clinical Genomics Research Group, Germans Trias i Pujol Research Institute (IGTP); Can Ruti Campus, Badalona, Barcelona, Spain
2) Genetics Department, Germans Trias i Pujol University Hospital (HUGTiP), Can Ruti Campus, Badalona, Barcelona, Spain
3) Medicine Regenerative Program, Institut d'Investigació Biomèdica de Bellvitge. IDIBELL, Hospital Duran i Reynals, Gran Via de L'Hospitalet, 199-203, 08908, L'Hospitalet de Llobregat, Barcelona, Spain.
4) Hereditary Cancer Group, Germans Trias i Pujol Research Institute (IGTP); Can Ruti Campus, Badalona, Barcelona, Spain
5) Otorhinolaryngology Department, Germans Trias i Pujol University Hospital (HUGTiP), Can Ruti Campus, Badalona, Barcelona, Spain

Corresponding authors:

Meritxell Carrió (mcarriol@igtp.cat) and Elisabeth Castellanos (ecastellanos@igtp.cat/ecastellanosp.germanstrias@gencat.cat)

ABSTRACT

NF2-related schwannomatosis is an autosomal dominant syndrome that predisposes to the development of benign tumors of the nervous system. Schwannomas, particularly bilateral vestibular schwannomas (VS), are the most characteristic features of the disease. These tumors are caused by the bi-allelic inactivation of the *NF2* gene in a cell of the Schwann cell lineage. Our current understanding of the molecular pathogenesis of the *NF2* gene, as well as the development of new effective therapies is hampered by the absence of human non-perishable cell-based bearing distinct *NF2* pathogenic variants. With this aim, we generated and characterized three isogenic paired induced pluripotent stem cell (iPSC) lines with single or bi-allelic inactivation of *NF2* by combining the direct reprogramming of VS cells with the use of CRISPR/Cas9 editing. Our results show a critical function of *NF2* for the maintenance of a stable pluripotent state. However, we were able to nudge them towards the Neural Crest-Schwann Cell (NC-SC) axis by applying a 3D Schwann cell differentiation protocol. *NF2*(+/-) and *NF2*(-/-) spheroids homogeneously expressed classical markers of the NC-SC lineage. In addition, *NF2*(-/-) SC-like spheroids showed dysregulation of multiple signaling pathways already described for merlin-deficient SC, and altered in human schwannomas. Therefore, *NF2*(+/-) and *NF2*(-/-) SC-like spheroids can represent a bona fide human *in vitro* cellular model to study the role of *NF2* pathogenesis.

KEYWORDS: NF2-related schwannomatosis; Induced Pluripotent Stem Cells; Schwann Cell; Neural Crest Stem Cell; Vestibular Schwannomas, Schwannomas

INTRODUCTION

NF2-related schwannomatosis (NF2-related SWN; OMIM 101000) is a dominantly inherited autosomal syndrome characterized by the development of multiple tumors of the nervous system (1,2). The incidence of the disease is estimated to be 1 in 28,000 live births with complete penetrance by the age of 60, with 50-60% of cases due to *de novo* mutations (3). The pathognomonic feature of the disease is the development of bilateral vestibular schwannomas (VS), well-circumscribed tumors that arise from Schwann cells (SC) in the vestibulocochlear nerve, due to the bi-allelic inactivation of *NF2*. VS are responsible for hearing loss, deafness and tinnitus(4). Affected individuals may also develop multiple schwannomas in other cranial, spinal and peripheral nerves as well as intracranial and intraspinal meningiomas, and spinal ependymomas (5–8). Cutaneous and ocular manifestations are also common (9,10). Due to their number and anatomical location, these tumors, although histologically benign, are responsible for high morbidity and reduced life expectancy (11). The gold standard treatment for NF2 patients includes surveillance, surgery, radiotherapy and bevacizumab for VS (4), although this last is associated with toxicity if used for extended periods (12), so there is an important need to develop new treatments.

NF2-related SWN is caused by pathogenic variants in the *NF2* tumor suppressor gene (22q12.2), which encodes for merlin (moesin-ezrin-radixin-like) protein, a scaffold protein that interacts with transmembrane receptors and intracellular effectors to mediate mitogenic, survival, proliferation and cell-cell contact signaling pathways (13–15). Changes in merlin result in the inhibition of cell proliferation and the dysregulation of a wide variety of signaling cascades from the cell surface to the nucleus. One of the most studied roles of merlin is its involvement in the Hippo signaling pathway, by repressing YAP/TAZ nuclear translocation. Once in the nucleus, YAP and TAZ associate with the DNA-binding TEAD proteins (TEADs 1-4) to activate the expression of regulators of the cell cycle and apoptosis (16). Merlin also controls cell proliferation by contact inhibition through mechanisms that include the formation of

stable adherens junctions, and the regulation of specific growth factor receptors, the focal

adhesion kinase (FAK) and phosphoinositide 3-kinase (PI3K)/AKT/mTOR pathways (16). In

addition, merlin is known to be involved in other signaling pathways such as the

Ras/Raf/MAPK, TP53 and the Rac1-Pak1 pathways (17,18).

Studying the pathogenesis of the *NF2* gene and developing potential therapies remains

challenging due to the lack of durable preclinical models that recapitulate the genetics and

pathophysiology of NF2-realted tumors. Initial attempts to generate *in vivo* mouse models

revealed the essential role of *NF2* during embryogenesis and early development (19). In

NF2(+/-) mutant mice, the absence of a single *NF2* allele causes the appearance of malignant

epithelial tumors and sarcomas with an aggressive tendency to metastasize (20). Conditional

KO murine models can generate schwannomas and VS (21–23). In addition, several patient-

derived xenograft (PDX) mouse models have also been developed, with varied levels of success

in recapitulating the biology of human NF2-related SWN traits (24). In addition, *in vitro*,

primary SC cultures derived from VS have been set up (25,26), although the limited lifespan of

primary cultures can severely constrain their use for extended experiments. Transformed and

immortalized VS-derived cell lines have been developed to overcome this limitation and have

been widely used as platforms for drug screening (HEI-93, JEI-001) (27,28), although these

models undergo physiological changes due to the immortalization process, severely reducing

the value of any comparison with primary cells.

Induced pluripotent stem cell (iPSC) (29) models have been used in a wide range of diseases

and represent an unlimited source of cells with multiple applications such as disease

modelling, drug development, regenerative medicine, or gene regulation (30–33). The first

NF2-related SWN model reported was an iPSC line harboring a homozygous variant in the *NF2*

gene (UMi031-A-2), obtained by CRISPR/Cas9 gene-editing of an unaffected pluripotent cell

line (34); and, more recently, an iPSC clone was derived from mononuclear bone marrow stem

cells from a mosaic NF2-related SWN patient (35).

One of the main defining characteristics of iPSCs is their ability to differentiate into the three

germ layers, and thus their potential to differentiate into any cell type involved in NF2-related

SWN traits. Protocols have been developed to differentiate iPSC towards the SC lineage,

passing through an intermediate multipotent neural crest stage (36–38). Using these

protocols, it has been shown that iPSC-derived SCs show homogeneous expression of SC

markers and the ability to myelinate axons when co-cultured with dorsal root ganglia neurons

(38). Therefore, the use of iPSCs appears to be a suitable human cellular model to study the

molecular pathogenesis of the *NF2* gene, and the differentiation capacity and growth behavior

of the *NF2* deficient SC-derived cells that originate NF2-related schwannomas.

In the present study, we generated three isogenic iPSC lines harboring single or bi-allelic loss-

of-function variants in the *NF2* gene by direct reprogramming of VS cells and the use of

CRISPR/Cas9 editing. For the first time, these cells were characterized and differentiated

towards the NC-SC lineage, resulting in *NF2*(+/-) and *NF2*(-/-) SC-like spheroids co-expressing

classic markers of the NC-SC axis. In addition, *NF2*(-/-) SC-like spheroids showed dysregulation

of several signaling pathways previously described in *NF2*(-/-) cells, providing a bona fide *in*

vitro cellular model to study the role of *NF2* pathogenesis.

MATERIALS AND METHODS

All procedures performed were in accordance with the ethical standards of the IGTP Human

Research Ethics Committee (CEIC) (PI-17-250), which approved this study, and with the 1964

Helsinki declaration and its later amendments. The *"Comisión de Garantías para la Donación y*

Utilización de Células y Tejidos Humanos, ISCIII") authorized the Project.

Patients and Vestibular Schwannoma samples: tumor samples were obtained from NF2-

related SWN patients clinically managed at the Spanish Reference Centre (CSUR) for

phacomatoses HUGTiP-ICO-IGTP, diagnosed according to standard diagnostic criteria

(**Supplemental Table S1**) (39). Written informed consent was obtained for iPSC generation and

genomic analysis.　Tumor samples were processed as described in Castellanos et al., 2018 (40).

Reprogramming of Vestibular Schwannomas: between $0.2 \cdot 10^6$-$1 \cdot 10^6$ of VS derived cells were

reprogrammed using the CytoTune™-iPS 2.0 Sendai Reprogramming Kit (Thermo Fisher

Scientific), a non-integrative cell reprogramming method holding the four Yamanaka factors

(OCT4, SOX2, KLF4, and cMYC), according to manufacturer's instructions.

Tumor and iPSC molecular characterization: genetic analysis was performed using the

customized I2HCP panel as previously described (41) or Whole Exome Sequencing (WES). The

latter was performed using KAPA HyperCap technology with KAPA HyperExome Probes (Roche)

according to manufacturer's instructions and sequenced in a NextSeq instrument (Illumina).

Analysis of small nucleotide variants was performed with Mutect2 and annotated with

Funcotator (42). Potential SNV and small indels were detected among the original cell lines and

the paired CRISPR lines, GATK 4.2.6.1 was used for the entire process (42) (**Supplemental**

Material and Methods). To detect *NF2* splicing variants, tumors were studied both at RNA and

DNA level. *NF2* variants were confirmed by Sanger Sequencing. SNP-array analysis was

assessed using Illumina HumanOmniExpress v1 BeadChips (730,525 SNPs) according to

manufacturers' instructions and as described in Supplemental Material and Methods.

CRISPR/Cas9 gene edition in iPSC lines: CRISPR/Cas9 editing was performed with the ArciTect

ribonucleoprotein (RNP) system (STEMCELL Technologies). The designed sgRNA targeted exon

2 of the *NF2* gene (GTACACAATCAAGGACACAG) using the Synthego - CRISPR Design Tools

(https://www.synthego.com/products/bioinformatics/crispr-design-tool).　　　TransIT-X2®

Dynamic Delivery System (Mirus) was used for transfection according to manufacturers'

instructions.

iPSC culture: iPSCs were grown on growth factor-reduced Matrigel (BD Biosciences)-coated 6-

well plates and cultured in mTESR Plus medium (STEMCELL Technologies). iPSCs were split

using Accutase (Merk) and cells were seeded with Rock Inhibitor (STEMCELL Technologies)

(1:1000) for 24h. When required, visibly differentiated cells were removed manually from the

culture of NF2(-/-) cells. In this study, the NF2(+/+) FiPS line (FiPS Ctrl 1-SV4F-7) was used for all

experiments as a control.

Neural Crest (NC) differentiation: differentiation of iPSC lines into NC was performed as

previously described in Menendez et al. (2013) (37) with minor modifications (38). $9x10^4$ iPSCs

were seeded on matrigel-coated dishes in mTESR Plus medium. The following day, the medium

was replaced with Neural Crest Differentiation Media (**Supplemental Material and Methods**)

and was replaced every day. NC were maintained in this medium and split with Accutase when

necessary.

Differentiation into SC in 2D: to establish SC differentiation, $0.4x10^6$ NC cells/well were plated

onto 0.1mg/mL poly-L-lysine (Sigma) and 4µg/mL laminin (Gibco) 6-well plates and cultured in

SC differentiation media (SCDM) (**Supplemental Material and Methods**) as previously

described (38).

Differentiation towards SC in 3D: NC cells were detached with Accutase and, $2.25 \cdot 10^6$

cells/well were seeded onto AggreWell TM800 24-well plates (Stem Cell Technologies) in 2mL

SCDM (described above). The medium was changed twice a week removing 1 mL and replacing

1 mL of fresh SCDM. On days 7, 14 and 30 spheroids were collected and processed for

subsequent analysis.

Flow cytometry: accutase-dissociated cells were resuspended in PBS-0.1% BSA and incubated

with p75 primary antibody and subsequently incubated with Alexa Fluor 568 secondary

antibody, followed by the incubation of Hnk1 primary antibody that was detected with Alexa

Fluor 488-conjugated secondary antibody. Antibodies were incubated 30 min on ice. Flow

cytometry analysis was performed using BD LSR Fortessa SORP and BD FACS Diva 6.2 software.

Immunocytochemistry: cells or spheroids were fixed in 4% paraformaldehyde in PBS for 15min

at room temperature (RT), permeabilized with 0.1%Triton-X 100 in PBS for 10 min at RT,

blocked in 10% FBS in PBS for 15 min at RT, and incubated with the indicated antibodies

(**Supplemental Table M1**) overnight at 4°C. Secondary antibodies Alexa Fluor 488 and Alexa

Fluor 568 were incubated for 1h at RT. Nuclei were stained with DAPI and images captured

using LEICA DMIL6000 and LASAF software.

RNA processing, sequencing and analysis: Total RNA extraction from iPSCs, NC cells and SC-

differentiating spheroids was extracted with the 16 LEV simplyRNA Purification kit (Promega),

following manufacturer's instructions. RNA was quantified with a Nanodrop 1000

spectrophotometer (Thermo Scientific). The polyA RNA libraries were sequenced on an

Illumina Novaseq 6000 in 150 bp pair-end mode. For RNAseq processing, genome GRCh37.p13

and gene annotation gencode version 19 were used. Reads were aligned with STAR version

2.7.10a (43) and count files were produced with QoRTs version 1.3.6 (44). These raw counts

were used for DESeq2 differential expression (DE) analysis with Wald significance test and

Benjamini & Hochberg correction (FDR) correction. A corrected p-value of 0.01 and an absolute

Log2-fold change of 1 was used for considering a gene as DE. Volcano plots were performed

using EnhancedVolcano (with the https://github.com/kevinblighe/EnhancedVolcano) outputs

from Deseq2. To account for more direct measurement, barplots in Figure 2 were instead

normalized by Trimmed Mean of M-values from EdgeR (45) and then RPKMs were calculated

for every gene. The *vst* normalization from DESeq2 (46) was used for unsupervised cluster

heatmaps with Euclidean metric and Ward method. PCA features were chosen using the top

2000 genes with the highest SD over vst normalization. Pathway enrichment analyses for DE

genes were performed using GSEA pre-ranked test (47) using the "stat" value from Deseq2. A

FDR <0.05 was considered as significant. Single sample GSEA (ssGSEA) with MSigDB version

2022.1.Hs (48) was used when comparing more than two groups of samples. Enrichr (49,50)

was used only with DE genes as input. The 41 raw fastq files schwannomas for comparison

were retrieved from the EGA repository with accession code EGAD00001002722, and were

processed with the same methodology described above. For re-analysis of this cohort together

with our set, vst values were recalculated for all of them without correction for batch.

RESULTS

1. <u>Generation of IPSC harboring single or bi-allelic pathogenic variants in the *NF2* gene</u>

Three VS from three independent NF2-related SWN patients that met standard diagnostic criteria (39) were used for reprogramming (VS-25, VS-245 and VS-267) with the aim of obtaining iPSC clones harboring pathogenic variants present in NF2 patients. Molecular characterization of these tumors through the I2HCP panel (41) and SNP-array analysis uncovered the *NF2* alterations reported in **Supplemental Table S1** and **Supplemental Figure S1**. Except for the alteration in the *NF2* locus, no pathogenic variants were identified in any known oncogene or tumor suppressor gene. The non-integrating Sendai virus vector (SeV) encoding the 4 Yamanaka factors (29) was used to induce pluripotency of VS-dissociated cells and we determined the *NF2* genotype of the generated single-cell clones through Sanger sequencing (**Figure 1A, Supplemental Table S2**).

From VS-25 and VS-267, only clones carrying the germline *NF2* pathogenic variant (*NF2*(+/-)) were obtained. One clone derived from each tumor was selected for further characterization. From VS-245, a VS coming from a mosaic patient, we obtained both *NF2*(-/-) and *NF2*(+/+) iPSC clones, named VSi-245. As VSi-245(+/+) clones were reprogrammed from an unaffected cell, these were not further characterized. In contrast, all nine VSi-245 *NF2*(-/-) iPSC clones analysed sustained the exogenous expression of Yamanaka factors by the Sendai (SeV) construct for more than 40 passages after reprogramming and, therefore, were also discarded for the rest of the study (**Supplemental Figure S2**).

With the aim of achieving iPSCs with complete *NF2* inactivation, and given the low efficiencies obtained in the generation of *NF2*(-/-) clones by VS reprogramming, we used the CRISPR/Cas9 technology to edit the *NF2* gene in both, the *NF2*(+/-) VSi-25 and VSi-267 iPSCs and a control *NF2*(+/+) iPSC line (FiPS Ctrl 1-SV4F-7, registered in the Spanish National Stem Cell Bank/ ESiO44C in https://hpscreg.eu/)). Thus, using CRISPR/Cas9 RNP complexes, we targeted exon 2

of the *NF2* gene and obtained single-cell *NF2*(-/-) iPSC clones from VS-derived lines (named 25-

CasD7, when derived from VSi-25, and 267-CasD2 when derived from VSi-267) and isogenic

NF2(+/-) and *NF2*(-/-) iPSC clones derived from the control line (named FiPS-CasB2 and FiPS-

CasH6, respectively). Pathogenic variants in *NF2* were fully characterized through PCR and

Sanger sequencing and the *NF2* coding region was cloned to confirm compound heterozygous

clones (**Table 1**). In addition, all *NF2* genotypes were confirmed through a merlin western blot

(**Figure 1B**). From this point on, we worked on the characterization of three pairs of isogenic

iPSC lines: three with one truncating variant in *NF2* and three lines carrying the bi-allelic

inactivation of *NF2* (two pairs VS-reprogrammed and one from a control iPSC line (**Table 1**)).

iPSCs: induced pluripotent stem cell; VS: vestibular schwannoma; FiPS; fibroblasts induced pluripotent

stem cell; Ctrl: Control; WT: wild-type

1.1. Characterization of *NF2* (+/-) and *NF2* (-/-) iPSC

All clones analysed expressed cell surface proteins and transcription factors associated with

pluripotency: Oct4, SSEA-3, SSEA-4, Nanog, Sox2, Tra-1-60 and Tra-1-81; were positive for

alkaline phosphatase staining; and showed karyotype stability after at least 20 passages (46,

XY) (**Supplemental Figure S3, A-C**). Although all iPSC clones expressed the typical pluripotent

markers, we could observe differences in clone morphology between *NF2*(+/-) and *NF2*(-/-) cell

lines. Whereas *NF2*(+/-) showed classical iPSC colony morphology, edited *NF2*(-/-) iPSCs

presented less compact colony formation, eventually aggregating in the center of the colony,

growing upwards and undergoing major spontaneous differentiation compared with the iPSC

control line (**Figure 1C**). Furthermore, *NF2*(+/-) iPSCs showed the capacity to differentiate into

the three primary germ layers *in vitro* (mesoderm, endoderm, and ectoderm) through

embryoid bodies (EBs) formation. In contrast, *NF2*(-/-) lines could not be differentiated using

standard EBs generation protocols and required direct differentiation to acquire expression of

the three germ layers (**Figure 1D, Supplemental Table S3**).

Genomic characterization of *NF2*(+/-) VSi-25 and VSi-267 iPSCs selected clones by SNP-array

analysis and WES showed no differences with respect to the tumor of origin, with the

exception of the absence of the the second hit in the *NF2*(+/-) lines (**Supplemental Figure S1**).

Generated *NF2*(-/-) iPSC lines revealed no pathogenic off-target alterations by WES analysis of

the edited lines (**Supplemental Table S4**).

We have therefore generated three *NF2* iPSCs isogenic pair lines (*NF2*(+/-) and *NF2*(-/-)) with

three different genomic backgrounds and cells of origin, providing an unlimited source of cells,

consisting of a biological triplicate system. The phenotype observed in all three *NF2*(-/-) lines,

which showed relevant phenotypic and differentiation abnormalities, suggests that the *NF2*

gene may play a role in maintaining a purely pluripotent state.

2.   *NF2* iPSC differentiation towards the NC-SC lineage

Given that the cells that initiate schwannoma formation in NF2-related SWN are *NF2*(-/-) cells

derived from the SC lineage, we applied a differentiation protocol towards the NC-SC axis. As a

first step, we generated NC by applying a chemically defined differentiation protocol as

previously reported (37). After ten days of NC differentiation, cells achieved NC morphology

(**Figure 2A**). Cell identity was assessed by flow cytometry of NC markers, NGFR (Nerve Growth

Factor Receptor) (p75) and HNK1 (CD57), at early passages of the NC differentiation process

(p4) and in a mature NC state (p8). *NF2(+/+)* an *NF2(+/-)* cells showed high expression of both

markers already at p4 which increased up to more than 80% at a mature NC state. *NF2(-/-)*

cells already showed some expression of p75 and Hnk1 at the iPSC stage which increased at

p4, but did not acquire the same percentages of p75 and Hnk1-positive cells at the mature NC

stage (65-80%), compared with their controls *NF2(+/+)* and *NF2(+/-)* counterparts (**Figure 2B**).

Immunocytochemistry and transcriptome analysis (RNA-seq) of NC derived cells showed that

all lines expressed NC markers Sox10, TFAP2A and p75 and showed no expression of

pluripotency markers POU5F1 (Oct4), Nanog and Sox2 (**Figure 2C and 2D**). However, some *NF2*

(-/-) NC cells were also expressing S100B at both the RNA and protein levels, which was

consistent with the heterogeneity detected by flow cytometry assays. Nonetheless, although

NF2-deficient lines exhibited a distinct phenotype, they could be differentiated towards a NC-

like identity, were able to maintain self-renewal capacity for at least 18 passages and could be

cultured after several freeze-thaw cycles.

A last step in the differentiation towards SC, consisted of inducing the differentiation of the

established NC cultures into a SC state. Thus, NC cells were cultured in SC differentiation

medium (SCDM) for 30 days, as previously described (37,38). However, five days after inducing

SC differentiation, *NF2*(+/-) and *NF2*(-/-) cells showed difficulties in maintaining adherence to

culture dishes, in contrast to the control line, which after seven days of differentiation already

showed a more elongated spindle morphology resembling SC-like cells (**Supplemental Figure**

S4). We reasoned that the poor adherence capacity observed compromised their ability to

differentiate towards a SC identity in the standard 2D stablished conditions, and therefore, we

tested their capacity to differentiate in 3D conditions, following a previously established

protocol (51) with minor modifications.

2.1 NC-SC lineage marker analysis on *NF2*(+/-) and *NF2*(-/-) spheroids

To evaluate the SC differentiation capacity of *NF2*(+/-) and *NF2*(-/-) cell lines in 3D, we studied

the transcriptome at three time points (7, 14 and 30 days) of the differentiation process.

Principal component analysis (PCA) revealed that iPSCs, NCs and SCs triplicates clustered

according to cell type, regardless of genotype or differentiation state (**Figure 3A**).

At the RNA level, we observed that *NF2*(+/+) differentiating spheroids expressed typical SC

lineage markers (S100B, MPZ or PLP1) already at day 7. At day 14 of differentiation, *NF2*(+/+)

spheroids were positive for p75 and S100B markers identified by immunostaining (**Figure 3B**)

and expressed SC markers closely resembling those expected in SCs (**Supplemental Figure S5**).

However, at day 30 of differentiation the expression of some SC markers decreased (SOX10,

PLP1) whereas some central nervous glial markers appeared (FABP7, ASTN1), suggesting loss of

SC commitment. For this reason, we decided to analyse samples at day 14 of differentiation

(**Supplemental Figure S5**). *NF2*(+/-) VSi-25 and VSi-267 cell lines formed spheroids that stained

positive for p75 and S100B markers, as did the three *NF2*(-/-) cell lines (**Figure 3B**). However,

we were unable to obtain spheroids from the FiPS-CasB2 *NF2*(+/-) edited cell line despite

several attempts (**Supplemental Figure S6**).

RNAseq analysis revealed that *NF2*(+/-) and *NF2*(-/-) cell lines expressed high levels of SC

markers such as CDH19, GAP43, MPZ, PLP1 or S100B at 14 days of differentiation, in a similar

way to the *NF2*(+/+) control line, indicating that NC cells were progressing from a precursor

state into a committed SC phenotype (**Figure 3C**). Finally, we used a recently published *in vitro*

NC-SC expression roadmap (50) to analyze the *NF2*(-/-) cell lines (51) and they displayed a very

similar NC-SC expression roadmap to that previously determined for *NF2*(+/+) differentiating

SCs in 2D (**Supplemental Figure S7**) (51). Overall, we successfully obtained *NF2*(+/-) and *NF2*(-/-

) SC-like spheroids, that express NC-SC lineage markers at the RNA and protein levels, by

applying 3D SC differentiation conditions in three isogenic paired lines.

3.   **Merlin-related pathway analysis in the *NF2*(-/-) SC-like cells**

To better characterize *NF2*(-/-) SC-like spheroids and the effect of the absence of *NF2* on them,

we performed differential expression analysis of the distinct genotypes at 14 days of SC

differentiation stage. Only 125 genes were differentially expressed (DEG) when comparing

NF2(+/+) and *NF2*(+/-) genotypes; most of them related to cellular polarity, cell adhesion and

to the mTOR-pI2K-Akt signalling pathway; and some of them directly regulated by merlin, such

as CHL1 (52) (**Supplemental Figure S8A-B**). A higher number of DEG were found when

comparing the *NF2*(+/+) and *NF2*(-/-) (**Supplemental Figure S8C**), as well as when comparing

NF2(+/-) and *NF2*(-/-) (2874 and 3447 respectively) (**Figure 4A**), indicating that the second hit

in *NF2* is the driver of the major expression changes in these cells.

As it was not possible to obtain *NF2*(+/+) lines from SWN-related NF2 patients, we focused on

comparing *NF2*(+/-) and *NF2*(-/-) SC-like spheroids and further investigated DEG by functional

enrichment analyses. These showed significantly enriched Hallmark signaling pathways in

NF2(-/-) SC-like spheroids, including mTORC1, NFKβ, p53, Hedgehog, and IL6-JAK-STAT3,

among others, most of them directly or indirectly regulated by merlin. Other altered signaling

pathways were Epithelial Mesenchymal Transition (EMT), cholesterol homeostasis and fatty

acids metabolism, ROS signaling, hypoxia and apoptosis, Wnt/β-cantenin and finally (α and γ)

interferon response (**Figure 4B**). Similar results were obtained when comparing *NF2*(+/+) vs the

NF2(-/-) SC-like spheroids (**Supplemental Figure S8D**).

Further analyses were performed using single sample gene set enrichment analysis (ssGSEA) to

test whether signaling pathways previously described in merlin-deficient cells were

systematically altered in all *NF2(-/-)* SC-like spheroids. First, we examined the levels of TAZ

target genes since their expression is negatively regulated by merlin. As expected from merlin

deficiency, TAZ target genes were upregulated in *NF2*(-/-) SC-like spheroids. Similar results

were observed for TEAD2 target genes, although we observed variability between cell lines

(**Figure 4C**). Expression levels of genes regulated by the focal adhesion kinase (FAK) and

phosphoinositide 3-kinase (PI3K)/AKT/mTOR pathways, activated in the absence of merlin,

were upregulated in all three *NF2*(-/-) SC-like spheroid lines. (**Figure 4C**).

We also examined the expression of specific genes known to be altered in both *NF2*(-/-)

primary SCs and schwannomas and compared their normalized expression levels between the

three genotypes. Alterations were found in laminin receptor α6 integrin (Itga6), α-dystroglycan

(Dag1), and collagen (Col1a2), consistent with an alteration of basal cytoskeletal organization

in *NF2*(-/-) SC-like spheroids when comparted to *NF2*(+/+) and *NF2*(+/-) spheroids (**Figure 4D**).

In addition, analysing the status of the Hippo pathway signaling, we were able to confirm that

some of the major YAP target genes described in SC, involved in cell growth, proliferation and

migration, were significantly up-regulated in SC-like spheroids lacking *NF2* (*CCND1, CYR61,*

ANKRD1, AXL) (**Figure 4D**).

Altogether, these findings showed that the alterations and differentiation properties of *NF2*(-/-

) iPSC-derived SC-like spheroids could be attributed to the lack of merlin. Moreover, these

results highlight a strong correlation between the previously described altered signaling

pathways and gene expression profile in merlin-deficient SC and that observed in *NF2*(-/-) iPSC-

derived SC-like spheroids at 14 days of differentiation, indicating that these cells could

represent a genuine *in vitro* human NF2 SC model system.

GSEA analysis comparing *NF2*(+/-) and *NF2*(-/-) SC-like cells revealed that other relevant merlin

targets previously found to be altered in schwannomas were also upregulated in the *NF2*(-/-)

spheroids, such as the previously mentioned MEK, PI3K-AKT/mTOR, TP53, TNFα-NFκB

signaling, or *ERBB2* signaling pathways, inflammatory response as well as the EMT (**Figure 4B**).

We confirmed this by specifically interrogating expression levels of *MAPK1*, *KRAS*, *AKT*, *TP53*,

NFKB and *VEGFA* (**Supplemental Figure S9A**). These analyses indicated that *NF2*(-/-) SC-like

cells could recapitulate profile signatures observed in NF2-related schwannomas.

To further explore this hypothesis, we compared the transcriptome of these cells at 14 days of

differentiation with published transcriptomes of 41 schwannomas (EGAD00001002722) (53).

PCA revealed that the second component was driven by the *NF2* genotype (**Supplemental**

Figure S9B), thus, we extracted the top 500 genes with higher weight for each feature, and

selected sub-clusters where schwannomas and *NF2*(-/-) SC-like cells showed a similar pattern

of expression compared to cells harboring at least one copy of *NF2* (**Figure 5A**). EnrichR

pathway analysis of these 500 genes showed 160 (out of 273) of them were upregulated in

NF2(+/-) and *NF2*(+/+) SC-like cells and were enriched in cell motility and extracellular

organization, therefore related to merlin's function (**Figure 5B**). On the other hand, 113 genes

in merlin deficient schwannomas and SC-like cells were enriched in axonogenesis and

neurogenesis (**Supplemental Figure S9C**). These results suggest that schwannomas and NF2(-/-

) SC-like spheroids share a common pattern of gene expression for key pathways related to

NF2 biology compared to wild type cells.

Overall, we successfully applied 3D SC differentiation conditions to NF2(+/-) and NF2(-/-) to

obtain spheroids expressing NC-SC lineage markers at RNA and protein level (**Figure 5C**). In

addition, gene expression of iPSC-derived merlin-deficient differentiating SCs showed that

these cells constitute a bona fide cellular model for the study of the NF2 role in Schwann cells,

and potentially in any cell type associated with NF2 pathogenesis.

DISCUSSION

In this work we show the generation of an unperishable human iPSC-based model for the

study of the molecular pathogenesis of the NF2 gene. We have differentiated the generated

isogenic iPSCs towards the NC-SC lineage and shown that NF2(-/-) SC-like spheroids share gene

expression patterns with NF2(-/-) cells and schwannomas.

We first aimed to generate NF2(-/-) iPSCs from VS reprogramming to obtain cell lines that

recapitulate the genetic background of a VS constituent cell. Our results showed low

efficiencies in reprogramming NF2-deficient VS cells and we were not able to generate NF2(-/-)

iPSCs through this process, hypothesizing that NF2 could have a relevant role in achieving

pluripotency due to its essential role in embryogenesis and early stages of development (19),

which is also consistent with the reported essential function of NF2 in haploid human iPSCs

and leukemia cells (54,55). Reprogramming resulted in the generation of two NF2(+/-) lines

(VSi-25 and VSi-267), and their NF2 deficient isogenic cell lines (25-CasD7, 267-CasD2) were

obtained by CRIPR/Cas editing. A third isogenic paired iPSC line was obtained by editing the

control FiPS iPSC line (NF2(+/-) FiPS_CasB2 and NF2(-/-) FiPS-CasH6). Thus, in this study we

generated a triplicate of paired NF2(+/-) and NF2(-/-) iPSC lines by combining direct

reprogramming of VS cells and *NF2* gene editing, as has been recently reported (34). In our

hands, the three *NF2*(-/-) iPSC lines required extensive experimental care, showed aberrant

colony morphology and were susceptible to spontaneous differentiation. No differences were

observed between the three lines, hinting that observed phenotypes are due to the lack of

merlin and not to the iPS cell of origin. All these factors taken together point to an essential

role for *NF2* in the maintenance of pluripotency of diploid cells. The three isogenic paired iPSC

lines generated in this study represent a valuable and non-perishable source of cells with

single or bi-allelic inactivation of *NF2*, which could be used as a NF2-cell based model to

develop further assays and to study the role of *NF2* in human cells (56).

Here, we showed that both *NF2*(+/-) and *NF2*(-/-) iPSCs have the capacity to differentiate into a

NC identity, although *NF2*(-/-) NC cultures exhibited an unstable NC phenotype, with some

cells undergoing spontaneous differentiation and expression of S100B and requiring a higher

confluence in culture to maintain their NC identity, consistent with the role of merlin in

cytoskeleton maintenance and control of proliferation by cell-cell contact inhibition (57–59).

Also, in line with these findings, in the process of differentiation of NCs into SCs, we observed

that both *NF2*(+/-) and *NF2*(-/-) cells were easily detached from the cell culture surface,

eventually leading to cell death and preventing the generation of SC-like cultures. However,

and most importantly, we overcame this limitation by successfully generating *NF2*(+/-) and

NF2(-/-) spheroids growing in suspension, as had been previously reported for other SC-based

spheroids (51). These spheroids expressed classical NC-SC markers after seven days in SCDM

and preserved a previously characterized *in vitro* NC-SC differentiation expression roadmap as

control iPSCs (51). Therefore, for the first time, we have achieved merlin-deficient SC-like cells.

NF2(-/-) SC-like spheroids showed deregulation of several signaling pathways as previously

described in *NF2*(-/-) SC. As expected for merlin deficiency the Hippo pathway was

downregulated and YAP/TAZ targets were overexpressed (*Cyr61, AXl1, ANKRD1, CCND1*) (60),

while mTORC1 signaling was upregulated (61), as previously described in schwannomas,

meningiomas and arachnoidal merlin- deficient cells, (60), consistent with the hypothesis that

deregulation of mTORC1 activation, in addition to the Hippo pathway, underlies the aberrant

growth and proliferation of NF2-deficient cells (62). Differentiated *NF2*(-/-) SC-like cells also

showed upregulation of well-known merlin-related cytoskeletal organization markers, such as

Itga6, Dag1 and Col1a2 (63). Furthermore, *NF2*(-/-) SC-like cells overexpressed inflammatory

response, ROS and NFkB-related pathways, most probably by Rac1 deregulation as described

in NF2-tumor models and merlin-deficient cells (53,64–67). Finally, merlin-deficient SC-like

cells also showed upregulation of TP53, VEGF and epithelial-to-mesenchymal transition

signaling, which is known to be deregulated in schwannomas (17,53,68–72). These results

confirmed the strong correlation between the described and observed changes in signaling

pathways and gene expression due to merlin deficiency, suggesting that the cells developed in

this study could be a reliable *in vitro* human NF2 SC model system to study the molecular

pathogenesis of *NF2*. In addition, because there are expression patterns described in

schwannomas that reproduce in merlin-deficient SC-like spheroids; if they were validated *in*

vivo, these cells would be an *in vitro* cellular model for schwannoma cells, which could be used

as a platform for drug screening.

CONCLUSION

In this study, we generated three isogenic paired *NF2*(+/-) and *NF2*(-/-) iPSC lines through

reprogramming VS and *NF2* editing. According to our results, the *NF2* gene seems to play a

relevant role in the maintenance of pluripotency. For the first time, iPSCs with different *NF2*

genotypes were differentiated into NC-SC lineage cells, although *merlin*-deficient iPSCs

exhibited an altered differentiation capacity. The application of a 3D SC differentiation

protocol resulted in the successful generation of *NF2*(+/-) and *NF2*(-/-) SC-like spheroids

expressing NC-SC lineage markers. In addition, *NF2*(-/-) SC-like spheroids showed dysregulation

of several signaling pathways previously described in merlin-deficient Schwann cells, such as

mTOR and Hippo-YAP, amongst others. Therefore, *NF2*(-/-) SC-like spheroids constitute a

valuable resource for studying *NF2* biology in Schwann cells and potentially in any cell type

associated with *NF2* pathogenesis.

CONFLICT OF INTEREST STATEMENT

The authors declare no competing interests

ACKNOWLEDGEMENTS

We thank the HGTP Clinical Services and staff for their collaboration in generating and

collecting patient samples. We thank the IGTP Flow Cytometry and IGTP Genomics core

facilities and their staff for their contribution and technical support. We would like to

acknowledge the constant support of the different NF lay associations: *Asociación de*

Afectados de Neurofibromatosis, Chromo22 and ACNefi. This study has been funded by the

Chromo22 Patients association; the *Instituto de Salud Carlos III* through the project PI20/00215

(Co-funded by European Regional Development Fund "A way to make Europe"), and through

the project AC22/00033, partner of the EJP RD. The EJP RD initiative has received funding from

the European Union's Horizon 2020 research and innovation program under grant agreement

N°825575"; funded also by *Fundació La Marató de TV3* (126/C/2020), the Children's Tumor

Foundation (CTF-2019-05-005, CTF-2022-05-005), Fundación Proyecto Neurofibromatosis, the

Catalan NF Association (AcNeFi), and the Government of Catalonia (SGR-Cat 2021 - 00967).

AUTHOR CONTRIBUTIONS

NC: Collection and/or assembly of data; Data analysis and interpretation; Manuscript writing;

MT-M, BG: Data analysis and interpretation; **AN, BK, IR, GC-S, BA:** Collection and/or assembly

of data; **HM, AV, AR, IB:** Scientific input; **FR-R, EA**: Provision of study material or patients; **ES,**

MC, EC: Conception and design; Financial support; Manuscript writing; Final approval of

manuscript.

DATA AVAILABILITY STATEMENT

The authors confirm that the data supporting the findings of this study are available within the

article [and/or] its supplementary materials. Raw data for WES and RNAseq is allocated at the

European Geno-Phenome Archive (EGA) with accession number (XXXXXXXXX) and will be made

available to external researchers upon reasonable requests.

REFERENCES

1. Evans DGR, Moran A, King A, Saeed S, Gurusinghe N, Ramsden R. Incidence of vestibular schwannoma and neurofibromatosis 2 in the North West of England over a 10-year period: Higher incidence than previously thought. Otology and Neurotology. 2005;26(1):93–7.

2. Plotkin SR, Messiaen L, Legius E, Pancza P, Avery RA, Blakeley JO, et al. Updated diagnostic criteria and nomenclature for neurofibromatosis type 2 and schwannomatosis: An international consensus recommendation. Genetics in Medicine. 2022;24(9):1967–77.

3. Evans DG, Bowers NL, Tobi S, Hartley C, Wallace AJ, King AT, et al. Schwannomatosis: A genetic and epidemiological study. J Neurol Neurosurg Psychiatry. 2018 Jun 14;

4. Lloyd SKW, Evans DGR. Neurofibromatosis type 2 (NF2): diagnosis and management. 1st ed. Vol. 115, Handbook of clinical neurology. Elsevier B.V.; 2013. 957–67 p.

5. Evans DGR, Huson SM, Donnai D, Neary W, Blair V, Newton V, et al. A genetic study of type 2 neurofibromatosis in the United Kingdom . II . Guidelines for genetic counselling. 1992;(3):847–52.

6. Parry DM, Eldridge R, Kaiser-Kupfer MI, Bouzas EA, Pikus A, Patronas N. Neurofibromatosis 2 (NF2): Clinical characteristics of 63 affected individuals and clinical evidence for heterogeneity. Am J Med Genet. 1994;52(4):450–61.

7. Baser ME, Makariou E V., Parry DM. Predictors of vestibular schwannoma growth in patients with neurofibromatosis Type 2. J Neurosurg. 2002 Feb;96(2):217–22.

8. Forde C, King AT, Rutherford SA, Hammerbeck-Ward C, Lloyd SK, Freeman SR, et al. Disease course of neurofibromatosis type 2: A 30-year follow-up study of 353 patients seen at a single institution. Neuro Oncol. 2021 Jul 1;23(7):1113–24.

9.   Ragge NK, Baser ME, Klein J, Nechiporuk A, Sainz J, Pulst SM, et al. Ocular abnormalities

in neurofibromatosis 2. Am J Ophthalmol [Internet]. 1995;120(5):634–41. Available

from: http://dx.doi.org/10.1016/S0002-9394(14)72210-X

10.   Plana-Pla A, Bielsa-Marsol I, Carrato-Moñino C. Diagnostic and Prognostic Relevance of

the Cutaneous Manifestations of Neurofibromatosis Type 2. Actas Dermosifiliogr. 2017

Sep 1;108(7):630–6.

11.   Hexter A, Jones A, Joe H, Heap L, Smith MJ, Wallace AJ, et al. Clinical and molecular

predictors of mortality in neurofibromatosis 2: A UK national analysis of 1192 patients. J

Med Genet. 2015 Aug 14;52(10):699–705.

12.   Plotkin SR, Duda DG, Muzikansky A, Allen J, Blakeley J, Rosser T, et al. Multicenter,

prospective, phase II and biomarker study of high-dose bevacizumab as induction

therapy in patients with neurofibromatosis type 2 and progressive vestibular

schwannoma. Journal of Clinical Oncology. 2019;37(35):3446–54.

13.   McClatchey AI, Giovannini M. Membrane organization and tumorigenesis - The NF2

tumor suppressor, Merlin. Genes Dev. 2005;19(19):2265–77.

14.   Gladden AB, Hebert AM, Schneeberger EE, McClatchey AI. The NF2 Tumor Suppressor,

Merlin, Regulates Epidermal Development through the Establishment of a Junctional

Polarity Complex. Dev Cell [Internet]. 2010;19(5):727–39. Available from:

http://dx.doi.org/10.1016/j.devcel.2010.10.008

15.   Petrilli AM, Merlin V, Nf I, Merlin I, Nf O. Role of Merlin / NF2 inactivation in tumor

biology. 2016;(December 2014):537–48.

16.   Petrilli AM, Fernández-Valle C. Role of Merlin/NF2 inactivation in tumor biology.

Oncogene. 2015 Feb;35(February):537–48.

17.   Chen H, Xue L, Huang H, Wang H, Zhang X, Zhu W, et al. Synergistic effect of Nutlin-3

553   combined with MG-132 on schwannoma cells through restoration of merlin and p53

554   tumour suppressors. EBioMedicine. 2018 Oct 1;36:252–65.

555   18.   Zhou L, Hanemann CO. Merlin, a multi-suppressor from cell membrane to the nucleus.

556   Vol. 586, FEBS Letters. 2012. p. 1403–8.

557   19.   McClatchey AI, Saotome I, Ramesh V, Gusella JF, Jacks T. The Nf2 tumor suppressor

558   gene product is essential for extraembryonic development immediately prior to

559   gastrulation. Genes Dev. 1997;11(10):1253–65.

560   20.   McClatchey AI, Saotome I, Mercer K, Crowley D, Gusella JF, Bronson RT, et al. Mice

561   heterozygous for a mutation at the Nf2 tumor suppressor locus develop a range of

562   highly metastatic tumors. Genes Dev. 1998;12(8):1121–33.

563   21.   Schulz A, Büttner R, Hagel C, Baader SL, Kluwe L, Salamon J, et al. The importance of

564   nerve microenvironment for schwannoma development. Acta Neuropathol. 2016 Aug

565   1;132(2):289–307.

566   22.   Stemmer-Rachamimov AO, Louis DN, Nielsen GP, Antonescu CR, Borowsky AD, Bronson

567   RT, et al. Comparative pathology of nerve sheath tumors in mouse models and humans.

568   Cancer Res. 2004;64(10):3718–24.

569   23.   Giovannini M, Robanus-Maandag E, Van Der Valk M, Niwa-Kawakita M, Abramowski V,

570   Goutebroze L, et al. Conditional biallelic Nf2 mutation in the mouse promotes

571   manifestations of human neurofibromatosis type 2 [Internet]. 2000. Available from:

572   www.genesdev.org

573   24.   Ren Y, Chari DA, Vasilijic S, Welling DB, Stankovic KM. New developments in

574   neurofibromatosis type 2 and vestibular schwannoma. Neurooncol Adv. 2021;3(1):1–

575   13.

25. Dilwali S, Patel PB, Roberts DS, Basinsky GM, Harris GJ, Emerick KS, et al. Primary

culture of human Schwann and schwannoma cells: Improved and simplified protocol.

Hear Res. 2014;315:25–33.

26. Rosenbaum C, Kluwe L, Mautner VF, Friedrich RE, Mü HW, Hanemann CO. Isolation and

Characterization of Schwann Cells from Neurofibromatosis Type 2 Patients. 1998.

27. Xue L, He W, Wang Z, Chen H, Wang Z, Wu H. Characterization of a newly established

schwannoma cell line from a sporadic vestibular schwannoma patient [Internet]. Vol.

13, Am J Transl Res. 2021. Available from: www.ajtr.org

28. Hung G, Rhim JS, Slattery W, Lim D. Establishment and characterization of a

schwannoma cell line from a patient with neurofibromatosis 2. Vol. 20, INTERNATIONAL

JOURNAL OF ONCOLOGY. 2002.

29. Takahashi K, Yamanaka S. Induction of Pluripotent Stem Cells from Mouse Embryonic

and Adult Fibroblast Cultures by Defined Factors. 2006;2:663–76.

30. Shi Y, Inoue H, Wu JC, Yamanaka S. Induced pluripotent stem cell technology: A decade

of progress. Nat Rev Drug Discov [Internet]. 2017;16(2):115–30. Available from:

http://dx.doi.org/10.1038/nrd.2016.245

31. Papapetrou EP. Patient-derived induced pluripotent stem cells in cancer research and

precision oncology. Vol. 22, Nature Medicine. Nature Publishing Group; 2016. p. 1392–

401.

32. Raya Á, Rodríguez-Piz I, Guenechea G, Vassena R, Navarro S, Barrero MJ, et al. Disease-

corrected haematopoietic progenitors from Fanconi anaemia induced pluripotent stem

cells. Nature. 2009;460(7251):53–9.

33.   Mungenast AE, Siegert S, Tsai LH. Modeling Alzheimer's disease with human induced

pluripotent stem (iPS) cells. Molecular and Cellular Neuroscience [Internet].

2016;73:13–31. Available from: http://dx.doi.org/10.1016/j.mcn.2015.11.010

34.   Nourbakhsh A, Gosstola NC, Fernandez-valle C, Dykxhoorn DM. Characterization of

UMi031-A-2 inducible pluripotent stem cell line with a neurofibromatosis type 2-

associated mutation. Stem Cell Res [Internet]. 2021;55(June):102474. Available from:

https://doi.org/10.1016/j.scr.2021.102474

35.   Ishi Y, Era T, Yuzawa S, Okamoto M, Sawaya R, Motegi H, et al. Analysis of induced

pluripotent stem cell clones derived from a patient with mosaic neurofibromatosis

type 2. Vol. 188, American journal of medical genetics. Part A. United States; 2022. p.

1863–7.

36.   Jessen KR, Mirsky R. The origin and development of glial cells in peripheral nerves. Nat

Rev Neurosci. 2005;6(9):671–82.

37.   Menendez L, Kulik MJ, Page AT, Park SS, Lauderdale JD, Cunningham ML, et al. Directed

differentiation of human pluripotent cells to neural crest stem cells. Nat Protoc. 2013

Jan;8(1):203–12.

38.   Carrió M, Mazuelas H, Richaud-Patin Y, Gel B, Terribas E, Rosas I, et al. Reprogramming

Captures the Genetic and Tumorigenic Properties of Neurofibromatosis Type 1

Plexiform Neurofibromas. Stem Cell Reports. 2019;12.

39.   Baser ME, Friedman JM, Wallace AJ, Ramsden RT, Joe H, Evans DGR. Evaluation of

clinical diagnostic criteria for neurofibromatosis 2. Neurology. 2002 Dec;59(11):1759–

65.

40.   Castellanos E, Plana A, Carrato C, Carri M, Rosas I, Amilibia E, et al. Early genetic

621   diagnosis of neurofibromatosis type 2 from skin plaque plexiform schwannomas in

622   childhood. JAMA Dermatol. 2018 Mar 1;154(3):341–6.

623   41.   Castellanos E, Gel B, Rosas I, Tornero E, Santín S, Pluvinet R, et al. A comprehensive

624   custom panel design for routine hereditary cancer testing: Preserving control,

625   improving diagnostics and revealing a complex variation landscape. Sci Rep. 2017;7.

626   42.   Van der Auwera GA, O'Connor BD. Genomics in the Cloud: Using Docker, GATK, and

627   WDL   in   Terra   [Internet].   O'Reilly   Media;   2020.   Available   from:

628   https://books.google.es/books?id=vsXaDwAAQBAJ

629   43.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

630   universal RNA-seq aligner. Bioinformatics. 2013 Jan;29(1):15–21.

631   44.   Hartley SW, Mullikin JC. QoRTs: a comprehensive toolset for quality control and data

632   processing of RNA-Seq experiments. BMC Bioinformatics. 2015 Jul;16(1):224.

633   45.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential

634   expression analysis of digital gene expression data. Bioinformatics. 2010 Jan;26(1):139–

635   40.

636   46.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

637   RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

638   47.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene

639   set enrichment analysis: A knowledge-based approach for interpreting genome-wide

640   expression   profiles   [Internet].   2005.   Available   from:

641   www.pnas.orgcgidoi10.1073pnas.0506580102

642   48.   Barbie DA, Tamayo P, Boehm JS, Kim SY, Moody SE, Dunn IF, et al. Systematic RNA

643   interference reveals that oncogenic KRAS-driven cancers require TBK1. Nature. 2009

644   Nov 5;462(7269):108–12.

645   49.   Kuleshov M V., Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. Enrichr: a

646   comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids

647   Res. 2016 Jul 8;44(1):W90–7.

648   50.   Prakash YS, Singh A, Shannon CP, Kim YW, Demarco ML, Gauvreau GM, et al. Identifying

649   Molecular Mechanisms of the Late-Phase Asthmatic Response by Integrating Cellular,

650   Gene, and Metabolite Levels in Blood Mucins and Their Sugars Critical Mediators of

651   Hyperreactivity and Inflammation [Internet]. Vol. 13, Ann Am Thorac Soc. 2016.

652   Available from: www.atsjournals.org.www.atsjournals.org.

653   51.   Mazuelas H, Magallón-Lorenz M, Fernández-Rodríguez J, Uriarte-Arrazola I, Richaud-

654   Patin Y, Terribas E, et al. Modeling iPSC-derived human neurofibroma-like tumors in

655   mice uncovers the heterogeneity of Schwann cells within plexiform neurofibromas. Cell

656   Rep   [Internet].   2022   Feb;38(7):110385.   Available   from:

657   https://linkinghub.elsevier.com/retrieve/pii/S2211124722001061

658   52.   Chen J, Jiang C, Fu L, Zhu CL, Xiang YQ, Jiang LX, et al. CHL1 suppresses tumor growth

659   and metastasis in nasopharyngeal carcinoma by repressing PI3K/AKT signaling pathway

660   via interaction with integrin β1 and merlin. Int J Biol Sci. 2019;15(9):1802–15.

661   53.   Agnihotri S, Jalali S, Wilson MR, Danesh A, Li M, Klironomos G, et al. The genomic

662   landscape of schwannoma. Nat Genet. 2016 Nov 1;48(11):1339–48.

663   54.   Yilmaz A, Peretz M, Aharony A, Sagi I, Benvenisty N. Defining essential genes for human

664   pluripotent stem cells by CRISPR-Cas9 screening in haploid cells. Nat Cell Biol [Internet].

665   2018;20(5):610–9. Available from: http://dx.doi.org/10.1038/s41556-018-0088-1

666 55. Blomen VA, Májek P, Jae LT, Bigenzahn JW, Nieuwenhuis J, Staring J, et al. Gene
667   essentiality and synthetic lethality in haploid human cells. Science (1979).
668   2015;350(6264):1092–6.

669 56. Rowe RG, Daley GQ. Induced pluripotent stem cells in disease modelling and drug
670   discovery. Vol. 20, Nature Reviews Genetics. Nature Publishing Group; 2019. p. 377–88.

671 57. Curto M, McClatchey AI. Nf2/Merlin: A coordinator of receptor signalling and
672   intercellular contact. Br J Cancer. 2008;98(2):256–62.

673 58. Curto M, Cole BK, Lallemand D, Liu CH, McClatchey AI. Contact-dependent inhibition of
674   EGFR signaling by Nf2/Merlin. Journal of Cell Biology. 2007 Jun 4;177(5):893–903.

675 59. Catasús N, Rosas I, Bonache S, Negro A, Torres-Martin M, Plana-Pla A, et al. Antisense
676   oligonucleotides targeting exon 11 are able to partially rescue the NF2-related
677   schwannomatosis phenotype in vitro. Mol Ther Nucleic Acids [Internet].
678   2022;30(December):493–505.    Available    from:
679   https://doi.org/10.1016/j.omtn.2022.10.026

680 60. Wang Y, Xu X, Maglic D, Dill MT, Mojumdar K, Ng PKS, et al. Comprehensive Molecular
681   Characterization of the Hippo Signaling Pathway in Cancer. Cell Rep. 2018 Oct
682   30;25(5):1304-1317.e5.

683 61. James MF, Han S, Polizzano C, Plotkin SR, Manning BD, Stemmer-Rachamimov AO, et al.
684   NF2/Merlin Is a Novel Negative Regulator of mTOR Complex 1, and Activation of
685   mTORC1 Is Associated with Meningioma and Schwannoma Growth. Mol Cell Biol. 2009
686   Aug;29(15):4250–61.

687 62. Giovannini M, Bonne NX, Vitte J, Chareyre F, Tanaka K, Adams R, et al. MTORC1
688   inhibition delays growth of neurofibromatosis type 2 schwannoma. Neuro Oncol.
689   2014;16(4):493–504.

63. Chiasson-MacKenzie C, Vitte J, Liu CH, Wright EA, Flynn EA, Stott SL, et al. Cellular

mechanisms of heterogeneity in NF2-mutant schwannoma. Nat Commun [Internet].

2023 Mar 21;14(1):1559. Available from: https://www.nature.com/articles/s41467-

023-37226-0

64. Shi Y, Bollam SR, White SM, Laughlin SZ, Graham GT, Wadhwa M, et al. Rac1-Mediated

DNA Damage and Inflammation Promote Nf2 Tumorigenesis but Also Limit Cell-Cycle

Progression. Dev Cell. 2016 Nov 21;39(4):452–65.

65. Kaempchen K, Mielke K, Utermark T, Langmesser S, Hanemann CO. Upregulation of the

Rac1/JNK signaling pathway in primary human schwannoma cells. Hum Mol Genet.

2003 Jun 1;12(11):1211–21.

66. Lallemand D, Manent J, Couvelard A, Watilliaux A, Siena M, Chareyre F, et al. Merlin

regulates transmembrane receptor accumulation and signaling at the plasma

membrane in primary mouse Schwann cells and in human schwannomas. Oncogene.

2009;28(6):854–65.

67. Scoles DR, Nguyen VD, Qin Y, Sun CX, Morrison H, Gutmann DH, et al.

Neurofibromatosis 2 (NF2) tumor suppressor schwannomin and its interacting protein

HRS   regulate   STAT   signaling   [Internet].   Available   from:

https://academic.oup.com/hmg/article/11/25/3179/578782

68. Wallesch M, Pachow D, Blücher C, Firsching R, Warnke JP, Braunsdorf WEK, et al.

Altered expression of E-Cadherin-related transcription factors indicates partial

epithelial-mesenchymal transition in aggressive meningiomas. J Neurol Sci. 2017 Sep

15;380:112–21.

69.   Takahashi E, Haga A, Tanihara H. Merlin regulates epithelial-to-mesenchymal transition

713   of ARPE-19 cells via TAK1-p38MAPK-mediated activation. Invest Ophthalmol Vis Sci.

714   2015;56(4):2449–58.

715   70.   Li X, Chen H, Xue L, Pang X, Zhang X, Zhu Z, et al. P53 performs an essential role in

716   mediating the oncogenic stimulus triggered by loss of expression of neurofibromatosis

717   type 2 during in vitro tumor progression. Oncol Lett. 2017;14(2):2223–31.

718   71.   Beltrami S, Kim R, Gordon J. Neurofibromatosis Type 2 Protein, NF2: An uncoventional

719   cell cycle regulator.

720   72.   Toshiyuki Nakayama, Yang Cheul Cho, Yuka Mine, Ayumi Yoshizaki, Shinji Naito, Chun

721   Yang Wen, et al. WJG-12-6182 (1). World J Gastroenterol. 2006;38(12):6182–7.

Table 1. iPSCs lines information

		Reprogramming VS		Editing iPSCs from VS		Editing FiPS Ctrl 1-SV4F-7	
Generation method							
iPS Cell line		VSi-25	VSi-267	VSi-25-CasD7	VSi-267-CasD2	FiPS-CasB2	FiPS-CasH6
NF2 Genotype	*Allele A* *Allele B*	c.1736A>G WT	c.784C>T WT	c.1736A>G c.211dup	c.784C>T c. 212_213del	c.212_213delCA WT	c.211dup c.212_213del
	Protein	p.(Lys579Arg)	p.(Arg262*)	p.(Lys579Arg) & p.(Thr71Asnfs*15)	p.(Arg262*) & p.(Thr71Serfs*14)	p.(Thr71Serfs*14)	p.(Thr71Asnfs*15) & p.(Thr71Serfs*14)
		NF2 (+/-)	*NF2* (+/-)	*NF2* (-/-)	*NF2* (-/-)	*NF2* (+/-)	*NF2* (-/-)
Effect **Affected exons**		Splicing 15	Nonsense 8	Splicing and frameshift 2 and 15	Nonsense and frameshift 2 and 8	Frameshift 2	Frameshift 2

A

Source
- 25
- FiPS
- 267
- Schwannomas

Cell Type
- Schwannomas
- SC-like Spheroid

Genotype
- NF2(+/-)
- NF2(+/+)
- NF2(-/-)

Color scale: 4 – 14

NF2(+/+)
NF2(+/-)   Schwannomas   NF2(-/-)

B

Biological Process

- Positive Regulation Of Cell Motility (GO:2000147)
- Regulation Of Cell Migration (GO:0030334)
- Negative Regulation Of Platelet Activation (GO:0010544)
- Regulation Of Plasminogen Activation (GO:0010755)
- Extracellular Structure Organization (GO:0043062)
- External Encapsulating Structure Organization (GO:0045229)
- Negative Regulation Of Plasminogen Activation (GO:0010757)
- Regulation Of Wound Healing (GO:0061041)
- Extracellular Matrix Organization (GO:0030198)
- Negative Regulation Of Angiogenesis (GO:0016525)

$-\log_{10}$(Adjusted P-value)

KEGG

- Focal adhesion
- PI3K-Akt signaling pathway
- Bladder cancer
- ECM-receptor interaction
- MAPK signaling pathway
- Primary immunodeficiency
- Fluid shear stress and atherosclerosis
- Glioma
- p53 signaling pathway
- Melanoma

$-\log_{10}$(Adjusted P-value)

C

VS
NF2 patient

Vestibular Schwannoma

NF2(+/-) cells
NF2(-/-) SC

NF2(+/+) cells
FiPS

Gene editing → NF2(-/-) iPSC

Reprogramming → NF2(+/-) iPSC

Gene editing

NF2(-/-) iPSC
SeV expression
(Discarded)

NC Differentiation → NF2(-/-) NC

NF2(+/-) NC

SC Differentiation in 3D → NF2(-/-) spheroid

NF2(+/-) spheroid