MERLIN-DEFICIENT iPSCs SHOW ALTERED PLURIPOTENCY AND CONSTITUTE A POTENTIAL *IN*

1

2   ***VITRO* MODEL FOR NF2-RELATED SCHWANNOMAS**

3

4   Núria Catasús[1], Miguel Torres-Martin[1,2], Alex Negro[1,2], Bernd Kuebler[3], Inma Rosas[1,2], Gemma

5   Casals[1], Helena Mazuelas[4], Francesc Roca-Ribas[5], Emilio Amilibia[5], Begoña Aran[3], Anna Veiga[3],

6   Ángel Raya[3], Bernat Gel[4], Ignacio Blanco[1,2], Eduard Serra[4*], Meritxell Carrió[4*] and Elisabeth

7   Castellanos[1,2*]

8   *Equally contributed

9

10   1)   Clinical Genomics Research Group, Germans Trias i Pujol Research Institute (IGTP); Can

11   Ruti Campus, Badalona, Barcelona, Spain

12   2)   Clinical Genomics Unit, Germans Trias i Pujol University Hospital (HUGTiP), Can Ruti

13   Campus, Badalona, Barcelona, Spain

14   3)   Medicine Regenerative Program, Institut d'Investigació Biomèdica de Bellvitge. IDIBELL,

15   Hospital Duran i Reynals, Gran Via de L'Hospitalet, 199-203, 08908, L'Hospitalet de

16   Llobregat, Barcelona, Spain.

17   4)   Hereditary Cancer Group, Germans Trias i Pujol Research Institute (IGTP); Can Ruti

18   Campus, Badalona, Barcelona, Spain

19   5)   Otorhinolaryngology Department, Germans Trias i Pujol University Hospital (HUGTiP), Can

20   Ruti Campus, Badalona, Barcelona, Spain

21

22

23   **Corresponding author:** Eduard Serra (eserra@igtp.cat); Meritxell Carrió (mcarriol@igtp.cat);

24   Elisabeth Castellanos (ecastellanos@igtp.cat/ecastellanosp.germanstrias@gencat.cat)

25

26

Abstract

Background: The appearance of bilateral vestibular schwannomas (VS) is one of the most

characteristic features of NF2-related schwannomatosis (NF2-related SWN), an autosomal

dominant syndrome that predisposes to the development of tumours of the nervous system.

VS are caused by the bi-allelic inactivation of the NF2 gene in a cell of the Schwann cell lineage.

Our current understanding of VS initiation and progression as well as the development of new

effective therapies is hampered by the absence of human non-perishable cell-based models.

Principal Findings: We generated and characterized induced pluripotent stem cell (iPSC) lines

with single or bi-allelic inactivation of *NF2* by combining the direct reprogramming of VS cells

with the use of CRISPR/Cas9 editing. Despite the difficulty of maintaining merlin-deficient

iPSCs, we were able to differentiate them into neural crest (NC) cells. At this stage, these cells

showed

spontaneous expression of the SC marker S100B and the impossibility of generating Schwann

cells in 2D cultures. Nevertheless, by applying a 3D Schwann cell differentiation protocol, we

successfully generated *NF2*(+/-) and *NF2*(-/-) spheroids homogeneously expressing classical

markers of the NC-SC axis.

Conclusions: Our results show a critical function of *NF2* for both reprograming and maintaining

a stable pluripotent state. In addition, merlin-deficient cultures also denoted an altered

differentiation capacity of merlin-deficient cells towards the NC-SC axis, in the *in vitro*

conditions used. Finally, the generated *NF2*(+/-) and *NF2*(-/-) spheroids show potential as a

genuine in vitro model of NF2-related tumours.

Keywords: Neurofibromatosis Type 2; iPSC; Schwann Cell; Neural crest stem cell; Vestibular

Schwannomas, Schwannomas

Introduction

NF2-related schwannomatosis (NF2-related SWN; OMIM 101000) is a dominantly inherited autosomal syndrome characterized by the development of multiple tumours of the nervous system (1,2). This condition is caused by pathogenic variants in the *NF2* tumour suppressor gene (22q12.2), which codes for merlin (moesin-ezrin-radixin-like protein); a scaffold protein that interacts with transmembrane receptors and intracellular effectors to mediate mitogenic, survival and cell-cell contact signalling pathways. Changes in merlin result in the inhibition of cell proliferation (3–5). The incidence of the disease is estimated at 1:28.000 live births with a complete penetrance by the age of 60, with 50–60% of cases being due to *de novo* mutations (6). The pathognomonic feature of the disease is the development of bilateral vestibular schwannomas (VS), well-circumscribed intracranial tumours attached to the vestibulocochlear nerve that arise from Schwann Cells (SC) as a result of the bi-allelic *NF2* inactivation. Affected individuals may also develop multiple schwannomas in other cranial, spinal and peripheral nerves as well as intracranial and intraspinal meningiomas, and spinal ependymomas (7–10). Cutaneous and ocular manifestations are also frequent (11,12). Although histologically benign, these tumours are responsible for high morbidity and a reduced life expectancy due to their number and anatomical location (13).

VS naturally grow over time (9,14–16) and are responsible for significant morbidity associated with NF2-related SWN, which includes hearing loss, deafness and tinnitus that can eventually progress to facial paraesthesia, ataxia and vertigo (17). The gold standard for treatment of VS include surveillance, surgery, radiotherapy and Bevacizumab (17). The latter, an anti-VEGF monoclonal antibody, that has been shown to induce reduction of tumour growth and an improvement in hearing loss in NF2-related SWN patients, although with associated toxicity when sustained over time (18). However, the remaining therapeutic options also present different adverse effects, for example, the resection of VS often results in hearing loss, patients then requiring cochlear or brainstem implants to preserve hearing whereas,

radiotherapy of these tumours has been shown to induce tumour malignancy over time

(19,20). For the reasons above, efforts have been made to develop alternative therapies for

NF2-related tumours, such as Losartan (antihypertensive), Lapatinib (tyrosine kinase inhibitor)

or AR-42 (Histone deacetylase (HDAC) inhibitor) (21–24). However, since the disease course for

these tumours can be extremely long, there is a need to develop novel treatments with

minimal side effects.

The study of the role of *NF2* in VS development and progression, and the development of

potential therapies, is still challenging due to the lack of non-perishable preclinical models that

reproduce patient genetics and the pathophysiology of NF2-related tumours. The first

attempts to generate *in vivo* murine models revealed the essentiality of *NF2* during

embryogenesis and early development (19). While homozygous *NF2* mutant mice fail to

develop vital extraembryonic structures, which impedes them from reaching a gastrulation

stage, mosaic *NF2*(-/-) embryos show defects in a variety of other embryonic tissues (19). In

NF2(+/-) mutant mice, the deficiency of a single *NF2* allele causes the appearance of malignant

epithelial tumours and sarcomas with an aggressive tendency to metastasize (25). (25).

Conditional KO murine models can generate schwannomas and, in some cases VS, although

histologically they differ from human schwannomas (26,27). In addition, several xenograft

mouse models have also been developed, with different levels of success in recapitulating the

biology of human NF2-related SWN traits (28).

In vitro, primary SC cultures derived from VS have been established and demonstrated to

preserve the pathophysiology of the tumour (29,30). However, the limited lifespan of primary

cultures can severely constrain their use for extended experiments, or when large numbers of

cells are required. Transformed and immortalized VS derived cell lines have been developed in

order to overcome this limitation, and have been widely used as platforms for drug screening

(HEI-93, JEI-001) (31,32), although these models may undergo physiological changes due to the

immortalisation process, severely reducing the value of any comparison with primary cells.

Induced pluripotent stem cell (iPSC) (33) models have been used in a wide range of diseases,

representing an unlimited source of cells with diverse applications such as disease modelling,

drug development, regenerative medicine or gene regulation (34–37). The first related model

reported for NF2-related SWN was an iPSC line harbouring a homozygous variant in the *NF2*

gene (UMi031-A-2), obtained through gene editing by CRISPR/Cas9 of a non-affected

pluripotent cell line (38); and more recently, an iPSC clone has been derived from mononuclear

bone marrow-derived stem cells of a mosaic NF2-related SWN patient (39). However, there is

still the need to develop relevant disease models from these iPSC, which fully recapitulate the

genetic and physiological status of *NF2* derived VS.

One of the main defining characteristics of iPSC is that they have the ability to differentiate to

the three germ layers. Protocols have been developed to differentiate iPSC towards the SC

lineage, passing through an intermediate multipotent neural crest stage (40–42). It has been

shown that when applying these protocols, iPSC-derived SCs show homogenous expression of

SC markers and have the capacity to myelinate axons when they are co-cultured with dorsal

root ganglia neurons (42). Therefore, the use of iPSC seems to represent a suitable human SC-

based cellular model to study the molecular pathogenesis of NF2-related schwannomas.

In the present study we have generated iPSC lines harbouring single or bi-allelic loss of

function variants in the *NF2* gene by direct reprogramming of VS cells and the use of

CRISPR/Cas9 editing. These cells were characterized and differentiated towards the NC-SC

lineage, obtaining *NF2*(+/-) and *NF2*(-/-) spheroids that co-express classical markers of the NC-

SC axis, thereby potentially representing a bona fide NF2-related schwannoma model.

Results

1.　**Generation of induced pluripotent stem cells harbouring single or bi-allelic mutations**

in the *NF2* gene

Three VS from NF2-related SWN patients that met standard diagnostic criteria (43) were used

for reprogramming (VS-25, VS-245 and VS-267). Molecular characterization of these tumours

through the I2HCP panel (44) uncovered the *NF2* alterations reported in Supplemental Table

S1. In addition, no pathogenic variants were identified in known oncogenes or tumour

suppressor genes, as has been previously reported for VS. SNP-array analysis showed loss of

heterozygosity (LOH) on chromosome 22q in VS-25, and no complex rearrangements were

observed in any of the tumours analysed (Supplemental Figure S1).

The non-integrating Sendai virus vector (SeV) encoding the 4 Yamanaka factors (33) was used

to induce pluripotency of VS-dissociated cells, with the aim of generating iPSC *NF2*(-/-) clones

that could recapitulate the genetic background of Schwann cells involved in VS tumorigenesis.

After VS reprogramming, we determined the *NF2* genotype through Sanger sequencing of 38

single-cell clones for VS-25, 12 for VS-267 and 10 for VS-245 (Supplemental Table S2). From VS-

25 and VS-267, only clones carrying the germline *NF2* pathogenic variant, *NF2*(+/-), were

obtained. One of these clones derived from each tumour was selected for further

characterization. From VS-245, a tumour coming from a mosaic patient, we obtained both

NF2(-/-) and *NF2*(+/+) iPSC clones. As VSi-245(+/+) clones were reprogrammed from an

unaffected cell, these were not further characterized. We focused on characterizing *NF2*(-/-)

clones that were already distinctive in culture due to their aberrant morphology, containing

cells that were piling up on top of others (Supplemental Figure S2). Surprisingly, all VSi-245

NF2(-/-) iPSC clones (n=9), proved to be dependent on the exogenous expression of Yamanaka

factors by the Sendai (SeV) construct. Although SeV is a non-integrating RNA virus widely used

for reprogramming, the expression of which is commonly lost due to dilution of the viral

genome, *NF2*(-/-) clones could only be established directly from VS in the presence of SeV, this

remained true 40 passages after reprogramming, or after submitting the iPSC clones to 39°C

for several passages to achieve complete clearance of the virus. These clones showed

sustained expression of reprogramming factors (Supplemental Figure S2). Therefore, we

concluded that these clones could not constitute a reliable model and were discarded for the

Table 1. iPSCs lines information

rest of the study.

Genomic characterization of *NF2*(+/-) VSi-25 and VSi-267 iPSCs selected clones by SNP-array

analysis and WES showed no differences with respect to the tumour of origin, with a logical

exception for VSi-25, in which the LOH on chromosome 22q as the second hit was not present

in the (+/-) clones obtained from this tumour (Supplemental Figure S1).

With the aim of achieving iPSCs with complete *NF2* inactivation, and given the low efficiencies

obtained and the SeV dependency in the generation of *NF2*(-/-) clones through

reprogramming, we decided to use CRISPR/Cas9 technology to edit the *NF2* gene in both, the

NF2(+/-) clones obtained from reprogrammed VS cells and a control *NF2*(+/+) iPSC line (FiPS

Ctrl 1-SV4F-7, registered in the Spanish National Stem Cell Bank/ ESiO44C in

https://hpscreg.eu/)). Thus, using CRISPR/Cas9 RNP complexes, we targeted exon 2 of the *NF2*

gene and obtained several single-cell *NF2*(-/-) iPSC clones from VS-derived lines (named 25-

CasD7, when derived from VSi-25, and 267-CasD2 when derived from VSi-267) and isogenic

NF2(+/-) and *NF2*(-/-) iPSC clones derived from the control line (named FiPS-CasB2 and FiPS-

CasH6, respectively). Pathogenic variants in *NF2* were fully characterized through PCR and

Sanger sequencing and for each of them, the *NF2* coding region was cloned to confirm

compound heterozygous clones (Table 1). In addition, WES analysis of the edited lines revealed

no pathogenic off-target alterations (Supplemental Table S3).

From this point on, we worked on the characterization of six iPSC lines: three with one

pathogenic variant in *NF2* (two VS-reprogrammed and one control iPSC-edited) and three

carrying the bi-allelic inactivation of *NF2* (Table 1).

| Generation method | Reprogramming VS | | Editing iPSCs from VS | | Editing FiPS Ctrl 1-SV4F-7 | |
iPS Cell line	VSi-25	VSi-267	VSi-25-CasD7	VSi-267-CasD2	FiPS-CasB2	FiPS-CasH6
NF2 Genotype	c.1736A>G	c.784C>T	Allele A: c.211dupA	c.212_213delCA	c.212_213delCA	c.211dupA
	WT *NF2* (+/-)	WT *NF2* (+/-)	Allele B: c.212_213delCA and c.1736A>G *NF2* (-/-)	c.784C>T *NF2* (-/-)	WT *NF2* (+/-)	c.212_213delCA *NF2* (-/-)
p.	p.(Lys529Asnfs*19)	p.(Arg262*)	Allele A: p.(Thr71Asnfs*15) Allele B: p.(Thr71Serfs*14) p.(Lys529Asnfs*19)	p.(Thr71Serfs*14) p.(Arg262*)	p.(Thr71Serfs*14)	p.(Thr71Asnfs*15) p.(Thr71Serfs*14)
Effect	Splicing	Truncating	Splicing and truncating	Truncating	Truncating	Truncating
	NF2 (+/-)	*NF2* (+/-)	*NF2* (-/-)	*NF2* (-/-)	*NF2* (+/-)	*NF2* (-/-)
Affected exons	15	8	2 and 15	2 and 8	2	2

iPSCs: induced pluripotent stem cell; VS: vestibular schwannoma; FiPS; fibroblasts induced pluripotent stem cell; Ctrl: Control; WT: wild-type

2. *NF2* is necessary for maintaining a stable pluripotent state

The combination of strategies between VS reprogramming and *NF2* gene editing allowed us to obtain clones with single (+/-) or double (-/-) mutations in the *NF2* gene (Figure 1A). All of them expressed cell surface proteins and transcription factors associated with pluripotency: Oct4, SSEA-3, SSEA-4, Nanog, Sox2, Tra-1-60 and Tra-1-81; were positive for alkaline phosphatase staining; and showed karyotype stability after at least 20 passages (46, XY) (Supplemental Figure S3). In addition to the constitutional and somatic *NF2* mutation characterization (Table 1), we confirmed all *NF2* genotypes by performing western blot of merlin (Figure 1B). *NF2*(+/-) cell lines exhibited an approximately 50% reduction in merlin expression compared to the control and *NF2*(-/-) iPSCs showed no expression of the protein

(Figure 1B). Although all iPSC clones expressed the typical pluripotent markers, we could

observe differences in clone morphology between *NF2*(+/-) and *NF2*(-/-) cell lines. Whereas

NF2(+/-) showed classical iPSC colony morphology, edited *NF2*(-/-) iPSC colonies presented less

compact colony formation, eventually aggregating in the centre of the colony and growing

upwards and undergoing major spontaneous differentiation compared with the iPSC control

line (Figure 1C), as we had previously observed for the reprogrammed *NF2*(-/-) iPSC line.

Notably, the 267-CasD2 line required removal of visibly differentiated cells before each

passage. Furthermore, the three *NF2*(+/-) iPSC lines showed the capacity to differentiate into

the three primary germ layers *in vitro* (mesoderm, endoderm and ectoderm) through

embryoid body (EBs) formation (Figure 1D). In contrast, *NF2*(-/-) clones showed a unique

response. For the FiPS-CasH6 line, only EBs thrived under conditions of differentiation to

ectoderm and stained positive for TUJ1 and GFAP markers, but they did not achieve

differentiation to endoderm. In addition, differentiation through EBs did not succeed for the

25-CasD7 and 267-CasD2 lines applying standard protocols. After attempting differentiation

through EBs formation, direct differentiation protocols were used. Positive staining for some

markers of the three primary germ layers was achieved, although the morphology of iPSCs and

their derivatives was abnormal (Figure 1D, Supplemental Table S4), indicating again that loss of

NF2 affects the pluripotency capacity of iPSCs. These observations, together with the data

previously compiled from VS cell reprogramming (low efficiency in the generation of *NF2*(-/-)

clones; dependency on SeV-expressed Yamanaka factors) and the phenotypic characteristics of

NF2(-/-) iPSC clones, suggest a critical role of the *NF2* gene in the maintenance of a

pluripotency state.

Figure 1. Absence of merlin in iPSCs results in an altered pluripotent phenotype. (A)

Schematic representation of the experimental procedure for obtaining merlin-null clones. (B)

Morphology of the cell colonies, Scale bar, 75µM; (C) Immunochemistry to demonstrate the

capacity of the lines to *in vitro* differentiate to the three primary germ layers of *NF2*(+/-) lines:

mesoderm (ASMA in green and ASA in red), ectoderm (TUJ1 in green and GFAP in red) and

endoderm (AFP in green and FOXA2 in red). For NF2(-/-), a directed differentiation was

performed due to the inability of these lines to generate EBs, for these: mesoderm (ASMA in

green and GATA4 in red), ectoderm (TUJ1 in green, GFAP in red and PAX6 in pink) and

endoderm (AFP in green, FOXA2 and SOX17 in red). Scale bar, 75μM (D) Merlin expression

analysed by Western Blot. The FiPS line generated from fibroblasts (NF2 +/+) was used as a

control cell line, NPE stands for Normalized Protein Expression.

3.   Merlin-deficient iPSCs have an altered capacity to acquire and retain a stable neural

crest identity

The next step in the development of an iPSC-based VS model was their differentiation into

neural crest (NC) cells, previous to a final differentiation towards SCs. To generate NCs, we

applied chemically defined differentiation conditions as previously reported [41], containing

activators of the Wnt signaling pathway and inhibitors of the TGFβ/activin/nodal pathway.

During the first ten days after the induction of NC differentiation, cell morphology shifted and

achieved a NC morphology in all cell lines tested (Figure 2A). Cell identity was first assessed by

flow cytometry of NC markers, NGFR (p75) and HNK1 (CD57), at early passages of the NC

differentiation process (p4) and in a mature NC state (p8) (Figure 2B). In our hands, *NF2*(+/-)

NC cell lines, derived either from reprogrammed VS cells or edited from isogenic control (FiPS-

CasB2), showed around 90% of cells expressing both markers in a similar way to, but a bit less

efficiently than the *NF2*(+/+) control line. Distinctively, *NF2*(-/-) cell lines already showed some

expression of p75 and Hnk1 already at the iPSC stage, but did not acquired the same

percentages of p75/Hnk1 positive cells at the mature NC stage, with an average of 76,3% for

the three cell lines (Figure 2B).

NC identity was further characterized through immunocytochemistry and transcriptome

analysis (RNA-seq), showing that all cell lines, *NF2* (+/+), (+/-) and (-/-), expressed the NC

markers Sox10, TFAP2A and p75 and showed no expression of the pluripotency markers

POU5F1 (Oct4), Nanog and Sox2 (Figure 2C and 2D). However, some *NF2*(-/-) NC cells were also

expressing S100B both at the RNA and protein levels (Figure 2C and 2D), which was consistent

with the heterogeneity detected by flow cytometry assays. Differences in merlin-deficient NC

cells were also observed in culture conditions, as these cells required culturing at higher cell

confluence conditions, compared to *NF2*(+/+) or *NF2*(+/-), to maintain a NC cell morphology.

Despite these differences, all generated lines were able to maintain a self-renewal capacity for

at least 18 passages while preserving a NC identity, and could be cultured after several freeze-

thaw cycles.

In summary, *NF2*(+/-) and *NF2*(-/-) iPSC lines could be differentiated towards a NC identity.

However, generation of *NF2*(-/-) NC cultures was less efficient, required special care to

maintain a NC morphology, and the resulting *NF2*(-/-) NC population expressed both NC

markers but also the S100B SC marker, suggesting that *NF2* function is important to acquire

and retain a stable neural crest identity.

Figure 2. Neural Crest Characterization. (A) NC Morphology of *NF2*(+/-) and *NF2*(-/-) cell lines;

Scale bar, 75µM; (B) Flow cytometry assays of the NC markers NGFR (p75) and Hnk1. The

percentage of p75 and Hnk1-positive cells is shown in pink. P4 and P8 stand for passages 4 and

8, respectively; (C) Immunocytochemistry of AP2 (green), p75 (green) and S100B (red), Scale

bars, 25µM; Oct4 (green) and SOX10 (red), Scale bar, 100µM; DAPI (blue) was used to stain cell

nuclei. (D) Expression levels of the genes related to a pluripotent stage (Nanog and Sox2), a NC

stage (Sox10 and AP2) and a SC marker (S100). Y-axis represents normalized FPKM expression.

4.   **Schwann Cell Differentiation of *NF2*(+/-) and *NF2*(-/-) from derived-NC cultures**

A final step consisted in inducing the differentiation of the established NC cultures towards a

SC state. Thus, NCs were cultured in SC differentiation medium (SCDM) for 30 days, as

previously stablished [41]. During the differentiation process, significant differences were

observed between *NF2*(+/+) control and merlin-deficient cell lines: five days after inducing SC

differentiation, *NF2*(+/-) and *NF2*(-/-) cell lines showed difficulties in maintaining adherence to

culture dishes, in contrast to the control line, which after seven days already showed a more

elongated spindle morpohlogy resembling SC-like cells (Supplemental Figure S5). Expression

analysis indicated that *NF2*(+/+) NC cultures in 2D were able to differentiate towards a

Schwann cell lineage, as they expressed classical NC-SC lineage markers and also the NC-SC

roadmap gene signature described in Mazuelas et al (2022) (Supplemental Figure S6) (45). In

contrast, we reasoned that the poor adherence capacity of merlin-deficient NCs compromised

their ability to differentiate towards a SC identity. Consequently, we tested the SC

differentiation capacity of these cells in 3D cultures, following a previously established

protocol (45) with minor modifications. In brief, NC cells were seeded in uncoated spheroid

AggreWell plates for up to 30 days in SC differentiation medium. This procedure was

monitored at three time points (7, 14 and 30 days) through immunocytochemistry and RNA-

seq analysis (Figure 3). We used *NF2*(+/+) differentiating cells as an experimental control. At

the RNA level, we observed that *NF2*(+/+) differentiating spheroids expressed typical lineage

markers (S100B, MPZ or PLP1) already at day 7, which were maintained after 14 days under

SCDM. The expression of SC markers was not equally sustained up to 30-days of differentiation

(Supplementary Figure S7), indicating that this differentiation protocol was most appropriate

up to 14 days of differentiation. *NF2*(+/+) spheroids were also positive for p75 and S100B

markers identified by immunostaining (Figure 3A). Then, we differentiated *NF2* mutant cell

lines in these conditions. Both *NF2*(+/-) VSi-25 and VSi-267 cell lines formed spheroids that

stained positive for p75 and S100B markers, as well as the three *NF2*(-/-) cell lines (Figure 3A).

However, unexpectedly we were unable to obtain spheroids from the edited FiPS-CasB2

NF2(+/-) cell line (Supplemental Figure S8).

In order to determine the SC differentiation capacity of NF2(+/-) and (-/-) cell lines, we studied

the transcriptome of the whole differentiation process. First we performed a principal

component analysis (PCA) of all cell lines, including the iPSC and NC stages and throughout the

SC differentiation process (Figure 3B). Cells at different differentiation stages clustered

together, indicating that we were able to reliably distinguish between the three cell types. All

iPSC cells grouped together, although the edited NF2-deficient 267-CasD2 line positioned

slightly apart from the rest. Also, all SCs differentiating in 3D clustered together independently

of the genotype or days of differentiation (Figure 3B). Secondly, we studied the expression

levels of classical NC-SC lineage markers (46). NF2(+/-) and merlin-deficient cell lines,

expressed high levels of SC markers such as CDH19, GAP43, MPZ, PLP1 or S100B at 14 days of

differentiation, in a similar way to NF2(+/+) control lines (Figure 3C), indicating that NC cells

were progressing from a precursor state into a committed SC phenotype, probably without

reaching a mature myelinating phenotype. Surprisingly, merlin-deficient differentiating SCs

maintained the expression of the NC-SC markers up to 30 days in 3D (data not shown).

Furthermore, and given that the cells that would make up a schwannoma cell model would be

NF2(-/-) cells, we also analysed the in vitro NC-SC expression roadmap (45) (Figure 3D). NF2(-/-

) cell lines displayed a very similar NC-SC expression roadmap to that previously determined

for NF2(+/+) differentiating SCs in 2D (Figure 3D and Supplemental Figure S6) (45).

Figure 3. Schwann Cell Differentiation in 3D cultures (A) Immunochemistry of p75 (green) and

S100B (red) after 14 days of SC differentiation, Scale bar, 250µM. (B) PCA analysis (C) Gene

expression at day 14 of the 3D SC differentiation protocol (D) In vitro NC-SC expression

roadmap of the WT line (FiPS, NF2(+/+)) in the 2D differentiation protocol and the merlin-

deficient lines.

Overall, we successfully applied 3D SC differentiation conditions to *NF2*(+/-) and *NF2*(-/-) to

obtain spheroids expressing NC-SC lineage markers at RNA and protein level. iPSC-derived

merlin-deficient differentiating SCs constitute a potential reliable model for VS cells (Figure 4).

Figure 4. Schematic representation of the experimental process.

Discussion

The development of bilateral vestibular schwannomas (VS) constitute the most distinctive trait

of NF2-related Schwannomatosis (NF2) (7). VS arise from a cell of the Schwann Cell (SC) lineage

at the vestibular nerve. SCs from VS can be expanded as primary cultures. However, the small

number of passages that primary cells can achieve limits their use in experiments requiring

large numbers of cells. In this study we generated *NF2*(+/-) and *NF2*(-/-) iPSC lines by

combining direct reprogramming of VS cells and *NF2* gene editing. Specifically, by

reprogramming VS cells it was possible to generate two *NF2*(+/-) lines (VSi-25 and VSi-267)

derived from two VS (Table 1). Also, a third *NF2*(+/-) cell line (FiPS_CasB2) was generated by

CRISPR/Cas9 editing of a control iPSC (FiPS). In addition, by editing of VSi-25 and VSi-267, we

obtained two *NF2*(-/-) isogenic cell lines (25-CasD7, 267-CasD2). A third *NF2*(-/-) iPSC line (FiPS-

CasH6) was obtained by editing the control FiPS iPSC, also isogenic to *NF2*(+/-) FiPS_CasB2.

In contrast to the generation of iPSCs by reprogramming cells derived from other benign

peripheral nervous system tumors, like plexiform neurofibromas (42), we were not able to

generate genuine iPSCs by reprogramming *NF2*-deficient VS cells. The only *NF2*(-/-) cell line

obtained was dependent on the exogenous expression of Yamanaka factors. We hypothesize

that *NF2* function is required during the reprogramming process and might be related to its

essential role in embryogenesis and early stages of development (19), which is also consistent

with the reported essential function of *NF2* in haploid human iPSCs and leukaemia cells

(47,48). However, it is possible that using different conditions for the reprogramming of VS

cells and/or the maintenance of *NF2*(-/-) iPSCs could produce a different result, so additional

evidence is needed to reach a definitive conclusion.

Once the cells were in a pluripotent state, we were indeed able to generate *NF2*(-/-) iPSCs

through gene editing, as has been recently reported (38). However, in our hands, all three of

the *NF2*(-/-) iPSC lines generated required extensive experimental care; showed aberrant

colony morphology with a tendency to pile up cells during growth; and were susceptible to

spontaneous differentiation in a similar way to the *NF2*(-/-) colonies obtained by reprograming

VS cells. Therefore, considering that the origins and the genetic backgrounds of all three *NF2*(-

/-) cell lines were different and that the only thing they shared was complete *NF2* inactivation,

our results indicate that *NF2* function is responsible for this aberrant phenotype, suggesting it

also has an essential role in maintaining pluripotency of diploid cells. However, despite all

these limitations, the iPSC lines generated in this study constitute a valuable and unperishable

source of cells harbouring single or bi-allelic inactivation of *NF2,* that under adequate culture

conditions, could be used as a NF2-cell based model to develop further assays (50).

We showed that both *NF2*(+/-) and *NF2*(-/-) iPSCs have the capacity to differentiate towards a

NC identity, although *NF2*(-/-) NC cultures exhibited an unstable NC phenotype, with some

cells undergoing spontaneous expression of S100B. In addition, we highlight that *NF2*-deficient

iPSC-derived NCs required a higher confluence in culture, thus an apparently closer cell-cell

contact to maintain NC identity, consistent with the role of merlin in cytoskeleton

maintenance and control of proliferation by cell-cell contact inhibition (51–53). Also, in line

with these findings, in the process of differentiation of NCs towards SCs, we observed that

both *NF2*(+/-) and *NF2*(-/-) cells were easily detached from the cell culture surface, eventually

leading to cell death and preventing the generation of SC-like cultures.

However, and most importantly, we overcame this limitation by successfully generating

NF2(+/-) and NF2(-/-) spheroids growing in suspension as has been previously reported for

other SC-based spheroids (45). These spheroids already expressed classical NC-SC markers, as

shown by immunostaining, after seven days in SC differentiation medium. In addition,

extensive transcriptome analysis of *NF2*(+/-) and *NF2*(-/-) spheroids was performed, evidencing

the expression of well characterized NC-SC lineage markers (40). Furthermore, *NF2*(-/-) cells

also preserved a previously characterized *in vitro* NC-SC differentiation expression roadmap as

control iPSCs (45).

Finally, in this study we have generated *NF2*(+/-) and *NF2*(-/-) iPSC lines through

reprogramming VS and *NF2* editing. According to our results, the *NF2* gene seems to have a

relevant role in the induction and maintenance of pluripotency. iPSCs with different *NF2*

genotypes were differentiated into NC cells, although *merlin*-deficient iPSCs exhibited an

altered differentiation capacity. *NF2*(-/-) NC cultures showed heterogeneity in the expression

of classical NC markers with some cells spontaneously expressing non-NC markers such as

S100B. In addition, they were unable to generate SC-like cells in 2D cultures due to a deficient

adherence. However, the application of a 3D SC differentiation protocol resulted in the

successful generation of *NF2*(+/-) and *NF2*(-/-) spheroids expressing NC-SC lineage markers

(Figure 4). These spheroids constitute a valuable resource for the investigation of the biology

of NF2-related schwannomas, the study of the impact of *NF2* loss in the molecular

pathogenesis of NF2 traits and for the testing of new therapeutic approaches for *NF2*-related

tumors. In addition, despite the challenges in the use of *NF2*-deficient iPSCs, this resource

could be used to obtain other cell types involved in other NF2-related tumors, such as

meningioma- and ependymoma-originating cells.

Materials and Methods

All procedures performed were in accordance with the ethical standards of the IGTP

Institutional Review Board, which approved this study, and with the 1964 Helsinki declaration

and its later amendments. This project was authorised by the "Comisión de Garantías para la

Donación y Utilización de Células y Tejidos Humanos, ISCIII")

Patients and Vestibular Schwannoma samples: tumour samples were obtained from NF2-

related SWN patients clinically managed at the Spanish Reference Centre (CSUR) on

Phacomatoses HUGTiP-ICO-IGTP, diagnosed according to standard diagnostic criteria (43).

Tumour samples were processed as described in Castellanos et al., 2018 (54).

Reprogramming of Vestibular Schwannomas: between $0.2 \cdot 10^6$-$1 \cdot 10^6$ of VS derived cells were

reprogrammed using the CytoTune™-iPS 2.0 Sendai Reprogramming Kit (Thermo Fisher

Scientific), a non-integrative cell reprogramming method containing the four Yamanaka factors

(OCT4, SOX2, KLF4, and cMYC), according to manufacturer's instructions.

Tumour and iPSC molecular characterization: genetic analysis was performed using the

customized I2HCP panel as previously described (44) or Whole Exome Sequencing (WES). The

latter was performed using KAPA HyperCap technology with KAPA HyperExome Probes (Roche)

according to manufacturer's instructions and sequenced in a NextSeq instrument (Illumina).

Analysis of small nucleotide variants was performed with Mutect2 and annotated with

Funcotator (55). Potential SNV and small indels were detected among the original cell lines and

the paired CRISPR lines, GATK 4.2.6.1 was used for the whole process (55) (Supplemental

Material and Methods). In order to detect *NF2* splicing variants, tumours were studied both at

RNA and DNA level. *NF2* variants were validated by Sanger Sequencing. SNP-array analysis was

assessed using Illumina HumanOmniExpress v1 BeadChips (730,525 SNPs) according to

manufacturers' instructions and as described in Supplemental Material and Methods.

CRISPR/Cas9 gene edition in iPSC lines: CRISPR/Cas9 editing was performed with the ArciTect

ribonucleoprotein (RNP) system (STEMCELL Technologies). The designed sgRNA targeted exon

2 of the *NF2* gene (GTACACAATCAAGGACACAG). TransIT-X2® Dynamic Delivery System (Mirus)

was used for transfection according to manufacturers' instructions.

iPSC culture: iPSCs were grown on growth factor-reduced Matrigel (23µg/cm2) (BD

Biosciences)-coated 6-well plates and cultured in mTESR Plus medium (STEMCELL

Technologies). Splits were performed using Accutase (Merk) and cells were seeded with Rock

Inhibitor (1:1000) for 24h (STEMCELL Technologies). When required, visibly differentiated cells

were removed manually from the culture of *NF2*(-/-) cells. In this study, the *NF2*(+/+) FiPS line

(FiPS Ctrl 1-SV4F-7) was used for all experiments as a control.

Neural Crest (NC) differentiation: differentiation of iPSCs lines towards NC was performed as

previously described in Menendez et al. (2013) (41) with minor modifications (42). $9x10^4$ iPS

cells were seeded on matrigel-coated dishes in mTESR Plus medium. The following day, the

medium was replaced with Neural Crest Differentiation Media (Supplemental Material and

Methods) and was replaced every day. NC were maintained in this medium and split with

Accutase when necessary.

Differentiation towards SC in 2D: to stablish SC differentiation, $0.4x10^6$ NC cells/well were

plated onto 0.1mg/mL poly-L-lysine (Sigma) and 4µg/mL laminin (Gibco) and cultured in SC

differentiation media (SCDM) (Supplemental Material and Methods) as previously described

(42).

Differentiation towards SC in 3D: NC cells were detached with Accutase and, $2.25 \cdot 10^6$ NC

cells/well were seeded onto AggreWell TM800 24-well plates (Stem Cell Technologies) in 2mL

SCDM (described above). The medium was changed twice a week removing 1 mL and replacing

1 mL of fresh SCDM. On days 7, 14 and 30 spheroids were collected and processed for

subsequent analysis.

Flow cytometry: cells in PBS-0.1% BSA were incubated with p75 primary antibody and

subsequently incubated with Alexa Fluor 568 secondary antibody, and Hnk1 primary antibody

and detected with Alexa Fluor 488-conjugated secondary antibody. Antibodies were incubated

30 min on ice. Cells were analysed by flow cytometry using BD LSR Fortessa SORP and BD FACS

Diva 6.2 software.

Immunocytochemistry: cells or spheroids were fixed in 4% paraformaldehyde in PBS for 15min

at room temperature (RT), permeabilized with 0.1%Triton-X 100 in PBS for 10 min at RT,

blocked in 10% FBS in PBS for 15 min at RT, and incubated with the indicated antibodies

(Supplemental Table M1) overnight at 4°C. Secondary antibodies Alexa Fluor 488 and Alexa

Fluor 568 were incubated for 1h at RT. Nuclei were stained with DAPI and images captured

using LEICA DMIL6000 and LASAF software.

RNA analysis and heatmaps: For RNAseq analysis, genome GRCh37.p13 and gene annotation

gencode version 19 were used. For the alignment, STAR version 2.7.10a (56)) and QoRTs

version 1.3.6 (57) for gene count and general QC were used. For heatmaps, vst normalization

from DESeq2 (58) was used. For unsupervised cluster heatmaps, Euclidean metric and Ward

method were selected. For barplots in Figure 2, normalization was done by Trimmed Mean of

M-values from EdgeR (59) and then RPKMs were calculated for every gene.

Acknowledgements

We thank the HGTP Clinical Services and staff for their collaboration in generating and

collecting patient samples and data. We thank the IGTP Flow Cytometry core facility and their

staff for their contribution and technical support. We are indebted to all members of the

Cytogenetics Platform, Haematology Department at Germans Trias i Pujol Hospital, Catalan

Institute of Oncology and Josep Carreras Leukaemia Research Institute (Badalona, Spain) for

providing karyotype studies and images. We would like to acknowledge the constant support

of the different NF lay associations: *Asociación de Afectados de Neurofibromatosis*, Chromo22

and ACNefi. This work has been supported by: *Associació Chromo 22*; The Children's Tumor

Foundation (CTF-2019-05-005), the Spanish Association for the NF Affected (AANF), *Fundación*

Proyecto Neurofibromatosis and the Catalan NF Association (AcNeFi). This study has been

funded by the *Instituto de Salud Carlos III* through the project PI20/00215 (Co-funded by

European Regional Development Fund "A way to make Europe"), *Fundació La Marató de TV3*

(126/C/2020) and the Government of Catalonia (2017 SGR 496).

Author contributions

ES, MC and EC conceived and supervised the study and wrote the manuscript that was revised,

corrected, and improved by all the authors. NC performed most of the experimental work,

analysed the data, generated the figures for the paper and wrote the manuscript. MTM and

BG performed bioinformatics analysis. AN, BK, IR, GC and BA performed experimental work.

FRR and EA contributed with sample collection. HM, AV, AR and IB provided scientific input. All

authors approved the final version of the manuscript.

Conflict of interest statement

483 The authors declare no conflicts of interest.

References

1.   Evans DGR, Moran A, King A, Saeed S, Gurusinghe N, Ramsden R. Incidence of vestibular

schwannoma and neurofibromatosis 2 in the North West of England over a 10-year

period: Higher incidence than previously thought. Otol Neurotol. 2005;26(1):93–7.

2.   Plotkin SR, Messiaen L, Legius E, Pancza P, Avery RA, Blakeley JO, et al. Updated

diagnostic   criteria   and   nomenclature   for   neurofibromatosis   type   2   and

schwannomatosis: An   international   consensus   recommendation.   Genet   Med.

2022;24(9):1967–77.

3.   McClatchey AI, Giovannini M. Membrane organization and tumorigenesis - The NF2

tumor suppressor, Merlin. Genes Dev. 2005;19(19):2265–77.

4.   Gladden AB, Hebert AM, Schneeberger EE, McClatchey AI. The NF2 Tumor Suppressor,

Merlin, Regulates Epidermal Development through the Establishment of a Junctional

Polarity   Complex.   Dev   Cell   [Internet].   2010;19(5):727–39.   Available   from:

http://dx.doi.org/10.1016/j.devcel.2010.10.008

5.   Petrilli AM, Merlin V, Nf I, Merlin I, Nf O. Role of Merlin / NF2 inactivation in tumor

biology. 2016;(December 2014):537–48.

6.   Evans DG, Bowers NL, Tobi S, Hartley C, Wallace AJ, King AT, et al. Schwannomatosis: A

genetic and epidemiological study. J Neurol Neurosurg Psychiatry. 2018 Jun 14;

7.   Evans DGR, Huson SM, Donnai D, Neary W, Blair V, Teare D, et al. A genetic study of

type 2 neurofibromatosis in the United Kingdom. I. Prevalence, mutation rate, fitness,

and confirmation of maternal transmission effect on severity. J Med Genet.

1992;29(12):841–6.

8.   Parry DM, Eldridge R, Kaiser-Kupfer MI, Bouzas EA, Pikus A, Patronas N.

Neurofibromatosis 2 (NF2): Clinical characteristics of 63 affected individuals and clinical

evidence for heterogeneity. Am J Med Genet. 1994;52(4):450–61.

9.   Baser ME, Makariou E V., Parry DM. Predictors of vestibular schwannoma growth in

patients with neurofibromatosis Type 2. J Neurosurg. 2002 Feb;96(2):217–22.

10.   Forde C, King AT, Rutherford SA, Hammerbeck-Ward C, Lloyd SK, Freeman SR, et al.

Disease course of neurofibromatosis type 2: A 30-year follow-up study of 353 patients

seen at a single institution. Neuro Oncol. 2021;23(7):1113–24.

11.   Ragge NK, Baser ME, Klein J, Nechiporuk A, Sainz J, Pulst SM, et al. Ocular abnormalities

in neurofibromatosis 2. Am J Ophthalmol [Internet]. 1995;120(5):634–41. Available

from: http://dx.doi.org/10.1016/S0002-9394(14)72210-X

12.   Plana-Pla A, Bielsa-Marsol I, Carrato-Moñino C. Diagnostic and Prognostic Relevance of

the Cutaneous Manifestations of Neurofibromatosis Type 2. Actas Dermosifiliogr.

2017;108(7):630–6.

13.   Hexter A, Jones A, Joe H, Heap L, Smith MJ, Wallace AJ, et al. Clinical and molecular

predictors of mortality in neurofibromatosis 2: A UK national analysis of 1192 patients. J

Med Genet. 2015;52(10):699–705.

14.   Dirks MS, Butman JA, Kim HJ, Wu T, Morgan K, Tran AP, et al. Long-term natural history

of neurofibromatosis Type 2-associated intracranial tumors: Clinical article. J

Neurosurg. 2012 Jul;117(1):109–17.

15. Peyre M, Goutagny S, Bah A, Bernardeschi D, Larroque B, Sterkers O, et al. Conservative

management of bilateral vestibular schwannomas in neurofibromatosis type 2 patients:

Hearing and tumor growth results. Neurosurgery. 2013 Jun;72(6):907–13.

16. Baser ME, Mautner VF, Parry DM, Evans DGR. Methodological issues in longitudinal

studies: Vestibular schwannoma growth rates in neurofibromatosis 2. Vol. 42, Journal

of Medical Genetics. BMJ Publishing Group; 2005. p. 903–6.

17. Lloyd SKW, Evans DGR. Neurofibromatosis type 2 (NF2): Diagnosis and management

[Internet]. 1st ed. Vol. 115, Handbook of Clinical Neurology. Elsevier B.V.; 2013. 957–

967 p. Available from: http://dx.doi.org/10.1016/B978-0-444-52902-2.00054-0

18. Plotkin SR, Duda DG, Muzikansky A, Allen J, Blakeley J, Rosser T, et al. Multicenter,

prospective, phase II and biomarker study of high-dose bevacizumab as induction

therapy in patients with neurofibromatosis type 2 and progressive vestibular

schwannoma. J Clin Oncol. 2019;37(35):3446–54.

19. McClatchey AI, Saotome I, Ramesh V, Gusella JF, Jacks T. The Nf2 tumor suppressor

gene product is essential for extraembryonic development immediately prior to

gastrulation. Genes Dev. 1997;11(10):1253–65.

20. Behling F, Bersali I, Santacroce A, Hempel J, Kandilaris K, Schittenhelm J, et al. Transition

of a vestibular schwannoma to a malignant peripheral nerve sheath tumor with loss of

H3K27 trimethylation after radiosurgery—a case report and review of the literature.

Neurosurg Rev [Internet]. 2022;45(1):915–22. Available from:

https://doi.org/10.1007/s10143-021-01620-3

21. Wu L, Vasilijic S, Sun Y, Chen J, Landegger LD, Zhang Y, et al. Losartan prevents tumor-

induced hearing loss and augments radiation efficacy in NF2 schwannoma rodent

models [Internet]. Vol. 13, Sci. Transl. Med. 2021. Available from:

https://www.science.org

22.   Karajannis MA, Legault G, Hagiwara M, Ballas MS, Brown K, Nusbaum AO, et al. Phase II

trial of lapatinib in adult and pediatric patients with neurofibromatosis type 2 and

progressive vestibular schwannomas. Neuro Oncol. 2012 Sep;14(9):1163–70.

23.   Osorio DS, Hu J, Mitchell C, Allen JC, Stanek J, Hagiwara M, et al. Effect of lapatinib on

meningioma growth in adults with neurofibromatosis type 2. J Neurooncol. 2018 Sep

1;139(3):749–55.

24.   Welling DB, Collier KA, Burns SS, Oblinger JL, Shu E, Miles-Markley BA, et al. Early phase

clinical studies of AR-42, a histone deacetylase inhibitor, for neurofibromatosis type 2-

associated vestibular schwannomas and meningiomas. Laryngoscope Investig

Otolaryngol. 2021 Oct 1;6(5):1008–19.

25.   McClatchey AI, Saotome I, Mercer K, Crowley D, Gusella JF, Bronson RT, et al. Mice

heterozygous for a mutation at the Nf2 tumor suppressor locus develop a range of

highly metastatic tumors. Genes Dev. 1998;12(8):1121–33.

26.   Stemmer-Rachamimov AO, Louis DN, Nielsen GP, Antonescu CR, Borowsky AD, Bronson

RT, et al. Comparative pathology of nerve sheath tumors in mouse models and humans.

Cancer Res. 2004;64(10):3718–24.

27.   Giovannini M, Robanus-Maandag E, Van Der Valk M, Niwa-Kawakita M, Abramowski V,

Goutebroze L, et al. Conditional biallelic Nf2 mutation in the mouse promotes

manifestations of human neurofibromatosis type 2 [Internet]. 2000. Available from:

www.genesdev.org

28.   Ren Y, Chari DA, Vasilijic S, Welling DB, Stankovic KM. New developments in

neurofibromatosis type 2 and vestibular schwannoma. Neuro-Oncology Adv.

2021;3(1):1–13.

29.   Dilwali S, Patel PB, Roberts DS, Basinsky GM, Harris GJ, Emerick KS, et al. Primary

culture of human Schwann and schwannoma cells: Improved and simplified protocol.

577   Hear Res. 2014;315:25–33.

578   30.   Rosenbaum C, Kluwe L, Mautner VF, Friedrich RE, Mü HW, Hanemann CO. Isolation and

579   Characterization of Schwann Cells from Neurofibromatosis Type 2 Patients. 1998.

580   31.   Xue L, He W, Wang Z, Chen H, Wang Z, Wu H. Characterization of a newly established

581   schwannoma cell line from a sporadic vestibular schwannoma patient. Am J Transl Res.

582   2021;13(8):8787–803.

583   32.   Hung G, Rhim JS, Slattery W, Lim D. Establishment and characterization of a

584   schwannoma cell line from a patient with neurofibromatosis 2. Vol. 20, INTERNATIONAL

585   JOURNAL OF ONCOLOGY. 2002.

586   33.   Takahashi K, Yamanaka S. Induction of Pluripotent Stem Cells from Mouse Embryonic

587   and Adult Fibroblast Cultures by Defined Factors. 2006;2:663–76.

588   34.   Shi Y, Inoue H, Wu JC, Yamanaka S. Induced pluripotent stem cell technology: A decade

589   of progress. Vol. 16, Nature Reviews Drug Discovery. Nature Publishing Group; 2017. p.

590   115–30.

591   35.   Papapetrou EP. Patient-derived induced pluripotent stem cells in cancer research and

592   precision oncology. Vol. 22, Nature Medicine. Nature Publishing Group; 2016. p. 1392–

593   401.

594   36.   Raya Á, Rodríguez-Piz I, Guenechea G, Vassena R, Navarro S, Barrero MJ, et al. Disease-

595   corrected haematopoietic progenitors from Fanconi anaemia induced pluripotent stem

596   cells. Nature. 2009;460(7251):53–9.

597   37.   Mungenast AE, Siegert S, Tsai LH. Modeling Alzheimer's disease with human induced

598   pluripotent stem (iPS) cells. Mol Cell Neurosci [Internet]. 2016;73:13–31. Available

599   from: http://dx.doi.org/10.1016/j.mcn.2015.11.010

600   38.   Nourbakhsh A, Gosstola NC, Fernandez-Valle C, Dykxhoorn DM, Liu XZ. Characterization

601   of UMi031-A-2 inducible pluripotent stem cell line with a neurofibromatosis type 2-

602   associated mutation. Stem Cell Res. 2021 Aug 1;55.

39.   Ishi Y, Era T, Yuzawa S, Okamoto M, Sawaya R, Motegi H, et al. Analysis of induced

pluripotent stem cell clones derived from a patient with mosaic neurofibromatosis

type 2. Vol. 188, American journal of medical genetics. Part A. United States; 2022. p.

1863–7.

40.   Jessen KR, Mirsky R. The origin and development of glial cells in peripheral nerves. Nat

Rev Neurosci. 2005;6(9):671–82.

41.   Menendez L, Kulik MJ, Page AT, Park SS, Lauderdale JD, Cunningham ML, et al. Directed

differentiation of human pluripotent cells to neural crest stem cells. Nat Protoc. 2013

Jan;8(1):203–12.

42.   Carrió M, Mazuelas H, Richaud-Patin Y, Gel B, Terribas E, Rosas I, et al. Reprogramming

Captures the Genetic and Tumorigenic Properties of Neurofibromatosis Type 1

Plexiform Neurofibromas. Stem Cell Reports. 2019;12.

43.   Baser ME, Friedman JM, Wallace AJ, Ramsden RT, Joe H, Evans DGR. Evaluation of

clinical diagnostic criteria for neurofibromatosis 2. Neurology. 2002 Dec;59(11):1759–

65.

44.   Castellanos E, Gel B, Rosas I, Tornero E, Santín S, Pluvinet R, et al. A comprehensive

custom panel design for routine hereditary cancer testing: Preserving control,

improving diagnostics and revealing a complex variation landscape. Sci Rep. 2017;7.

45.   Mazuelas H, Magallón-Lorenz M, Fernández-Rodríguez J, Uriarte-Arrazola I, Richaud-

Patin Y, Terribas E, et al. Modeling iPSC-derived human neurofibroma-like tumors in

mice uncovers the heterogeneity of Schwann cells within plexiform neurofibromas. Cell

Rep   [Internet].   2022   Feb;38(7):110385.   Available   from:

https://linkinghub.elsevier.com/retrieve/pii/S2211124722001061

46.   Jessen KR, Mirsky R, Lloyd AC. Schwann cells: Development and role in nerve repair.

Cold Spring Harb Perspect Biol. 2015 Jul 1;7(7):1–15.

47.   Yilmaz A, Peretz M, Aharony A, Sagi I, Benvenisty N. Defining essential genes for human

pluripotent stem cells by CRISPR-Cas9 screening in haploid cells. Nat Cell Biol [Internet].

2018;20(5):610–9. Available from: http://dx.doi.org/10.1038/s41556-018-0088-1

48. Blomen VA, Májek P, Jae LT, Bigenzahn JW, Nieuwenhuis J, Staring J, et al. Gene

essentiality and synthetic lethality in haploid human cells. Science (80-).

2015;350(6264):1092–6.

49. Nourbakhsh A, Gosstola NC, Fernandez-Valle C, Dykxhoorn DM, Liu XZ. Characterization

of UMi031-A-2 inducible pluripotent stem cell line with a neurofibromatosis type 2-

associated mutation. Stem Cell Res [Internet]. 2021;55(June):102474. Available from:

https://doi.org/10.1016/j.scr.2021.102474

50. Rowe RG, Daley GQ. Induced pluripotent stem cells in disease modelling and drug

discovery. Vol. 20, Nature Reviews Genetics. Nature Publishing Group; 2019. p. 377–88.

51. Curto M, McClatchey AI. Nf2/Merlin: A coordinator of receptor signalling and

intercellular contact. Br J Cancer. 2008;98(2):256–62.

52. Curto M, Cole BK, Lallemand D, Liu CH, McClatchey AI. Contact-dependent inhibition of

EGFR signaling by Nf2/Merlin. J Cell Biol. 2007 Jun 4;177(5):893–903.

53. Catasús N, Rosas I, Bonache S, Negro A, Torres-Martin M, Plana-Pla A, et al. Antisense

oligonucleotides targeting exon 11 are able to partially rescue the NF2-related

schwannomatosis phenotype in vitro. Mol Ther - Nucleic Acids [Internet].

2022;30(December):493–505. Available from:

https://doi.org/10.1016/j.omtn.2022.10.026

54. Castellanos E, Plana A, Carrato C, Carri M, Rosas I, Amilibia E, et al. Early genetic

diagnosis of neurofibromatosis type 2 from skin plaque plexiform schwannomas in

childhood. JAMA Dermatology. 2018 Mar 1;154(3):341–6.

55. Van der Auwera GA, O'Connor BD. Genomics in the Cloud: Using Docker, GATK, and

WDL in Terra [Internet]. O'Reilly Media; 2020. Available from:

https://books.google.es/books?id=vsXaDwAAQBAJ

56.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

656   universal RNA-seq aligner. Bioinformatics. 2013 Jan;29(1):15–21.

657   57.   Hartley SW, Mullikin JC. QoRTs: a comprehensive toolset for quality control and data

658   processing of RNA-Seq experiments. BMC Bioinformatics. 2015 Jul;16(1):224.

659   58.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

660   RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

661   59.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential

662   expression analysis of digital gene expression data. Bioinformatics. 2010 Jan;26(1):139–

663   40.

NF2$^{+/+}$ cells
FiPS

Gene editing

NF2$^{-/-}$ iPSC

NF2$^{-/-}$ NC

SC Differentiation
in 3D

NF2$^{-/-}$ spheroid

Gene editing

NC Differentiation

NF2$^{+/-}$ iPSC

NF2$^{+/-}$ NC

NF2$^{+/-}$ spheroid

VS

NF2 patient

Vestibular Schwannoma

NF2$^{+/-}$ cells

Reprogramming

NF2$^{-/-}$ SC

NF2$^{-/-}$ iPSC
SeV - dependent
(Discarded)