Habitual ground nesting in the Bugoma Forest chimpanzees (*Pan troglodytes schweinfurthii*),

Uganda.

Catherine Hobaiter[1,2]

Harmonie Klein[1]

Thibaud Gruber[2,3]

[1] Wild Minds Lab; School of Psychology and Neuroscience, University of St Andrews,

Scotland KY16 8LT

[2] Bugoma Primate Conservation Project, Bugoma Central Forest Reserve, Hoima, Uganda

[3] Faculty of Psychology and Educational Sciences and Swiss Center for Affective Sciences,

University of Geneva, Geneva CH-1202

clh42@st-andrews.ac.uk; hk96@st-andrews.ac.uk; thibaud.gruber@unige.ch

correspondence: Dr Catherine Hobaiter, School of Psychology and Neuroscience, University

of St Andrews, Scotland KY16 8LT; +44 (0)1334 462053; clh42@st-andrews.ac.uk

Running Title: Ground nesting in Bugoma chimpanzees

Abstract

We report the presence of habitual ground nesting in a newly studied East African

chimpanzee *(Pan troglodytes schweinfurthii)* population in the Bugoma Central Forest

Reserve, Uganda. Across a 2-year period we encountered 989 night-nests, 205 of which were

classified as ground nests, a rate of ~21%. We find no preliminary evidence of socio-

ecological factors that would promote its use and highlight local factors that appear to make

its use disadvantageous. While further study is required to establish this behaviour as meeting

the strict criteria for non-human animal culture, we support the argument that the wider use

of population and group-specific behavioural repertoires in flagship species, such as

chimpanzees, offers a tool to promote the urgent conservation action needed to protect

threatened ecosystems, including the Bugoma forest.

Keywords: Chimpanzee, Culture, Nesting, Uganda

Introduction

All apes construct overnight arboreal nests by interweaving branches and other vegetation

(Fruth et al., 2018; Anderson et al., 2019). Gorillas, while able to climb while foraging, are

the most terrestrial of the great apes and build overnight nests primarily on the ground using

available herbaceous vegetation, although they increase the use of arboreal nesting

seasonally, or when terrestrial vegetation is limited (Tutin et al., 1995; Brugiere & Sakom,

2000; Mfossa et al., 2022). In contrast, orangutans, bonobos, and chimpanzees primarily

build arboreal nests (Goodall 1962; Fruth & Hohmann, 1993; Prasetyo et al., 2009; Hicks,

2010; van Casteren et al., 2012; Fruth et al., 2018), using multiple interwoven supporting

branches, filling in with smaller softer material (Furuichi & Hashimoto, 2000; Prasetyo et al.,

2009; Samson & Hunt, 2012; Fruth et al., 2018). There appears to be an effect of learning,

with younger individuals constructing more robust nests earlier if exposed to adults who

build nests (Lethmate, 1977; McGrew, 2004; Videan, 2006). Each independent individual

constructs their own nest, but individuals will typically nest in a party with others in the same

or near-by trees (Fruth, 1995; Schaller, 1963; Goodall, 1968). Nests are constructed each

evening for overnight use, but they are also built during the day for a range of reasons from

sleeping, to play, or sexual solicitation (Boesch, 1995; Plumptre and Reynolds, 1997; Fruth

and Hohmann, 1996; Brownlow et al., 2001; McGrew, 2010; Fruth et al., 2018). with

bonobos also regularly constructing additional full nests during the day (Wessling & Surbeck

2022). Chimpanzees and bonobos typically construct nests at between 8 to 20m in height, but

they may be as low as 3-4m typically where taller trees are limited (Pruetz et al., 2008;

Samson & Hunt, 2012; Fruth et al., 2018). Full nests – night or day – are distinguished in

their construction from other flimsier resting structures such as 'day beds' and 'cushions',

which typically involve only 1-2 bent branches, loosely interwoven, and may be as simple as

a single small sapling bent over or a clump of ferns (Boesch, 1995; Furuichi & Hashimoto,

2000; Brownlow et al., 2001; Koops et al., 2007).

Despite the use of arboreal night nesting in every chimpanzee community studied to

date, our understanding of their function remains limited, with – potentially complementary –

hypotheses currently including seasonal patterns (Samson & Hunt, 2014; van Casteren et al.,

2012), thermoregulation (Koops et al., 2012; McGrew, 2004; Stewart et al., 2018), pathogen

and parasite avoidance (Anderson, 1998; Lacroux et al., 2022), and predation avoidance

(Koops et al., 2012; Kortlandt, 1992; Pruetz et al., 2008; Stewart & Pruetz, 2013). Some

aspects of the local environment appear to impact the choice of nest location, for example a

preference for slopes over flatter ground (Issa: Hernandez-Aguilar 2009; Mahale: Izawa and

Itani, 1966; Suzuki, 1969; Assirik: Baldwin, 1979). However, in habitats where trees are

limited this tends to lead to a concentration of nests in available trees (Hernandez-Aguilar

2009), rather than alternative strategies such as ground nesting.

The very occasional use of overnight ground nests is widely reported in all chimpanzee sub-

species, including in Uganda (see below and c.f. Tagg et al., 2013). Excluding the minimally

constructed day beds and cushions (Furuichi & Hashimoto, 2000; Koops et al., 2007), the use

of fully constructed overnight nests on the ground is typically rare (5 to 10% of nests

constructed in a given population; Matsuzawa & Yamakoshi, 1996; Koops et al. 2007),

perhaps as a fall-back by sick or injured individuals (<1%; Furuichi & Hashimoto, 2000).

Slightly more frequent use of overnight ground nests was reported by male West African

chimpanzees (*Pan troglodytes verus*) of Seringbara in the Nimba Mountains in Guinea (~3%

to 5% of night nests; Koops et al., 2007; 2012) and of Fongoli long term field site (~12% of

night nests; Stewart 2011). Frequent ground nests have also been recently observed in

Cameroun in Nigeria-Cameroun chimpanzees (*Pan troglodytes ellioti*) of one (i.e., Andu) of

the two field sites of the Lebialen-Mone Forest (~32% of night nests; Last & Muh, 2013) and

in Central African chimpanzees (*Pan troglodytes troglodytes)* of La Belgique research field

site in the Dja Biosphere Reserve (~3% to ~9% of night nests; Guislain & Duplain, 2005;

Tagg et al., 2013). However, despite ~ 250 years of continuous observations across long-term

research sites (Gombe, >60 years; Mahale, >60 years; Kibale-Kanyawara, >30 years; Kibale-

Ngogo, >20 years; Kalinzu, >30 years; Budongo-Sonso, >30 years, Budongo-Waibira, >10

years); the customary use of overnight ground nests has only been reported in East African

chimpanzee communities in the Northern DRC (~11% of nesting sites; Hicks et al., 2019),

approximately 600km from the Bugoma Forest Reserve, and separated by a large body of

water (i.e., Lake Albert).

Here we report on the presence of apparently habitual ground nesting in the newly

studied East African chimpanzee *(Pan troglodytes schweinfurthii)* population in the Bugoma

Central Forest Reserve, in Uganda. We find no preliminary evidence of specific socio-

ecological factors that would promote its use and highlight local factors that appear to make

its use disadvantageous. Establishing a case for cultural behaviour on the basis of either

evidence of social transmission or the exclusion of alternative hypotheses, such as genetic or

ecological variation may take decades in long-lived species. However, we argue that the use

of apparently cultural group-differences in the behavioural repertoires in flagship species,

including chimpanzees – irrespective of the socio-ecological or cognitive mechanism by

which they are produced – offers an effective tool to promote the urgent conservation action

needed to protect threatened ecosystems, such as the Bugoma forest.

Methods

Site

The Bugoma Central Forest Reserve includes 400km^2 of semi-deciduous tropical rain forest

(located at 01°15'N 30°58'E and at 990-1300m elevation). Located between the Central

Forest Reserve of Budongo (425km^2) and Kibale National Park (776km^2), Bugoma

represents the largest contiguous forest habitat for chimpanzees in East Africa in which there

has been no long-term research activity. In 2006, chimpanzee density was estimated at 1.9

chimpanzees per km^2 (giving an approximate population estimate of ~760 chimpanzees;

Plumptre and Cox, 2006) but given widespread primate population declines (Estrada et al.,

2017) a current population estimate of ~600 chimpanzees may be more appropriate.

Following surveys in 2015, in 2016 the Bugoma Primate Conservation Project (BPCP,

www.bugomaprimates.com) initiated daily habituation and monitoring activities across

several chimpanzee communities. Our data are concentrated on the Mwera-South

community, where regular contacts were initiated in 2016 and were typically occurring on

most days by 2018. Chimpanzees in the Mwera-North community were also encountered, as

their range overlaps with that of Mwera-South, but they are not intentionally followed and

they still tend to flee when our presence is detected allowing us to distinguish between the

groups. The current estimated range for the Mwera-South community is ~10km2 (see Figure

1). During the study period, chimpanzees in the Mwera-South community were followed on

regular basis (several days a week for several hours per day) with an estimate of 60-80

individuals based on local East African chimpanzee community sizes (Wilson et al., 2014).

However, as it was not possible to nest or denest them, direct observations of night nest

building were not available.

Data collection

In addition to direct observations of chimpanzee behavioural and health data, project staff

record the type and GPS location of signs of chimpanzee activity (prints, dung, feeding

remains, discarded tools, and nests), and of illegal human activity (primarily hunting and

logging) whenever they are encountered ad libitum.

A 'nesting site' was defined as one or more nests of the same age in which no two

nests were more than 30m apart (Furuichi & Hashimoto, 2000). Following increasing

encounters with overnight ground nests, in 2019, BPCP staff started to collect detailed

information on overnight nests. However, data collection was impacted by the Covid19

outbreak with 1) a stop of data collection from February 2020 to July 2020 and 2) a change of

data collection protocol after this period. Therefore, here, we provide the following nesting

data: GPS location, approximate age (1 day, 2 days, under a week, over a week), the nest type

(ground, tree) and the number of ground and tree nests in the nesting site from August 2019

to June 2022. Other nesting data, including: density of surrounding forest (Open, Medium,

Dense), forest type (climax, secondary-mixed, disturbed, water-swamp, grassland), the height

from the ground of all tree nests (m), the primary tree species used, the width and length of

ground nests (cm), and the use of bent or detached branches and the inclusion of Terrestrial

Herbaceous Vegetation (THV) in ground nest construction, are only available for the pre-

Covid19 outbreak, i.e. from August 2019 to February 2020.

Data availability

All data used for the analyses and figures in this manuscript are available in a publicly

accessible GitHub repository: https://github.com/Wild-Minds/GroundNesting_Bugoma

Ethical statement

This study was observational and did not involve any interventions, apart from daily visits to

the chimpanzee communities' territories. BPCP staff follow strict hygiene and observation

distance rules to prevent disease transmission. Following the Covid19 outbreak, from March

2020 data collection moved from chimpanzee follows to biomonitoring only. Permissions to

collect data were provided by the National Forest Authority of Uganda, the Uganda Wildlife

Authority, and the Ugandan National Council of Science and Technology, and all data

collection adhered to national and international guidelines, including the American Society of

Primatologists Principles for the Ethical Treatment of Non-Human Primates, and the Code of

Best Practices for Field Primatology.

Results

Presence of ground and tree nests

Between August 2019 and July 2022, we recorded 989 night-nests within 336 nesting sites,

number of nests per site ranged from 1 to 20 (Figure 1). We recorded 784 tree nests and 205

ground nests (21%). Ground nests occurred in 82 of the 336 sites (24%) and co-occurred

with tree nests in 29 of these sites (6% all nest sites; 35% of sites with a ground nest; up to 15

ground nests within a site). Ground nests were observed year-round (57% in the dry season;

43% in wet season).

Between August 2019 and February 2020 we recorded additional information on nest

construction and local habitat for 138 nesting sites containing 310 night-nests(269 tree nests,

41 ground nests). Tree nests varied in height from 1-40m; and n=11 (4%) were at 2m or less

from the ground. All ground nests that could be measured (n=39) contained detached

interwoven branches, and most (n=34, 87%) incorporated bent branches as well. A small

number (n=2) incorporated THV into the construction. The majority of nest sites were

located in medium-density forest (n=117) as opposed to open (n=17) or high-density (n=4);

similarly, the majority of nest sites were located in mixed-secondary forest (n=121) rather

than disturbed (n=8) or savannah (n=1; Unknown n=8).

Discussion

Bugoma chimpanzees regularly construct ground nests overnight and, in addition,

construct tree nests at a height that would be vulnerable to human threat (<2m). In addition to

their observed frequency, we find ground nests most often appeared as clusters suggesting

that they were not built by a single individual or a single family. As these chimpanzees are

not fully habituated, direct observations of overnight ground nest building were not possible.

The estimated number of independent individuals in the community (~40) is similar to the

total number of ground nests encountered, so it remains possible that each ground nest

observed was built by a separate individual on only one occasion; however, we would

suggest that this is less likely than the repeated building of ground nests by at least some

individuals. If so, ground nesting in the Mwera-South community of chimpanzees meets the

criteria for a 'habitual' behaviour (Whiten et al., 1999).

It is possible that local ecological factors promote the use of ground nesting in the

Bugoma communities. For example: a limited number of suitable trees for tree nesting

(McCarthy et al. 2016). While systematic research is needed to confirm this, we believe that

this is highly unlikely both because of the regular co-occurrence of tree and ground nests in

the same nesting party; and because the secondary mixed-forest habitat in which the majority

of nest sites are located is similar in structure to that of the neighbouring Budongo Forest

Reserve, in which no overnight ground nests have been reported in over 40-years of

cumulative observation across two communities. Another explanation could be seasonality

(Pruetz et al. 2018; Tagg et al., 2013), but Bugoma chimpanzees seem to nest on the ground

year round (see also Koops et al. 2012). An alternative might be that large numbers of

individuals are unable to construct tree nests due to injury or illness. Again, we believe that

this is unlikely. Our long-term data collection includes health monitoring of the chimpanzees

and while there are outbreaks of respiratory infection in some individuals, these appear

similar to those reported in other Uganda forest populations (Scully et al., 2018). A more

likely pressure would be the impact of snare wires which regularly maim chimpanzees who

trap their hands and feet in them – these injuries can include wasted and amputated limbs.

However, once again, the neighbouring Budongo communities also have high numbers of

snare-injured chimpanzees (Reynolds, 2005; Stokes & Byrne, 2006; Waller & Reynolds,

2001), including individuals missing hands and feet (including individuals where more than

one limb is amputated), and do not report any use of ground nesting. In contrast, ground

nesting has sometimes been explained by scholars as resulting from an absence of predators

or human activity (Pruetz et al. 2008; Hicks, 2010; McCarthy et al., 2012; Last & Muh,

2013). However, it's unlikely the case at Bugoma as a number of features of the forest make

ground nesting potentially risky. While there are no recent confirmed sightings of

chimpanzees' main nonhuman predator (leopard, *Panthera pardus*) in Bugoma, other large

animals that could present potential risk were regularly directly or indirectly (through

camera-traps) observed over the study period, including elephant (*Loxodonta cyclotis*),

golden cat (*Caracal aurata*), python (*Python sebae*), and rhinoceros viper (*Bitis nasicornis*;

Figure 2a). Elephant presence has been previously described as impacting ground nesting at

Lope (Tutin et al., 1995) and they are also present other sites where ground nesting has been

observed (i.e., La Belgique, Tagg et al., 2013; Bili-Uele, Hicks 2010). Perhaps the most

significant risk to chimpanzees in Bugoma is human activity. Hunting is prominent; as well

as the use of snare wires designed to target duiker and pig species, hunters in Bugoma are

regularly encountered with large packs of dogs in the chimpanzees' range (Figure 2b). The

butchered remains of chimpanzees, apparently following human hunting, have been

discovered by the project on two occasions, with additional unconfirmed reports (Bugoma

Primate Conservation Project, unpublished data). Interestingly, while it remains counter-

intuitive, Tagg and colleagues (2013) also found that chimpanzee ground nesting appeared to

increase with human pressure. Given the lack of obvious ecological or physical drivers,

together with the combination of significant risk to life by, at least, human hunters, we

suggest that there is likely a cultural component to the Bugoma chimpanzee ground nesting.

Importantly, this highlights that, as in other species (Laland & Williams, 1998; Franz &

Matthews, 2010), some socially acquired chimpanzee behavioural variants may not have a

positive effect on individual fitness. Behavioural variants linked to improving vital rates,

such as survival, were suggested as being of particular conservation importance (Brakes et

al., 2021), but establishing these connections is challenging (Carvalho et al., 2022) and

association with individual fitness may vary with fluctuating local pressures, for example

human hunting.

Chimpanzee cultures have been shown to be inflexible, with resistance to the

modification or acquisition of new behaviour even where it could be advantageous (Boesch

& Boesch-Achermann, 2000; Gruber et al., 2011; but see Hobaiter et al., 2014), suggesting

that they may be similarly resistant to its loss where it appears to be disadvantageous. With

increasing human pressure on the forest, it is unclear how the Bugoma chimpanzees will react

with respect to their nest building behaviour. Closely monitoring ground nest building in

Bugoma forest may provide a direct assessment of the impact of anthropogenic activities on

wild semi-habituated chimpanzees.

The importance of community and population specific cultural behaviour from a

conservation perspective has started to receive significant attention (Brakes et al., 2019;

Brakes et al., 2021). The loss of a specific community involves the irreversible loss of a

unique culture and the information or individual knowledge that it is founded on. The

importance of these cultural markers in chimpanzee conservation was recently recognised by

the United Nations Convention on the Conservation of Migratory Species, who dedicated a

Concerted Action to the conservation of West African chimpanzee (*Pan troglodytes verus*)

nut cracking, the first time a non-human animal behaviour was recognised in this way. The

wider Bugoma forest habitat is under considerable pressure, not only from illegal activities

such as hunting and logging, but also from clear-felling for crop production, from local

energy resource extraction, and from increased infrastructure and anthropogenic activities

(McLennan et al., 2020; Plumptre et al., 2020). Non-human cultures are sensitive to the

impact of human anthropogenic activity on the transformation, fragmentation, and

degradation of habitats (Gruber et al., 2019; Kalan et al., 2020). Widespread urgent calls to

action throughout 2019 to 2022 for its protection by Ugandan and international organisations,

as well as planned sustainable resource generation from eco-tourism projects may benefit

from a clear and strong case for the unique behavioural repertoire of the chimpanzees in this

area.

Acknowledgements

We thank the staff of the Bugoma Primate Conservation Project, and the Ugandan National

Forestry Authority and the Jane Goodall Institute Switzerland for their support of the field

site. Fieldwork for this research received funding from the European Union's 8[th] Framework

Programme, Horizon 2020, under grant agreement no 802719; National Geographic (Grant

GS-63895R-19), the Swiss National Science Foundation (Grant PCEFP1_186832). Funders

had no role in the study design, writing, or decision to publish.

References

Anderson, J. R. (1998). Sleep, sleeping sites, and sleep-related activities: awakening to their

significance. *American Journal of Primatology,* 46, 63-75.

Anderson J.R., Ang M.Y.L., Lock M.C., & Weiche, I. (2019). Nesting, sleeping, and

nighttime behaviors in wild and captive great apes. *Primates,* 60, 321-332.

https://doi.org/10.1007/s10329-019-00723-2

Baldwin, P.J. (1979). *The natural history of the chimpanzee (Pan troglodytes verus) at Mt.*

Assrik, Senegal. (Doctoral dissertation, University of Stirling).

Boesch, C. (1995). Innovation in wild chimpanzees (*Pan troglodytes*). *International Journal*

of Primatology, 16(1), 1-16.

Boesch, C., & Boesch-Achermann, H. (2000). *The chimpanzees of the Taï forest:*

Behavioural ecology and evolution. Oxford: Oxford University Press.

Brakes, P., Dall, S. R., Aplin, L. M., Bearhop, S., Carroll, E. L., Ciucci, P., … Rutz, C.

(2019). Animal cultures matter for conservation. *Science,* 363, 1032-1034.

Brakes, P., Carroll, E. L., Dall, S. R., Keith, S.A., McGregor, P. K., Mesnick, S. L., …

Garland, E. C. (2021). A deepening understanding of animal culture suggests lessons for

conservation. *Proceedings of the Royal Society B: Biological Sciences,* 288,

rspb.2020.2718

Brownlow, A. R., Plumptre, A. J., Reynolds, V., & Ward, R. (2001). Sources of variation in

the nesting behavior of chimpanzees (*Pan troglodytes schweinfurthii*) in the Budongo

Forest, Uganda. *American Journal of Primatology*, 55(1), 49-55.

Brugiere, D., & Sakom, D. (2001). Population density and nesting behaviour of lowland

gorillas (Gorilla gorilla gorilla) in the Ngotto forest, Central African Republic. *Journal of*

Zoology, 255(2), 251-259.

Carvalho, S., Wessling, E. G., Abwe, E. E., Almeida☐Warren, K., Arandjelovic, M., Boesch,

C., ... & Koops, K. (2022). Using nonhuman culture in conservation requires careful and

concerted action. *Conservation Letters*, *15*(2), e12860.

Estrada, A., Garber, P. A., Rylands, A. B., Roos, C., Fernandez-Duque, E., Di Fiore, A., …

Li, B. (2017). Impending extinction crisis of the world's primates: Why primates matter.

Science Advances, 3(1), e1600946.

Franz, M., & Matthews, L. J. (1998) Social enhancement can create adaptive, arbitrary, and

maladaptive cultural traditions. *Proceedings of the Royal Society B: Biological Sciences*

doi:10.1098/rspb.2010.0705

Fruth, B. (1995). Nests and nest groups in wild bonobos (*Pan paniscus*): Ecological and

behavioural correlates. Aachen: Verlag Shaker.

Fruth, B., & Hohmann, G. (1993). Ecological and behavioral aspects of nest building in wild

bonobos (*Pan paniscus*). *Ethology*, *94*(2), 113-126.

Fruth, B., & Hohmann, G. (1996). Comparative analyses of nest building behavior in

Bonobos and Chimpanzees. In: Wrangham, R.W, W.C. McGrew, F.B.M. de Wall & P.G.

Heltne (eds). Chimpanzee Cultures. Harvard University Press: xxii pp 424.

Fruth, B., Tagg, N., & Stewart, F. (2018). Sleep and nesting behavior in primates: A

review. *American Journal of Physical Anthropology*, 166(3), 499-509.

Furuichi, T., & Hashimoto, C. (2000). Ground beds of chimpanzees in the Kalinzu Forest,

Uganda. *Pan Africa News,* 7, 26-28.

Galef, B. G., & Whiten, A. (2017). The comparative psychology of social learning. In J. Call,

G. M. Burghardt, I. M. Pepperberg, C. T. Snowdon, T. Zentall (Eds.), *APA handbook of*

comparative psychology: perception, learning, and cognition. pp. 411-439. APA.

Goodall, J. (1962). Nest building behavior in the free ranging chimpanzees. *Annals of the*

New York Academy of Sciences, 102 (2): 455-467.

Guislain, P., & Dupain, J. (2005). Scientific report: determinants of habitat use by sympatric

chimpanzee and gorilla populations at the periphery of the Dja Faunal Reserve, Cameroon.

Report for the L.S.B. Leakey Foundation, PGS Cameroon.

Gruber, T., Muller, M. N., Reynolds, V., Wrangham, R. W., & Zuberbühler, K. (2011).

Community-specific evaluation of tool affordances in wild chimpanzees. *Scientific*

Reports, *1,* 1-7.

Gruber, T., Luncz, L. V., Mörchen, J., Schuppli, C., Kendal R. L., & Hockings, K. J. (2019).

Cultural change in animals: a flexible behavioural adaptation to human disturbance.

Palgrave Communications 5, 1-9. doi: 10.1057/s41599-019-0271-4

Hernandez-Aguilar, R. A. (2009). Chimpanzee nest distribution and site reuse in a dry

habitat: implications for early hominin ranging. *Journal of Human Evolution*, *57*(4), 350-

364.

Hicks, T. C. (2010). A chimpanzee mega-culture? Exploring behavioral continuity in Pan

troglodytes schweinfurthii across northern DR Congo. *African Primates*, *7*(1), 1-18.

Hicks, T. C., Kühl, H. S., Boesch, C., Dieguez, P., Ayimisin, A. E., Fernandez, R. M., …

Roessingh, P. (2019). Bili-Uéré: A chimpanzee behavioural realm in Northern Democratic

Republic of Congo. *Folia Primatologica,* 90, 3-64.

Hobaiter, C., Poisot, T., Zuberbühler, K., Hoppitt, W., & Gruber, T. (2014). Social network

analysis shows direct evidence for social transmission of tool use in wild chimpanzees.

PloS Biology, 12(2), e1001960.

Izawa, K., & Itani, J. (1966). Chimpanzees in Kasakati Basin, Tanganyika (1) Ecological

study in the rainy season 1963–1964. *Kyoto University African Studies* 1, 73–156.

Kalan, A. K., Kulik, L., Arandjelovic, M., Boesch, C., Haas, F., Dieguez, P., … Kühl, H. S.

(2020). Environmental variability supports chimpanzee behavioural diversity. *Nature*

Communications, 11.1, 1-10. doi: 10.1038/s41467-020-18176-3

Koops, K., McGrew, W. C., de Vries, H., & Matsuzawa, T. (2012). Nest-building by

chimpanzees (*Pan troglodytes verus*) at Seringbara, Nimba Mountains: antipredation,

thermoregulation, and antivector hypotheses. *International Journal of Primatology*, 33,

356-380.

Koops, K., Humle, T., Sterck, E. H. M., & Matsuzawa, T. (2007). Ground-nesting by the

chimpanzees of the Nimba Mountains, Guinea: environmentally or socially determined?

American Journal of Primatology, 69, 407-419.

Kortlandt, A. (1992). On Chimpanzee Dormitories and Early Hominid Home Sites. *Current*

Anthropology, 33, 399-401.

Lacroux, C., Pouydebat, E., Rossignol, M., Durand, S., Aleeje, A., Asalu, E., ... & Krief, S.

(2022). Repellent activity against Anopheles gambiae of the leaves of nesting trees in the

Sebitoli chimpanzee community of Kibale National Park, Uganda. *Malaria journal*, 21(1),

1-11.

Laland, K. N., & Williams, K. (1998). Social transmission of maladaptive information in the

guppy. *Behavioural Ecology,* 9, 493-499.

Last, C., & Muh, B. (2013). Effects of human presence on chimpanzee nest location in the

Lebialem-Mone Forest landscape, Southwest Region, Cameroon. *Folia Primatologica,* 84,

51–63

Lethmate, J. (1977). Nest-building behaviour of a young Orang-utan raised in

isolation. *Primates*, *18*(3), 545-554.

Matsuzawa, T., & Yamakoshi, G. (1996). Comparison of chimpanzee material culture

between Bossou and Nimba, West Africa. In A. E. Russon, K. A. Bard, & S. T. Parker

(Eds.), *Reaching into thought: The minds of the great apes* (pp. 211–232). Cambridge

University Press.

McCarthy, M. S., Lester, J. D., & Stanford, C. B. (2017). Chimpanzees (Pan troglodytes)

flexibly use introduced species for nesting and bark feeding in a human-dominated

habitat. *International Journal of Primatology*, *38*(2), 321-337

McGrew, W. C. (2004). *The cultured chimpanzee: Reflections on cultural primatology*.

Cambridge: Cambridge University Press.

McGrew, W. C. (2010). Chimpanzee technology. *Science*, *328*(5978), 579-580.

McLennan, M. R., Hintz, B., Kiiza, B. V., Rohen, J., Lorenti, G. A., & Hockings, K. J.

(2020). Surviving at the extreme: chimpanzee ranging is not restricted in a deforested

human-dominated landscape in Uganda. *African Journal of Ecology*,

doi:10.1111/aje.12803

Mfossa, D. M., Gazagne, E., Gray, R. J., Ketchen, M. E., Abwe, E. E., Beudels-Jamar, R. C.,

... & Brotcorne, F. (2022). Montane Grassland Resources Drive Gorilla (Gorilla Gorilla)

Nesting Behaviours in the Ebo Forest, Littoral Region, Cameroon. *ResearchSquare* DOI:

https://doi.org/10.21203/rs.3.rs-2082431/v1

Plumptre, A. J., & Reynolds, V. (1997). Nesting behavior of chimpanzees: implications for

censuses. *International journal of primatology*, *18*(4), 475-485

Plumptre, A. J., & Cox, D. (2006). Counting primates for conservation: primate surveys in

Uganda. *Primates,* 47, 65-73.

Plumptre, A. J., Ayebare, S., Kujirakwinja, D., & Segan, D. (2020). Conservation planning

for Africa's Albertine Rift: conserving a biodiverse region in the face of multiple threats.

Oryx, 55, 302-310.

Prasetyo, D., Ancrenaz, M., Morrogh-Bernard, H.C., Utami Atmoko, S.S., Wich, S.A., van

Schaik, C.P. (2009). *Nest building in orangutans*. In: Wich, S.A., Utami Atmoko, S.S.,

Mitra Setia, T., van Schaik, C.P. Orangutans: geographic variation in behavioral ecology

and conservation. New York, US: Oxford University Press, 269-277.

Pruetz, J., Fulton, S. J., Marchant, L. F., McGrew, W. C., Schiel, M., & Waller, M. (2008).

Arboreal nesting as anti-predator adaptation by savanna chimpanzees (*Pan troglodytes*

verus) in southeastern Senegal. *American Journal of Primatology*, 70, 393-401.

Reynolds, V. (2005). *The chimpanzees of the Budongo forest: Ecology, behaviour and*

conservation. Oxford University Press, Oxford.

Samson, D. R., & Hunt, K. D. (2012). A thermodynamic comparison of arboreal and

terrestrial sleeping sites for dry-habitat chimpanzees (*Pan troglodytes schweinfurthii*) at

the Toro-Semliki Wildlife Reserve, Uganda. *American Journal of Primatology,* 4, 811-

818.

Samson, D. R., & Hunt, K. D. (2014). Chimpanzees preferentially select sleeping platform

construction tree species with biomechanical properties that yield stable, firm, but

compliant nests. *PLoS One,* 9, e95361.

Scully, E. J., Basnet, S., Wrangham, R. W., Muller, M. N., Otali, E., Hyeroba, D., …

Goldberg, T. L. (2018). Lethal respiratory disease associated with Human Rhinovirus C in

wild chimpanzees, Uganda, 2013. *Emerging Infectious Diseases,* 24, 267-274.

Schaller, G. E. (1963). The Mountain gorilla: Ecology and behavior. Oxford, England:

University of Chicago Press.

Stewart, F. A. (2011). *The evolution of shelter: ecology and ethology of chimpanzee nest*

building (Doctoral dissertation, University of Cambridge).

Stewart, F. A., & Pruetz, J. D. (2013). Do chimpanzee nests serve and anti-predatory

function? *American Journal of Primatology,* 75, 593-604.

Stewart, F. A., Piel, A. K., Azkarate, J. C., & Pruetz, J. D. (2018). Savanna chimpanzees

adjust sleeping nest architecture in response to local weather conditions. *American Journal*

of Physical Anthropology, *166*(3), 549-562.

Stokes, E. J., & Byrne, R. W. (2006). Effect of Snare Injuries on the Fig-Feeding Behavior of

Chimpanzees of the Budongo Forest, Uganda. In *Primates of western Uganda* (pp. 281-

297). Springer, New York, NY.

Sylla, S. F., Ndiaye, P., Lindshield, S. M., bogart, s. L., & pruetz, j. D. (2022). The western

chimpanzee (pan troglodytes verus) in the antenna zone (niokolo koba national park,

senegal): nesting ecology and sympatrics with other mammals. *Applied ecology and*

environmental research, 20(3), 2663-2681.

Suzuki, A. (1969). An ecological study of chimpanzees in a savanna

woodland. *Primates*, *10*(2), 103-148.

Tagg, N., Willie, J., Petre, C.A., Haggis, O. (2013). Ground night nesting in chimapnzees:

new insights from central chimpanzees (*Pan troglodytes troglodytes*) in south-east

Cameroon. *Folia Primatologica*, 84(6), 362-383.

Tutin CEG, Fernandez M (1984). Nationwide census of gorilla *(Gorilla g. gorilla)* and

chimpanzee *(Pan t. troglodytes)* populations in Gabon. *American Journal of Primatology,*

6, 313–336.

Tutin, C. E., Ham, R., & Wrogemann, D. (1995). Tool-use by chimpanzees (Pan t.

troglodytes) in the Lopé Reserve, Gabon. *Primates*, *36*(2), 181-192.

van Lawick-Goodall, J. (1968). The behaviour of free-living chimpanzees in the Gombe

Stream Reserve. *Animal Behaviour Monographs,* 1, 161–311.

van Casteren, A., Sellers, W. I., Thorpe, S. K., Coward, S., Crompton, R.H., Myatt, J. P., &

Ennos, A. R. (2012). Nest-building orangutans demonstrate engineering know-how to

produce safe, comfortable beds. *Proceedings of the National Academy of Sciences,* 109,

6873–6877.

Videan EN (2006) Bed-building in captive chimpanzees (*Pan troglodytes*): the importance of

early rearing. *American Journal of Primatology,* 68, 745–751.

Waller, J. C., & Reynolds, V. (2001). Limb injuries resulting from snares and traps in

chimpanzees (Pan troglodytes schweinfurthii) of the Budongo Forest,

Uganda. *Primates*, *42*(2), 135-139.

Wessling, E. G., & Surbeck, M. (2021). Failure to account for behavioral variability

overestimates bonobo populations and compromises accuracy in population

monitoring. *bioRxiv*.

Whiten, A., Goodall, J., McGrew, W. C., Nishida, T., Reynolds, V., Sugiyama, Y., …

Boesch, C. (1999). Culture in chimpanzees. *Nature*, 399, 682-685.

Wilson, M. L., Boesch, C., Fruth, B., Furuichi, T., Gilby, I. C., Hashimoto, C., ... &

Wrangham, R. W. (2014). Lethal aggression in Pan is better explained by adaptive

strategies than human impacts. *Nature*, *513*(7518), 414-417.

Figure 1: Location of nesting sites within the current area of research in the Bugoma
Central Forest Reserve. White = tree nests only; Grey = tree and ground nests together;
Black = Ground nests only. Inset shows full boundaries of Forest Reserve.

472
473

474 **Figure 2: Large primarily terrestrial mammals and reptiles present in the Bugoma**
475 **Central Forest Reserve that represent a potential threat to chimpanzees.** Pictures
476 captured directly or by camera trap within the home range of the Mwera community. From
477 Left to right: African elephant (*Loxodonta cyclotis)*; Rhinoceros viper (*Bitis nasicornis)*;
478 Golden cat (*Caracal aurata).*

479
480
481 **Figure 3: Human activities in the Bugoma Central Forest Reserve.** Pictures captured
482 directly or by camera trap in the home range of the Mwera community. Left to right: illegal
483 timber felling; overlapping presence of hunting dogs in key resource locations for
484 chimpanzees; use of wire and nylon snares for trapping small game.
485

486