CAAStools, a toolbox to identify and test Convergent Amino Acid Substitutions.

Fabio Barteri* [a,d], Alejandro Valenzuela* [a], Xavier Farré [g], David de Juan [a], Gerard Muntané [a,e,f], Borja Esteve-Altava [h] and Arcadi Navarro [a,b,c,d].

Authors affiliations

a.   IBE, Institute of Evolutionary Biology (UPF-CSIC), Department of Medicine and Life Sciences, Universitat Pompeu Fabra. PRBB, C. Doctor Aiguader N88, 08003 Barcelona, Spain
b.   Institució Catalana de Recerca i Estudis Avançats (ICREA) and Universitat Pompeu Fabra. Pg. Lluís Companys 23, 08010, Barcelona, Spain
c.   Center for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology, Av. Doctor Aiguader, N88, 08003 Barcelona, Spain
d.   BarcelonaBeta Brain Research Center, Pasqual Maragall Foundation, C. Wellington 30, 08005, Barcelona, Spain
e.   Hospital Universitari Institut Pere Mata, Institut d'Investigació Sanitària Pere Virgili (IISPV), Universitat Rovira i Virgili, Reus, Spain.
f.   Centro de investigación biomédica en red en salud mental (CIBERSAM), Spain.
g.   Genomes for Life-GCAT lab. Germans Trias i Pujol Research Institute (IGTP), Badalona, Spain
h.   European Molecular Biology Laboratory. Meyerhofstraße 1, 69117 Heidelberg, Germany

* Equal contribution

Abstract.

Background.
Upon phylogenetic data, coincidence of Convergent Amino Acid Substitutions (CAAS) with phenotypic convergences allow pinpointing genes that are likely to be associated with trait variation. Such findings can provide useful insights into the genetic architecture of complex phenotypes. Here we introduce CAAStools, a set of bioinformatics tools to identify and validate CAAS in orthologous protein alignments for pre-defined groups of species representing the phenotypic values targeted by the user.

Implementation and availability
CAAStools source code is available at http://github.com/linudz/caastools, along with documentation and examples.

Supplementary information

40   **Supplementary data are available.**

Introduction.

Convergent Amino Acid Substitutions (CAAS) can provide important insights into the
genetic changes underlying phenotypic variation (Zhang and Kumar, 1997; Ray *et al.*,
2015). Recent examples include the identification of genes potentially involved in marine
adaptation in mammals (Foote *et al.*, 2015), and the convergent evolution of
mitochondrial genes in deep-sea fish species (Sheng *et al.*, 2019). Notably, in 2018,
Muntané *et al.* identified a set of 25 genes involved in longevity in primates (Muntané *et*
al., 2018). A few years later, a similar analysis for a wider phylogeny retrieved 996 genes
associated with lifespan determination in mammals (Farré *et al.*, 2021). While these
analyses often need to be adapted for each particular phenotype and phylogeny, the CAAS
detection and validation strategies reported in the literature share some common steps
(Rey *et al.*, 2019). First, researchers select the species to compare for CAAS analysis, and
split them into two or more groups according to the phenotype. The criteria to select these
groups can be quite diverse. Groups can be formed, for instance, by species having
diverging values of a given continuous trait, or by species sharing different adaptations,
like terrestrial and marine mammals (Foote *et al.*, 2015). The second step consists in
linking amino acid substitutions with each group. For that step, different approaches can
be used, such as identifying identical substitutions for the same amino acid (Besnard *et*
al., 2009; Chabrol *et al.*, 2018), detecting topological incongruencies (Li *et al.*, 2008),
variations in amino acid profiles (Rodrigue *et al.*, 2010; Rey *et al.*, 2018), or relying on
consistent patterns of groups of amino acids in different groups of species (Zhang et al,
2014; Muntané *et al.*, 2018; Farré *et al.*, 2021). The third step consists in testing the
significance of the results. Molecular convergence is a noisy process, as apparent CAAS
may randomly take place under no selective force (Shahoua *et al.*, 2017). To overcome
this, researchers have adopted different strategies, mostly based on the idea that adaptive
CAAS tend to exceed convergent noise. The delta Site-Specific log-Likelihood Score
(ΔSSLS), for instance, is a method that consists in comparing the CAAS likelihood for
different phylogenetic topologies (Castoe *et al.*, 2009; Wang, Susko and Roger, 2012;
Parker *et al.*, 2013). Another approach is instead based on bootstrap resampling tests to
evaluate whether the number of detected CAAS is larger than expected by chance
(Muntané *et al.*, 2018; Farré *et al.*, 2021). Alternatively, some authors adopted a strategy
that consists in quantifying the convergent noise and focus on the detection of
Convergence on Conservative Sites (CCS) (Xu *et al.*, 2017; He *et al.*, 2020). In spite of
all these contributions, there is still no consensus approach. Some authors question
whether phenotypic convergence coincides with genome-wide molecular convergence
(Zou and Zhang, 2015b), or that adaptive substitutions outnumber random CAAS (Zou
and Zhang, 2015a; Gregg *et al.*, 2015). Further methodological developments can surely
benefit from the availability of software tools that are specifically designed to retrieve
CAAS, allowing to compare different validation strategies.

Here we present CAAStools, a toolbox to identify and validate CAAS. CAAStools is
based on the strategy applied in our previous studies (Muntané et al., 2018; Farré et al.,
2021), and implements different testing strategies through bootstrap analysis. CAAStools
is designed to be included in parallel workflows and is optimized to allow scalability at
proteome level.

Implementation.

CAAStools is a multi-modular python application organized into three tools. The outline of the suite is presented in **Figure 1**. The discovery tool is based on the protocol used in Muntané *et al*., 2018 and Farré *et al*., 2021. This approach identifies CAAS between two groups of species in an amino-acid Multiple Sequence Alignment (MSA) of orthologous proteins. These groups are named Foreground Group (FG) and Background Group (BG). Collectively, the two groups are called Discovery Groups (DG), as they represent the base for CAAS discovery. The CAAS identification algorithm scans each MSA and returns those positions that meet the following conditions: First, the FG and the BG species must share no amino acids in that position. Second, all the species in at least one of the two discovery groups (FG or BG) must share the same amino acid. The combination of these two conditions determines a set of different mutation patterns that the tool identifies as CAAS. Details on these patterns are provided in Supplementary Table 1.

Finally, CAAStools calculates the probability of obtaining a CAAS in a given position compared to randomized DGs, corresponding to the empirical p-value of the predicted CAAS in that position. This p-value represents a quantification of the convergent noise (Shahoua *et al.*, 2017) that is associated with a specific position. The details of this calculation are presented in Supplementary section 3. The Resample tool sorts species into *n* virtual DGs (re-samplings) for bootstrap analysis according to different combination strategies. This tool is thought to enable bootstrap analyses based on CAAS excess or likelihood (Muntané et al., 2018; Farré et al., 2021; Castoe *et al*., 2009). In a *Naive* modality, the probability of every species being included in a DG is considered identical and independent. This feature allows for bootstrap analyses aimed at quantifying convergent noise. However, species are phylogenetically related, biasing their probability of sharing a phenotype or amino acid. To address these phylogenetic dependencies CAAStools includes two other testing strategies. In the *Phylogeny-restricted* modality, the randomization can be restricted to some taxonomic orders or defined clades. These clades will match the ones of the species included in the DGs. In the *Brownian motion* modality, resampling is based on Brownian Motion simulations. This latter is inspired by the "permulation" strategy for trait randomization (Saputra et al., 2021), and its implementation relies on the *simpervec()* function from the RERconverge package (Kowalczyk *et al*., 2019). Finally, the *bootstrap* tool determines the iterations returning a CAAS for each position in a MSA to establish the corresponding empirical p-value for the detection of a CAAS in that position. Both the discovery and the bootstrap tools are designed to be launched on single MSAs, in order to allow the user to parallelize the workflow for large protein sets.

Usage and testing

We tested CAAStools on the dataset from Farré *et al*., (2021). The details of this test are reported in Supplementary 3. The full dataset is available in the /test folder within the CAAStools repository.

Figures

Figure 1. CAAStools layout. The three tools of the CAAStools suite rely on three pieces of information; a phylogenetic tree, the trait information, and an amino acid MSA. The discovery tool (A) detects the CAAS between two groups of species that are defined by the user on the basis of trait values. The resample tool (B) performs *n* trait resamplings in different modalities, on the bases of the phylogeny and the trait value distributions. The output of this resampling is processed by the bootstrap tool (C) that elaborates a bootstrap distribution from the MSA. All the tools can be executed independently.

Acknowledgements

We'd like to thank our collaborators at the Institute of Evolutionary Biology of Pompeu Fabra University for supporting us and sharing their ideas in scientific discussions, with particular mention of Dr. Tomas Marques Bonet and members of his group. This work was supported by the I+D+i project PID2021-127792NB-I00 funded by MCIN/AEI/10.13039/501100011033 (FEDER Una manera de hacer Europa)" and by "Unidad de Excelencia María de Maeztu", funded by the AEI (CEX2018-000792-M) and Departament de Recerca i Universitats de la Generalitat de Catalunya (GRC 2021 SGR 0467).

Author Contributions

Fabio Barteri (FB) has been in charge of the development of CAAStools. Alejandro Valenzuela (AV) carried on the beta-testing and code debugging. FB and AV have contributed equally to this manuscript. Xavi Farré (XF), Gerard Muntané (GM) and Arcadi Navarro (AN) designed the CAAS identification and validation protocol. XF wrote a set of scripts to identify CAAS that served as template for CAAStools implementation. AN and GM conceptualized the method. AN, David de Juan (DJ) and Borja Esteve-Altava (BEA) - along with GM - participated in the scientific discussion

156 and supervision of this project. DJ's contribution was particularly helpful for the
157 optimization of CAAStools code.

Bibliography

Besnard, Guillaume, A. Muthama Muasya, Flavien Russier, Eric H. Roalson, Nicolas
Salamin, y Pascal-Antoine Christin. «Phylogenomics of C4 Photosynthesis in
Sedges (Cyperaceae): Multiple Appearances and Genetic Convergence». Molecular
Biology and Evolution 26, n.º 8 (1 de agosto de 2009): 1909-19.
https://doi.org/10.1093/molbev/msp103.
Castoe, Todd A., A. P. Jason de Koning, Hyun-Min Kim, Wanjun Gu, Brice P. Noonan,
Gavin Naylor, Zhi J. Jiang, Christopher L. Parkinson, y David D. Pollock. «Evidence
for an ancient adaptive episode of convergent molecular evolution». Proceedings of
the National Academy of Sciences 106, n.º 22 (2 de junio de 2009): 8986-91.
https://doi.org/10.1073/pnas.0900233106.
Chabrol, Olivier, Manuela Royer-Carenzi, Pierre Pontarotti, y Gilles Didier. «Detecting
the Molecular Basis of Phenotypic Convergence». Methods in Ecology and
Evolution 9, n.º 11 (2018): 2170-80. https://doi.org/10.1111/2041-210X.13071.
Farré, Xavier, Ruben Molina, Fabio Barteri, Paul R H J Timmers, Peter K Joshi,
Baldomero Oliva, Sandra Acosta, Borja Esteve-Altava, Arcadi Navarro, y Gerard
Muntané. «Comparative Analysis of Mammal Genomes Unveils Key Genomic
Variability for Human Life Span». Molecular Biology and Evolution 38, n.º 11 (1 de
noviembre de 2021): 4948-61. https://doi.org/10.1093/molbev/msab219.
Foote, Andrew D., Yue Liu, Gregg W. C. Thomas, Tomáš Vinař, Jessica Alföldi, Jixin
Deng, Shannon Dugan, et al. «Convergent Evolution of the Genomes of Marine
Mammals». Nature Genetics 47, n.º 3 (marzo de 2015): 272-75.
https://doi.org/10.1038/ng.3198.
He, Ziwen, Shaohua Xu, Zhang Zhang, Wuxia Guo, Haomin Lyu, Cairong Zhong, David
E Boufford, Norman C Duke, The International Mangrove Consortium, y Suhua Shi.
«Convergent adaptation of the genomes of woody plants at the land–sea interface».
National Science Review 7, n.º 6 (1 de junio de 2020): 978-93.
https://doi.org/10.1093/nsr/nwaa027.
Kowalczyk, Amanda, Wynn K Meyer, Raghavendran Partha, Weiguang Mao, Nathan L
Clark, y Maria Chikina. «RERconverge: an R package for associating evolutionary
rates with convergent traits». Bioinformatics 35, n.º 22 (1 de noviembre de 2019):
4815-17. https://doi.org/10.1093/bioinformatics/btz468.
Li, Gang, Jinhong Wang, Stephen J. Rossiter, Gareth Jones, James A. Cotton, y Shuyi
Zhang. «The hearing gene Prestin reunites echolocating bats». Proceedings of the
National Academy of Sciences 105, n.º 37 (16 de septiembre de 2008): 13959-64.
https://doi.org/10.1073/pnas.0802097105.
Muntané, Gerard, Xavier Farré, Juan Antonio Rodríguez, Cinta Pegueroles, David A.
Hughes, João Pedro de Magalhães, Toni Gabaldón, y Arcadi Navarro. «Biological
Processes Modulating Longevity across Primates: A Phylogenetic Genome-
Phenome Analysis». Molecular Biology and Evolution 35, n.º 8 (1 de agosto de
2018): 1990-2004. https://doi.org/10.1093/molbev/msy105.
Parker, Joe, Georgia Tsagkogeorga, James A. Cotton, Yuan Liu, Paolo Provero, Elia
Stupka, y Stephen J. Rossiter. «Genome-Wide Signatures of Convergent Evolution
in Echolocating Mammals». Nature 502, n.º 7470 (octubre de 2013): 228-31.
https://doi.org/10.1038/nature12511.

Rey, Carine, Laurent Guéguen, Marie Sémon, y Bastien Boussau. «Accurate Detection of Convergent Amino-Acid Evolution with PCOC». Molecular Biology and Evolution 35, n.º 9 (1 de septiembre de 2018): 2296-2306. https://doi.org/10.1093/molbev/msy114.

Rey, Carine, Vincent Lanore, Philippe Veber, Laurent Guéguen, Nicolas Lartillot, Marie Sémon, y Bastien Boussau. «Detecting adaptive convergent amino acid evolution». Philosophical Transactions of the Royal Society B: Biological Sciences 374, n.º 1777 (22 de julio de 2019): 20180234. https://doi.org/10.1098/rstb.2018.0234.

Rodrigue, Nicolas, Hervé Philippe, y Nicolas Lartillot. «Mutation-selection models of coding sequence evolution with site-heterogeneous amino acid fitness profiles». Proceedings of the National Academy of Sciences 107, n.º 10 (9 de marzo de 2010): 4629-34. https://doi.org/10.1073/pnas.0910915107.

Saputra, Elysia, Amanda Kowalczyk, Luisa Cusick, Nathan Clark, y Maria Chikina. «Phylogenetic Permulations: A Statistically Rigorous Approach to Measure Confidence in Associations in a Phylogenetic Context». Molecular Biology and Evolution 38, n.º 7 (1 de julio de 2021): 3004-21. https://doi.org/10.1093/molbev/msab068.

Shen, Xuejuan, Zhiqing Pu, Xiao Chen, Robert W. Murphy, y Yongyi Shen. «Convergent Evolution of Mitochondrial Genes in Deep-Sea Fishes». Frontiers in Genetics 10 (2019). https://www.frontiersin.org/articles/10.3389/fgene.2019.00925.

Wang, Huai-Chun, Edward Susko, y Andrew J. Roger. «The Site-Wise Log-Likelihood Score Is a Good Predictor of Genes under Positive Selection». Journal of Molecular Evolution 76, n.º 5 (mayo de 2013): 280-94. https://doi.org/10.1007/s00239-013-9557-0.

Xu, Shaohua, Ziwen He, Zixiao Guo, Zhang Zhang, Gerald J. Wyckoff, Anthony Greenberg, Chung-I. Wu, y Suhua Shi. «Genome-Wide Convergence during Evolution of Mangroves from Woody Plants». Molecular Biology and Evolution 34, n.º 4 (1 de abril de 2017): 1008-15. https://doi.org/10.1093/molbev/msw277.

Zhang, Guojie, Cai Li, Qiye Li, Bo Li, Denis M. Larkin, Chul Lee, Jay F. Storz, et al. «Comparative genomics reveals insights into avian genome evolution and adaptation». Science 346, n.º 6215 (12 de diciembre de 2014): 1311-20. https://doi.org/10.1126/science.1251385.

Zhang, J., y S. Kumar. «Detection of Convergent and Parallel Evolution at the Amino Acid Sequence Level». Molecular Biology and Evolution 14, n.º 5 (mayo de 1997): 527-36. https://doi.org/10.1093/oxfordjournals.molbev.a025789.

Zou, Zhengting, y Jianzhi Zhang. «Are Convergent and Parallel Amino Acid Substitutions in Protein Evolution More Prevalent Than Neutral Expectations?» Molecular Biology and Evolution 32, n.º 8 (1 de agosto de 2015): 2085-96. https://doi.org/10.1093/molbev/msv091.

———. «No Genome-Wide Protein Sequence Convergence for Echolocation». Molecular Biology and Evolution 32, n.º 5 (1 de mayo de 2015): 1237-41. https://doi.org/10.1093/molbev/msv014.

CAAStools, a toolbox to identify and test Convergent Amino Acid Substitutions.

Supplementary information.

Supplementary 1. CAAS Discovery algorithm.

Given two Discovery Groups (DGs, Foreground and Background groups, FG and BG, respectively), the discovery tool recognizes as CAAS all those substitutions that meet two requirements. Let A be an MSA of q sequences of length t. We can describe A as an array of t positions [1]. Each position (pos_i) will consist of a set of N different amino acids, a, with absolute frequency (or count), f, where NS is the total number of symbols in the alignment [1].

[1] $A = (pos_1\ pos_2\ \dots\ pos_t)\ ;\ pos_i = a_1\ f_1\ a_2\ f_2\ \dots\ a_{NS}\ \dots\ f_{NS}\ ;\ \sum_1^{NS}\ f = q$

The FG and the BG are formalized as sets of different species s_{FG} and s_{BG}, with no intersection and size l_{FG} and l_{BG} [2].

[2] $s_{FG} = \left(s_1\ s_2\ \dots\ s_{l_{FG}}\right)\ ;\ s_{BG} = \left(s_1\ s_2\ \dots\ s_{l_{BG}}\right)$

$s_{FG} \cap s_{BG} = \emptyset$

In each alignment position, s_{FG} and s_{BG} are associated with two sets of amino acids, $fg(pos_i)$ and $bg(pos_i)$, with length w_{FG} and w_{BG}.

[3] $fg(pos_i) = a_1\ f_1\ a_2\ f_2\ \dots\ a_{w_{FG}}\ \dots\ f_{w_{FG}}\ ;\ bg(pos_i) = a_1\ f_1\ a_2\ f_2\ \dots\ a_{w_{BG}}\ \dots\ f_{w_{BG}}$

CAAStools identifies a CAAS when three conditions are met [4]. First, the two groups must share no amino acids. This means that all the species in the FG need to have different AAs than the species in the BG. Second, at least one of the two DGs must share (or "converge to") the same amino acid. Also, the CAAS is detected if at least one amino acid is associated to both DGs

[4] $CAAS_i\{fg(pos_i) \cap bg(pos_i) = \emptyset\ w_{FG} == 1\ or\ w_{BG} == 1\ w_{FG} > 0\ and\ w_{BG} > 0$

The combination of these three rules defines 3 different mutation *patterns*. We define *pattern 1* when the DGs converge to two different amino acids ($w_{FG} = 1$; $w_{BG} = 1$). The *pattern 2* will be verified as the FG converges to one amino acid, but the BG will be associated with different amino acids ($w_{FG} = 1$; $w_{BG} > 1$). *Pattern 3* will consist in the opposite situation, or else when the FG is associated with different amino acids, whilst the BG converges to a single amino acid ($w_{FG} > 1$; $w_{BG} = 1$). **Supplementary Table**

295 **1** summarizes the different mutation patterns and the meeting of requirements for CAAS
296 identification.
297
298 ***Supplementary Table 1.*** *Mutation patterns and associated program decisions on CAAS*
299 *assignment.*

Discovery Groups		Difference between DGs	Convergence in		Pattern
FG	**BG**		**FG**	**BG**	
KV	K	NO	NO	YES	Not a CAAS (No difference)
M	TM	NO	YES	NO	Not a CAAS (No difference)
MK	VE	YES	NO	NO	Not a CAAS (No convergence)
K	V	YES	YES	YES	**Pattern 1** (Both convergent)
K	VM	YES	YES	NO	**Pattern 2** (FG convergent, BG multiple)
KE	W	YES	NO	YES	**Pattern 3** (FG multiple, BG convergent)

300
301

Supplementary 2. CAAS discovery statistical testing

303
CAAStools calculates an empirical p-value for each CAAS prediction. This p-value is
equal to the probability of obtaining a CAAS with random species, and under the same
conditions as the CAAS discovery (size of the DGs, maximum permitted gaps and
missing species). Following the MSA description in [1], we'll consider a couple of DG
(*FG* and *BG*) of size l_{FG} and l_{BG}, as formalized in [2]. The probability to obtain a CAAS
from random species is calculated as the probability of extracting concomitantly k_{FG}
and k_{BG} objects from a population of size N over a number of extractions *n*, provided the
conditions in [4], i.e. $k_{FG} \cap k_{BG} = \emptyset$ and $wk_{FG} == 1 \ or \ wk_{BG} == 1$ where *wk* is the
number of symbols in the resampling *k*. This probability can be calculated through the
probability mass function from the hypergeometric distribution [5].
314

315 [5] $P(k) = \dfrac{\binom{K}{k}\binom{N-k}{n-k}}{\binom{N}{n}} = Hyp(N, K, k, n)$

316     $P(CAAS) = P(FG) * P(BG)$
317

318 $\begin{cases} P(FG) = Hyp(N_{FG}, K_{FG}, k_{FG}, n_{FG}) \\ \quad N_{FG} = q \ ; \\ \quad k_{FG} = l_{FG} - null_{FG} \\ \quad n_{FG} = k_{FG} \end{cases}$

320 $\begin{cases} P(BG) = Hyp(N_{BG}, K_{BG}, k_{BG}, n_{NG}) \\ \quad N_{BG} = q - l_{FG} \\ \quad k_{BG} = l_{BG} - null_{BG} \\ \quad n_{BG} = k_{BG} \end{cases}$

321
Note that the size of the population N in *P(FG)* differs from the one considered in *P(BG)*.
In the first case, the probability of obtaining a convergence in the *FG* is calculated on the
total number of sequences in the alignment. In *BG*, the size considered is the difference
between the total number of sequences in the alignment *q* and the size of the other group
$(q - l_{BG})$, $(q - l_{FG})$, as the two events are concomitant but not independent. The
number of extractions k_{FG} and k_{BG} are equal to the number of the difference between the
size of the DGs and the number of indels and missing species allowed by the user (*null*).
The terms K_{FG} and K_{BG} represent the number of successes in the population. In [6], [7]

and [8], we see how this value can be calculated considering all the possible combinations of amino acid symbols that meet the requirements for CAAS detection [4].

[6] $\quad C_{P1,2} = \{K_{FG} = [f_j]; \, K_{BG} = [q - f_j] \, \forall \, a_j \in pos_i\}$

$\qquad C_{P1,2} = [(K_{FG_1}; K_{BG_1}), (K_{FG_2}; K_{BG_2}) \dots (K_{FG_z}; K_{BG_z})]$

[7] $\quad C_{P1,3} = \{K_{FG} = [q - f_j]; \, K_{BG} = [f_j] \, \forall \, a_j \in pos_i\}$

$\qquad C_{P1,3} = [(K_{FG_1}; K_{BG_1}), (K_{FG_2}; K_{BG_2}) \dots (K_{FG_z}; K_{BG_z})]$

[8] $\quad C_{P1} = \{K_{FG} = [f_j]; \, K_{BG} = [f_h] \, \forall \, a_j, a_k \in pos_i\}$

$\qquad C_{P1} = [(K_{FG_1}; K_{BG_1}), (K_{FG_2}; K_{BG_2}) \dots (K_{FG_z}; K_{BG_z})]$

These combinations are based on patterns (P). Note that $C_{P1,2}$ and $C_{P1,3}$ overlap, and that the intersection coincides with C_{P1}. We can now calculate the CAAS probability separately for each pattern [9].

[9] $\quad P(CAAS_{P1,3}) = \sum_{x=1}^{z} \quad Hyp(N_{FG}, K_{FG_x}, k_{FG}, n_{FG}) *$
$Hyp(N_{BG}, K_{BG_x}, k_{BG}, n_{NG})$

$\qquad P(CAAS_{P1,2}) = \sum_{x=1}^{z} \quad Hyp(N_{FG}, K_{FG_x}, k_{FG}, n_{FG}) *$
$Hyp(N_{BG}, K_{BG_x}, k_{BG}, n_{NG})$

$\qquad P(CAAS_{P1}) = \sum_{x=1}^{z} \quad Hyp(N_{FG}, K_{FG_x}, k_{FG}, n_{FG}) *$
$Hyp(N_{BG}, K_{BG_x}, k_{BG}, n_{NG})$

The probability to obtain a CAAS in position pos_i is hence calculated as it follows:

[10] $\quad pvalue_{pos_i} = P(CAAS_{pos_i}) = P(CAAS_{P1,3}) + P(CAAS_{P1,2}) - P(CAAS_{P1})$

2.1 Correction for discovery groups of equal size.

If the species found in the alignment are the same for FG and BG sizes ($l_{FG} = l_{BG}$), the probability of retrieving pattern 2 and pattern 3 are equal. In this case, the p-value is equal to the $P(CAAS_{P1,2})$.

[11] $\quad \begin{cases} pvalue_{pos_i} = P(CAAS_{pos_i}) = P(CAAS_{P1,2}) \\ \qquad\qquad if \, l_{FG} = l_{BG} \end{cases}$

Supplementary 3. CAAS discovery from Farré et al., 2021.

As a test run for CAAStools, we repeated the CAAS discovery from the results published by Farré et al., in 2021 and entitled "*Comparative Analysis of Mammal Genomes Unveils Key Genomic Variability for Human Life Span*" (DOI: 10.1093/molbev/msab219). In this work, 13,035 MSA from UCSC public database (https://genome.ucsc.edu/, accessed August, 2019) were scanned to find CAAS between two groups of species with divergent maximum lifespan. The "long lived" group is formed by *Homo sapiens* (hg38), *Nomascus leucogenys* (nomLeu3), *Heterocephalus glaber* (hetGla2), *Myotis davidii* (myoDav1), *Myotis lucifugus* (myoLuc2), *Eptesicus fuscus* (eptFus1). The "short lived" group is

formed by *Mesocricetus auratus* (mesAur1), *Rattus norvegicus* (rn6), *Pantholops*
hodgsonii (panHod1), *Sorex araneus* (sorAra2), *Condylura cristata* (conCri1),
Monodelphis domestica (monDom5). Farré et al., filtered the results from CAAS
discovery to those CAAS having no gaps or missing species, and focused their analysis
on the CAAS of scenarios 1 and 2, which correspond to patter 1 and 2 in CAAStools
terminology.
382
383   We have repeated this analysis under the same conditions, filtering for pattern 1 and 2
384   and for no gaps in foreground (*short-lived* group) and background (*long-lived* group). The
385   results (*Supplementary dataset 1*) and the phenotype configuration (*Supplementary*
386   *dataset 2*) are available in the supplementary.material.xls spreadsheet. Our analysis
387   confirmed the identification of 2737 mutations in 2004 MSA.
388

Supplementary 4. An example of p-value calculation from *Farré et al.* dataset.

389
390
391
The gene AKAP13 (RefSeq code: NM_006738) is part of the dataset published by Farré
et al. in 2021. In that analysis, authors selected only those positions that were associated
with no gaps in both Foreground and Background, obtaining 3 CAAS from this gene.
Here, we repeated the CAAS detection without any filtering for gaps or missing species.
The result is shown in *Supplementary Table 2*. The extended CAAStools discovery
output can be found in *Supplementary Dataset 3*.
398
399   ***Supplementary Table 2****: Resume of the CAAStools discovery analysis for AKAP13 with*
400   *phenotype information from Farré et al., 2021.*
401

Gene	Position	Substitution	P-value	Pattern	Gaps FG	Gaps BG
AKAP13	392	T/AIS	0.0009037398709	2	0	1
AKAP13	451	M/GLST	0.001177073702	2	0	1
AKAP13	577	A/INST	0.0005992501486	2	0	1
AKAP13	657	D/AEGSV	0.0002559580225	2	0	0
AKAP13	**931**	**V/AMRT**	**0.0001223275131**	**2**	**0**	**0**
AKAP13	1180	D/EPRS	0.007729389679	2	3	0
AKAP13	1185	V/ADP	0.0152264739	2	2	3
AKAP13	1204	D/AENP	0.001044565793	2	1	1
AKAP13	1210	G/CDV	0.0121208014	2	3	3
AKAP13	1212	R/DK	0.02023676039	2	3	3
AKAP13	1213	E/HKQ	0.01388995848	2	3	3
AKAP13	**1219**	**GP/V**	**0.1087036943**	**3**	**3**	**4**
AKAP13	1298	K/EHQST	0.0001593229323	2	0	0
AKAP13	1315	E/GIKR	0.0008881559737	2	1	1
AKAP13	1827	S/GN	0.00135352871	2	0	1
AKAP13	2685	R/CHQ	0.004566143925	2	1	2
AKAP13	2692	H/NRY	0.0007280567925	2	0	1

402
In order to show the details of the p-value calculation through the hypergeometric
function, we decided to focus on the position 931 and 1219. Both positions present a

CAAS, but their p-values are largely different. In position 931, the p-value is
0.0001223275131, while the p-value associated with position 1219 is 0.1087036943. The
details of the p-value calculations are presented in *Supplementary Table 3*. For each
amino acid combination, the algorithm calculates the probability to obtain a convergence
in the *FG* (*P(FG)*) and the probability to obtain a *BG* with different amino-acids (*P(BG)*).
410
411 **Supplementary Table 3.** *P-value calculation for positions 931 and 1219 in AKAP13 gene*
412 *(NM_006738).*
413

P 931	FG	BG	P(FG)	P(BG)	P(CAAS)	Patterns
	V	L,G,T,A,R,M	0.0731423047	0.00167128469	0.000122241614	1,2
	A	L,V,G,T,R,M	1.14E-07	0.8928571429	1.02E-07	1,2
				P-value	**1.22E-04**	

P 1219	FG	BG	P(FG)	P(BG)	P(CAAS)	Patterns
	A	I,P,S,G,V,T	0.03024193548	0.5310344828	0.01605951057	1,2
	I,P,S,G,V,T	A	0.2822580645	0.1264367816	0.03568780126	1,3
	P	I,A,S,G,V,T	0.06671554252	0.3931034483	0.02622610982	1,2
	I,A,S,G,V,T	P	0.1776026393	0.2091954023	0.03715365558	1,3
	V	I,A,P,S,G,T	0.0007331378299	0.9333333333	0.0006842619746	1,2
	I,A,P,S,G,T	V	0.6697214076	0.01379310345	0.009237536657	1,3
	A	P	0.03024193548	0.2091954023	0.00632647386	1
	A	V	0.03024193548	0.01379310345	0.0004171301446	1
	P	V	0.06671554252	0.01379310345	0.0009202143796	1
	P	A	0.06671554252	0.1264367816	0.00843529848	1
	V	A	0.0007331378299	0.1264367816	9.27E-05	1
	V	P	0.0007331378299	0.2091954023	0.0001533690633	1
				P-value	**0,1087**	

Inputs

Outputs