Can we improve our ability to interpret ectotherm thermal tolerance?

Jacinta D. Kong[1*], Nicholas C. Wu[2]

[1] *School of Natural Sciences, Trinity College Dublin, Dublin 2, Ireland*

[2]*Hawkesbury Institute for the Environment, Western Sydney University, NSW 2753, Australia*

Keywords: amphibia, critical temperature maxima, life stage, macrophysiology, ontogeny

***Correspondence**: jacintadkong@gmail.com

ORCID: JDK (0000-0002-1085-8612), NCW (0000-0002-7130-1279)

Author contributions: JDK: Conceptualization, Investigation, Data Curation, Software, Writing - Original Draft. NCW: Resources, Software, Formal analysis, Visualization. Both authors reviewed and edited the work and gave final permission for submission.

Competing Interest: The authors declare no competing interests.

Data availability statement: Data and code are available on Zenodo (https://github.com/jacintak/frog_ctmax)

Article type: Research article

Acknowledgements: The authors would like to thank researchers C. Filho (Universidade Estadual de Santa Cruz) and R. Percinz (Ruth Percino) who provided data for this study.

Abstract

Thermal tolerances, such as critical temperatures, are important indices for understanding an organism's vulnerability to changing environmental temperature. Differences in thermal tolerance over ontogeny may generate a 'thermal bottleneck' that sets the climate vulnerability for organisms with complex life cycles. However, a species' microhabitat preference and methodological differences among studies can generate confounding variation in thermal tolerance that may mask trends in large-scale comparative studies and may hinder our ability to assess climate change vulnerability within and among species. Here, we evaluated two approaches to resolving ontogenetic and environmental drivers of thermal tolerance and methodological variation: mathematical standardisation of thermal tolerance and classifying microhabitat preferences. Using phylogenetically informed, multi-level models with a global dataset of upper critical temperatures from 438 Anuran species, we found ontogenetic trends in thermal tolerance were similar across microhabitat preferences and standardising critical temperatures against common methodological variation had little impact on our conclusions. Our results suggested thermal bottlenecks are not strongly present in Anurans but instead, implied strong developmental or genetic conservatism of thermal tolerance within families and ecotypes. We discussed

36   considerations for resolving confounding variation to interpret thermal tolerance at a
37   macrophysiological scale.

Introduction

Thermal tolerance metrics, such as critical temperatures, have endured in ecological studies
despite their criticisms because they are simple to conceptualise (e.g. impediment to
movement), are generalisable across taxa, and can be readily assayed in the laboratory
(Sinclair et al., 2016). These metrics facilitate our ability to link environmental temperature
to organismal responses (Bates & Morley, 2020), synthesise general trends in physiological
diversity from local to macro scales (Chown & Gaston, 2016), and index vulnerability or
sensitivity to climate change via metrics such as thermal safety margins (Deutsch et al.,
2008). For example, differences in upper critical temperature (CT_{max}) across life stages may
indicate a thermal bottleneck at the most temperature-sensitive life stage (Dahlke, Wohlrab,
Butzin, & Pörtner, 2020; Pandori & Sorte, 2019). For organisms with a complex life cycle
such as frogs, juvenile or larval stages in aquatic environments are generally expected to
have lower thermal tolerance and thus be more vulnerable to climate change than adult
stages (Lenoir, Hattab, & Pierre, 2017).

Identifying a thermal bottleneck represents a significant amount of work that is limited to
commonly studied species (e.g., Ruthsatz, Dausmann, Peck, and Glos (2022)). Thermal
bottlenecks can also be identified using a macrophysiological approach which has described
stage-specific differences in thermal sensitivities and tolerances in marine invertebrates
(Collin, Rebolledo, Smith, & Chan, 2021; Pandori & Sorte, 2019; Truebano, Fenner, Tills,
Rundle, & Rezende, 2018), fish (Dahlke et al., 2020; Moyano et al., 2017), insects (Kingsolver
& Buckley, 2020; Klockmann, Günter, & Fischer, 2017) and reptiles (Levy et al., 2015).
Among species comparisons in frogs are often limited to a single geographic region and are
often limited to either adults or larva even if both are collected from the same region (Carilo
Filho et al., 2021; Cheng et al., 2022; Pintanel et al., 2022; Pintanel, Tejedo, Ron, Llorente, &
Merino-Viteri, 2019; Ruthsatz et al., 2022). Thus, whether thermal bottlenecks occur and
how consistent they are across Anuran species is not well understood. A macrophysiological
analyses of thermal tolerance throughout ontogeny will vastly improve our understanding of
the presence and generality of thermal bottlenecks across an environmentally sensitive
taxonomic group.

Comparative macrophysiological analyses present two key challenges towards assessing and
interpreting thermal tolerances. Microhabitat preferences and among-study variation may
generate confounding variation in thermal tolerances that may mask or accentuate the
presence of a thermal bottleneck. Here, we analysed the CT_{max} of 438 Anuran species to 1)
identify whether Anurans have a thermal bottleneck in their life cycle and their potential
evolutionary drivers and ecological consequences at a macrophysiological scale, and 2)
evaluate whether proposed solutions to two challenges of comparative analyses of
ectotherm thermal tolerance improved our ability to identify thermal bottlenecks. We
acknowledge there is a diversity of definitions and uses of thermal tolerance that would
influence their interpretation. In this study, we only consider CT_{max} as the end point of a
temperature ramping assay with a constant heating rate, or dynamic ramping protocol, and

we define heating tolerance as the temperature range between starting temperature and
CT_{max} during the ramping assay.

Do microhabitat preferences improve identification of thermal bottlenecks?

The key challenge to interpret thermal tolerance over the life cycle is to disentangle
confounding variation deriving from microhabitat selection from inherent developmental or
physiological drivers. A diversity of microhabitats may mask broader adaptive covariation
between thermal tolerance and microclimate selection to explain, in part, an observed lack
of variation in CT_{max} at a global scale or drive strong associations with thermal tolerance
(González-del-Pliego et al., 2020; Pincebourde & Woods, 2020). However, distinguishing
between these patterns requires corresponding microclimate information for each
ontogenetic stage that is not always collected for thermal tolerance studies or are available,
but see Pintanel et al. (2022). If differences in thermal environments are consistent within
broad microhabitat preferences in Anurans, then microhabitat preferences may be a
reliable proxy for local environmental selection on thermal tolerance. Among species, frogs
show strong microhabitat preferences that are associated with phenotypic syndromes
encompassing behaviour, morphology and physiology and can broadly be categorised into
ecological types ("ecotype") (Moen, Cabrera-Guzman, Caviedes-Solis, Gonzalez-Bernal, &
Hanna, 2022; Moen & Wiens, 2017). In turn, the animal's ecotype may mediate the
confounding interaction between life stages and their respective microenvironment (Carilo
Filho et al., 2021; Pincebourde & Casas, 2015). We tested whether ecotypes are reliable
proxies of environmental adaptation to resolve ontogeny by environment interaction of
CT_{max} and identify thermal bottlenecks.

Does removing nuisance heterogeneity change thermal tolerance trends?

A large literature on ectotherm thermal tolerances has emerged because of the duality of
critical temperatures as characteristics of a thermal performance curve or as physiological
endpoints of a thermal regime (Bates & Morley, 2020; Sinclair et al., 2016). For comparative
purposes, protocols to assay thermal tolerance are generally not standardised across taxa
for either definition, and variation in the starting temperature and heating rate used will
affect the measured CT_{max} among individuals in a ramping assay (Peck, Clark, Morley,
Massey, & Rossetti, 2009; Terblanche, Deere, Clusella-Trullas, Janion, & Chown, 2007). Such
methodological variation or "nuisance heterogeneity" generates uncertainty in thermal
tolerances that may mask signals of interest (Noble et al., 2022). The significant nuisance
heterogeneity of CT_{max} complicates how accurately this metric can be translated from the
laboratory to the field regardless of any ecological relevance and undermines prediction
accuracy and interpretation of climate vulnerability metrics with confidence (Pinsky, Eikeset,
McCauley, Payne, & Sunday, 2019; Terblanche et al., 2011). One solution is to correct for
variation induced by methodology to a standard protocol (Cupp, 1980; Lebenzon, Des
Marteaux, & Sinclair, 2020). Resolving confounding variation will improve our ability to link
environmental gradients to trait variation and test hypotheses of thermal tolerance
evolution (Bates & Morley, 2020; Kovacevic, Latombe, & Chown, 2019). We applied a
mathematical model that standardises thermal tolerances to the same heating rate and test

whether standardising methodological variation in thermal tolerance metrics improves our
ability to resolve biological trends of interest while accounting for among-study variation in
methodology (Kong et al., 2022).

Methods

Data compilation

We started with a published dataset of upper critical temperatures of Anurans complied
from the literature (1962 – 2015) (Rohr et al., 2018). We supplemented this data with
studies from 2016 to 2022 using Web of Science (accessed 25th February) and the keywords
amphibian* AND ("critical thermal max*" OR CT_{max} OR "thermal toleran*" OR "critical
temp*" OR "upper temp*")". The terms returned 90 studies that were screened for
inclusion by one person (J.D.K.). Studies that measured CT_{max} using ramping assays were
retained. Studies that estimated CT_{max} from fitted thermal performance curves or compiled
values from literature were excluded. Only control animals were included from studies that
used multiple stressors (e.g. dehydration or infection). The screening process yielded 36
papers that provided extractable data. Provided data were used over published values
where possible. Populations were considered independent observations. Multiple endpoints
of critical temperatures on an individual were recorded but only one CT_{max} was used per
individual (loss of righting response where possible). We added studies that did not return
from keywords from forward searching. We distinguished between temperatures animals
were held at prior to experimentation (acclimation temperature) and the starting
temperature of the ramping assay. We assume that observed CT_{max} reflected their
acclimated CT_{max} as acclimation in frogs is reported on scales of hours (Brattstrom, 1970).

We obtained ecological types (hereby ecotype), from all Anuran species listed on the
International Union for Conservation of Nature's Red List of Threatened Species (IUCN,
2022). Ecotypes were classified from ecological descriptions from IUCN based on adult
behaviour and microhabitat preferences outside the breeding season (Moen & Wiens,
2017). We grouped species by ecotype even if the larval stage may have differing ecological
types (e.g. lotic vs lentic systems). This is because the ecotype diversity of the larvae stage is
broader than adults and is less described in the detail from the literature. The resulting
dataset represented 32 families spanning the globe (Fig. S1). Thermal tolerance records
were first matched to the ecotype dataset by binomial names used in the original source,
then unmatched records were joined by IUCN names based on synonyms accepted by IUCN
or Catalogue of Life. CT_{max} records that could not be matched with ecotype (56
observations) were excluded.

Standardisation of thermal tolerance metrics

The starting temperature and heating rate used affects the measured CT_{max} among
individuals in a ramping assay (Peck et al., 2009; Terblanche et al., 2007). The temperature
range between the start and end of a heating assay with a constant heating rate, the
heating tolerance, describes an elapsed duration of time which scales with temperature
similarly to heating tolerance. Heating rate has a consistent and predictable effect on the

effective duration of a heating assay, which accounts for a non-linear rate-temperature
relationship, that in turn influences heating tolerance (Kong et al., 2022). This predictable
relationship between heating rate, heating duration and consequently heating tolerance can
be used to standardise heating tolerance to a given heating rate. To standardise CT_{max} to a
heating rate of $1°C$ min^{-1}, we rescale heating tolerance into the effective duration of a
heating assay, apply the correction for the effect of heating rate on effective heating
duration, then rescale corrected heating duration back into a corrected heating tolerance
(detailed in the Supplementary Information). A heating rate of $1°C$ min^{-1} was the most
common heating rate and is considered ecologically relevant (e.g. (Leong, Tsang, & Guenard,
2022)). We did not standardise starting temperature because temperatures used may have
reflected local or field temperatures and indicate local adaptation. There were 1318
observations from 438 species with CT_{max}, 1316 observations from 438 species with heating
tolerance and 1238 observations from 432 species with corrected CT_{max} or corrected heating
tolerance. In effect, we reduced potential methodological "nuisance heterogeneity" that
may occur within and among ramping assays while retaining potential adaptive variation
attributed with starting temperature that may reflect acclimation or developmental,
physiological, genetic, or environmental constraints.

Relationship between CT_{max} and heating tolerance with ecotype and ontogeny

We constructed a phylogenetically corrected multi-level Bayesian model to investigate the
relationship between CT_{max}, corrected CT_{max}, heating tolerance or corrected heating
tolerance across ontogenetic stage and ecotype (see Supplementary Information for detail).
We included starting temperature of the assay as a co-variate. Ontogenetic stages 1-5
correspond to larva, 6 to juveniles, and 7 to adults (Ruthsatz et al., 2022). We used an
amphibian phylogeny dataset comprising of 7,238 species to account for phylogenetic
relatedness (Jetz & Pyron, 2018). We pruned the phylogenetic tree to matched binomial
names in the thermal tolerance data (438 species, Fig. 1). Species names based on Jetz and
Pyron (2018) were used in analysis to account for duplicate names matching with the IUCN.
Records that were not in the phylogenetic tree (n = 1) were excluded. We converted the
pruned tree to a phylogenetic relatedness correlation matrix for subsequent analysis. For
the CT_{max} model, we used a quadratic equation to model change in CT_{max} over ontogeny and
the quadratic equation provided a better fit than the linear model (Table S1). All model
outputs (Tables S2 – S5) and posterior predictive checks (Figs. S4 & S6) are presented in the
Supplementary Information. All model parameters are reported as mean Bayesian posterior
estimates ± 95% credible intervals (95% CI), which is equivalent to 95% confidence intervals
for frequentist statistics. Summary statistics are presented as mean ± standard deviation.

Figure 1. Phylogenetic patterns of CT$_{max}$ across 438 Anurans based on Jetz & Pyron (2018). Colours represents species mean CT$_{max}$. Example families are indicated by outer black lines. Representative silhouettes obtained from phylopic.org and background colours within the circles represent mean CT$_{max}$ at the Family level.

Results

CT_max

CT$_{max}$ ranged between 20.4 °C and 46.1 °C (mean 37.1 ± 4.1 °C) across all ecotypes. On average, fossorial frogs have the highest mean CT$_{max}$ (40.1 ± 3.1 °C) and semi-aquatic frogs have the lowest CT$_{max}$ (35.4 ± 3.54 °C). Larval CT$_{max}$ (Stage 1-5) was on average 3.5 °C higher than adult (Stage 7) and juvenile (Stage 6) CT$_{max}$ combined; the mean difference ranged from 1.5 °C for semi-aquatic frogs and 5.2 °C for stream-dwelling frogs. However, data on CT$_{max}$ for early larval (Stage 1) and juvenile (Stage 6) stages were low (n = 41 and n = 33, respectively) relative to later larval stages (Stage 3, n = 207) and adults (Stage 7, n = 733).

Variation in CT$_{max}$ was strongly influenced by ontogeny where CT$_{max}$ generally decreased across ontogeny (Table S2; Fig. 2). CT$_{max}$ did not differ substantially between ecotypes, but on average, stream-dwelling frogs have lower CT$_{max}$ (-0.74 [-2.63: 1.11]), while fossorial frogs generally had higher CT$_{max}$ (1.96 [-0.06:4.05]). There was a strong phylogenetic effect on CT$_{max}$ (λ = 0.85 [0.75:0.92]). Starting temperature also had a strong effect on CT$_{max}$, where the higher the starting temperature, the higher the CT$_{max}$ (0.16 [0.14:0.18]).

Figure 2. CT$_{max}$ (°C) decreases over ontogeny (stage 1–7) across all Anuran ecotypes (438 species). Quadratic regression line and shaded area represents ecotype-specific model estimate ± 95% CI.

Corrected CT$_{max}$

CT$_{max}$ corrected to 1°C min^{-1} was on average 1.1 °C higher (range difference between -1.6 – 6.8 °C) than observed CT$_{max}$ (Fig. S3). Variation in corrected CT$_{max}$ was strongly influenced by ontogeny where CT$_{max}$ generally decreased across ontogeny (Table S3; Fig. 3). Corrected CT$_{max}$ between ecotypes showed similar trends to the uncorrected CT$_{max}$, where stream-dwelling frogs had lower CT$_{max}$ (-0.2 [-2.37: 1.95]), while fossorial frogs had higher CT$_{max}$ (2.08 [-0.31:4.51]). There was a strong phylogenetic effect on corrected CT$_{max}$ (λ = 0.85 [0.74:0.92), and starting temperature also influenced corrected CT$_{max}$ (0.15 [0.13:0.17]).

Figure 3. Corrected CT$_{max}$ (°C) decreases over ontogeny (stage 1–7) across all ecotypes (432 species). Quadratic regression line and shaded area represent ecotype-specific model estimate ± 95% CI. CT$_{max}$ was corrected to a heating rate of 1 °C min^{1}.

Heating tolerance

Heating tolerances ranged between 2 °C and 36.5 °C (16.7 ± 6.34 °C) across all ecotypes. Mean heating tolerance varied between ecotypes with the highest mean in fossorial frogs (18.6 ± 6.1 °C) and lowest in stream-dwelling frogs (16.0 ± 6.2 °C). Heating tolerance was also influenced by ontogenetic stage (-0.42 [-0.49:-0.35]; Fig. 4), with marginal influence by ecotypes (Table S4). There was a strong phylogenetic effect on heating tolerance (λ = 0.86 [0.76:0.92]) and an effect of starting temperature where heating tolerance was lower at higher starting temperature (-0.83 [-0.85:-0.82]). Results were also consistent for the corrected heating tolerance (Table S5; Fig. S5).

Figure 4. Heating tolerance (°C) decreases across ontogeny (stage 1–7, colours). Regression line and shaded area represents overall model estimate + 95% CI. Heating tolerance is CT_{max} – starting temperature of the ramping assay (438 species).

Discussion

No strong evidence for thermal bottlenecks in Anurans

Amphibians are an excellent system in which to investigate ontogenetic variation in thermal tolerance because they have a biphasic life cycle with life stages clearly delineated by metamorphosis. The delineation between life stages set by metamorphosis is key as metamorphosis may compartmentalise the life cycle and allow for a potential thermal bottleneck at one stage with minimal detrimental effects of temperature across a life cycle (Enriquez-Urzelai et al., 2019; Lowe, Martin, Skelly, & Woods, 2021).

We find striking differences between adults and larva across ecotypes, even after correcting for experimental variation, in the opposite direction than expected if larva represented a thermal bottleneck. A similar general negative relationship between thermal tolerance over ontogeny is widely reported for frogs (Cupp, 1980; Enriquez-Urzelai et al., 2019; Ruthsatz et al., 2022). Although lower thermal tolerances may be expected for aquatic larva, a higher thermal tolerance may be advantageous for small bodied larva that have reduced capacity to escape temperature extremes (Lenoir et al., 2017). Similarly, a wider selection of

microhabitats in terrestrial environments may allow adults to compensate for a lower
thermal tolerance through behavioural site selection than aquatic larva which grow in a
thermally homogeneous environment (Enriquez-Urzelai et al., 2020).

We find ecotype mediates the magnitude of ontogenetic differences in thermal tolerances
but is not an accurate proxy for environmental selection or constraint. The discrete nature
of complex life cycles with metamorphosis and the need to survive thermal stress at one
stage until the transition to the next stage may place high importance on the interaction
between traits and the environment where a thermal bottleneck is a selective disadvantage.
Microhabitat preferences has been demonstrated to strongly correlate with thermal biology
in reptiles (Greer, 1980; Michelangeli, Goulet, Kang, Wong, & Chapple, 2018), bees (Wilson,
Murphy, Rinehart, Yocum, & Bowsher, 2020), frogs (Carilo Filho et al., 2021; Duarte et al.,
2012), and ants (Bujan, Roeder, de Beurs, Weiser, & Kaspari, 2020; Leahy, Scheffers,
Williams, & Andersen, 2022). We found strong phylogenetic conservatism in CT_{max} which
likely reflects how ecotypes strongly correlate within Family (Moen & Wiens, 2017). For
example, Hylidae have a high proportion of arboreal species (Vidal-García & Keogh, 2015).
The use of ecotype as a proxy is complicated by variation in developmental characteristics
that may not be consistent within Family or ecotypes to make generalisations. For example,
direct-developing larva (not available in our dataset) may occupy more similar microhabitats
to adults than species with an aquatic larva and fully terrestrial adults as discussed with the
limitations of larvae ecotypes. Egg thermal tolerance is rare in the Amphibian literature, but
eggs may also occupy vastly different microhabitats to larva and adults. For instance,
arboreal frogs may create foam nests in the canopy but the larva inhabit puddles on the
forest floor (Haddad & Prado, 2005). While it would be possible to classify larval ecotypes or
egg microhabitats (Carilo Filho et al., 2021), this information is not readily available at the
scale of this study, and we encourage further investigation into these modulators. Doing so
may reveal whether diversity in developmental traits within ecotypes masks local scale
signals of environmental adaption via interactions with ecotype. Nevertheless, we found
fossorial species, on average, had higher thermal tolerance than other ecotypes.

The large overlap in thermal tolerance across ecotypes suggests developmental or genetic
constraints have a greater influence on the evolution of thermal tolerances than
environmental factors. Developmental or genetic constraints on climate niches or thermal
traits are suggested for reptiles and amphibians (Bodensteiner et al., 2021; Carilo Filho et
al., 2021; Pie, Campos, Meyer, & Duran, 2017). Several inferences support developmental or
genetic constraints underlie consistent differences between larva and adult thermal
tolerance. First, an absence for a bottleneck may indicate metamorphosis modulates
thermal tolerances over ontogeny (Lowe et al., 2021). Minimal carry over effects of
development temperature on CT_{max} are found between larva and adults (Enriquez-Urzelai et
al., 2019). Physiological remodelling and morphological development during metamorphosis
are hypothesised to trade-off with physiological processes and explain why CT_{max} decreases
at metamorphic climax (Cupp, 1980; Ruthsatz et al., 2022).

Second, if juveniles and adults are not drastically different in their physiology (i.e.,
biochemical processes are not affected by metamorphosis), then differences in thermal

tolerance may indicate differences in energy allocation between life stages. An energetics
interpretation depends on whether capacity to respond to a thermal challenge is considered
part of maintenance metabolic costs and the flexibility of reallocating energy. Perhaps, the
allocation of energy to reproduction at the adult stage trade-offs with the capacity to
increase maintenance metabolic rates in response to heating, whereas larva without a
reproductive metabolic cost have greater scope to rapidly respond to a thermal challenge at
the cost of allocating energy to growth. This explanation is in line with how the temperature
size rule emerges from differences between development and growth rates (Forster & Hirst,
2012; Gahm, Arietta, & Skelly, 2021; Rubalcaba & Olalla-Tárraga, 2020).

A higher physiological capacity to respond to change (i.e., plasticity) in larva is a
parsimonious explanation from an energetics perspective. In contrast with studies that
show decreasing acclimation capacity with increasing size (Burton, Lakka, & Einum, 2020;
Rohr et al., 2018), *Rana temporaria* larva have lower acclimation capacity than adults
(Ruthsatz et al., 2022). Assuming the cost of having a given thermal tolerance is the same
between life stages, low plasticity in thermal tolerance suggests larva continuously express
physiological mechanisms that permit higher CT_{max}, which is very costly (i.e., in maintenance
metabolic cost) during a time when they should be allocating energy to growth. One
explanation for low thermal plasticity is tadpoles have a narrow range of temperatures to
select for behavioural thermoregulation and thus primarily rely on physiological
thermoregulation via a higher CT_{max} to avoid heat stress (Casterlin & Reynolds, 1978). As
larva would have lower absolute metabolic rate based on size allometry alone (in general),
whether or not they can offset an increased energetic cost with increased food intake or by
other means can be explored.

Third, if physiological mechanisms are different between life stages, then the differences
may reflect developmental constraints. Perhaps the physiological machinery during larval
growth covaries with greater capacity to respond to an environmental challenge, or that
larva have lower energetic costs for responding to thermal challenges, which allows them to
continuously express high CT_{max} at minimal cost to growth, in contrast with adults. This
explanation would resolve the paradox of the high metabolic costs of physiological
thermoregulation.

Standardisation reduces confounding interaction across life cycles but does not change our conclusions

Minimal difference between observed CT_{max} and corrected CT_{max} likely underpin the
absence of significant changes to our inferences. There is a relative consistency in methods
to assay CT_{max} in amphibians and the range of heating rates is narrow with 1 °C min^{-1} as the
mode (Hutchison, 1961). Most studies in our literature search (2016-2022) assayed CT_{max}
using a ramping protocol rather than static protocols (zero studies described) or from fitted
thermal performance curves (three studies). Applying a correction to CT_{max} is likely more
effective for other taxa or in broader scale interspecific comparisons. For example, heating
rate varied between 0.0007 and 4 °C min^{-1} (mean 0.5 °C min^{-1}) in a global dataset of
ectotherm CT_{max} that included tropical to polar organisms (Morley, Peck, Sunday, Heiser, &

Bates, 2018), and rescaling this data for non-linear rate-temperature dynamics produced
high precision estimates of CT_{max} (Kong et al., 2022). In studies on frogs, ontogenetic stage is
confounded with heating rates as lower heating rates are typically used for larva than adults
or juveniles (Fig. S3). Thus, applying a correction to CT_{max} may be useful to infer true effects
of ontogenetic stage and we suggest it a reasonable practice to account for nuisance
heterogeneity in comparative study.

To our knowledge, our study is the first to apply a theoretic model incorporating heating
rate and starting temperature to a comparative dataset of this scale (Pottier, Burke,
Drobniak, & Nakagawa, 2022). Previous studies have used statistical regression to correct
for variation in acclimation temperature on CT_{max} (e.g. Sunday et al. (2014)). In other words,
rescaling of heating tolerances represents a null model of how CT_{max} may vary with heating
rate and starting temperature, based on physiological and non-linear rate-time dynamics.
However, missing covariates of CT_{max} reduced the effectiveness of applying a mathematical
correction. We were not able to correct CT_{max} for studies that did not report acclimation
temperature, starting temperature or heating rate (80 observations). As heating rate and
starting temperature are important sources of variation in CT_{max}, we suggest these common
covariates should be reported as best practice (Schwanz et al., 2022; Sunday et al., 2014).

Applying any correction relies on making generalisations about the nature of the
relationship between CT_{max} and experimental covariates, and its consistency across taxa.
The rescaling approach used here assumes that the underlying processes that generate
what we observe as CT_{max} is modelled by a Boltzmann-Arrhenius function with an activation
energy of 0.6eV, and this relationship is insensitive to change (e.g., acclimation or plasticity).
This relationship may represent whole-organism metabolic rate encompassing the general
rate-dependent physiology of animals, for example, and may be appropriate to describe
thermal tolerances as emergent properties of several interacting processes at the sub-
individual level, but this model may differ from other hypotheses about how thermal
tolerance arises (MacMillan, 2019).

We suggest using a general model of thermal tolerance is more appropriate for large scale
comparative studies where it is inappropriate to assume consistent or homogeneous
mechanisms across taxa or large scales. Several other models of CT_{max} incorporating starting
temperature and heating rate invoke more specific mechanisms, such as injury
accumulation and damage repair (Jørgensen, Malte, Ørsted, Klahn, & Overgaard, 2021;
Rezende, Bozinovic, Szilágyi, & Santos, 2020). In our dataset, CT_{max} endpoints are relatively
consistent across studies and within ontogenetic stage, but ontogenetic stage is confounded
by endpoint definition. For example, onset of spasms or heat rigour were more common for
larva and loss of righting response is most common for adults. A model invoking a
hypothesised mechanism could be relevant for investigating specific endpoints or
definitions of upper critical temperatures. Overall, the appropriateness of a model for its
use in comparative study depends on the underlying assumptions, and the utility of the
model is a trade-off between the requirements for model parameterisation and its intended
use.

Conclusions

Dealing with high among-study variation or problems of scale and resolution are resounding challenges of macrophysiology (Chown & Gaston, 2016). This necessity becomes critical when assessing vulnerability to future climate change because understanding trends in physiological diversity across taxa and environmental gradients forms a basis to predict or forecast climate change responses or vulnerability (Urban et al., 2016). For broad-scale comparisons, we suggest standardising against common experimental variation is a meaningful exercise regardless of any conclusion because potentially confounding variation may be tied with an organisms' life cycle, and uncertainty in CT_{max} is passed onto the derived climate vulnerability metrics (Garcia, Allen, & Clusella-Trullas, 2019; Pincebourde & Casas, 2015).

References

Bates, A. E., & Morley, S. A. (2020). Interpreting empirical estimates of experimentally derived physiological and biological thermal limits in ectotherms. *Canadian Journal of Zoology, 98*(4), 237-244. doi:10.1139/cjz-2018-0276

Bodensteiner, B. L., Agudelo-Cantero, G. A., Arietta, A. Z. A., Gunderson, A. R., Muñoz, M. M., Refsnider, J. M., & Gangloff, E. J. (2021). Thermal adaptation revisited: How conserved are thermal traits of reptiles and amphibians? *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology, 335*(1), 173-194. doi:10.1002/jez.2414

Brattstrom, B. H. (1970). Thermal acclimation in Australian amphibians. *Comparative Biochemistry and Physiology, 35*(1), 69-103. doi:10.1016/0010-406X(70)90915-1

Bujan, J., Roeder, K. A., de Beurs, K., Weiser, M. D., & Kaspari, M. (2020). Thermal diversity of North American ant communities: Cold tolerance but not heat tolerance tracks ecosystem temperature. *Global Ecology and Biogeography, 29*(9), 1486-1494. doi:10.1111/geb.13121

Burton, T., Lakka, H.-K., & Einum, S. (2020). Acclimation capacity and rate change through life in the zooplankton Daphnia. *Proceedings of the Royal Society B: Biological Sciences, 287*(1924), 20200189. doi:10.1098/rspb.2020.0189

Carilo Filho, L. M., de Carvalho, B. T., Azevedo, B. K. A., Gutiérrez-Pesquera, L. M., Mira-Mendes, C. V., Solé, M., & Orrico, V. G. D. (2021). Natural history predicts patterns of thermal vulnerability in amphibians from the Atlantic Rainforest of Brazil. *Ecology and Evolution, 11*(23), 16462-16472. doi:10.1002/ece3.7961

Casterlin, M. E., & Reynolds, W. W. (1978). Behavioural thermoregulation in Rana pipiens tadpoles. *Journal of Thermal Biology, 3*(3), 143-145. doi:10.1016/0306-4565(78)90009-8

Cheng, C.-T., Chuang, M.-F., Haramura, T., Cheng, C.-B., Kim, Y. I., Borzée, A., . . . Kam, Y.-C. (2022). Open habitats increase vulnerability of amphibian tadpoles to climate warming across latitude. *Global Ecology and Biogeography, n/a*(n/a). doi:https://doi.org/10.1111/geb.13602

Chown, S. L., & Gaston, K. J. (2016). Macrophysiology – progress and prospects. *Functional Ecology, 30*(3), 330-344. doi:10.1111/1365-2435.12510

Collin, R., Rebolledo, A. P., Smith, E., & Chan, K. Y. K. (2021). Thermal tolerance of early
development predicts the realized thermal niche in marine ectotherms. *Functional
Ecology, 35*(8), 1679-1692. doi:10.1111/1365-2435.13850
Cupp, P. V. (1980). Thermal Tolerance of Five Salientian Amphibians during Development
and Metamorphosis. *Herpetologica, 36*(3), 234-244.
Dahlke, F. T., Wohlrab, S., Butzin, M., & Pörtner, H.-O. (2020). Thermal bottlenecks in the life
cycle define climate vulnerability of fish. *Science, 369*(6499), 65-70.
doi:10.1126/science.aaz3658
Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., &
Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms across
latitude. *Proceedings of the National Academy of Sciences of the United States of
America, 105*(18), 6668-6672. doi:10.1073/pnas.0709472105
Duarte, H., Tejedo, M., Katzenberger, M., Marangoni, F., Baldo, D., Beltrán, J. F., . . .
Gonzalez-Voyer, A. (2012). Can amphibians take the heat? Vulnerability to climate
warming in subtropical and temperate larval amphibian communities. *Global Change
Biology, 18*(2), 412-421. doi:10.1111/j.1365-2486.2011.02518.x
Enriquez-Urzelai, U., Sacco, M., Palacio, A. S., Pintanel, P., Tejedo, M., & Nicieza, A. G.
(2019). Ontogenetic reduction in thermal tolerance is not alleviated by earlier
developmental acclimation in Rana temporaria. *Oecologia, 189*(2), 385-394.
doi:10.1007/s00442-019-04342-y
Enriquez-Urzelai, U., Tingley, R., Kearney, M. R., Sacco, M., Palacio, A. S., Tejedo, M., &
Nicieza, A. G. (2020). The roles of acclimation and behaviour in buffering climate
change impacts along elevational gradients. *Journal of Animal Ecology, 89*(7), 1722-
1734. doi:10.1111/1365-2656.13222
Forster, J., & Hirst, A. G. (2012). The temperature-size rule emerges from ontogenetic
differences between growth and development rates. *Functional Ecology, 26*(2), 483-
492. doi:10.1111/j.1365-2435.2011.01958.x
Gahm, K., Arietta, A. Z. A., & Skelly, D. K. (2021). Temperature-mediated trade-off between
development and performance in larval wood frogs (Rana sylvatica). *Journal of
Experimental Zoology Part A: Ecological and Integrative Physiology, 335*(1), 146-157.
doi:10.1002/jez.2434
Garcia, R. A., Allen, J. L., & Clusella-Trullas, S. (2019). Rethinking the scale and formulation of
indices assessing organism vulnerability to warmer habitats. *Ecography, 42*(5), 1024-
1036. doi:10.1111/ecog.04226
González-del-Pliego, P., Scheffers, B. R., Freckleton, R. P., Basham, E. W., Araújo, M. B.,
Acosta-Galvis, A. R., . . . Edwards, D. P. (2020). Thermal tolerance and the importance
of microhabitats for Andean frogs in the context of land use and climate change.
Journal of Animal Ecology, 89(11), 2451-2460. doi:10.1111/1365-2656.13309
Greer, A. (1980). Critical Thermal Maximum Temperatures in Australian Scincid Lizards:
Their Ecological and Evolutionary Significance. *Australian Journal of Zoology, 28*(1),
91-102. doi:10.1071/ZO9800091
Haddad, C. F. B., & Prado, C. P. A. (2005). Reproductive Modes in Frogs and Their
Unexpected Diversity in the Atlantic Forest of Brazil. *Bioscience, 55*(3), 207-217.
doi:10.1641/0006-3568(2005)055[0207:RMIFAT]2.0.CO;2
Hutchison, V. H. (1961). Critical Thermal Maxima in Salamanders. *Physiological Zoology,
34*(2), 92-125. doi:10.1086/physzool.34.2.30152688

IUCN. (2022). *The IUCN Red List of Threatened Species*. Retrieved from:
https://www.iucnredlist.org

Jetz, W., & Pyron, R. A. (2018). The interplay of past diversification and evolutionary
isolation with present imperilment across the amphibian tree of life. *Nature Ecology*
& Evolution, 2(5), 850-858. doi:10.1038/s41559-018-0515-5

Jørgensen, L. B., Malte, H., Ørsted, M., Klahn, N. A., & Overgaard, J. (2021). A unifying model
to estimate thermal tolerance limits in ectotherms across static, dynamic and
fluctuating exposures to thermal stress. *Scientific Reports, 11*(1), 12840.
doi:10.1038/s41598-021-92004-6

Kingsolver, J. G., & Buckley, L. B. (2020). Ontogenetic variation in thermal sensitivity shapes
insect ecological responses to climate change. *Current Opinion in Insect Science, 41*,
17-24. doi:10.1016/j.cois.2020.05.005

Klockmann, M., Günter, F., & Fischer, K. (2017). Heat resistance throughout ontogeny: body
size constrains thermal tolerance. *Global Change Biology, 23*(2), 686-696.
doi:10.1111/gcb.13407

Kong, J. D., Arnoldi, J.-F., Jackson, A. L., Bates, A. E., Morley, S. A., Smith, J. A., & Payne, N. L.
(2022). Heating tolerance of ectotherms is explained by temperature's non-linear
influence on biological rates. *bioRxiv*, 2022.2012.2006.519315.
doi:10.1101/2022.12.06.519315

Kovacevic, A., Latombe, G., & Chown, S. L. (2019). Rate dynamics of ectotherm responses to
thermal stress. *Proceedings of the Royal Society B: Biological Sciences, 286*(1902),
20190174. doi:doi:10.1098/rspb.2019.0174

Leahy, L., Scheffers, B. R., Williams, S. E., & Andersen, A. N. (2022). Arboreality drives heat
tolerance while elevation drives cold tolerance in tropical rainforest ants. *Ecology,*
103(1), e03549. doi:10.1002/ecy.3549

Lebenzon, J. E., Des Marteaux, L. E., & Sinclair, B. J. (2020). Reversing sodium differentials
between the hemolymph and hindgut speeds chill coma recovery but reduces
survival in the fall field cricket, Gryllus pennsylvanicus. *Comparative Biochemistry*
and Physiology Part A: Molecular & Integrative Physiology, 244, 110699.
doi:10.1016/j.cbpa.2020.110699

Lenoir, J., Hattab, T., & Pierre, G. (2017). Climatic microrefugia under anthropogenic climate
change: implications for species redistribution. *Ecography, 40*(2), 253-266.
doi:10.1111/ecog.02788

Leong, C. M., Tsang, T. P. N., & Guenard, B. (2022). Testing the reliability and ecological
implications of ramping rates in the measurement of Critical Thermal maximum.
PLOS ONE, 17(3), e0265361. doi:10.1371/journal.pone.0265361

Levy, O., Buckley, L. B., Keitt, T. H., Smith, C. D., Boateng, K. O., Kumar, D. S., & Angilletta, M.
J., Jr. (2015). Resolving the life cycle alters expected impacts of climate change.
Proceedings: Biological Sciences, 282(1813), 20150837. doi:10.1098/rspb.2015.0837

Lowe, W. H., Martin, T. E., Skelly, D. K., & Woods, H. A. (2021). Metamorphosis in an Era of
Increasing Climate Variability. *Trends in Ecology & Evolution, 36*(4), 360-375.
doi:10.1016/j.tree.2020.11.012

MacMillan, H. A. (2019). Dissecting cause from consequence: a systematic approach to
thermal limits. *Journal of Experimental Biology, 222*(4). doi:10.1242/jeb.191593

Michelangeli, M., Goulet, C. T., Kang, H. S., Wong, B. B. M., & Chapple, D. G. (2018).
Integrating thermal physiology within a syndrome: Locomotion, personality and

habitat selection in an ectotherm. *Functional Ecology, 32*(4), 970-981. doi:10.1111/1365-2435.13034

Moen, D. S., Cabrera-Guzman, E., Caviedes-Solis, I. W., Gonzalez-Bernal, E., & Hanna, A. R. (2022). Phylogenetic analysis of adaptation in comparative physiology and biomechanics: overview and a case study of thermal physiology in treefrogs. *Journal of Experimental Biology, 225*(Suppl1), jeb243292. doi:10.1242/jeb.243292

Moen, D. S., & Wiens, J. J. (2017). Microhabitat and Climatic Niche Change Explain Patterns of Diversification among Frog Families. *The American Naturalist, 190*(1), 29-44. doi:10.1086/692065

Morley, S. A., Peck, L. S., Sunday, J., Heiser, S., & Bates, A. E. (2018). *Acclimation potential of global ectothermic species, collated from literature, 1960 to 2015*.

Moyano, M., Candebat, C., Ruhbaum, Y., Álvarez-Fernández, S., Claireaux, G., Zambonino-Infante, J.-L., & Peck, M. A. (2017). Effects of warming rate, acclimation temperature and ontogeny on the critical thermal maximum of temperate marine fish larvae. *PLOS ONE, 12*(7), e0179928. doi:10.1371/journal.pone.0179928

Noble, D. W. A., Pottier, P., Lagisz, M., Burke, S., Drobniak, S. M., O'Dea, R. E., & Nakagawa, S. (2022). Meta-analytic approaches and effect sizes to account for 'nuisance heterogeneity' in comparative physiology. *Journal of Experimental Biology, 225*(Suppl_1), jeb243225. doi:10.1242/jeb.243225

Pandori, L. L. M., & Sorte, C. J. B. (2019). The weakest link: sensitivity to climate extremes across life stages of marine invertebrates. *Oikos, 128*(5), 621-629. doi:10.1111/oik.05886

Peck, L. S., Clark, M. S., Morley, S. A., Massey, A., & Rossetti, H. (2009). Animal temperature limits and ecological relevance: effects of size, activity and rates of change. *Functional Ecology, 23*(2), 248-256. doi:10.1111/j.1365-2435.2008.01537.x

Pie, M. R., Campos, L. L. F., Meyer, A. L. S., & Duran, A. (2017). The evolution of climatic niches in squamate reptiles. *Proceedings: Biological Sciences, 284*(1858). doi:10.1098/rspb.2017.0268

Pincebourde, S., & Casas, J. (2015). Warming tolerance across insect ontogeny: influence of joint shifts in microclimates and thermal limits. *Ecology, 96*(4), 986-997. doi:10.1890/14-0744.1

Pincebourde, S., & Woods, H. A. (2020). There is plenty of room at the bottom: microclimates drive insect vulnerability to climate change. *Current Opinion in Insect Science, 41*, 63-70. doi:10.1016/j.cois.2020.07.001

Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L., & Sunday, J. M. (2019). Greater vulnerability to warming of marine versus terrestrial ectotherms. *Nature, 569*(7754), 108-111. doi:10.1038/s41586-019-1132-4

Pintanel, P., Tejedo, M., Merino-Viteri, A., Almeida-Reinoso, F., Salinas-Ivanenko, S., Lopez-Rosero, A. C., . . . Gutierrez-Pesquera, L. M. (2022). Elevational and local climate variability predicts thermal breadth of mountain tropical tadpoles. *Ecography, 2022*(5), e05906. doi:10.1111/ecog.05906

Pintanel, P., Tejedo, M., Ron, S. R., Llorente, G. A., & Merino-Viteri, A. (2019). Elevational and microclimatic drivers of thermal tolerance in Andean Pristimantis frogs. *Journal of Biogeography, 46*(8), 1664-1675. doi:10.1111/jbi.13596

Pottier, P., Burke, S., Drobniak, S. M., & Nakagawa, S. (2022). Methodological inconsistencies define thermal bottlenecks in fish life cycle: a comment on Dahlke et al. 2020. *Evolutionary Ecology, 36*(2), 287-292. doi:10.1007/s10682-022-10157-w

Rezende, E. L., Bozinovic, F., Szilágyi, A., & Santos, M. (2020). Predicting temperature
mortality and selection in natural *Drosophila* populations. *Science, 369*(6508), 1242-
1245. doi:10.1126/science.aba9287
Rohr, J. R., Civitello, D. J., Cohen, J. M., Roznik, E. A., Sinervo, B., & Dell, A. I. (2018). The
complex drivers of thermal acclimation and breadth in ectotherms. *Ecology Letters,
21*(9), 1425-1439. doi:10.1111/ele.13107
Rubalcaba, J. G., & Olalla-Tárraga, M. Á. (2020). The biogeography of thermal risk for
terrestrial ectotherms: Scaling of thermal tolerance with body size and latitude.
Journal of Animal Ecology, n/a(n/a), 1277-1285. doi:10.1111/1365-2656.13181
Ruthsatz, K., Dausmann, K. H., Peck, M. A., & Glos, J. (2022). Thermal tolerance and
acclimation capacity in the European common frog (Rana temporaria) change
throughout ontogeny. *J Exp Zool A Ecol Integr Physiol, 337*(5), 477-490.
doi:10.1002/jez.2582
Schwanz, L. E., Gunderson, A., Iglesias-Carrasco, M., Johnson, M. A., Kong, J. D., Riley, J., &
Wu, N. C. (2022). Best practices for building and curating databases for comparative
analyses. *Journal of Experimental Biology, 225*(Suppl_1), jeb243295.
doi:10.1242/jeb.243295
Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S., . . .
Huey, R. B. (2016). Can we predict ectotherm responses to climate change using
thermal performance curves and body temperatures? *Ecology Letters, 19*(11), 1372-
1385. doi:10.1111/ele.12686
Sunday, J. M., Bates, A. E., Kearney, M. R., Colwell, R. K., Dulvy, N. K., Longino, J. T., & Huey,
R. B. (2014). Thermal-safety margins and the necessity of thermoregulatory behavior
across latitude and elevation. *Proceedings of the National Academy of Sciences of the
United States of America, 111*(15), 5610-5615. doi:10.1073/pnas.1316145111
Terblanche, J. S., Deere, J. A., Clusella-Trullas, S., Janion, C., & Chown, S. L. (2007). Critical
thermal limits depend on methodological context. *Proceedings of the Royal Society
B: Biological Sciences, 274*(1628), 2935-2943. doi:doi:10.1098/rspb.2007.0985
Terblanche, J. S., Hoffmann, A. A., Mitchell, K. A., Rako, L., le Roux, P. C., & Chown, S. L.
(2011). Ecologically relevant measures of tolerance to potentially lethal
temperatures. *Journal of Experimental Biology, 214*(Pt 22), 3713-3725.
doi:10.1242/jeb.061283
Truebano, M., Fenner, P., Tills, O., Rundle, S. D., & Rezende, E. L. (2018). Thermal strategies
vary with life history stage. *Journal of Experimental Biology, 221*(8), jeb171629.
doi:10.1242/jeb.171629
Urban, M. C., Bocedi, G., Hendry, A. P., Mihoub, J. B., Pe'er, G., Singer, A., . . . Travis, J. M.
(2016). Improving the forecast for biodiversity under climate change. *Science,
353*(6304). doi:10.1126/science.aad8466
Vidal-García, M., & Keogh, J. S. (2015). Convergent evolution across the Australian
continent: ecotype diversification drives morphological convergence in two distantly
related clades of Australian frogs. *Journal of Evolutionary Biology, 28*(12), 2136-2151.
doi:10.1111/jeb.12746
Wilson, E. S., Murphy, C. E., Rinehart, J. P., Yocum, G., & Bowsher, J. H. (2020). Microclimate
Temperatures Impact Nesting Preference in Megachile rotundata (Hymenoptera:
Megachilidae). *Environmental Entomology*. doi:10.1093/ee/nvaa012

611