Microhabitat and ontogenetic variation of thermal tolerance in complex ectotherm life cycles

Jacinta D. Kong[1*], Nicholas C. Wu[2]

[1] *Department of Biology, Carleton University, ON, Canada*

[2]*Hawkesbury Institute for the Environment, Western Sydney University, NSW 2753, Australia*

Keywords: amphibia, critical temperature maxima, life stage, macrophysiology, ontogeny

***Correspondence**: jacintadkong@gmail.com

ORCID: JDK (0000-0002-1085-8612), NCW (0000-0002-7130-1279)

Author contributions: JDK: Conceptualization, Investigation, Data Curation, Software, Writing - Original Draft. NCW: Resources, Software, Formal analysis, Visualization. Both authors reviewed and edited the work and gave final permission for submission.

Competing Interest: The authors declare no competing interests.

Data availability statement: Data and code are available on Zenodo (https://github.com/jacintak/frog_ctmax)

Article type: Research article

Acknowledgements: The authors would like to thank researchers C. Filho (Universidade Estadual de Santa Cruz) and R. Percinz (Ruth Percino) who provided data for this study.

Abstract

Thermal tolerances, such as critical temperatures, are important indices for understanding an organism's vulnerability to changing environmental temperature. Differences in thermal tolerance over ontogeny may generate a 'thermal bottleneck' that sets the climate vulnerability for organisms with complex life cycles. However, a species' microhabitat preference and life history can hinder our ability to assess climate change vulnerability in

large-scale comparative studies as these reflect local adaptation. Here, we used

phylogenetically informed, multi-level models with a global dataset of upper critical

temperatures from 438 Anuran species that explicitly included microhabitat preferences

and ontogenetic stage to examine variation in upper critical temperatures. We found

ontogenetic trends in thermal tolerance were similar across microhabitat preferences. We

then used microclimate-driven models of the heat and water budget of a thermoregulating

frog to show the contribution of behavioural thermoregulation towards mitigating exposure

to thermal stress. Our results suggested thermal bottlenecks are not strongly present in

Anurans but instead implied strong developmental or genetic conservatism of thermal

tolerance within families and ecotypes, and that behavioural thermoregulation has a strong

potential for buffering frogs against extreme temperatures.

Introduction

Introduction

Thermal tolerance metrics, such as critical temperatures, have endured in ecological studies

despite their criticisms because they are simple to conceptualise (e.g. impediment to

movement), are generalisable across taxa, and can be readily assayed in the laboratory

(Sinclair et al. 2016). These metrics facilitate our ability to link environmental temperature

to organismal responses (Bates and Morley 2020), synthesise general trends in physiological

diversity from local to macro scales (Chown and Gaston 2016), and index vulnerability or

sensitivity to climate change via metrics such as thermal safety margins (Deutsch et al.

2008). For example, differences in upper critical temperature (CT_{max}) across life stages may

indicate a thermal bottleneck at the most temperature-sensitive life stage (Pandori and

Sorte 2019, Dahlke et al. 2020). For organisms with a complex life cycle such as frogs,

juvenile or larval stages in aquatic environments are generally expected to have lower

thermal tolerance and thus be more vulnerable to climate change than adult stages (Lenoir

et al. 2017).

Identifying a thermal bottleneck represents a significant amount of work that is limited to

commonly studied species (e.g., Ruthsatz et al. (2022)). Thermal bottlenecks can also be

identified using a macrophysiological approach which has described stage-specific

differences in thermal sensitivities and tolerances in marine invertebrates (Truebano et al.

2018, Pandori and Sorte 2019, Collin et al. 2021), fish (Moyano et al. 2017, Dahlke et al.

2020), insects (Klockmann et al. 2017, Kingsolver and Buckley 2020) and reptiles (Levy et al.

2015). Among species comparisons in frogs are often limited to a single geographic region

and are often limited to either adults or larva even if both are collected from the same

region (Pintanel et al. 2019, Carilo Filho et al. 2021, Cheng et al. 2022, Pintanel et al. 2022,

Ruthsatz et al. 2022). Thus, whether thermal bottlenecks occur and how consistent they are

across Anuran species is not well understood. A macrophysiological analyses of thermal

tolerance throughout ontogeny will vastly improve our understanding of the presence and

generality of thermal bottlenecks across an environmentally sensitive taxonomic group.

Comparative macrophysiological analyses present two key challenges towards assessing and

interpreting thermal tolerances. Microhabitat preferences and among-study variation may

generate confounding variation in thermal tolerances that may mask or accentuate the

presence of a thermal bottleneck. The key challenge to interpret thermal tolerance over the

life cycle is to disentangle confounding variation deriving from microhabitat selection from

inherent developmental or physiological drivers. A diversity of microhabitats may mask

broader adaptive covariation between thermal tolerance and microclimate selection to

explain, in part, an observed lack of variation in CT_{max} at a global scale or drive strong

associations with thermal tolerance (González-del-Pliego et al. 2020, Pincebourde and

Woods 2020). However, distinguishing between these patterns requires corresponding

microclimate information for each ontogenetic stage that is not always collected for thermal

tolerance studies or are available, but see Pintanel et al. (2022). If differences in thermal

environments are consistent within broad microhabitat preferences in Anurans, then

microhabitat preferences may be a reliable proxy for local environmental selection on

thermal tolerance. Among species, frogs show strong microhabitat preferences that are

associated with phenotypic syndromes encompassing behaviour, morphology and

physiology and can broadly be categorised into ecological types ("ecotype") (Moen and

Wiens 2017, Moen et al. 2022). In turn, the animal's ecotype may mediate the confounding

interaction between life stages and their respective microenvironment (Pincebourde and

Casas 2015, Carilo Filho et al. 2021).

Thermal safety margins (TSM) link upper critical temperatures to environmental

temperature to index the vulnerability of ectotherms to temperature extremes (Deutsch et

al. 2008). Most macrophysiological assessments of thermal safety margins use global

datasets of climate and mean critical temperatures for species but the resolution of climatic

data can have large impact on estimated TSM that leads to an underestimation of TSM

(Garcia et al. 2019, Pinsky et al. 2019). Behavioural thermoregulation is important in this

context as ectotherms that are able to behaviourally thermoregulate may be able to

mitigate the effects of warming (Sunday et al. 2014). Thus, assessments of climate

vulnerability using TSM should consider both the resolution of climate data and the innate

capacity of species to respond to temperature variation.

Here, we tested whether ecotypes are reliable proxies of environmental adaptation to

resolve ontogeny by environment interaction of CT_{max} and identify thermal bottlenecks. We

analysed the CT_{max} of 438 Anuran species to 1) identify whether Anurans have a thermal

bottleneck in their life cycle and their potential evolutionary drivers and ecological

consequences at a macrophysiological scale, and 2) evaluate whether proposed solutions to

two challenges of comparative analyses of ectotherm thermal tolerance improved our

ability to identify thermal bottlenecks. We acknowledge there is a diversity of definitions

and uses of thermal tolerance that would influence their interpretation. In this study, we

only consider CT_{max} as the end point of a temperature ramping assay with a constant

heating rate, or dynamic ramping protocol, and we define heating tolerance as the

temperature range between starting temperature and CT_{max} during the ramping assay.

Figure 1 Schematic diagram of ecotype microclimate and key variables used to calculate representative microclimate.
Arboreal frogs were modelled at 1m 60% shaded air temperature, fossorial frogs at 10cm soil depth in 90% shaded soil, and
ground-dwelling frogs were modelled at 1cm 0% shaded (open) air temperature. Thermoregulating ground-dwelling frogs
could move to 90% shaded 1cm air temperatures, indicated by the arrows.

Methods

Data compilation

We started with a published dataset of upper critical temperatures of Anurans complied from the literature (1962 – 2015) (Rohr et al. 2018). We supplemented this data with studies from 2016 to 2022 using Web of Science (accessed 25th February) and the keywords amphibian* AND ("critical thermal max*" OR CT_{max} OR "thermal toleran*" OR "critical temp*" OR "upper temp*")". The terms returned 90 studies that were screened for inclusion by one person (J.D.K.). Studies that measured CT_{max} using ramping assays were retained. Studies that estimated CT_{max} from fitted thermal performance curves or compiled values from literature were excluded. Only control animals were included from studies that used multiple stressors (e.g. dehydration or infection). The screening process yielded 36 papers that provided extractable data. Provided data were used over published values

where possible. Populations were considered independent observations. Multiple endpoints

of critical temperatures on an individual were recorded but only one CT_{max} was used per

individual (loss of righting response where possible). We added studies that did not return

from keywords from forward searching. We distinguished between temperatures animals

were held at prior to experimentation (acclimation temperature) and the starting

temperature of the ramping assay. We assume that observed CT_{max} reflected their

acclimated CT_{max} as acclimation in frogs is reported on scales of hours (Brattstrom 1970).

We obtained ecological types (hereby ecotype), from all Anuran species listed on the

International Union for Conservation of Nature's Red List of Threatened Species (IUCN

2022). The data were reported in (Wu et al. 2024). Ecotypes were classified from ecological

descriptions from IUCN based on adult behaviour and microhabitat preferences outside the

breeding season (Moen and Wiens 2017). We grouped species by ecotype even if the larval

stage may have differing ecological types (e.g. lotic vs lentic systems) because the ecotype

diversity of the larvae stage is broader than adults and is less described in the detail from

the literature. The resulting dataset represented 32 families spanning the globe (Fig. S1).

Thermal tolerance records were first matched to the ecotype dataset by binomial names

used in the original source, then unmatched records were joined by IUCN names based on

synonyms accepted by IUCN or Catalogue of Life. CT_{max} records that could not be matched

with ecotype (56 observations) were excluded.

143   Relationship between CT_{max} and heating tolerance with ecotype and

144   ontogeny

We constructed a phylogenetically corrected multi-level Bayesian model to investigate the

relationship between CT_{max} or heating tolerance across ontogenetic stage and ecotype (see

Supplementary Information for detail). We included starting temperature of the assay as a

co-variate. Ontogenetic stages 1-5 correspond to larva, 6 to juveniles, and 7 to adults

(Ruthsatz et al. 2022). We used an amphibian phylogeny dataset comprising of 7,238 species

to account for phylogenetic relatedness (Jetz and Pyron 2018). We pruned the phylogenetic

tree to matched binomial names in the thermal tolerance data (438 species, Fig. 2). Species

names based on Jetz and Pyron (2018) were used in analysis to account for duplicate names

matching with the IUCN. Records that were not in the phylogenetic tree (n = 1) were

excluded. We converted the pruned tree to a phylogenetic relatedness correlation matrix

for subsequent analysis. For the CT_{max} model, we used a quadratic equation to model

change in CT_{max} over ontogeny and the quadratic equation provided a better fit than the

linear model (Table S1). All model outputs (Tables S2 – S5) and posterior predictive checks

(Figs. S4 & S6) are presented in the Supplementary Information. All model parameters are

reported as mean Bayesian posterior estimates ± 95% credible intervals (95% CI), which is

equivalent to 95% confidence intervals for frequentist statistics. Summary statistics are

presented as mean ± standard deviation.

Thermal safety margins at macro- and micro-climate scales

To calculate thermal safety margins we extracted mean annual temperature and maximum

temperature of the warmest month from WorldClim 2.1 (resolution: 0.5 min of a degree) for

each observation (Fick and Hijmans 2017). TSM was calculated as upper critical temperature

(CTmax) - maximum temperature of the warmest month. Observations that could not be

queried on WorldClim were removed from analysis. To illustrate the effect of spatial

resolution on estimates of TSM for representative ecotypes, we chose frogs representing

three ecotypes (Fig. 1): fossorial (*Neobatrachus pictus*), arboreal (*Litoria peronii*) and

ground-dwelling (*Crinia signifera*). These species are sympatric in north-eastern New South

Wales, Australia and were measured in the same study using the same conditions (5°C

acclimation temperature and 1°C min^{-1} heating rate) (Brattstrom 1970). Because of the

additional challenges of modelling microclimates involving water and the potentially large

temperature difference between air and water, we were not able to conduct simulations for

aquatic, stream-dwelling and semi-aquatic ecotypes We used NicheMapR to simulate

representative microclimates for a fossorial (90% shaded 0.1 m soil temperature), arboreal

(60% shaded 1m air temperature) and ground-dwelling species (0% shaded 0.01m air

temperature) (Kearney and Porter 2017). These microclimates informed a biophysical

energy and mass balance model of a hypothetical non-thermoregulating 5g frog

(approximating a leopard frog) that we used to estimate operative temperature (maximum

body temperature) for each species (Kearney and Porter 2019, Enriquez-Urzelai et al. 2020,

Wu et al. 2024). To demonstrate the role of behavioural thermoregulation for estimating

TSM, we ran the biophysical model twice for the ground-dwelling frog allowing for or

restricting microhabitat selection by shuttling between shaded and unshaded microhabitats.

For each ecotype, we extracted maximum temperature at the locality of the individual and

maximum body temperature, representing operative temperature, from the year-worth of

output (Fig. 1).

Figure 2. Phylogenetic patterns of CT$_{max}$ across 438 Anurans based on Jetz & Pyron (2018). Colours represents species mean

CT$_{max}$. Example families are indicated by outer black lines. Representative silhouettes obtained from phylopic.org and

background colours within the circles represent mean CT$_{max}$ at the Family level.

Results

CT$_{max}$

CT$_{max}$ ranged between 20.4 °C and 46.1 °C (mean 37.1 ± 4.1 °C) across all ecotypes. On average, fossorial frogs have the highest mean CT$_{max}$ (40.1 ± 3.1 °C) and semi-aquatic frogs have the lowest CT$_{max}$ (35.4 ± 3.54 °C). Larval CT$_{max}$ (Stage 1-5) was on average 3.5 °C higher than adult (Stage 7) and juvenile (Stage 6) CT$_{max}$ combined; the mean difference ranged from 1.5 °C for semi-aquatic frogs and 5.2 °C for stream-dwelling frogs. However, there are fewer data on CT$_{max}$ for early larval (Stage 1, n = 41) and juvenile (Stage 6, n = 33) stages than later larval stages (Stage 3, n = 207) and adults (Stage 7, n = 733).

Variation in CT$_{max}$ was strongly influenced by ontogeny where CT$_{max}$ generally decreased across ontogeny (Table S2; Fig. 3). CT$_{max}$ did not differ substantially between ecotypes, but on average, stream-dwelling frogs have lower CT$_{max}$ (-0.74 [-2.63: 1.11]), while fossorial frogs generally had higher CT$_{max}$ (1.96 [-0.06:4.05]). There was a strong phylogenetic effect on CT$_{max}$ (λ = 0.85 [0.75:0.92]). Starting temperature also had a strong effect on CT$_{max}$, where the higher the starting temperature, the higher the CT$_{max}$ (0.16 [0.14:0.18]).

Figure 3. CT_max (℃) decreases over ontogeny (stage 1–7) across all Anuran ecotypes (438 species). Quadratic regression line and shaded area represents ecotype-specific model estimate ± 95% CI.

Heating tolerance

Heating tolerances ranged between 2 °C and 36.5 °C (16.7 ± 6.34 °C) across all ecotypes.
Mean heating tolerance varied between ecotypes with the highest mean in fossorial frogs
(18.6 ± 6.1 °C) and lowest in stream-dwelling frogs (16.0 ± 6.2 °C). Heating tolerance was
also influenced by ontogenetic stage (-0.42 [-0.49:-0.35]; Fig. 4), with marginal influence by
ecotypes (Table S4). There was a strong phylogenetic effect on heating tolerance (λ = 0.86
[0.76:0.92]) and an effect of starting temperature where heating tolerance was lower at
higher starting temperature (-0.83 [-0.85:-0.82]).

*Figure 4. Heating tolerance (°C) decreases across ontogeny (stage 1–7, colours). Regression line and shaded area represents
overall model estimate + 95% CI. Heating tolerance is CT_{max} – starting temperature of the ramping assay (438 species).*

Thermal safety margin

Estimates of thermal safety margin vary with the spatial resolution of the environmental
temperature used in calculations, microhabitat characteristics, and the presence of
behavioural thermoregulation (Table 1, Fig. 5). Using macroclimate underestimates TSM for
fossorial and ground-dwelling frogs because the microclimate of these ecotypes are not well
captured by weather stations. Higher TSM using microclimate than macroclimate for a
fossorial species are generated by a lower soil temperature than air temperature. There is
little difference between using operative temperatures and microclimate because fossorial
frogs do not leave their burrow. TSM are similar using all spatial resolutions for arboreal
frogs because arboreal microhabitats represented by 1m shaded environments (60% canopy

cover) are similar to weather stations (e.g. 1.2m 90% canopy cover). A ground-dwelling frog

in an open, sunny habitat will encounter maximum air temperatures and exhibit maximum

body temperatures exceeding their upper critical temperature (negative TSM) at the

relevant microclimatic scale of the individual (1cm above ground in our simulations). When

allowed to select cooler microhabitats, predicted body temperatures of simulated ground-

dwelling frogs never exceed CTmax (positive TSM).

Table 1 Thermal Safety Margin (TSM; CTmax – maximum temperature; ⁰C) calculated using maximum temperatures of
varying spatial resolution for sympatric species representing three ecotypes: fossorial (Neobatrachus pictus), arboreal
(Litoria peronii) and ground-dwelling (Crinia signifera) species. Macroclimate: Maximum temperature of the warmest
month (WorldClim, BIO5). Microclimate: Maximum temperature from downscaled climate to representative microhabitat
conditions of the ecotype (NicheMapR). Operative temperature: maximum body temperature for a simulated 5g frog in
their preferred microhabitat.

Temperature	Fossorial	Arboreal	Ground-dwelling
Current macroclimate	-0.88	7.40	6.70
Future macroclimate	-4.58	3.60	2.90
Microclimate	5.62	6.14	-7.90
Operative temperature (no thermoregulation)	5.56	4.78	-12.36
Operative temperature (thermoregulating)	NA	NA	2.32

Figure 5 Thermal Safety Margin (TSM; CTmax – maximum temperature; ⁰C) calculated using maximum temperatures of
varying spatial resolution for sympatric species representing three ecotypes: fossorial (Neobatrachus pictus), arboreal
(Litoria peronii) and ground-dwelling (Crinia signifera) species. Macroclimate: Maximum temperature of the warmest
month (WorldClim, BIO5). Microclimate: Maximum temperature from downscaled climate to representative microhabitat
conditions of the ecotype (NicheMapR). Operative temperature: maximum body temperature for a simulated 5g frog in
their preferred microhabitat.

Discussion

Amphibians are an excellent system in which to investigate ontogenetic variation in thermal tolerance because they have a biphasic life cycle with life stages clearly delineated by metamorphosis. The delineation between life stages set by metamorphosis is key as metamorphosis may compartmentalise the life cycle and allow for a potential thermal bottleneck at one stage with minimal detrimental effects of temperature across a life cycle (Enriquez-Urzelai et al. 2019, Lowe et al. 2021).

We found striking differences between adults and larva across ecotypes in the opposite direction than expected if larva represented a thermal bottleneck. A similar general negative relationship between thermal tolerance over ontogeny is widely reported for frogs (Cupp 1980, Enriquez-Urzelai et al. 2019, Ruthsatz et al. 2022). Although lower thermal tolerances may be expected for aquatic larva, a higher thermal tolerance may be advantageous for small bodied larva that have reduced capacity to escape temperature extremes (Lenoir et al. 2017). Similarly, a wider selection of microhabitats in terrestrial environments may allow adults to compensate for a lower thermal tolerance through behavioural site selection than aquatic larva which grow in a thermally homogeneous environment (Enriquez-Urzelai et al. 2020).

We found ecotype mediates the magnitude of ontogenetic differences in thermal tolerances but is not an accurate proxy for environmental selection or constraint. The discrete nature of complex life cycles with metamorphosis and the need to survive thermal stress at one stage until the transition to the next stage may place high importance on the interaction between traits and the environment where a thermal bottleneck is a selective disadvantage (Lowe et al. 2021). Microhabitat preferences has been demonstrated to strongly correlate with thermal biology in reptiles (Greer 1980, Michelangeli et al. 2018), bees (Wilson et al. 2020), frogs (Duarte et al. 2012, Carilo Filho et al. 2021), and ants (Bujan et al. 2020, Leahy et al. 2022). We found strong phylogenetic conservatism in CT_{max} which likely reflects how ecotypes strongly correlate within Family (Moen and Wiens 2017). For example, Hylidae have a high proportion of arboreal species (Vidal-García and Keogh 2015). The use of ecotype as a proxy is complicated by variation in developmental characteristics that may not be consistent within Family or ecotypes to make generalisations. For example, direct-

developing larva (not available in our dataset) may occupy more similar microhabitats to

adults than species with an aquatic larva and fully terrestrial adults as discussed with the

limitations of larvae ecotypes. Egg thermal tolerance is rare in the Amphibian literature, but

eggs may also occupy vastly different microhabitats to larva and adults. For instance,

arboreal frogs may create foam nests in the canopy but the larva inhabit puddles on the

forest floor (Haddad and Prado 2005). While it would be possible to classify larval ecotypes

or egg microhabitats (Carilo Filho et al. 2021), this information is not readily available at the

scale of this study, and we encourage further investigation into these modulators. Doing so

may reveal whether diversity in developmental traits within ecotypes masks local scale

signals of environmental adaption via interactions with ecotype. Nevertheless, we found

fossorial species, on average, had higher thermal tolerance than other ecotypes.

The large overlap in thermal tolerance across ecotypes suggests developmental or genetic

constraints have a greater influence on the evolution of thermal tolerances than

environmental factors. Developmental or genetic constraints on climate niches or thermal

traits are suggested for reptiles and amphibians (Pie et al. 2017, Bodensteiner et al. 2021,

Carilo Filho et al. 2021). Several inferences support developmental or genetic constraints

underlie consistent differences between larva and adult thermal tolerance. First, an absence

for a bottleneck may indicate metamorphosis modulates thermal tolerances over ontogeny

(Lowe et al. 2021). Minimal carry over effects of development temperature on CT_{max} are

found between larva and adults (Enriquez-Urzelai et al. 2019). Physiological remodelling and

morphological development during metamorphosis are hypothesised to trade-off with

physiological processes and explain why CT_{max} decreases at metamorphic climax (Cupp

1980, Ruthsatz et al. 2022).

Second, if juveniles and adults are not drastically different in their physiology (i.e.,

biochemical processes are not affected by metamorphosis), then differences in thermal

tolerance may indicate differences in energy allocation between life stages. An energetics

interpretation depends on whether capacity to respond to a thermal challenge is considered

part of maintenance metabolic costs and the flexibility of reallocating energy. Perhaps, the

allocation of energy to reproduction at the adult stage trade-offs with the capacity to

increase maintenance metabolic rates in response to heating, whereas larva without a

reproductive metabolic cost have greater scope to rapidly respond to a thermal challenge at

the cost of allocating energy to growth. This explanation is in line with how the temperature

size rule emerges from differences between development and growth rates (Forster and

Hirst 2012, Rubalcaba and Olalla-Tárraga 2020, Gahm et al. 2021).

A higher physiological capacity to respond to change (i.e., plasticity) in larva is a

parsimonious explanation from an energetics perspective. In contrast with studies that

show decreasing acclimation capacity with increasing size (Rohr et al. 2018, Burton et al.

2020), *Rana temporaria* larva have lower acclimation capacity than adults (Ruthsatz et al.

2022). Assuming the cost of having a given thermal tolerance is the same between life

stages, low plasticity in thermal tolerance suggests larva continuously express physiological

mechanisms that permit higher CT_{max}, which is very costly (i.e., in maintenance metabolic

cost) during a time when they should be allocating energy to growth. One explanation for

low thermal plasticity is tadpoles have a narrow range of temperatures to select for

behavioural thermoregulation and thus primarily rely on physiological thermoregulation via

a higher CT_{max} to avoid heat stress (Casterlin and Reynolds 1978). As larva would have lower

absolute metabolic rate based on size allometry alone (in general), whether or not they can

offset an increased energetic cost with increased food intake or by other means can be

explored.

Third, if physiological mechanisms are different between life stages, then the differences

may reflect developmental constraints. Perhaps the physiological machinery during larval

growth covaries with greater capacity to respond to an environmental challenge, or that

larva have lower energetic costs for responding to thermal challenges, which allows them to

continuously express high CTmax at minimal cost to growth, in contrast with adults. This

explanation would resolve the paradox of the high metabolic costs of physiological

thermoregulation.

A notable characteristic of the amphibian CT_{max} literature is that there is a relative

consistency in methods to assay CT_{max} and the range of heating rates is narrow with a mode

of 1 °C min^{-1} (Hutchison 1961). Most studies in our literature search (2016-2022) assayed

CT_{max} using a ramping protocol rather than static protocols (zero studies described) or from

fitted thermal performance curves (three studies). In contrast, there is a much larger

diversity in heating rates reported for global ectotherm CT_{max} varying between 0.0007 and 4

°C min^{-1} (mean 0.5 °C min^{-1}) that reflects the tropical to polar origin of species (Morley et al.

2018). In studies on frogs, ontogenetic stage is confounded with heating rates as lower

heating rates are typically used for larva than adults or juveniles (Fig. S3). This different may

contribute towards nuisance heterogeneity in comparative study but the effects may be

minimal compared to other taxa (Noble et al. 2022). In our dataset, CT_{max} endpoints are

relatively consistent across studies and within ontogenetic stage, but ontogenetic stage was

confounded by endpoint definition. For example, onset of spasms or heat rigour were more

common for larva and loss of righting response is most common for adults. A model

invoking a hypothesised mechanism could be relevant for investigating specific endpoints or

definitions of upper critical temperatures.

Simulating body temperatures trades-off with making significant assumptions about the

microclimate characteristics of each ecotype and their thermal biology. For example,

arboreal frogs bask but we simulated they did not (Tracy et al. 2010). Using operative

temperature is best suited to species with well characterised thermal biology to derive

accurate TSM estimates and activity patterns at a local scale, for species whose body

temperatures are expected to be significantly different to environmental temperature, or

for large-scale studies that trade-off accuracy for scale. If the thermal biology of species is

known, then a microclimate-driven biophysical models of ectotherm body temperatures can

provide a fine-scale analysis of estimated activity times under warming scenarios and

hypothetical capacities to mitigate potential heat stress (Anderson et al. 2022).

A size-dependency of CTmax may explain why heating tolerances are relatively conserved

over ontogeny. Larger sizes may buffer adult frogs against higher heating rates via inertia

effects, thus a lower CTmax for an adult is equivalent to higher CTmax in smaller larva when

rate-time dynamics are considered. A lower size limit for thermal inertia in dry-skinned

ectotherms is approximately 1kg, which is larger than most frogs. A similar threshold for

frogs is unclear. While tadpoles are reported to be at thermal equilibrium with water

temperature, a small basking arboreal frog (<10g) can elevate their body temperature above

ambient temperature. Rates of heat exchange that influence body temperature changes in

frogs (a wet-skinned ectotherm) depends on resistance to water loss, size, and the

difference in thermal properties between water and air.

Microclimate and microhabitat preference strongly modulate estimates of climate

vulnerability. We demonstrate that spatial resolution of temperature strongly influences

estimated metrics of climate vulnerability. The magnitude of under- or over-prediction was

highly dependent on the microhabitat preferences of species (Table 1). Specifically, canopy

cover, which intercepts direct solar radiation, in the microclimate model set the ecological

accuracy of the simulated microhabitat and may differ from the selected microhabitats in

the field. In general, we suggest using simulated microclimates to estimate TSM than

macroclimate, particularly for large scale studies for which field measurements of

temperature are not feasible. Doing so relies on fewer assumptions than using a biophysical

model of the heat balance of a generalised frog to estimate operative temperatures,

requires one model instead of two for deriving environmental temperature to calculate

TSM, and may be a reasonable proxy for body temperature for an ecotype. It may be

reasonable to use soil temperature as a proxy for body temperature for fossorial frogs that

do not frequently leave their burrow.

Our simplified simulations clearly illustrate the importance of behavioural thermoregulation

(Bogert effect) and the availability of thermal refugia in the context of assessing thermal

safety margins (Kearney 2013, Sunday et al. 2014). At face value, our results suggest for a

given location, ground dwelling frogs are highly vulnerable to climate change, but the frog

was not simulated to avoid temperatures above their activity thresholds. Several studies

demonstrated microclimate is a better predictor of frog CTmax than microclimate

(Gutiérrez-Pesquera et al. 2016, Katzenberger et al. 2018, Pintanel et al. 2019), and that

variation in tadpole CTmax is explained by local scale water temperatures measured using

data loggers (Bonino et al. 2020, Pintanel et al. 2022), or simulated microclimate

(Oyamaguchi et al. 2018). Understanding environmental tolerance across all life stages is

crucial to identify vulnerability in for organisms with complex life cycles (Lowe et al. 2021).

The differences in thermal tolerance between aquatic larval stages and mobile adult stages

suggested that there is an evolutionary capacity to maximise performance to the

environment for a given life stage. However, the potential to fully decouple detrimental

environmental effects on fitness at one life stage to the next (i.e. carry-over effects) in

amphibians needs to be explored at a comparative global scale.

References

References

Anderson, R. O., S. Meiri, and D. G. Chapple. 2022. The biogeography of warming tolerance
in lizards. Journal of Biogeography **n/a**:1274-1285.
Bates, A. E., and S. A. Morley. 2020. Interpreting empirical estimates of experimentally
derived physiological and biological thermal limits in ectotherms. Canadian Journal
of Zoology **98**:237-244.
Bodensteiner, B. L., G. A. Agudelo-Cantero, A. Z. A. Arietta, A. R. Gunderson, M. M. Muñoz, J.
M. Refsnider, and E. J. Gangloff. 2021. Thermal adaptation revisited: How conserved
are thermal traits of reptiles and amphibians? Journal of Experimental Zoology Part
A: Ecological and Integrative Physiology **335**:173-194.
Bonino, M. F., F. B. Cruz, and M. G. Perotti. 2020. Does temperature at local scale explain
thermal biology patterns of temperate tadpoles? Journal of Thermal Biology
94:102744.
Brattstrom, B. H. 1970. Thermal acclimation in Australian amphibians. Comparative
Biochemistry and Physiology **35**:69-103.
Bujan, J., K. A. Roeder, K. de Beurs, M. D. Weiser, and M. Kaspari. 2020. Thermal diversity of
North American ant communities: Cold tolerance but not heat tolerance tracks
ecosystem temperature. Global Ecology and Biogeography **29**:1486-1494.
Burton, T., H.-K. Lakka, and S. Einum. 2020. Acclimation capacity and rate change through
life in the zooplankton Daphnia. Proceedings of the Royal Society B: Biological
Sciences **287**:20200189.
Carilo Filho, L. M., B. T. de Carvalho, B. K. A. Azevedo, L. M. Gutiérrez-Pesquera, C. V. Mira-
Mendes, M. Solé, and V. G. D. Orrico. 2021. Natural history predicts patterns of
thermal vulnerability in amphibians from the Atlantic Rainforest of Brazil. Ecology
and Evolution **11**:16462-16472.
Casterlin, M. E., and W. W. Reynolds. 1978. Behavioural thermoregulation in Rana pipiens
tadpoles. Journal of Thermal Biology **3**:143-145.
Cheng, C.-T., M.-F. Chuang, T. Haramura, C.-B. Cheng, Y. I. Kim, A. Borzée, C.-S. Wu, Y.-H.
Chen, Y. Jang, N. C. Wu, and Y.-C. Kam. 2022. Open habitats increase vulnerability of
amphibian tadpoles to climate warming across latitude. Global Ecology and
Biogeography **n/a**.
Chown, S. L., and K. J. Gaston. 2016. Macrophysiology – progress and prospects. Functional
Ecology **30**:330-344.
Collin, R., A. P. Rebolledo, E. Smith, and K. Y. K. Chan. 2021. Thermal tolerance of early
development predicts the realized thermal niche in marine ectotherms. Functional
Ecology **35**:1679-1692.
Cupp, P. V. 1980. Thermal Tolerance of Five Salientian Amphibians during Development and
Metamorphosis. Herpetologica **36**:234-244.
Dahlke, F. T., S. Wohlrab, M. Butzin, and H.-O. Pörtner. 2020. Thermal bottlenecks in the life
cycle define climate vulnerability of fish. Science **369**:65-70.
Deutsch, C. A., J. J. Tewksbury, R. B. Huey, K. S. Sheldon, C. K. Ghalambor, D. C. Haak, and P.
R. Martin. 2008. Impacts of climate warming on terrestrial ectotherms across
latitude. Proceedings of the National Academy of Sciences of the United States of
America **105**:6668-6672.
Duarte, H., M. Tejedo, M. Katzenberger, F. Marangoni, D. Baldo, J. F. Beltrán, D. A. Martí, A.
Richter-Boix, and A. Gonzalez-Voyer. 2012. Can amphibians take the heat?

Vulnerability to climate warming in subtropical and temperate larval amphibian
communities. Global Change Biology **18**:412-421.

Enriquez-Urzelai, U., M. Sacco, A. S. Palacio, P. Pintanel, M. Tejedo, and A. G. Nicieza. 2019.
Ontogenetic reduction in thermal tolerance is not alleviated by earlier
developmental acclimation in Rana temporaria. Oecologia **189**:385-394.

Enriquez-Urzelai, U., R. Tingley, M. R. Kearney, M. Sacco, A. S. Palacio, M. Tejedo, and A. G.
Nicieza. 2020. The roles of acclimation and behaviour in buffering climate change
impacts along elevational gradients. Journal of Animal Ecology **89**:1722-1734.

Fick, S. E., and R. J. Hijmans. 2017. WorldClim 2: new 1-km spatial resolution climate
surfaces for global land areas. International Journal of Climatology **37**:4302-4315.

Forster, J., and A. G. Hirst. 2012. The temperature-size rule emerges from ontogenetic
differences between growth and development rates. Functional Ecology **26**:483-492.

Gahm, K., A. Z. A. Arietta, and D. K. Skelly. 2021. Temperature-mediated trade-off between
development and performance in larval wood frogs (Rana sylvatica). Journal of
Experimental Zoology Part A: Ecological and Integrative Physiology **335**:146-157.

Garcia, R. A., J. L. Allen, and S. Clusella-Trullas. 2019. Rethinking the scale and formulation of
indices assessing organism vulnerability to warmer habitats. Ecography **42**:1024-
1036.

González-del-Pliego, P., B. R. Scheffers, R. P. Freckleton, E. W. Basham, M. B. Araújo, A. R.
Acosta-Galvis, C. A. Medina Uribe, T. Haugaasen, and D. P. Edwards. 2020. Thermal
tolerance and the importance of microhabitats for Andean frogs in the context of
land use and climate change. Journal of Animal Ecology **89**:2451-2460.

Greer, A. 1980. Critical Thermal Maximum Temperatures in Australian Scincid Lizards: Their
Ecological and Evolutionary Significance. Australian Journal of Zoology **28**:91-102.

Gutiérrez-Pesquera, L. M., M. Tejedo, M. Á. Olalla-Tárraga, H. Duarte, A. Nicieza, and M.
Solé. 2016. Testing the climate variability hypothesis in thermal tolerance limits of
tropical and temperate tadpoles. Journal of Biogeography **43**:1166-1178.

Haddad, C. F. B., and C. P. A. Prado. 2005. Reproductive Modes in Frogs and Their
Unexpected Diversity in the Atlantic Forest of Brazil. Bioscience **55**:207-217.

Hutchison, V. H. 1961. Critical Thermal Maxima in Salamanders. Physiological Zoology **34**:92-
125.

IUCN. 2022. The IUCN Red List of Threatened Species.

Jetz, W., and R. A. Pyron. 2018. The interplay of past diversification and evolutionary
isolation with present imperilment across the amphibian tree of life. Nature Ecology
& Evolution **2**:850-858.

Katzenberger, M., J. Hammond, M. Tejedo, and R. Relyea. 2018. Source of environmental
data and warming tolerance estimation in six species of North American larval
anurans. Journal of Thermal Biology **76**:171-178.

Kearney, M. R. 2013. Activity restriction and the mechanistic basis for extinctions under
climate warming. Ecology Letters **16**:1470-1479.

Kearney, M. R., and W. P. Porter. 2017. NicheMapR - an R package for biophysical modelling:
the microclimate model. Ecography **40**:664-674.

Kearney, M. R., and W. P. Porter. 2019. NicheMapR – an R package for biophysical
modelling: the ectotherm and Dynamic Energy Budget models. Ecography **43**:85-96.

Kingsolver, J. G., and L. B. Buckley. 2020. Ontogenetic variation in thermal sensitivity shapes
insect ecological responses to climate change. Current Opinion in Insect Science
41:17-24.

Klockmann, M., F. Günter, and K. Fischer. 2017. Heat resistance throughout ontogeny: body
size constrains thermal tolerance. Global Change Biology **23**:686-696.
Leahy, L., B. R. Scheffers, S. E. Williams, and A. N. Andersen. 2022. Arboreality drives heat
tolerance while elevation drives cold tolerance in tropical rainforest ants. Ecology
103:e03549.
Lenoir, J., T. Hattab, and G. Pierre. 2017. Climatic microrefugia under anthropogenic climate
change: implications for species redistribution. Ecography **40**:253-266.
Levy, O., L. B. Buckley, T. H. Keitt, C. D. Smith, K. O. Boateng, D. S. Kumar, and M. J.
Angilletta, Jr. 2015. Resolving the life cycle alters expected impacts of climate
change. Proceedings: Biological Sciences **282**:20150837.
Lowe, W. H., T. E. Martin, D. K. Skelly, and H. A. Woods. 2021. Metamorphosis in an Era of
Increasing Climate Variability. Trends in Ecology & Evolution **36**:360-375.
Michelangeli, M., C. T. Goulet, H. S. Kang, B. B. M. Wong, and D. G. Chapple. 2018.
Integrating thermal physiology within a syndrome: Locomotion, personality and
habitat selection in an ectotherm. Functional Ecology **32**:970-981.
Moen, D. S., E. Cabrera-Guzman, I. W. Caviedes-Solis, E. Gonzalez-Bernal, and A. R. Hanna.
2022. Phylogenetic analysis of adaptation in comparative physiology and
biomechanics: overview and a case study of thermal physiology in treefrogs. Journal
of Experimental Biology **225**:jeb243292.
Moen, D. S., and J. J. Wiens. 2017. Microhabitat and Climatic Niche Change Explain Patterns
of Diversification among Frog Families. The American Naturalist **190**:29-44.
Morley, S. A., L. S. Peck, J. Sunday, S. Heiser, and A. E. Bates. 2018. Acclimation potential of
global ectothermic species, collated from literature, 1960 to 2015. Natural
Environment Research Council, Cambridge, UK, Polar Data Centre.
Moyano, M., C. Candebat, Y. Ruhbaum, S. Álvarez-Fernández, G. Claireaux, J.-L. Zambonino-
Infante, and M. A. Peck. 2017. Effects of warming rate, acclimation temperature and
ontogeny on the critical thermal maximum of temperate marine fish larvae. PLOS
ONE **12**:e0179928.
Noble, D. W. A., P. Pottier, M. Lagisz, S. Burke, S. M. Drobniak, R. E. O'Dea, and S. Nakagawa.
2022. Meta-analytic approaches and effect sizes to account for 'nuisance
heterogeneity' in comparative physiology. Journal of Experimental Biology
225:jeb243225.
Oyamaguchi, H. M., P. Vo, K. Grewal, R. Do, E. Erwin, N. Jeong, K. Tse, C. Chen, M. Miyake, A.
Lin, and M. Gridi-Papp. 2018. Thermal sensitivity of a Neotropical amphibian
(Engystomops pustulosus) and its vulnerability to climate change. Biotropica **50**:326-
337.
Pandori, L. L. M., and C. J. B. Sorte. 2019. The weakest link: sensitivity to climate extremes
across life stages of marine invertebrates. Oikos **128**:621-629.
Pie, M. R., L. L. F. Campos, A. L. S. Meyer, and A. Duran. 2017. The evolution of climatic
niches in squamate reptiles. Proceedings: Biological Sciences **284**.
Pincebourde, S., and J. Casas. 2015. Warming tolerance across insect ontogeny: influence of
joint shifts in microclimates and thermal limits. Ecology **96**:986-997.
Pincebourde, S., and H. A. Woods. 2020. There is plenty of room at the bottom:
microclimates drive insect vulnerability to climate change. Current Opinion in Insect
Science **41**:63-70.

Pinsky, M. L., A. M. Eikeset, D. J. McCauley, J. L. Payne, and J. M. Sunday. 2019. Greater
vulnerability to warming of marine versus terrestrial ectotherms. Nature **569**:108-
111.
Pintanel, P., M. Tejedo, A. Merino-Viteri, F. Almeida-Reinoso, S. Salinas-Ivanenko, A. C.
Lopez-Rosero, G. A. Llorente, and L. M. Gutierrez-Pesquera. 2022. Elevational and
local climate variability predicts thermal breadth of mountain tropical tadpoles.
Ecography **2022**:e05906.
Pintanel, P., M. Tejedo, S. R. Ron, G. A. Llorente, and A. Merino-Viteri. 2019. Elevational and
microclimatic drivers of thermal tolerance in Andean Pristimantis frogs. Journal of
Biogeography **46**:1664-1675.
Rohr, J. R., D. J. Civitello, J. M. Cohen, E. A. Roznik, B. Sinervo, and A. I. Dell. 2018. The
complex drivers of thermal acclimation and breadth in ectotherms. Ecology Letters
21:1425-1439.
Rubalcaba, J. G., and M. Á. Olalla-Tárraga. 2020. The biogeography of thermal risk for
terrestrial ectotherms: Scaling of thermal tolerance with body size and latitude.
Journal of Animal Ecology **n/a**:1277-1285.
Ruthsatz, K., K. H. Dausmann, M. A. Peck, and J. Glos. 2022. Thermal tolerance and
acclimation capacity in the European common frog (Rana temporaria) change
throughout ontogeny. J Exp Zool A Ecol Integr Physiol **337**:477-490.
Sinclair, B. J., K. E. Marshall, M. A. Sewell, D. L. Levesque, C. S. Willett, S. Slotsbo, Y. Dong, C.
D. Harley, D. J. Marshall, B. S. Helmuth, and R. B. Huey. 2016. Can we predict
ectotherm responses to climate change using thermal performance curves and body
temperatures? Ecology Letters **19**:1372-1385.
Sunday, J. M., A. E. Bates, M. R. Kearney, R. K. Colwell, N. K. Dulvy, J. T. Longino, and R. B.
Huey. 2014. Thermal-safety margins and the necessity of thermoregulatory behavior
across latitude and elevation. Proceedings of the National Academy of Sciences of
the United States of America **111**:5610-5615.
Tracy, C. R., K. A. Christian, and C. R. Tracy. 2010. Not just small, wet, and cold: effects of
body size and skin resistance on thermoregulation and arboreality of frogs. Ecology
91:1477-1484.
Truebano, M., P. Fenner, O. Tills, S. D. Rundle, and E. L. Rezende. 2018. Thermal strategies
vary with life history stage. Journal of Experimental Biology **221**:jeb171629.
Vidal-García, M., and J. S. Keogh. 2015. Convergent evolution across the Australian
continent: ecotype diversification drives morphological convergence in two distantly
related clades of Australian frogs. Journal of Evolutionary Biology **28**:2136-2151.
Wilson, E. S., C. E. Murphy, J. P. Rinehart, G. Yocum, and J. H. Bowsher. 2020. Microclimate
Temperatures Impact Nesting Preference in Megachile rotundata (Hymenoptera:
Megachilidae). Environmental Entomology.
Wu, N. C., R. P. Bovo, U. Enriquez-Urzelai, S. Clusella-Trullas, M. R. Kearney, C. A. Navas, and
J. D. Kong. 2024. Global exposure risk of frogs to increasing environmental dryness.
Nature Climate Change **14**.

586