T cell landscape definition by multi-omics identifies Galectin-9 as novel immunotherapy target in chronic lymphocytic leukemia

Llaó Cid L[1,*], Wong JKL[1,*], Fernandez Botana I[2,3], Paul Y[1], Wierz M[2], Flörchinger, A[1,4], Gonder, S[2,3], Pagano G[2,3], Chazotte M[5], Bestak K[5], Schifflers C[1], Iskar M[1], Roider T[6,7,8], Mallm JP[9], Cosma, A[10], Campton DE[11], Gerhard-Hartmann E[12], Rosenwald A[12], Colomer D[13], Campo E[13], Schapiro D[5,14], Dietrich S[6], Lichter P[1], Moussay E[2,#], Paggetti J[2,#], Zapatka M[1,#], Seiffert M[1,#,§]

[1]Division of Molecular Genetics, German Cancer Research Center, Heidelberg, Germany;
[2]Tumor Stroma Interactions, Department of Cancer Research, Luxembourg Institute of Health, Luxembourg, Luxembourg;
[3]Faculty of Science, Technology and Medicine, University of Luxembourg, Esch-sur-Alzette, Luxembourg;
[4]Faculty of Biosciences, University of Heidelberg, Heidelberg, Germany;
[5]Institute for Computational Biomedicine, Heidelberg University Hospital, Heidelberg, Germany;
[6]European Molecular Biology Laboratory (EMBL), Heidelberg, Germany;
[7]Molecular Medicine Partnership Unit (MMPU), Heidelberg, Germany;
[8]Department of Medicine V, Hematology, Oncology and Rheumatology, University of Heidelberg, Heidelberg, Germany;
[9]scOpenLab, German Cancer Research Center, Heidelberg, Germany;
[10]National Cytometry Platform, Luxembourg Institute of Health, Luxembourg, Luxembourg;
[11]RareCyte, Seattle, WA, USA;
[12]Institute for Pathology, University of Würzburg, Würzburg, Germany;
[13]Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Hematopathology Unit, Hospital Clinic, CIBERONC, Barcelona, Spain;
[14]Institute for Pathology, Heidelberg University Hospital, Heidelberg, Germany;

* equal contribution
# equal contribution
§ corresponding author

Current address of Marina Wierz: MVZ für Histologie, Zytologie und Molekulare Diagnostik Trier, 54296 Trier, Germany

Contact information for correspondence:

Martina Seiffert

Molecular Genetics (B060), German Cancer Research Center (DKFZ)

Im Neuenheimer Feld 280, 69120 Heidelberg, Germany

Tel.: +49-6221-42 4586, Fax: +49-6221-42 4639

Email: m.seiffert@dkfz.de

Keywords: CLL; single-cell RNA-seq; mass cytometry; T cell exhaustion; tumor

microenvironment; Galectin-9; TIM3

Abstract

Failure of immunotherapy after applying checkpoint inhibitors or CAR-T cells is linked to T cell exhaustion. Here, we explored the T cell landscape in chronic lymphocytic leukemia (CLL) by single-cell omics analyses of blood, bone marrow and lymph node samples of patients and spleen samples of a CLL mouse model. By single-cell RNA-sequencing, mass cytometry (CyTOF), and multiplex image analysis of tissue microarrays, we defined the spectrum of phenotypes and transcriptional programs of T cells and and their differentiation state trajectories. We identified disease-specific accumulation of distinct regulatory T cell subsets and T cells harboring an exhausted phenotype exclusively in the CLL lymph node tissue. Integration of TCR data revealed a clonal expansion of CD8[+] precursor exhausted T cells, suggesting their reactivity for CLL cells. Interactome analyses identified the TIM3 ligand Galectin-9 as novel immunoregulatory molecule in CLL. Blocking of Galectin-9 in CLL-bearing mice slowed down disease development and reduced the number of TIM3 expressing T cells. Galectin-9 expression correlated with shorter survival of CLL patients. Thus, Galectin-9 contributes to immune escape in CLL and represents a novel target for immunotherapy.

Introduction

Failure of response to immune checkpoint inhibitors (ICI) is commonly seen in cancer patients with unclear causes and predictors which limits the use of these novel immunotherapy approaches. B cell lymphomas develop in lymph nodes (LNs), the site of T cell priming and activation in infections and cancer. Even though there is a constant interaction of malignant B cells with CD4[+] and CD8[+] T cells in this tissue, adaptive immune control fails and response to ICI is very limited in patients with B cell lymphomas. Very low response rates to anti-PD1 antibodies were observed in patients with chronic lymphocytic leukemia (CLL) (Ding et al., 2017; Xu-Monette et al., 2018), which is a non-Hodgkin lymphoma of mature B cells that accumulate in LNs and blood. In contrast to patients with acute lymphocytic leukemia (ALL), also response rates to CD19 CAR-T cell therapy remain below the expectations in CLL (Cappell et al., 2020). A lack of fit and functional T cells is discussed as a main limitation explaining the failure of immunotherapy responses in CLL and beyond (Fraietta et al., 2018; van Bruggen et al., 2019). Chronic exposure of T cells to tumor (neo)antigens and an immune suppressive microenvironment lead to their exhaustion and dysfunction (Blank et al., 2019). Whereas terminally exhausted T cells, characterized by high PD1 expression levels and expression of TOX, fail to reactivation by ICI, their precursor state that expresses lower levels of PD1 and is positive for TCF-7, harbors self-renewal capacity and the ability to control tumor development upon ICI treatment (Im et al., 2016; Scott et al., 2019; Utzschneider et al., 2016). The presence of these cell states has been confirmed in CLL patients and mouse models (Hanna et al., 2021), even though a characterization of T cells in CLL LN is lacking due to the limited access to such samples. Current efforts aim at characterizing mechanisms of T cell exhaustion in cancer and at identifying novel therapeutic targets to overcome this limitation.

Here, we used single-cell RNA sequencing (scRNA-seq) and mass cytometry (CyTOF) of LNs, peripheral blood (PB) and bone marrow (BM) samples of patients with CLL, as well as tumor-free reactive lymph nodes (rLN) from healthy controls, combined with multiplex imaging of LN tissue sections and microarrays to characterize the distribution, phenotype, and function of T cells under chronic exposure to cancer. We show that CLL LNs constitute a unique niche where clonally expanded CD8[+] T cells expressing CD39 and with an exhausted phenotype are enriched in comparison to healthy LNs. We further defined cells with a precursor exhausted signature that accumulate in CLL LNs, as well as several types of regulatory T cells. The results of interactome analyses led us to target Galectin-9 in a CLL mouse model which resulted in improved T cell function and an attenuated tumor development.

Results

Definition of the T cell landscape in CLL using mass cytometry

In order to comprehensively characterize the T cell compartment associated with CLL, a

large-scale and high-dimensional analysis of T cells from CLL patients as well as healthy

controls (HC) was conducted using mass cytometry, scRNA-seq and multiplex imaging of

tissue sections followed by integrative data analysis (Figure 1A). The patient cohort reflects

the heterogeneity of CLL in respect to cell-of-origin, including both *IGHV* mutated (n=10) and

unmutated (n=10) cases, as well as patients with the major genetic aberrations (Suppl. Table

1). We performed mass cytometry profiling of T cells from 22 CLL LNs, plus 7 paired PB and

3 paired BM samples, as well as 13 reactive LNs (rLNs) from HC using a panel of 42

antibodies (Key Resource Table) designed to identify and characterize naïve, memory,

effector, regulatory and exhausted T cells. The analysis comprised 5.29 x10^6 T cells, with a

median of 51,937 cells per sample (Suppl. Figure 1A-C). Unsupervised graph-based

clustering based on the differential expression of the analyzed proteins grouped cells into 30

clusters which are presented as uniform manifold approximation and projection (UMAP) plot

(Figure 1B, C, Suppl. Figure 2A). This approach led to the identification of 15 CD4$^+$ T cell

clusters, 9 CD8$^+$ T cell clusters, 4 double-negative T cell clusters, and 2 clusters with a

mixture of CD4$^+$ and CD8$^+$ T cells (Figure 1D, Suppl. Figure 2B, 3A, and Suppl. Table 2). The

expression of CD45RA, CD45RO, and CCR7 allowed for a general classification of T cells

into naïve (T$_N$, CD45RA$^+$ CD45RO$^-$ CCR7$^+$), central memory (T$_{CM}$, CD45RA$^-$ CD45RO$^+$

CCR7$^+$), effector memory (T$_{EM}$, CD45RA$^-$ CD45RO$^+$ CCR7$^-$) and effector cells (T$_{EF}$, CD45RA$^+$

CD45RO$^-$ CCR7$^-$). The CD8$^+$ T cell subsets comprised 1 naïve (CD8 T$_N$), 1 central memory

(CD8 T$_{CM}$), 3 subsets with a short-lived effector cell (SLEC) phenotype expressing TBET and

KLRG1 (CD8 T$_{EF}$ TBET, CD8 T$_{EM}$ TBET, and CD8 T$_{EF}$ NCAM) (Joshi et al., 2007; Obar et al.,

2011), and 4 effector memory subsets with a higher expression of the cytotoxic molecule

granzyme K (GZMK) and an increasing expression of exhaustion-related molecules such as

the inhibitory receptors PD1 and TIGIT, the TFs EOMES and TOX, and the ectoenzymes

CD38 and CD39 (CD8 T$_{EM}$ GZMK, CD8 T$_{EM}$ TBET GZMK, CD8 T$_{EX}$ CD38, and CD8 T$_{EX}$

CD39) (McLane et al., 2019). The CD4$^+$ T cell subsets comprised 1 naïve (CD4 T$_N$), 3 central

memory (CD4 T$_{CM1}$ CCR7, CD4 T$_{CM2}$ CD25, and CD4 T$_{CM3}$), 1 follicular helper (T$_{FH}$), and 6

effector memory subsets. Of these, two expressed TBET and KLRG1, resembling SLEC

subsets (CD4 T$_{EM}$ NCAM, and CD4 T$_{EM}$ TBET), while 3 of them expressed higher levels of

PD1 and TIGIT (CD4 T$_{EM}$ PD1), CTLA4 and CD38 (CD4 T$_{EM}$ CTLA4), or CD39 (CD4 T$_{EM}$

CD39). The last CD4 T$_{EM}$ subset expressed EOMES, GZMK and PD1 (CD4 T$_{EM}$ GZMK), and

shares therefore characteristics of T regulatory type 1 cells (T$_R$1) and CD4$^+$ T cells with

cytotoxic properties (Oh and Fong, 2021; Roessner et al., 2021). In addition, FOXP3

expression identified 4 T$_{REG}$ clusters within the CD4$^+$ T cells, 2 with a central memory

phenotype ($T_{REG-CM1}$ and $T_{REG-CM2}$), and 2 subsets with an activated inhibitory phenotype

(T_{REG} PD1 and T_{REG} CD39). CD4 and CD8 double negative (DN) $CD3^+$ T cells comprised 2

clusters with a phenotype similar to CD8 SLEC subsets (DN T_{EM} KLRG1 and DN T_{EF} TBET),

and 2 cytotoxic clusters expressing GZMK and EOMES as well as CD38 (DN T_{EF} CD38) or

the transcription factor HELIOS (DN T_{EM} HELIOS). A $KI67^+$ proliferative subset (T_{PR}), as well

as a cluster with high expression of ICOS (T_{EM} ICOS) were both composed of $CD4^+$ and

$CD8^+$ T cells.

In order to identify relationships between the T cell populations present in the CLL LNs,

we correlated the subset frequencies of all individual CLL LN samples (n = 20, excluding 2

duplicate samples, Figure 1E). Naïve $CD4^+$ and $CD8^+$ T cells as well as $CD4^+$ and $CD8^+$ T

cells with a SLEC phenotype were significantly associated with each other, indicating that the

LN tumor microenvironment (TME) is similarly influencing the differentiation of these cell

states both in $CD4^+$ and $CD8^+$ T cells. Spatial analyses of immune cell populations by

multiplex immunofluorescence staining of a tissue microarray (TMA) of 42 CLL LN samples

confirmed the presence and colocalization of malignant B cells with T cells in the tissue

(Figure 1F, Suppl. Figure 3B-C). Quantification of the frequencies of the major immune cell

subsets in the tissues revealed a positive correlation for $CD8^+$ $PD1^-$ and CD8 $PD1^+$ T cells,

for myeloid cells and T_{REG}, and the strongest correlation for T_{REG} and $PD1^+$ $CD8^+$ T cells

(Figure 1G, Suppl. Figure 3D) which is in line with a role of T_{REG} in promoting the

accumulation of dysfunctional $CD8^+$ T cells (Sawant et al., 2019).

CLL LNs are a distinct milieu enriched with immunosuppressive T cell phenotypes

We next examined the distribution of T cell immunophenotypes across the three different

tissues analyzed in CLL. The sample composition of PB and BM was similar, but differed

from LN samples (Figure 2A, B, Suppl. Figure 4A), suggesting that tissue cues significantly

determine T cell composition. PB and BM samples were enriched in SLEC T cells, such as

CD8 T_{EM} TBET, CD4 T_{EM} TBET, and DN T_{EF} TBET, while CLL LNs contained higher

frequencies of exhausted T cells (CD8 T_{EX} CD39 and CD8 T_{EX} CD38), T_R1 cells, T_{FH} cells,

several T_{REG} subsets ($T_{REG-CM1}$, T_{REG} PD1, and T_{REG} CD39), as well as proliferating cells (T_{PR})

(Figure 2C). Even though the abundancies among cell clusters within the tissue types

correlated, no significant associations of the frequencies of cell subsets between PB and LN

samples of the same patient were identified, underscoring a differential composition of T cell

subsets between tissues (Suppl. Figure 4B).

The comparison of cell subset frequencies between CLL LNs and rLNs revealed that

higher percentages of cytotoxic CD8 T_{EM} GZMK as well as exhausted $CD8^+$ T cells (CD8 T_{EX}

CD39 and CD8 T_{EX} CD38) were present in CLL LNs (Figure 2D). Within the CD4

compartment, T_R1 cells and all T_{REG} clusters ($T_{REG-CM1}$, $T_{REG-CM2}$, T_{REG} PD1, and T_{REG} CD39)

were also enriched in CLL LNs, in addition to proliferating T_{PR} cells and DN T_{EM} HELIOS cells

(Figure 2D). Collectively, these results suggest that CLL LNs constitute a distinct niche

where malignant B cell accumulation induces an immunosuppressive microenvironment with

an enrichment of regulatory T cells and exhausted cytotoxic T cells.

Next, we examined the relation of clinical features to a distinct T cell composition in the

CLL LNs. We detected a strong negative association between age of the patients and the

frequency of naïve $CD8^+$ T cells (Suppl. Figure 4C, Suppl. Table 3) which is in line with

published data (Gupta et al., 2004). In addition, the frequencies of 2 T_{REG} clusters ($T_{REG\ CM2}$

and T_{REG} PD1) showed a positive correlation with age (Suppl. Figure 4D, E). We did not

observe significant associations between the frequencies of T cell clusters with sex or the

clinical stage of the patients (Suppl. Table 3).

Definition of T cell exhaustion states in CLL LNs by single-cell RNA-sequencing

Next, single-cell RNA-seq of T cells from CLL LNs was performed to characterize their

transcriptional profile and clonal diversity. $CD3^+$ T cells and CLL cells (as small spike-in

population) from 5 CLL LN samples were FACS-sorted and subjected to scRNA-seq using

the 10X Genomics platform, yielding 8,825 cells with an average of 4,577 reads per cell and

a median of 1,475 genes per cell (STAR methods, Suppl. Figure 5A). Using unsupervised

graph-based clustering, we identified 13 clusters presented as UMAP plot (Figure 3A, STAR

methods), for which we assigned an identity based on the differential gene expression of

each cluster compared to the rest of the cells (Figure 3B). All cell clusters were shared

among patients although at different proportions (Figure 3C, Suppl. Figure 5B). Clusters

were defined as CLL cells, a small cluster of dendritic cells (DCs), 6 $CD4^+$ T cell subsets, 4

$CD8^+$ T cell subsets, and a $KI67^+$ proliferating T cell cluster (T_{PR}) which contained both $CD4^+$

and $CD8^+$ T cells (Figure 3A, B). More specifically, both $CD4^+$ and $CD8^+$ naïve (CD4 T_N and

CD8 T_N) T cells were defined based on the expression of marker genes such as *SELL*,

CCR7 and *IL7R*; regulatory $CD4^+$ T cells (T_{REG}) were identified by high expression of *FOXP3*,

IL2RA and *IKZF2*; a heterogeneous cluster of cells containing mainly follicular helper $CD4^+$ T

cells (T_{FH}) was defined by the expression of *CXCR5*, *CD200*, *TOX2* and *TOX* (Suppl. Figure

5C, D), the small proportion of $CD8^+$ T cells in this cluster is in line with a shared

transcriptional program of exhausted $CD8^+$ T cells with T_{FH} (Im et al., 2016); and

conventional $CD4^+$ T cells with a central memory phenotype were identified by the high

expression of *IL7R, CD40LG, KLRB1* as well as *CD69* (CD4 T_H CD69) or *ITGB1* and *KLF2*

(CD4 T_H KLF2) in accordance with published data of other B cell lymphoma (Roider et al.,

2022). A heterogeneous cluster mostly comprising $CD4^+$ T cells expressed memory marker

and effector molecule genes such as *GZMK*, resembling the CD4 T_{EM} GZMK subset that

shares features with T_R1-like cells, identified by mass cytometry (Figure 1B). $CD8^+$ effector

memory subsets were characterized by a higher expression of effector molecule genes like
CCL5, NKG7, PRF1, *GZMA* and *GZMK*, with one subset expressing generally lower levels of
these genes and high levels of *LYAR* (CD8 T_{EM} LYAR), another subset expressing higher
levels of *CCL5* (CD8 T_{EM} CCL5) which resembles the CD8 T_{EM} GZMK cluster identified by
mass cytometry (Figure 1B), and a third showing elevated expression of genes related to
exhaustion such as *HAVCR2* (coding for TIM3), *LAG3, ENTPD1* (CD39), *PDCD1* (PD1),
CTLA4 and *TIGIT* (CD8 T_{EX}), and therefore resembling the two T_{EX} clusters defined in the
CyTOF data by high expression of these exhaustion markers (Figure 1B). Notably, the later
subset expressed highest levels of the exhaustion signature from Zheng *et al.* (Zheng et al.,
2021) (Figure 3D) confirming its dysfunctional state, while CD8 T_{EM} CCL5 cells presented the
highest score of the precursor exhausted signatures from Guo *et al.* (Guo et al., 2018)
(Figure 3E) and Andreatta M *et al.* (Andreatta et al., 2021) (Suppl. Figure 5E) suggesting that
these cells constitute a precursor state of exhausted CD8$^+$ T cells. Subset clustering and
definition were in agreement with the annotation from the pbmc3k signatures from Hao *et al.*
(Hao et al., 2021) supporting the accuracy of our annotations (Suppl. Figure 5F). In addition,
the frequency of the main T cell clusters was comparable and positively correlated between
scRNA-seq and mass cytometry in the 4 samples analyzed by both techniques (Suppl.
Figure 5G, H), underscoring the robustness of our data.

To delineate the relationship between T cell states, we performed trajectory analyses of
CD4$^+$ and CD8$^+$ T cells by destiny (Angerer et al., 2016). When defining CD8 T_N cells as the
starting point of the differentiation path, we observed a divarication into two branches, one
containing CD8 T_{EM} LYAR cells and the other CD8 T_{EM} CCL5 cells, from which T_{EX} cells
diverged and constituted a terminal state (Figure 3F, Suppl. Figure 5I). When inspecting the
differentiation path from CD8 T_N to CD8 T_{EM} LYAR, four clusters of differentially expressed
genes (DEGs) were identified along the trajectory. Naïve marker genes (*LEF1, SELL*) were
highly expressed in the first cluster, the second included the transcription factor *KLF2* and
IL6R, the third activation marker genes (*CD69* and *DUSP1*), and the fourth effector molecule
genes (*GZMH, NKG7, GZMA and FAS*) (Figure 3G). The differentiation path from CD8 T_N to
CD8 T_{EM} CCL5 and CD8 T_{EX} was also comprised of four clusters of DEGs, with naïve marker
genes (*SELL* and *LEF1*, and *IL7R* and *CCR7*, respectively) expressed at the beginning,
followed by genes associated with activation (such as *CD69*) and cytotoxic function (*GZMK*),
and genes related to exhaustion (*LAG3, TIGIT, EOMES, CD38* and *HAVCR2*) that were
highest at the end of the trajectory (Figure 3G). This data suggests GZMK$^+$ CD8$^+$ T cells as a
precursor state of terminally exhausted CD8$^+$ T cells which is in line with previously published
data (Liu et al., 2022). Together with our CyTOF results that showed an accumulation of the
CD8 T_{EM} GZMK precursor state and terminally exhausted CD8$^+$ T cells in CLL LNs (Figure

2E), these findings define cancer-driven exhaustion of CD8$^+$ T cells in the lymphatic tissue of
patients with CLL.

The CD4$^+$ T cell trajectory followed a similar pattern as CD8$^+$ T cells, starting with naïve
genes (*SELL, TCF7* and *IL7R*), followed by genes for activation and effector molecules
(*CD44, CXCR3,* and *GZMM*), and then by exhaustion-related genes (*LAG3, EOMES, LAYN,*
BTLA, and *IRF4*) (Figure 3H, I; Suppl. Figure 5J), indicating a cancer-driven exhaustion also
of CD4$^+$ T cells. The T$_{FH}$ and T$_{REG}$ cell clusters were diverting into two branches of the
trajectory already at an early stage (Figure 3H).

Effector memory CD8 T cells resembling precursor exhausted cells are clonally
expanded in CLL LNs

The reconstruction of the T cell receptor (TCR) sequences at the single-cell level allowed
us to examine the clonal diversity of T cells within the identified subsets. T cells that shared
the same TCR sequence with at least one other cell were defined as clonal. This data
identified a major clonal expansion of CD8$^+$ T cells (ranging from 18% to 69%), but not CD4$^+$
T cells (4% to 26%) in the 5 CLL LN samples, and no correlation between the expansion rate
of CD4$^+$ and CD8$^+$ T cells (Figure 4A). A comparison of the TCR sequences of the major
clones with VDJdb verified that non of these TCRs were linked to viral antigens (Suppl. Table
4). The clonal composition within the CD8$^+$ T cell compartment was highly variable among
patients, with most samples showing mostly small and medium sized clones, and one
sample (BC1) presenting one strongly dominating, hyperexpanded clone. Notably, CD8$^+$ T
cell clonal expansion was enriched in the two CD8 T$_{EM}$ subsets and in CD8 T$_{EX}$ cells (Figure
4B, C) which explains the relative enrichment of the CD8 T$_{EM}$ LYAR cluster in the BC1
sample (Figure 3C) and suggests that these cells are tumor-reactive. Most TCR clones were
confined to a specific T cell cluster, but some were shared between clusters with similar
functional phenotypes, such as CD8 T$_{EM}$ CCL5 and CD8 T$_{EM}$ LYAR (Figure 4D, Suppl. Figure
6A). This suggests that either a dividing activated T cell produces cells with distinct
transcriptional states, or that cells dynamically transition between two transcriptional states.

CLL cells and T cells establish a unique interactome including Galectin-9 and TIM3 as
ligand-receptor pair

Next, we predicted receptor-ligand interactions between T cells and CLL cells in the LNs
by using the public repository of ligand-receptor interactions from OmniPath (Türei et al.,
2016) to perform differential interaction analysis using CellChat (Jin et al., 2021). We
included published scRNA-seq data from 5 rLNs (Aoki et al., 2020)(Suppl. Figure 7A, B) in
this analysis to predict CLL-specific interactions. T cell subsets in both datasets were
grouped into major functional types (CD4 T$_N$, CD8 T$_N$, T$_{FH}$, CD4 T$_{CM}$, T$_{REG}$, CD4 T$_{EM}$ and CD8

T_{EM}) to facilitate comparison. Predicted receptor-ligand interactions were prioritized by the
probability of differential interaction across CLL LNs and rLNs. Overall, the number of
inferred interactions was higher in the CLL LNs compared to the rLNs (1828 vs. 1622; Suppl.
Figure 7C). The highest number of predicted interactions were found for CLL cells and CD8
T_{EM} cells, which act as senders and receivers of signals, both in an autocrine and paracrine
manner (Figure 5A, B). These two cell types were predicted to strongly interact with each
other and also with all other cell subsets. Inferred CLL-specific interactions included ligand-
receptor pairs with known relevance in CLL, such as CD74-MIF (Reinart et al., 2013), and
the chemokines CCL4 and CCL5 and their receptors (Burger et al., 2009; Kiaii et al., 2013)
(Figure 5C). In addition, collagen and adhesion molecules and their binding partners
(COL9A2/3 and ICAM-ITGAL) were predicted as novel interactions. Numerous inferred
interactions between T cells and CLL cells were related to T cell inhibitory signals, such as
HLA-LAG3 (Wierz et al., 2018), BTLA-CD247 (Karabon et al., 2020) and LGALS9-related
circuits. Galectin-9 (encoded by *LGALS9*) is a known ligand of the inhibitory receptor TIM3
and its binding to TIM3 induces T cell death and thus contributes to tumor immune escape
(Yang et al., 2008; Zhu et al., 2005). In CLL, Galectin-9 levels are increased in serum of
patients (Pang et al., 2021b; Wdowiak et al., 2019; Xierenguli et al., 2020). Our scRNA-seq
data identified CLL cells as a source of Galectin-9, and dendritic cells showing an increased
LGALS9 expression in CLL compared to rLNs (Figure 5D). The Galectin-9 binding partner
TIM3 (encoded by *HAVCR2*) was detected in CD8 T_{EX} cells in CLL LNs, but was absent in
rLNs (Figure 5D). Multiplex immunofluorescence staining of CLL LN sections further
confirmed the presence of TIM3 expressing CD4[+] and CD8[+] T cells in the tissue (Figure 5E).
To estimate the relevance of Galectin-9 in CLL development, we analyzed a previously
published proteome data set of CLL cells for Galectin-9 expression (Herbst et al., 2022).
Dividing patients into two groups based on the median level of Galectin-9 expression clearly
demonstrated a significantly shorter treatment-free survival in cases with low Galectin-9
expression (Figure 5F). Altogether this suggests Galectin-9 as novel potential target for
immunotherapy in CLL.
311

Galectin-9 blockade reduces CLL development in mice
We next aimed to preclinically test the potential of Galectin-9 blockade in a mouse model
of CLL. As the Eμ-TCL1 mouse line (Bichi et al., 2002) is the most explored and used *in vivo*
model of CLL, we first performed single-cell RNA-seq to characterize the T cell compartment
of mice that developed CLL-like disease after adoptive transfer of malignant B cells from this
mouse line (TCL1 AT). This allowed us to identify 13 clusters of T cells in the spleen of
leukemic mice, including naïve, effector, effector memory, and regulatory T cell subsets
(Suppl. Figure 8A, B, C). When integrating TCR-seq data, it became clear that one of the two

analyzed samples (#107) harbored a hyperexpanded CD4$^+$ T cell clone showing an effector
phenotype and *Gzmk* expression, which was not the case for the second sample (#110) that
contained a large CD8$^+$ effector memory T cell cluster (Suppl. Figure 8D, E). The clonally
expanded CD4$^+$ effector T cells in sample #107 contained a cluster of cells expressing
exhaustion-related genes. This data suggests the occurrence of CD4$^+$ T cell-driven adaptive
immune control in TCL1 AT mice which is in line with observations in other tumor mouse
models and patients with cancer (Oh and Fong, 2021; Oh et al., 2020; Quezada et al., 2010;
Xie et al., 2010).

Flow cytometry analyses showed that the vast proportion of malignant B cells in the
spleen and bone marrow of TCL1 AT mice expressed Galectin-9 which was not the case for
B cells from wild-type (WT) mice (Figure 6A, Suppl. Figure 8F). Also higher frequencies of
Galectin-9-positive CLL cells were detected in lymph nodes and PB of TCL1 AT mice
compared to WT mice (Suppl. Figure 8F). We further observed TIM3-positive T cells in
tumor-bearing TCL1 AT but not WT mice, with the highest frequencies detected in CD8 T_{EF}
and T_{REG} cells (Figure 6B).

To test the potential of Galectin-9 as novel therapy target for CLL, we treated CLL-bearing
TCL1 AT mice with Galectin-9-blocking antibodies for 20 days (Figure 6C). By a weekly
assessment of CLL cell counts in the blood of the mice, a deceleration of leukemia
development was observed by anti-Galectin-9 compared to isotype control treatment (Figure
6D). At the endpoint of the experiment, 7 weeks after TCL1 AT, a reduced spleen weight and
lower numbers of CLL cells were detected in the spleen and lymph nodes of the mice
receiving anti-Galectin-9 (Figure 6E, F). We further analyzed T cells in the spleen and lymph
nodes at the endpoint and did not observe major differences in the numbers of CD8$^+$ T cells,
conventional CD4$^+$ T cells, or T_{REG} (Suppl. Figure 9A, B). We detected however a lower
frequency of CD39$^+$ and a higher frequency of CD107A$^+$ CD8$^+$ T cells in the treated mice
(Figure 6G, H), suggesting a higher functionality of CD8$^+$ T cells upon anti-Galectin-9
treatment. The most drastic differences were observed in the frequencies of TIM3$^+$ T cells,
which were drastically decreased for CD8$^+$ T cells, conventional CD4$^+$ T cells and PD1$^+$ T_{REG}
cells in the spleen and to a lower degree also in LNs (Figure 6I-K, Suppl. Figure 9C) which is
in line with the proposed role of Galectin-9 in preventing apoptosis of TIM3$^+$ T cells (Zhu et
al., 2005). Based on these data, we suggest that anti-Galectin-9 treatment results in a better
immune control of CLL via suspending the immune suppressive activity of TIM3 on CLL-
associated T cells.

Discussion

T cell exhaustion is considered as one of the main reasons for failure of immunotherapies.
However, its contribution to immunotherapy resistance of patients with B cell malignancies to
such treatments has not been explored so far. Our study provides, for the first time, a single-
cell-resolved analysis of the T cell landscape in blood and tissue samples of CLL, the most
common leukemia in the Western world. CLL is a disease that arises in the lymph nodes, but
the malignant B cells accumulate also in the peripheral blood. Most previous studies
characterized disease-associated alterations in the T cell compartment in blood samples.
Nevertheless, our data clearly show that exhaustion of cytotoxic T cells and accumulation of
regulatory T cells is very prominent in the lymph nodes but not in blood, suggesting the
lymphoid tissue as the site of cancer-directed immune responses in CLL.

In addition to terminally exhausted CD8 T cells, the lymph nodes contain a population of
cells that resembled the previously described precursor cells of exhaustion (Im et al., 2016;
Utzschneider et al., 2016), that share features with memory T cells and have a high *GZMK*
expression (Liu et al., 2022). These cells have self-renewal capacity which likely explains our
observation of a strong accumulation of TCR clonotypes in this population. Our data
identifying such cells in CLL LNs is in line with recent studies demonstrating their presence in
tumor-draining LNs of solid cancer where they are essential for efficient response to ICI
therapy (Connolly et al., 2021; Huang et al., 2022).

CLL cells in lymph nodes are densely packed together with T cells and interactome
analyses using the scRNA-seq data suggested a list of known and novel molecular
interactions between the two cell types. It is well-accepted that CLL cells depend on
microenvironment-derived signals for survival and proliferation, and the identified interactions
including MIF-CD74 (Reinart et al., 2013), CCL4 and CCL5 (Kiaii et al., 2013), and INFG
(Buschle et al., 1993) are part of this support. Further, the interactome study inferred several
inhibitory molecular interactions including BTLA-CD247, CTLA4 and Galectin-9 (encoded by
LGALS9). Galectin-9 is a known ligand of TIM3, an inhibitory receptor that is expressed on
exhausted T cells (Wolf et al., 2020). And even though we detected very low levels of
HAVCR2 (encoding TIM3) by scRNA-seq, we confirmed the presence of TIM3-positive T
cells in CLL LN tissue slides via immunofluorescence staining, and assumed an involvement
of this interaction in immune escape as Galectin-9 binding to TIM3 promotes T_{REG}
development in CLL (Pang et al., 2021a).

TIM3 targeting antibodies enter now clinical trials and show first promising anti-tumor
activity in patients with advanced solid tumors (Curigliano et al., 2021). These antibodies
block the interaction of TIM3 with phosphatidylserine and CEACAM1, but only partially the
binding to Galectin-9 (Sabatos-Peyton et al., 2020; Sabatos-Peyton et al., 2018). Blocking
antibodies targeting Galectin-9 can overcome this limitation and suppress tumor growth in

combination with chemotherapy in a breast cancer mouse model (de Mingo Pulido et al.,

2018). This activity was dependent on CD8$^+$ T cells, which is in line with our observation of

an enhanced CD8$^+$ effector function in anti-Galectin-9 treated TCL1 mice.

Binding of Galectin-9 to TIM3 on T cells induces cell death thereby limiting adaptive

immunity (Zhu et al., 2005). This raises the question of why terminally exhausted T cells

harboring TIM3 expression persist in the tumor microenvironment. Recent data showed that

PD1, via physically interacting with Galectin-9 and TIM3, protects T cells from apoptosis

(Yang et al., 2021). As a consequence, PD1$^+$ TIM3$^+$ T cells accumulate in tumors even in the

presence of enhanced levels of Galectin-9. Blockade of Galectin-9 *in vitro* and in mouse

models results in enhanced T cell survival and improved anti-tumor immunity, especially in

combination with co-stimulatory anti-GITR treatment (Yang et al., 2021). In accordance, anti-

Galectin-9 treatment of TCL1 mice diminished PD1$^+$ TIM3$^+$ T cells along with a reduction in

tumor development.

Besides improving T cell responses, antibody-mediated targeting of Galectin-9 on B-ALL

cells induces DNA damage, alters cell cycle progression, and promotes apoptosis *in vitro*

and significantly extends the survival of mice with aggressive B-ALL (Lee et al., 2022).

Accordingly, we cannot exclude that a similar direct activity on CLL cells contributes to the

delayed leukemia development in TCL1 mice. Further, ligand activation of Galectin-9 in the

tumor microenvironment of pancreatic ductal adenocarcinoma results in tolerogenic

macrophage programming and adaptive immune suppression (Daley et al., 2017). As CLL is

dominated by immunosuppressive myeloid cells (Hanna et al., 2019), Galectin-9 blockade

potentially ameliorates this milieu which likely contributes to a better T cell function.

As our study focused on an in-depth characterization of the T cell landscape in CLL, we

excluded myeloid cells from the analysis, which represents a limitation of our analysis.

Immunosuppressive interactions of CLL-associated myeloid cells and T cells contribute to

immune escape in CLL (Hanna et al., 2016; Jitschin et al., 2014), but their analysis requires

fresh tissue samples to prevent a biased loss of myeloid cell subsets by freezing and

thawing, and such samples are rarely available. As suppression of anti-tumor immunity is

complex and multifactorial, obtaining a complete picture of all cell types and their interactions

with the cancer cells will be necessary for the development of combinatorial treatment

targeting multiple immune cell types and checkpoints to avoid immune escape and

resistance to therapy. The data provided here suggest that Galectin-9 is an attractive novel

therapy target for immunotherapy in CLL and likely beyond in other cancers.

Figure 1: Profiling of the T cell landscape in CLL tissues at the single-cell resolution

A) Graphical overview of the study design. Mass cytometry analyses were performed on 13 rLNs, 7 CLL PB, 3 CLL BM and 22 CLL LNs, 15-color multiplex immunostaining was applied in an additional data set of 42 CLL LNs, and 5 CLL LNs were analyzed using scRNA-seq and TCR-seq. **B)** UMAP plot of 5.2×10^6 CD3$^+$ T cells from 13 rLNs, 7 CLL PB, 3 CLL BM and 22 CLL LNs analyzed by mass cytometry identifying 30 clusters, including 15 for CD4$^+$ T cells, 9 for CD8$^+$ T cells, 2 clusters containing both CD4$^+$ and CD8$^+$ T cells, and 4 DN T cells. **C)** Projection of a selection of protein markers identifying T cell states. Cells are colored based on the normalized protein expression. **D)** Dot plot of the expression of marker genes in the 30 cell clusters. **E)** Heatmap showing the Pearson correlation coefficient and its associated p-value of cell subset proportions from the 20 CLL LNs, corresponding to the worst performing subset (for which the p-value was the highest) of all leave-one-out patient sample sets. **F)** Representative CLL LN-derived tissue core stained with 15-color Orion multiplex with cells colored according to subset identification. CLL cells are displayed as B$_{PR}$: Ki-67$^+$ proliferating B cells, and Other: mainly non-proliferating B cells. **G)** Pearson correlations of PD1$^+$ CD8$^+$ T cells and T$_{REG}$ frequencies determined by multiplex staining of 42 CLL LN cores. Each dot represents an individual patient.

Figure 2: The T cell composition of CLL LNs is distinct and enriched in regulatory and exhausted subsets

A) Principal component analysis of all samples analyzed by mass cytometry based on cell subset frequencies. **B)** UMAP plots of T cells from rLNs, CLL PB, CLL BM and CLL LNs overlaid with a contour plot indicating the cell density. **C)** Boxplot showing cell subset abundances in LNs (n = 20), PB (n = 7) and BM (n = 3) of CLL patients. Statistical significance between CLL LNs and CLL PBs was assessed by limma on normalized cell counts. **D)** Boxplot showing cell subset abundances in CLL LNs (n = 20) and rLN (n = 13). Each symbol represents an individual patient sample and statistical significance was tested by limma on normalized cell counts, $*p<0.05$, $**<0.01$, $***<0.001$.

Figure 3: Single-cell RNA-seq defines T cell states in CLL LNs

A) UMAP plot of 8,576 cells from 5 CLL LNs analyzed by scRNA-seq identifying 13 clusters, including 6 for CD4$^+$ T cells, 4 CD8$^+$ T cells, proliferating CD4$^+$ and CD8$^+$ T cells, DCs and spiked in CLL cells. **B)** Dot plot of the expression of marker genes in the 13 cell clusters. **C)** Cell subset abundance in each sample, with colors and numbers corresponding to the identified cell clusters. **D-E)** UMAP plot colored by average expression levels of the

exhaustion gene signature adapted from Zheng L, *et al* Science, 2021 (D), and the precursor

exhaustion gene signature from Guo X, *et al* Nat Med, 2018 (E). **F)** Pseudotime trajectory

across the 4 CD8$^+$ T cell subsets. **G)** Heatmap showing genes with significant expression

changes along the trajectory from CD8 T_N to CD8 T_{EM} LYAR (left), and from T_N to CD8 T_{EX}

(right). Color represents z-scores. **H)** Pseudotime trajectory across the 6 CD4$^+$ T cell

subsets. **I)** Heatmap showing genes with significant expression changes along the trajectory

from CD4 T_N to CD4 T_{FH}. Color represents z-scores.

Figure 4: T cell clonal expansion in CLL is mainly restricted to CD8 T_{EM} cells

A) Bar plot indicating the percentage (rounded values are indicated) of single, small,

medium, large and hyperexpanded sized clones in CD8$^+$ (left) and CD4$^+$ (right) T cells for

each patient analyzed. **B)** UMAP plot colored according to the T cell clone size based on the

TCR-seq data. **C)** Graph showing the TCR Shannon diversity index for each T cell subset

identified by scRNA-seq. **D)** Chord diagram showing clone sharing from CD8 T_{EM} CCL5 and

from CD8 T_{EM} LYAR. Only cells carrying expanded TCR sequence are presented in the

circle. Cells sharing same TCR sequences across clusters are summed and connected by

bands.

Figure 5: Interactome analyses of CLL LNs identify Galectin-9 as a potential therapeutic

target

A) Circle plot depicting the differential number of interactions between cell subsets in CLL

LNs compared to rLNs. Thickness of bands represent the number of differential interactions

between the two data sets, and increased interactions are depicted in red, decreased

interactions in blue. **B)** Heatmap showing the differential number of interactions of CLL LNs

versus rLNs. Rows and columns represent source and target subsets, respectively. Bar plots

represent the total outgoing (right) and incoming (top) interactions scores respectively. **C)**

Heatmap plot depicting a curated list of ligand-receptor pairs differentially upregulated in CLL

LNs compared to rLNs as identified via CellChat (Jin et al., 2021). The dot color and size

represent the calculated communication probability and p-values of differential

communication, respectively. **D)** Violin plots showing the expression distribution of *LGALS9*

(top two plots) and *HAVCR2* (bottom plots) genes in the CLL LN data set and the rLN data

set from Aoki *et* al. (Aoki et al., 2020). **E)** Representative microscopy images showing CD3,

CD8, TIM3 and DAPI in CLL LN tissue sections. **F)** Time-to-treatment in CLL patients with

high or low Galectin-9 protein levels.

Figure 6: Blocking of Galectin-9 controls tumor growth in the TCL1 mouse model

A) Representative contour plot and percentage of Galectin-9$^+$ B cells and CLL cells from spleen of WT and TCL1 AT mice, respectively. **B)** Percentage of TIM3$^+$ cells out of CD8 T_N, CD8 T_M, CD8 T_{EF}, CD4 T_{CONV} and T_{REG} T cells in spleen of WT and TCL1 AT mice. **C)** Schematic diagram of Galectin-9-blocking antibody treatment in TCL1 AT mice. Analyses of T, myeloid and CLL cells were performed 7 weeks after treatment start. **D)** Absolute number of CD19$^+$ CD5$^+$ CLL cells in blood of isotype antibody- and anti-Galectin-9-treated mice. **E)** Spleen weight, and **F)** absolute number of CD19$^+$ CD5$^+$ CLL cells in spleen and lymph nodes of isotype antibody- and anti-Galectin-9-treated mice. **G)** Representative contour plot and percentage of CD39$^+$ cells out of CD8$^+$ T cells from spleen of isotype antibody- and anti-Galectin-9-treated mice. **H)** Representative contour plot and percentage of CD107A$^+$ out of CD8$^+$ T cells from spleen of isotype antibody- and anti-Galectin-9-treated mice. **I-K)** Representative contour plot and percentage of PD1$^+$ TIM3$^+$ cells out of CD8$^+$ T cells (I), CD4$^+$ T_{CONV} (J), and T_{REG} (K) cells from spleen of isotype antibody- and anti-Galectin-9-treated mice. Each symbol represents an individual mouse, statistical significance was tested by unpaired t-test with Welch approximation, *p<0.05, **<0.01, ***<0.001. Bars indicate mean ± SEM.

Supplementary Figure Legends

Supplementary Figure 1 (related to Figure 1):

A) Gating strategy used to select live, CD45$^+$ CD3$^+$ T cells: gates were placed on cells

(Beads vs. Ir191), singlets (Ir193 vs. Ir191) and live cells (Pt195$^-$), which were gated and

exported as .fcs files. CD3$^+$ cells were further analyzed. **B)** Number of T cells per analyzed

sample. **C)** UMAP plot depicting sample distribution.

Supplementary Figure 2 (related to Figure 1):

A) Projection of all protein markers analyzed by mass cytometry. Cells are colored based on

the normalized protein expression. **B)** Heatmap showing median protein expression of the 33

markers considered (columns) for each cluster (rows) identified. Both markers and clusters

were grouped based on hierarchical clustering.

Supplementary Figure 3 (related to Figure 1):

A) Cell subset frequency for each cluster. Each dot represents one sample. Clusters are

ordered from highest to lowest median frequency. **B)** Representative images of 15-color

multiplex staining of tissue cores of CLL LNs. **C)** Frequency of cell types identified using the

15-color multiplex staining in each core (n = 42). **D)** Heatmap showing the Pearson

correlation coefficient and its associated p-value of the multiplex staining cell subset

proportions from the 42 CLL LN cores, corresponding to the worst performing subset of all

leave-one-out patient samples. The correlation for each cell subset pair was then defined as

the result among subsets for which the p-value was highest. Significant results are depicted

with black borders.

Supplementary Figure 4 (related to Figure 2):

A) Hierarchical clustering (with Pearson correlation and complete linkage) of samples based

on the T cell subset abundances in each sample. Cell subsets are color-coded as defined in

Fig. 1B. Row annotations indicate sample condition (BM, bone marrow, LN, lymph node, and

PB, peripheral blood), age (years) and tumor load (TL). **B)** Heatmap showing the Pearson

correlation coefficient and its associated p-value of cell subset proportions from the 7-paired

CLL LN and PB samples, corresponding to the worst performing subset of all leave-one-out

patient samples. The correlation for each cell subset pair was then defined as the result

among subsets for which the p-value was highest. **C-E)** Pearson correlation plot between

age and frequency of CD8 T_N cells (C), $T_{REG\ CM2}$ cells (D), and T_{REG} PD1 cells (E). Each

symbol represents an individual patient sample.

Supplementary Figure 5 (related to Figure 3):

A) Gating strategy used to isolate live, CD45$^+$ CD3$^+$ T cells and CLL cells: gates were placed on cells (SSC-A vs FSC-A), singlets (FSC-H vs FSC-A) and live cells (Negative for viability dye). CD45$^+$ cells were further gated for CLL cells (CD19$^+$ CD5$^+$) or T cells (CD19$^-$ CD3$^+$). **B)** UMAP plot depicting sample distribution. **C)** UMAP plot of T$_{FH}$ cells. **D)** Heatmap of differentially expressed genes of the 4 clusters of T$_{FH}$ cells. **E)** UMAP plot colored by average expression of the precursor exhaustion gene signature from Andreatta *et al* (Andreatta et al., 2021), and **F)** according to PBMC3K data set (Hao et al., 2021) applied to the data set of 5 CLL LNs. **G)** Cell subset distribution of the major T cell subsets in the 4 CLL samples analyzed with scRNA-seq and mass cytometry. **H)** Pearson correlation of major T cell subset frequencies determined with scRNA-seq and the mass cytometry data for the 4 CLL patient samples analyzed with both techniques. **I-J)** CD8$^+$ (I) and CD4$^+$ (J) T cell subsets ordered by diffusion pseudotime.

Supplementary Figure 6 (related to Figure 4):

A) Chord diagram showing the expanded TCR clone sharing between the T cell subsets identified. Cells carrying identical TCR sequences are connected by bands.

Supplementary Figure 7 (related to Figure 5):

A) UMAP plot of 20,673 cells from 5 rLNs analyzed by scRNA-seq from Aoki *et* al. (Aoki et al., 2020) identifying 21 clusters, including 9 T cell clusters, 8 B cell clusters, proliferating cells, NK cells and DCs. **B)** UMAP plot where PBMC3K data set (Hao et al., 2021) was mapped to the 5 rLNs data set from Aoki *et* al. (Aoki et al., 2020). **C)** Number of inferred interactions between T cells and B cells in rLN samples, or T cells and CLL cells in CLL LN samples.

Supplementary Figure 8 (related to Figure 6):

A) UMAP plot of 6,201 cells from 2 TCL1 AT spleens analyzed by scRNA-seq identifying 13 clusters, including 11 T cells clusters, CLL cells and myeloid cells. **B)** UMAP plot depicting sample distribution. **C)** Dot plot of the expression of marker genes in the 13 cell clusters**. D)** UMAP plot colored according to the T cell clone size based on the TCR-seq data**. E)** Bar plot indicating the percentage (rounded values are indicated) of single, small, medium, large and hyperexpanded sized clones in CD8$^+$ (left) and CD4$^+$ (right) T cells for the two mouse samples analyzed**. F)** Percentage of Galectin-9$^+$ B cells and CLL cells from bone marrow, lymph nodes and peripheral blood of WT and TCL1 AT mice. Each symbol represents an individual mouse, statistical significance was tested by unpaired t-test with Welch approximation, *p<0.05, **<0.01, ***<0.001.

Supplementary Figure 9 (related to Figure 6):

A-B) Absolute number of CD4$^+$ T$_{CONV}$, T$_{REG}$ and CD8$^+$ T cells in spleen (A) and lymph nodes (B) of isotype antibody- and anti-Galectin-9-treated mice. **C)** Percentage of PD1$^+$ TIM3$^+$ cells out of CD8$^+$, CD4$^+$ T$_{CONV}$ and T$_{REG}$ cells in lymph nodes of isotype antibody- and anti-Galectin-9-treated mice. Each symbol represents an individual mouse, and statistical significance was tested by unpaired t-test with Welch approximation, *p<0.05, **<0.01. Bars indicate mean ± SEM. Ns = not significant.

Methods

Patient samples

Lymph node (LN, n = 21), peripheral blood (PB, n = 7), and bone marrow (BM, n = 3)

samples from CLL patients, and reactive lymph node (rLN, n = 13) samples from heathy

controls were obtained after informed consent and according to the guidelines of the Hospital

Clínic Ethics Committee, the Ethics Committee of the University of Heidelberg, and the

Declaration of Helsinki. Patients with CLL were diagnosed following the World Health

Organization (WHO) classification criteria. All clinical information of the patients analyzed in

this work is provided in Suppl. Table 1.

Mice and tumor models

C57BL/6 wild-type (WT) mice were purchased from Charles River Laboratories (Germany)

or Janvier Labs (France). Adoptive transfer of TCL1 leukemia cells was performed as

previously described (Hanna et al., 2016; Wierz et al., 2018). In short, malignant B cells were

enriched from splenocytes of Eμ-TCL1 mice using EasySep Mouse Pan-B Cell Isolation Kit

(StemCell Technologies, Inc., Cologne, Germany). The CD5$^+$ CD19$^+$ content of purified cells

was typically above 95%, as measured by flow cytometry. 1 x 10^7 malignant TCL1

splenocytes were transplanted by intravenous (i.v) injection into 8-12 weeks-old C57BL/6 WT

female mice. Leukemia progression was monitored weekly by assessment of the percentage

of CD5$^+$CD19$^+$ cells in the PB.

Antibody treatments

For Galectin-9 blockade experiments, mice were first transplanted with TCL1 leukemia

cells. After 2 weeks, mice were assigned to different treatment arms according to the

percentage of CD5$^+$CD19$^+$ (CLL) cells out of CD45$^+$ cells in PB. Subsequently, mice were

injected i.p. with 6 mg/kg of anti-Galectin-9 (clone: RG9-1) or rat IgG2b isotype control

antibody (clone: LTF-2), 3 times per week for another 7 weeks. All antibodies for in vivo

experiments were acquired from BioXcell (West Lebanon, NH).

Tissue sample collection and preparation of cell suspensions

Mice were euthanized by increasing concentrations of carbon dioxide (CO_2) or by cervical

dislocation before reaching the established humane end-point. PB was drawn from the

submandibular vein or via cardiac puncture and collected in ethylenediaminetetraacetic acid

(EDTA)-coated tubes (Sarstedt, Numbrecht, Germany). Single-cell suspensions from

spleens, bone marrow (BM) and inguinal LNs were prepared as previously described (Hanna

et al., 2016; McClanahan et al., 2015). BM cells were flushed from femurs with 5 mL of

phosphate-buffered saline (PBS)/5% fetal calf serum (FCS). Spleen single-cell suspensions

were generated by using the gentleMACS tissue dissociator with Gentle MACS tubes C
(Miltenyi Biotec). Single cell suspensions from LNs were prepared by grinding the tissue
through 70 mm cell strainers (BD Biosciences, Heidelberg, Germany). Erythrocytes were
lysed by using Red blood cell lysis buffer (Mitenyi Biotec).
641
Flow cytometry
Single-cell suspensions were immunostained with antibodies against cell surface proteins
(Key Resource Table) in FACS buffer containing 0.1 % fixable viability dye (Thermo Fisher
Scientific, 65- 0866-14) for 30 min at 4 °C. Cells were subsequently washed twice in FACS
buffer, fixed with IC fixation buffer (Thermo Fisher Scientific, 00-8222-49) for 30 min at room
temperature, washed twice with FACS buffer, and stored at 4 °C in dark conditions until
being analyzed.
For transcription factor or intracellular cytokine staining, cell surface-stained cells were
fixed for 30 min at room temperature with Foxp3 fixation/permeabilization buffer (Thermo
Fisher Scientific, 00-5523-00 or Miltenyi Biotec, 130-093-142). After a washing step with
FACS buffer, cells were permeabilized for 30 min at room temperature with 1X
permeabilization buffer (Thermo Fisher Scientific, 00- 8333-56). Intracellular staining with
antibodies against transcription factors or cytokines was performed in 1X permeabilization
buffer for 30 min at room temperature. Excess antibodies were washed twice with 1X
permeabilization buffer and cells were resuspended in 1X permeabilization buffer and stored
at 4 °C in dark conditions until they were analyzed by flow cytometry.
Cell fluorescence was assessed using a BD LSRFortessa[TM] (BD Biosciences) or a
Novocyte Quanteon (Agilent) flow cytometer, and data was analyzed using FlowJo X 10.0.7
software (FlowJo). In each experiment, single fluorochrome stainings were used to
compensate for spectral overlap. Fluorescence minus one (FMO) controls were employed for
proper gating of positive cell populations. FMO-normalized Mean Fluorescence Intensity
(nMFI) for unimodal distributions or percentage of positive cells for bimodal distributions were
determined for populations of interest. The standard gating strategy for analysis of CLL cells
and T cells is depicted in Suppl. Figure 4A.
666
Immunofluorescence of whole lymph node sections
Paraffin-embedded tissue sections of lymph nodes were obtained from the National
Center for Tumor Disease (NCT), Hospital Clínic de Barcelona, and the University of
Würzburg.
After validation of primary and secondary HRP-conjugated antibodies using DAB
chromogenic detection, incubation times and appropriate antibody concentrations were
optimized using the Opal™ 5-color kit (Akoya Biosciences, NEL840001KT). Epitope stability

was assessed in order to determine the order of each antibody in sequence, and antigen
stripping efficiency was confirmed in all used primary antibodies.

Deparaffinization and rehydration of lymph node sections was performed by immersion in
xylene for 10 minutes twice, followed by immersion in a series of descending ethanol
concentrations prior to distilled water. Slides were then cooked with antigen retrieval buffer
pH9 (Akoya Biosciences, AR900250ML) at 100 °C for 25 minutes in a steam cooker. Slides
were allowed to cool at room temperature and subsequently washed in tris-buffered saline
containing 0.1 % tween 20 (TBS-T). Next, slides were incubated with blocking buffer (Akoya
Biosciences, NEL840001KT) for 10 minutes at room temperature, and incubated with the
primary antibody for the indicated time and concentration (Key Resource Table). After three
washing steps with TBS-T, slides were incubated with anti-rabbit HRP-conjugated secondary
antibody (Akoya Biosciences, NEL840001KT). Sections underwent rounds of antigen
stripping via incubation at 100 °C in AR9 buffer (Akoya Biosciences, AR900250ML) for 25
minutes, allowing removal of the excess of antibody complex formed. For multi-color staining,
tissue sections went through multiple rounds of blocking, primary and secondary antibody
labelling, each round with a different Opal^{TM} dye (Akoya Biosciences, NEL840001KT).
Finally, background signal reduction through incubation for 10 minutes at RT in 0.1 % Sudan
Black (Sigma-Aldrich, 199664) took place and sections were washed thoroughly in TBS-T.
Finally, nuclei were stained with DAPI (Akoya Biosciences, NEL840001KT) and slides were
mounted using ProLong Diamond Antifade Mountant (Life Technologies, P36965).
Slides were imaged using a Carl Zeiss Axioscan 7 slide scanner at 20X.
695

Immunofluorescence staining and imaging of CLL TMA
TMA 159/2 was de-paraffinized and subjected to antigen retrieval for 5 minutes at 95°C
followed by 5 minutes at 107°C, using pH8.5 EZ-AR 2 Elegance buffer (BioGenex). To
reduce tissue autofluorescence, slides were placed in a transparent reservoir containing
4.5% H_2O_2 and 24 mM NaOH in PBS and illuminated with white light for 60 minutes followed
by 365 nm light for 30 minutes at room temperature as previously described (Lin et al.,
2018). Slides were rinsed with surfactant wash buffer (0.025% Triton X-100 in PBS), placed
in a humidified stain tray, and incubated in Image-iT™ FX Signal Enhancer (Thermo Fisher)
for 15 minutes at room temperature. After rinsing with surfactant wash buffer, the slides were
placed in a humidity tray and stained with the panel of fluor- and hapten-labeled primary
antibodies in PBS-Antibody Stabilizer (CANDOR Bioscience GmbH) containing 5% mouse
serum and 5% rabbit serum for 2 hours at room temperature. Slides were then rinsed again
with surfactant wash buffer and placed in a humidified stain tray and incubated with SYTOX
Blue (Thermo Fisher), ArgoFluor™ 845 mouse-anti-DIG in PBS-Antibody Stabilizer
containing 10% goat serum for 30 minutes at room temperature. The slides were then rinsed

a final time with surfactant wash buffer and PBS, coverslipped with ArgoFluor™ Mounting

Media (RareCyte, Inc.) and dried overnight (Lin et al., 2022).

Slides were imaged using an Orion instrument (RareCyte, Inc.) at 20X. Raw image files

were processed to correct for system aberrations; then signals from individual targets were

isolated to separate channels using the Spectral Matrix obtained with control samples,

followed by stitching of FOVs to generate a continuous open microscopy environment (OME)

pyramid TIFF image.

Image analysis

Preprocessing

TMA 159/2 with 18-color staining was analyzed using the multiple-choice microscopy

pipeline MCMICRO (Schapiro et al., 2022) (Commit

91fcd5eefd69da112414b06cf3a65a9a66afeccf). Dearraying was performed by Coreograph

(Schapiro et al., 2022) 2.2.8, Unmicst (Yapp et al., 2022) 2.7.0 segmented the cells based on

the nuclear channel Sytox, S3segmenter (Saka et al., 2019) 1.3.12 generated single-cell

masks and Mcquant (Schapiro et al., 2022) 1.5.1 quantified the signal of each of the 18

channels in each cell. The corresponding parameters (params.yml) for full reproducibility of

this MCMICRO runs are available on Github

(https://github.com/SchapiroLabor/ImageAnalysisTMA159_2).

Following visual inspection of Coreograph's QC file, all cores that were damaged were

filtered out. Additionally, the 4 muscle tissue cores – which were used as control – were

filtered. In total we performed the downstream analysis with 42 cores. Channel numbers 2

(Sytox), 14 (E-cadherin), 17 (Pan-CK) and 18 (AF-Tissue) were removed for downstream

analysis since (i) Sytox and AF-Tissue represent nuclei and autofluorescence which are not

relevant for cell type calling; and (ii) E-cadherin and Pan-CK expression are not expected in

LN tissue and therefore, failed our visual QC due to unspecific staining patterns.

Batch correction

Cell type clustering on the raw data results in clusters consisting mainly of cells of a single

core of origin. This is due to preanalytical variability, which requires batch correction on core-

level. Therefore, marker intensities were normalized with Mxnorm (Harris et al., 2022)

0.0.0.9000 (transform: log10_mean_divide, method: none) before using Rphenograph

(Levine et al., 2015) 0.99.1 to cluster the individual cell types.

Cell type assignment

Cell types were assigned based on marker thresholding (positive / negative) and
comparison to an expert curated list of markers associated with the specific cell types (Suppl.
Table 2).

Statistical analysis

The subsequently calculated cell type frequencies in each core were used to perform two-
sided t-tests (base package stats) for changes in cell type frequencies between patients that
were positive or negative for a variety of binary clinical parameters (sex, IGHV, del13q,
del17p, died, treated, del17p_tp53 (del17p and/or tp53 positive)). Additionally, Cox
regression (survivalAnalysis 0.3.0) was calculated on overall survival and time to next
treatment. Finally, correlation with the Pearson method was checked (cor.test() in base
package stats) between overall survival, time to next treatment and all cell type frequencies.
Up to the statistical analysis, calculations were performed under R version 4.1.0 (2021-05-
18) on platform x86_64-apple-darwin17.0 (64-bit) running under macOS 13.0. Statistical
analysis was done on R version 4.0.5 (2021-03-31) on platform: x86_64-apple-darwin17.0
(64-bit) running under: macOS Big Sur 10.16.
All mentioned steps of image processing were performed on the BWforCluster.

Mass cytometry

A panel of 42 heavy metal-labeled antibodies for detection of both surface and
intracellular proteins was adopted from Bengsch et al. and modified based on literature
research in order to characterize diverse T-cell phenotypes (Bengsch et al., 2018). The
complete list of proteins detected and the heavy metal-conjugated antibodies used are listed
in the Key Resource Table. For most of the markers, heavy metal-conjugated antibodies
were commercially available and purchased from Fluidigm. Where no heavy metal-
conjugated antibodies were commercially available, coupling of heavy metal to respective
antibodies was performed in-house using the Maxpar® X8 Multimetal Labeling kit (Fluidigm)
following the manufacturer's instructions. ^{113}In, ^{115}In and ^{139}La heavy metal isotopes (#) were
not available from Fluidigm and were purchased from Trace Sciences International (^{113}In)
and Sigma (^{115}In, #203440-1G; ^{139}La, #211605-100G) Following the preparation of a 1M
solution, ^{113}In, ^{115}Ln and ^{139}La heavy-metal isotopes (#) were conjugated to monoclonal
purified IgG antibodies using the Maxpar® X8 Multimetal Labeling kit (Fluidigm, #201300).

Frozen cell suspensions were thawed and washed prior to incubation in 15 mL of pre-
heated RPMI 1640 containing 10 % FBS in a roller incubator for 30 min at room temperature.
Cells were filtered to remove dead cells and B-cell depletion was performed using human
CD19 microbead (Miltenyi Biotec) labeling according to the manufacturer's protocol.

Unlabeled CD19$^-$ cells were collected, washed, filtered, and counted prior to mass cytometry

processing.

CD19-depleted single-cell suspensions were stained as previously described (Wierz et al.,

2018). Briefly, cells were stained with 5 µM cisplatin (Cell-ID Cisplatin, Fluidigm) for 5 min in

order to label dead cells. Cells were then washed once and cell surface staining was

performed for 30 min at room temperature. After a washing step, cells were fixed with

Fixation/Permeabilization buffer (Thermo Fisher Scientific) following the manufacturer's

instructions. Intracellular staining was performed by incubating the antibody- cocktail for 30

min at room temperature. After a washing step, cells were stained with cell-ID Intercalator-Ir

(Fluidigm) in fixation and permeabilization solution, followed by another two washing steps

with PBS and ddH$_2$O, respectively. Prior to acquisition, cells were resuspended at a

concentration of 5 x 10^5 cells/mL in ddH$_2$O with 1:10 calibration beads (EQ Four Element

Calibration Beads, Fluidigm). Samples were analyzed at a flow rate of 0.03 mL/min with the

Helios mass cytometer (Fluidigm) of the National Cytometry Platform (LIH). Initial data

processing and quality control were performed. Flow cytometry standard (FCS) files were

normalized with EQ beads using the HELIOS instrument acquisition software (Fluidigm).

As a first analysis step, samples were pre-gated as follows: gates were placed on cells

(Beads vs. Ir191), singlets (Ir193 vs. Ir191) and live cells (Pt195$^-$). Next, non-B immune cells

(CD19$^-$ CD45$^+$) and CD3$^+$ T-cells (CD3$^+$ NCAM$^-$) were selected exported as .fcs files.

Raw data .fcs files were merged into a flowSet object using flowCore package (Hahne et

al., 2009). Signal intensities for each marker were arcsinh-transformed using a co-factor of 5

(default) (Bruggner et al., 2014). Sample quality was assessed based on cell counts per

sample and samples with less than 1,500 cells were excluded. Multi-dimension scaling

(MDS) plotting using median arcsinh-transformed marker expression for all cells in each

sample was used for sample clustering analysis.

Cytometry data analysis tools (CATALYST) R package (Chevrier et al., 2018) containing

FlowSOM (Van Gassen et al., 2015) and ConsensusClusterPlus (Wilkerson and Hayes,

2010) metaclustering methods was used for cell cluster identification using all cells from all

samples. Clustering was performed using arcsinh-transformed expression of 33 markers, i.e.,

excluding the markers DNA content and cisplatin viability, which were used to select viable,

single cells, and CD19 and HLA-DR, which were used to exclude B cells (Key Resource

Table). TIM3, LAG3, GITR, CD47 and 4-1BB were excluded from the analysis due to low

signal and likely adding noise in the cluster generation process.

The maximum number of clusters allowed to be evaluated was set to maxK = 30, after

biological relevance of obtained clusters was assessed, and k < 30 and k > 30 were verified

to be underfitting or overfitting, respectively. Cell clustering was visualized using the UMAP

algorithm, displaying 1 x10^3 random cells from each sample. The robustness of this approach

was evaluated by repeatedly plotting an increasing number of randomly picked cells from

each sample, with a range of 200 to 1,000 cells, and evaluating the similarity of the 30

subpopulations recognized in the respective UMAP plots. Sample #HD3 was excluded as an

outlier.

Cell sorting for single-cell RNA and TCR sequencing

Single-cell suspensions from LNs of CLL patients were retrieved by partially thawing vials

of cryopreserved cells in order to preserve unused cells. Samples were stained for cell

surface proteins for 30 min. After washing with PBS, cells were resuspended in PBS + 5 %

FBS containing 0.2 µg/mL DAPI prior cell sorting. The gating strategy for $CD3^+$ T-cell and

CLL cell sorting is depicted in Suppl. Figure 5A. Cells were sorted in PBS + 2 % FBS using a

BD FACSAria^{TM} II or BD FACSArisa Fusion (both from BD Biosciences) cell sorters. The

purity of cells after sorting was above 95 %.

Single-cell RNA sequencing library construction using the 10x Genomics Chromium platform

Five CLL LN samples were processed for single-cell TCR and 5' gene expression profiling

of CD3 T-cells and CLL cells using the Chromium Next GEM Single Cell V(D)J Solution from

10x Genomics following the manufacturer's instructions. Briefly, for a target cell recovery of

5,000 cells, the concentration of T-cells was adjusted to 1 x10^3 cells/µL and tumor cells were

added at a ratio of 1:20. Cells were then loaded into the Single Cell A Chip (10x Genomics).

GEX libraries were sequenced on a HiSeq 4000 machine (Illumina) or on a NovaSeq 6000

(Illumina), and V(D)J-enriched libraries were sequenced on a NextSeq 550 (Illumina).

Single cell RNA-Seq data processing

Sequencing reads were aligned by Cell Ranger(5.0.1) to reference version hg38(2020-A)

for human data and to reference version mm10(2020-A) for mouse data. Raw count matrices

were processed by the quality control pipeline to determine the optimal cut-offs for filtering.

Cells were filtered by read counts and mitochondrial RNA content. Lower cut-off for read

counts were determined by read-counts distribution for samples in each dataset and were set

at 350 for human and 500 for mouse, mitochondrial RNA content should be lower than 10%.

Raw gene count matrices were processed by Seurat(v4.1.0) (Hao et al., 2021),

expression values were normalized and scaled by SCTransform. Doublets were filtered by

DoubletFinder(2.0.3) (McGinnis et al., 2019). Dimensional reduction was performed by

Principle Component Analysis (PCA). Datasets were integrated by HARMONY (Korsunsky et

al., 2019) to reduce batch bias on calculated principal components. Sample identity was

used as the batch variable for HARMONY integration. UMAP was used for dimensional

reduction on first 23 HARMONY computed cell-embeddings.

Cluster were defined by the Louvain method at the resolution of 0.5, clusters defined at
this step were used for annotation by manual curation. Markers for each cluster were
produced by differential expression analysis searching for overexpressing markers of each
cluster. Top 10 overexpressing markers of each cluster were illustrated by a heatmap.
860

Pseudotime analysis

Diffusion pseudotime analysis was performed using destiny(3.8.1) (Angerer et al., 2016).
Cells expressing CD4 and CD8 were analysed separately. To retrieve the pseudotime of
each marker differentiated cell groups, "DPT" function from destiny was used for computing
the diffusion pseudotime on each cell. For each cell type, the root nodes were placed within
the corresponding naïve cell type cluster.

To performed differential gene expression analysis on pseudotime, the diffusion
pseudotime values of cells in trajectories were analysed together with their transcript
expression values by tradeSeq (Van den Berge et al., 2020). Differentially expressed genes
were plotted against ranked diffusion pseudotime for illustration.
871

TCR data analysis

V(D)J transcripts from single cells were aligned and counted using the Cell Ranger
pipeline(5.0.1). GRCm38 VDJ Reference 5.0.0 from 10X genomics was used for mouse data
and GRCh38 VDJ Reference 5.0.0 from 10X genomics was used for human data. In
addition, an output file containing TCR α- and β-chain CDR3 nucleotide sequences and a cell
barcode for all single cells was generated. Only productive rearrangements were evaluated,
and 3 or more cells containing the same α- and β-chain CDR3 consensus nucleotide
sequences were considered cell clones. scRepertoire (Borcherding et al., 2020) was used to
process contig and clonotype information from Cell Ranger. The summarized TCR
information per cell were merged with Seurat object using the shared cell barcodes between
the TCR library and the RNA-Seq library for integrated analysis. Shannon diversity index was
computed to represent the diversity of the TCR repertoire in each single cell clusters. Cells
with expanded TCRs are categorized by their expansion level and then illustrated by umap
plots. To illustrate expanded TCRs shared between different clusters, cells with expanded
TCRs were represented by two types of chord diagrams. The first type depicts how other
clusters share expanded TCRs with CD8 TEM clusters, where a chord diagram was created
for each CD8 TEM cluster. Circular fragments in these chord diagrams represent cells
carrying an expanded TCR in each cell type, arcs connect cells that share the same TCRs
with the CD8 TEM of interest. Width of the arcs are scaled by the number of cells sharing
same TCRs. The second type of chord diagram present a similar concept, it depicts TCR
sharing across all cell types instead of one.

The VDJdb browser (Bagaev et al., 2020) was used in order to identify CDR3 regions
from the V(D)J single-cell dataset whose antigen-specificity has been reported. Both TCR α-
and β- chain CDR3 amino acid sequences from each sample were used for the literature
analysis aimed at identifying previously described binding epitopes.

Ligand-receptor interaction analysis

To find differential ligand-receptor interactions between datasets, a control dataset derived
from 5 reactive lymph nodes (Aoki et al., 2020) was added for comparison. The reactive
lymph node dataset was independently processed by the same quality control pipeline and
single cell analysis pipeline. CellChat was used for differential cell-cell communication
analysis. Ligand-receptor interactions from OmnipathR database (Türei et al., 2021) were
included as candidates. CellChat analysis was first performed independently on the two
datasets using LIANA (Dimitrov et al., 2022). The results were then contrasted by CellChat
for differential interactions. Labels from the two comparing datasets were harmonized to 8
major cell types (CLL/B, CD4 T_{CM}, T_{REG}, CD4 T_{EM}, CD8 T_{EM}, CD4 T_N, CD8 T_N, T_{FH}) for
comparison. Differential Interactions with p-values below 0.01 were visualized by the dot plot
functions of CellChat.

Sub-cluster analyses and cell-signature annotation

To reveal potential substructure among cell types of interest, we performed a subcluster
analysis on the T_{FH} cluster. UMAP analysis was repeated on the harmony corrected
embeddings of cells from the T_{FH} cluster to reveal substructures. To identify the potential cell
types, we applied a T_{FH} signature (Roider et al., 2020) on each cell for in-silico staining.
Signature scores were computed by the AddModuleScore function of Seurat and then
visualized by a UMAP plot.

Data availability

Human single-cell RNA- and TCR-sequencing data are deposited to European Genome-
phenome Archive (EGA) under the accession ID (Pending) and are available on request.
Mouse single-cell RNA- and TCR-sequencing data are available on Gene Expression
Omnibus under the accession ID (Pending).

References

References

Andreatta, M., J. Corria-Osorio, S. Müller, R. Cubas, G. Coukos, and S.J. Carmona. 2021.
Interpretation of T cell states from single-cell transcriptomics data using reference
atlases. *Nat Commun*. 12:2965.
Angerer, P., L. Haghverdi, M. Büttner, F.J. Theis, C. Marr, and F. Buettner. 2016. destiny:
diffusion maps for large-scale single-cell data in R. *Bioinformatics*. 32:1241-1243.
Aoki, T., L.C. Chong, K. Takata, K. Milne, M. Hav, A. Colombo, E.A. Chavez, M. Nissen, X.
Wang, T. Miyata-Takata, V. Lam, E. Viganò, B.W. Woolcock, A. Telenius, M.Y. Li, S.
Healy, C. Ghesquiere, D. Kos, T. Goodyear, J. Veldman, A.W. Zhang, J. Kim, S.
Saberi, J. Ding, P. Farinha, A.P. Weng, K.J. Savage, D.W. Scott, G. Krystal, B.H.
Nelson, A. Mottok, A. Merchant, S.P. Shah, and C. Steidl. 2020. Single-Cell
Transcriptome Analysis Reveals Disease-Defining T-cell Subsets in the Tumor
Microenvironment of Classic Hodgkin Lymphoma. *Cancer discovery*. 10:406-421.
Bagaev, D.V., R.M.A. Vroomans, J. Samir, U. Stervbo, C. Rius, G. Dolton, A. Greenshields-
Watson, M. Attaf, E.S. Egorov, I.V. Zvyagin, N. Babel, D.K. Cole, A.J. Godkin, A.K.
Sewell, C. Kesmir, D.M. Chudakov, F. Luciani, and M. Shugay. 2020. VDJdb in 2019:
database extension, new analysis infrastructure and a T-cell receptor motif
compendium. *Nucleic Acids Res*. 48:D1057-d1062.
Bengsch, B., T. Ohtani, O. Khan, M. Setty, S. Manne, S. O'Brien, P.F. Gherardini, R.S.
Herati, A.C. Huang, K.M. Chang, E.W. Newell, N. Bovenschen, D. Pe'er, S.M.
Albelda, and E.J. Wherry. 2018. Epigenomic-Guided Mass Cytometry Profiling
Reveals Disease-Specific Features of Exhausted CD8 T Cells. *Immunity*. 48:1029-
1045.e1025.
Bichi, R., S.A. Shinton, E.S. Martin, A. Koval, G.A. Calin, R. Cesari, G. Russo, R.R. Hardy,
and C.M. Croce. 2002. Human chronic lymphocytic leukemia modeled in mouse by
targeted TCL1 expression. *Proc Natl Acad Sci U S A*. 99:6955-6960.
Blank, C.U., W.N. Haining, W. Held, P.G. Hogan, A. Kallies, E. Lugli, R.C. Lynn, M. Philip, A.
Rao, N.P. Restifo, A. Schietinger, T.N. Schumacher, P.L. Schwartzberg, A.H. Sharpe,
D.E. Speiser, E.J. Wherry, B.A. Youngblood, and D. Zehn. 2019. Defining 'T cell
exhaustion'. *Nat Rev Immunol*. 19:665-674.
Borcherding, N., N.L. Bormann, and G. Kraus. 2020. scRepertoire: An R-based toolkit for
single-cell immune receptor analysis. *F1000Res*. 9:47.
Bruggner, R.V., B. Bodenmiller, D.L. Dill, R.J. Tibshirani, and G.P. Nolan. 2014. Automated
identification of stratifying signatures in cellular subpopulations. *Proc Natl Acad Sci U
S A*. 111:E2770-2777.
Burger, J.A., M.P. Quiroga, E. Hartmann, A. Bürkle, W.G. Wierda, M.J. Keating, and A.
Rosenwald. 2009. High-level expression of the T-cell chemokines CCL3 and CCL4 by
chronic lymphocytic leukemia B cells in nurselike cell cocultures and after BCR
stimulation. *Blood*. 113:3050-3058.
Buschle, M., D. Campana, S.R. Carding, C. Richard, A.V. Hoffbrand, and M.K. Brenner.
1993. Interferon gamma inhibits apoptotic cell death in B cell chronic lymphocytic
leukemia. *J Exp Med*. 177:213-218.
Cappell, K.M., R.M. Sherry, J.C. Yang, S.L. Goff, D.A. Vanasse, L. McIntyre, S.A.
Rosenberg, and J.N. Kochenderfer. 2020. Long-Term Follow-Up of Anti-CD19
Chimeric Antigen Receptor T-Cell Therapy. *J Clin Oncol*. 38:3805-3815.
Chevrier, D.M., L. Raich, C. Rovira, A. Das, Z. Luo, Q. Yao, A. Chatt, J. Xie, R. Jin, J. Akola,
and P. Zhang. 2018. Molecular-Scale Ligand Effects in Small Gold-Thiolate
Nanoclusters. *J Am Chem Soc*. 140:15430-15436.
Connolly, K.A., M. Kuchroo, A. Venkat, A. Khatun, J. Wang, I. William, N.I. Hornick, B.L.
Fitzgerald, M. Damo, M.Y. Kasmani, C. Cui, E. Fagerberg, I. Monroy, A. Hutchins,
J.F. Cheung, G.G. Foster, D.L. Mariuzza, M. Nader, H. Zhao, W. Cui, S.
Krishnaswamy, and N.S. Joshi. 2021. A reservoir of stem-like CD8(+) T cells in the
tumor-draining lymph node preserves the ongoing antitumor immune response.
Science immunology. 6:eabg7836.

Curigliano, G., H. Gelderblom, N. Mach, T. Doi, D. Tai, P.M. Forde, J. Sarantopoulos, P.L. Bedard, C.C. Lin, F.S. Hodi, S. Wilgenhof, A. Santoro, C.A. Sabatos-Peyton, T.A. Longmire, A. Xyrafas, H. Sun, S. Gutzwiller, L. Manenti, and A. Naing. 2021. Phase I/Ib Clinical Trial of Sabatolimab, an Anti-TIM-3 Antibody, Alone and in Combination with Spartalizumab, an Anti-PD-1 Antibody, in Advanced Solid Tumors. *Clin Cancer Res.* 27:3620-3629.

Daley, D., V.R. Mani, N. Mohan, N. Akkad, A. Ochi, D.W. Heindel, K.B. Lee, C.P. Zambirinis, G.S.B. Pandian, S. Savadkar, A. Torres-Hernandez, S. Nayak, D. Wang, M. Hundeyin, B. Diskin, B. Aykut, G. Werba, R.M. Barilla, R. Rodriguez, S. Chang, L. Gardner, L.K. Mahal, B. Ueberheide, and G. Miller. 2017. Dectin 1 activation on macrophages by Galectin 9 promotes pancreatic carcinoma and peritumoral immune tolerance. *Nat Med.* 23:556-567.

de Mingo Pulido, Á., A. Gardner, S. Hiebler, H. Soliman, H.S. Rugo, M.F. Krummel, L.M. Coussens, and B. Ruffell. 2018. TIM-3 Regulates CD103⁺ Dendritic Cell Function and Response to Chemotherapy in Breast Cancer. *Cancer cell.* 33:60-74.e66.

Dimitrov, D., D. Türei, M. Garrido-Rodriguez, P.L. Burmedi, J.S. Nagai, C. Boys, R.O. Ramirez Flores, H. Kim, B. Szalai, I.G. Costa, A. Valdeolivas, A. Dugourd, and J. Saez-Rodriguez. 2022. Comparison of methods and resources for cell-cell communication inference from single-cell RNA-Seq data. *Nat Commun.* 13:3224.

Ding, W., B.R. LaPlant, T.G. Call, S.A. Parikh, J.F. Leis, R. He, T.D. Shanafelt, S. Sinha, J. Le-Rademacher, A.L. Feldman, T.M. Habermann, T.E. Witzig, G.A. Wiseman, Y. Lin, E. Asmus, G.S. Nowakowski, M.J. Conte, D.A. Bowen, C.N. Aitken, D.L. Van Dyke, P.T. Greipp, X. Liu, X. Wu, H. Zhang, C.R. Secreto, S. Tian, E. Braggio, L.E. Wellik, I. Micallef, D.S. Viswanatha, H. Yan, A.A. Chanan-Khan, N.E. Kay, H. Dong, and S.M. Ansell. 2017. Pembrolizumab in patients with CLL and Richter transformation or with relapsed CLL. *Blood.* 129:3419-3427.

Fraietta, J.A., S.F. Lacey, E.J. Orlando, I. Pruteanu-Malinici, M. Gohil, S. Lundh, A.C. Boesteanu, Y. Wang, R.S. O'Connor, W.-T. Hwang, E. Pequignot, D.E. Ambrose, C. Zhang, N. Wilcox, F. Bedoya, C. Dorfmeier, F. Chen, L. Tian, H. Parakandi, M. Gupta, R.M. Young, F.B. Johnson, I. Kulikovskaya, L. Liu, J. Xu, S.H. Kassim, M.M. Davis, B.L. Levine, N.V. Frey, D.L. Siegel, A.C. Huang, E.J. Wherry, H. Bitter, J.L. Brogdon, D.L. Porter, C.H. June, and J.J. Melenhorst. 2018. Determinants of response and resistance to CD19 chimeric antigen receptor (CAR) T cell therapy of chronic lymphocytic leukemia. *Nat Med.* 24:563-571.

Guo, X., Y. Zhang, L. Zheng, C. Zheng, J. Song, Q. Zhang, B. Kang, Z. Liu, L. Jin, R. Xing, R. Gao, L. Zhang, M. Dong, X. Hu, X. Ren, D. Kirchhoff, H.G. Roider, T. Yan, and Z. Zhang. 2018. Global characterization of T cells in non-small-cell lung cancer by single-cell sequencing. *Nat Med.* 24:978-985.

Gupta, S., R. Bi, K. Su, L. Yel, S. Chiplunkar, and S. Gollapudi. 2004. Characterization of naïve, memory and effector CD8+ T cells: effect of age. *Exp Gerontol.* 39:545-550.

Hahne, F., N. LeMeur, R.R. Brinkman, B. Ellis, P. Haaland, D. Sarkar, J. Spidlen, E. Strain, and R. Gentleman. 2009. flowCore: a Bioconductor package for high throughput flow cytometry. *BMC Bioinformatics.* 10:106.

Hanna, B.S., L. Llaó-Cid, M. Iskar, P.M. Roessner, L.C. Klett, J.K.L. Wong, Y. Paul, N. Ioannou, S. Öztürk, N. Mack, V. Kalter, D. Colomer, E. Campo, J. Bloehdorn, S. Stilgenbauer, S. Dietrich, M. Schmidt, R. Gabriel, K. Rippe, M. Feuerer, A.G. Ramsay, P. Lichter, M. Zapatka, and M. Seiffert. 2021. Interleukin-10 receptor signaling promotes the maintenance of a PD-1(int) TCF-1(+) CD8(+) T cell population that sustains anti-tumor immunity. *Immunity.* 54:2825-2841.e2810.

Hanna, B.S., F. McClanahan, H. Yazdanparast, N. Zaborsky, V. Kalter, P.M. Rößner, A. Benner, C. Dürr, A. Egle, J.G. Gribben, P. Lichter, and M. Seiffert. 2016. Depletion of CLL-associated patrolling monocytes and macrophages controls disease development and repairs immune dysfunction in vivo. *Leukemia.* 30:570-579.

Hanna, B.S., S. Öztürk, and M. Seiffert. 2019. Beyond bystanders: Myeloid cells in chronic lymphocytic leukemia. *Molecular Immunology.* 110:77-87.

Hao, Y., S. Hao, E. Andersen-Nissen, W.M. Mauck, 3rd, S. Zheng, A. Butler, M.J. Lee, A.J.
Wilk, C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E.P. Mimitou, J.
Jain, A. Srivastava, T. Stuart, L.M. Fleming, B. Yeung, A.J. Rogers, J.M. McElrath,
C.A. Blish, R. Gottardo, P. Smibert, and R. Satija. 2021. Integrated analysis of
multimodal single-cell data. *Cell*. 184:3573-3587.e3529.
Harris, C.R., E.T. McKinley, J.T. Roland, Q. Liu, M.J. Shrubsole, K.S. Lau, R.J. Coffey, J.
Wrobel, and S.N. Vandekar. 2022. Quantifying and correcting slide-to-slide variation
in multiplexed immunofluorescence images. *Bioinformatics*. 38:1700-1707.
Herbst, S.A., M. Vesterlund, A.J. Helmboldt, R. Jafari, I. Siavelis, M. Stahl, E.C. Schitter, N.
Liebers, B.J. Brinkmann, F. Czernilofsky, T. Roider, P.M. Bruch, M. Iskar, A. Kittai, Y.
Huang, J. Lu, S. Richter, G. Mermelekas, H.M. Umer, M. Knoll, C. Kolb, A. Lenze, X.
Cao, C. Österholm, L. Wahnschaffe, C. Herling, S. Scheinost, M. Ganzinger, L.
Mansouri, K. Kriegsmann, M. Kriegsmann, S. Anders, M. Zapatka, G. Del Poeta, A.
Zucchetto, R. Bomben, V. Gattei, P. Dreger, J. Woyach, M. Herling, C. Müller-Tidow,
R. Rosenquist, S. Stilgenbauer, T. Zenz, W. Huber, E. Tausch, J. Lehtiö, and S.
Dietrich. 2022. Proteogenomics refines the molecular classification of chronic
lymphocytic leukemia. *Nat Commun*. 13:6226.
Huang, Q., X. Wu, Z. Wang, X. Chen, L. Wang, Y. Lu, D. Xiong, Q. Liu, Y. Tian, H. Lin, J.
Guo, S. Wen, W. Dong, X. Yang, Y. Yuan, Z. Yue, S. Lei, Q. Wu, L. Ran, L. Xie, Y.
Wang, L. Gao, Q. Tian, X. Zhou, B. Sun, L. Xu, Z. Tang, and L. Ye. 2022. The
primordial differentiation of tumor-specific memory CD8(+) T cells as bona fide
responders to PD-1/PD-L1 blockade in draining lymph nodes. *Cell*. 185:4049-
4066.e4025.
Im, S.J., M. Hashimoto, M.Y. Gerner, J. Lee, H.T. Kissick, M.C. Burger, Q. Shan, J.S. Hale,
J. Lee, T.H. Nasti, A.H. Sharpe, G.J. Freeman, R.N. Germain, H.I. Nakaya, H.-H.
Xue, and R. Ahmed. 2016. Defining CD8+ T cells that provide the proliferative burst
after PD-1 therapy. *Nature*. 537:417.
Jin, S., C.F. Guerrero-Juarez, L. Zhang, I. Chang, R. Ramos, C.H. Kuan, P. Myung, M.V.
Plikus, and Q. Nie. 2021. Inference and analysis of cell-cell communication using
CellChat. *Nat Commun*. 12:1088.
Jitschin, R., M. Braun, M. Büttner, K. Dettmer-Wilde, J. Bricks, J. Berger, M.J. Eckart, S.W.
Krause, P.J. Oefner, K. Le Blanc, A. Mackensen, and D. Mougiakakos. 2014. CLL-
cells induce IDOhi CD14+HLA-DRlo myeloid-derived suppressor cells that inhibit T-
cell responses and promote TRegs. *Blood*. 124:750-760.
Joshi, N.S., W. Cui, A. Chandele, H.K. Lee, D.R. Urso, J. Hagman, L. Gapin, and S.M.
Kaech. 2007. Inflammation directs memory precursor and short-lived effector CD8(+)
T cell fates via the graded expression of T-bet transcription factor. *Immunity*. 27:281-
295.
Karabon, L., A. Partyka, L. Ciszak, E. Pawlak-Adamska, A. Tomkiewicz, A. Bojarska-Junak,
J. Roliński, D. Wołowiec, T. Wrobel, I. Frydecka, and A. Kosmaczewska. 2020.
Abnormal Expression of BTLA and CTLA-4 Immune Checkpoint Molecules in Chronic
Lymphocytic Leukemia Patients. *J Immunol Res*. 2020:6545921.
Kiaii, S., P. Kokhaei, F. Mozaffari, E. Rossmann, F. Pak, A. Moshfegh, M. Palma, L.
Hansson, K. Mashayekhi, M. Hojjat-Farsangi, A. Österborg, A. Choudhury, and H.
Mellstedt. 2013. T cells from indolent CLL patients prevent apoptosis of leukemic B
cells in vitro and have altered gene expression profile. *Cancer immunology,
immunotherapy : CII*. 62:51-63.
Korsunsky, I., N. Millard, J. Fan, K. Slowikowski, F. Zhang, K. Wei, Y. Baglaenko, M.
Brenner, P.R. Loh, and S. Raychaudhuri. 2019. Fast, sensitive and accurate
integration of single-cell data with Harmony. *Nat Methods*. 16:1289-1296.
Lee, M., J.A.G. Hamilton, G.R. Talekar, A.J. Ross, L. Michael, M. Rupji, B. Dwivedi, S.S.
Raikar, J. Boss, C.D. Scharer, D.K. Graham, D. DeRyckere, C.C. Porter, and C.J.
Henry. 2022. Obesity-induced Galectin-9 is a therapeutic target in B-cell acute
lymphoblastic leukemia. *Nat Commun*. 13:1157.
Levine, J.H., E.F. Simonds, S.C. Bendall, K.L. Davis, A.D. Amir el, M.D. Tadmor, O. Litvin,
H.G. Fienberg, A. Jager, E.R. Zunder, R. Finck, A.L. Gedman, I. Radtke, J.R.

Downing, D. Pe'er, and G.P. Nolan. 2015. Data-Driven Phenotypic Dissection of AML
Reveals Progenitor-like Cells that Correlate with Prognosis. *Cell*. 162:184-197.
Lin, J.-R., Y.-A. Chen, D. Campton, J. Cooper, S. Coy, C. Yapp, J.B. Tefft, E. McCarty, K.L.
Ligon, S.J. Rodig, S. Reese, T. George, S. Santagata, and P.K. Sorger. 2022. Multi-
modal digital pathology for colorectal cancer diagnosis by high-plex
immunofluorescence imaging and traditional histology of the same tissue section.
bioRxiv.
Lin, J.R., B. Izar, S. Wang, C. Yapp, S. Mei, P.M. Shah, S. Santagata, and P.K. Sorger.
2018. Highly multiplexed immunofluorescence imaging of human tissues and tumors
using t-CyCIF and conventional optical microscopes. *Elife*. 7.
Liu, B., X. Hu, K. Feng, R. Gao, Z. Xue, S. Zhang, Y. Zhang, E. Corse, Y. Hu, W. Han, and Z.
Zhang. 2022. Temporal single-cell tracing reveals clonal revival and expansion of
precursor exhausted T cells during anti-PD-1 therapy in lung cancer. *Nat Cancer*.
3:108-121.
McGinnis, C.S., L.M. Murrow, and Z.J. Gartner. 2019. DoubletFinder: Doublet Detection in
Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst*.
8:329-337.e324.
McLane, L.M., M.S. Abdel-Hakeem, and E.J. Wherry. 2019. CD8 T Cell Exhaustion During
Chronic Viral Infection and Cancer. *Annual review of immunology*. 37:457-495.
Obar, J.J., E.R. Jellison, B.S. Sheridan, D.A. Blair, Q.-M. Pham, J.M. Zickovich, and L.
Lefrançois. 2011. Pathogen-Induced Inflammatory Environment Controls Effector and
Memory CD8$^+$ T Cell Differentiation. *The Journal of Immunology*. 187:4967-4978.
Oh, D.Y., and L. Fong. 2021. Cytotoxic CD4(+) T cells in cancer: Expanding the immune
effector toolbox. *Immunity*. 54:2701-2711.
Oh, D.Y., S.S. Kwek, S.S. Raju, T. Li, E. McCarthy, E. Chow, D. Aran, A. Ilano, C.S. Pai, C.
Rancan, K. Allaire, A. Burra, Y. Sun, M.H. Spitzer, S. Mangul, S. Porten, M.V. Meng,
T.W. Friedlander, C.J. Ye, and L. Fong. 2020. Intratumoral CD4(+) T Cells Mediate
Anti-tumor Cytotoxicity in Human Bladder Cancer. *Cell*. 181:1612-1625.e1613.
Pang, N., X. Alimu, R. Chen, M. Muhashi, J. Ma, G. Chen, F. Zhao, L. Wang, J. Qu, and J.
Ding. 2021a. Activated Galectin-9/Tim3 promotes Treg and suppresses Th1 effector
function in chronic lymphocytic leukemia. *Faseb j*. 35:e21556.
Pang, N., X. Alimu, R. Chen, M. Muhashi, J. Ma, G. Chen, F. Zhao, L. Wang, J. Qu, and J.
Ding. 2021b. Activated Galectin-9/Tim3 promotes Treg and suppresses Th1 effector
function in chronic lymphocytic leukemia. *The FASEB Journal*. 35:e21556.
Quezada, S.A., T.R. Simpson, K.S. Peggs, T. Merghoub, J. Vider, X. Fan, R. Blasberg, H.
Yagita, P. Muranski, P.A. Antony, N.P. Restifo, and J.P. Allison. 2010. Tumor-reactive
CD4(+) T cells develop cytotoxic activity and eradicate large established melanoma
after transfer into lymphopenic hosts. *J Exp Med*. 207:637-650.
Reinart, N., P.H. Nguyen, J. Boucas, N. Rosen, H.M. Kvasnicka, L. Heukamp, C. Rudolph, V.
Ristovska, T. Velmans, C. Mueller, K.S. Reiners, E.P. von Strandmann, G. Krause,
M. Montesinos-Rongen, B. Schlegelberger, M. Herling, M. Hallek, and G. Fingerle-
Rowson. 2013. Delayed development of chronic lymphocytic leukemia in the absence
of macrophage migration inhibitory factor. *Blood*. 121:812-821.
Roessner, P.M., L. Llaó Cid, E. Lupar, T. Roider, M. Bordas, C. Schifflers, L. Arseni, A.C.
Gaupel, F. Kilpert, M. Krötschel, S.J. Arnold, L. Sellner, D. Colomer, S. Stilgenbauer,
S. Dietrich, P. Lichter, A. Izcue, and M. Seiffert. 2021. EOMES and IL-10 regulate
antitumor activity of T regulatory type 1 CD4(+) T cells in chronic lymphocytic
leukemia. *Leukemia*. 35:2311-2324.
Roider, T., M.A. Baertsch, D. Fitzgerald, H. Voehringer, B.J. Brinkmann, F. Czernilofsky, M.
Knoll, L. Llao-Cid, P.-M. Bruch, N. Liebers, C.M. Schuerch, V. Passerini, A. Brobeil,
G. Mechtersheimer, C. Mueller-Tidow, O. Weigert, M. Seiffert, G.P. Nolan, W. Huber,
and S. Dietrich. 2022. Multimodal and spatially resolved profiling identifies distinct
patterns of T-cell infiltration in nodal B-cell lymphoma entities.
bioRxiv:2022.2011.2004.514366.
Roider, T., J. Seufert, A. Uvarovskii, F. Frauhammer, M. Bordas, N. Abedpour, M. Stolarczyk,
J.P. Mallm, S.A. Herbst, P.M. Bruch, H. Balke-Want, M. Hundemer, K. Rippe, B.

Goeppert, M. Seiffert, B. Brors, G. Mechtersheimer, T. Zenz, M. Peifer, B. Chapuy, M. Schlesner, C. Muller-Tidow, S. Frohling, W. Huber, S. Anders, and S. Dietrich. 2020. Dissecting intratumour heterogeneity of nodal B-cell lymphomas at the transcriptional, genetic and drug-response levels. *Nature cell biology*. 22:896-+.

Sabatos-Peyton, C., T. Longmire, L. Baker, N. Patel, A.-S. Wavreille, M. Verneret, P. Jayaraman, X. Jiang, S. Schwartz, V. Cremasco, H. Lu, S. Qiu, F. Sharp, M. Rinne, and G. Dranoff. 2020. 439 Dual modes of action for anti-TIM-3 antibody MBG453 in myelodysplastic syndromes (MDS) and acute myeloid leukemia (AML): preclinical evidence for immune-mediated and anti-leukemic activity. *Journal for ImmunoTherapy of Cancer*. 8:A267-A267.

Sabatos-Peyton, C.A., J. Nevin, A. Brock, J.D. Venable, D.J. Tan, N. Kassam, F. Xu, J. Taraszka, L. Wesemann, T. Pertel, N. Acharya, M. Klapholz, Y. Etminan, X. Jiang, Y.H. Huang, R.S. Blumberg, V.K. Kuchroo, and A.C. Anderson. 2018. Blockade of Tim-3 binding to phosphatidylserine and CEACAM1 is a shared feature of anti-Tim-3 antibodies that have functional efficacy. *Oncoimmunology*. 7:e1385690.

Saka, S.K., Y. Wang, J.Y. Kishi, A. Zhu, Y. Zeng, W. Xie, K. Kirli, C. Yapp, M. Cicconet, B.J. Beliveau, S.W. Lapan, S. Yin, M. Lin, E.S. Boyden, P.S. Kaeser, G. Pihan, G.M. Church, and P. Yin. 2019. Immuno-SABER enables highly multiplexed and amplified protein imaging in tissues. *Nature biotechnology*. 37:1080-1090.

Sawant, D.V., H. Yano, M. Chikina, Q. Zhang, M. Liao, C. Liu, D.J. Callahan, Z. Sun, T. Sun, T. Tabib, A. Pennathur, D.B. Corry, J.D. Luketich, R. Lafyatis, W. Chen, A.C. Poholek, T.C. Bruno, C.J. Workman, and D.A.A. Vignali. 2019. Adaptive plasticity of IL-10(+) and IL-35(+) T(reg) cells cooperatively promotes tumor T cell exhaustion. *Nat Immunol*. 20:724-735.

Schapiro, D., A. Sokolov, C. Yapp, Y.A. Chen, J.L. Muhlich, J. Hess, A.L. Creason, A.J. Nirmal, G.J. Baker, M.K. Nariya, J.R. Lin, Z. Maliga, C.A. Jacobson, M.W. Hodgman, J. Ruokonen, S.L. Farhi, D. Abbondanza, E.T. McKinley, D. Persson, C. Betts, S. Sivagnanam, A. Regev, J. Goecks, R.J. Coffey, L.M. Coussens, S. Santagata, and P.K. Sorger. 2022. MCMICRO: a scalable, modular image-processing pipeline for multiplexed tissue imaging. *Nat Methods*. 19:311-315.

Scott, A.C., F. Dündar, P. Zumbo, S.S. Chandran, C.A. Klebanoff, M. Shakiba, P. Trivedi, L. Menocal, H. Appleby, S. Camara, D. Zamarin, T. Walther, A. Snyder, M.R. Femia, E.A. Comen, H.Y. Wen, M.D. Hellmann, N. Anandasabapathy, Y. Liu, N.K. Altorki, P. Lauer, O. Levy, M.S. Glickman, J. Kaye, D. Betel, M. Philip, and A. Schietinger. 2019. TOX is a critical regulator of tumour-specific T cell differentiation. *Nature*. 571:270-274.

Türei, D., T. Korcsmáros, and J. Saez-Rodriguez. 2016. OmniPath: guidelines and gateway for literature-curated signaling pathway resources. *Nat Methods*. 13:966-967.

Türei, D., A. Valdeolivas, L. Gul, N. Palacio-Escat, M. Klein, O. Ivanova, M. Ölbei, A. Gábor, F. Theis, D. Módos, T. Korcsmáros, and J. Saez-Rodriguez. 2021. Integrated intra- and intercellular signaling knowledge for multicellular omics analysis. *Mol Syst Biol*. 17:e9923.

Utzschneider, D.T., M. Charmoy, V. Chennupati, L. Pousse, D.P. Ferreira, S. Calderon-Copete, M. Danilo, F. Alfei, M. Hofmann, D. Wieland, S. Pradervand, R. Thimme, D. Zehn, and W. Held. 2016. T Cell Factor 1-Expressing Memory-like CD8(+) T Cells Sustain the Immune Response to Chronic Viral Infections. *Immunity*. 45:415-427.

van Bruggen, J.A.C., A.W.J. Martens, J.A. Fraietta, T. Hofland, S.H. Tonino, E. Eldering, M.D. Levin, P.J. Siska, S. Endstra, J.C. Rathmell, C.H. June, D.L. Porter, J.J. Melenhorst, A.P. Kater, and G.J.W. van der Windt. 2019. Chronic lymphocytic leukemia cells impair mitochondrial fitness in CD8(+) T cells and impede CAR T-cell efficacy. *Blood*. 134:44-58.

Van den Berge, K., H. Roux de Bezieux, K. Street, W. Saelens, R. Cannoodt, Y. Saeys, S. Dudoit, and L. Clement. 2020. Trajectory-based differential expression analysis for single-cell sequencing data. *Nat Commun*. 11:1201.

Van Gassen, S., B. Callebaut, M.J. Van Helden, B.N. Lambrecht, P. Demeester, T. Dhaene, and Y. Saeys. 2015. FlowSOM: Using self-organizing maps for visualization and interpretation of cytometry data. *Cytometry A*. 87:636-645.

Vardi, A., E. Vlachonikola, M. Karypidou, E. Stalika, V. Bikos, K. Gemenetzi, C. Maramis, A. Siorenta, A. Anagnostopoulos, S. Pospisilova, N. Maglaveras, I. Chouvarda, K. Stamatopoulos, and A. Hadzidimitriou. 2017. Restrictions in the T-cell repertoire of chronic lymphocytic leukemia: high-throughput immunoprofiling supports selection by shared antigenic elements. *Leukemia*. 31:1555-1561.

Wdowiak, K., E. Gallego Colon, T. Francuz, P. Czajka Francuz, N. Ruiz Agamez, M. Kubeczko, I. Grochoła, M.T. Wybraniec, J. Chudek, and J. Wojnar. 2019. Increased serum levels of Galectin 9 in patients with chronic lymphocytic leukemia. *Oncol Lett*. 17:1019-1029.

Wierz, M., S. Pierson, L. Guyonnet, E. Viry, A. Lequeux, A. Oudin, S.P. Niclou, M. Ollert, G. Berchem, B. Janji, C. Guérin, J. Paggetti, and E. Moussay. 2018. Dual PD1/LAG3 immune checkpoint blockade limits tumor development in a murine model of chronic lymphocytic leukemia. *Blood*. 131:1617-1621.

Wilkerson, M.D., and D.N. Hayes. 2010. ConsensusClusterPlus: a class discovery tool with confidence assessments and item tracking. *Bioinformatics*. 26:1572-1573.

Wolf, Y., A.C. Anderson, and V.K. Kuchroo. 2020. TIM3 comes of age as an inhibitory receptor. *Nat Rev Immunol*. 20:173-185.

Wu, T.D., S. Madireddi, P.E. de Almeida, R. Banchereau, Y.J. Chen, A.S. Chitre, E.Y. Chiang, H. Iftikhar, W.E. O'Gorman, A. Au-Yeung, C. Takahashi, L.D. Goldstein, C. Poon, S. Keerthivasan, D.E. de Almeida Nagata, X. Du, H.M. Lee, K.L. Banta, S. Mariathasan, M. Das Thakur, M.A. Huseni, M. Ballinger, I. Estay, P. Caplazi, Z. Modrusan, L. Delamarre, I. Mellman, R. Bourgon, and J.L. Grogan. 2020. Peripheral T cell expansion predicts tumour infiltration and clinical response. *Nature*. 579:274-278.

Xie, Y., A. Akpinarli, C. Maris, E.L. Hipkiss, M. Lane, E.K. Kwon, P. Muranski, N.P. Restifo, and P.A. Antony. 2010. Naive tumor-specific CD4(+) T cells differentiated in vivo eradicate established melanoma. *J Exp Med*. 207:651-667.

Xierenguli, A., X. Zeng, N. Pang, R. Zhagn, J. Ma, Y. Zhao, and J. Qu. 2020. [TIM-3/Galectin-9 is involved in negative regulation of T cells in patients with chronic lymphocytic leukemia]. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi*. 36:1021-1025.

Xu-Monette, Z.Y., J. Zhou, and K.H. Young. 2018. PD-1 expression and clinical PD-1 blockade in B-cell lymphomas. *Blood*. 131:68-83.

Yang, R., L. Sun, C.F. Li, Y.H. Wang, J. Yao, H. Li, M. Yan, W.C. Chang, J.M. Hsu, J.H. Cha, J.L. Hsu, C.W. Chou, X. Sun, Y. Deng, C.K. Chou, D. Yu, and M.C. Hung. 2021. Galectin-9 interacts with PD-1 and TIM-3 to regulate T cell death and is a target for cancer immunotherapy. *Nat Commun*. 12:832.

Yang, R.Y., G.A. Rabinovich, and F.T. Liu. 2008. Galectins: structure, function and therapeutic potential. *Expert Rev Mol Med*. 10:e17.

Yapp, C., E. Novikov, W.D. Jang, T. Vallius, Y.A. Chen, M. Cicconet, Z. Maliga, C.A. Jacobson, D. Wei, S. Santagata, H. Pfister, and P.K. Sorger. 2022. UnMICST: Deep learning with real augmentation for robust segmentation of highly multiplexed images of human tissues. *Commun Biol*. 5:1263.

Zheng, L., S. Qin, W. Si, A. Wang, B. Xing, R. Gao, X. Ren, L. Wang, X. Wu, J. Zhang, N. Wu, N. Zhang, H. Zheng, H. Ouyang, K. Chen, Z. Bu, X. Hu, J. Ji, and Z. Zhang. 2021. Pan-cancer single-cell landscape of tumor-infiltrating T cells. *Science*. 374:abe6474.

Zhu, C., A.C. Anderson, A. Schubart, H. Xiong, J. Imitola, S.J. Khoury, X.X. Zheng, T.B. Strom, and V.K. Kuchroo. 2005. The Tim-3 ligand Galectin-9 negatively regulates T helper type 1 immunity. *Nat Immunol*. 6:1245-1252.

Acknowledgements

We would like to thank all members of the High Throughput Sequencing Unit of the Genomics and Proteomics Core Facility, of the Light Microscopy Core Facility, and of the Single-cell Open Lab (scOpenLab) at the German Cancer Research Center (DKFZ, Heidelberg) for their technical assistance. We also thank the National Cytometry Platform (NCP) of the Luxembourg Institute of Health (LIH, Luxembourg) for assistance with the generation of cytometry data, in particular Maira Konstantinou and Dominique Revets. The NCP is supported by funding from Luxembourg's Ministry of Higher Education and Research (MESR). Tissue microarrays and lymph node sections were provided by the Tissue Bank of the National Center for Tumor Diseases (NCT) Heidelberg, Germany in accordance with the regulations of the tissue bank and the approval of the ethics committee of Heidelberg University.

This study was supported by a research grant of the German Cancer Aid (Deutsche Krebshilfe) to MS, by grants from FNRS-Télévie to IFB (7.4529.19, 7.6603.21), MW (7.6504.18), GP (7.4501.18, 7.6518.20), SG (7.4502.19, 7.6604.21), and from the Luxembourg National Research Fund (FNR), Fondation Cancer and Plooschter Projet to EM and JP (C20/BM/14582635, and C20/BM/14592342). The authors acknowledge support by the state of Baden-Württemberg through bwHPC and the German Research Foundation (DFG) through grant INST 35/1597-1 FUGG. MC, KB and DS are supported by the German Federal Ministry of Education and Research (BMBF 01ZZ2004).

The authors acknowledge support by the state of Baden-Württemberg through bwHPC and the German Research Foundation (DFG) through grant INST 35/1597-1 FUGG. M.C., K.B and D.S are supported by the German Federal Ministry of Education and Research (BMBF 01ZZ2004).

Author information

L.L.C. designed the study, performed experiments, analyzed data, generated figures, and wrote the paper. J.K.L.W. performed bioinformatical analyses, generated figures, and wrote the paper. I.F.B. and M.W. conceptualized part of the study, performed experiments, and analyzed data. Y.P. analyzed scRNA-seq data and CyTOF data. A.F. conducted experiments, analyzed data, and generated figures. S.G., G.P., C.S. A.C., and D.E.C. performed experiments and analyzed data. M.C., K.B. and D.S. analyzed multiplex imaging data, and generated figures. M.I. performed bioinformatical analyses and generated figures. T.R. and S.D. provided input regarding cluster annotations and performed correlation analyses. J.P.M. established the pipeline for single-cell RNA-sequencing. E.G.-H. and A.R. supported the study with lymph node sections and expertise in interpreting imaging data of

1294 lymph nodes. D.C., E.C., and S.D. provided clinical samples and information. P.L. provided
1295 logistic and budget support and was involved in scientific discussions. E.M., J.P., M.Z., and
1296 M.S., conceptualized the study, guided experiments and data analysis, provided logistic and
1297 budget support, and wrote the paper.
1298

Ethics declarations
Samples from patients and heathy controls were obtained after informed consent and
according to the guidelines of the Ethics Committees of the involved University hospitals and
the Declaration of Helsinki.
All experiments involving laboratory animals were conducted in pathogen-free animal
facilities at the German Cancer Research Center in Heidelberg or the Luxembourg Institute
of Health in Luxembourg with the approval of the Regierungspräsidium Karlruhe (G-77/19
and G-112/21) and the Luxembourg Ministry for Agriculture (#LUPA 2019/21), respectively.
Mice were treated in accordance with the European guidelines.
1308

1309 **Supplemental Excel Table Titles**
1310 **Suppl. Table 1:** Information on patients and donors
1311 **Suppl. Table 2:** Cell cluster annotations
1312 **Suppl. Table 3:** Correlations with clinical data
1313 **Suppl. Table 4:** VDJdb results of TCR-seq data