1

4 **Title**

5 A reductionist paradigm for high-throughput behavioural

6 fingerprinting in *Drosophila melanogaster*

11 **Authors and affiliations**

12 Hannah Jones[1], Jenny A Willis[2], Lucy C Firth[2], Carlo N G Giachello[2], and Giorgio F Gilestro[1#]

14 *# To whom correspondence should be addressed. E-mail: giorgio@gilest.ro*

16 [1]Department of Life Sciences, Imperial College London, London, UK

17 [2]Syngenta, Jealott's Hill International Research Centre, Bracknell, UK

Abstract

Understanding how the brain encodes behaviour is the ultimate goal of neuroscience and the ability to

objectively and reproducibly describe and quantify behaviour is a necessary milestone on this path.

Recent technological progresses in machine learning and computational power have boosted the

development and adoption of systems leveraging on high-resolution video recording to track an animal

pose and describe behaviour in all four dimensions. However, the high temporal and spatial resolution

that these systems offer must come as a compromise with their throughput and accessibility. Here we

describe *coccinella*, an open-source reductionist framework combining high-throughput analysis of

behaviour using real-time tracking on a distributed mesh of microcomputers (ethoscopes) with

resource-lean statistical learning (HCTSA/Catch22). Coccinella is a reductionist system, yet

outperforms state-of-the-art alternatives when exploring the pharmacobehaviour in *Drosophila*

melanogaster.

Introduction

The nervous system integrates stimuli, internal states, expectations, and previous experience to regulate

behavioural output. Describing, quantifying, and modulating behaviour are critical aspects of modern

neuroscience and, ever since its inception, the field has spent considerable effort into building and

sharing paradigms or tools aimed at objectively and reproducibly quantify behaviours in the most

disparate animal models, to the point that this exercise is now recognised as an exciting subfield of

neuroscience in its own right: ethomics[1,2]. As the portmanteau name itself suggests, ethomics is not just

about describing behaviour (*etho-*) but also about doing so in a high-throughput fashion (*-omics*),

collecting data simultaneously from a large number of individuals, which can remain undisturbed

throughout recording. Irrespective of the behaviour or the animal model to be analysed, the first

compromise a researcher will face when choosing a tool for behavioural quantification will always be

between throughput and resolution: a high-throughput analysis will allow for powerful experimental

manipulations – such as genetics or pharmacological screens – offering unbiased approaches in

identifying neuronal circuits, genes, molecules underpinning behaviour; high-resolution analysis, on

the other hand, promises to identify and discriminate even minuscule differences that may not be

immediately visible to the human eye, and to label behaviours into identifiable classes (*e.g.:*

"grooming", "courting", "shaking") that may be more relevant to researchers interested in modelling

disease or in anthropomorphic descriptions. In the past years, the field has generally converged towards

the adoption of high-resolution video recording of activity, in some cases adopting cameras that have

milliseconds temporal resolution or developing setups that provide depth information for three-

dimensional reconstruction of motion or posture[3–7]. Given the recent evolution in machine learning and

progresses in computational power, even these high-resolution analyses can be at least in part

compatible with high-throughput approaches[8], especially when employed on small invertebrate animal

models[9–12] or when aided by robotic handling[13]. These systems, however, can still be prohibitively

expensive for most laboratories, and not easily compatible with throughput in the '*omics* scale.

Moreover, besides the technical urge of removing entry barrier and make ethomics an accessible tool,

an equally important underlying question concerns what is the minimal amount of information that

needs to be extracted to identify and classify behaviour. Do we always necessarily gain information

from extracting micro-postural features or by analysing activity in three dimensions? To what extent

this may actually add counterproductive biological noise to some assays?

Here we introduce *coccinella*: a new experimental framework that combines high-throughput,

inexpensive, real-time ethomics[14] with state-of-the-art statistical analysis[15,16] to characterise and

discriminate complex behaviours using a reductionist approach based solely on one simple feature

(https://lab.gilest.ro/coccinella). *Coccinella* builds on ethoscopes[14], an accessible open-source platform,

to extract, in real-time, minimalist information from flies. Despite its minimalist nature, *coccinella*

outperforms state-of-the-art alternatives in recognising the pharmacobehavioural space, providing

better discernibility at a fraction of the cost, thus opening a new path to high-throughput ethomics.

Results

Drosophila ethomics studies generally rely on image acquisition through so called industrial-cameras,

able to collect videos with high temporal and spatial resolution and featuring mounts for a large

selection of lenses. Some of the cameras commonly used for these purposes (*e.g.:* FLIR, Point Grey,

Basler)[17] are expensive and normally employed in close-up imaging that microscopically highlights the

smallest anatomical features of the animal but at the same time greatly limits the number of

experimental subjects that can be recorded by a single device. Normally, one or few more camera

would be connected to a dedicated powerful computer for acquisition and storage of videos. The cost

and physical footprint of these setups makes them incompatible, at least for most laboratories, with

high-throughput simultaneous acquisition. To lower this barrier, we created a framework that employs

the distributing computing power of ethoscopes[14], thus allowing for inexpensive analysis of activity in

hundreds or thousands of flies at once. Ethoscopes are open source and can be manufactured by a

skilled end-user at a cost of about £75 per machine, mostly building on two off-the-shelf component: a

Raspberry Pi microcomputer and a Raspberry Pi NoIR camera overlooking a bespoke 3D printed arena

hosting freely moving flies. The temporal and spatial resolution of the collected images depends on the

working modality the user chooses. When operating in offline mode, ethoscopes are capable to acquire

720p videos at 60 fps, which is a convenient option with fast moving animals. In this study, we instead

opted for the default ethoscope working settings, providing online tracking and real-time parametric

extraction, meaning that images are analysed by each raspberry Pi at the very moment they are

acquired (Figure 1b). This latter modality limits the temporal resolution of information being processed

(one frame every 444 ms $\pm$ 127 ms, equivalent to 2.2 fps on a Raspberry Pi3 at a resolution of

1280x960 pixels with each animal being constricted in an ellipse measuring 25.8 $\pm$ 1.4 x 9.85 $\pm$ 1.4

pixels – Figure 1a) but provides the most affordable and high-throughput solution, dispensing the

researcher from organising video storage or asynchronous video processing for tracking the animals. In

the work here described, flies moved freely in a circular two-dimensional space with a diameter of

11.5 mm designed to maintain the animal in the walking position[18] while venturing on solidified agar

providing nutrients alone or nutrients and drugs. In previous analysis of activity and sleep[14,19], we found

that the maximal velocity of the fly over a period of 10 seconds best described the basic motion

features of the animal, allowing us to accurately differentiate between different activity patterns, such

as walking, grooming, feeding[14]. We therefore adopted this measure for *coccinella* too, ultimately

producing monodimensional time-series of a behavioural correlate, which were then digested using

highly comparative time-series analysis (HCTSA)[15], a computational framework that effortlessly

subjects time-series to more than 7700 literature-relevant statistical tests, looking for meaningful

discriminative features. Features successfully extracted through HCTSA were then used to classify

behaviour using a linear support vector machine (SVM$_{linear}$)[20] (Figure 1b) and here presented and

compared using confusion matrices (Figure 1 - Figure Supplement 1a). To test the ability of this system

to discriminate behaviour, we started by exploring the pharmacobehavioral space of flies fed with a

panel of known or putative neurotropic chemicals, comprising molecules previously described in the

literature along with uncharacterised ones being considered as potential insecticides (Figure 1c, 1d and

Supplementary Table 1). Using an initial panel of 17 treatments (16 drugs and one solvent control) we

were able to discern compounds with an accuracy of 71.4% (vs. 5.8% of a random classifier – Figure

1c). Some compounds induced behaviours with a particularly high predictive fingerprint: dieldrin, for

instance, was predicted with an accuracy of 94% and flonicamid with an accuracy of 87%. For others,

our system fared more poorly (*e.g.* tetramethrin showed 41% accuracy). In all cases, however, the

relative confusion was negligible, with all compounds being correctly identified as the first choice and

with the first predicted compounds having, on average, a score that was 15-times greater compared to

the second-best choices (Figure 1c – min.: 3.3x; max.: 65x). To validate our framework and exclude

artifacts operated by overfitting biologically irrelevant information we followed two lines of control.

Firstly, we fed flies with lower concentrations of the same compounds (Figure 1c). Feeding flies with

different concentrations of drugs unsurprisingly showed a different effect on short term lethality

(Figure 1 - Figure Supplement 1b), with several compounds hitting a 25% lethality rate before the end

of the experiment when fed at the highest concentration (1000ppm – Figure 1 - Figure Supplement 1b).

Lowering the compound concentration, the predictive accuracy of the system decreased from 71.4%

(1000ppm) to 61.8% (100ppm), falling to 36.1% with the lowest concentrations (1ppm), indicating that

the system does operate on pharmacologically induced, biologically meaningful behavioural correlates

(Figure 1c). A similar drop in accuracy was observed using a smaller panel of 12 treatments (Figure 1 -

Figure Supplement 1c). As a second line of work to test specificity, we obtained genetic mutants known

to be resistant to specific pharmacological treatments: the *para*^{L1029F} allele encodes for a version of the

α-subunit of voltage-gated sodium channel conferring resistance to DDT and pyrethroids[21]; the *Rdl*^{A301S}

allele encodes for a version of the ligand-gated chloride channel conferring resistance to dieldrin and

fiproles[22]. Challenging these mutants with their respective compounds created confusion in the

clustering algorithm for which discerning between drug treatment and solvent control became a harder

task, especially in the case of DDT and *para*^{L1029F} (Figure 1 - Figure Supplement 2a). The observed drop

in accuracy suggests again that *coccinella* is working on biologically relevant behavioural signatures

and the fact that some discrimination can still be observed with targets harbouring point mutations –

which should severely affect compound efficacy – is indicative of high sensitivity.

Having established the accuracy and sensitivity of the system, we next wanted to test its usefulness in a

genuine high-throughput scenario. We subjected a total of 2192 flies to a panel of 40 treatments

(Figure 1d), mostly featuring known compounds but also two unexplored molecules (Supplementary

Table 1). Given that the 100ppm intermediate concentrations showed the best compromise between

accuracy and lethality in the previous pilot experiment (Figure 1 - Figure Supplement 1b), we

performed this larger screen using compounds diluted at 100ppm only. Even with such a large panel,

the system was able to first-guess 39 out of 40 of the tested treatments (the only exception being the

Syngenta Compound #5) with an overall accuracy of 44.5% vs 2.5% of the random classifier.

A reductionist approach served us well so far, identifying with remarkable accuracy even subtle

changes when we explored the pharmacobehavioural space of a large number of neuroactive

compounds in wild type and mutant flies. But how does it compare to other more established

paradigms? *Coccinella* is arguably to be preferred in terms of accessibility and throughput, but what is

the amount of useful information that we are sacrificing by adopting a reductionist approach? To

quantify any possible loss in information content, we ran a series of parallel experiments in which flies

were fed with a selected panel of 12 treatments (11 drugs and a solvent control, Figure 2) and their

behaviour analysed either using *coccinella* or using other widely adopted state-of-the-art methods,

which started with high-resolution imaging and employed supervised machine learning for pose

estimation (DeepLabCut[17]) followed by unsupervised identification of behavioural grammar (B-

SOiD[4]). To widen the range of comparisons, data were then either immediately clustered using different

common clustering algorithms (K-nearest neighbours, random forest) or first processed through a

smaller, selected subsample of the HCTSA array (Catch22[16]) before being clustered using SVM_{linear}

(Figure 2a). In this challenge, *coccinella* unambiguously identified 10 out of 12 compounds, with poor

performance only for two of them (flubendiamide and tetramethrin) and an overall accuracy of 42.4%,

vs 8.3% of the random classifier (Figure 2b). Surprisingly, none of the state-of-the-art high-resolution

paths did better than this. The combination of pose-estimation → grammar extraction → random forest

classification scored as the second best, with an accuracy of 25.4% but with only three compounds

being unambiguously identified (Figure 2c). The same experimental dataset clustered with even poorer

performance when using K-nearest neighbours (Figure 2e) and even the application of HCTSA features

extraction to the B-SoID output still could not match the accuracy observed with *coccinella*'s

reductionist approach (Figure 2d). This analysis is not meant to be conclusive. We expect that some

alternative combination of state-of-the-art approaches will probably manage to match or likely improve

over *coccinella*'s performance, yet the fact we could obtain such an impressive result with a system that

is arguably unmatched in terms of throughput and economic cost is, alone, an argument that gives new

weight to this (and future) reductionist approaches.

Finally, to push the system to its limit, we asked *coccinella* to find qualitative differences not in
pharmacologically induced changes in activity, but in a type of spontaneous behaviour mostly
characterised by lack of movement: sleep. In particular, we wondered whether *coccinella* could provide
biological insights comparing conditions of sleep rebound observed after different regimes of sleep
deprivation. *Drosophila melanogaster* is known to show a strong, conserved homeostatic regulation of
sleep that forces flies to recover at least in part lost sleep, for instance after a night of forceful sleep
deprivation[23,24]. We previously showed that the extent of sleep rebound observed after sleep deprivation
only loosely correlates with the amount of lost sleep[25] and it remains an open question whether similar
amounts of sleep rebound may in fact differ from each other in some inscrutable feature that would
underpin a different "sleep depth"[26,27], similarly to what it is believed to happen in mammals. Here, we
analysed a dataset of 727 flies that experienced different regimes of mechanically enforced sleep
deprivation during the 12 hours of the night (Figure 3). Flies were housed in tubes that would rotate
after a set time of inactivity ranging from 20 to 1000 seconds leading to different degrees of sleep
restriction (Figure 3a, dataset from[25]). In this experimental paradigm, all treatments led to a statistically
significant rebound compared to the undisturbed control animals (Figure 3b). We then ran *coccinella* on
the two subsets of the panel: the baseline data, acquired the morning before the sleep deprivation
(Figure 3c), and the rebound data, on the morning after (Figure 3d). Unsurprisingly, we could not detect
any internal biological difference in the pre sleep deprivation control set, featuring flies of identical
genotype and age housed in different tubes before the sleep deprivation treatment. In these conditions,
coccinella could not discern, and performed exactly as a random classifier would (9% vs 9% –
Figure 3c). However, analysis of those same animals during rebound after sleep deprivation showed a
clear clustering, segregating the samples in two subsets with separation around the 300 seconds
inactivity trigger (Figure 3d). This result is important for two reasons: on one hand, it provides, for the
third time, strong evidence that the system is not simply overfitting data of nought biological
significance, given that it could not perform any better than a random classifier on the baseline control.
On the other hand, *coccinella* could find biologically relevant differences on rebound data after
different regimes of sleep deprivation. Interestingly enough, the 300 seconds threshold that *coccinella*
independently identified has a deep intrinsic significance for the field, for it is considered to be the
threshold beyond which flies lose arousal response to external stimuli, defining a "sleep quantum" (*i.e.:*
the minimum amount of time required for transforming inactivity bouts into sleep bouts[23,24,28]).
Coccinella's analysis ran agnostic of the arbitrary 5-minutes threshold and yet identified the same value

as the one able to segregate the two clusters, thus providing an independent confirmation of the five-

minutes rule in *D. melanogaster*.

Discussion

System neuroscience is living a period of renaissance, and *Drosophila* is driving this revolution strong

of the first full-brain connectome, a plethora of new genetic reagents that allow thermo- and opto-

genetic manipulations, a galore of genetic transformants for circuit tracking and manipulation, and

multiple tools for large-scale quantification of behaviour. Progresses in machine learning and computer

power have had a massive impact on the field of ethomics, especially in achieving levels of anatomical

tracking that allow mapping of the tiniest movements on an experimental animal model with the

highest temporal resolution and with little human supervision[6,17]. Most of these systems, however, rely

on relatively expensive setups and do not scale easily to high-throughput experimental paradigms. They

are ideal – and irreplaceable – to identify behavioural patterns and study fine motor control but may be

undue for other uses. Here we introduce a new framework, *coccinella*, that merges an open-source,

economically accessible hardware platform (ethoscopes[14,29]) with a powerful toolbox for statistical

analysis and clustering (HCTSA[15] / Catch22[16]). *Coccinella* is a reductionist tool, not meant to replace

the behavioural categorization that other tools can offer but to complement it. It relies on raspberry PIs

as main acquisition devices, with associated advantages and limitations. Ethoscopes are inexpensive

and versatile but are limited in terms of computing power and acquisition rates. Their online acquisition

speed is fast enough to successfully capture the motor activity of different species of Drosophilae[28], but

may not be sufficient for other animals moving more swiftly, such as zebrafish larvae. Moreover,

coccinella cannot – and is not meant to – apply labels to behaviour ("courting", "lounging", "sipping",

"jumping" *etc.*) but it can successfully identify large behavioural phenotypes and generate unbiased

hypothesis on how behaviour, and a nervous system at large, can be influenced by chemicals, genetics,

artificial manipulations in general. Here, we provided evidence that *coccinella* can be used to

successfully explore and compartmentalise the pharmacobehavioural space and also showed that a

reductionist approach can be employed to discern otherwise invisible shades of a very subtle naturally

occurring behavior: sleep. The success of *Drosophila* as experimental model was built on the many

genetic screens of the 1900s. We propose *coccinella* as an accessible, pivotal tool to boost again this

important line of work in any laboratory, without funding or access to technology being a
discriminative factor.

Acknowledgements

We thank the Gilestro lab at Imperial College London and Robert Lind at Syngenta for useful
discussions. HJ was supported by a BBSRC/CASE studentship in partnership with Syngenta (project
reference BB/M011178/1/1958700). Stocks obtained from the Bloomington Drosophila Stock Center
(NIH P40OD018537) were used in this study.

Methods

Drosophila **rearing**

Fly lines were maintained on a 12-hour light: 12-hour dark (LD) cycle and raised on polenta and yeast-
based fly media (agar 96 g, polenta 240 g, fructose 960 g and Brewer's yeast 1,200 g in 12 litres of
water). Canton-Special (CS) *Drosophila melanogaster* were used as the wild-type line for all
experiments.

Drug resistant mutants

Rdl^{A301S} is derived from Rdl^{MDRR} (RRID:BDSC_1675), an *Rdl* allele isolated from a natural population in
Maryland[30], and underwent isogenization and selection on dieldrin to eliminate the metabolic resistance
and maintain the dieldrin target site resistance[31]. The Rdl^{MDRR} was obtained from the Bloomington
Drosophila Stock Center. For *para:* the L1029F mutation, located in the voltage gated sodium channel
paralytic, has been extensively reported to confer resistance to DDT and pyrethroids in many other
insect species (called kdr, knockdown resistance, reviewed in[32]). In the *Drosophila* gene, kdr maps to
L1029F and is equivalent to the often cited L1014F in other insects (e.g. *Musca domestica*[33]). The kdr
L1029F mutation in *Drosophila* Para was introduced via CRISPR/Cas9-mediated genome editing (see
below). This genome edited generated mutation resulted in a similar resistance to DDT as previously
reported[34].

Generation of Para^{L1029F} via CRISPR-CAS9 based genome editing

CRISPR/Cas9-mediated genome editing was used to introduce a point mutation L1029F in para-PBG isoform, CTT to TTT, L to F by homology-dependent repair (HDR) using one guide RNA and a dsDNA plasmid donor. The strategy design, molecular biology and screening was completed by Well Genetics Inc., Taiwan (R.O.C.). The cassette PBacDsRed contains Piggy Bac 3' terminal repeats, the selection marker 3xP3-DsRed, and Piggy Bac 5' terminal repeats. The selection marker 3xP3-DsRed contains Piggy Bac 3' terminal repeats, 3x Pax3 and hsp70 promoter, DsRed2, SV40 3'UTR, and Piggy Bac 5' terminal repeats. The dsRed marker facilitates the genetic screening and was excised by Piggy Bac transposase. Only one TTAA motif was left after transposition embedded in mutated intron sequence, and create a mutation G to A on X:16,486,649; X:16,486,649~ X:16,486,646, CTAA to T TAA in intron. The CRISPR Target Site [PAM]: CACAAGATTGCCGATGACAA[CGG]. Guide RNA Primers: Sense oligo5'-CTTCGCACAAGATTGCCGATGACAA and Antisense oligo5' - AAACTTGTCATCGGCAATCTTGTGC. Upstream Homology Arm: 1,027bp,the +34,097nt to +35,123 nt from ATG of para. Forward Oligo5'-GTTCACCAAACTCGGAATCG; Reverse Oligo5'- GTGGCCAAGAAGAAGGGAAT. Downstream Homology Arm: 1,022 bp, the +35,128nt to +3,6149 nt from ATG of para Forward Oligo: 5'-CCATGGCTTTAAGCATCGCA; Reverse Oligo: 5'- TTATGACGGATACGGTTACGG. Synthesis fragment: 5'– GGTTGTCATCGGCAATTTTGTGgtgagtactcttatcgaactgctgacttgtaaacgatgtttactggctataatgctgacttatcgcct

The *Drosophila* injection strain was *white*[1118]. 206 embryos were injected. 36 G0 crosses were established. Of the 78 positive lines crosses, in the F1 screen 25 positive lines were identified. Seven lines were positively validated by PCR and 1 line was sequenced confirming no unexpected changes in *para*. Lines were isogenized and balanced. DsRed was excised using PiggyBac (PBac) Transposase Bloomington Stock RRID:BDSC_8285. Excision was validated by genomic PCR and sequencing. Resulting lines were hemizygous viable. The line used in study had internal identifier: 20256ex1.

Choice, handling and preparation of drugs

The initial preliminary analysis was conducted using a group of 12 compounds "proof of principle" compounds and a solvent control. These compounds were initially used to compare both the video method and ethoscope method. After testing these initial compounds, it was found that the ethoscope methodology was more successful, and then the compound list was expanded to 17 (including the control) only using the ethoscope method. As a final test, we included additional compounds for a single concentration, bringing up the total to 40 (including control), also for the ethoscope method. All

insecticide compounds were supplied by Syngenta Ltd. from their in-house stock (see Supplementary

Table 1 for a full list of compounds used). Compounds were received in solid form and diluted in

solvent containing 5% ethanol (VWR, 20821), 5% acetone (Sigma,179124) and 10% dimethylsulfoxide

(DMSO) (D2650, Sigma) in distilled water to 1000ppm initially, then further diluted in the solvent

mixture to 100ppm and 1ppm (where 1mL/L = 1000ppm). For insecticide assays, 0.5mL of 5% sucrose

(Sigma, S0389), 1% agarose (Sigma, A6236) solution was pipetted into each well and allowed to set.

Following this, 2µL of compound solution were placed on the surface and allowed to dry for 30

minutes or more. Male flies were then placed on the surface with a small glass cover slip placed on top

(13mm circular cover slip, VWR631-0150). Flies were briefly anaesthetised (>1 minute) before being

placed onto the surface of the plate. Once each well had been filled with a single male fly, arenas were

placed into the ethoscope and recorded for a minimum of two days. All experiments were started

between ZT0 and ZT1 and within 30 minutes of the flies being placed in the wells. For each compound

in Figures 1c and Figure 1 - Figure Supplement 1c, three repeats were done at different time points. For

Figure 1D, two repeats to three biological repeats per compound.

Data acquisition and processing

301   Ethoscope data were first processed in R using rethomics[29]. Each time series was exported (.csv) and

302   converted using Python to individual time series in a file format (.dat) compatible with MatLab. A

303   metadata (.txt) file served as a reference file of each individual time series with keywords outlining

304   compound groups and concentrations for processing data using HCTSA.

HCTSA / Catch22

Following this process, HCTSA feature extraction was performed on the time-series data (for figures

1c, 1d, 3c, 3d, Figure 1 – Figure Supplements 1 and 2a). After the features were extracted, outputs of

error-producing operations were removed through a normalization process using a sigmoidal

transformation. HCTSA inbuilt functions were then used to classify data using a linear SVM classifier

and a confusion matrix comparing the time series was generated. For some of the time-series data (that

in Figure 2B, 2D, Figure 1 - Figure Supplement 2b), a smaller feature set of HCTSA, Catch-22 was

used for feature extraction. Due to the smaller number of features used with this method, normalization

was not required before using a linear SVM classifier to generate a confusion matrix comparing the

results. A time series of 12 hours was used for the HCTSA analysis in Figure 1 and Figure 1 - Figure

Supplement 1,2. A length of 3 hours was used for the HCTSA analysis in Figure 3. All video data in

Figure 2 are from time series of 15 minutes. By always running the full set of features on aggregate to
train a classifier (e.g., TS_Classify in HCTSA), no post-hoc correction is necessary because the trained
classifier only ever makes a single prediction (only one test is performed).

Video generation for flies on insecticides

Custom 3D-printed squares were designed using the online CAD software Onshape and printed using
Ultimaker 2+ 3D printers using PLA plastic. For insecticide assays, 0.5mL of 5% sucrose (Sigma,
S0389), 1% agarose (Sigma, A6236) solution was pipetted into each well and allowed to set. Following
this, 2µL of compound solution were placed on the surface and allowed to dry for 30 minutes or more
before male flies were placed on the surface with a small glass cover slip placed on top (13mm circular
cover slip, VWR631-0150). Flies were briefly anaesthetised (>1 minute) before being placed onto the
surface of the square. Once each well had been filled with a single male fly, squares were placed in the
arena and a video was recorded for a minimum of 12 hours using an ELP 8 megapixel camera with an
IMX179 Sensor and 2.8-12 mm variable focus manual lens. All video recordings were started between
ZT0 and ZT1. Recordings of flies exposed to compounds were done in a randomised manner. Video
data was then broken down into shorter segments of 15-minute videos for processing. The first 15-
minutes following 1 hour of fly exposure to compound or control was used for pose extraction.

Deeplabcut / B-SoID

The use of DeepLabCut (version 2.1) followed the detailed protocol outlined by[5]. Briefly, frames for
labelling were extracted from 3 representative videos using a K-means algorithm and frames were
labelled with 22 unique body parts (head, left eye, right eye, thorax top, thorax bottom, abdomen top,
abdomen middle, abdomen bottom, left wing tip, right wing tip, left foreleg tip, left foreleg middle,
right foreleg tip, right foreleg middle, left middle leg tip, left middle leg middle, right middle leg tip,
right middle leg middle, left back leg tip, left back leg middle, right back leg tip, right back leg middle).
These frames were labelled locally with a DeepLabCut graphical user interface (GUI) before the
project file was uploaded to Google Drive for training and video analysis to be done using Google
Colab. The data was split into a 9:1 test:train dataset and training was run for more than 150,000
iterations before the average Euclidean error was computed between labels and predictions. The model
at the best performing checkpoint was used to predict pose in novel videos. Following this, B-SOID
was used to de-structure behaviour using the output of DeepLabCut and generate fly-specific time-
series of behavioural grammar as in[4].

Sleep deprivation

Maximal velocity time series data generated from recording flies exposed to either control conditions (no SD) or incrementally increasing immobility-triggered SD conditions were taken from the dataset generated by Geissmann *et al*[25]. Only data for female flies were taken, and where groups consisted of more than 60 individuals, a sample of 60 random individual flies were chosen to take time series data from. Time series data were processed in the same way as described in section 4.2 using the raw ethoscope data.

Survival

Flies were fed with the specified compounds at the desired concentrations and concomitantly analysed in ethoscopes for 24 hours. Time of death was calculated as the last moment of detected motion.

Code and data availability

A notebook version of the source code used to generate all figures is available on the Zenodo public repository, along with all the metadata and the raw data collected in this study (DOIs: 10.5281/zenodo.7335575 and 10.5281/zenodo.7393689). Data were analysed using rethomics[35] and ethoscopy[36]. Two notebooks showing how to use the system for multiple uses are provided as supplementary material.

362 **References**

1. Brown, A. E. X. & Bivort, B. de. Ethology as a physical science. *Nat. Phys.* 1 (2018) doi:10.1038/s41567-018-0093-0.

2. Datta, S. R., Anderson, D. J., Branson, K., Perona, P. & Leifer, A. Computational Neuroethology: A Call to Action. *Neuron* **104**, 11–24 (2019).

3. Wiltschko, A. B. *et al.* Mapping Sub-Second Structure in Mouse Behavior. *Neuron* **88**, 1121–1135 (2015).

4. Hsu, A. I. & Yttri, E. A. B-SOiD, an open-source unsupervised algorithm for identification and fast prediction of behaviors. *Nat. Commun.* **12**, 5188 (2021).

5. Nath, T. *et al.* Using DeepLabCut for 3D markerless pose estimation across species and behaviors. *Nat. Protoc.* **14**, 2152–2176 (2019).

6. Pereira, T. D. *et al.* Fast animal pose estimation using deep neural networks. *Nat. Methods* **16**, 117–125 (2019).

7. Gosztolai, A. *et al.* LiftPose3D, a deep learning-based approach for transforming two-dimensional to three-dimensional poses in laboratory animals. *Nat. Methods* **18**, 975–981 (2021).

8. Wiltschko, A. B. *et al.* Revealing the structure of pharmacobehavioral space through motion sequencing. *Nat. Neurosci.* **23**, 1433–1443 (2020).

9. McDermott-Rouse, A. *et al.* Behavioral fingerprints predict insecticide and anthelmintic mode of action. *Mol. Syst. Biol.* **17**, e10267 (2021).

10. Ayroles, J. F. *et al.* Behavioral idiosyncrasy reveals genetic control of phenotypic variability. *Proc. Natl. Acad. Sci.* **112**, 6706–6711 (2015).

11. Branson, K., Robie, A. A., Bender, J., Perona, P. & Dickinson, M. H. High-throughput ethomics in large groups of Drosophila. *Nat. Methods* **6**, 451–457 (2009).

12. Kabra, M., Robie, A. A., Rivera-Alba, M., Branson, S. & Branson, K. JAABA: interactive machine learning for automatic annotation of animal behavior. *Nat. Methods* **10**, 64–67 (2013).

13. Alisch, T., Crall, J. D., Kao, A. B., Zucker, D. & de Bivort, B. L. MAPLE (modular automated platform for large-scale experiments), a robot for integrated organism-handling and phenotyping. *eLife* **7**, e37166 (2018).

14. Geissmann, Q. *et al.* Ethoscopes: An Open Platform For High-Throughput Ethomics. *PLOS Biol.* **15**, e2003026 (2017).

15. Fulcher, B. D. & Jones, N. S. hctsa: A Computational Framework for Automated Time-Series Phenotyping Using Massive Feature Extraction. *Cell Syst.* **5**, 527-531.e3 (2017).

16. Lubba, C. H. *et al.* catch22: CAnonical Time-series CHaracteristics. *Data Min. Knowl. Discov.* **33**, 1821–1852 (2019).

17. Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat. Neurosci.* **21**, 1281–1289 (2018).

18. Simon, J. C. & Dickinson, M. H. A New Chamber for Studying the Behavior of Drosophila. *PLOS ONE* **5**, e8793 (2010).

19. Geissmann, Q. High-throughput recording, analysis and manipulation of sleep in drosophila. (2018) doi:10.25560/69514.

20. Chatterjee, S., Dey, D. & Munshi, S. Chapter 4 - Feature selection and classification. in *Recent Trends in Computer-Aided Diagnostic Systems for Skin Diseases* (eds. Chatterjee, S., Dey, D. & Munshi, S.) 95–135 (Academic Press, 2022). doi:10.1016/B978-0-323-91211-2.00001-9.

21. Kaduskar, B. *et al.* Reversing insecticide resistance with allelic-drive in Drosophila melanogaster. *Nat. Commun.* **13**, 291 (2022).

22. Remnant, E. J. *et al.* The role of Rdl in resistance to phenylpyrazoles in Drosophila melanogaster. *Insect Biochem. Mol. Biol.* **54**, 11–21 (2014).

23. Shaw, P. J., Cirelli, C., Greenspan, R. J. & Tononi, G. Correlates of Sleep and Waking in Drosophila melanogaster. *Science* **287**, 1834–1837 (2000).

24. Hendricks, J. C. *et al.* Rest in Drosophila Is a Sleep-like State. *Neuron* **25**, 129–138 (2000).

25. Geissmann, Q., Beckwith, E. J. & Gilestro, G. F. Most sleep does not serve a vital function: Evidence from Drosophila melanogaster. *Sci. Adv.* **5**, eaau9253 (2019).

26. French, A. S., Geissmann, Q., Beckwith, E. J. & Gilestro, G. F. Sensory processing during sleep in Drosophila melanogaster. *Nature* (tbc) doi:10.1038/s41586-021-03954-w.

27. Wiggin, T. D. *et al.* Covert sleep-related biological processes are revealed by probabilistic analysis in Drosophila. *Proc. Natl. Acad. Sci.* **117**, 10024–10034 (2020).

28. Joyce, M. *et al.* Divergent evolution of sleep functions. 2023.05.27.541573 Preprint at https://doi.org/10.1101/2023.05.27.541573 (2023).

29. Geissmann, Q., Rodriguez, L. G., Beckwith, E. J. & Gilestro, G. F. Rethomics: an R framework to analyse high-throughput behavioural data. *bioRxiv* 305664 (2018) doi:10.1101/305664.

30. Ffrench-Constant, R. H., Roush, R. T., Mortlock, D. & Dively, G. P. Isolation of Dieldrin Resistance from Field Populations of Drosophila melanogaster (Diptera: Drosophilidae). *J. Econ. Entomol.* **83**, 1733–1737 (1990).

31. Blythe, J. *et al.* The mode of action of isocycloseram: A novel isoxazoline insecticide. *Pestic. Biochem. Physiol.* **187**, 105217 (2022).

32. Arena, J. P., Liu, K. K., Paress, P. S., Schaeffer, J. M. & Cully, D. F. Expression of a glutamate-activated chloride current in Xenopus oocytes injected with Caenorhabditis elegans RNA: evidence for modulation by avermectin. *Mol. Brain Res.* **15**, 339–348 (1992).

33. Dong, K. Insect sodium channels and insecticide resistance. *Invert. Neurosci.* **7**, 17 (2007).

34. Samantsidis, G.-R. *et al.* 'What I cannot create, I do not understand': functionally validated synergism of metabolic and target site insecticide resistance. *Proc. R. Soc. B Biol. Sci.* **287**, 20200838 (2020).

35. Geissmann, Q., Rodriguez, L. G., Beckwith, E. J. & Gilestro, G. F. Rethomics: An R framework to analyse high-throughput behavioural data. *PLOS ONE* **14**, e0209331 (2019).

36. Blackhurst, L. & Gilestro, G. F. ethoscopy & ethoscope-lab: a framework for behavioural analysis to lower entrance barrier and aid reproducibility. 2022.11.28.517675 Preprint at https://doi.org/10.1101/2022.11.28.517675 (2022).

Figure 1 – *Coccinella* successfully classifies pharmacobehaviours. a) Bespoke 3D printed arena can house 24 individual flies in a two-dimensional circular space. Each arena measures 11.5 mm in diameter and allows for back illumination either by visible or infrared light sources embedded in the ethoscope base. The red ellipse shows the data being extracted by the ethoscope in real time. w, *h*: width and height of the ellipse inscribing the animal. φ: the angle of the ellipse in reference to the region of interest. *max. vel.*: maximal velocity over the last 10 seconds. **b)** Flowchart describing the analytical pipeline and the tools that compose *coccinella*. **c)** Confusion matrices for treatments with 16 compounds and a solvent control, with drugs used at concentrations of 1000ppm (left), 100ppm (center), 1ppm (right). The target icon indicates calculated accuracy while the rolling die indicate the accuracy of a random classifier. **d)** Confusion matrix for the largest panel of 40 treatments at 100ppm.

Figure 2 – Comparison between *coccinella* and the state-of-the-art. a) Experimental pipeline illustrating the four experimental analysis. **b)** Confusion matrix for 12 treatments (11 drugs at 1000ppm and one solvent control) analysed using *coccinella*. **c)** Same experimental treatments as in b, analysed using the DeepLabCut → B-SoID → Random forest pipeline starting from high resolution images. The Random forest classifier was trained on a 4:1 training:testing ratio. **d)** Same as c but with Catch22 identification and SVM clustering after B-SoID grammar dissection. This is a hybrid treatment combining HCTSA feature extraction to the high resolution pipeline. **e)** Same as c but using K-nearest neighbours (KNN) as cluster algorithm. KNN required a much higher training:testing ratio of 9:1, dramatically reducing the size of the testing dataset. The accuracy of a random classifier for all matrices on this figure would be 8.3% (not shown on figures for lack of space).

Figure 3 – *Coccinella* finds differences in type of sleep rebound after sleep deprivation. a) Sleep profile of flies over the period of three days. A 12 hours sleep deprivation regime starts at the beginning of the dark phase of day 0 (purple bar). The three-hours windows labelled with green boxes were analysed by *coccinella* in search of meaningful differences. The letters above refer to the panels using data in those time windows **b)** Extent of rebound as observed following sleep deprivation as performed in a. Panel a and b reproduce data from[25]. **c)** Confusion matrix showing the classification using *coccinella* of the baseline time-series. No accuracy gain compared to the random classifier. **d)** Confusion matrix of the rebound data. The classification finds two clusters, separated by the 300 seconds threshold (thick black lines).

Figure 1 - Figure Supplement 1. a) Explanatory drawing of how confusion matrices present the data. The blue diagonal boxes indicate the number of flies correctly classified (true positives). The vertical cells indicate the number of flies wrongly classified (false positives). The horizontal cells indicate the number of flies that should have been classified (false negatives). **b)** Survival curves upon treatment with a panel of 12 drugs administered at different concentrations, indicated above each figure (1ppm, 100ppm, 1000ppm). **c)** Confusion matrices for treatments with 11 compounds and a solvent control, with drugs used at concentrations of 1000ppm (left), 100ppm (middle), 1ppm (right). The target icon indicates calculated accuracy. Accuracy of a random classifier would be 8.3%.

Figure 1 - Figure Supplement 2. a) Confusion matrix classifying the action of two selected drugs (DDT 100ppm and Dieldrin 100ppm) or solvent on wild type flies (WT) or mutants with known resistance to those drugs. The LD100 of Dieldrin is 0.1-1ppm and the LD100 of DDT is 10-100ppm. The three purple dotted boxes highlight the three experimental clusters (wild-type flies, *para* mutants, *Rdl* mutants)

Supplementary Table 1. List of all the compounds used in this study with bibliographic reference.

Supplementary Notebooks. Two Jupyter notebooks guiding the user through the integration of ethoscope data with HCTSA (notebook 1) and Catch22 (notebook2).

Figure 1

Figure 2

Figure 3

Figure 1 - Figure Supplement 1

Figure 1 - Figure Supplement 2

Compound	Mode of action	Ref	Figure
Syngenta Cpd 3	Mitochondrial ATP synthesis		1D
DDT	Voltage-gated sodium channel activator	1	1C
Juglone	ROS	2,3	1D
Diflubenzuron	Chitin synthesis	4,5	1D
Rotenone	Complex I of the mitochondrial respiratory chain (MRC)	6	1D
Methotrexate	Dihydrofolate reductase inhibitor	7	1D
Diafenthiuron	Mitochondrial ATPase	8	1D
Fenoxycarb	Juvenile hormone mimic	9	1D
DCCD	Mitochondrial FIFO ATP synthase	10	1D
Lufenuron	Chitin synthesis inhibitor	11	1C
Chlorfenapyr	Mitochondrial oxidative phosphorylation	12	1D
Thiamethoxam	NACh-R13	13	1D
Tebfenozide	Ecdysone receptor	14	1D
Flonicamid	Sodium channels/TRP-V channels	15	1C
Syngenta Cpd 2	NACh-R16	16	1D
Quinuclidine Hycl.	Muscarinic Ach-R	17	1D
Indoxacarb	Voltage-gated sodium channels	18	1D
Fluazinam	Mitochondrial ATP synthesis	19,20	1D
Syngenta Cpd 5	Unknown		1D
Syngenta Cpd 7	Octopamine receptor		1D
Methoprene	Juvenile hormone mimic	21,22	1D
Permethrin	Voltage-gated sodium channels	23	1C
Tubulin	Unknown		1D
Metaflumizone	Voltage-gated sodium channels	24	1D
Syngenta Cpd 12	Unknown		1D
Cyenopyrafen	Succinate dehydrogenase	25,26	1D
Syngenta Cpd 1	GABAA receptors		1C
Syngenta Cpd 10	Octopamine receptor	27	1D
Abamectin	Glutamate-gated chloride channels	28,29	1C,1D,2
Diamide Syn Cpd	Ryanodine receptors		1C,1D,2
Chlorantraniliprole	Ryanodine receptors	30	1C,1D,2
Cyantraniliprole	Ryanodine receptors	31	1C,1D,2
Dieldrin	GABAA receptors	32	1C,1D,2

Flubendiamide	Ryanodine receptors	33	1C,1D,2
Imidacloprid	NACh-R	34,35	1C,1D,2
Pymetrozine	TRP-V channels	36	1C,1D,2
Ryanodine	Ryanodine receptors	37	1C,1D,2
Spiroindoline	Cholinergic transmission	38	1C,1D,2
Tetramethrin	Sodium channel modulator	39	1C,1D,2

Bibliography for references listed in supplementary table 1 only (see column "Ref").

1.   O'Brien, R. D. & Matsumura, F. DDT: A New Hypothesis of Its Mode of Action. Science 146, 657–658 (1964).
2.   Islam, A. K. M. M. & Widhalm, J. R. Agricultural Uses of Juglone: Opportunities and Challenges. Agronomy 10, 1500 (2020).
3.   Maleita, C. et al. Juglone and 1,4-Naphthoquinone—Promising Nematicides for Sustainable Control of the Root Knot Nematode Meloidogyne luci. Front. Plant Sci. 13, 867803 (2022).
4.   Cohen, E. & Casida, J. E. Inhibition of Tribolium gut chitin synthetase. Pestic. Biochem. Physiol. 13, 129–136 (1980).
5.   Meola, S. M. & Mayer, R. T. Inhibition of Cellular Proliferation of Imaginal Epidermal Cells by Diflubenzuron in Pupae of the Stable Fly. Science 207, 985–987 (1980).
6.   Palmer, G., Horgan, D. J., Tisdale, H., Singer, T. P. & Beinert, H. Studies on the Respiratory Chain-linked Reduced Nicotinamide Adenine Dinucleotide Dehydrogenase. J. Biol. Chem. 243, 844–847 (1968).
7.   Stone, S. R., Montgomery, J. A. & Morrison, J. F. Inhibition of dihydrofolate reductase from bacterial and vertebrate sources by folate, aminopterin, methotrexate and their 5-deaza analogues. Biochem. Pharmacol. 33, 175–179 (1984).
8.   Ruder, F. J. & Kayser, H. The Carbodiimide Product of Diafenthiuron Inhibits Mitochondria in Vivo. Pestic. Biochem. Physiol. 46, 96–106 (1993).
9.   Grenier, S. & Grenier, A.-M. Fenoxycarb, a fairly new Insect Growth Regulato: review of its effects on insects. Ann. Appl. Biol. 122, 369–403 (1993).
10.   Ruder, F. & Kayser, H. The thiourea insecticide diafenthiuron inhibits mitochondrial ATPase in vitro and in vivo by its carbodiimide product. Biochem. Soc. Trans. 22, 241–244 (1994).
11.   Wilson, T. G. & Cryan, J. R. Lufenuron, a chitin-synthesis inhibitor, interrupts development ofDrosophila melanogaster. J. Exp. Zool. 278, 37–44 (1997).
12.   Black, B. C., Hollingworth, R. M., Ahammadsahib, K. I., Kukel, C. D. & Donovan, S. Insecticidal Action and Mitochondrial Uncoupling Activity of AC-303,630 and Related Halogenated Pyrroles. Pestic. Biochem. Physiol. 50, 115–128 (1994).
13.   Wiesner, P. & Kayser, H. Characterization of nicotinic acetylcholine receptors from the insectsAphis craccivora,Myzus persicae, andLocusta migratoria by radioligand binding assays: Relation to thiamethoxam action. J. Biochem. Mol. Toxicol. 14, 221–230 (2000).
14.   Retnakaran, A. et al. Mode of action of the ecdysone agonist tebufenozide (RH-5992), and an exclusion mechanism to explain resistance to it. Pest Manag. Sci. 57, 951–957 (2001).
15.   Taylor-Wells, J. et al. Toxicity, mode of action, and synergist potential of flonicamid against mosquitoes. Pestic. Biochem. Physiol. 151, 3–9 (2018).
16.   Chambers, C. et al. Insecticidal spider toxins are high affinity positive allosteric modulators of the nicotinic acetylcholine receptor. FEBS Lett. 593, 1336–1350 (2019).

17.   Dick, M. R., Dripps, J. E. & Orr, N. Muscarinic Agonists as Insecticides and Acaricides. Pestic. Sci. 49, 268–276 (1997).

18.   Lapied, B., Grolleau, F. & Sattelle, D. B. Indoxacarb, an oxadiazine insecticide, blocks insect neuronal sodium channels: DCJW blocks neuronal Na + channels. Br. J. Pharmacol. 132, 587–595 (2001).

19.   Lee, J. E. et al. Fluazinam targets mitochondrial complex I to induce reactive oxygen species-dependent cytotoxicity in SH-SY5Y cells. Neurochem. Int. 60, 773–781 (2012).

20.   Guo, Z., Miyoshi, H., Komyoji, T., Haga, T. & Fujita, T. Uncoupling activity of a newly developed fungicide, fluazinam [3-chloro-N-(3-chloro-2,6-dinitro-4-trifluoromethylphenyl)-5-trifluoromethyl-2-pyridinamine]. Biochim. Biophys. Acta BBA - Bioenerg. 1056, 89–92 (1991).

21.   Miura, K., Oda, M., Makita, S. & Chinzei, Y. Characterization of the Drosophila Methoprene -tolerant gene product: Juvenile hormone binding and ligand-dependent gene regulation. FEBS J. 272, 1169–1178 (2005).

22.   Jindra, M., Palli, S. R. & Riddiford, L. M. The Juvenile Hormone Signaling Pathway in Insect Development. Annu. Rev. Entomol. 58, 181–204 (2013).

23.   Vais, H., Williamson, M. S., Devonshire, A. L. & Usherwood, P. N. The molecular interactions of pyrethroid insecticides with insect and mammalian sodium channels. Pest Manag. Sci. 57, 877–888 (2001).

24.   Salgado, V. L. & Hayashi, J. H. Metaflumizone is a novel sodium channel blocker insecticide. Vet. Parasitol. 150, 182–189 (2007).

25.   Nakahira, K. Strategy for discovery of a novel miticide Cyenopyrafen which is one of electron transport chain inhibitors. J. Pestic. Sci. 36, 511–515 (2011).

26.   Wei, P. et al. Downregulation of carboxylesterase contributes to cyflumetofen resistance in Tetranychus cinnabarinus (Boisduval): Downregulated esterase genes in cyflumetofen resistance in mites. Pest Manag. Sci. 75, 2166–2173 (2019).

27.   O'Sullivan, A. C. et al. Synthesis and Insecticidal Activity of New Benzyl- and Indanyl-Oxazolines, Thiazolines and Alkoxy-Alkyl-Imidazolines. in ACS Symposium Series (eds. Maienfisch, P. & Stevenson, T. M.) vol. 1204 411–430 (American Chemical Society, 2015).

28.   Cully, D. F., Paress, P. S., Liu, K. K., Schaeffer, J. M. & Arena, J. P. Identification of a Drosophila melanogaster Glutamate-gated Chloride Channel Sensitive to the Antiparasitic Agent Avermectin. J. Biol. Chem. 271, 20187–20191 (1996).

29.   Arena, J. P., Liu, K. K., Paress, P. S., Schaeffer, J. M. & Cully, D. F. Expression of a glutamate-activated chloride current in Xenopus oocytes injected with Caenorhabditis elegans RNA: evidence for modulation by avermectin. Mol. Brain Res. 15, 339–348 (1992).

30.   Cordova, D. et al. Anthranilic diamides: A new class of insecticides with a novel mode of action, ryanodine receptor activation. Pestic. Biochem. Physiol. 84, 196–214 (2006).

31.   Selby, T. P. et al. Discovery of cyantraniliprole, a potent and selective anthranilic diamide ryanodine receptor activator with cross-spectrum insecticidal activity. Bioorg. Med. Chem. Lett. 23, 6341–6345 (2013).

32.   ffrench-Constant, R. H., Rocheleau, T. A., Steichen, J. C. & Chalmers, A. E. A point mutation in a Drosophila GABA receptor confers insecticide resistance. Nature 363, 449–451 (1993).

33.   Masaki, T. et al. Flubendiamide, a Novel Ca 2+ Channel Modulator, Reveals Evidence for Functional Cooperation between Ca 2+ Pumps and Ca 2+ Release. Mol. Pharmacol. 69, 1733–1739 (2006).

34.   Bodereau-Dubois, B. et al. Transmembrane Potential Polarization, Calcium Influx, and Receptor Conformational State Modulate the Sensitivity of the Imidacloprid-Insensitive Neuronal Insect Nicotinic Acetylcholine Receptor to Neonicotinoid Insecticides. J. Pharmacol. Exp. Ther. 341, 326–339 (2012).

35.   Perry, T. et al. Effects of mutations in Drosophila nicotinic acetylcholine receptor subunits on sensitivity to insecticides targeting nicotinic acetylcholine receptors. Pestic. Biochem. Physiol. 102, 56–60 (2012).

36.   Nesterov, A. et al. TRP Channels in Insect Stretch Receptors as Insecticide Targets. Neuron 86, 665–671 (2015).

37.   Fleischer, S., Ogunbunmi, E. M., Dixon, M. C. & Fleer, E. A. Localization of Ca2+ release channels with ryanodine in junctional terminal cisternae of sarcoplasmic reticulum of fast skeletal muscle. Proc. Natl. Acad. Sci. 82, 7256–7259 (1985).

38.   Sluder, A. et al. Spiroindolines Identify the Vesicular Acetylcholine Transporter as a Novel Target for Insecticide Action. PLoS ONE 7, e34712 (2012).

39.   Yamamoto, D., Quandt, F. N. & Narahashi, T. Modification of single sodium channels by the insecticide tetramethrin. Brain Res. 274, 344–349 (1983).