The Kelch13 compartment is a hub of highly divergent vesicle trafficking proteins in malaria parasites

Sabine Schmidt[1,§], Jan Stephan Wichers-Misterek[1,§], Hannah Michaela Behrens[1], Jakob Birnbaum[1], Isabelle G. Henshall[1], Jana Dröge[1], Ernst Jonscher[1], Sven Flemming[1], Carolina Castro-Peña[1], Paolo Mesén-Ramírez[1], Tobias Spielmann[1,#]

[1] Bernhard Nocht Institute for Tropical Medicine, 20359 Hamburg, Germany
§ contributed equally, order determined alphabetically
# Corresponding author: spielmann@bnitm.de

ORCID:
Sabine Schmidt:	0000-0002-6958-9832
Jan Stephan Wichers-Misterek:	0000-0002-0599-1742
Hannah Michaela Behrens:	0000-0002-7406-7086
Jakob Birnbaum:	0000-0002-1419-522X
Isabelle Henshall:	0000-0002-5906-0687
Jana Dröge:	0000-0002-4018-6175
Ernst Jonscher:	0000-0002-8480-6102
Carolina Castro-Peña:	0000-0002-6878-4951
Paolo Mesén-Ramírez	0000-0001-7842-5867
Tobias Spielmann:	0000-0002-3968-4601

ABSTRACT

Single amino acid changes in the parasite protein Kelch13 (K13) result in reduced susceptibility of *P. falciparum* parasites to artemisinin and its derivatives (ART). Recent work indicated that K13 and other proteins co-localising with K13 (K13 compartment proteins) are involved in the endocytic uptake of host cell cytosol (HCCU) and that a reduction in HCCU results in ART resistance. HCCU is critical for parasite survival but is poorly understood, with the K13 compartment proteins among the few proteins so far functionally linked to this process. Here we further defined the composition of the K13 compartment by analysing more hits from a previous BioID, showing that MyoF and MCA2 as well as Kelch13 interaction candidate (KIC) 11 and 12 are found at this site. Functional analyses, tests for ART susceptibility as well as comparisons of structural similarities using AlphaFold2 predictions of these and previously identified proteins showed that canonical vesicle trafficking and endocytosis domains were frequent in proteins involved in resistance or endocytosis (or both), comprising one group of K13 compartment proteins. While this strengthened the link of the K13 compartment to endocytosis, many proteins of this group showed unusual domain combinations and large parasite-specific regions, indicating a high level of taxon-specific adaptation of this process. Another group of K13 compartment proteins did not influence endocytosis or ART susceptibility and lacked detectable vesicle trafficking domains. We here identified the first protein of this group that is important for asexual blood stage development and showed that it likely is involved in invasion. Overall, this work identified novel proteins functioning in endocytosis and at the K13 compartment. Together with comparisons of structural predictions it provides a repertoire of functional domains at the K13 compartment that indicate a high level of adaption of endocytosis in malaria parasites.

INTRODUCTION

Malaria, caused by protozoan parasites of the genus *Plasmodium*, is one of the deadliest infectious diseases, responsible for an estimated 627 000 deaths and 241 million cases in 2020 [1]. Malaria deaths have been declining in the last two decades [1,2], but this trend has reversed in the last few years [1]. One of the factors that contributed to the reduction of cases is efficient treatment which currently relies on artemisinin and its derivatives (ART) administered together with a partner drug [3]. However, parasites with a reduced susceptibility to ART have become widespread in South East Asia [4–12] and have now also emerged in Africa [13–16], Papa New Guinea [17] and South America [18]. The reduced effectivity of ART is linked to a delayed parasite clearance in patients that is due to a decreased susceptibility of ring-stage parasites to ART [19,20], leading to recrudescence and treatment failure in patients [21–24]. Experimental human infections indicate that a doubled clearance half-life in parasites with reduced ART susceptibility results in a >10 fold increase in viable parasite half-life compared to the sensitive parasites [25]. Decreased susceptibility of ring-stage parasites can be measured *in vitro* using the ring stage survival assay (RSA) [26]. We here refer to this *in vitro* measurable phenomenon as "*in vitro* ART resistance" when the survival of parasites is above 1% in an RSA as previously defined [26]. Point mutations in the gene encoding the parasite protein Kelch13 (K13) are the main cause for the reduced ART susceptibility of laboratory and field parasite isolates [27,28] and was shown to lead to decreased K13 protein levels [29–33]. This results in reduced hemoglobin uptake via endocytosis, which is assumed to cause less ART activation through hemoglobin degradation products, resulting in the reduced parasite ART susceptibility [29,34]. Beside K13 mutations in other genes, such as Coronin [35] UBP1 [29,36–38] or AP2µ [38,39] have also been linked with reduced ART susceptibility. In contrast to K13 which is only needed for endocytosis in ring stages (the stage relevant for in vitro ART resistance), some of these proteins (AP2µ and UBP1) are also needed for endocytosis in later stage parasites [29]. At least in the case of UBP1, this is associated with a higher fitness cost but lower resistance compared to K13 mutations [34,40]. Hence, the stage-specificity of endocytosis functions is relevant for *in vitro* ART resistance: proteins influencing endocytosis in trophozoites are expected to have a high fitness cost whereas proteins not needed for endocytosis in rings would not be expected to influence resistance. Endocytosis is a critical process for blood stage growth of malaria parasites. During its asexual development in erythrocytes the parasite uses this process to take up more than two-thirds of the soluble host cell content, which consists almost exclusively of hemoglobin [41–43]. The endocytosed host cell cytosol is transported to the acidic lysosome-like food vacuole (FV), where digestion of hemoglobin results in generation

of free heme, toxic to the parasite, which is further converted into nontoxic hemozoin
[44,45]. Host cell cytosol uptake (HCCU) provides both space and building blocks
(amino acids from digested hemoglobin) for parasite growth [42]. As a consequence,
ART resistant parasites are hypersensitive to amino acid restriction, highlighting the
importance of endocytosis for parasite growth and its connection to ART resistance
[46,47]. Hemoglobin trafficking to the parasite food vacuole is believed to be initiated
at membrane invaginations called cytostomes [30,42,48,49] followed by vesicular
transport from the parasite plasma membrane (PPM) to the food vacuole [42,50], likely
in an actin-myosin motor dependent manner [48,51–53]. The molecular effectors
involved in this process remain poorly characterized. So far VPS45 [54], Rbsn5 [55],
Rab5b [55], the phosphoinositide-binding protein PX1 [56], the host enzyme
peroxiredoxin 6 [57] and K13 and some of its compartment proteins (Eps15, AP2μ,
KIC7, UBP1) [29] have been reported to act at different steps in the endocytic uptake
pathway of hemoglobin. While inactivation of VPS45, Rbsn5, Rab5b, PX1 or actin
resulted in an accumulation of hemoglobin filled vesicles [53–56], indicative of a block
during endosomal transport (late steps in endocytosis), no such vesicles were
observed upon inactivation of K13 and its compartment proteins [29], suggesting a
role of these proteins during initiation of endocytosis (early steps in endocytosis).
107
In a previous work we used a quantitative BioID approach to identify proteins in close
proximity to K13 and its interactor Eps15 [29]. We designated these proteins as K13
interaction candidates (KICs) [29]. As the proteins identified by proximity labelling
approaches not only include interactors but also proteins that are located in a close
spatial position of the bait [58,59], we will here refer to the collective of these proteins
as the "proxiome". We reasoned that due to the high number of proteins that turned
out to belong to the K13 compartment when validating the top hits of the K13 [29], the
remaining hits of these experiments might contain further proteins belonging to the
K13 compartment. Here we identified and functionally analysed further K13
compartment proteins from the K13 proxiome [29] and classify these and previously
confirmed K13 compartment proteins. Location and functional data as well as
comparisons of alpha fold predicted structural elements revealed that the K13
compartment contains proteins suggestive of a highly divergent endocytosis
mechanism in malaria parasites but also a second group of proteins with other
functions.
123
RESULTS
To identify novel K13 compartment proteins we further exploited previously described
proximity-dependent biotinylation experiments that had used K13 or the K13
compartment protein Eps15 as bait, selecting enriched proteins not characterized in

our previous work [29] (Table S1). In order to do this we excluded proteins that (i) had
previously been analysed, (ii) were either linked with or had been shown to localise to
the inner membrane complex (IMC) or the basal complex (PF3D7_1345600 [60];
PF3D7_0109000 (PhIL1) [61–63]; PF3D7_0717600 (IMC32) [64]; PF3D7_0822900
(PIC2) [65], PF3D7_1018200 (PPP8) [65–67], (iii) are considered typical DiQ-BioID
'contaminants' (PF3D7_0708400 (HSP90) and PF3D7_1247400 (FKBP35) [29,68]),
(iv) localised to the apical polar ring in *P. berghei* (PF3D7_1141300 (APR1) [69]), (v)
localised to the nucleus PF3D7_1247400 (FKBP35) [70,71], (vi) were linked with the
apicoplast (PF3D7_0721100 [72]) or (vi) were also present in BioID experiments using
Clathrin heavy chain (CHC) as bait [29] (PF3D7_0408100). These selection criteria
resulted in a candidate list of thirteen proteins (PF3D7_1438400 (MCA2),
PF3D7_1243400,   PF3D7_1365800,   PF3D7_1447800,   PF3D7_1142100,
PF3D7_0103100   (VPS51),   PF3D7_1329100   (MyoF),   PF3D7_1329500,
PF3D7_0405700 (UIS14), PF3D7_0907200, PF3D7_0204300, PF3D7_1117900 and
PF3D7_1016200), of which ten were chosen for further characterization in this
manuscript (Table 1, Table S1).
144
MyosinF is involved in host cell cytosol uptake and associated with the K13
compartment
The presence of MyosinF (MyoF; PF3D7_1329100 previously also MyoC), in the K13
proxiome could indicate an involvement of actin/myosin in endocytosis in malaria
parasites. We therefore analysed its location in the cell. For this, we first generated
transgenic parasites expressing a C-terminally 2xFKBP-GFP-2xFKBP tagged version
of MyoF expressed from its original locus using selection-linked integration (SLI) [73]
(Figure S1A). Expression and localisation of the fusion protein was analysed by
fluorescent microscopy. The tagged MyoF was detectable as foci close to the food
vacuole from the stage parasites turned from late rings to young trophozoite stage
onwards, while in schizonts multiple MyoF foci were visible (Figure 1A, S2A). This
expression pattern is in agreement with its transcriptional profile [74,75]. A similar
localisation was observed in a cell line expressing MyoF with a smaller tag (MyoF-
2xFKBP-GFP) from its endogenous locus (Figure S1B, Figure S2B). The proximity of
the MyoF foci to the food vacuole in trophozoites was also evident by co-localisation
with an episomally expressed food vacuole marker (P40PXmCherry) [54,76] (Figure
1B). Next, we episomally co-expressed mCherry-K13 in the MyoF-2xFKBP-GFP-
2xFKBP[endo] parasites to assess whether MyoF is found at the K13 compartment
(Figure 1C, S2A). Quantification from live fluorescence images of late ring and
trophozoite stage parasites revealed that 8% of MyoF foci overlapped with K13 foci,
12% showed a partial overlap, 36% were close together (touching but not
overlapping), while 44% of MyoF foci were further away from K13 foci (Figure 1D). As

the GFP-tagging of MyoF appeared to have some effect on the parasite (see below),
we validated the MyoF localisation in respect to the K13 compartment by generating
parasites with an endogenously 3xHA-tagged MyoF (MyoF-3xHA[endo] parasites) using
SLI (Figure S1C) and episomally co-expressed mCherry-K13. IFA confirmed the focal
localisation of MyoF and its spatial association with mCherry-K13 foci with a tendency
for a higher overlap with K13, which might be due to the partial inactivation of the GFP-
tagged MyoF (Figure S2C). We also detected MyoF signal in ring stage parasites
(Figure S2C). As we did not detect the GFP-tagged version in rings, this likely indicates
a low level of expression of MyoF in ring stage parasites as IFA is more sensitive. In
late schizonts and merozoite the MyoF-GFP signal was not associated with K13, but
was present in elongated foci (Figure 1C, S2A) reminiscent of the MyoE signal
previously reported in *P. berghei* schizonts [77]. Taken together these results show
that MyoF is in foci that are frequently close or overlapping with K13, indicating that
MyoF is found in a regular close spatial association with the K13 compartment and at
times overlaps with that compartment.

During routine *in vitro* culturing we noticed that MyoF-2xFKBP-GFP-2xFKBP[endo]
parasites grew poorly and subsequent flow cytometry-based proliferation assays
revealed a mean relative growth of 36.7% compared to 3D7 wild type parasites after
two replication cycles (Figure 1E). These results indicated that C-terminal 2xFKBP-
GFP-2xFKBP tagging of MyoF impaired its function and that this protein has an
important role for the growth of asexual blood stages. We therefore generated MyoF-
2xFKBP-GFP-2xFKBP[endo] parasites episomally expressing a nuclear mislocaliser
(1xNLS-FRB-mCherry) in order to conditionally inactivate it using knock-sideways.
This system allows the conditional mislocalisation of a protein from its site of action
into the nucleus upon addition of rapalog [73,78,79]. Assessment of mislocalisation
efficacy by fluorescent microscopy at timepoints between 1 hour and 22 hours post
induction revealed partial mislocalisation of MyoF-2xFKBP-GFP-2xFKBP to the
nucleus with some MyoF remaining at foci close to the food vacuole (Figure 1F, S2D).
Despite the only partial inactivation of MyoF by knock sideways, flow cytometry-based
proliferation assays revealed a 40.5% reduced growth after two replication cycles upon
addition of rapalog compared to control parasites without rapalog, while such an effect
was not observed for 3D7 wild type parasites upon addition of rapalog (Figure 1G,
Figure S2E). Hence, conditional inactivation of MyoF further reduced growth despite
the fact that the tag on MyoF already led to a substantial growth defect, indicating an
important role for MyoF during asexual blood stage development.

Inspection of the MyoF-2xFKBP-GFP-2xFKBP[endo], compared to 3D7 wild type-
trophozoites, revealed an increased number of vesicles in the parasite cytoplasm

(Figure 1H), resembling the phenotype observed after inactivation of VPS45 or Rbsn5,
two proteins involved in the transport of host cell cytosol to the parasite's food vacuole
[54,55]. This was even more pronounced upon conditional inactivation of MyoF by
knock-sideways (Figure 1H), suggesting this is due to a reduced function of MyoF. Of
note, C-terminal tagging with the small 3xHA epitope had no significant effect on
parasite proliferation (Figure S2F), and accordingly showed only a small, but still
significant increase of the number of vesicles in the parasite cytoplasm (Figure S2G),
indicating only a minor impairment by C-terminal tagging with the small 3xHA epitope.
These findings indicated that inactivation of MyoF might impair the transport of host
cell cytosol to the food vacuole.
216
We therefore directly tested for an effect of MyoF inactivation on HCCU using a
'bloated food vacuole assay' [54]. For this we incubated parasites upon MyoF
inactivation (and control parasites without rapalog) with the protease inhibitor E64 [80].
In the presence of E64, newly internalized hemoglobin cannot be degraded and
accumulates in the food vacuole, resulting in bloated food vacuoles if endocytosis is
operational. In 3D7 wild type parasites 98% of the cells showed a bloated food vacuole
in both rapalog treated and control parasites (Figure S2H-J). In contrast, 88% of the
MyoF-2xFKBP-GFP-2xFKBP^{endo} control (without rapalog) parasites developed
bloated food vacuoles, and after further inactivation of MyoF by knock sideway (with
rapalog) only 52% of the cells showed a bloated food vacuole, indicating reduced
HCCU with increasing inactivation of MyoF (Figure 1I, Figure S2H-J). As the effect of
MyoF inactivation was only partial, we measured the parasite and food vacuole size
to obtain a finer readout for HCCU. This analysis revealed a significantly reduced food
vacuole size in the parasites with MyoF inactivated by knock sideways, while there
was no significant effect on parasite size (Figure 1J). Plotting the values of the
individual parasites showed that the food vacuoles of similarly sized parasites were
consistently smaller in the MyoF knock sideways compared to the parasites without
knock sideways inactivation of MyoF (Figure 1K), indicating that the reduced
hemoglobin delivery to the food vacuole upon inactivation of MyoF was not an indirect
effect due to parasite growth impairment during the assay time. Next, we plotted the
endocytosis phenotypes (vesicles and bloating) and growth of 3D7 (wild type MyoF
activity), MyoF-3xHA (small degree of MyoF inactivation), MyoF-2xFKBP-GFP-
2xFKBP (more MyoF inactivation) and MyoF-2xFKBP-GFP-2xFKBP knock sideways
(even more MyoF inactivation) which showed that the degree of the endocytosis
phenotypes correlated with growth and hence level of MyoF inactivation (Figure 1L).
Overall, this indicates that MyoF function is needed for hemoglobin to reach the food
vacuole.

Finally, we also tested if inactivation of MyoF has an effect on *in vitro* ART resistance.
RSAs [26] revealed no significant increase in parasite survival upon MyoF inactivation
with neither 12 h nor 24 h rapalog pre-incubation to inactivate MyoF by knock sideways
(Figure 1M, Figure S2K). Of note, due to the partial inactivation due to the tagging of
MyoF we would have expected the control (MyoF-2xFKBP-GFP-2xFKBP parasites
without rapalog) to already have elevated RSA survival levels if MyoF impairment
reduces endocytosis in rings, which was not the case (Figure 1M). Similarly, also
incubation with the actin destabilising agent Cytochalasin D [81], had no effect on RSA
survival in 3D7 or K13^{C580Y} [29] parasites, indicating an actin/myosin independent
endocytosis pathway in ring stage parasites (Figure 1M) and speaking against other
myosins taking over the MyoF endocytosis function in rings.
255
256   Overall, our results indicate a close association of MyoF foci with the K13 compartment
257   and a role of MyoF in endocytosis albeit not in rings and at a step in the endocytosis
258   pathway when hemoglobin-filled vesicles had already formed and hence is
259   subsequent to the function of the other so far known K13 compartment proteins.
260

261   KIC11 is a K13 compartment protein important for asexual parasite proliferation
262   due to a function not involving endocytosis
PF3D7_1142100, currently annotated as 'conserved Plasmodium protein, unknown
function', was renamed K̲13 i̲nteracting c̲andidate 11 (KIC11), following the previously
established nomenclature for potential K13 compartment proteins [29]. In order to test
whether KIC11 is a member of the K13 compartment or not, we first generated
transgenic parasites expressing it as a C-terminally 2xFKBP-GFP-2xFKBP tagged
fusion protein from the original genomic locus using SLI (KIC11-2xFKBP-GFP-
2xFKBP^{endo} parasites). Correct genomic modification of the *kic11* locus was verified
by PCR (Figure S1D). The tagged KIC11 showed multiple foci in ring and trophozoite
stage parasites (mean 2.2 per ring stage and 4.0 per trophozoite), while in schizonts
many KIC11-2xFKBP-GFP-2xFKBP foci were visible (mean 14.3), which are not co-
localizing with apical organelle marker proteins (Figure 2A, Figure S3A, B). Next, we
generated KIC11-2xFKBP-GFP-2xFKBP^{endo} parasites episomally co-expressing
mCherry-K13 to assess presence at the K13 compartment. Fluorescence microscopy
revealed more KIC11 than K13 foci per cell in ring and trophozoite stage parasites
(Figure S3B). Quantification showed that in ring stage parasites 16% of K13 foci
overlapped with KIC11 foci, 45% showed a partial overlap, 26% were close together
(touching but not overlapping), while 13% were further away. In trophozoites 23% of
K13 foci overlapped with KIC11 foci, 46% showed a partial overlap, 15% were close
together, while 15% were further away (Figure 2B,C Figure S3C). We conclude that

KIC11 is a protein located at the K13 compartment with an additional protein pool
located at foci elsewhere.

In order to assess the importance of KIC11 for parasite proliferation, we generated
KIC11-2xFKBP-GFP-2xFKBP[endo] parasites episomally co-expressing the nuclear
mislocaliser 1xNLS-FRB-mCherry, enabling conditional inactivation by knock-
sideways. Addition of rapalog resulted in efficient mislocalisation of KIC11-2xFKBP-
GFP-2xFKBP into the nucleus as evaluated 4 and 16 hours post induction (Figure 2D).
Assessment of parasite proliferation by flow-cytometry over two developmental cycles
revealed a mean relative growth of 10.3% compared to control parasites, indicating an
important function of KIC11 for asexual parasite proliferation (Figure 2E, Figure S3D).
This interpretation was supported by several unsuccessful attempts to generate a cell
line with a truncated *kic11* using the SLI-TGD system [73].

To better characterize the growth phenotype of the KIC11 knock-sideways, we added
rapalog to tightly synchronised parasites at different time points (4, 24, and 32 hpi)
and monitored parasite growth by flow cytometry. Additionally, we quantified parasite
stage in Giemsa smears at 6, 24, 32, 40, 48, 72, and 96 hpi. While no effect on
parasitemia and stage distribution was observed during growth in the first cycle, a
reduced number of newly formed ring stage parasites was obvious at the beginning of
the second cycle, indicating an invasion defect or an effect on parasite viability in
merozoites or early rings but no effect on other parasite stages (Figure 2F-H, Figure
S3F-G). As KIC11 appeared to be important at the schizont to ring transition, we
further analysed the phenotype at this stage. KIC11 inactivation did not affect egress
but resulted in fewer detectable rings, suggesting a defect in invasion or possibly very
early ring stage development (Figure 2H, S3G). No consistent overlap of KIC11 foci
with rhoptry (ARO) or microneme (AMA1) marker proteins was observed, while as
expected [65] for a K13 compartment protein, an overlap with the IMC marker (IMC1c)
was apparent. In addition, KIC11 inactivation did not result in a detectable alteration
of these markers, excluding major structural disorganisation of the micronemes,
rhoptries or IMC as a reason for the observed phenotype (Figure S3A).

As trophozoites with inactivated KIC11 developed into schizonts without
morphological differences to controls (Figure 2G, Figure S3A), KIC11 does not appear
to be needed for endocytosis. To directly address this we tested if inactivation of KIC11
influences *in vitro* ART resistance (based on RSA) or endocytosis (using vesicle
accumulation and bloated food vacuole assays), but no significant differences were
observed (Figure 2I-K).

Overall, our results indicate that KIC11 is part of the K13 compartment but also in additional foci and that it has an important role in invasion or very early ring stage development, which is in contrast to previously characterised essential K13 compartment proteins [29]. These findings indicate that there are also proteins found at the K13 compartment that have important functions not related to endocytosis or ART resistance.

KIC12 is located in the nucleus and at the K13 compartment and is involved in endocytosis but not in ART resistance

PF3D7_1329500, currently annotated as 'conserved protein, unknown function' was renamed K13 interacting candidate 12 (KIC12). In order to test whether KIC12 is a member of the K13 compartment or not, we first generated transgenic parasites expressing C-terminally 2xFKBP-GFP-2xFKBP tagged KIC12 from its original genomic locus (KIC12-2xFKBP-GFP-2xFKBP[endo] parasites) (Figure S1E). Expression and localisation of tagged KIC12 was analysed by fluorescent microscopy. KIC12 was detectable in the nucleus in ring stage parasites. In trophozoites foci in the parasite periphery (Figure 3A, white arrows) were observed in addition to the signal in the nucleus (Figure 3A, light blue arrows). In schizonts these foci were not present anymore. Instead, only the nuclear signal and a faint uniform cytoplasmic GFP signal was detected in late trophozoites and early schizonts and these signals were absent in later schizonts and merozoites (Figure 3A, Figure S4A,B). In line with this expression pattern, the *kic12* transcriptional profile indicated mRNA levels peak in merozoites and early ring-stage parasites and no RNA expression in trophozoites and schizonts [74,75,82]. Next, we generated KIC12-2xFKBP-GFP-2xFKBP[endo] parasites episomally co-expressing mCherry-K13 which revealed 49% of parasites with overlap, 46% with partial overlap, while 5% showed no overlap of the KIC12 foci with the K13 foci in early trophozoites (Figure 3B, Figure S4A). In rings only 4% of parasites showed an overlap of KIC12 with K13 foci, while in late trophozoites no overlap was observed, in agreement with the exclusively nuclear localisation of KIC12 in these stages (Figure 3B, S4A). We conclude that KIC12 is a protein with a dual location in the nucleus and the K13 compartment in trophozoites.

In order to assess the importance of the K13-located pool of KIC12 for parasite proliferation, we generated KIC12-2xFKBP-GFP-2xFKBP[endo] parasites enabling conditional KIC12 inactivation by knock-sideways using a nuclear (1xNLS-FRB-mCherry) mislocaliser. Efficient mislocalisation of KIC12-2xFKBP-GFP-2xFKBP into the nucleus and absence of KIC12 foci in the cytoplasm of trophozoites upon addition of rapalog was confirmed by microscopy at 4 and 16 hours post induction (Figure 3C). Assessing parasite proliferation after knock sideways of KIC12 showed a mean

relative growth of 37.0% compared to control parasites after two development cycles,
indicating an important function of the K13 compartment located pool of KIC12 for
asexual parasite proliferation (Figure 3D; Figure S4C).

Due to the dual localisation of KIC12 we also generated KIC12-2xFKBP-GFP-
2xFKBP[endo] parasites episomally co-expressing an alternative mislocaliser (Lyn-FRB-
mCherry) [73], enabling conditional inactivation of the nuclear pool of KIC12-2xFKBP-
GFP-2xFKBP by mislocalisation to the parasite plasma membrane (PPM), an
approach previously shown to be suitable for efficient inactivation of nuclear proteins
[73,83,84]. Induction of KIC12 mislocalisation to the PPM resulted in a loss of KIC12
in the nucleus 4 hours post induction (Figure 3E). Foci were still detected in the
parasite periphery and it is unclear whether these remained with the K13 compartment
or were also in some way affected by the Lyn-mislocaliser. Parasite proliferation
assays revealed a growth defect of 50.8%, compared with a defect of 63.0% of the
nuclear mislocalisation approach (Figure 3D; Figure S4D). Due to the possible
influence on the K13 compartment located foci of KIC12 with the Lyn mislocaliser, a
clear interpretation in regards to the functional importance of the nuclear pool of KIC12
other than that it confirms the importance of this protein for asexual blood stages is
not possible. In contrast, the results with the nuclear mislocaliser indicate that the K13
located pool of KIC12 is important for efficient parasite growth. The general importance
of KIC12 for asexual parasite proliferation was further supported by failure to obtain a
cell line with a truncated *kic12* using the SLI-TGD system.

Based on the presence at the K13 compartment in trophozoites, we tested the effect
of KIC12 inactivation on endocytosis. Bloated food vacuole assays showed that >99%
of control parasites developed bloated food vacuoles, while only 49,3% (NLS
mislocaliser) or 72.9% (Lyn mislocaliser) of the cells with inactivated KIC12 showed a
bloated food vacuole (Figure 3F, Figure S4E), indicating an effect on endocytosis of
host cell cytosol. As the effect was only partial, we decided to measure the parasite
and food vacuole size. This analysis with the KIC12 NLS mislocaliser parasites
revealed a significantly reduced food vacuole size in the parasites with inactivated
KIC12, while there was no effect on parasite size (Figure 3G; Figure S4F). Plotting the
values of the individual parasites showed that the food vacuoles of similarly sized
parasites were consistently smaller in the KIC12 NLS knock sideways compared to
controls (Figure 3H), indicating that the effect on hemoglobin delivery to the food
vacuole upon inactivation of KIC12 was not an indirect effect due to parasite growth
impairment during the assay time. Similar results were obtained using the KIC12 Lyn
mislocaliser line (Figure 3I,J, S4G). Quantification of the number of vesicles in
trophozoites upon inactivation (with either the NLS or Lyn mislocaliser) of KIC12

revealed no difference compared to control (Figure 3K), indicating that KIC12 acted in
the early phase of endocytosis, similar to the previously studied K13 compartment
proteins [29]. While again the data with the Lyn mislocaliser is more difficult to interpret,
the results with the NLS mislocaliser indicate that the K13 compartment located pool
of KIC12 is important for endocytosis of host cell cytosol.
404
405 The lacking evidence for a K13 compartment location of KIC12 in rings indicates that
406 KS using the NLS mislocaliser would not affect ART susceptibility. Indeed, rapalog-
407 induction of the KIC12 NLS mislocaliser KS line showed no significantly decreased
408 ART susceptibility in RSAs (Figure 3L, Figure S4H), congruent with the lacking co-
409 localisation of KIC12 with K13 in rings.
410
411 Overall, our results indicate the presence of KIC12 at the K13 compartment in
412 trophozoites, a role in HCCU and an additional pool of KIC12 in the nucleus.
413
MCA2 is part of the K13 compartment and its truncation confers *in vitro* ART
resistance
We previously identified metacaspase-2 (MCA2) as one of the top 5 most enriched
proteins of the K13 proxiome, but it was so far not localised, as endogenous C-terminal
tagging with a 2xFKBP-GFP-2xFKBP tag had not been achieved despite several
attempts [29]. By screening of the MalariaGEN *Plasmodium falciparum* Community
Project database [85] a single-nucleotide polymorphism (SNP) at amino acid position
1344 of MCA2 was identified that on its own leads to a stop codon, thereby removing
the metacaspase (MCA) domain. According to MalariaGEN, this SNP is found in Africa
with a mean prevalence of 52% and in South East Asia with 5% prevalence [85]. As a
previously generated truncation of MCA2 at amino acid position 57 (MCA2^{TGD}-GFP
(targeted gene disruption (TGD)) reduced the parasites' ART susceptibility as judged
by RSAs [29], we generated a second cell line with MCA2 disrupted at the Y1344
position (MCA2^{Y1344Stop}-GFP^{endo} parasites) (Figure 4A), originally to assess if this
change could mediate resistance in the field. Subsequent analyses of African parasite
isolates [86] revealed that this change was always accompanied by a second change
in the following base, reverting the stop codon, indicating that this change is not of
relevance for resistance in endemic settings. However, as we had not so far been able
to localise MCA2, we took advantage of these parasites with the first 1344 amino acids
of MCA2 (66.5% of the protein) fused to GFP to study its localisation. Correct
integration of the construct into the *mca2* genomic locus was confirmed by PCR
(Figure S1F) and expression of the truncated fusion protein was verified by Western
blot (Figure S1F). Live cell imaging of MCA2^{Y1344Stop}-GFP^{endo} parasites revealed

expression of MCA2 throughout the intraerythrocytic development cycle, appearing as
a single focus in ring stage parasites while two or more foci were detectable in
trophozoites and schizonts (Figure 4B), in contrast to the predominantly cytosolic
localisation of the severely truncated MCA2^{TGD}-GFP (Figure S5A). As this localisation
is reminiscent of K13 and KIC localisation [29,73] we generated MCA2^{Y1344Stop}-GFP^{endo}
parasite episomally co-expressing mCherry-K13 and the spatial relationship of
MCA2^{Y1344Stop}-GFP and K13 foci was quantified in ring and trophozoite stage parasites
based on fluorescence microscopy (Figure 4C). This analysis showed that 59% of the
MCA2^{Y1344Stop} foci overlapped with K13 foci, 21% of the foci showed partial overlap
and 20% of the MCA2^{Y1344Stop} foci did not overlap with K13 foci (Figure 4C), indicating
a presence of MCA2 at the K13 compartment. In order to exclude an effect of the
truncation on MCA2 localisation we generated parasites with an endogenously 3xHA-
tagged full length MCA2 (MCA2-3xHA^{endo}) (Figure S1G) that episomally co-expressed
mCherry-K13 and performed immunofluorescence assays (IFA). These experiments
confirmed the focal localisation of MCA2. Quantification of the relative localisation of
MCA2 and K13 revealed that 44% of MCA2 foci overlapped with K13 (Figure S5B).
Partial overlap was observed for 24% of MCA2 foci, whereas 32% MCA2 foci did not
overlap with K13 (Figure S5B). Overall, these findings indicated that MCA2 is a K13
compartment protein but is also found in additional, non-K13 compartment foci.
456
MCA2 was recently found to be enriched in BioID experiments with the inner
membrane complex (IMC) marker protein PhIL1 [65]. We generated MCA2^{Y1344Stop}-
GFP^{endo} parasites episomally co-expressing PhIL1mCherry and analysed schizont
stage parasites. Fluorescent imaging with these parasites revealed a close association
of MCA2^{Y1344Stop}-GFP foci with the IMC at the periphery of the newly formed
merozoites (Figure 4D), as previously observed for K13 [65]. This data indicated that
MCA2 and the K13 compartment - as previously suggested [30,65] - are found
proximal to the IMC in schizonts.
465
We previously showed that a truncation of MCA2 at amino acid position 57 (2.8% of
the protein left) results in significantly reduced *in vitro* parasite proliferation [29].
Proliferation assays with the MCA2^{Y1344Stop}-GFP^{endo} parasites which express a much
larger portion of this protein (66.5% of the protein left), yet still lack the MCA domain
(Figure 4A), indicated no growth defect in these parasites compared to 3D7 wild type
parasites (Figure 4E). Hence, the MCA domain in MCA2 does not appear to be needed
for efficient *in vitro* blood stage growth.
473

Next, we tested the effect of MCA2 truncation on endocytosis. For this we adapted the
bloated food vacuole assays to compare MCA2-TGD and MCA2^{Y1344Stop}-GFP^{endo} with
3D7 parasites. The assay showed that in all three cell lines >95% of parasites
developed bloated food vacuoles (Figure S5C). Additionally, there was no apparent
difference in the size of food vacuoles of similarly sized parasites between the three
lines (Figure S5D-E). Of note, in line with the reduced parasite proliferation rate [29],
an effect on parasite and food vacuole size was observed for the MCA2-TGD parasites
(Figure S5D-E). Overall, with this assay, there was no detectable effect on endocytosis
in trophozoites in these parasite lines.
483
Last, we performed RSAs which revealed that the truncation of MCA2 at amino acid
position Y1344 resulted in a reduced sensitivity to ART (Figure 4F). The mean parasite
survival rate was 1.64%, which is above the defined ART resistance cut-off value of
1% [26], but lower than the ~5% survival rate of the MCA2-TGD parasites [29]. This
considerable reduction in ART susceptibility in the parasites with the truncation at
MCA2 position 57 compared to the parasites still expressing 1344 amino acids of
MCA2, despite both versions of the protein lacking the MCA domain, indicates that the
influence on ART resistance is not, or only partially due to the MCA domain.
492

Candidate proteins not detected at the K13 compartment
We also generated parasites expressing endogenously C-terminally 2xFKBP-GFP-
2xFKBP tagged UIS14, PF3D7_1365800, PF3D7_1447800, PF3D7_0907200 and
VPS51 and episomally co-expressed mCherry-K13 (Figure S1H-L). Fluorescence
microscopy revealed no clear association with the K13 compartment in rings and
trophozoites in any of these parasite lines. Instead UIS14, PF3D7_1447800 and
VPS51 showed GFP foci in the parasite cytosol without consistent overlap with
mCherry-K13 foci (Figure S6A,D,F), PF3D7_0907200 showed a weak cytosolic GFP
signal and PF3D7_1365800 showed cytosolic GFP signal with additional foci closely
associated with the nucleus without consistent overlap of the main foci with mCherry-
K13 foci (Figure S6G and Figure S7A-B). Several attempts to generate
PF3D7_1243400-2xFKBP-GFP-2xFKBP^{endo} or 2xFKBP-GFP-2xFKBP-
PF3D7_1243400^{endo} parasites remained unsuccessful, indicating the gene might be
refractory to C- and N-terminal modification and hence might be essential. For *vps51*
and *uis14* we additionally were able to generate targeted gene disruption cell lines
(Figure S1M-N, Figure S6B,E), indicating these candidates are dispensable for *in vitro*
asexual parasite proliferation, although growth assays indicated a need of UIS14 for
optimal growth (Figure S6C).
511

Structural homology search [87] revealed the presence of a N-terminal arfaptin homology (AH) domain in PF3D7_1365800 (Figure S8C), a domain known to promote binding of arfaptin to Arf and Rho family GTPases important for vesicle budding at the Golgi [88,89]. Given that in *Toxoplasma* an intersection of endocytosis and secretion was observed at the trans Golgi [90], we tested the potential (indirect) influence of this protein on endocytosis related processes. We performed conditional inactivation using knock-sideways, but despite efficient loss of the PF3D7_1365800 foci, no growth defect was observed (Figure S7C-D) and co-expression of the Golgi marker GRASP [91] revealed no consistent overlap between the foci of these two proteins (Figure S7E).

Based on this analysis we did not classify UIS14, PF3D7_1365800, PF3D7_1447800, PF3D7_0907200, and VPS51 as K13 compartment proteins and the location of PF3D7_1243400 remains unknown.

The domain repertoire in K13-compartment proteins
With the extended complement of K13 compartment proteins from this and previous work [29], we assessed the repertoire of functional domains at this site. For this we took advantage of recent advances in protein structure prediction to identify structural similarities in K13-compartment members for which no information could be inferred from sequence homology. We compared their structures predicted with the AlphaFold algorithm [87,92] with experimentally determined protein structures in the Protein Data Bank and identified 25 domains, 15 of which were not previously identified according to PlasmoDB and Interpro (Figure 5A,B and S8).

The largest number of recognisable folds were detected in KIC4, a protein for which we previously detected some similarity to α-adaptins [29]. KIC4 contained an N-terminal VHS domain (IPR002014), followed by a GAT domain (IPR004152) and an Ig-like clathrin adaptor α/β/γ adaptin appendage domain (IPR008152) (Figure 5A-C, Figure S8). This is an arrangement typical for GGAs (Golgi-localised gamma ear-containing Arf-binding proteins) which are vesicle adaptors first found to function at the trans-Golgi [93,94]. Surprisingly, KIC4 however also contains an additional domain at its C-terminus, a β-adaptin appendage C-terminal subdomain (IPR015151) which is a part of the ear domain of β-adaptins and not found in GGAs. Together with the preceding clathrin adaptor α/β/γ adaptin domain, the C-terminus of KIC4 therefore resembles a β-adaptin. The region without detectable fold after the GAT domain likely corresponds to the hinge region in GGAs. This suggests that KIC4 is a hybrid between GGAs and an AP complex subunit beta (Figure 5C), the two known types of adaptors.

Based on Interpro [95], such a domain organization has to date not been observed in
any other protein.

KIC5 also contains a clathrin adaptor α/β/γ-adaptin domain (IPR008152) and one of
the two subdomains of the ear domain of α-adaptins (clathrin adaptor α-
adaptin_appendage C-terminal subdomain, IPR003164) (Figure 5 A,B,D). More than
97% of proteins containing these domains also contain an Adaptin_N (IPR002553)
domain [95] and in this combination typically function in vesicle adaptor complexes as
subunit α [96,97] (Figure 5D) but no such domain was detectable in KIC5. KIC5 thus
displays some similarities to AP complex subunit α but similar to KIC4, there appear
to be profound difference to the canonical adaptors.

Based on the identified adaptor domains in KIC4 and KIC5 we decided to test for an
effect on endocytosis in previously established parasites with truncations in these
genes [29]. But neither KIC4-TGD nor KIC5-TGD parasites showed a difference in
bloated food vacuole assay compared to 3D7 parasites (Figure S9). Of note, as for
MCA2-TGD, a reduced parasite and food vacuole size was observed for KIC5-TGD,
which is in line with the reported reduced parasite proliferation [29]. While this overall
indicates no role of these proteins in endocytosis in trophozoites, we note that TGD
lines might not be ideal to capture non-absolute effects on endocytosis. Also, if an
endocytosis defect were the reason for the slowed parasite growth in the KIC5- and
MCA2-TGD lines, our assay would not show this as an endocytosis defect.

KIC7 contains an ArfGAP domain, as recently also predicted for its *Toxoplasma*
homolog AGFG (TGME49_257070) [98]. ArfGAPs regulate the activity of Arfs which
are small GTP binding proteins coordinating vesicle trafficking [99,100]. UBP1
contains a ubiquitin specific protease (USP) domain at its C-terminus which previously
led to its name ubiquitin carboxyl-terminal hydrolase 1 (UBP1) [101]. Here we also
identified a VHS domain in its centre. VHS domains occur in vesicular trafficking
proteins, particularly in endocytosis proteins, but typically are found at the N-terminus
of proteins (in over 99.8% of cases according to annotations by Interpro). A
combination with a USP domain has not been observed so far. If the VHS domain is
functional in UBP1 despite its central position, it is the first structural domain that would
support the functional data showing this K13 compartment protein has a role in
endocytosis [29].

KIC2 contains a GAR domain which typically bind the cytoskeleton [102] and KIC3
contains a PAH domain which can serve as a scaffold for transcription factors
[103,104]. While not consistently detected at the K13 compartment, it is interesting

that we here found that PF3D7_1365800 contains an AH domain, a member of the
AH/BAR domain family (Figure S8C). According to Interpro, no AH or BAR domain
proteins have so far been detected in malaria parasites, rendering this an interesting
protein. However, we found this protein as being likely dispensable for intra-
erythrocytic parasite development and no colocalisation with K13 was observed,
suggesting PF3D7_1365800 is not needed for endocytosis.
595
Overall this analysis revealed that most of the proteins involved in endocytosis or *in*
vitro ART resistance contain regions with structural homology to vesicle trafficking
domains, often specific for endocytosis (Figure 5E). However, apart from AP2 the
domain arrangements are unusual, the conservation on the primary sequence level is
low (which precluded initial detection) and there are large regions without any
resemblance to other proteins, altogether indicating strong parasite-specific
adaptations. In contrast, the K13 compartment proteins where no role in ART
resistance (based on RSA) or endocytosis was detected, KIC1, KIC2, KIC6, KIC8,
KIC9 and KIC11, do not contain such domains (Figure 5E). This analysis suggests
that proteins detected at the K13 compartment can be classified into at least two
groups of which one comprises proteins involved in endocytosis or *in vitro* ART
resistance whereas the other group might have different functions yet to be
discovered.
609
DISCUSSION
The BioID proxiome of K13 and Eps15 revealed the first proteins involved in the initial
steps of endocytosis in malaria parasites, a process that in blood stage parasites leads
to the uptake of large quantities of host cell cytosol. As demonstrated in model
organisms, endocytosis is a complex and highly regulated process involving a
multitude of proteins [105–107], but in Apicomplexan parasites is not well studied [42].
Thus, the K13 compartment proxiome represents an opportunity to identify proteins
involved in this process in malaria parasites and will be important to understand it on
a mechanistic level. Understanding how and if HCCU differs from the canonical
endocytic processes in human cells, will not only help to understand this critical
process in parasite biology but might also reveal parasite-specific aspects that permit
specific inhibition and could be targets for drug development efforts.
622
Here we expanded the repertoire of K13 compartment proteins and functionally
analysed several of them. An assessment of structural similarities indicated an
abundance of vesicle trafficking - and more specifically - endocytosis domains, in the
confirmed K13 compartment proteins. The proteins with such domains now comprise
KIC4, KIC5, KIC7, Eps15, AP2µ and UBP1, all of which reduce *in vitro* ART

susceptibility when inactivated, indicating that this led to a reduced endocytosis in ring
stages [29]. A role in endocytosis has been experimentally shown for all of these
except for KIC4 and KIC5, which are both non-essential for the parasites asexual
blood stage development and for which only gene disruptions were available [29]. We
here nevertheless attempted to assess endocytosis in the corresponding TGD lines
but failed to detect an effect. This indicates that either these proteins are not needed
for endocytosis (which would be unexpected given the altered RSA response of these
TGD lines), or that we did not detect it for technical reasons. This is possible because
a comparably small reduction in protein abundance (or activity) of endocytosis proteins
can already lead to a substantial increase in RSA survival [29,40]. Hence, it is possible
that the effect was too small for our assays to measure it. We also note that if the
growth defect in the TGD lines is due to endocytosis, the food vacuole size reduction
may be proportional to parasite size reduction which would not be classified as an
endocytosis defect in the assay. Finally, it is also possible that these proteins are only
important for endocytosis in ring stages. Further work will be needed to clarify this.
643
With KIC4 and KIC5 malaria parasites contain two additional K13 compartment
proteins (besides the AP2 complex) that based on our structural similarity analysis
contain domains found in adaptor subunits. KIC4 and KIC5 likely have additional rather
than redundant roles to the AP2 complex, as the AP2 complex is essential for blood
stage growth and on its own is needed for HCCU [29,108]. While KIC4 disruption did
not lead to a growth defect, disruption of KIC5 impaired parasite growth [29], indicating
that there cannot be full redundancy between these two proteins either. Of note, in *T.*
gondii a homologue of KIC4 (*Tg*KAE, TGME49_272600) was detected at the
micropore and shown to be essential, while no homologue of KIC5 could be identified
[98,109]. Altogether these considerations support specific individual roles of all the
known K13 compartment proteins with adaptor domains.
655
In model organisms, the α/β/γ-adaptin appendage domain, the α-adaptin domain and
the β-adaptin domain, in the AP2 complex and in GGAs, act as interaction hubs for
more than 15 accessory proteins, including Eps15, Arfs, amphiphysin, epsins and
rabaptins as well as lipids involved in vesicle budding [110]. This fits with the presence
of Eps15 and KIC7 (which contains an ArfGAP domain) at the K13 compartment. The
interaction between the α/β/γ-adaptin appendage domain and Eps15 has been
captured by X-ray crystallography (2I9V) [110], highlighting a possible functional
connection between Eps15 and AP-2α, AP-2β, KIC4 or KIC5, which all contain this
domain. Indeed, the *Toxoplasma* homologue of KIC4 has recently been shown to bind
Eps15, while *Tg*AP-2α did not [98].
666

Despite the detection of various domains in K13 compartment proteins it is noteworthy
that most of these proteins (e.g. EPS15, UBP1, KIC7) still contain large regions without
any homology to other proteins. The parasite-specific nature of the initial steps of
endocytosis is also evident from the difficulty of primary sequence-based detection of
the vesicular trafficking domains in K13 compartment proteins and from the unusual
domain combinations. Furthermore it is peculiar that despite the presence of the
clathrin adaptor AP2, clathrin itself does not seem to be involved [29,108], further
indicating parasite-specific features of HCCU. Overall, this indicates a strongly
adapted mechanism of the first steps in endocytosis for HCCU in malaria parasites.
One protein typically involved in endocytosis that did not appear in the list of highly
enriched proteins of the K13 and Eps15 proxiome is a dynamin [111] which might
indicate further differences to the canonical mechanism in model organism. However,
at least in *Toxoplasma* an association of K13 and its compartment with a dynamin was
reported [98,112], indicating that an equivalent is likely also present in malaria
parasites.
682
A protein that has a more canonical structure is MyoF, a Class XXII myosin [77], which
we here link with endocytosis and found in foci that were often close or overlapping
with the K13 compartment. The only partial overlap could indicate that either it i) only
transiently associates with the K13 compartment, or ii) is in a separate compartment
that is close to the K13 compartment or iii) is in a region of the K13 compartment that
is close, but non-overlapping, with that defined by the other so far known K13
compartment proteins. A number of conclusions can be drawn from our MyoF
characterisation. Firstly, its inactivation resulted in the appearance of vesicles, similar
to the phenotype of VPS45 inactivation [54], indicating it has a role in endosomal
transport, downstream of the initial steps of endocytosis. This is in contrast to the other
K13 compartment proteins and might explain the only partial overlap in location of
MyoF with the K13 compartment. Secondly, its involvement suggests a role of
actin/myosin in endosomal transport which is well known from other organisms
[113,114] and supports the observation that the actin inhibitor CytochalasinD leads to
vesicles similar to MyoF inactivation [53]. Hence, myosin may generate force needed
for the transport of host cell cytosol filled vesicles to the food vacuole. Thirdly, there
was no effect of MyoF inactivation or CytochalasinD treatment on ART resistance,
which might indicate that there is no need for actin/myosin for endocytosis in rings.
Although there were some limitations to our system to study MyoF, the substantial
inactivation caused by simply tagging MyoF would already have led to a decreased
susceptibility to ART in RSAs (as seen with other K13 compartment proteins [29]), if
ring stage endocytosis had been affected, which was not the case. It is of note that
the endocytosis function of MyoF is reciprocal to that of K13 which is only required in

rings, indicating that there appear to be stage-specific differences in endocytosis
between these stages. However, the majority of components are at the K13
compartment throughout the cycle and for instance KIC7 is needed for HCCU in both
rings and trophozoites [29]. Of note, mutations in *myoF* have previously been found to
be associated with reduced ART susceptibility [115], but 12 mutations tested in the
laboratory strain 3D7 did not result in increased RSA survival [40].

Attempts to generate a knockout of *P. berghei myof* were reported to be unsuccessful
[77], while a genome-wide mutagenesis screen predicted *Pf*MyoF to be dispensable
for asexual blood stage development [116]. *Tg*MyoF has previously been implicated
in Golgi and rhoptry positioning [117], trafficking of dense granule [118] or nuclear
encoded apicoplast proteins [119] or centrosomes positioning and apicoplast
inheritance [120]. We cannot exclude that beside the effect on HCCU, a part of the
growth defect we see after MyoF inactivation arises from similar functions in malaria
blood stages.

MCA2 has been identified as putative K13 interaction candidate, but its location was
unknown [29]. Here, we establish MCA2 as member of the K13 compartment by
endogenously tagging it with HA as well as by using a cell line with a truncated MCA2
fused to GFP. MCA2 was located in foci overlapping with the K13 compartment and
in schizonts the foci were in proximity of the IMC in line with previous data linking the
K13 compartment and the IMC [65,121]. A close association of the K13 compartment
and the IMC is also supported by recent work in *T. gondii* [98,109,122] showing that
the K13 compartment and its proteins are present at the IMC embedded micropore
[98,109].

We previously showed that gene disruption of *mca2* results in reduced *in vitro* parasite
proliferation and *in vitro* ART resistance [29], which is also supported by a knockout of
the *mca2* homologue in the rodent infecting *P. berghei* [123,124]. Truncation of MCA2
at amino acid position Y1344 still resulted in decreased ART susceptibility, even
though the susceptibility reduction was less than a disruption at amino acid position
57. In contrast to the full disruption, truncation at residue 1344 did not lead to a growth
defect, indicating that the predicted metacaspase domain (AA 1527-1848) – not
present in this truncated MCA2 protein – is dispensable for asexual parasite
proliferation. This finding is in contrast to results indicating an important role of the
metacaspase domain for parasite proliferation using MCA2 inhibitors [125–127].

Here we also identified the first protein, KIC11, at the K13 compartment that had an
important function for the growth of blood stage parasites but did not appear to function

in endocytosis. Instead, KIC11 had an important function in invasion or possibly early ring stage development. KIC11 was predicted to be non-essential by a genome-wide mutagenesis screen [116], while the orthologue in the rodent malaria parasite *P. berghei* (PBANKA_0906900) was classified as essential for asexual proliferation in the PlasmoGem screen [128].

Overall the classification of K13 compartment proteins presented in this work indicates that there are two main groups. The first group includes proteins that define a highly unusual endocytic pathway to internalise host cell cytosol. These proteins are predominately parasite-specific (exceptions being AP2μ, MyoF and in part K13 which however lacks vesicle trafficking domains). This group includes proteins that are involved in *in vitro* ART resistance (K13, MCA2, UBP1, KIC7), but also proteins such as KIC12 or MyoF that do not confer *in vitro* ART resistance when inactivated, presumably because they are not needed for endocytosis in rings. In the case of MyoF this idea is supported by low expression in rings whereas KIC12 was not at the K13 compartment in rings. Reciprocal to this, we found in previous work that K13 is needed for endocytosis in ring stages only [29]. Hence, there is heterogeneity in the stage-specificity of the endocytosis proteins at the K13 compartment. In terms of function, most of the endocytosis and *in vitro* ART resistance group of K13 compartment proteins are involved in the initial phase of HCCU. In contrast to VPS45 [54] their inactivation does not result in the generation of endosomal intermediates in the parasite cytoplasm. The exception is MyoF which (of the proteins here classified as K13 compartment associated) generated vesicles when inactivated. MyoF showed the lowest spatial overlap with K13 and may form a link to downstream steps of endosomal transport. The second group of K13 compartment proteins neither affects RSA survival nor endocytosis and does not contain vesicle trafficking domains (as based on our domain identification here). These proteins likely serve other functions but the homogeneity of functions of this group is unclear. KIC11, the first essential protein of this group, might help to shed light on the function of the non-endocytosis related K13 compartment proteins. However, it should be noted that KIC11 showed an additional protein pool independent of the K13 compartment, most prominently in schizonts. Hence, it is possible that KIC11's essential function is not related to the K13 compartment but is carried out by the pool of KIC11 located elsewhere.

Overall, this work strengthens the notion that the K13 compartment is involved in endocytosis, reveals novel essential HCCU proteins and provides a classification of its members that might inform future studies to understand the unusual mechanism of endocytosis in apicomplexans.

METHODS

P. falciparum culture

Blood stages of *P. falciparum* 3D7 [129] were cultured in human red blood cells (O+ ; University Medical Center Hamburg, Eppendorf (UKE)). Cultures were maintained at 37°C in an atmosphere of 1 % O_2, 5 % CO_2 and 94 % N_2 using RPMI complete medium containing 0.5 % Albumax according to standard protocols [130].

In order to obtain highly synchronous parasite cultures, late schizonts were isolated by percoll gradient [131] and cultured with fresh erythrocytes for 4 hours. Afterwards sorbitol synchronization [132] was applied in order to remove remaining schizonts resulting in a highly synchronous ring stage parasite culture with a four-hour age window.

Cloning of plasmid constructs for parasite transfection

For endogenous C-terminal 2x-FKBP-GFP-2xFKBP tagging using the SLI system [73] a homology region of 321-1044 bp (1044 bp for *Pfmyof* (PF3D7_1329100), 690 bp for *Pfkic11* (PF3D7_1142100), 780 bp for PF3D7_1447800, 695 bp for *Pfkic12* (PF3D7_1329500), 411 bp for PF3D7_1243400, 321 bp for PF3D7_1365800, 674 bp for *Pfuis14* (PF3D7_0405700), 698 bp for PF3D7_0907200, 756 bp for *Pfvps51* (PF3D7_0103100)) was amplified from 3D7 gDNA and cloned into pSLI-sandwich [73] using the NotI/AvrII restriction sites.

For endogenous C-terminal 2x-FKBP-GFP tagging using the SLI system [73] a homology region of 1044 bp for *Pfmyof* (PF3D7_1329100), was amplified using 3D7 gDNA and cloned into pSLI-2xFKBP-GFP [73] using the NotI/AvrII restriction sites.

For endogenous C-terminal 3xHA tagging using the SLI system [73] a homology region of 948-999 bp (999 bp for *Pfmca2* (PF3D7_1438400), 948 bp for *Pfmyof* (PF3D7_1329100)) was amplified using 3D7 gDNA and cloned into pSLI-3xHA [133] using the NotI/XhoI restriction sites.

For endogenous N-terminal 2x-FKBP-GFP-2xFKBP tagging of PF3D7_1243400 using the SLI system [73] a homology region of 276 bp was amplified from 3D7 gDNA and cloned into p-N-SLI-sandwich-loxp [73] using the NotI/PmeI restriction sites. A functional and codon-changed version of PF3D7_1243400 including the C-terminal loxp site was synthesised (BioCat GmbH, Germany) and cloned into the construct using the AvrII/XhoI restriction sites.

For generating *Pf*MCA2^{Y1344Stop}-GFP^{endo} a 984 bp homology region was amplified using 3D7 gDNA and cloned into the pSLI-TGD plasmid [73] using NotI and MluI restriction sites.

For targeted gene disruption (TGD) a 429-617 bp (534 bp for *kic12* (PF3D7_1329500), 615 bp for *Pfkic11* (PF3D7_1142100), 617 bp for *Pfuis14* (PF3D7_0405700), 429 bp

for *Pfvps51* (PF3D7_0103100)) was amplified using 3D7 gDNA and cloned into the
pSLI-TGD plasmid [73] using NotI and MluI restriction sites.

All oligonucleotides used to generate DNA fragments as well as those used for
genotyping PCRs are listed in table S2.

For co-localisation experiments the plasmids p40PX-mCherry [54], pmCherry-
K13_DHODH*nmd3* [29], *ama1*PhIL1mCherry [134] and pGRASPmCherry-BSD*nmd3* [29]
were used.

In order to generate the plasmids for co-localisation experiments while simultaneously
carrying out knock sideways, *ama1*AMA1mCherry, *ama1*AROmCherry and
*ama1*IMC1cmCherry were amplified from pARL*ama1*AMA1mCherry,
pARL*ama1*AROmCherry or pARL*ama1*IMC1cmCherry [65] by PCR and cloned into
ama1-p40-mSca_*nmd3'*-NLS-FRB-T2A-yDHODH, which was generated by inserting
the *ama1* promoter using the NotI/KpnI restriction sites into *sf3a2*-p40-mSca_*nmd3'*-
NLS-FRB-T2A-yDHODH [55], using the KpnI/XmaI restriction sites, resulting in
*ama1*AROmCherry_*nmd3'*-NLS-FRB-T2A-yDHODH, *ama1*AMA1mCherry_*nmd3'*-NLS-
FRB-T2A-yDHODH and *ama1*IMC1cmCherry_*nmd3'*-NLS-FRB-T2A-yDHODH,

Transfection of *P. falciparum*
For transfection, Percoll-purified [131] parasites at late schizont stage were
transfected with 50 µg plasmid DNA using Amaxa Nucleofector 2b (Lonza,
Switzerland) as previously described [135]. Transfectants were selected using either
4 nM WR99210 (Jacobus Pharmaceuticals), 0.9 µM DSM1 [136] (BEI Resources) or
2 mg/ml for Blasticidin (BSD) (Invitrogen). In order to select for parasites carrying the
genomic modification via the SLI system [73], G418 (ThermoFisher, USA) at a final
concentration of 400 µg/mL was added to a culture with about 5 % parasitemia. The
selection process and integration test were performed as previously described [73].

Imaging
All fluorescence images were captured using a Zeiss Axioskop 2plus microscope with
a Hamamatsu Digital camera (Model C4742-95).

Microscopy of live parasite-infected erythrocytes was performed as previously
described [137]. Approximately 5 µL of infected erythrocytes were added on a glass
slide and covered with a cover slip. Nuclei were stained with 1 µg/mL Hoechst-33342
(Invitrogen) or 1µg/mL 4',6'-diamidine-2'-phenylindole dihydrochloride (DAPI)
(Roche).

Immunofluorescence assays (IFA) were performed as previously described [133].
Asynchronous parasite cultures with 5% parasitemia were harvested, washed twice
with PBS, air-dried as thin monolayers on 10-well slides (Thermo Fischer) and fixed in
100% acetone for 30 min at room temperature. After rehydration with PBS, the cells

were incubated with primary antibody solution containing rabbit a-HA (1:500) (Cell Signalling) and rat a-RFP (1:500) (Chromotek) diluted in 3% BSA in PBS. After three wash steps with PBS, incubation with corresponding secondary antibodies (Molecular probes) was performed. Slides were sealed with a coverslip using mounting medium (Dako).

Parasite proliferation assay

For parasite proliferation assays a flow cytometry based assay, adapted from previously published assays [54,138], was performed to measure multiplication over five days. These assays were either preformed using synchronous or non-synchronous parasites, as indicated in the corresponding figure legends. For synchronous proliferation assays schizonts were isolated by percoll gradient [131] and incubated with fresh erythrocytes at 37°C for 4h followed by sorbitol synchronization [132] resulting in ring stage parasite cultures with a 4h time window.

For knock sideways experiments, the same culture was split into control and + 250 nM rapalog at the start of the proliferation assay and growth of each of these two cultures followed over the course of the assay. In order to determine parasitemia, parasite cultures were resuspended and 20 μL samples were transferred to an Eppendorf tube containing 80 μL RPMI. Hoechst-33342 and dihydroethidium (DHE) were added to obtain final concentrations of 5 μg/mL and 4.5 μg/mL, respectively. Samples were incubated for 20 min (protected from UV light) at room temperature, and parasitemia was determined using an LSRII flow cytometer by counting 100,000 events using the FACSDiva software (BD Biosciences).

Immunoblotting

Protein samples were resolved by SDS-PAGE and transferred to Amersham Protran membranes (GE Healthcare) in a tankblot device (Bio-Rad) using transfer buffer (0.192 M Glycine, 0.1% SDS, 25 mM Tris) with 20% methanol. Next, membranes were blocked for 30 minutes with 5% skim milk, and incubated with primary antibodies diluted in PBS containing 5% skim milk for 2h or overnight, followed by three washing steps with PBS and 2h incubation with horseradish peroxidase-conjugated secondary antibodies diluted in PBS containing 5% skim milk. Detection was performed using the Clarity Western ECL kit (Bio-Rad), and signals were recorded with a ChemiDoc XRS imaging system (Bio-Rad) equipped with Image Lab software 5.2 (Bio-Rad).

Antibodies were applied in the following dilutions: mouse a-GFP (1:1000) (Roche), rat a-HA (1:2000) (Roche), rabbit anti-aldolase (1.2000) [139], goat α-rat (1:2000) (Dianova), goat α-mouse (1:2000) (Dianova) and donkey α-rabbit (1:2000) (Dianova).

Conditional inactivation via knock-sideways

For knock-sideways cell lines were transfected with plasmids encoding the nuclear mislocaliser (NLS-FRB-mCherry) of the PPM mislocaliser (Lyn-FRB-mCherry) [73]. The knock-sideways approach was performed as described previously [73]. Briefly, cultures were split into two 2-ml cultures of which one was supplemented with 250 nM rapalog (Clontech). Mislocalisation of the target protein was verified by live-cell microscopy.

Ring stage survival assay (RSA)

RSAs were done as described previously [26,29]. Schizonts were purified from an asynchronous parasite culture using a percoll gradient [131] and were allowed to invade fresh RBCs at 37°C for 3 hours after which they were synchronised with 5% sorbitol [132] to obtain 0 - 3 hour old rings. These rings were washed 3 times with medium and challenged with 700 nM DHA for 6 hours. Afterwards, the cells were washed three times in RPMI medium and the parasites were grown for another 66 hours. Finally, Giemsa smears were prepared in order to determine the parasite survival rate (parasitemia of viable parasites after DHA compared to parasitemia of non-DHA treated control). A survival rate of 1% parasite was considered the threshold for *in vitro* ART resistance [26]. Number of cells counted are indicated in the corresponding Figure legends. Addition of 250 nM rapalog or cytochalasin D (10µM final concentration) were added before or with the start of the RSA as indicated in the corresponding schemes and figure legends and lifted after the DHA pulse as described [29].

Vesicle accumulation assay

The vesicle accumulation assay was adapted from [54]. Briefly, the number of vesicles per parasite were determined based on DIC images of synchronised trophozoites. For this, parasite cultures were two times synchronised (6-8 hours apart) using 5% sorbitol [132], split into two 2ml dishes of which one received rapalog to a final concentration of 250 nM, while the other culture served as control without rapalog and then grown for 16-24 hours to obtain trophozoite stages. Parasites were imaged in the DIC channel and the vesicles in the DIC images were counted. The assay was performed blinded and in at least three independent experiments (n of analysed cells indicated in the corresponding Figure legend).

Bloated food vacuole assay / E64 hemoglobin uptake assay

The bloated food vacuole assay was performed as previously described [54]. Briefly, ring stage parasite cultures with an 8h time window were obtained using double 5% sorbitol synchronisation. Parasites were either split into two 2ml dishes of which one received rapalog to a final concentration of 250 nM, while the other culture served as

control without rapalog for *Pf*KIC11 and *Pf*KIC12 (see experimental setup scheme in
Figure S4C, S5D) and incubated at 37°C overnight, first incubated at 37°C overnight
and split at the start of the assay for *Pf*MyoF (see experimental setup scheme in Figure
S3G), or not split in case for bloated food vacuole assays with TGD cell lines. When
parasites reached the young trophozoites stage, the medium was aspirated and 1 ml
medium containing 33 µM E64 protease inhibitor (Sigma Aldrich) was added. The cells
were cultured for 8 hours and then imaged. The DIC image was used for scoring
bloated food vacuoles and determination of parasite and food vacuole size. For
visualisation of bloated food vacuoles, the cells were stained with 4.5 mg/ml
dihydroethidium (DHE) for 15 minutes at 37°C. The experiment was performed blinded
and in at least three independent experiments.
950

Invasion and egress assay
The invasion/egress assay was adapted from previously published assays [68,74].
Briefly, KIC11-2xFKBP-GFP-2xFKBP late schizonts were purified using Percoll
followed by an incubation period of 30 minutes at 37°C in the shaking incubator (800
rpm) allowing the parasites to reinvade new erythrocytes. Subsequently, the cells were
transferred to static standard culture conditions, incubated for 3.5 hours and sorbitol
synchronization was performed resulting in tightly synchronized parasite culture with
a time window of 4h. Parasites were divided into two dishes and knock-sideways was
induced by addition of rapalog to one dish. Both dishes were cultured at 37°C until
Giemsa-stained smears were prepared at the time points 38-42 hpi ('pre-egress') and
46-50 hpi ('post-egress'). The numbers of ring and schizont stage parasites were
determined by counting parasites in randomly selected fields of sight and used to
calculate the percentage of ruptured schizonts and number of new rings per ruptured
schizont in the 'post-egress' samples.
965

Domain identification using AlphaFold
The predicted protein structures of all known K13-compartment members were
downloaded from the AlphaFold Protein Structure Database (alphafold.ebi.ac.uk) [87],
except for UBP1 which was not available. The structure for UBP1 was predicted in 6
parts, coving resides 1-640, 641-1280, 1281-1920, 1921-2560, 2561-2880 and 2881-
3499   using   ColabFold   [140].   VAST   searches
(ncbi.nlm.nih.gov/Structure/VAST/vastsearch.html) [141] were performed on all
structures. The top 3 hits for each protein and protein part were aligned with the search
model using the PyMol command cealign (Schrödinger, USA). Similarities with
RMSDs of under 4 Å over more than 60 amino acids are listed in the results. Domains
that were not previously annotated in Interpro (as of April 2022) [95] or PlasmoDB v.57
[142] were considered as newly identified.

Software

Statistical analyses were performed with GraphPad Prism version 8 (GraphPad Software, USA), microscopy images were processed in Corel Photo-Paint X6-X8 (https://www.coreldraw.com) or Fiji [143], plasmids and oligonucleotides were designed using ApE [144]. Protein structures were analysed and visualized using PyMol (Schrödinger, USA). Figures were arranged in CorelDraw X6-8.

Acknowledgements

We thank Jacobus Pharmaceuticals for WR99210. DSM1 (MRA-1161) was obtained from MR4/BEI Resources, NIAID, NIH. We thank Tim Gilberger for providing the PhIL1mcherry plasmid.

Author contribution

Conceptualization: TS

Validation: CCP

Formal Analysis: JSWM, SS, HB, TS

Investigation: JSWM, SS, HB, JB, IH, JD, EJ, SS, SF

Resources: JB, EJ, SF, PMR, TS

Writing – original Draft: JSWM, HB, TS

Writing – Review & Editing: JSWM, HB, IH, TS

Visualization: JSWM, SS, HB, IH, TS

Supervision: TS

Funding Acquisition: TS

All authors read and approved the manuscript.

The authors declare that they have no conflict of interest.

Funding

HMB acknowledges funding by the Ortrud Mührer Fellowship of the Vereinigung der Freunde des Tropeninstituts Hamburg e.V, CCP thanks DAAD for funding. This work was funded by the European Research Council (ERC, grant 101021493).

Figures:

Table 1: Selection of putative K13 compartment proteins for further characterization in this manuscript

Figure 1: MyoF is involved in host cell cytosol uptake and associated with the K13 compartment

(A) Localisation of MyoF-2xFKBP-GFP-2xFKBP expressed from the endogenous locus by live-cell microscopy across the intra-erythrocytic development cycle. Nuclei were stained with DAPI. Scale bar, 5 μm. **(B)** Live cell microscopy images of parasites expressing the MyoF-2xFKBP-GFP-2xFKBP fusion protein with an episomally expressed FV marker P40PX-mCherry. Scale bar, 5 μm. **(C)** Live cell microscopy images of parasites expressing the MyoF-2xFKBP-GFP-2xFKBP fusion protein with episomally expressed mCherry-K13. Scale bar, 5 μm. Arrows indicate categories from Figure 1D **(D)** Foci were categorized into 'overlap' (black), 'partial overlap' (dark grey), 'close foci' (=less than one focus radius apart) (grey) and 'non overlap' (light grey). Three independent live microscopy sessions with each n=14 analysed parasites. **(E)** Relative growth of synchronised MyoF-2xFKBP-GFP-2xFKBP^{endo} compared with 3D7 wild type parasites after two growth cycles. Each dot shows one of four independent growth experiments. P-values determined by one-sample t-test. **(F)** Live-cell microscopy of knock sideways (rapalog) and control MyoF-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser parasites 1 hour or 22 hours after the induction by addition of rapalog. Scale bar, 5 μm. **(G)** Relative growth of asynchronous 3D7 and asynchronous MyoF-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser parasites (+ rapalog) compared with the corresponding control parasites (without rapalog) after five days. Three (3D7) and eight (MyoF-2xFKBP-GFP-2xFKBP^{endo}) independent experiments (individual experiments shown in Figure S2D). Error bars, mean ± SD. P-values determined by Welch's t-test. **(H)** Number of vesicles per parasite in trophozoites determined by live-cell fluorescence microscopy (DIC) in 3D7 and MyoF-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser parasites with and without rapalog addition. Three (3D7) and six (MyoF-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser) independent experiments with each time n=16-25 (mean 20.9) parasites analysed per condition. Mean of each independent experiment indicated by coloured triangle, individual datapoints by grey dots. Data presented according to SuperPlot guidelines [145]; Error bars represent mean ± SD. P-value for ± rapalog determined by paired t-test and for 3D7 vs MyoF by Mann-Whitney. Representative DIC images are displayed below. **(I)** Bloated food vacuole assay with MyoF-2xFKBP-GFP-2xFKBP^{endo} parasites 8 hours after inactivation of MyoF (+ rapalog) compared with controls (- rapalog). Cells were categorized as with 'bloated FV' or 'non-bloated FV' and percentage of cells with bloated FV is displayed; n = 4 independent experiments with each n=33-40 (mean 34.6) parasites analysed per condition. P-values determined by Welch's t-test. Representative DIC and fluorescence microscopy images shown on the right. Parasite cytoplasm was visualized with DHE. Experimental setup shown in Figure S2G. **(J)** Area of the FV, area of the parasite and area of FV divided by area of the corresponding parasite of MyoF-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser parasites analysed in Figure 1I. Mean of each independent experiment indicated by

coloured triangle, individual data points by grey dots. Data presented according to
SuperPlot guidelines [145]; Error bars represent mean ± SD. P-value determined by
paired t-test. Representative DIC and fluorescence microscopy images are shown in
the Figure S2J. **(K)** Area of FV of individual cells plotted versus the area of the
corresponding parasite in MyoF-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser
parasites of the experiments shown in Figure 1H-I. Line represents linear regression
with error indicated by dashed line. **(L)** Summary heatmap showing the effect of C-
terminal tagging with 3xHA or 2xFKBP-GFP-2xFKBP and knock-sideways on MyoF
functionality depicted as growth [%], bloated FV [%] and number of induced vesicles.
(M) Parasite survival rate (% survival compared to control without DHA) 66 h after 6 h
DHA (700 nM) treatment in standard RSA. MyoF-2xFKBP-GFP-2xFKBP^{endo} parasites
+ 1xNLSmislocaliser were pre-treated with rapalog either 12 or 24 hours prior to the
assay. Three independent experiments, P-value determined by Wilcoxon test. Green
dashed line indicates 1% ART resistance cut-off [26]. 2988-8392 (mean 5102) cells
for control and 22704-44038 (mean 32077) cells for DHA treated samples were
counted. Experimental setup shown in Figure S2K. 3D7 and 3D7-K13^{C580Y}[29]
parasites were incubated with CytochalasinD for 6h in parallel with DHA pulse. 850-
13180 (mean 4677) cells for control and 1065-38515 (mean 15532) cells for DHA
treated samples were counted. Three (3D7) or five (3D7-K13^{C580Y}) independent
experiments, P-value determined by Wilcoxon test.
1076
Figure 2: KIC11 is a K13 compartment protein important for asexual parasite
proliferation, but not involved in endocytosis or *in vitro* ART resistance
(A) Localisation of KIC11-2xFKBP-GFP-2xFKBP expressed from the endogenous
locus by live-cell microscopy across the intra-erythrocytic development cycle. Nuclei
were stained with DAPI. Scale bar, 5 µm. (**B)** Live cell microscopy images of parasites
expressing KIC11-2xFKBP-GFP-2xFKBP with episomally expressed mCherry-K13.
Nuclei were stained with DAPI. Scale bar, 5 µm. Arrows indicate categories from
Figure 2C. **(C)** Foci were categorized into 'overlap' (black), 'partial overlap' (dark grey),
'close foci' (=less than one focus radius apart) (grey) and 'non overlap' (light grey) for
ring (n=35) and trophozoite (n=25) stage parasites. **(D)** Live-cell microscopy of knock
sideways (+ rapalog) and control (without rapalog) KIC11-2xFKBP-GFP-
2xFKBP^{endo}+1xNLSmislocaliser parasites 4 and 16 hours after the induction of knock-
sideways by addition of rapalog. Scale bar, 5 µm. **(E)** Relative growth of asynchronous
KIC11-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser plus rapalog compared with
control parasites over five days. Five independent experiments were performed
(depicted by different symbols) and mean relative parasitemia ± SD is shown
(individual experiments shown in Figure S4A). **(F)** Parasite stage distribution in
Giemsa smears at the time points (average hours post invasion, h) indicated above

each bar in tightly synchronised (±4h) KIC11-2xFKBP-GFP-
2xFKBP^{endo}+1xNLSmislocaliser parasites (rapalog addition at 4 hpi, 20 hpi, or 32 hpi
and control) assayed over two consecutive cycles (last time point in cycle 3). A second
replicate is shown in Figure S3F. **(G)** Giemsa smears of control and at 4 hpi, 20 hpi,
or 32 hpi rapalog-treated KIC11-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser
parasites shown in (E). **(H)** Quantification of rings per ruptured schizont at 'post-
egress' time point compared to 'pre-egress' time point in knock-sideways (+rapalog)
compared to control (-rapalog) KIC11-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser
parasites from n=3 independent experiments (indicated by different symbol shapes).
P values were determined with ratio-paired t-test, data displayed as mean ±SD.
Quantification of corresponding ruptured schizonts is shown in Figure S3G. **(I)** Number
of vesicles per parasite in trophozoites determined by live-cell fluorescence
microscopy (DIC) in KIC11-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser parasites
with and without rapalog. Four independent experiments with n=16-59 (mean 30.1)
parasites analysed per condition per experiment. Mean of each independent
experiment indicated by coloured triangle, individual datapoints by grey dots. Data
presented according to SuperPlot guidelines [145]; Error bars represent mean ± SD.
P-value determined by paired t-test. Representative DIC images are displayed. **(J)**
Bloated food vacuole assay with KIC11-2xFKBP-GFP-2xFKBP^{endo} parasites 8 hours
after inactivation of KIC11 (+rapalog) compared with control (without rapalog). Cells
were categorized as with 'bloated FV' or 'non-bloated FV' and displayed as percentage
of cells with bloated FV; n=3 independent experiments with each n=19-36 (mean 26.5)
parasites analysed per condition. P-values determined by Welch's t-test.
Representative DIC and fluorescence microscopy images are shown in the right panel.
Parasite cytoplasm was visualized with DHE. Experimental setup shown in Figure
S3H. **(K)** Parasite survival rate (% survival compared to control without DHA) 66 h
after 6 h DHA treatment in standard RSA. Three independent experiments, P-value
determined by paired t-test. Green dashed line indicates 1% ART resistance cut-off
[26]. 2896-7135 (mean 4502) cells for control and 23183-32455 (mean 28496) cells
for DHA treated samples were counted. Experimental setup shown in Figure S3I.
1125
Figure 3: KIC12 shows a dual localisation in the nucleus and at the K13
compartment and is involved in endocytosis but not in ART resistance
(A) Localisation of KIC12-2xFKBP-GFP-2xFKBP expressed from the endogenous
locus by live-cell microscopy across the intra-erythrocytic development cycle. Arrow
heads indicate foci (white) and nuclear (light blue) signal. Nuclei were stained with
Hoechst. Scale bar, 5 μm. **(B)** Live cell microscopy images of parasites expressing
KIC12-2xFKBP-GFP-2xFKBP with episomally expressed mCherry-K13. Scale bar, 5
μm. Black arrowheads indicate overlapping foci. Nuclei were stained with DAPI. Scale

bar, 5 µm. Extended panel shown in Figure S4A. **(C)** Live-cell microscopy of knock

sideways (+ rapalog) and control (without rapalog) KIC12-2xFKBP-GFP-

2xFKBP^{endo}+1xNLSmislocaliser parasites 4 and 16 hours after the induction of knock-

sideways by addition of rapalog. Scale bar, 5 µm. **(D)** Relative growth of asynchronous

KIC12-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser (blue) and KIC12-2xFKBP-

GFP-2xFKBP^{endo}+Lyn-mislocaliser (red) parasites plus rapalog compared with the

corresponding untreated control parasites over five days. Five independent growth

experiments were performed and mean relative parasitemia +/- SD is shown (all

replicas shown in Figure S4C and D). **(E)** Live-cell microscopy of knock sideways (+

rapalog) and control (without rapalog) KIC12-2xFKBP-GFP-2xFKBP^{endo}+Lyn-

mislocaliser parasites 4 hours after the induction of knock-sideways by addition of

rapalog. Scale bar, 5 µm. **(F)** Bloated food vacuole assay with KIC12-2xFKBP-GFP-

2xFKBP^{endo} parasites 8 hours after inactivation of KIC12 (+rapalog) by NLS-

mislocaliser or Lyn-mislocaliser compared with corresponding control (without

rapalog). Cells were categorized as 'bloated FV' and 'non-bloated FV'. Results are

displayed as percentage of cells with bloated FV. n=3 (NLS) or n=4 (LYN) independent

experiments were performed with each n=19-80 (mean 41.3) parasites analysed per

condition. P-values determined by Welch's t-test. Representative DIC and

fluorescence microscopy images are shown in the right panel. Parasite cytoplasm was

visualized with DHE. Experimental setup shown in Figure S4E. Scale bar, 5 µm. **(G/I)**

Area of the FV, area of the parasite and area of FV divided by area of the

corresponding parasite of the FV of KIC12-2xFKBP-GFP-

2xFKBP^{endo}+1xNLSmislocaliser (G) and KIC12-2xFKBP-GFP-

2xFKBP^{endo}+LYNmislocaliser (I) parasites analysed in 4F. Mean of each independent

experiment indicated by coloured triangle, individual datapoints by grey dots. Data

presented according to SuperPlot guidelines [145]; Error bars represent mean ± SD.

P-value determined by paired t-test. **(H/J)** Area of FV of individual cells plotted versus

the area of the corresponding parasite in KIC12-2xFKBP-GFP-

2xFKBP^{endo}+1xNLSmislocaliser and KIC12-2xFKBP-GFP-2xFKBP^{endo}+Lyn-

mislocaliser parasites of the experiments shown in (F,G and I). Line represents linear

regression with error indicated by dashed line. Ten representative DIC images of each

independent experiment are shown in Figure S4F,G **(K)** Number of vesicles per

parasite in trophozoites determined by live-cell fluorescence microscopy (DIC) in

KIC12-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser and KIC12-2xFKBP-GFP-

2xFKBP^{endo}+Lyn-mislocaliser parasites with and without rapalog addition. Three

independent experiments with n=10-56 (mean 28) parasites analysed per condition

and experiment Data presented according to SuperPlot guidelines [145]; Mean of each

independent experiment indicated by coloured triangle, individual data points by grey

dots. Error bars represent mean ± SD. P-value determined by paired t-test.

Representative DIC images are displayed. **(L)** Parasite survival rate of KIC12-
2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser (% survival compared to control
without DHA) 66 h after 6 h DHA treatment in standard RSA. Three independent
experiments, P-value determined by paired t-test. Green dashed line indicates 1%
ART resistance cut-off [26]. 3151-4273 (mean 3690) cells for control and 6209-18941
(mean 12290) cells for DHA treated samples were counted. Experimental setup shown
in Figure S4H.
1180
Figure 4: MCA2 is part of the K13 compartment and its truncation reduces ART
susceptibility
(A) Schematic representation and nomenclature of MCA2 fusion proteins (not to scale)
expressed in the parasite by modification of the endogenous gene. GFP in green,
3xHA in red and metacaspsase domain in grey. Amino acid positions are indicated by
numbers. **(B)** Localisation of MCA2^{Y1344STOP}-GFP by live-cell microscopy across the
intra-erythrocytic development cycle. Nuclei were stained with DAPI. Scale bar, 5 µm.
(**C**) Live cell microscopy images of parasites expressing the truncated MCA2^{Y1344STOP}-
GFP fusion protein with an episomally expressed mCherry-K13 fusion protein. Foci
from rings and trophozoite stage parasites were categorized into 'overlap' (black),
'partial overlap' (dark grey) and 'no overlap' (light grey) and shown as frequencies in
the pie chart (n = 46 cells were scored from a total of three independent experiments).
Scale bar, 5 µm. **(D)** Live cell microscopy images of parasites expressing the truncated
MCA2^{Y1344STOP}-GFP fusion protein with the IMC marker protein PhIL1mCherry. Zoom,
enlarged region (factor 400%). Nuclei were stained with Hoechst-33342; scale bars, 5
µm and for zoom 1 µm. **(E)** Relative growth of synchronised MCA2^{Y1344STOP} compared
with 3D7 wild type parasites after two growth cycles. Each dot shows one of four
independent experiments. P-values determined by one-sample t-test. **(F)** Parasite
survival rate (% survival compared with control without DHA) 66 h after 6 h DHA
treatment in standard RSA. Two (3D7) or three (MCA2^{Y1344STOP}) independent
experiments, p-value determined by unpaired t-test. Green dashed line indicates 1%
ART resistance cut-off [26]; 3419-3828 (mean 3632) erythrocytes for control and
10364-11734 (mean 11254) cells for DHA treated samples were counted.
1204
Figure 5: Type of domain found in K13 compartment proteins coincide with
functional group
(A) K13-compartment members shown to scale with domains indicated that were
identified from AlphaFold structure predictions. Newly identified domains (i.e. domains
not previously identified by sequence homology) are shown in colours, previously
known domains are shown in grey. A brief summary of the function of each newly
identified domain and Interpro IDs of all domains can be found in supplementary figure

S8. **(B)** Alphafold prediction of each newly identified domain is shown in the same colour as in A and aligned with the most similar domain from the PDB. PDB ID and alignment details are indicated beneath each set of aligned domains. Root mean square deviations (RMSD) for all comparisons are given. **(C)** Domain organizations of human GGA1 (hGGA1), KIC4 and human AP-2 subunit beta 1 (hAP2B1) are shown (not to scale). Domains are coloured as in A and B. **(D)** Domain organizations of KIC5 and human AP-2 subunit alpha 1 (hAP2A1) are shown (not to scale). Domains are coloured as in A, B and C. (**E**) Table summarising K13 compartment proteins according to property (presence of vesicle trafficking domain (VTD)) and function.

Figure S1: Validation of generated transgenic cell lines

Confirmatory PCR of unmodified wildtype (WT) and transgenic knock-in (KI) / targeted-gene-disruption (TGD) cell lines to confirm correct genomic integration at the 3'- and 5'-end of the locus. Oligonucleotides used are listed in Table S2. **(A)** MCA2^{Y1344STOP}-GFP^{endo}; **(B)** MCA2-3xHA^{endo}; **(C)** MyoF-2xFKBP-GFP-2xFKBP^{endo}; **(D)** MyoF-2xFKBP-GFP^{endo}; **(E)** MyoF-3xHA^{endo}; **(F)** KIC11-2xFKBP-GFP-2xFKBP^{endo}; **(G)** KIC12-2xFKBP-GFP-2xFKBP^{endo}; **(H)** PF3D7_0907200-2xFKBP-GFP-2xFKBP^{endo}; **(I)** VPS51-2xFKBP-GFP-2xFKBP^{endo}; **(J)** PF3D7_1365800-2xFKBP-GFP-2xFKBP^{endo}; **(K)** UIS14-2xFKBP-GFP-2xFKBP^{endo}; **(L)** PF3D7_1447800-2xFKBP-GFP-2xFKBP^{endo}; **(M)** VPS51-TGD^{endo}; **(N)** UIS14-TGD^{endo}. Right panel in **(F and G)** Western Blot analysis of **(F)** MCA2^{Y1344STOP}-GFP^{endo} cell line using mouse anti-GFP to detect the tagged fusion protein (upper panel) and rabbit anti-aldolase to control for equal loading (lower panel)(expected molecular weight for MCA2^{Y1344STOP}-GFP fusion proteins: 187 kDa) and **(G)** wildtype (3D7) and knock-in MCA2-3xHA^{endo} cell line using mouse anti-HA to detect the tagged full-length protein (upper panel) and rabbit anti-aldolase to control for equal loading (lower panel) (expected molecular weight for MCA2-3xHA fusion protein: 281 kDa). Protein size is indicated in kDa.

Figure S2: Additional data for MyoF

(A) Extended panel of live cell microscopy images of parasites expressing the MyoF-2xFKBP-GFP-2xFKBP fusion protein from the endogenous locus with episomally expressed mCherry-K13. Nuclei were stained with Hoechst. Scale bar, 5 µm. **(B)** Localisation of MyoF-2xFKBP-GFP by live-cell microscopy across the intra-erythrocytic development cycle. Nuclei were stained with DAPI. Scale bar, 5 µm. **(C)** IFA microscopy images of acetone-fixed parasites expressing MyoF-3xHA with episomally expressed mCherry-K13 across the intra-erythrocytic development cycle. Nuclei were stained with DAPI. Scale bar, 5µm. Foci were categorized into 'overlap' (black), 'partial overlap' (dark grey), close foci (light blue) and 'non overlap' (light grey) in n=31 parasites. Scale bar, 5µm. **(D)** Live-cell microscopy of knock sideways (+

rapalog) and control (without rapalog) MyoF-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser parasites at 0, 1, 2, 4 and 22 hours after the induction of knock-sideways by addition of rapalog. **(E)** Individual growth curves of MyoF-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser with (red) or without (blue) addition of rapalog shown in Figure 1F. Summary of individual growth curves shown as percentage of control parasites (without rapalog). Three (3D7, black) and eight (MyoF-2xFKBP-GFP-2xFKBP^{endo}, red) independent experiments. Error bars, mean ± SD. **(F)** Relative growth of synchronised MyoF-3xHA^{endo} compared with 3D7 wild type parasites after two growth cycles. Each dot shows one of six independent growth experiments. P-values determined by one-sample t-test. **(G)** Number of vesicles per parasite in trophozoites determined by live-cell fluorescence microscopy (DIC) in 3D7 and MyoF-3xHA^{endo} parasites. Three independent experiments with each time n=27-38 (mean 31.7) parasites analysed per condition. Representative DIC images shown on the right. **(H)** Experimental setup of the bloated food vacuole assay shown in Figure 1H. **(I)** Bloated food vacuole assay with 3D7 parasites 8 hours after rapalog addition compared with controls (- rapalog). Cells were categorized as with 'bloated FV' or 'non-bloated FV' and percentage of cells with bloated FV is displayed; n = 2 independent experiments with each n=26-40 (mean 31,3) parasites analysed per condition. Representative DIC and fluorescence microscopy images are shown. Parasite cytoplasm was visualized with DHE. **(J)** Representative images of bloated food vacuole assays with knock sideways (+ rapalog; bottom row) and control (without rapalog; top row) MyoF-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser parasites shown in Figure 1I-J. Scale bar, 5 µm. **(K)** Experimental setup of the RSA shown in Figure 1I. Schematic representation of the cell lines depicted above the corresponding panel.

Figure S3: Additional data for KIC11

(A) Live-cell microscopy of knock sideways (+ rapalog) and control (without rapalog) KIC11-2xFKBP-GFP-2xFKBP^{endo} ^{ama1}IMC1c-mSca_$nmd3$'-NLS-FRB-T2A-DHODH, KIC11-2xFKBP-GFP-2xFKBP^{endo}+ ^{ama1}ARO-mSca_$nmd3$'-NLS-FRB-T2A-DHODH or KIC11-2xFKBP-GFP-2xFKBP^{endo}+ ^{ama1}AMA1-mSca_$nmd3$'-NLS-FRB-T2A-DHODH parasites 40 hours after the induction of knock-sideways by addition of rapalog. Nuclei were stained with Hoechst; scale bar, 5 µm. **(B)** Quantification of KIC11 and K13 foci per cell in ring, trophozoite and schizont stage KIC11-2xFKBP-GFP-2xFKBP^{endo} with episomally expressed mCherry-K13 parasites. N indicates the number of analysed KIC11-2xFKBP-GFP-2xFKBP parasites with episomally expressed mCherry-K13 per stage. **(C)** Extended panel of live cell microscopy images of parasites expressing KIC11-2xFKBP-GFP-2xFKBP^{endo} with episomally expressed mCherry-K13 shown in Figure 2B. Nuclei were stained with DAPI. Scale bar, 5 µm except last column (zoom

merozoite image): Scale bar = 1.25µm. Arrows are indicating categories from Figure
2C **(D)** Individual growth curves of knock sideways (+ rapalog) and control (without
rapalog) of KIC11-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser shown in Figure 2E.
(E) Live-cell microscopy of knock sideways (+ rapalog) and control (without rapalog)
KIC11-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser parasites 80 hours after the
induction of knock-sideways by addition of rapalog. Scale bar, 5 µm. **(F)** Parasite stage
distribution in Giemsa smears at the time points (average hours post invasion, h)
indicated above each bar in tightly synchronised (±4h) KIC11-2xFKBP-GFP-
2xFKBP^{endo}+1xNLSmislocaliser parasites (rapalog addition at 4 hpi, 20 hpi, or 32 hpi
and control) parasite cultures over two consecutive cycles (last time point in cycle 3).
A second replicate is shown in Figure 2F. **(G)** Quantification of ruptured schizonts at
'post-egress' time point compared to 'pre-egress' time point in knock-sideways
(+rapalog) compared to control (-rapalog) in KIC11-2xFKBP-GFP-
2xFKBP^{endo}+1xNLSmislocaliser parasites from n=3 independent experiments
(indicated by different symbol shapes). P values were determined with ratio-paired t-
test, data displayed as mean ±SD. Corresponding quantification of ring per ruptured
schizont is shown in Figure 2H. **(H)** Experimental setup of the bloated food vacuole
assay shown in Figure 2J. **(I)** Experimental setup of the RSA shown in Figure 2K.
Schematic representation of the cell lines depicted above the corresponding panel.
1309
Figure S4: Additional data for KIC12
(A) Extended panel of KIC12-2xFKBP-GFP-2xFKBP^{endo} localisation by live-cell
microscopy across the intra-erythrocytic development cycle. Nuclei were stained with
Hoechst. Scale bar, 5 µm. Expanded panel of Figure 3B Foci were categorized into
'overlap' (black), 'partial overlap' (grey) and 'no overlap' (white) and shown as
frequencies in the pie chart (n = 97 cells were scored from a total of five independent
experiments). **(B)** Absence of KIC12-2xFKBP-GFP-2xFKBP^{endo} in free merozoites by
live-cell microscopy. Nuclei were stained with Hoechst. Scale bar, 5 µm. **(C)** Individual
growth curves of knock sideways (+ rapalog; red) and control (without rapalog; blue)
KIC12-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser shown in Figure 3D. **(D)**
Individual growth curves of knock sideways (+ rapalog; red) and control (without
rapalog; blue) KIC12-2xFKBP-GFP-2xFKBP^{endo}+LYNmislocaliser shown in Figure 3D.
(E) Experimental setup of the bloated food vacuole assay shown in Figure 3F. **(F)**
Representative images of bloated food vacuole assay knock sideways (+ rapalog;
bottom row) and control (without rapalog; top row) KIC12-2xFKBP-GFP-
2xFKBP^{endo}+1xNLSmislocaliser Scale bar, 5 µm. **(G)** Representative images of knock
sideways (+ rapalog, top row) and control (without rapalog; bottom row) KIC12-
2xFKBP-GFP-2xFKBP^{endo}+LYNmislocaliser parasites from bloated food vacuole

assay. Scale bar, 5 µm. **(H)** Experimental setup of the RSA shown in Figure 3L.
Schematic representation of the cell lines depicted above the corresponding panel.
1330
Figure S5: Additional data for MCA2
(A) Localisation of MCA2-TGD-GFP by live-cell microscopy across the intra-
erythrocytic development cycle. Nuclei were stained with DAPI. Scale bar, 5 µm. **(B)**
IFA microscopy images of acetone-fixed parasites expressing MCA2-3xHA with
episomally expressed mCherry-K13 across the intra-erythrocytic development cycle.
Nuclei were stained with DAPI. Scale bar, 5µm. Foci were categorized into 'overlap'
(black), 'partial overlap' (dark grey) and 'no overlap' (light grey) in n=30 parasites.
Schematic representation of the cell lines depicted above the corresponding panel.
(C) Bloated food vacuole assay with 3D7, MCA2^{Y1344STOP} and MCA2-TGD parasites.
Cells were categorized as with 'bloated FV' or 'non-bloated FV' and percentage of
cells with bloated FV is displayed; n = 4 (3D7, MCA2^{Y1344STOP}) or n=3 (MCA2-TGD)
independent experiments with each n=10-34 (mean 25.8) parasites analysed per
condition. **(D)** Area of the FV, area of the parasite and area of FV divided by area of
the corresponding parasites were determined. Mean of each independent experiment
indicated by coloured symbols, individual data points by grey dots. Data presented
according to SuperPlot guidelines [145]; Error bars represent mean ± SD. P-value
determined by un-paired t-test. Area of FV of individual cells plotted versus the area
of the corresponding parasite. Line represents linear regression with error indicated
by dashed line. **(E)** Representative DIC images from S2C-D are displayed. Schematic
representation of the cell lines depicted above the corresponding panel.
1351
Figure S6: Candidate proteins not associated with the K13 compartment
(A) Live cell microscopy images of parasites endogenously expressing VPS51-
2xFKBP-GFP-2xFKBP with episomally expressed mCherry-K13 across the intra-
erythrocytic development cycle. Scale bar, 5 µm. **(B)** Localisation of truncated
VPS51TGD-GFP fusion protein by live-cell microscopy across the intra-erythrocytic
development cycle. Nuclei were stained with DAPI. Scale bar, 5 µm. Schematic
representation of the truncation strategy depicted above the panel, numbers indicating
AA. **(C)** Relative growth of VPS51TGD and UIS14TGD parasites compared to 3D7
wild type parasites after two cycles. Four independent growth experiments. P-values
determined by one-sample t-test. **(D)** Live cell microscopy images of parasites
endogenously expressing UIS14-2xFKBP-GFP-2xFKBP with episomally expressed
mCherry-K13 across the intra-erythrocytic development cycle. Scale bar, 5 µm. **(E)**
Localisation of truncated UIS14TGD-GFP fusion protein by live-cell microscopy across
the intra-erythrocytic development cycle. Nuclei were stained with DAPI. Scale bar, 5
µm. Schematic representation of the truncation strategy depicted above the panel,

numbers indicating AA. **(F)** Live cell microscopy images of parasites endogenously
expressing PF3D7_1447800-2xFKBP-GFP-2xFKBP with episomally expressed
mCherry-K13 across the intra-erythrocytic development cycle. Nuclei were stained
with DAPI. Scale bar, 5 µm. **(G)** Live cell microscopy images of parasites
endogenously expressing PF3D7_0907200-2xFKBP-GFP-2xFKBP with episomally
expressed mCherry-K13 across the intra-erythrocytic development cycle. Scale bar, 5
µm. Schematic representation of relevant features of each cell line depicted above
the corresponding panel.
1375

Figure S7: 3D7_1365800 an AH domain containing protein is dispensable for
asexual parasite development
(A) Live cell microscopy images of parasites endogenously expressing
PF3D7_1365800-2xFKBP-GFP-2xFKBP by live-cell microscopy across the intra-
erythrocytic development cycle. Nuclei were stained with DAPI. Scale bar, 5 µm. **(B)**
Expression of PF3D7_1365800-2xFKBP-GFP-2xFKBP with episomally expressed
mCherry-K13. Scale bar, 5 µm. **(C)** Live-cell microscopy of knock sideways (+ rapalog)
and control (without rapalog) PF3D7_1365800-2xFKBP-GFP-
2xFKBP^{endo}+1xNLSmislocaliser parasites or PF3D7_1365800-2xFKBP-GFP-
2xFKBP^{endo}+LYNmislocaliser parasites 16 hours after the induction of knock-sideways
by addition of rapalog. Scale bar, 5 µm. **(D)** Individual growth curves of
PF3D7_1365800-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser or PF3D7_1365800-
2xFKBP-GFP-2xFKBP^{endo}+LYNmislocaliser parasites with (red) or without (blue)
addition of rapalog. Relative growth of knock sideways (+ rapalog) compared to control
(without rapalog) PF3D7_1365800-2xFKBP-GFP-2xFKBP^{endo}+1xNLSmislocaliser
(blue) or PF3D7_1365800-2xFKBP-GFP-2xFKBP^{endo}+LYNmislocaliser (red)
parasites over five days. Three (NLSmisloc) or six (LYNmisloc) independent growth
experiments were performed and mean relative parasitemia ± SD is shown. **(E)** Live
cell microscopy images of parasites expressing PF3D7_1365800-2xFKBP-GFP-
2xFKBP with episomally expressed Golgi marker GRASP-mCherry. Schematic
representation of the cell lines depicted above the corresponding panel. Scale bar, 5
µm.
1398

Figure S8: Type of domain found in K13 compartment proteins coincide with
functional group
(A) Full domain names and Interpro domain numbers for each domain in Figure 5 with
indication in which K13-compartment members these proteins occur. For each newly
identified domain a brief summary of its reported function is given. Colours are as in
Figure 5. **(B)** Full length AlphaFold predictions for each K13-compartment member in
which new domains were identified. New domains are coloured as in A and Figure 5.

For UBP1 no prediction was available in the EMBL-AlphaFold database, and the structure was predicted in fragments as described in the methods. The predicted fragment containing the newly identified domain is shown. **(C)** Full length structure of PF3D7_1365800, with AH domain in green. PF3D7_1365800 AH domain in green aligned with the most similar domain from the PDB. PDB ID and alignment details are indicated beneath each set of aligned domains. Brief summary about the AH domain and its Interpro ID are given.

Figure S9: Targeted gene disruption of KIC4 and KIC5 has no effect in bloated food vacuole assays

Bloated food vacuole assay with 3D7, KIC4-TGD and KIC5-TGD parasites. Cells were categorized as with 'bloated FV' or 'non-bloated FV' and percentage of cells with bloated FV is displayed; n = 3 independent experiments with each n=20-38 (mean 26.7) parasites analysed per condition. Representative DIC are displayed. Area of the FV, area of the parasite and area of FV divided by area of the corresponding parasites were determined. Mean of each independent experiment indicated by coloured symbols, individual data points by grey dots. Data presented according to SuperPlot guidelines [145]; Error bars represent mean ± SD. P-value determined by paired t-test. Area of FV of individual cells plotted versus the area of the corresponding parasite. Line represents linear regression with error indicated by dashed line.

Table S1: List of putative K13 compartment proteins, with proteins selected for further characterization in this manuscript highlighted.

Ranked by enrichment: from high to low by the average log2 ratio normalized of four published Kelch13 BioID experiments [29].

Table S2: Oligonucleotides used for cloning and diagnostic genotyping PCR

References

1.	WHO. World Malaria Report 2021. 2021.

2.	Weiss DJ, Lucas TCD, Nguyen M, Nandi AK, Bisanzio D, Battle KE, et al. Mapping the global prevalence, incidence, and mortality of Plasmodium falciparum, 2000–17: a spatial and temporal modelling study. Lancet. 2019;394: 322–331. doi:10.1016/S0140-6736(19)31097-9

3.	WHO. WHO Guidelines for malaria. WHO Guidelines for malaria. 2021.

4.	Dondorp AM, Nosten F, Yi P, Das D, Phyo AP, Tarning J, et al. Artemisinin Resistance in Plasmodium falciparum Malaria. N Engl J Med. 2009;361: 455–467. doi:10.1056/NEJMoa0808859

5.	Ashley EA, Dhorda M, Fairhurst RM, Amaratunga C, Lim P, Suon S, et al. Spread of Artemisinin Resistance in *Plasmodium falciparum* Malaria. N Engl J Med. 2014;371: 411–423. doi:10.1056/NEJMoa1314981

6.   Amaratunga C, Sreng S, Suon S, Phelps ES, Stepniewska K, Lim P, et al.
Artemisinin-resistant Plasmodium falciparum in Pursat province, western
Cambodia: A parasite clearance rate study. Lancet Infect Dis. 2012;12: 851–
858. doi:10.1016/S1473-3099(12)70181-0

7.   Duru V, Khim N, Leang R, Kim S, Domergue A, Kloeung N, et al. Plasmodium
falciparum dihydroartemisinin-piperaquine failures in Cambodia are associated
with mutant K13 parasites presenting high survival rates in novel piperaquine
in vitro assays: Retrospective and prospective investigations. BMC Med.
2015;13. doi:10.1186/s12916-015-0539-5

8.   Woodrow CJ, White NJ. The clinical impact of artemisinin resistance in
Southeast Asia and the potential for future spread. FEMS Microbiology
Reviews. Oxford University Press; 2017. pp. 34–48.
doi:10.1093/femsre/fuw037

9.   Spring MD, Lin JT, Manning JE, Vanachayangkul P, Somethy S, Bun R, et al.
Dihydroartemisinin-piperaquine failure associated with a triple mutant including
kelch13 C580Y in Cambodia: An observational cohort study. Lancet Infect Dis.
2015;15: 683–691. doi:10.1016/S1473-3099(15)70049-6

10. Noedl H, Se Y, Schaecher K, Smith BL, Socheat D, Fukuda MM. Evidence of
Artemisinin-Resistant Malaria in Western Cambodia. N Engl J Med. 2008;359:
2619–2620. doi:10.1056/NEJMc0805011

11. Wasakul V, Disratthakit A, Mayxay M, Chindavongsa K, Sengsavath V, Thuy-
Nhien N, et al. Malaria outbreak in Laos driven by a selective sweep for
Plasmodium falciparum kelch13 R539T mutants: a genetic epidemiology
analysis. Lancet Infect Dis. 2022;0. doi:10.1016/S1473-3099(22)00697-1

12. Phyo AP, Nkhoma S, Stepniewska K, Ashley EA, Nair S, McGready R, et al.
Emergence of artemisinin-resistant malaria on the western border of Thailand:
a longitudinal study. Lancet. 2012;379: 1960–1966. doi:10.1016/S0140-
6736(12)60484-X

13. Uwimana A, Legrand E, Stokes BH, Ndikumana JLM, Warsame M, Umulisa N,
et al. Emergence and clonal expansion of in vitro artemisinin-resistant
Plasmodium falciparum kelch13 R561H mutant parasites in Rwanda. Nat Med.
2020;26. doi:10.1038/s41591-020-1005-2

14. Uwimana A, Umulisa N, Venkatesan M, Svigel SS, Zhou Z, Munyaneza T, et
al. Association of Plasmodium falciparum kelch13 R561H genotypes with
delayed parasite clearance in Rwanda: an open-label, single-arm, multicentre,
therapeutic efficacy study. Lancet Infect Dis. 2021;21: 1120–1128.
doi:10.1016/S1473-3099(21)00142-0

15. Stokes BH, Dhingra SK, Rubiano K, Mok S, Straimer J, Gnädig NF, et al.
Plasmodium falciparum k13 mutations in africa and asia impact artemisinin
resistance and parasite fitness. Elife. 2021;10. doi:10.7554/eLife.66277

16. Tumwebaze PK, Conrad MD, Okitwi M, Orena S, Byaruhanga O, Katairo T, et
al. Decreased susceptibility of Plasmodium falciparum to both
dihydroartemisinin and lumefantrine in northern Uganda. Nat Commun.
2022;13: 1–12. doi:10.1038/s41467-022-33873-x

17. Miotto O, Sekihara M, Tachibana SI, Yamauchi M, Pearson RD, Amato R, et
al. Emergence of artemisinin-resistant Plasmodium falciparum with kelch13
C580Y mutations on the island of New Guinea. PLoS Pathog. 2020;16:
e1009133. doi:10.1371/journal.ppat.1009133

18.   Mathieu LC, Cox H, Early AM, Mok S, Lazrek Y, Paquet JC, et al. Local
emergence in amazonia of plasmodium falciparum K13 C580Y mutants
associated with in vitro artemisinin resistance. Elife. 2020;9: 1–21.
doi:10.7554/eLife.51015

19.   Klonis N, Xie SC, McCaw JM, Crespo-Ortiz MP, Zaloumis SG, Simpson JA, et
al. Altered temporal response of malaria parasites determines differential
sensitivity to artemisinin. Proc Natl Acad Sci U S A. 2013;110: 5157–5162.
doi:10.1073/pnas.1217452110

20.   Saralamba S, Pan-Ngum W, Maude RJ, Lee SJ, Tarning J, Lindegårdh N, et
al. Intrahost modeling of artemisinin resistance in Plasmodium falciparum.
Proc Natl Acad Sci U S A. 2011;108: 397–402. doi:10.1073/pnas.1006113108

21.   Saunders DL, Vanachayangkul P, Lon C. Dihydroartemisinin–Piperaquine
Failure in Cambodia. N Engl J Med. 2014;371: 484–485.
doi:10.1056/nejmc1403007

22.   Amaratunga C, Lim P, Suon S, Sreng S, Mao S, Sopha C, et al.
Dihydroartemisinin-piperaquine resistance in Plasmodium falciparum malaria
in Cambodia: A multisite prospective cohort study. Lancet Infect Dis. 2016;16:
357–365. doi:10.1016/S1473-3099(15)00487-9

23.   Thanh NV, Thuy-Nhien N, Tuyen NTK, Tong NT, Nha-Ca NT, Dong LT, et al.
Rapid decline in the susceptibility of Plasmodium falciparum to
dihydroartemisinin-piperaquine in the south of Vietnam. Malar J. 2017;16: 1–
10. doi:10.1186/s12936-017-1680-8

24.   Watts RE, Odedra A, Marquart L, Webb L, Abd-Rahman AN, Cascales L, et al.
Safety and parasite clearance of artemisinin-resistant Plasmodium falciparum
infection: A pilot and a randomised volunteer infection study in Australia. PLoS
Med. 2020;17. doi:10.1371/JOURNAL.PMED.1003203

25.   Rebelo M, Pawliw R, Gower J, Webb L, Mitchell H, Pava Z, et al. Parasite
Viability as a Superior Measure of Antimalarial Drug Activity in Humans. J
Infect Dis. 2021;223: 2154–2163. doi:10.1093/infdis/jiaa678

26.   Witkowski B, Amaratunga C, Khim N, Sreng S, Chim P, Kim S, et al. Novel
phenotypic assays for the detection of artemisinin-resistant Plasmodium
falciparum malaria in Cambodia: In-vitro and ex-vivo drug-response studies.
Lancet Infect Dis. 2013;13: 1043–1049. doi:10.1016/S1473-3099(13)70252-4

27.   Ariey F, Witkowski B, Amaratunga C, Beghain J, Langlois AC, Khim N, et al. A
molecular marker of artemisinin-resistant Plasmodium falciparum malaria.
Nature. 2014;505: 50–55. doi:10.1038/nature12876

28.   Straimer J, Gnädig NF, Witkowski B, Amaratunga C, Duru V, Ramadani AP, et
al. K13-propeller mutations confer artemisinin resistance in Plasmodium
falciparum clinical isolates. Science (80-). 2015;347: 428–431.
doi:10.1126/science.1260867

29.   Birnbaum J, Scharf S, Schmidt S, Jonscher E, Hoeijmakers WAM, Flemming
S, et al. A Kelch13-defined endocytosis pathway mediates artemisinin
resistance in malaria parasites. Science (80-). 2020;367: 51–59.
doi:10.1126/science.aax4735

30.   Yang T, Yeoh LM, Tutor M V., Dixon MW, McMillan PJ, Xie SC, et al.
Decreased K13 Abundance Reduces Hemoglobin Catabolism and Proteotoxic
Stress, Underpinning Artemisinin Resistance. Cell Rep. 2019;29: 2917-

2928.e5. doi:10.1016/j.celrep.2019.10.095

31.   Schumann R, Bischoff E, Klaus S, Möhring S, Flock J, Keller S, et al. Protein abundance and folding rather than the redox state of Kelch13 determine the artemisinin susceptibility of Plasmodium falciparum. Redox Biol. 2021;48. doi:10.1016/j.redox.2021.102177

32.   Siddiqui G, Srivastava A, Russell AS, Creek DJ. Multi-omics based identification of specific biochemical changes associated with PfKelch13-mutant artemisinin-resistant plasmodium falciparum. J Infect Dis. 2017;215: 1435–1444. doi:10.1093/infdis/jix156

33.   Mok S, Stokes BH, Gnädig NF, Ross LS, Yeo T, Amaratunga C, et al. Artemisinin-resistant K13 mutations rewire Plasmodium falciparum's intra-erythrocytic metabolic program to enhance survival. Nat Commun. 2021;12. doi:10.1038/s41467-020-20805-w

34.   Behrens HM, Schmidt S, Spielmann T. The newly discovered role of endocytosis in artemisinin resistance. Med Res Rev. 2021; med.21848. doi:10.1002/med.21848

35.   Demas AR, Sharma AI, Wong W, Early AM, Redmond S, Bopp S, et al. Mutations in Plasmodium falciparum actin-binding protein coronin confer reduced artemisinin susceptibility. Proc Natl Acad Sci. 2018; 201812317. doi:10.1073/pnas.1812317115

36.   Borrmann S, Straimer J, Mwai L, Abdi A, Rippert A, Okombo J, et al. Genome-wide screen identifies new candidate genes associated with artemisinin susceptibility in Plasmodium falciparum in Kenya. Sci Rep. 2013;3. doi:10.1038/srep03318

37.   Simwela N V., Hughes KR, Roberts AB, Rennie MT, Barrett MP, Waters AP. Experimentally engineered mutations in a ubiquitin hydrolase, UBP-1, modulate in vivo susceptibility to artemisinin and chloroquine in plasmodium berghei. Antimicrob Agents Chemother. 2020;64. doi:10.1128/AAC.02484-19

38.   Henrici RC, van Schalkwyk DA, Sutherland CJ. Modification of pfap2μ and pfubp1 Markedly Reduces Ring-Stage Susceptibility of Plasmodium falciparum to Artemisinin in Vitro. Antimicrob Agents Chemother. 2020;64. doi:10.1128/AAC.01542-19

39.   Henriques G, Hallett RL, Beshir KB, Gadalla NB, Johnson RE, Burrow R, et al. Directional selection at the pfmdr1, pfcrt, pfubp1, and pfap2mu loci of Plasmodium falciparum in Kenyan children treated with ACT. J Infect Dis. 2014;210: 2001–2008. doi:10.1093/infdis/jiu358

40.   Behrens HM, Schmidt S, Peigney D, Sabitzki R, Henshall I, May J, et al. Impact of different mutations on Kelch13 protein levels, ART resistance and fitness cost in Plasmodium falciparum parasites. bioRxiv. 2023; 2022.05.13.491767. doi:10.1101/2022.05.13.491767

41.   Francis SE, Sullivan DJ, Goldberg DE. Hemoglobin metabolism in the malaria parasite Plasmodium falciparium. Annual Review of Microbiology. Annual Reviews 4139 El Camino Way, P.O. Box 10139, Palo Alto, CA 94303-0139, USA; 1997. pp. 97–123. doi:10.1146/annurev.micro.51.1.97

42.   Spielmann T, Gras S, Sabitzki R, Meissner M. Endocytosis in Plasmodium and Toxoplasma Parasites. Trends in Parasitology. Elsevier Ltd; 2020. pp. 520–532. doi:10.1016/j.pt.2020.03.010

43. Krugliak M, Zhang J, Ginsburg H. Intraerythrocytic Plasmodium falciparum
utilizes only a fraction of the amino acids derived from the digestion of host cell
cytosol for the biosynthesis of its proteins. Mol Biochem Parasitol. 2002;119:
249–256. doi:10.1016/S0166-6851(01)00427-3

44. Sherman IW. Amino acid metabolism and protein synthesis in malarial
parasites. Bulletin of the World Health Organization. World Health
Organization; 1977. pp. 265–276.

45. Matz JM. Plasmodium's bottomless pit: properties and functions of the malaria
parasite's digestive vacuole. Trends in Parasitology. Elsevier Ltd; 2022. pp.
525–543. doi:10.1016/j.pt.2022.02.010

46. Mesén-Ramírez P, Bergmann B, Elhabiri M, Zhu L, Thien H von, Castro-Peña
C, et al. The parasitophorous vacuole nutrient pore is critical for drug access in
malaria parasites and modulates the fitness cost of artemisinin resistance. Cell
Host Microbe. 2021;0: 283. doi:10.1016/j.chom.2021.11.002

47. Bunditvorapoom D, Kochakarn T, Kotanan N, Modchang C, Kümpornsin K,
Loesbanluechai D, et al. Fitness Loss under Amino Acid Starvation in
Artemisinin-Resistant Plasmodium falciparum Isolates from Cambodia. Sci
Rep. 2018;8: 1–9. doi:10.1038/s41598-018-30593-5

48. Milani KJ, Schneider TG, Taraschi TF. Defining the morphology and
mechanism of the hemoglobin transport pathway in Plasmodium falciparum-
infected erythrocytes. Eukaryot Cell. 2015;14: 415–426.
doi:10.1128/EC.00267-14

49. Tutor M V, Shami GJ, Siddiqui G, Creek DJ, Tilley L, Ralph SA. The
Plasmodium falciparum artemisinin resistance-associated protein Kelch 13 is
required for formation of normal cytostomes. bioRxiv. 2023;
2023.03.30.534478. doi:10.1101/2023.03.30.534478

50. Bakar NA, Klonis N, Hanssen E, Chan C, Tilley L. Digestive-vacuole genesis
and endocytic processes in the early intraerythrocytic stages of Plasmodium
falciparum. J Cell Sci. 2010;123: 441–450. doi:10.1242/jcs.061499

51. Smythe WA, Joiner KA, Hoppe HC. Actin is required for endocytic trafficking in
the malaria parasite Plasmodium falciparum. Cell Microbiol. 2008;10: 452–464.
doi:10.1111/j.1462-5822.2007.01058.x

52. Zhou HC, Gao YH, Zhong X, Wang H. Dynamin like protein 1 participated in
the hemoglobin uptake pathway of Plasmodium falciparum. Chin Med J (Engl).
2009;122: 1686–1691. doi:10.3760/cma.j.issn.0366-6999.2009.14.015

53. Lazarus MD, Schneider TG, Taraschi TF. A new model for hemoglobin
ingestion and transport by the human malaria parasite Plasmodium falciparum.
J Cell Sci. 2008;121: 1937–1949. doi:10.1242/jcs.023150

54. Jonscher E, Flemming S, Schmitt M, Sabitzki R, Reichard N, Birnbaum J, et al.
PfVPS45 Is Required for Host Cell Cytosol Uptake by Malaria Blood Stage
Parasites. Cell Host Microbe. 2019;25: 166-173.e5.
doi:10.1016/j.chom.2018.11.010

55. Sabitzki R, Schmitt M, Flemming S, Jonscher E, Hoehn K, Froehlke U, et al.
Identification of a Rabenosyn-5 like protein and Rab5b in host cell cytosol
uptake reveals conservation of endosomal transport in malaria parasites.
bioRxiv. 2023; 2023.04.05.535711. doi:10.1101/2023.04.05.535711

56. Mukherjee A, Crochetière M-È, Sergerie A, Amiar S, Thompson LA,

Ebrahimzadeh Z, et al. A Phosphoinositide-Binding Protein Acts in the
Trafficking Pathway of Hemoglobin in the Malaria Parasite Plasmodium
falciparum. Soldati-Favre D, editor. MBio. 2022;13. doi:10.1128/mbio.03239-21

57. Wagner MP, Formaglio P, Gorgette O, Dziekan JM, Huon C, Berneburg I, et
al. Human peroxiredoxin 6 is essential for malaria parasites and provides a
host-based drug target. Cell Rep. 2022;39: 110923.
doi:10.1016/j.celrep.2022.110923

58. Roux KJ, Kim DI, Raida M, Burke B. A promiscuous biotin ligase fusion protein
identifies proximal and interacting proteins in mammalian cells. J Cell Biol.
2012;196: 801–810. doi:10.1083/jcb.201112098

59. Kimmel J, Kehrer J, Frischknecht F, Spielmann T. Proximity-dependent
biotinylation approaches to study apicomplexan biology. Mol Microbiol. 2021.
doi:10.1111/mmi.14815

60. Kono M, Herrmann S, Loughran NB, Cabrera A, Engelberg K, Lehmann C, et
al. Evolution and architecture of the inner membrane complex in asexual and
sexual stages of the malaria parasite. Mol Biol Evol. 2012;29: 2113–2132.
doi:10.1093/molbev/mss081

61. Parkyn Schneider M, Liu B, Glock P, Suttie A, McHugh E, Andrew D, et al.
Disrupting assembly of the inner membrane complex blocks Plasmodium
falciparum sexual stage development. Gilberger TW, editor. PLOS Pathog.
2017;13: e1006659. doi:10.1371/journal.ppat.1006659

62. Saini E, Zeeshan M, Brady D, Pandey R, Kaiser G, Koreny L, et al.
Photosensitized INA-Labelled protein 1 (PhIL1) is novel component of the
inner membrane complex and is required for Plasmodium parasite
development. Sci Rep. 2017;7: 15577. doi:10.1038/s41598-017-15781-z

63. Saini E, Sheokand PK, Sharma V, Agrawal P, Kaur I, Singh S, et al.
Plasmodium falciparum PhIL1-associated complex plays an essential role in
merozoite reorientation and invasion of host erythrocytes. PLOS Pathog.
2021;17: e1009750. doi:10.1371/JOURNAL.PPAT.1009750

64. Torres JA, Pasquarelli RR, Back PS, Moon AS, Bradley PJ. Identification and
Molecular Dissection of IMC32, a Conserved Toxoplasma Inner Membrane
Complex Protein That Is Essential for Parasite Replication. Boyle JP, editor.
MBio. 2021;12. doi:10.1128/mBio.03622-20

65. Wichers JS, Wunderlich J, Heincke D, Pazicky S, Strauss J, Schmitt M, et al.
Identification of novel inner membrane complex and apical annuli proteins of
the malaria parasite Plasmodium falciparum. Cell Microbiol. 2021;23: e13341.
doi:10.1111/cmi.13341

66. Rudlaff RM, Kraemer S, Streva VA, Dvorin JD. An essential contractile ring
protein controls cell division in Plasmodium falciparum. Nat Commun. 2019;10:
2181. doi:10.1038/s41467-019-10214-z

67. Morano AA, Rudlaff RM, Dvorin JD. A PPP-type pseudophosphatase is
required for the maintenance of basal complex integrity in Plasmodium
falciparum. Nat Commun. 2023;14: 3916. doi:10.1038/s41467-023-39435-z

68. Geiger M, Brown C, Wichers JS, Strauss J, Lill A, Thuenauer R, et al.
Structural Insights Into PfARO and Characterization of its Interaction With
PfAIP. J Mol Biol. 2020;432: 878–896. doi:10.1016/j.jmb.2019.12.024

69. Koreny L, Zeeshan M, Barylyuk K, Tromer EC, van Hooff JJE, Brady D, et al.

Molecular characterization of the conoid complex in Toxoplasma reveals its
conservation in all apicomplexans, including Plasmodium species. Duffy M,
editor. PLOS Biol. 2021;19: e3001081. doi:10.1371/journal.pbio.3001081

70. Thommen BT, Dziekan JM, Achcar F, Tjia S, Passecker A, Buczak K, et al.
FKBP35 secures ribosome homeostasis in Plasmodium falciparum. bioRxiv.
2022; 2022.12.09.519720. doi:10.1101/2022.12.09.519720

71. Kumar R, Adams B, Musiyenko A, Shulyayeva O, Barik S. The FK506-binding
protein of the malaria parasite, Plasmodium falciparum, is a FK506-sensitive
chaperone with FK506-independent calcineurin-inhibitory activity. Mol Biochem
Parasitol. 2005;141: 163–73. doi:10.1016/j.molbiopara.2005.02.007

72. Boucher MJ, Ghosh S, Zhang L, Lal A, Jang SW, Ju A, et al. Integrative
proteomics and bioinformatic prediction enable a high-confidence apicoplast
proteome in malaria parasites. Striepen B, editor. PLoS Biol. 2018;16:
e2005895. doi:10.1371/journal.pbio.2005895

73. Birnbaum J, Flemming S, Reichard N, Soares AB, Mesén-Ramírez P,
Jonscher E, et al. A genetic system to study Plasmodium falciparum protein
function. Nat Methods. 2017;14: 450–456. doi:10.1038/nmeth.4223

74. Wichers JS, Scholz JAM, Strauss J, Witt S, Lill A, Ehnold L-I, et al. Dissecting
the Gene Expression, Localization, Membrane Topology, and Function of the
Plasmodium falciparum STEVOR Protein Family. Boothroyd JC, editor. MBio.
2019;10: e01500-19. doi:10.1128/mBio.01500-19

75. Subudhi AK, O'Donnell AJ, Ramaprasad A, Abkallo HM, Kaushik A, Ansari
HR, et al. Malaria parasites regulate intra-erythrocytic development duration
via serpentine receptor 10 to coordinate with host rhythms. Nat Commun.
2020;11. doi:10.1038/s41467-020-16593-y

76. Tawk L, Chicanne G, Dubremetz J-F, Richard V, Payrastre B, Vial HJ, et al.
Phosphatidylinositol 3-Phosphate, an Essential Lipid in Plasmodium, Localizes
to the Food Vacuole Membrane and the Apicoplast. Eukaryot Cell. 2010;9:
1519–1530. doi:10.1128/EC.00124-10

77. Wall RJ, Zeeshan M, Katris NJ, Limenitakis R, Rea E, Stock J, et al.
Systematic analysis of Plasmodium myosins reveals differential expression,
localisation, and function in invasive and proliferative parasite stages. Cell
Microbiol. 2019;21. doi:10.1111/cmi.13082

78. Robinson MS, Sahlender DA, Foster SD. Rapid Inactivation of Proteins by
Rapamycin-Induced Rerouting to Mitochondria. Dev Cell. 2010;18: 324–331.
doi:10.1016/j.devcel.2009.12.015

79. Haruki H, Nishikawa J, Laemmli UK. The Anchor-Away Technique: Rapid,
Conditional Establishment of Yeast Mutant Phenotypes. Mol Cell. 2008;31:
925–932. doi:10.1016/j.molcel.2008.07.020

80. Sijwali PS, Rosenthal PJ. Gene disruption confirms a critical role for the
cysteine protease falcipain-2 in hemoglobin hydrolysis by Plasmodium
falciparum. Proc Natl Acad Sci U S A. 2004;101: 4384–4389.
doi:10.1073/pnas.0307720101

81. Casella JF, Flanagan MD, Lin S. Cytochalasin D inhibits actin polymerization
and induces depolymerization of actin filaments formed during platelet shape
change. Nature. 1981;293: 302–305. doi:10.1038/293302a0

82. Otto TD, Wilinski D, Assefa S, Keane TM, Sarry LR, Böhme U, et al. New

insights into the blood-stage transcriptome of Plasmodium falciparum using
RNA-Seq. Mol Microbiol. 2010;76: 12–24. doi:10.1111/j.1365-
2958.2009.07026.x

83. Kimmel J, Schmitt M, Sinner A, Jansen PWTC, Mainye S, Ramón-Zamorano
G, et al. Gene-by-gene screen of the unknown proteins encoded on
Plasmodium falciparum chromosome 3. Cell Syst. 2023;14: 9-23.e7.

84. Hoeijmakers WAM, Miao J, Schmidt S, Toenhake CG, Shrestha S, Venhuizen
J, et al. Epigenetic reader complexes of the human malaria parasite,
Plasmodium falciparum. Nucleic Acids Res. 2019;47: 11574–11588.
doi:10.1093/nar/gkz1044

85. MalariaGEN, Ahouidi A, Ali M, Almagro-Garcia J, Amambua-Ngwa A,
Amaratunga C, et al. An open dataset of Plasmodium falciparum genome
variation in 7,000 worldwide samples. Wellcome open Res. 2021;6: 42.
doi:10.12688/wellcomeopenres.16168.2

86. Wichers JS, Tonkin-Hill G, Thye T, Krumkamp R, Kreuels B, Strauss J, et al.
Common virulence gene expression in adult first-time infected malaria patients
and severe cases. Elife. 2021;10. doi:10.7554/eLife.69040

87. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al.
Highly accurate protein structure prediction with AlphaFold. Nature. 2021;596:
583–589. doi:10.1038/s41586-021-03819-2

88. Cherfils J. Structural mimicry of DH domains by Arfaptin suggests a model for
the recognition of Rac-GDP by its guanine nucleotide exchange factors. FEBS
Lett. 2001;507: 280–284. doi:10.1016/S0014-5793(01)02970-2

89. Tarricone C, Xiao B, Justin N, Walker PA, Rittinger K, Gamblin SJ, et al. The
structural basis of Arfaptin-mediated cross-talk between Rac and Arf signalling
pathways. Nature. 2001;411: 215–219. doi:10.1038/35075620

90. McGovern OL, Rivera-Cuevas Y, Kannan G, Narwold AJ, Carruthers VB.
Intersection of endocytic and exocytic systems in Toxoplasma gondii. Traffic.
2018;19: 336–353. doi:10.1111/tra.12556

91. Struck NS, de Souza Dias S, Langer C, Marti M, Pearce JA, Cowman AF, et
al. Re-defining the Golgi complex in Plasmodium falciparum using the novel
Golgi marker PfGRASP. J Cell Sci. 2005;118: 5603–5613.
doi:10.1242/jcs.02673

92. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al.
AlphaFold Protein Structure Database: Massively expanding the structural
coverage of protein-sequence space with high-accuracy models. Nucleic Acids
Res. 2022;50: D439–D444. doi:10.1093/nar/gkab1061

93. Hirst J, Lui WWY, Bright NA, Totty N, Seaman MNJ, Robinson MS. A family of
proteins with γ-adaptin and VHS domains that facilitate trafficking between the
trans-golgi network and the vacuole/lysosome. J Cell Biol. 2000;149: 67–79.
doi:10.1083/jcb.149.1.67

94. Dell'Angelica EC, Puertollano R, Mullins C, Aguilar RC, Vargas JD, Hartnell
LM, et al. GGAs: A family of ADP ribosylation factor-binding proteins related to
adaptors and associated with the Golgi complex. J Cell Biol. 2000;149: 81–93.
doi:10.1083/jcb.149.1.81

95. Blum M, Chang HY, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et
al. The InterPro protein families and domains database: 20 years on. Nucleic

Acids Res. 2021;49: D344–D354. doi:10.1093/nar/gkaa977

96.   Traub LM, Downs MA, Westrich JL, Fremont DH. Crystal structure of the α
appendage of AP-2 reveals a recruitment platform for clathrin-coat assembly.
Proc Natl Acad Sci U S A. 1999;96: 8907–8912. doi:10.1073/pnas.96.16.8907

97.   Hirst J, Robinson MS. Clathrin and adaptors. Biochim Biophys Acta - Mol Cell
Res. 1998;1404: 173–193. doi:10.1016/S0167-4889(98)00056-1

98.   Koreny L, Mercado-Saavedra BN, Klinger CM, Barylyuk K, Butterworth S, Hirst
J, et al. Stable endocytic structures navigate the complex pellicle of
apicomplexan parasites. Nat Commun. 2023;14: 2167. doi:10.1038/s41467-
023-37431-x

99.   Tanna CE, Goss LB, Ludwig CG, Chen PW. Arf GAPs as regulators of the
actin cytoskeleton—An update. International Journal of Molecular Sciences.
MDPI AG; 2019. doi:10.3390/ijms20020442

100.   Kjos I, Vestre K, Guadagno NA, Borg Distefano M, Progida C. Rab and Arf
proteins at the crossroad between membrane transport and cytoskeleton
dynamics. Biochimica et Biophysica Acta - Molecular Cell Research. Elsevier
B.V.; 2018. pp. 1397–1409. doi:10.1016/j.bbamcr.2018.07.009

101.   Hunt P, Afonso A, Creasey A, Culleton R, Sidhu ABS, Logan J, et al. Gene
encoding a deubiquitinating enzyme is mutated in artesunate- and
chloroquine-resistant rodent malaria parasites. Mol Microbiol. 2007;65: 27–40.
doi:10.1111/j.1365-2958.2007.05753.x

102.   Sun D, Leung CL, Liem RK. Characterization of the microtubule binding
domain of microtubule actin crosslinking factor (MACF): identification of a
novel group of microtubule associated proteins. J Cell Sci. 2001;114: 161–172.
doi:10.1242/jcs.114.1.161

103.   Sahu SC, Swanson KA, Kang RS, Huang K, Brubaker K, Ratcliff K, et al.
Conserved Themes in Target Recognition by the PAH1 and PAH2 Domains of
the Sin3 Transcriptional Corepressor. J Mol Biol. 2008;375: 1444–1456.
doi:10.1016/j.jmb.2007.11.079

104.   Brubaker K, Cowley SM, Huang K, Loo L, Yochum GS, Ayer DE, et al.
Solution structure of the interacting domains of the mad-sin3 complex:
Implications for recruitment of a chromatin-modifying complex. Cell. 2000;103:
655–665. doi:10.1016/S0092-8674(00)00168-9

105.   Doherty GJ, McMahon HT. Mechanisms of Endocytosis. Annu Rev Biochem.
2009;78: 857–902. doi:10.1146/annurev.biochem.78.081307.110540

106.   Kaksonen M, Roux A. Mechanisms of clathrin-mediated endocytosis. Nature
Reviews Molecular Cell Biology. Nature Publishing Group; 2018. pp. 313–326.
doi:10.1038/nrm.2017.132

107.   Thottacherry JJ, Sathe M, Prabhakara C, Mayor S. Spoiled for choice: Diverse
endocytic pathways function at the cell surface. Annual Review of Cell and
Developmental Biology. Annual Reviews Inc.; 2019. pp. 55–84.
doi:10.1146/annurev-cellbio-100617-062710

108.   Henrici RC, Edwards RL, Zoltner M, van Schalkwyk DA, Hart MN, Mohring F,
et al. The plasmodium falciparum artemisinin susceptibility-associated ap-2
adaptin μ subunit is clathrin independent and essential for schizont maturation.
MBio. 2020;11. doi:10.1128/mBio.02918-19

109.   Wan W, Dong H, Lai D-H, Yang J, He K, Tang X, et al. The Toxoplasma

micropore mediates endocytosis for selective nutrient salvage from host cell compartments. Nat Commun. 2023;14: 977. doi:10.1038/s41467-023-36571-4

110. Schmid EM, Ford MGJ, Burtey A, Praefcke GJK, Peak-Chew S-Y, Mills IG, et al. Role of the AP2 β-Appendage Hub in Recruiting Partners for Clathrin-Coated Vesicle Assembly. Hughson F, editor. PLoS Biol. 2006;4: e262. doi:10.1371/journal.pbio.0040262

111. Perrais D. Cellular and structural insight into dynamin function during endocytic vesicle formation: a tale of 50 years of investigation. Biosci Rep. 2022;42. doi:10.1042/bsr20211227

112. Heredero-Bermejo I, Varberg JM, Charvat R, Jacobs K, Garbuz T, Sullivan WJ, et al. TgDrpC, an atypical dynamin-related protein in *Toxoplasma gondii,* is associated with vesicular transport factors and parasite division. Mol Microbiol. 2019;111: 46–64. doi:10.1111/mmi.14138

113. DePina AS, Langford GM. Vesicle transport: The role of actin filaments and myosin motors. Microsc Res Tech. 1999;47: 93–106. doi:10.1002/(SICI)1097-0029(19991015)47:2<93::AID-JEMT2>3.0.CO;2-P

114. Chakrabarti R, Lee M, Higgs HN. Multiple roles for actin in secretory and endocytic pathways. Current Biology. Cell Press; 2021. pp. R603–R618. doi:10.1016/j.cub.2021.03.038

115. Cerqueira GC, Cheeseman IH, Schaffner SF, Nair S, McDew-White M, Phyo AP, et al. Longitudinal genomic surveillance of Plasmodium falciparum malaria parasites reveals complex genomic architecture of emerging artemisinin resistance. Genome Biol. 2017;18: 78. doi:10.1186/s13059-017-1204-4

116. Zhang M, Wang C, Otto TD, Oberstaller J, Liao X, Adapa SR, et al. Uncovering the essential genes of the human malaria parasite Plasmodium falciparum by saturation mutagenesis. Science. 2018;360: eaap7847. doi:10.1126/science.aap7847

117. Carmeille R, Schiano Lomoriello P, Devarakonda PM, Kellermeier JA, Heaslip AT. Actin and an unconventional myosin motor, TgMyoF, control the organization and dynamics of the endomembrane network in Toxoplasma gondii. Gubbels M-J, editor. PLOS Pathog. 2021;17: e1008787. doi:10.1371/journal.ppat.1008787

118. Heaslip AT, Nelson SR, Warshaw DM. Dense granule trafficking in *Toxoplasma gondii* requires a unique class 27 myosin and actin filaments. Steinberg G, editor. Mol Biol Cell. 2016;27: 2080–2089. doi:10.1091/mbc.E15-12-0824

119. Devarakonda PM, Sarmiento V, Heaslip AT. Trafficking of nuclear-encoded apicoplast proteins depends on F-actin and Myosin F in Toxoplasma gondii. bioRxiv. 2023; 2023.01.01.521342. doi:10.1101/2023.01.01.521342

120. Jacot D, Daher W, Soldati-Favre D. Toxoplasma gondii myosin F, an essential motor for centrosomes positioning and apicoplast inheritance. EMBO J. 2013;32: 1702–1716. doi:10.1038/emboj.2013.113

121. Liffner B, Cepeda Diaz AK, Blauwkamp J, Anaguano D, Frölich S, Muralidharan V, et al. Atlas of Plasmodium falciparum intraerythrocytic development using expansion microscopy. bioRxiv Prepr Serv Biol. 2023. doi:10.1101/2023.03.22.533773

122. Chern JH, Pasquarelli RR, Moon AS, Chen AL, Sha J, Wohlschlegel JA, et al.

A Novel Toxoplasma Inner Membrane Complex Suture-Associated Protein Regulates Suture Protein Targeting and Colocalizes with Membrane Trafficking Machinery . MBio. 2021. doi:10.1128/mbio.02455-21

123. Kumari V, Prasad KM, Kalia I, Sindhu G, Dixit R, Rawat DS, et al. Dissecting the role of Plasmodium Metacaspase-2 in malaria gametogenesis and sporogony. Emerg Microbes Infect. 2022; 1–50. doi:10.1080/22221751.2022.2052357

124. Fonager J, Franke-Fayard BMD, Adams JH, Ramesar J, Klop O, Khan SM, et al. Development of the piggyBac transposable system for Plasmodium berghei and its application for random mutagenesis in malaria parasites. BMC Genomics. 2011;12. doi:10.1186/1471-2164-12-155

125. Kumari V, Singh AP, Singh J, Sharma R, Akhter M, Mishra PK, et al. Biochemical characterization of unusual cysteine protease of P. falciparum, metacaspase-2 (MCA-2). Mol Biochem Parasitol. 2018;220: 28–41. doi:10.1016/j.molbiopara.2018.01.001

126. Vandana, Shankar S, Prasad KM, Kashif M, Kalia I, Rai R, et al. A nonpeptidyl molecule modulates apoptosis-like cell death by inhibiting P. Falciparum metacaspase-2. Biochem J. 2020;477: 1323–1344. doi:10.1042/BCJ20200050

127. Vandana , Pandey R, Srinivasan E, Kalia I, Singh AP, Saxena A, et al. Plasmodium falciparum metacaspase-2 capture its natural substrate in a non-canonical way . J Biochem. 2021;170: 639–653. doi:10.1093/jb/mvab086

128. Bushell E, Gomes AR, Sanderson T, Anar B, Girling G, Herd C, et al. Functional Profiling of a Plasmodium Genome Reveals an Abundance of Essential Genes. Cell. 2017;170: 260-272.e8. doi:10.1016/j.cell.2017.06.030

129. Walliker D, Quakyi IA, Wellems TE, McCutchan TF, Szarfman A, London WT, et al. Genetic analysis of the human malaria parasite Plasmodium falciparum. Science (80-). 1987;236: 1661–6.

130. Trager W, Jensen JB. Continuous culture of Plasmodium falciparum: its impact on malaria research. Int J Parasitol. 1997;27: 989–1006. doi:10.1016/S0020-7519(97)00080-5

131. Rivadeneira E, Wasserman M, Espinal C. Separation and Concentration of Schizonts of Plasmodium falciparum by Percoll Gradients. J Protozool. 1983;30: 367–370. doi:10.1111/j.1550-7408.1983.tb02932.x

132. Lambros C, Vanderberg JP. Synchronization of Plasmodium falciparum Erythrocytic Stages in Culture. J Parasitol. 1979;65: 418. doi:10.2307/3280287

133. Mesén-Ramírez P, Bergmann B, Tran TT, Garten M, Stäcker J, Naranjo-Prado I, et al. EXP1 is critical for nutrient uptake across the parasitophorous vacuole membrane of malaria parasites. de Koning-Ward TF, editor. PLoS Biol. 2019;17: e3000473. doi:10.1371/journal.pbio.3000473

134. Wichers-Misterek JS, Binder AM, Mesén-Ramírez P, Dorner LP, Safavi S, Fuchs G, et al. A Microtubule-Associated Protein Is Essential for Malaria Parasite Transmission. Miller LH, editor. MBio. 2023. doi:10.1128/mbio.03318-22

135. Moon RW, Hall J, Rangkuti F, Ho YS, Almond N, Mitchell GH, et al. Adaptation of the genetically tractable malaria pathogen Plasmodium knowlesi to continuous culture in human erythrocytes. Proc Natl Acad Sci. 2013;110: 531–536. doi:10.1073/pnas.1216457110

136. Ganesan SM, Morrisey JM, Ke H, Painter HJ, Laroiya K, Phillips MA, et al.
Yeast dihydroorotate dehydrogenase as a new selectable marker for
Plasmodium falciparum transfection. Mol Biochem Parasitol. 2011;177: 29–34.
doi:10.1016/j.molbiopara.2011.01.004
137. Grüring C, Spielmann T. Imaging of live malaria blood stage parasites.
Methods Enzymol. 2012;506: 81–92. doi:10.1016/B978-0-12-391856-7.00029-
9
138. Malleret B, Claser C, Ong ASM, Suwanarusk R, Sriprawat K, Howland SW, et
al. A rapid and robust tri-color flow cytometry assay for monitoring malaria
parasite development. Sci Rep. 2011;1: 118. doi:10.1038/srep00118
139. Mesén-Ramírez P, Reinsch F, Blancke Soares A, Bergmann B, Ullrich AK,
Tenzer S, et al. Stable Translocation Intermediates Jam Global Protein Export
in Plasmodium falciparum Parasites and Link the PTEX Component EXP2 with
Translocation Activity. PLoS Pathog. 2016;12: e1005618.
doi:10.1371/journal.ppat.1005618
140. Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M.
ColabFold: making protein folding accessible to all. Nat Methods. 2022;19:
679–682. doi:10.1038/s41592-022-01488-1
141. Gibrat JF, Madej T, Bryant SH. Surprising similarities in structure comparison.
Curr Opin Struct Biol. 1996;6: 377–385. doi:10.1016/S0959-440X(96)80058-3
142. Aurrecoechea C, Brestelli J, Brunk BP, Dommer J, Fischer S, Gajria B, et al.
PlasmoDB: a functional genomic database for malaria parasites. Nucleic Acids
Res. 2009;37: D539–D543. doi:10.1093/nar/gkn814
143. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et
al. Fiji: an open-source platform for biological-image analysis. Nat Methods.
2012;9: 676–682. doi:10.1038/nmeth.2019
144. Davis MW, Jorgensen EM. ApE, A Plasmid Editor: A Freely Available DNA
Manipulation and Visualization Program. Front Bioinforma. 2022;2: 5.
doi:10.3389/fbinf.2022.818619
145. Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK. SuperPlots: Communicating
reproducibility and variability in cell biology. J Cell Biol. 2020;219.
doi:10.1083/jcb.202001064
1950

Figure 1

Figure 2

A

KIC2 701-771 vs.
GAR domain (1V5R)
RMSD 2.8 Å over 72 aa

KIC3 183-249 vs.
PAH domain (1G1E)
RMSD 2.3 Å over 64 aa

KIC4 1-169 vs.
VHS domain (1ELK)
RMSD 3.3 Å over 144 aa

KIC4 170-268 vs.
GAT domain (1WRD)
RMSD 2.5 Å over 88 aa

KIC4 661-764 vs.
clathrin adaptor α/β/γ (2IV9)
RMSD 3.8 Å over 69 aa

KIC4 774-888 vs.
β adaptin app. (2IV9)
RMSD 2.7 Å over 69 aa

KIC5 1-88 vs.
GAT domain (1YD8)
RMSD 1.6 Å over 80 aa

KIC5 342-453 vs.
clathrin adaptor α/β/γ (1KYF)
RMSD 4.0 Å over 88 aa

KIC5 1712-1717+1791-1870 vs.
clathrin adaptor α (1KYF)
RMSD 3.9 Å over 72 aa

KIC7 8-83 vs.
ArfGAP domain (2D9L)
RMSD 3.4 Å over 72 aa

UBP1 1741-1853 vs.
VHS domain (1VDY)
RMSD 3.6 Å over 104 aa

Legend:
GAR
PAH
VHS
GAT
Clathrin adapter α/β/γ-adaptin, appendage, Ig-like
β-adaptin appendage, C-terminal subdomain
Clathrin adaptor, α-adaptin, appendage, C-terminal
ArfGAP
metacaspase
EH
USP
Kelch propeller
BTB/POZ
Clathrin adaptor, μ
μ homology domain
Myosin SH3
myosin head, motor
Ras-like-GTPase
WD40 repeat
TPR
DnaJ

C
hGGA1: 3 4 hinge 5
PfKIC4: 3 4 unfolded 5 6
hAP2B1: Clathrin/coatomer adapt-like_N 5 6

D
PfKIC5: 4 5 7
hAP2A1: Clathrin/coatomer adapt-like_N 5 7

E
K13 compartment proteins

link endocytosis or ART resistance

yes		no	
Name	VTD	Name	VTD
KIC4	+	KIC1	–
KIC5	+	KIC2	–
KIC7	+	KIC3	–
KIC12	–	KIC6	–
AP2μ	+	KIC8	–
EPS15	+	KIC9	–
MCA2	+	KIC11	–
UBP1	+		
K13	–		
MyoF	–		

Table 1

Gene ID	PlasmoDB annotation	Gene name
PF3D7_1438400	metacaspase-like protein	*mca2*
PF3D7_1243400	conserved Plasmodium protein, unknown function	
PF3D7_0204300	conserved Plasmodium protein, unknown function	
PF3D7_1365800	conserved Plasmodium protein, unknown function	
PF3D7_1447800	calponin homology domain-containing protein, putative	
PF3D7_1142100	conserved Plasmodium protein, unknown function	*kic11*
PF3D7_0103100	vacuolar protein sorting-associated protein 51, putative	*vps51*
PF3D7_1117900	conserved Plasmodium protein, unknown function	
PF3D7_1016200	Rab3 GTPase-activating protein non-catalytic subunit, putative	
PF3D7_1329100	myosin F	*myof*
PF3D7_1329500	conserved Plasmodium protein, unknown function	*kic12*
PF3D7_0405700	lysine decarboxylase, putative	*uis14*
PF3D7_0907200	GTPase-activating protein, putative	