A ligand-receptor interactome atlas of the zebrafish

Milosz Chodkowski[1,2], Andrzej Zielezinski[1], Savani Anbalagan[1,2#]

[1]Institute of Molecular Biology and Biotechnology, Faculty of Biology, Adam Mickiewicz University in Poznań, Poznań, Poland.

[2]Authors contributed equally

[#]Correspondence: savani.anbalagan@amu.edu.pl

Running title: Zebrafish ligand-receptor interactome atlas

Keywords: Secretome, Receptome, Interactome, synapse, PSD, glia, pituicyte, oxytocin, Zebrafish

Summary

Studies in zebrafish can unravel the functions of cellular communication and thus identify novel bench-to-bedside drugs targeting cellular communication signalling molecules. Due to the incomplete annotation of zebrafish proteome, the knowledge of zebrafish receptome, secretome and tools to explore their interactome is limited. To address this gap, we *de novo* predicted the cellular localization of zebrafish reference proteome using deep learning algorithm. We combined the predicted and existing annotations on cellular localization of zebrafish proteins, and created repositories of zebrafish secretome, receptome, and interactome as well as associated diseases and targeting drugs. Unlike other tools, our interactome atlas is primarily based on physical interaction data of zebrafish proteome. The resources are available as R and Python scripts (https://github.com/*DanioTalk*). *DanioTalk* provides a novel resource for researchers interested in targeting cellular communication in zebrafish, as we demonstrate in applications studying synapse and axo-glial interactome.

Introduction

Ligand-receptor-mediated cellular communication is essential for diverse processes such as differentiation, homeostasis, tissue morphogenesis, animal development, and behaviour (Blockus and Chédotal, 2016; Dufour et al., 2020; Pires-daSilva and Sommer, 2003; Steinhart and Angers, 2018). Dysregulated cellular communication can lead to a plethora of phenotypes ranging from uncontrolled cellular proliferation, developmental defects to disease-states and drug resistance (Kowalczyk et al., 2022; Nisar et al., 2022; Straussman et al., 2012). Hence, receptors and ligands are among the leading drug targets for treating human disease conditions (Attwood et al., 2020; Hauser et al., 2017).

With advances in cryo-electron microscopy and machine learning approaches that enable protein structure prediction, the number of drug targets is expected to increase (Congreve et al., 2020; Lees et al., 2021; Tunyasuvunakool et al., 2021). These developments can aid drug screenings directly using animal model organisms. Due to challenges in using mammalian models in drug screening and limitations of *in vitro* drug screens, larval zebrafish are being increasingly used in high throughput drug screening (Cagan et al., 2019; Hoffman et al., 2016; Patton et al., 2021; Ségalat, 2007). However, despite the importance of zebrafish in drug discovery, there are no actively maintained datasets of zebrafish secreted factors, receptors and their interactions. Few existing predictions of zebrafish secretome and membrane proteome are based on incomplete annotation data (Csályi et al., 2016; Goldberg et al., 2014; Klee, 2008; Nauroy et al., 2018).

With the extensive single-cell RNA-seq datasets of zebrafish, cellular communication in zebrafish can be explored at a holistic level even for rare and transient cells (Dimitrov et al., 2022; Schier, 2020). Several tools built on mammalian interactome data has been previously applied to in zebrafish-based studies to map the ligand-receptor interactions in (Campbell et al., 2021; Holler et al., 2021; Hu et al., 2022). However, only about half of the zebrafish secretome and receptome data have orthologous interaction records for human genes. Tools that depend on such ortholog records are likely to ignore zebrafish genes lacking mammalian orthologs or genes with incomplete orthologue

records. Hence, it is necessary to build a zebrafish secretome and receptome database that also includes information on ortholog-less genes and genes with incomplete annotation data.

Here, we describe *DanioTalk*, a repository of zebrafish secretome, receptome, and interactome and tools to explore them from zebrafish-based omics datasets. To address the limitations of existing tools, *DanioTalk* secretome and receptome datasets are based on deep learning-based subcellular localization predictions allowing the incorporation of orphan zebrafish proteins. *DanioTalk* can identify secretome-receptome pairs primarily based on physical interaction records of zebrafish proteome. *DanioTalk* scripts are available on GitHub and allow exploration of ligand-receptor interactions in any zebrafish-based -omics datasets. Finally, we present the application of *DanioTalk* on mapping the secretome-receptome interactions between synaptosome and postsynaptic density (PSD) and axo-glial interactions between oxytocin neurons and glial pituicytes from RNA-seq datasets. We identified novel ligand-receptor interaction pairs with potential functions in axo-glial interactions. *DanioTalk* provides a novel tool and opportunity for the zebrafish research community to explore ligand-receptor interactions.

Results

De novo prediction of zebrafish secretome and membrane proteome

A bottleneck in using gene ontology (GO) terms or UniprotKB-based cellular localization records of the zebrafish proteome is the incompleteness of such annotations (**Figure 1 – figure supplement 1A**) (Gene Ontology Consortium, 2021; The UniProt Consortium, 2021). To address this issue, we used DeepLoc 2.0, a sequence-based deep learning algorithm that can predict cellular localization of proteins with high accuracy (Thumuluri et al., 2022) (**Figure 1A**). We subjected the current zebrafish reference proteome (UP000000437) to DeepLoc 2.0-based prediction (Nevers et al., 2022). Majority of proteins were predicted to be cytoplasmic (n = 12186; 26%), nuclear (n = 7905; 17%) and multilocalizing (n = 12211; 26%) (**Figure 1B-C, Figure 1 – figure**

supplement 1B & Supplementary file 1). DeepLoc 2.0 predicted that 3699 (8%) and 6481 (14%) of proteins were exclusively extracellular- and cell membrane-localized, respectively (**Figure 1B & Supplementary file 1**). This corresponds to 2571 (10%) and 4002 (15%) genes that code extracellular and membrane-localized proteins, respectively (**Figure 1B'**). In comparison to previous records, DeepLoc 2.0 identified an additional 1104 and 1953 genes coding for extracellular and membranal proteins, respectively (**Figure 1C** & **D**) (Goldberg et al., 2014; Nauroy et al., 2018; The UniProt Consortium, 2021).

Next, we manually assessed how reliable are DeepLoc 2.0 predictions for a subset of proteins. DeepLoc 2.0 successfully predicted the most abundant secreted proteins comprising of pancreatic enzymes (Prss, Cela, Cpa, Amy2a family) as extracellular proteins (coded by 24 genes) (Uhlén et al., 2015) (**Supplementary file 2**). DeepLoc 2.0 prediction efficiency was also high for Wnt family of secreted proteins (coded by 24 genes). Next, we assessed the DeepLoc 2.0 predictions for the Fgf protein family, which includes signal peptide-less secreted (Fgf9, Fgf16, Fgf20) and non-secreted Fgf11 subfamily (Ornitz and Itoh, 2015). DeepLoc 2.0 successfully predicted 17 genes coding for canonical signal peptide-containing extracellular Fgf proteins (Fgf 3-8,10, 17-19, 21-23) and all 7 of the intracellular Fgf11 subfamily proteins as non-extracellular proteins. However, when classifying signal peptide-less extracellular Fgf's, DeepLoc 2.0 only classified Fgf9 as extracellular and misclassified Fgf16, Fgf20a and Fgf20b as cytoplasmic proteins (**Supplementary file 2**). Hence, although DeepLoc 2.0 is highly reliable in predicting zebrafish secretome, manual curation is required.

Curation of zebrafish secretome and membrane receptome

To create a list of zebrafish secretome and receptome, we first combined our DeepLoc 2.0 predictions with annotations from other existing records (UniprotKB CC, GO terms, and Matrisome) (Gene Ontology Consortium, 2021; Nauroy et al., 2018; The UniProt Consortium, 2021). In addition, we added the zebrafish orthologues of mammalian secretome- and receptor-coding genes based on Cell-Cell interaction

(baderlab.org/CellCellInteractions) and CellTalkDB databases (**Figure 2A**) (Shao et al., 2021; Ximerakis et al., 2019) and performed manual curation (as detailed in Material sections). Our curated secretome (ligands) and cell membrane receptome database consist of 2788 (10.5%) and 2343 (8.8%) of protein-coding genes respectively (**Figure 2B** & **Supplementary file 3**). In comparison to zebrafish matrisome dataset, our curated ligand lists contain 1848 additional genes (1566 DeepLoc 2 predictions and 282 manually curated extracellular-coding genes) (**Figure 2C**).

Disease-associated secretome and membrane receptome

Next, we explored the identity of human diseases in which zebrafish models can be used for understanding disease etiology and/or in drug screening. We cross-referenced our ligand and membrane receptor list with the human orthologue information available in ZFIN database (Bradford et al., 2022). We identified 51% and 43% of the ligand- and cell membrane receptor-coding genes having human orthologue gene records (**Figure 2D & Supplementary file 4**). Next, we cross-referenced our dataset with the zebrafish orthologues of human disease-associated genes in Online Mendelian Inheritance in Man (omim.org) database. We identified 549 ligand- and 397 receptor-coding genes containing records of human orthologue genes that are disease-associated (**Fig. 2E**). Some of the major diseases associated with ligands and cell membrane receptors are related to bone diseases, macular degeneration, diabetes mellitus, and cancer (**Fig. 2F**). Nevertheless, several of the ligands and receptors were also associated with numerous syndromes (e.g., Ehlers-Danlos syndrome, Loeys-Dietz syndrome and Long QT syndrome) (**Fig. 2F**).

Next, we explored the drug-targetable status of the zebrafish secretome and receptome. We cross-referenced our ligand and receptome list with DrugCentral database that contains an up-to-date list of drugs and their targets (Avram et al., 2021). We identified 149 ligands and 392 receptors that can be potentially targeted by drugs (**Fig. 2G & Supplementary file 5**). With respect to the choice of drugs, 311 (13%) and

1023 (44%) of the drugs can potentially target zebrafish ligands and cell membrane receptors respectively (**Fig. 2H**).

DanioTalk - putative *secretome-membrane receptome interactome in zebrafish*

Next, to map the ligand-receptome interactions, we identified all pairwise interactions between our ligand and receptome proteins in the STRING (v11.5) physical protein-protein interaction records of zebrafish (**Figure 3A**), a subset of which is shown in (**Figure 3A'**) (Szklarczyk et al., 2021). The experimental score in STRING database predicts the confidence in the physical proximity of two proteins. Ligand-receptor pairs with scores with high-confidence scores are very likely to physically interact and result in associated downstream signalling activities. Our interactome database contains data for 370 ligands and 342 receptors at high confidence (**Figure 3B**). As we observed highly variable STRING physical interaction scores for experimentally validated orthologous human ligand-receptor pairs, we also incorporated medium-ranked pairs and orthologous human ligand-receptors pairs available in protein-protein interaction database IID (**Figure 3 – figure supplement 3**) (Kotlyar et al., 2022). This resulted in 10,446 ligand-receptor pairs (**Figure 3C, 3C' & Supplementary file 6**). Based on our analysis, at least 1681 ligands and 1455 receptors lack a high-ranking interacting receptor or ligand, respectively (**Figure 3B**).

We next probed the accuracy of our ligand-receptor interactome database. We focused on the Nodal-related TGFβ ligands and WNT ligands, key regulators of development (Schier and Shen, 2000; Shen, 2007; Steinhart and Angers, 2018). Our interaction database identified all the previously reported receptors for Ndr2/Cyclops, Ndr1/Squint and Spaw/Southpaw ligands namely Acvr1b/Alk4 (rank 6 and 13), Acvr2a (rank 2 and 11), Acvr2b (rank 1 and 10) respectively (**Figure 3D** & **Supplementary file 7**). We also noticed Acvr1c, Tgfbr1 and Tgfbr2 as top ranked receptors. Next, we probed Wnt signalling ligand-receptor interactome. Our interaction database identified all the Wnt receptors (Lrp5, Ryk, Fzd, Ror1 and Lrp6) that can interact with the major Wnt

ligands (**Figure 3D'** & **Supplementary file 7**). Altogether, *DanioTalk* is able to accurately identify major ligand-receptor interactions in zebrafish.

Zebrafish brain synaptic secretome-membrane receptome interactome

Next, we tested the application of *DanioTalk* on existing datasets. We first focused on the zebrafish adult brain synaptosomal and postsynaptic density (PSD) proteome dataset (**Figure 4A**) (Bayés et al., 2017). We identified protein products of 148 ligand- and 184 receptor-coding genes in the synaptosomal fraction. While in the PSD fraction, we identified protein products from 75 ligand- and 88 receptor-coding genes (**Figure 4 – figure supplement 1** & **Supplementary file 8**). Between synaptosome ligands and PSD receptors, we identified 4 and 78 interactions at high and medium confidence rankings, respectively (**Supplementary file 9**). The top-ranked ligand-receptor interacting pairs were Ntn1b-Neo1a, Ntn1b-Dcc, Bdnf-Ntrk3a (**Figure 4B**). While in the case of PSD ligand and synaptosome receptor interactions, we identified 21 and 123 interactions at high and medium confidence ranking respectively (**Supplementary file 9**). The top-ranked PSD ligand- synaptosome receptor pairs were Efnb1-Ephb1/Ephb2b, Agrn/Dag1 (**Figure 4B'**).

Zebrafish Oxytocin Neuron-Glial Pituicytes Axo-Glial Interactome

Axo-glial interactions contribute to synaptic morphogenesis and plasticity (Allen and Eroglu, 2017; Allen and Lyons, 2018). However, the identify of all the axo-glial ligand-receptor interactions of neurohypophysis (NH), a major neuroendocrine interface has not been addressed (Pearson and Placzek, 2013; Wittkowski, 1998). The NH-projecting axonal tracts, NH axo-glial cellular components and their ultrastructure are conserved in zebrafish (Anbalagan et al., 2018; Grinevich and Neumann, 2021; Gutnick et al., 2011; Herget et al., 2017). We mapped the axo-glial interactome of Oxytocin (OXT) neurons and glial pituicytes. Towards this, we used the glial pituicyte bulk RNA-seq data and recently identified OXT neuron scRNA-seq data (**Figure 4C**) (Anbalagan et al., 2018; Shafer et al., 2022). We identified 104 enriched ligands and 66 enriched receptors in OXT neurons, while in glial pituicytes we identified 172 enriched ligands and 82

enriched receptors (**Supplementary file 10**). The high ranked pituicyte ligand-OXT neuron receptor interactions were Pomca-Mc4r/Mc3r, Slit3-Robo2, Fn1a-Itgb1a while the high ranked OXT neuron ligand-pituicyte receptors were Ihhb-Cdon/Boc, Bdnf-Sort1a, Wnt3a-Fzd2/Fzd8a and Nts-Sort1a (**Figure 4D** & **Supplementary file 11**). We also observed several autocrine signalling interactions in glial pituicytes (Fgf10a-Fgfr2, Vegfab-Nrp2b) and in OXT neurons (Slit3/Slit2-Robo2) (**Figure 4D** & **Supplemental file 11**).

Discussion

Zebrafish single-cell RNAseq datasets are used to explore the developmental ligand-receptor interactome (Campbell et al., 2021; Herpelinck et al., 2022; Holler et al., 2021; Hu et al., 2022). However, a bottleneck in using such tools is in the dependence on orthologous mammalian ligand-receptor interactome data while ignoring zebrafish-specific data (Farnsworth et al., 2020; Farrell et al., 2018). Thus, tools to study zebrafish secretome, receptome and interactome independent of orthologue data are essential.

In the lack of experimental records, recent studies reported the relatively high success of machine learning-based tools in predicting protein structures and cellular localizations (Jumper et al., 2021; Lin et al., 2022; Thumuluri et al., 2022). Here, we applied a deep learning algorithm to predict the cellular localization of the zebrafish proteome and created a curated repository of the zebrafish secretome and membrane receptome. This allows the inclusion of orphan zebrafish secretome or membranal receptor proteins that lack cellular localization or ortholog records. Also, our database of secretome and receptome-linked diseases will aid researchers interested in developing zebrafish models of human disease.

Generally, creating disease models of zebrafish may require the creation of mutations in paralogous genes (Hinman et al., 2021; Howe et al., 2013). Our results suggest that this is not the case for all diseases. For instance, secreted factor-coding genes such as *col13a1, cpe, dll4* and *tgfb2* can be potentially used to model congenital myasthenic

syndrome 19, BDV syndrome, Adam-Oliver syndrome and Loeys-Dietz syndrome 4 respectively (Bosch et al., 2021; Fontana et al., 2014; Logan et al., 2015; Meester et al., 2015). Whereas receptor coding genes such as *gpc4* and *ccln4* can be potentially used for studying Keipert syndrome and Raynaud syndrome respectively (Amor et al., 2019; Palmer et al., 2018).

One of the strengths of research on larval zebrafish is in the pharmacological perturbation experiments which we have also taken advantage in the past (Anbalagan et al., 2018). Some subscription-based tools built for mammalian model organisms can list drugs targeting mammalian proteins (Krämer et al., 2014; Stelzer et al., 2016). Our *DanioTalk* repository and output files contain similar information on drugs that can likely target the zebrafish secretome and receptome.

The potential of *DanioTalk* was revealed in our analysis of the existing dataset on adult zebrafish brain synaptosome-PSD proteomes (Bayés et al., 2017). Several top ligand-receptor pairs were previously studied for their synaptic role in other model organisms. For example, orthologues of our top-ranking Ntn1b-Dcc ligand-receptor pair are well known for their functions in synaptogenesis and synaptic plasticity (Goldman et al., 2013; Horn et al., 2013; Manitt et al., 2009). Another top ligand-receptor pair was Contactin 2 (Cntn2)-Cntnap1 (CASPR). In mice studies, Contactin has been shown to promote synaptic Cntnap1 surface localization via *Cis*-interaction (Murai et al., 2002). But the fact that Cntn2 and Cntnap1 are enriched in synaptosome and PSD fractions suggests an experimental artefact or additional trans-synaptic roles for the Cntn2-Cntnap1 interaction (Dubessy et al., 2019; Ruegg et al., 1989).

The presence of several conserved ligand-receptor pairs in the zebrafish synapses further stresses the importance of zebrafish in understanding the role of ligand-receptor in synapse biology and in drugs targeting synapses (Hoffman et al., 2016; Hutson and Chien, 2002; Oprişoreanu et al., 2021). Although our synaptosome and PSD ligand-receptor pairs lack spatiotemporal accuracy, *DanioTalk* can be further combined with zebrafish single-cell brain transcriptome records to identify synaptosome-PSD ligand-

receptor interactions for neuronal cell types of interest (Farnsworth et al., 2020; Raj et al., 2020).

In the absence of cell-specific proteome data, RNA-seq data in combination with protein interaction records are largely used to explore ligand-receptor-based cellular interactions (Efremova et al., 2020; Türei et al., 2021). However, such data were not explored to analyse axo-glial ligand-receptor interactions in NH, a major neuroendocrine interface critical in maintaining water balance and reproductive functions (Anbalagan et al., 2018; Brown, 2016; Romanov et al., 2017). Except for few molecular players, the axo-glial interactions in NH were largely unknown (Brown, 2016; Leng et al., 2015; Miyata, 2017; Rosso and Mienville, 2009).

Our analysis validated the Slit3-Robo2 glial pituicyte-OXT neuron interaction, which has previously been reported to promote synaptic OXT levels (Anbalagan et al., 2019). However, we also observed autocrine interactions of Slit2-Robo2 and Slit3-Robo2 in OXT neurons. Slit-Robo autocrine signalling in mice motor neurons has been shown to promote axon fasciculation; however, such defects were not reported for NH-projecting OXT axons in zebrafish (Anbalagan et al., 2019; Jaworski and Tessier-Lavigne, 2012).

Apart from glia to neuronal signalling, neuron to glial signalling are also major players in neuro-glial signalling (Allen and Eroglu, 2017). We observed OXT-neuron-derived Indian hedgehog homolog ligand potentially signalling to glial pituicytes via Cell adhesion molecule-related/down-regulated by oncogenes (Cdon) and Brother of CDO (Boc) receptors. Apart from the role of Hedgehog signalling in midline axonal guidance (Charron and Tessier-Lavigne, 2007; Evans and Bashaw, 2010; Liu and Placzek, 2014), Neuron-derived Hedgehog proteins can regulate astrocyte reactivity and synaptogenesis during mice forebrain development (Chu et al., 2006; Garcia et al., 2010; Xie et al., 2022). It is unknown whether OXT-neuron-derived hedgehog signalling regulates glial pituicytes development and synaptic morphogenesis.

We also observed potential Fgf10a-Fgfr2- and Fgf3-Fgfr2-based autocrine interactions in glial pituicytes (Zhang et al., 2006). This suggests that in addition to the previously reported paracrine role of Fgf10a in neurovascular morphogenesis of NH, Fgf10a and Fgf3 could have additional autocrine functions in glial pituicytes such as reactivity and/or proliferation (Gómez-Pinilla et al., 1995; Goodman et al., 2020; Haan et al., 2013, p. 10; Liu et al., 2013).

Overall, *DanioTalk* offers a comprehensive repository of zebrafish secretome, receptome, secretome-receptome interaction atlas, and a novel data exploratory tool for zebrafish-based researchers.

Methods

Zebrafish dataset:

Zebrafish reference proteome (UP000000437) sequences along with GO-term and cellular localization annotations were downloaded from UniprotKB (The UniProt Consortium, 2021). Gene symbols, orthologue gene symbols, aliases/synonyms, gene descriptions, and ZFIN IDs were downloaded from the ZFIN database (Bradford et al., 2022). The extracellular protein-coding gene was identified based on the annotation terms 'extracellular' or 'secreted' or 'ligands'. Cell membrane protein-coding genes were identified based on terms "plasma membrane".

Subcellular localization prediction:

Zebrafish reference proteome cellular localization was predicted using DeepLoc 2.0 with default parameters (Thumuluri et al., 2022).

Curated database of the zebrafish secretome and membrane receptome:

Human receptors and ligands were obtained from the Cell-Cell Interaction Database (Gary Bader group, Univ. of Toronto) and the CellTalkDB receptor database (Shao et al., 2021; Ximerakis et al., 2019). In addition human ligands were obtained from MatrisomeDB (Shao et al., 2020). The initial list of zebrafish ligands was obtained by combining i) zebrafish matrisome, ii) zebrafish orthologues from the human ligand

database (see above) and iii) extracellular proteins in DeepLoc 2.0-based extracellular protein coding genes. Initial zebrafish membrane-localized receptors list were obtained by combining zebrafish orthologues of the human receptor database (see above), and DeepLoc 2.0-based membranal protein-coding genes. The list was manually curated based on gene descriptions and localization records of human orthologues available in the Compartments database and the Human Protein Atlas (Binder et al., 2014; Uhlén et al., 2015).

Ligand-receptor interactome database:

The ligand-receptor interaction database was generated based on STRING physical interaction database for zebrafish (v11.5, Ref) and zebrafish orthologues of human protein interaction data available in the IID database (version 2021-05) (Kotlyar et al., 2022; Szklarczyk et al., 2021). Pairs were ranked according to the STRING physical interaction score (High ≥ 700, Medium ≥ 400). Orthologous proteins with experimentally validated interactions in the IID database (*H. sapiens*) database were given a score of 500 for ranking purpose.

Application of *DanioTalk* on zebrafish datasets:

Zebrafish synapse and PSD proteome datasets were extracted from supplementary files in (Bayés et al., 2017). Zebrafish OXT neuron dataset was extracted from Neuronal_04_2 cluster data of Supplemental_data/4-pseudobulk_expression/Subclusters-Danio_rerio.csv reported in (Shafer et al., 2022). Glial pituicytes were extracted from the supplemental table (Log2FC > 1, p-adj. < 0.05, read count ≥ 50) provided in (Anbalagan et al., 2018). Gene symbols were updated based on ZFIN alias database. The ligand and receptors were compared and ranked based on *DanioTalk* interaction database and plotted using Circlize R package (Gu et al., 2014; Team, 2013).

Code availability:

The scripts used in this manuscript are available at https://github.com/*DanioTalk*.

Authors' contributions

S.A. obtained funding, conceived and designed the study. A.Z. contributed in study design and performed DeepLoc 2.0 predictions. M.C. wrote all the scripts and performed all analysis. S.A. performed manual curation of ligands and receptors, wrote the manuscript and prepared figures. All authors read and approved of the manuscript.

Acknowledgments

We thank Emilia Wysocka, Iwona Kanonik-Jędrzejak and Arleta Kucz for technical and administrative support. We thank Dr. Damian Szklarczyk (Swiss Institute of Bioinformatics, Zurich) for discussion and advice. We thank BioRender.com team. S.A. is supported by National Science Centre grants (SONATA-BIS 2020/38/E/NZ3/00090 and SONATA 2021/43/D/NZ3/01798).

Competing interests

The authors declare no competing interests exist.

References

Allen NJ, Eroglu C. 2017. Cell Biology of Astrocyte-Synapse Interactions. *Neuron* **96**:697–708. doi:10.1016/j.neuron.2017.09.056

Allen NJ, Lyons DA. 2018. Glia as architects of central nervous system formation and function. *Science* **362**:181–185. doi:10.1126/science.aat0473

Amor DJ, Stephenson SEM, Mustapha M, Mensah MA, Ockeloen CW, Lee WS, Tankard RM, Phelan DG, Shinawi M, de Brouwer APM, Pfundt R, Dowling C, Toler TL, Sutton VR, Agolini E, Rinelli M, Capolino R, Martinelli D, Zampino G, Dumić M, Reardon W, Shaw-Smith C, Leventer RJ, Delatycki MB, Kleefstra T, Mundlos S, Mortier G, Bahlo M, Allen NJ, Lockhart PJ. 2019. Pathogenic Variants in GPC4 Cause Keipert Syndrome. *Am J Hum Genet* **104**:914–924. doi:10.1016/j.ajhg.2019.02.026

Anbalagan S, Blechman J, Gliksberg M, Gordon L, Rotkopf R, Dadosh T, Shimoni E, Levkowitz G. 2019. Robo2 regulates synaptic oxytocin content by affecting actin dynamics. *Elife* **8**:e45650. doi:10.7554/eLife.45650

Anbalagan S, Gordon L, Blechman J, Matsuoka RL, Rajamannar P, Wircer E, Biran J, Reuveny A, Leshkowitz D, Stainier DYR, Levkowitz G. 2018. Pituicyte Cues Regulate the Development of Permeable Neuro-Vascular Interfaces. *Dev Cell* **47**:711-726.e5. doi:10.1016/j.devcel.2018.10.017

Attwood MM, Jonsson J, Rask-Andersen M, Schiöth HB. 2020. Soluble ligands as drug targets. *Nat Rev Drug Discov* **19**:695–710. doi:10.1038/s41573-020-0078-4

Avram S, Bologa CG, Holmes J, Bocci G, Wilson TB, Nguyen D-T, Curpan R, Halip L, Bora A, Yang JJ, Knockel J, Sirimulla S, Ursu O, Oprea TI. 2021. DrugCentral 2021 supports drug discovery and repositioning. *Nucleic Acids Res* **49**:D1160–D1169. doi:10.1093/nar/gkaa997

Bayés À, Collins MO, Reig-Viader R, Gou G, Goulding D, Izquierdo A, Choudhary JS, Emes RD, Grant SGN. 2017. Evolution of complexity in the zebrafish synapse proteome. *Nat Commun* **8**:14613. doi:10.1038/ncomms14613

Binder JX, Pletscher-Frankild S, Tsafou K, Stolte C, O'Donoghue SI, Schneider R, Jensen LJ. 2014. COMPARTMENTS: unification and visualization of protein subcellular localization evidence. *Database* **2014**:bau012. doi:10.1093/database/bau012

Blockus H, Chédotal A. 2016. Slit-Robo signaling. *Development* **143**:3037–3044. doi:10.1242/dev.132829

Bosch E, Hebebrand M, Popp B, Penger T, Behring B, Cox H, Towner S, Kraus C, Wilson WG, Khan S, Krumbiegel M, Ekici AB, Uebe S, Trollmann R, Woelfle J, Reis A, Vasileiou G. 2021. BDV Syndrome: An Emerging Syndrome With Profound Obesity and Neurodevelopmental Delay Resembling Prader-Willi Syndrome. *J Clin Endocrinol Metab* **106**:3413–3427. doi:10.1210/clinem/dgab592

Bradford YM, Van Slyke CE, Ruzicka L, Singer A, Eagle A, Fashena D, Howe DG, Frazer K, Martin R, Paddock H, Pich C, Ramachandran S, Westerfield M. 2022. Zebrafish information network, the knowledgebase for Danio rerio research. *Genetics* **220**:iyac016. doi:10.1093/genetics/iyac016

Brown CH. 2016. Magnocellular Neurons and Posterior Pituitary Function. *Compr Physiol* **6**:1701–1741. doi:10.1002/cphy.c150053

Cagan RL, Zon LI, White RM. 2019. Modeling Cancer with Flies and Fish. *Dev Cell* **49**:317–324. doi:10.1016/j.devcel.2019.04.013

Campbell NR, Rao A, Hunter MV, Sznurkowska MK, Briker L, Zhang M, Baron M, Heilmann S, Deforet M, Kenny C, Ferretti LP, Huang T-H, Perlee S, Garg M, Nsengimana J, Saini M, Montal E, Tagore M, Newton-Bishop J, Middleton MR, Corrie P, Adams DJ, Rabbie R, Aceto N, Levesque MP, Cornell RA, Yanai I, Xavier JB, White RM. 2021. Cooperation between melanoma cell states promotes metastasis through heterotypic cluster formation. *Developmental Cell* **56**:2808-2825.e10. doi:10.1016/j.devcel.2021.08.018

Charron F, Tessier-Lavigne M. 2007. The Hedgehog, TGF-beta/BMP and Wnt families of morphogens in axon guidance. *Adv Exp Med Biol* **621**:116–133. doi:10.1007/978-0-387-76715-4_9

Chu T, Chiu M, Zhang E, Kunes S. 2006. A C-Terminal Motif Targets Hedgehog to Axons, Coordinating Assembly of the Drosophila Eye and Brain. *Developmental Cell* **10**:635–646. doi:10.1016/j.devcel.2006.03.003

Congreve M, de Graaf C, Swain NA, Tate CG. 2020. Impact of GPCR Structures on Drug Discovery. *Cell* **181**:81–91. doi:10.1016/j.cell.2020.03.003

Csályi K, Fazekas D, Kadlecsik T, Türei D, Gul L, Horváth B, Módos D, Demeter A, Pápai N, Lenti K, Csermely P, Vellai T, Korcsmáros T, Varga M. 2016. SignaFish: A Zebrafish-Specific Signaling Pathway Resource. *Zebrafish* **13**:541–544. doi:10.1089/zeb.2016.1277

Dimitrov D, Türei D, Garrido-Rodriguez M, Burmedi PL, Nagai JS, Boys C, Ramirez Flores RO, Kim H, Szalai B, Costa IG, Valdeolivas A, Dugourd A, Saez-Rodriguez J. 2022. Comparison of methods and resources for cell-cell communication inference from single-cell RNA-Seq data. *Nat Commun* **13**:3224. doi:10.1038/s41467-022-30755-0

Dubessy A, Mazuir E, Rappeneau Q, Ou S, Abi Ghanem C, Piquand K, Aigrot M, Thétiot M, Desmazières A, Chan E, Fitzgibbon M, Fleming M, Krauss R, Zalc B, Ranscht B, Lubetzki C, Sol-Foulon N. 2019. Role of a Contactin multi-molecular complex secreted by oligodendrocytes in nodal protein clustering in the CNS. *Glia* **67**:2248–2263. doi:10.1002/glia.23681

Dufour S, Quérat B, Tostivint H, Pasqualini C, Vaudry H, Rousseau K. 2020. Origin and Evolution of the Neuroendocrine Control of Reproduction in Vertebrates, With Special Focus on Genome and Gene Duplications. *Physiological Reviews* **100**:869–943. doi:10.1152/physrev.00009.2019

Efremova M, Vento-Tormo M, Teichmann SA, Vento-Tormo R. 2020. CellPhoneDB: inferring cell–cell communication from combined expression of multi-subunit ligand–receptor complexes. *Nat Protoc* **15**:1484–1506. doi:10.1038/s41596-020-0292-x

Evans TA, Bashaw GJ. 2010. Axon guidance at the midline: of mice and flies. *Curr Opin Neurobiol* **20**:79–85. doi:10.1016/j.conb.2009.12.006

Farnsworth DR, Saunders LM, Miller AC. 2020. A single-cell transcriptome atlas for zebrafish development. *Dev Biol* **459**:100–108. doi:10.1016/j.ydbio.2019.11.008

Farrell JA, Wang Y, Riesenfeld SJ, Shekhar K, Regev A, Schier AF. 2018. Single-cell reconstruction of developmental trajectories during zebrafish embryogenesis. *Science* **360**:eaar3131. doi:10.1126/science.aar3131

Fontana P, Genesio R, Casertano A, Cappuccio G, Mormile A, Nitsch L, Iolascon A, Andria G, Melis D. 2014. Loeys-Dietz syndrome type 4, caused by chromothripsis, involving the TGFB2 gene. *Gene* **538**:69–73. doi:10.1016/j.gene.2014.01.017

Garcia ADR, Petrova R, Eng L, Joyner AL. 2010. Sonic Hedgehog Regulates Discrete Populations of Astrocytes in the Adult Mouse Forebrain. *J Neurosci* **30**:13597–13608. doi:10.1523/JNEUROSCI.0830-10.2010

Gene Ontology Consortium. 2021. The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids Res* **49**:D325–D334. doi:10.1093/nar/gkaa1113

Goldberg T, Hecht M, Hamp T, Karl T, Yachdav G, Ahmed N, Altermann U, Angerer P, Ansorge S, Balasz K, Bernhofer M, Betz A, Cizmadija L, Do KT, Gerke J, Greil R, Joerdens V, Hastreiter M, Hembach K, Herzog M, Kalemanov M, Kluge M, Meier A, Nasir H, Neumaier U, Prade V, Reeb J, Sorokoumov A, Troshani I, Vorberg S, Waldraff S, Zierer J, Nielsen H, Rost B. 2014. LocTree3 prediction of localization. *Nucleic Acids Res* **42**:W350-355. doi:10.1093/nar/gku396

Goldman JS, Ashour MA, Magdesian MH, Tritsch NX, Harris SN, Christofi N, Chemali R, Stern YE, Thompson-Steckel G, Gris P, Glasgow SD, Grutter P, Bouchard J-F, Ruthazer ES, Stellwagen D, Kennedy TE. 2013. Netrin-1 promotes excitatory synaptogenesis between cortical neurons by initiating synapse assembly. *J Neurosci* **33**:17278–17289. doi:10.1523/JNEUROSCI.1085-13.2013

Gómez-Pinilla F, Vu L, Cotman CW. 1995. Regulation of astrocyte proliferation by FGF-2 and heparan sulfate in vivo. *J Neurosci* **15**:2021–2029. doi:10.1523/JNEUROSCI.15-03-02021.1995

Goodman T, Nayar SG, Clare S, Mikolajczak M, Rice R, Mansour S, Bellusci S, Hajihosseini MK. 2020. Fibroblast growth factor 10 is a negative regulator of postnatal neurogenesis in the mouse hypothalamus. *Development* **147**:dev180950. doi:10.1242/dev.180950

Grinevich V, Neumann ID. 2021. Brain oxytocin: how puzzle stones from animal studies translate into psychiatry. *Mol Psychiatry* **26**:265–279. doi:10.1038/s41380-020-0802-9

Gu Z, Gu L, Eils R, Schlesner M, Brors B. 2014. circlize Implements and enhances circular visualization in R. *Bioinformatics* **30**:2811–2812. doi:10.1093/bioinformatics/btu393

Gutnick A, Blechman J, Kaslin J, Herwig L, Belting H-G, Affolter M, Bonkowsky JL, Levkowitz G. 2011. The hypothalamic neuropeptide oxytocin is required for formation of the neurovascular interface of the pituitary. *Dev Cell* **21**:642–654. doi:10.1016/j.devcel.2011.09.004

Haan N, Goodman T, Najdi-Samiei A, Stratford CM, Rice R, El Agha E, Bellusci S, Hajihosseini MK. 2013. Fgf10-expressing tanycytes add new neurons to the appetite/energy-balance regulating centers of the postnatal and adult hypothalamus. *J Neurosci* **33**:6170–6180. doi:10.1523/JNEUROSCI.2437-12.2013

Hauser AS, Attwood MM, Rask-Andersen M, Schiöth HB, Gloriam DE. 2017. Trends in GPCR drug discovery: new agents, targets and indications. *Nat Rev Drug Discov* **16**:829–842. doi:10.1038/nrd.2017.178

Herget U, Gutierrez-Triana JA, Salazar Thula O, Knerr B, Ryu S. 2017. Single-Cell Reconstruction of Oxytocinergic Neurons Reveals Separate Hypophysiotropic and Encephalotropic Subtypes in Larval Zebrafish. *eNeuro* **4**:ENEURO.0278-16.2016. doi:10.1523/ENEURO.0278-16.2016

Herpelinck T, Ory L, Nasello G, Barzegari M, Bolander J, Luyten FP, Tylzanowski P, Geris L. 2022. An integrated single-cell atlas of the skeleton from development through adulthood. doi:10.1101/2022.03.14.484345

Hinman MN, Richardson JI, Sockol RA, Aronson ED, Stednitz SJ, Murray KN, Berglund JA, Guillemin K. 2021. Zebrafish mbnl mutants model physical and molecular phenotypes of myotonic dystrophy. *Dis Model Mech* **14**:dmm045773. doi:10.1242/dmm.045773

Hoffman EJ, Turner KJ, Fernandez JM, Cifuentes D, Ghosh M, Ijaz S, Jain RA, Kubo F, Bill BR, Baier H, Granato M, Barresi MJF, Wilson SW, Rihel J, State MW, Giraldez AJ. 2016. Estrogens Suppress a Behavioral Phenotype in Zebrafish Mutants of the Autism Risk Gene, CNTNAP2. *Neuron* **89**:725–733. doi:10.1016/j.neuron.2015.12.039

Holler K, Neuschulz A, Drewe-Boß P, Mintcheva J, Spanjaard B, Arsiè R, Ohler U, Landthaler M, Junker JP. 2021. Spatio-temporal mRNA tracking in the early zebrafish embryo. *Nat Commun* **12**:3358. doi:10.1038/s41467-021-23834-1

Horn KE, Glasgow SD, Gobert D, Bull S-J, Luk T, Girgis J, Tremblay M-E, McEachern D, Bouchard J-F, Haber M, Hamel E, Krimpenfort P, Murai KK, Berns A, Doucet G, Chapman CA, Ruthazer ES, Kennedy TE. 2013. DCC expression by neurons regulates synaptic plasticity in the adult brain. *Cell Rep* **3**:173–185. doi:10.1016/j.celrep.2012.12.005

Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, Collins JE, Humphray S, McLaren K, Matthews L, McLaren S, Sealy I, Caccamo M, Churcher C, Scott C, Barrett JC, Koch R, Rauch G-J, White S, Chow W, Kilian B, Quintais LT, Guerra-Assunção JA, Zhou Y, Gu Y, Yen J, Vogel J-H, Eyre T, Redmond S, Banerjee R, Chi J, Fu B, Langley E, Maguire SF, Laird GK, Lloyd D, Kenyon E, Donaldson S, Sehra H, Almeida-King J, Loveland J, Trevanion S, Jones M, Quail M, Willey D, Hunt A, Burton J, Sims S, McLay K, Plumb B, Davis J, Clee C, Oliver K, Clark R, Riddle C, Elliott D, Threadgold G, Harden G, Ware D, Begum S, Mortimore B, Kerry G, Heath P, Phillimore B, Tracey A, Corby N, Dunn M, Johnson C, Wood J, Clark S, Pelan S, Griffiths G, Smith M, Glithero R, Howden P, Barker N, Lloyd C, Stevens C, Harley J, Holt K, Panagiotidis G, Lovell J, Beasley H, Henderson C, Gordon D, Auger K, Wright D, Collins J, Raisen C, Dyer L, Leung K, Robertson L, Ambridge K, Leongamornlert D, McGuire S, Gilderthorp R, Griffiths C, Manthravadi D, Nichol S, Barker G, Whitehead S, Kay M, Brown J, Murnane C, Gray E, Humphries M, Sycamore N, Barker D, Saunders D, Wallis J, Babbage A, Hammond S, Mashreghi-Mohammadi M, Barr L, Martin S, Wray P, Ellington A, Matthews N, Ellwood

M, Woodmansey R, Clark G, Cooper JD, Tromans A, Grafham D, Skuce C, Pandian R, Andrews R, Harrison E, Kimberley A, Garnett J, Fosker N, Hall R, Garner P, Kelly D, Bird C, Palmer S, Gehring I, Berger A, Dooley CM, Ersan-Ürün Z, Eser C, Geiger H, Geisler M, Karotki L, Kirn A, Konantz J, Konantz M, Oberländer M, Rudolph-Geiger S, Teucke M, Lanz C, Raddatz G, Osoegawa K, Zhu B, Rapp A, Widaa S, Langford C, Yang F, Schuster SC, Carter NP, Harrow J, Ning Z, Herrero J, Searle SMJ, Enright A, Geisler R, Plasterk RHA, Lee C, Westerfield M, de Jong PJ, Zon LI, Postlethwait JH, Nüsslein-Volhard C, Hubbard TJP, Crollius HR, Rogers J, Stemple DL. 2013. The zebrafish reference genome sequence and its relationship to the human genome. *Nature* **496**:498–503. doi:10.1038/nature12111

Hu B, Lelek S, Spanjaard B, El-Sammak H, Simões MG, Mintcheva J, Aliee H, Schäfer R, Meyer AM, Theis F, Stainier DYR, Panáková D, Junker JP. 2022. Origin and function of activated fibroblast states during zebrafish heart regeneration. *Nat Genet* **54**:1227–1237. doi:10.1038/s41588-022-01129-5

Hutson LD, Chien C-B. 2002. Wiring the zebrafish: axon guidance and synaptogenesis. *Curr Opin Neurobiol* **12**:87–92. doi:10.1016/s0959-4388(02)00294-5

Jaworski A, Tessier-Lavigne M. 2012. Autocrine/juxtaparacrine regulation of axon fasciculation by Slit-Robo signaling. *Nat Neurosci* **15**:367–369. doi:10.1038/nn.3037

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**:583–589. doi:10.1038/s41586-021-03819-2

Klee EW. 2008. The zebrafish secretome. *Zebrafish* **5**:131–138. doi:10.1089/zeb.2008.0529

Kotlyar M, Pastrello C, Ahmed Z, Chee J, Varyova Z, Jurisica I. 2022. IID 2021: towards context-specific protein interaction analyses by increased coverage, enhanced annotation and enrichment analysis. *Nucleic Acids Res* **50**:D640–D647. doi:10.1093/nar/gkab1034

Kowalczyk W, Romanelli L, Atkins M, Hillen H, Bravo González-Blas C, Jacobs J, Xie J, Soheily S, Verboven E, Moya IM, Verhulst S, de Waegeneer M, Sansores-Garcia L, van Huffel L, Johnson RL, van Grunsven LA, Aerts S, Halder G. 2022. Hippo signaling instructs ectopic but not normal organ growth. *Science* **378**:eabg3679. doi:10.1126/science.abg3679

Krämer A, Green J, Pollard J, Tugendreich S. 2014. Causal analysis approaches in Ingenuity Pathway Analysis. *Bioinformatics* **30**:523–530. doi:10.1093/bioinformatics/btt703

Lees JA, Dias JM, Han S. 2021. Applications of Cryo-EM in small molecule and biologics drug design. *Biochem Soc Trans* **49**:2627–2638. doi:10.1042/BST20210444

Leng G, Pineda R, Sabatier N, Ludwig M. 2015. 60 YEARS OF NEUROENDOCRINOLOGY: The posterior pituitary, from Geoffrey Harris to our present understanding. *J Endocrinol* **226**:T173-185. doi:10.1530/JOE-15-0087

Lin Z, Akin H, Rao R, Hie B, Zhu Z, Lu W, Smetanin N, Verkuil R, Kabeli O, Shmueli Y, Costa A dos S, Fazel-Zarandi M, Sercu T, Candido S, Rives A. 2022. Evolutionary-scale prediction of atomic level protein structure with a language model. doi:10.1101/2022.07.20.500902

Liu F, Placzek M. 2014. Axon guidance effects of classical morphogens Shh and BMP7 in the hypothalamo-pituitary system. *Neurosci Lett* **562**:108–113. doi:10.1016/j.neulet.2013.11.046

Liu F, Pogoda H-M, Pearson CA, Ohyama K, Löhr H, Hammerschmidt M, Placzek M. 2013. Direct and indirect roles of Fgf3 and Fgf10 in innervation and vascularisation of the vertebrate hypothalamic neurohypophysis. *Development* **140**:1111–1122. doi:10.1242/dev.080226

Logan CV, Cossins J, Rodríguez Cruz PM, Parry DA, Maxwell S, Martínez-Martínez P, Riepsaame J, Abdelhamed ZA, Lake AVR, Moran M, Robb S, Chow G, Sewry C, Hopkins PM, Sheridan E, Jayawant S, Palace J, Johnson CA, Beeson D. 2015. Congenital Myasthenic Syndrome Type 19 Is Caused by Mutations in COL13A1, Encoding the Atypical Non-fibrillar Collagen Type XIII α1 Chain. *Am J Hum Genet* **97**:878–885. doi:10.1016/j.ajhg.2015.10.017

Manitt C, Nikolakopoulou AM, Almario DR, Nguyen SA, Cohen-Cory S. 2009. Netrin participates in the development of retinotectal synaptic connectivity by modulating axon arborization and synapse formation in the developing brain. *J Neurosci* **29**:11065–11077. doi:10.1523/JNEUROSCI.0947-09.2009

Meester JAN, Southgate L, Stittrich A-B, Venselaar H, Beekmans SJA, den Hollander N, Bijlsma EK, Helderman-van den Enden A, Verheij JBGM, Glusman G, Roach JC, Lehman A, Patel MS, de Vries BBA, Ruivenkamp C, Itin P, Prescott K, Clarke S, Trembath R, Zenker M, Sukalo M, Van Laer L, Loeys B, Wuyts W. 2015. Heterozygous Loss-of-Function Mutations in DLL4 Cause Adams-Oliver Syndrome. *Am J Hum Genet* **97**:475–482. doi:10.1016/j.ajhg.2015.07.015

Miyata S. 2017. Advances in Understanding of Structural Reorganization in the Hypothalamic Neurosecretory System. *Front Endocrinol (Lausanne)* **8**:275. doi:10.3389/fendo.2017.00275

Murai KK, Misner D, Ranscht B. 2002. Contactin Supports Synaptic Plasticity Associated with Hippocampal Long-Term Depression but Not Potentiation. *Current Biology* **12**:181–190. doi:10.1016/S0960-9822(02)00680-2

Nauroy P, Hughes S, Naba A, Ruggiero F. 2018. The in-silico zebrafish matrisome: A new tool to study extracellular matrix gene and protein functions. *Matrix Biol* **65**:5–13. doi:10.1016/j.matbio.2017.07.001

Nevers Y, Jones TEM, Jyothi D, Yates B, Ferret M, Portell-Silva L, Codo L, Cosentino S, Marcet-Houben M, Vlasova A, Poidevin L, Kress A, Hickman M, Persson E, Piližota I, Guijarro-Clarke C, the OpenEBench team the Quest for Orthologs Consortium, Iwasaki W, Lecompte O, Sonnhammer E, Roos DS, Gabaldón T, Thybert D, Thomas PD, Hu Y, Emms DM, Bruford E, Capella-Gutierrez S, Martin MJ, Dessimoz C, Altenhoff A. 2022. The Quest for Orthologs orthology benchmark service in 2022. *Nucleic Acids Research* **50**:W623–W632. doi:10.1093/nar/gkac330

Nisar S, Bhat AA, Masoodi T, Hashem S, Akhtar S, Ali TA, Amjad S, Chawla S, Bagga P, Frenneaux MP, Reddy R, Fakhro K, Haris M. 2022. Genetics of glutamate and its receptors in autism spectrum disorder. *Mol Psychiatry* **27**:2380–2392. doi:10.1038/s41380-022-01506-w

Oprişoreanu A-M, Smith HL, Krix S, Chaytow H, Carragher NO, Gillingwater TH, Becker CG, Becker T. 2021. Automated in vivo drug screen in zebrafish identifies synapse-stabilising drugs with relevance to spinal muscular atrophy. *Dis Model Mech* **14**:dmm047761. doi:10.1242/dmm.047761

Ornitz DM, Itoh N. 2015. The Fibroblast Growth Factor signaling pathway. *Wiley Interdiscip Rev Dev Biol* **4**:215–266. doi:10.1002/wdev.176

Palmer EE, Stuhlmann T, Weinert S, Haan E, Van Esch H, Holvoet M, Boyle J, Leffler M, Raynaud M, Moraine C, van Bokhoven H, Kleefstra T, Kahrizi K, Najmabadi H, Ropers H-

H, Delgado MR, Sirsi D, Golla S, Sommer A, Pietryga MP, Chung WK, Wynn J, Rohena L, Bernardo E, Hamlin D, Faux BM, Grange DK, Manwaring L, Tolmie J, Joss S, DDD Study, Cobben JM, Duijkers F a. M, Goehringer JM, Challman TD, Hennig F, Fischer U, Grimme A, Suckow V, Musante L, Nicholl J, Shaw M, Lodh SP, Niu Z, Rosenfeld JA, Stankiewicz P, Jentsch TJ, Gecz J, Field M, Kalscheuer VM. 2018. De novo and inherited mutations in the X-linked gene CLCN4 are associated with syndromic intellectual disability and behavior and seizure disorders in males and females. *Mol Psychiatry* **23**:222–230. doi:10.1038/mp.2016.135

Patton EE, Zon LI, Langenau DM. 2021. Zebrafish disease models in drug discovery: from preclinical modelling to clinical trials. *Nat Rev Drug Discov* **20**:611–628. doi:10.1038/s41573-021-00210-8

Pearson CA, Placzek M. 2013. Development of the medial hypothalamus: forming a functional hypothalamic-neurohypophyseal interface. *Curr Top Dev Biol* **106**:49–88. doi:10.1016/B978-0-12-416021-7.00002-X

Pires-daSilva A, Sommer RJ. 2003. The evolution of signalling pathways in animal development. *Nat Rev Genet* **4**:39–49. doi:10.1038/nrg977

Raj B, Farrell JA, Liu J, El Kholtei J, Carte AN, Navajas Acedo J, Du LY, McKenna A, Relić Đ, Leslie JM, Schier AF. 2020. Emergence of Neuronal Diversity during Vertebrate Brain Development. *Neuron* **108**:1058-1074.e6. doi:10.1016/j.neuron.2020.09.023

Romanov RA, Zeisel A, Bakker J, Girach F, Hellysaz A, Tomer R, Alpár A, Mulder J, Clotman F, Keimpema E, Hsueh B, Crow AK, Martens H, Schwindling C, Calvigioni D, Bains JS, Máté Z, Szabó G, Yanagawa Y, Zhang M-D, Rendeiro A, Farlik M, Uhlén M, Wulff P, Bock C, Broberger C, Deisseroth K, Hökfelt T, Linnarsson S, Horvath TL, Harkany T. 2017. Molecular interrogation of hypothalamic organization reveals distinct dopamine neuronal subtypes. *Nat Neurosci* **20**:176–188. doi:10.1038/nn.4462

Rosso L, Mienville J-M. 2009. Pituicyte modulation of neurohormone output. *Glia* **57**:235–243. doi:10.1002/glia.20760

Ruegg MA, Stoeckli ET, Kuhn TB, Heller M, Zuellig R, Sonderegger P. 1989. Purification of axonin-1, a protein that is secreted from axons during neurogenesis. *EMBO J* **8**:55–63. doi:10.1002/j.1460-2075.1989.tb03348.x

Schier AF. 2020. Single-cell biology: beyond the sum of its parts. *Nat Methods* **17**:17–20. doi:10.1038/s41592-019-0693-3

Schier AF, Shen MM. 2000. Nodal signalling in vertebrate development. *Nature* **403**:385–389. doi:10.1038/35000126

Ségalat L. 2007. Invertebrate animal models of diseases as screening tools in drug discovery. *ACS Chem Biol* **2**:231–236. doi:10.1021/cb700009m

Shafer MER, Sawh AN, Schier AF. 2022. Gene family evolution underlies cell-type diversification in the hypothalamus of teleosts. *Nat Ecol Evol* **6**:63–76. doi:10.1038/s41559-021-01580-3

Shao X, Liao J, Li C, Lu X, Cheng J, Fan X. 2021. CellTalkDB: a manually curated database of ligand-receptor interactions in humans and mice. *Brief Bioinform* **22**:bbaa269. doi:10.1093/bib/bbaa269

Shao X, Taha IN, Clauser KR, Gao Y (Tom), Naba A. 2020. MatrisomeDB: the ECM-protein knowledge database. *Nucleic Acids Research* **48**:D1136–D1144. doi:10.1093/nar/gkz849

Shen MM. 2007. Nodal signaling: developmental roles and regulation. *Development* **134**:1023–1034. doi:10.1242/dev.000166

Steinhart Z, Angers S. 2018. Wnt signaling in development and tissue homeostasis. *Development* **145**:dev146589. doi:10.1242/dev.146589

Stelzer G, Rosen N, Plaschkes I, Zimmerman S, Twik M, Fishilevich S, Stein TI, Nudel R, Lieder I, Mazor Y, Kaplan S, Dahary D, Warshawsky D, Guan-Golan Y, Kohn A, Rappaport N, Safran M, Lancet D. 2016. The GeneCards Suite: From Gene Data Mining to Disease Genome Sequence Analyses. *Current Protocols in Bioinformatics* **54**:1.30.1-1.30.33. doi:10.1002/cpbi.5

Straussman R, Morikawa T, Shee K, Barzily-Rokni M, Qian ZR, Du J, Davis A, Mongare MM, Gould J, Frederick DT, Cooper ZA, Chapman PB, Solit DB, Ribas A, Lo RS, Flaherty KT, Ogino S, Wargo JA, Golub TR. 2012. Tumour micro-environment elicits innate resistance to RAF inhibitors through HGF secretion. *Nature* **487**:500–504. doi:10.1038/nature11183

Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT, Legeay M, Fang T, Bork P, Jensen LJ, von Mering C. 2021. The STRING database in 2021: customizable protein-protein networks, and functional characterization of user-uploaded gene/measurement sets. *Nucleic Acids Res* **49**:D605–D612. doi:10.1093/nar/gkaa1074

Team RC. 2013. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. *http://www R-project org/*.

The UniProt Consortium. 2021. UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Research* **49**:D480–D489. doi:10.1093/nar/gkaa1100

Thumuluri V, Almagro Armenteros JJ, Johansen AR, Nielsen H, Winther O. 2022. DeepLoc 2.0: multi-label subcellular localization prediction using protein language models. *Nucleic Acids Res* gkac278. doi:10.1093/nar/gkac278

Tunyasuvunakool K, Adler J, Wu Z, Green T, Zielinski M, Žídek A, Bridgland A, Cowie A, Meyer C, Laydon A, Velankar S, Kleywegt GJ, Bateman A, Evans R, Pritzel A, Figurnov M, Ronneberger O, Bates R, Kohl SAA, Potapenko A, Ballard AJ, Romera-Paredes B, Nikolov S, Jain R, Clancy E, Reiman D, Petersen S, Senior AW, Kavukcuoglu K, Birney E, Kohli P, Jumper J, Hassabis D. 2021. Highly accurate protein structure prediction for the human proteome. *Nature* **596**:590–596. doi:10.1038/s41586-021-03828-1

Türei D, Valdeolivas A, Gul L, Palacio-Escat N, Klein M, Ivanova O, Ölbei M, Gábor A, Theis F, Módos D, Korcsmáros T, Saez-Rodriguez J. 2021. Integrated intra- and intercellular signaling knowledge for multicellular omics analysis. *Molecular Systems Biology* **17**:e9923. doi:10.15252/msb.20209923

Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C, Sjöstedt E, Asplund A, Olsson I, Edlund K, Lundberg E, Navani S, Szigyarto CA-K, Odeberg J, Djureinovic D, Takanen JO, Hober S, Alm T, Edqvist P-H, Berling H, Tegel H, Mulder J, Rockberg J, Nilsson P, Schwenk JM, Hamsten M, von Feilitzen K, Forsberg M, Persson L, Johansson F, Zwahlen M, von Heijne G, Nielsen J, Pontén F. 2015. Proteomics. Tissue-based map of the human proteome. *Science* **347**:1260419. doi:10.1126/science.1260419

Wittkowski W. 1998. Tanycytes and pituicytes: morphological and functional aspects of neuroglial interaction. *Microsc Res Tech* **41**:29–42. doi:10.1002/(SICI)1097-0029(19980401)41:1<29::AID-JEMT4>3.0.CO;2-P

Xie Y, Kuan AT, Wang W, Herbert ZT, Mosto O, Olukoya O, Adam M, Vu S, Kim M, Tran D, Gómez N, Charpentier C, Sorour I, Lacey TE, Tolstorukov MY, Sabatini BL, Lee W-CA, Harwell CC. 2022. Astrocyte-neuron crosstalk through Hedgehog signaling mediates cortical synapse development. *Cell Rep* **38**:110416. doi:10.1016/j.celrep.2022.110416

Ximerakis M, Lipnick SL, Innes BT, Simmons SK, Adiconis X, Dionne D, Mayweather BA, Nguyen L, Niziolek Z, Ozek C, Butty VL, Isserlin R, Buchanan SM, Levine SS, Regev A, Bader GD, Levin JZ, Rubin LL. 2019. Single-cell transcriptomic profiling of the aging mouse brain. *Nat Neurosci* **22**:1696–1708. doi:10.1038/s41593-019-0491-3

Zhang X, Ibrahimi OA, Olsen SK, Umemori H, Mohammadi M, Ornitz DM. 2006. Receptor Specificity of the Fibroblast Growth Factor Family: THE COMPLETE MAMMALIAN FGF FAMILY *. *Journal of Biological Chemistry* **281**:15694–15700. doi:10.1074/jbc.M601252200

Fig. 1 Chodkowski M. *et al.,*

Figure Legends:

Figure 1. Prediction of cellular localization of reference zebrafish proteome

(A) Graphic describing the DeepLoc 2.0-based prediction of cellular location of the reference zebrafish proteome.

(B) Zebrafish reference proteome cellular localization. Donut chart showing the major DeepLoc2.0 predictions of zebrafish proteome. The number of proteins (**B**) and protein-coding genes (**B′**) are shown.

(C) Venn diagram comparing the number of genes with existing extracellular annotation records and DeepLoc 2.0-based 'Extracellular' predictions.

(D) Venn diagram comparing genes with existing cell membrane localization annotation records and DeepLoc-based 'cell membrane' predictions.

A

Cellular localization annotation status

19281
(41.3%)

27406
(58.7%)

Annotated (UniprotKB or GO term)
Unannotated

B

DeepLoc 2.0 - Multilocalizing proteins

Figure 1 - figure supplement 1
Chodkowski M. *et al.,*

Figure 1 – figure supplement 1. Annotation status and DeepLoc 2.0 multilocalizing protein localizations

(A) Current status of the cellular localization annotation records of the reference proteome from zebrafish. Donut chart showing number of proteins with and without GO-term based or UniprotKB cellular localization records.

(B) Circle plot showing the number of proteins predicted to localize to multiple cellular regions and organelles.

Fig. 2 Chodkowski M. *et al.,*

Figure 2. Curated zebrafish ligands and membrane receptors

(A) Scheme describing the steps taken for the creation of zebrafish ligands and membrane receptors list.

(B) Donut chart showing the number and percentage of zebrafish genes coding for curated receptor- and ligand-coding genes.

(C) Venn diagram comparing curated ligands with DeepLoc2.0 extracellular predicted proteins and previously reported Matrisome dataset.

(D) Conservation between zebrafish and human ligand- and membrane receptor-coding genes based on ZFIN records.

(E) Venn diagram comparing the number of curated ligand- and receptor- coding genes with zebrafish orthologues of human disease-linked genes in the OMIM database.

(F) Dot plot showing the diseases associated with human orthologues of some of the zebrafish ligands and membrane receptors.

(G) Venn diagram showing the number and percentage of zebrafish membrane receptors and ligands targetable by drugs.

(H) Donut chart showing the number and percentage of drugs that can potentially target zebrafish receptors and/or ligands.

Fig. 3 Chodkowski M. *et al.,*

Figure 3. Zebrafish ligand-receptor interactome

(A) Scheme describing steps undertaken to map the zebrafish ligands and membrane receptors interactome (A). Circle plot showing the top 25 ranked ligand-receptor pairs (A').

(B) Orphan status of ligands and membrane receptors. Donut chart showing the number and percentage of genes coding for ligands and receptors with or without potential receptors and ligands respectively.

(C) Zebrafish ligand-receptor interactome stats. Venn diagram comparing ligand-receptor pairs across different ranking categories (C). Graph showing # of putative receptors/ligand and ligands/receptor (C').

(D) Receptors for Nodal-related TGF-beta ligands and Wnt ligands. Circle plot showing the top ranked receptors for Ndr2, Ndr1 and Spaw ligands (D) and Wnt ligands (D').

Figure 3 - figure supplement 1
Chodkowski M. *et al.,*

Figure 3 – figure supplement 1.

Conservation of zebrafish ligand-receptor interactions. Violin plot showing STRING database zebrafish ligand-receptor interactions scores to their human orthologue ligand-receptor records in IID database. 'Pred' denotes predicted interactions.

Fig. 4 Chodkowski M. *et al.,*

Figure 4. Application of *DanioTalk*

(A) Scheme showing synaptosome-PSD from adult zebrafish brain.

(B) Circle plot showing the top-ranked ligand-receptor interactions between the synaptosome and PSD (**B** and **B'**).

(C) Scheme showing axo-glial interaction between OXT neuron and glial pituicytes.

(D) Circle plot showing top-ranked ligand-receptor interactions between OXT neurons and glial pituicytes.

A Synaptasome

3241
(91%)   147
(4%)   0
        0
182   0
(5%)
        0

Protein coding genes
- Synaptosome
- Ligands
- Receptors

B PSD

1551
(91%)   74
(4%)   0
        0
86   0
(5%)
        0

Protein coding genes
- PSD
- Ligands
- Receptors

C OXT neurons

2129
(93%)   104
(4.5%) 0
        0
66   0
(3%)
        0

Protein coding genes
- OXT neurons
- Ligands
- Receptors

D Glial pituicytes

1084
(81%)   172
(13%)   0
        0
82   0
(6%)
        0

Protein coding genes
- Glial pituicytes
- Ligands
- Receptors

Figure 4 – figure supp 1
Chodkowski M. *et al.,*

Figure 4 – figure supplement 1. Application of *DanioTalk*

(A) Venn diagram showing the number and percentage of genes coding for synaptosome-enriched ligands and receptors.

(B) Venn diagram showing the number and percentage of genes coding for PSD-enriched ligands and receptors.

(C) Venn diagram showing the number and percentage of genes coding for OXT neuron-enriched ligands and receptors.

(D) Venn diagram showing the number and percentage of genes coding for Glial pituicyte-enriched ligands and receptors.