Prenatal maternal stress promotes neural stem cell proliferation in the ependymal-subventricular zone of adult male offspring

Z. Wang[1,*], L.Q. Zhou[1,*,†], C.M. Lee[2, *, ψ], J.Y. Shi[1], L. Liu[2], X.J. Yang[1], Y.X. Deng[1], J.P. Liu[3], J.B. Wang[3], W.M. Zhu[1,3], Y.E. Sun[1,3,†], and Q. Lin[1,2,†]

[1] Shanghai Institute of Stem Cell Research and Clinical Translation, Shanghai East Hospital, School of Medicine, Tongji University, Shanghai, China.

[2] Department of Psychiatry and Behavioral Sciences, Intellectual Development and Disabilities Research Center, University of California, Los Angeles, CA 90095

[3] Stem Cell Translational Research Center, Tongji Hospital, Tongji University, Shanghai, China.

[*] Equal contribution

[ψ] Current address: Des Moines University, 3200 Grand Ave. Des Moines, IA 50312

[†] To whom correspondence may be addressed. Q.L. qlin@g.ucla.edu, Y.E.S. yievesun@gmail.com, or jason_li_zhou@126.com

Abstract

Perturbations during critical time windows in the normal series of development can lead to adverse functional consequences that manifest later in life. Here, we report that prenatal stress (PNS) during the last week of gestation (E14-delivery) dramatically increased the number of proliferating neural stem/progenitor cells (NSC/NPCs) in the ependymal-subventricular zone (E-SVZ) of male mouse offspring. PNS did not cause significant cell death or a deficit in neuroblast migration to the olfactory bulb (OB). Olfactory behavioral tests showed that while prenatally stressed male mice displayed normal olfactory function in differentiating nonsocial odors, these mice showed impairment in discriminating different social smells. Bulk and single nucleus transcriptomic analyses combined with rescue assays using mitogen-activated protein kinases (MAPK) phosphorylation inhibitors revealed that PNS exposure during the critical period of neurogenesis promoted NSC/NPC proliferation in adult male offspring by sustaining MAPK3/1 activity. Compared to prenatally stressed males, their female littermates showed less change in the number of proliferating cells in the E-SVZ.

Introduction

The prenatal period represents a critical time window open to neural plasticity changes and is most vulnerable to the influence of external disturbances. In humans and model animals, environmental disturbances during the critical period have been shown to cause neural morphological, neurobiological, and behavioral deficits[1-11].

Since the evidence of cell mitosis in the SVZ of adult rats was published over 100 years ago[12], it has now been widely accepted that adult neurogenesis occurs in mammals[13-17]. Numerous studies have reported that the ependymal-subventricular zone (E-SVZ) and the dentate gyrus (DG) are the main brain regions that harbor NSC/NPCs to give rise to neurons and glial cells throughout life[14, 15, 18-22]. Mechanistically, EGF and VEGF are two key growth factors to activate dormant/quiescent neural stem cell and neural progenitor cell proliferation in the E-SVZ of the lateral and/or the 4th ventricles in mice[14, 19]. MAPK (a.k.a. extracellular signal-regulated kinase (ERK)), as a major transducer of growth factors like EGF and VEGF, plays a vital role in mediating their intracellular signaling[23, 24]. MAPK(ERK) has a TxY motif in its activation domain, which requires dual phosphorylation by upstream MAPK kinases (e.g., MAP2K1/2 (MEK1/2)) for catalytic activation. In cancer cells, MAPK3 and 1 (ERK1/2) have been shown to interact with wide-range signaling pathways, such as EGF/FGF/VEGF, WNT, BMP/TGFB, Estrogen, SHH[25-29] to regulate cell proliferation and to facilitate epithelial-mesenchymal transition (EMT)[30, 31].

In the past decade or so, studies reported that prenatal maternal environmental disturbances have profound impact on hippocampal neurogenesis but spare OB neurogenesis in postnatal offspring[32-35]. Although tremendous progress has been made to understand the long-lasting effects of PNS on brain development, little is known about the molecular programs that govern adult NSC/NPC proliferation and neurogenesis in response to the prenatal

stimuli. In the current study, by employing daily mild restraint stress during the last week of gestation, we found that PNS increased the numbers of BrdU$^+$ and MKI67$^+$ proliferating cells in the E-SVZ and the rostral migratory stream (RMS), and PNS also increased BrdU$^+$ cells in the OB. To reveal the underlying molecular program that mediates PNS triggered cell proliferation in adult brain, we carried out bulk and single nucleus transcriptomic analyses. Combined with rescue assays, we showed that PNS exposure at the late stage of embryonic development sustained postnatal phosphorylation levels of MAPK3/1, which in turn promoted NSC/NPC proliferation in the E-SVZ of adult male offspring. By olfactory behavioral tests, we found that prenatally stressed male mice had an intact olfactory function to locate food pellets and discriminate nonsocial odors, whereas they showed impairment in differentiation of different social smells.

Results

PNS triggers cell proliferation in the lateral E-SVZ. To evaluate the effects of the prenatal aversive stimulus on cell proliferation in the E-SVZ, we employed a widely used stress-inducing method, the restraint stress on pregnant dams during the third week of gestation (60min/day, 0800h-0900h)[36]. The third week of gestation in mice roughly corresponds to the second trimester of human gestation, which is considered to be a critical period for the development of psychiatric disorders[37, 38]. To eliminate the postnatal influence from stressed mothers, the prenatally stressed (STR) pups were fostered by unstressed dams at postnatal day 0 (P0). As a control (CTL), pups from unstressed dams also underwent the fostering process. BrdU was administered by intraperitoneal (IP) injection one hour before tissue collection (Fig.1A).

Compared to the CTL, the ratio of proliferating cells to the total number of cells in the lateral E-SVZ (MKI67$^+$ cells/total E-SVZ cells) was dramatically increased in STR male offspring at P60 and P180, whereas similar increases were not detected in neonatal or juvenile mice (P0.5-21) (Fig.1B; Sup.Fig.1). We found that the numbers of BrdU$^+$/PROM1$^+$/GFAP$^-$ and MKI67$^+$/PROM1$^+$/GFAP$^-$ ependymal cells, and MKI67$^+$/DCX$^+$, BrdU$^+$/DCX$^+$, and MKI67$^+$/TUBB3$^+$ neuroblasts increased dramatically in STR male mice (Figs.1C, D, 2A; Sup.Fig.3A). There were more BrdU$^+$ and MKI67$^+$ migrating neuroblasts in the rostral migratory stream (RMS) of stressed male mice than that of the CTL mice (Fig.2B). We did not observe statistically significant changes in the numbers of GFAP$^+$/BrdU$^+$ cells in either female or male offspring (Sup. Figs.2D, 3B).

In the female littermates, aforementioned changes were statistically insignificant, suggesting that female offspring were resilient to PNS (Sup.Figs.2, 4A,B). We found the ratio of proliferating cells to the total number of cells in the E-SVZ in CTL female mice were higher than that of their male littermates (BrdU$^+$/PROM1$^+$ cells: female = 2.1 ± 0.4% vs. male = 0.9 ± 0.2%; MKI67$^+$/PROM1$^+$ cells: female = 4.1 ± 0.4% vs. male = 2.7 ± 0.3%. mean ± SEM). A similar finding was demonstrated in the subgranular zone of female mice by Tanapat et al[39]. It seems that the influence of PNS exposure is highly sex dependent. Similar findings have been observed in both rats and humans. In rats, PNS significantly increased anxiety- and depression-related behaviors in male offspring,

but not in female offspring[40, 41]. In humans, it has been shown that prenatal disturbances preferentially affect males in a late-onset manner[42, 43].

E-SVZ NSC/NPC progenies migrate to and mature in the OB. To observe adult neurogenesis in the OB of STR offspring, we labeled proliferating cells with two BrdU administrations separated by a 12hr interval. The animals were sacrificed six weeks later (Fig.3A). BrdU labeled cells migrated to the OB and mainly resided in the granule cell layer (GrO) with a few cells in the glomerular layer (Gl) and the external plexiform layer (EPl) in both CTL and STR offspring (8-9-month-old). As expected, the number of BrdU-labeled, newborn RBFOX3$^+$ neurons increased in the OB of stressed mice (Fig.3B, C). We did not observe substantial cell death in the OB of either the CTL or STR mice (Fig.3D).

PNS promotes E-SVZ cell proliferation via regulating MAPK pathway. To reveal the underlying molecular programs of PNS-increased cell proliferation, we carried out transcriptomic analyses. 49 eight-month-old STR male and female mice were sacrificed, the brains were sectioned, and bilateral E-SVZ regions were dissected out, rapidly frozen using liquid nitrogen, and stored at -80°C. Tissues from left and right sides were stored separately: one for bulk RNA sequencing, and one for single nucleus RNA sequencing (snRNAseq) (Fig.4A).

Weighted gene co-expression network analysis (WGCNA) and gene ontology (GO) enrichment analysis of bulk RNAseq data implied that PNS dysregulated gene expression networks related to the regulation of three biological functions: transferase activity (*Dusps1, 6, 5, Rgs, Spry2*; the purple module), cellular response to morphogens (*Shisa2*, *Ccn1*, *Spry2*, *Bambi*, *Smad7*; the purple module), and gliogenesis (*Mag*, *Trf*, *Myrf*, *Plp1*, *Gjc2*; the magenta module) in adult male offspring (Fig.4B, C). Many genes in the purple module are critical components of RAS-RAF-MEK-MAPK(ERK) signaling cascade, which has been shown to be the major downstream of growth factor signaling (e.g., EGF, FGF, VEGF) to promote cell proliferation[14, 18, 44-47]. GO enrichment analysis of the greenyellow and pink modules did not identify any signaling pathways. The major sex differences between males and females were the regulation of the immune effector process and cytokine production (the yellow module) (Fig.4B,C, Sup.Fig.6A)[48].

To further examine PNS elicited changes in the E-SVZ, we carried out snRNAseq. The nuclei isolated from the E-SVZ cells of different litters were pooled, purified, and subjected to droplet-based snRNAseq analysis (Fig.4A). Quality control filtering and integration of CTL and STR datasets were performed using the Seurat package[49] and the SingCellaR package[50, 51], which generated a merged dataset consisting of 7,407 cells and 20,634 genes for further analysis. According to marker gene expression, we annotated the 18 clusters to 10 different types of cells: 1) *Foxj1*$^+$ ependymal cell (EpC, cluster 6); 2) *Prom1*$^+$ endothelial-like cell (ElC, cluster 17); 3) dorsal neural progenitor cell (NPCd, cluster 14); 4) neuroblasts (NB, clusters 7 and 11); 5) *Drd1*$^+$ spiny projection neurons (D1 SPN, clusters 2, 3, 10, 12); 6) *Drd2*$^+$ SPN (D2 SPN, clusters 1, 8, 13); 7) ventral neural progenitor cell (NPCv, cluster 16); 8) astrocytes (AC, clusters 4, 5); 9) oligodendrocytes (ODC, cluster 9) and

oligodendrocyte precursor cells (OPC, cluster 18); 10) microglia (MG, cluster 15) (Fig.4D,Sup.Fig.6C). Cluster 11 NBs are dividing cells expressing *Mki67* and *Top2a*, while cluster 7 NBs are postmitotic cells. Cluster 14 NPCs express *Egfr*, *Mki67*, and *Btg2*, but not the NB markers *Dcx* and *Tubb3*. Astrocytes are separated into *Aqp4⁻* (cluster 4) and *Aqp4⁺* (cluster 5) cells. We detected a significant number of D1 and D2 SPNs that reside close to the E-SVZ (Sup.Fig.6B,C). Compared to the CTL, there were no obvious cell type changes in STR samples (Sup. Fig.5C).

To determine potential intrinsic programs that control NSC/NPC proliferation, we reanalyzed proliferating neural lineage cells. These cells express at least one of the cell cycle markers, *Mki67*, *Mcm2*, *Top2a*, and *Pcna* (UMI>0). We identified eight types of cells: EpC (cluster 7), ElC (cluster 8), NPC (cluster 3), NB (clusters 4 and 6), AC (cluster 5), D1 SPN (cluster 1), D2 SPN (cluster 2), and 'eccentric' eSPN (cluster 9) (Fig.4E, Sup.Fig.6C)[52, 53]. Cell cycle position estimation analysis (estimating a continuous cell-cycle pseudotime by principal component analysis of cell-cycle genes) showed that most cells in cluster 3 (NPC), cluster 4 (NB), and some cells in cluster 8 (ELC) were actively dividing cells, while cells of other clusters were predicted to be in the G1-phase (Sup.Fig.6D, cell cycle score)[54]. Cells in cluster 3 (NPC, *Egfr⁺, GFAP⁻*) were likely to be the previously described activated NSC (aNSC) and/or transit-amplifying cells (TACs, C cells)[55]. Cluster 5 astrocytes expressed *Gfap* and *Pdgfrb*, but not *Egfr* and *Prom1*, suggesting an astroglial lineage (Sup.Fig.6B). It remains to be determined why a small subset of spiny projection neurons (D1, D2, and eSPN) expresses low levels of cell cycle markers. It seems that in addition to olfactory bulb neurons, this region could produce new striatal neurons[56]. Another possibility is that cells in or close to the E-SVZ could sense some environmental cues, and thus express cell cycle markers.

We next employed the bigSCale2 algorithm to construct gene regulatory networks for the proliferating cells[57] (Fig. 4F). Compared to the CTL network, the STR network showed a lower edge to node density (CTL = 8.17, STR = 7.63) and a higher modularity (CTL = 0.41, STR = 0.54), suggesting weaker intramodular connectivity but higher gene expression heterogeneity in the STR network. The analysis implicated two major network modules that regulate cell cycle progression and neural maturation (Fig.4F). GO analysis of differentially expressed hub genes showed that PNS promoted expression of genes that regulated cell cycle progression (e.g., *Atad2*, *Top2a*, *Brd8*, *Atad5*, *Smchd1*, *Ncapg2*, *Pola1*, *Ezh2*) and neuronal differentiation (e.g., *Sox2*, *Sox5*, *Nfib*, Zmiz1, *Zeb2*), while it downregulated expression levels of genes modulating synapse organization and membrane potential (e.g., *Slc8a2*, *Kcnd2*, *Grm5*, *Nfia*, *Ntrk3, Slit3*) (Fig.4F,G). More specifically, PNS enhanced the expression of bromodomain containing genes *Atad2* (ATPase family AAA domain containing 2) and *Brd8* (bromodomain containing 8) that have been shown to be involved in ESR1 (estrogen receptor alpha) mediated transcriptional regulation of cell proliferation in cancer cell lines[58-60]. PNS increased expression levels of Wnt pathway modulators *Ezh2, Tcf4*, *Ephb2*, and *Sox2*, which play important roles in sustaining cell stemness and promoting cell proliferations[61-65]. The Wnt and the estrogen signaling pathways are tightly associated with

MAPK3/1 signaling to stimulate tumorigensis[26, 66, 67]. Moreover, MAP4K4 (serine/threonine kinase mitogen-activated protein 4 kinase 4) has been identified as a positive regulator of MAPK(ERK) signaling in cancer cells[68]. In addition, previous studies by our group and others have shown that MAPK3/1 (ERK1/2) are essential in maintaining proliferation of adult NSC/NPCs[68-70]. Considering the above transcriptomic analysis, we hypothesized that PNS promoted cell proliferation via the activation of the MAPK(ERK) signaling in the E-SVZ NSC/NPCs of adult male offspring.

PNS regulates MAPK3/1 phosphorylation. To test this hypothesis, we first employed immunostaining approach to examine phosphorylation levels of MAPK3/1 in the E-SVZ of both CTL and STR mice. The antibody was developed to recognize phosphorylated MAPK3/1 at threonine 202/185 and tyrosine 204/187 loci specifically (ThermoFisher 44-680G). We found that immunostaining of phosphorylated MAPK (p-MAPK) showed a punctate pattern (Sup.Fig.8A). This pattern was affirmed by using another p-MAPK3/1 specific antibody from a different vendor (Cell Signaling, 4370) and an antibody against unphosphorylated MAPK(ERK) (Cell Signaling, 4695) (Sup.Fig.7A,B). Furthermore, when treating the section with lambda protein phosphatase (Lambda PP) before primary antibody incubation, the immunoreactivity of the p-MAPK3/1 antibody was completely abolished, suggesting the high specificity of the antibody in recognizing p-MAPK (Sup.Fig.7C).

Since both phosphorylated and unphosphorylated MAK3/1 staining displayed punctate patterns, we reasoned that the number of puncta correlates with the status of phosphorylation. By employing this quantitative methodology, we next examined the phosphorylation of MAPK3/1 in the E-SVZ at different postnatal developmental time points (P0.5-180). Concomitant with the cell proliferation finding (Fig.1B, Sup.Fig.1), we found phosphorylation level of MAPK3/1 was considerably upregulated in the E-SVZ of stressed adult offspring (P60, P180) but not at earlier stages (Sup.Fig.8A,B).

To definitively verify the role of the MAPK3/1 pathway in regulating cell proliferation in stressed male mice, we carried out rescue assays by treating the STR and CTL animals with a selective MAPK3/1 inhibitor, LY3214996, and an inhibitor of upstream protein kinases MAP2K1/2 (MEK1/2), GSK1120212. We found that administration of these inhibitors effectively reduced the numbers of p-MAPK3/1 puncta and proliferating cells in the E-SVZ of both CTL and STR mice (Fig.5). Taken together, the rescue assays suggested that MAPK(ERK) signaling played a critical role in promoting NSC/NPC proliferation in STR adult male mice.

Discussion

Psychosocial, cultural, and environmental stressors experienced during gestation can be detrimental to pregnancy as well as maternal and fetal health. Recent studies have suggested that prenatal stress can shape neural circuitry, neural cell fate, and behavioral development in offspring, which may set the stage for postnatal disease vulnerability[7, 9-11]. In the current study, we showed that prenatal restraint stress sustained MAPK3/1 phosphorylation levels, thereby increasing the numbers of adult-stage BrdU[+]/PROM1[+] and DCX[+]/TUBB3[+]

NSC/NPCs in the lateral E-SVZ and BrdU$^+$/RBFOX3$^+$ newborn neurons in the OB of male offspring. This finding is in line with a previous study that showed 'mild' prenatal stress (30min/day, E15-17) enhanced neurogenesis in the DG of neonatal rat brains[71]. Nevertheless, other studies showed that PNS inhibited neurogenesis in the DG under severe stress protocols[32-35], suggesting that different stress levels ranging from mild to severe can have different negative influences on the neural development of offspring.

Interestingly, snRNAseq data analysis suggested that PNS upregulated the expression level of *Zeb2* (zinc finger E-box binding homeobox 2) and its downstream target *Sox6*[72] (Fig.4F). ZEB2 is one of the essential transcription factors that induces epithelial-mesenchymal transition (EMT) during pluripotent stem cell differentiation, embryonic development, wound healing, and tumorigenesis[73-75]. Studies by Miquelajauregui et al. and Deryckere et al. found that ZEB2 plays a critical role in controlling neurogenesis in prenatal and postnatal brains, repsectively[72, 76]. Moreover, Wnt and MAPK(ERK) pathways have been shown to induce *Zeb2* expression and promote EMT in cancer cells[73, 74]. MAPK(ERK) is a potent signaling transducer of EGF and VEGF, which are essential for promoting dormant/quiescent neural stem cell activation and neural progenitor cell proliferation[14, 19, 24, 45, 69, 70, 77]. Therefore, it is possible that PNS could activate dormant or quiescent neural stem cell proliferation via promoting MAPK(ERK)-mediated EMT in the germinal zones of adult brain.

To examine the possible association between PNS and basic olfactory function, we carried out the buried food test and the habituation/dishabituation test. Our initial study showed that STR male mice (5-8 months old) had a largely intact olfactory function to locate food pellets and discriminate nonsocial odors, while these animals showed impairment in differentiation of different social odors and habituation of social odor 2 (STR Social 1-3 median=2.98 sec to Social 2-1 median=2.58sec; CTL Social 1-3 median=1.34sec to Social 2-1 median=4.66sec) (Sup.Fig.5B-D). Calbindin 2 (CALB2, a.k.a. Calretinin) staining showed a disorganized glomerular layer (Gl) and a slightly increased number of CALB2$^+$ cells in the external plexiform layer (EPl) and the granule cell layer (GrO) of the OB in stressed male mice (Sup.Fig.5A). Glomeruli and granular cells have been shown to play essential roles in the regulation of olfactory habituation behavior[78, 79]. Moreover, scRNAseq analysis showed that PNS attenuated expression of genes modulating synapse organization and membrane potential (Fig.4G). For example, PNS upregulated the expression of glutamate metabotropic Receptor 5 (*Grm5*, a.k.a. mGluR5), which has been shown to regulates granule cell-mediated inhibition and odor discrimination[80, 81]. Taken together, immunostaining and snRNAseq analyses showed that environmental perturbation during the peak of prenatal neurogenesis elicited great changes in neuronal cytoarchitecture and gene expression in the OB of postnatal offspring. These changes may directly contribute to the behavioral deficit in discriminating different social odors[82-85].

Materials and Methods
Experimental Animals. Mice were kept and fed in standard conditions on a 12hr light/dark cycle. Experimental

procedures on animals were performed in accordance with the guidelines of UCLA Institutional Animal Care and Use Committee and UCLA Animal Research Committee, and the Regional Committee for Medical Research Ethics of Tongji University. The inbred strain C57BL⁄6NCrlVr mice were used in this study. Pregnancy was timed by daily monitoring for vaginal plugs.

Restraint stress and tissue collection. To induce stress between gestational days 14 and delivery (~E18.5), the pregnant dam was placed individually in a plastic restrainer fitted closely to body size for 60 min/day (session at 0800h). Control pregnant females were left undisturbed in their home cages. After delivery, the prenatally stressed (STR) and the CTL pups were fostered by unstressed, same strain mothers at P0. All pups were weaned at P21. For tissue collection, the animals were sacrificed, and brains were removed for immunostaining or RNA sequencing.

Buried food and olfactory habituation/dishabituation tests.

The tests were described by Yang and Crawley[86]. Details are given in SI Appendix.

Bromodeoxyuridine (BrdU) administration. The BrdU injection was described in our previous study[49]. Details are given in SI Appendix.

Immunohistochemistry. The primary antibodies used in this study were BrdU (Abcam ab6326, 1:100; ab1893 1:250), MKI67 (Abcam ab15580, 1:200), GFAP (ThermoFisher PA1-10004, 1:500), TUBB3 (R&D MAB1195, 1:500), CALB2 (Synaptic Systems 214111, 1:200), DCX (Santa Cruz sc-8066, 1:400), RBFOX3 (Abcam ab177487, 1:500), p-MAPK3/1(ThermoFisher 44-680G, 1:100), p-MAPK3/1 (Cell Signaling 4370, 1:200), and MAPK3/1 (Cell Signaling 4695, 1:250). Fluorophore-conjugated secondary antibodies were purchased from ThermoFisher Scientific and Jackson ImmunoResearch.

Terminal deoxynucleotidyl transferase dUTP Nick-End Labeling (TUNEL) assay. TUNEL assay was carried out following manufacturer's instructions (*In Situ* Cell Death Detection Kit, Roche-11684795910).

Tissue processing for BrdU immunostaining. Details are given in SI Appendix.

Transcriptomic analysis. The brains of 7- or 8-month-old STR and CTL offspring were sectioned at 300μm thickness in artificial cerebrospinal fluid supplied with actinomysin D (3μM, Sigma-Aldrich). The lateral walls of ependymal SVZ (E-SVZ) were dissected, and snap frozen in liquid nitrogen, and stored in air-tight tubes in -80°C freezer.

Bulk RNAseq. Total RNA was extracted and purified using RNeasy Mini Kit (Qiagen 74104). 100ng of qualified RNA was subjected to VAHTS™ mRNA-seq V3 Library Prep Kit for Illumina® (Vazyme NR611) for library preparation, followed by quality assessment using Agilent Bioanalyzer 2100, and sequencing on Illumina NovaSeq 6,000. The mRNA sequencing reads were mapped to the reference Mouse reference genome GRCm38 using HISAT2 (version 2.2.1) and StringTie (version 2.1.4), and the mapped reads were assigned using featureCounts function of R package Rsubread (version 2.0.1). The counts were normalized using the DESeq2 R package (version 1.30.0)[87]. Pseudogenes (gene names with the prefix "Gm") were filtered out, and top 10,000 highly variable genes were kept for weighted gene co-expression network analysis (WGCNA) using the R package WGCNA (version 1.69)[88]. Specifically, a soft-power of 12 was chosen to construct a topological overlap matrix from the gene correlation network. Modules were determined by the Dynamic Hybrid Cut algorithm using a "deepSplit" parameter of 2. Highly correlated modules (Pearson correlation of module eigengene > 0.9) were merged as one module using a "mergeCutHeight" parameter of 0.25. Gene ontology (GO) and KEGG pathway enrichment analyses of genes in each identified module were performed using the R package clusterProfiler (version 4.4.4)[89] set at default parameters.

Single nucleus RNAseq (snRNAseq). Single nuclei were extracted from pooled frozen lateral walls of ependymal SVZ (E-SVZ) from CTL and STR mice using a previously described ultracentrifugation method[90] with some modifications. The single nuclei RNA sequencing library construction was performed using the Chromium Next GEM Single Cell 3' Reagent Kits v3.1 (10× Genomics) according to the user guide. The quality assessement of the libraries was conducted using Agilent Bioanalyzer 2100, and the sequencing was performed on a Illumina Novaseq 6000. The alignment, filtering, barcode assignment, and UMI counting of the sequenced reads were processed using the Cell Ranger pipeline (version 5.0.1, 10xGenomics). Quality control (QC) of cells and genes was performed using R package Seurat (version 4.1.1). Doublets were removed using R package scDblFinder (version 1.10.0). Additional downstream workflow (including further QC, sample integration, highly variable gene selection, dimensional reduction, clustering, cluster marker gene identification, and visualization) was processed via SingCellaR R package (version 1.2.1)[50, 51]. According to expression of the marker genes, along with well-established celltype specific gene sets, the 18 clusters were assigned to 11 major celltypes: dorsal neural progenitor cells (NPCd), ventral neural progenitor cells (NPCv), neuroblasts (NB), astrocytes (AC), ependymal cells (EpC), endothelia-like cells (ElC), *Drd1*+ SPN (D1), *Drd2*+ SPN (D2), oligodendrocyte progenitor cells (OPC), oligodendrocytes (ODC), and microglia (MG).

Cells (except for clusters without neuronal lineage potential, OPC, ODC and MG) expressing at least one of the four cell cycle related genes (*Mki67*, *Top2a*, *Mcm2*, and *Pcna*) were isolated and re-clustered via SingCellaR

workflow. The potentially dividing cells were clustered into 9 clusters, and annotated as neural progenitor cells (NPC), AC, NB, EpC, ElC, D1, D2, "eccentric" spiny projection neuron (eSPN), respectively. The cell-cycle position of each potentially dividing cell was estimated using the tricycle R package (version 1.4.0)[54]. The putative gene-to-gene correlated network of the potential dividing cells was inferred using the bigSCale2 algorithm (version 2.0)[57]. The network centrality PageRank (PR) was chosen to represent gene essentiality. Gene communities in each network were identified, and those communities consisting of over 5 genes were considered as main communities. Hub genes with maximal biological relevance in each main community were defined as the intersection of three gene sets: 1) top 500 genes with the biggest change in PR absolute value between STR and CTL; 2) genes with over 0.75 quantile of PR value; 3) DEGs between STR and CTL determined using SingCellaR package (version 1.2.1)[50, 51]. GO enrichment analysis of the hub genes was performed using R package clusterProfiler (version 4.4.4)[89] set at default parameters. Details are gaven in SI Appendix.

Rescue assay. A MAPK3/1 inhibitor LY3214996 (Selleck, 16 mg/kg) and a MAP2K1/2 inhibitor GSK1120212 (Selleck, 0.35 mg/kg) and the vehicle solution were daily intraperitoneally (IP) injected to 9- to 10-month-old prenatally stressed male offspring for 27 days (except for the 11th and 22nd days), respectively. Details are given in SI Appendix.

Lambda Protein Phosphatase (Lambda PP) reaction. The assay was carried out following manufacturer's instructions (NEB, P0753S). Details are given in SI Appendix.

Imaging and Statistical Analysis. At least three mice of each gender from different litters at different postnatal stages were randomly selected for immunostaining and quantification analysis. Stained sections were imaged with Zeiss LSM800 confocal microscope as tiled or single images. Images obtained were processed and quantified with Imaris software and ImageJ. Statistical analyses of the data were carried out with R-package, GraphPad Prism, and SAS (Statistical Analysis Software). Mann-Whitney test was used to assess statistical significance between independent experimental groups. All reported significant levels represent two-tailed values. Details are given in SI Appendix.

Figure legends
Fig. 1. PNS promoted PROM1+ cell proliferation in the lateral E-SVZ of prenatally stressed male offspring.
(**A**) The pregnant dams were stressed in a plastic restrainer from E14 until delivery. Samples were collected at postnatal different time points. To label proliferating cells, BrdU was IP injected 1hr before sacrifice. (**B**) PNS significantly increased E-SVZ NSC proliferation in prenatally stressed adult offspring (P60 and P180), but not neonatal and juvenile mice (P0.5-21). CPu, caudate putamen; LV, lateral ventricle; Sept, septum. Scale bar, 30μm.

(C) MKI67 (green color), PROM1 (red color), and GFAP (white color) immunostaining. The enlarged images at the bottom showed MKI67$^+$/PROMl$^+$ ependymal cells are GFAP negative (MKI67$^+$ cells: CTL = 24.5 ± 1.6%, n = 6; STR = 31.8 ± 2.8%, n=6. P = 0.026. MKI67$^+$/PROM1$^+$ cells: CTL = 2.7 ± 0.3%, n = 13; STR= 5.5 ± 0.8%, n = 10. P= 0.0002). The box plot showed the minimum, the maximum, the median, the mean (the + symbol), and the first and the third quartiles. Scale bars, 10μm, 2μm. **(D)** BrdU (green color), PROM1 (red color), and GFAP (white color) immunostaining (BrdU$^+$ cells: CTL = 13.3 ± 0.9%, n = 28; STR =21.4 ± 1.4%, n = 26. P<0.0001. BrdU$^+$/PROM1$^+$ cells: CTL = 0.9 ± 0.2%, n = 26; STR = 2.9 ± 0.5%, n = 18. P = 0.0002). The enlarged images on the right showed BrdU/PROMl double positive ependymal cells were GFAP negative. Scale bars, 10μm, 2μm. CTL, control; STR, stressed. Mann-Whitney test. All data were presented as mean ± SEM. *p < 0.05,***p < 0.001, ****p < 0.0001.

Fig. 2. PNS promotes neuroblast proliferation in the E-SVZ and the RMS of male offspring. (A) MKI67/BrdU/DCX immunostaining shows that the ratio of neuroblasts increased dramatically in the lateral E-SVZ (BrdU$^+$/DCX$^+$ cells: CTL = 3.5 ± 0.4%, n = 21; STR = 7.0 ± 0.6%, n = 23. P<0.0001. MKI67$^+$/DCX$^+$ cells: CTL = 11.8 ± 1.3%, n = 21; STR = 17.3 ± 1.3%, n = 23. P=0.0071). Scale bars, 30μm. **(B)** The ratio of proliferating A-cell (BrdU$^+$ or MKI67$^+$) increased in the RMS (MKI67$^+$ cells: CTL = 27.3 ± 1.8%, n = 38; STR = 35.1 ± 1.4%, n = 40. P = 0.0021. BrdU$^+$ cells: CTL = 14.6 ± 1.1%, n = 38; STR = 17.3 ± 0.6%, n = 40. P = 0.021). Scale bars, 100μm. Mann-Whitney test. All data were presented as mean ± SEM. *p < 0.05, **p < 0.01, ****p < 0.0001.

Fig. 3. PNS disturbs bulbar neurogenesis. **(A)** To determine fates of lateral E-SVZ NSCs in prenatally stressed male mice, NSCs were labeled with two BrdU IP injections. After 6 weeks, the brain samples were collected. **(B)** BrdU immunostaining (white color). The enlarged images on the right show BrdU staining in the GrO (BrdU$^+$ cells/OB section: CTL = 92 ± 8, n = 8. STR = 166 ± 19, n =8 . P = 0.0011. **p < 0.01. mean ± SEM). Mann-Whitney test. Scale bar, 200μm. **(C)** BrdU labeled cells migrated to the OB and matured to become RBFOX3$^+$ neurons. The arrow heads show BrdU/RBFOX3 double positive cells in the GrO. Scale bars, 200μm. **(D)** TUNEL assay shows that PNS does not cause cell apoptosis in the OB. Scale bars, 300μm. GrO, granular cell layer of OB; Mi, mitral cell layer; EPl, external plexiform layer; Gl, glomerular layer.

Fig. 4. Bulk and single nucleus transcriptomic analyses of cells in the lateral E-SVZ of STR and CTL mice. **(A)** Schematic overview of tissue and single nucleus isolation for bulk RNA sequencing and single nucleus RNA sequencing (snRNAseq). **(B)** Module-trait correlation analysis revealed 3 modules in male offspring were corelated with PNS (Magenta, Greenyellow, Purple). **(C)** GO analysis shows top 7 GO terms enriched in major gene modules (Magenta and Purple), excluding redundant terms. The top 5 enriched genes of each term are listed.

(D) Force-directed graph is used to visualize clusters of the total single nucleus dataset (the left panel). EpC, ependymal cell; ElC, endothelial-like cell; NPCd, dorsal neural progenitor cell; NPCv, ventral neural progenitor cell; NB, neuroblast; D1, dopamine receptor D1, $Drd1^+$ spiny projection neuron; D2, $Drd2^+$ spiny projection neuron; AC, astrocyte; OPC, oligodendrocyte progenitor cell; OD, oligodendrocyte; MG, microglia. Violin plot (the right panel) shows the top cell-type-specific genes expressed in the clusters defined in the left panel. **(E)** Force-directed graph drawing of the potentially dividing cells dataset (the left panel), and the violin plot (the right panel) of the expression of cell-type marker genes. MCM2, minichromosome maintenance complex component 2; TOP2A, DNA topoisomerase II alpha; PCNA, proliferating cell nuclear antigen; eSPN, 'eccentric' spiny projection neuron. The expression of top cell-type-specific genes of each cluster is showed in the right panel. **(F)** Gene regulatory network analysis of proliferating cells (E). The node size is proportional to its PageRank (PR) centrality. Nodes highlighted in darkorange/cadetblue colors in the network indicate genes in the intersection of three gene sets: 1) top 500 genes with the biggest changes in PR absolute value between STR and CTL; 2) genes with over 0.75 quantile of PR value in each main community (community consisting of over 5 genes) detected in the network; 3) DEGs between STR and CTL. The Nodes with text labels refer to hub genes with maximal biological relevance. **(G)** Bar plots show the GO enrichment results of up- and down- regulated hub genes in (F).

Fig. 5. Inhibition of MAPK activity counteracted PNS triggered E-SVZ NSC/NPC proliferation. (A, B) Administration of MAPK3/1 inhibitor LY3214996 (LY) and MAP2K1/2 inhibitor GSK1120212 (GSK) decreases the total phosphorylated MAPK3/1 puncta in the E-SVZ (CTL = 6.2 ± 0.3, n = 18; CTL/LY = 5.1 ± 0.5, n = 16; CTL/GSK = 5.3 ± 0.2, n = 17; STR = 8.7 ± 0.5, n = 16; STR/LY = 4.2 ± 0.3, n = 16; STR/GSK = 7.2 ± 0.3, n = 15. CTL vs LY, P = 0.0056; CTL vs GSK, P = 0.0032; CTL vs STR, P = 0.0002; STR vs LY, P < 0.0001 ; STR vs GSK, P = 0.0405). Scale bars, 30μm. **(C)** Inhibition of MAPK pathway reduces the number of both $PROM1^+/MKI67^+$ ependymal cells and $TUBB3^+/MKI67^+$ neuroblasts. $MKI67^+$cells: CTL = $20.7 \pm 0.8\%$, n = 18; CTL/LY = $17.3 \pm 1.0\%$, n = 15; CTL/GSK = $14.1 \pm 1.6\%$, n = 12; STR = $29.2 \pm 1.7\%$, n = 24; STR/LY = $20.3 \pm 0.8\%$, n = 38; STR/GSK = $22.4 \pm 0.9\%$, n = 38. CTL vs LY, P = 0.0229; CTL vs GSK, P = 0.0008; STR vs LY, P < 0.0001; STR vs GSK, P = 0.0013. $PROM1^+/MKI67^+$ cells: CTL = $1.8 \pm 0.3\%$, n = 16; CTL/LY = $1.4 \pm 0.2\%$, n = 15; CTL/GSK = $1.4 \pm 0.5\%$, n = 12; STR = $2.9 \pm 0.4\%$, n = 20; STR/LY = $0.8 \pm 0.2\%$, n = 17; STR/GSK = $1.0 \pm 0.2\%$, n = 14. CTL vs LY, P = 0.5263; CTL vs GSK, P = 0.3770; STR vs LY, P < 0.0001; STR vs GSK, P < 0.0001. $TUBB3^+/MKI67^+$ cells: CTL = $10.6 \pm 0.6\%$, n = 18; CTL/LY = $7.9 \pm 0.7\%$, n = 15; CTL/GSK = $6.0 \pm 0.5\%$, n = 12; STR = $14.3 \pm 0.8\%$, n = 20; STR/LY = $11.7 \pm 0.7\%$, n = 17; STR/GSK = $9.1 \pm 1.0\%$, n = 14. CTL vs LY, P = 0.0091; CTL vs GSK, P < 0.0001; STR vs LY, P < 0.0151; STR vs GSK, P < 0.0010. Scale bars, 30μm. Mann-Whitney test. All data were presented as mean ± SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

Acknowledgements

The study was supported by National Key Research and Development Program of China (2020YFC2002804); NIH Grants 5R21MH115382, U54HD087101, and P50HD103557, State Key Program of the National Natural Science Foundation of China (82030035), the Foundation of Shanghai Municipal Education Commission (2019-01-07-00-07-E00055), Peak Disciplines (Type IV) of Institutions of Higher Learning in Shanghai, China Postdoctoral Science Foundation funded project (2022M712418), the 2021 Shanghai "Super Postdoctoral" Incentive Program, and Key R&D Program of Jiangsu (BE2020026), National Natural Science Foundation of China (31872761). We thank the Intellectual and Developmental Disabilities Research Center at the University of California, Los Angeles.

Conflict of Interest

All authors declare that they have no conflicts of interest.

References

1.   Marco EM, Macri S, Laviola G. Critical age windows for neurodevelopmental psychiatric disorders: evidence from animal models. *Neurotox Res* 2011; **19**(2)**:** 286-307.

2.   Dehorter N, Vinay L, Hammond C, Ben-Ari Y. Timing of developmental sequences in different brain structures: physiological and pathological implications. *Eur J Neurosci* 2012; **35**(12)**:** 1846-1856.

3.   Weinstock M. Prenatal stressors in rodents- Effects on behavior. *Neurobiology of Stress* 2017; **6:** 3-13.

4.   Stevens HE, Su T, Yanagawa Y, Vaccarino FM. Prenatal stress delays inhibitory neuron progenitor migration in the developing neocortex. *Psychoneuroendocrinology* 2013; **38**(4)**:** 509-521.

5.   Rothenberger SE, Moehler E, Reck C, Resch F. Prenatal stress: course and interrelation of emotional and physiological stress measures. *Psychopathology* 2011; **44**(1)**:** 60-67.

6.   Uchida T, Furukawa T, Iwata S, Yanagawa Y, Fukuda A. Selective loss of parvalbumin-positive GABAergic interneurons in the cerebral cortex of maternally stressed Gad1-heterozygous mouse offspring. *Transl Psychiatry* 2014; **4:** e371.

7.   Mychasiuk R, Gibb R, Kolb B. Prenatal stress alters dendritic morphology and synaptic connectivity in the prefrontal cortex and hippocampus of developing offspring. *Synapse* 2012; **66**(4)**:** 308-314.

8.   Sadler TR, Nguyen PT, Yang J, Givrad TK, Mayer EA, Maarek JM *et al.* Antenatal maternal stress alters functional brain responses in adult offspring during conditioned fear. *Brain Res* 2011; **1385:** 163-174.

9.   Li Y, Sjölander A, Song H, Cnattingius S, Fang F, Yang Q *et al.* Associations of parental and perinatal factors with subsequent risk of stress-related disorders: a nationwide cohort study with sibling comparison. *Molecular Psychiatry* 2022.

10.   Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci* 2009; **10**(6)**:** 434-445.

11.   Robinson R, Lahti-Pulkkinen M, Heinonen K, Reynolds RM, Räikkönen K. Fetal programming of neuropsychiatric disorders by maternal pregnancy depression: a systematic mini review. *Pediatric Research* 2019; **85**(2)**:** 134-145.

12.   Allen E. The cessation of mitosis in the central nervous system of the Albino rat. *Journal of Comparative Neurology* 1912; **22**(6)**:** 547-568.

13.   Altman J, Das GD. Autoradiographic and histological evidence of postnatal hippocampal neurogenesis in rats. *Journal of Comparative Neurology* 1965; **124**(3)**:** 319-335.

14.   Luo Y, Coskun V, Liang A, Yu J, Cheng L, Ge W *et al.* Single-cell transcriptome analyses reveal signals to activate dormant neural stem cells. *Cell* 2015; **161**(5)**:** 1175-1186.

15.   Doetsch F, Caille I, Lim DA, Garcia-Verdugo JM, Alvarez-Buylla A. Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. *Cell* 1999; **97**(6)**:** 703-716.

16.   Eriksson PS, Perfilieva E, Bjork-Eriksson T, Alborn AM, Nordborg C, Peterson DA *et al.* Neurogenesis in the adult human hippocampus. *Nat Med* 1998; **4**(11)**:** 1313-1317.

17.   Gould E, Tanapat P, McEwen BS, Flugge G, Fuchs E. Proliferation of granule cell precursors in the dentate gyrus of adult monkeys is diminished by stress. *Proc Natl Acad Sci U S A* 1998; **95**(6)**:** 3168-3171.

18.   Johansson CB, Momma S, Clarke DL, Risling M, Lendahl U, Frisén J. Identification of a neural stem cell in the adult mammalian central nervous system. *Cell* 1999; **96**(1)**:** 25-34.

19.   Delgado AC, Maldonado-Soto AR, Silva-Vargas V, Mizrak D, von Kanel T, Tan KR *et al.* Release of stem cells from quiescence reveals gliogenic domains in the adult mouse brain. *Science* 2021; **372**(6547)**:** 1205-1209.

20.   Lois C, Alvarez-Buylla A. Proliferating subventricular zone cells in the adult mammalian forebrain can differentiate into neurons and glia. *Proceedings of the National Academy of Sciences of the United States of America* 1993; **90**(5)**:** 2074-2077.

21.   Pilz GA, Bottes S, Betizeau M, Jörg DJ, Carta S, Simons BD *et al.* Live imaging of neurogenesis in the adult mouse hippocampus. *Science* 2018; **359**(6376)**:** 658-662.

22.   Altman J, Das GD. Autoradiographic and histological evidence of postnatal hippocampal neurogenesis in rats. *J Comp Neurol* 1965; **124**(3)**:** 319-335.

23.   Zhang W, Liu HT. MAPK signal pathways in the regulation of cell proliferation in mammalian cells. *Cell Research* 2002; **12**(1)**:** 9-18.

24.   Lavoie H, Gagnon J, Therrien M. ERK signalling: a master regulator of cell behaviour, life and fate. *Nat Rev Mol Cell Biol* 2020; **21**(10)**:** 607-632.

25.   Rovida E, Stecca B. Mitogen-activated protein kinases and Hedgehog-GLI signaling in cancer: A crosstalk providing therapeutic opportunities? *Semin Cancer Biol* 2015; **35:** 154-167.

26. Atanaskova N, Keshamouni VG, Krueger JS, Schwartz JA, Miller F, Reddy KB. MAP kinase/estrogen receptor cross-talk enhances estrogen-mediated signaling and tumor growth but does not confer tamoxifen resistance. *Oncogene* 2002; **21**(25)**:** 4000-4008.

27. Yun MS, Kim SE, Jeon SH, Lee JS, Choi KY. Both ERK and Wnt/beta-catenin pathways are involved in Wnt3a-induced proliferation. *J Cell Sci* 2005; **118**(Pt 2)**:** 313-322.

28. Katz M, Amit I, Yarden Y. Regulation of MAPKs by growth factors and receptor tyrosine kinases. *Biochim Biophys Acta* 2007; **1773**(8)**:** 1161-1176.

29. Hartsough MT, Frey RS, Zipfel PA, Buard A, Cook SJ, McCormick F *et al.* Altered transforming growth factor signaling in epithelial cells when ras activation is blocked. *J Biol Chem* 1996; **271**(37)**:** 22368-22375.

30. Shin S, Dimitri CA, Yoon S-O, Dowdle W, Blenis J. ERK2 but Not ERK1 Induces Epithelial-to-Mesenchymal Transformation via DEF Motif-Dependent Signaling Events. *Molecular Cell* 2010; **38**(1)**:** 114-127.

31. Gonzalez DM, Medici D. Signaling mechanisms of the epithelial-mesenchymal transition. *Sci Signal* 2014; **7**(344)**:** re8.

32. Lemaire V, Koehl M, Le Moal M, Abrous DN. Prenatal stress produces learning deficits associated with an inhibition of neurogenesis in the hippocampus. *Proc Natl Acad Sci U S A* 2000; **97**(20)**:** 11032-11037.

33. Belnoue L, Grosjean N, Ladeveze E, Abrous DN, Koehl M. Prenatal stress inhibits hippocampal neurogenesis but spares olfactory bulb neurogenesis. *PloS one* 2013; **8**(8)**:** e72972.

34. Lin YL, Wang S. Prenatal lipopolysaccharide exposure increases depression-like behaviors and reduces hippocampal neurogenesis in adult rats. *Behav Brain Res* 2014; **259:** 24-34.

35. Lemaire V, Lamarque S, Le Moal M, Piazza PV, Abrous DN. Postnatal stimulation of the pups counteracts prenatal stress-induced deficits in hippocampal neurogenesis. *Biol Psychiatry* 2006; **59**(9)**:** 786-792.

36. Weinstock M. Prenatal stressors in rodents: Effects on behavior. *Neurobiology of stress* 2017; **6:** 3-13.

37. Glover V. Annual Research Review: Prenatal stress and the origins of psychopathology: an evolutionary perspective. *J Child Psychol Psychiatry* 2011; **52**(4)**:** 356-367.

38. Otis EM, Brent R. Equivalent ages in mouse and human embryos. *Anat Rec* 1954; **120**(1)**:** 33-63.

39. Tanapat P, Hastings NB, Reeves AJ, Gould E. Estrogen stimulates a transient increase in the number of new neurons in the dentate gyrus of the adult female rat. *J Neurosci* 1999; **19**(14)**:** 5792-5801.

40. Van den Hove DLA, Kenis G, Brass A, Opstelten R, Rutten BPF, Bruschettini M *et al.* Vulnerability versus resilience to prenatal stress in male and female rats; Implications from gene expression profiles in the hippocampus and frontal cortex. *European Neuropsychopharmacology* 2013; **23**(10)**:** 1226-1246.

41.   Nugent BM, O'Donnell CM, Epperson CN, Bale TL. Placental H3K27me3 establishes female resilience to prenatal insults. *Nat Commun* 2018; **9**(1)**:** 2555.

42.   Aleman A, Kahn RS, Selten JP. Sex differences in the risk of schizophrenia: evidence from meta-analysis. *Arch Gen Psychiatry* 2003; **60**(6)**:** 565-571.

43.   Baron-Cohen S, Knickmeyer RC, Belmonte MK. Sex Differences in the Brain: Implications for Explaining Autism. *Science* 2005; **310**(5749)**:** 819-823.

44.   Kidger AM, Sipthorp J, Cook SJ. ERK1/2 inhibitors: New weapons to inhibit the RAS-regulated RAF-MEK1/2-ERK1/2 pathway. *Pharmacol Ther* 2018; **187:** 45-60.

45.   Doetsch F, Petreanu L, Caille I, Garcia-Verdugo JM, Alvarez-Buylla A. EGF converts transit-amplifying neurogenic precursors in the adult brain into multipotent stem cells. *Neuron* 2002; **36**(6)**:** 1021-1034.

46.   Huang C-Y, Tan T-H. DUSPs, to MAP kinases and beyond. *Cell & Bioscience* 2012; **2**(1)**:** 24.

47.   Coskun V, Wu H, Blanchi B, Tsao S, Kim K, Zhao J *et al.* CD133+ neural stem cells in the ependyma of mammalian postnatal forebrain. *Proc Natl Acad Sci U S A* 2008; **105**(3)**:** 1026-1031.

48.   Klein SL, Flanagan KL. Sex differences in immune responses. *Nature Reviews Immunology* 2016; **16**(10)**:** 626-638.

49.   Lee C-M, Zhou L, Liu J, Shi J, Geng Y, Liu M *et al.* Single-cell RNA-seq analysis revealed long-lasting adverse effects of tamoxifen on neurogenesis in prenatal and adult brains. *Proceedings of the National Academy of Sciences* 2020; **117**(32)**:** 19578.

50.   Wang G, Wen WX, Mead AJ, Roy A, Psaila B, Thongjuea S. Processing single-cell RNA-seq datasets using SingCellaR. *STAR Protoc* 2022; **3**(2)**:** 101266.

51.   Roy A, Wang G, Iskander D, O'Byrne S, Elliott N, O'Sullivan J *et al.* Transitions in lineage specification and gene regulatory networks in hematopoietic stem/progenitor cells over human development. *Cell Rep* 2021; **36**(11)**:** 109698.

52.   Martin A, Calvigioni D, Tzortzi O, Fuzik J, Warnberg E, Meletis K. A Spatiomolecular Map of the Striatum. *Cell Rep* 2019; **29**(13)**:** 4320-4333 e4325.

53.   Gokce O, Stanley Geoffrey M, Treutlein B, Neff Norma F, Camp JG, Malenka Robert C *et al.* Cellular Taxonomy of the Mouse Striatum as Revealed by Single-Cell RNA-Seq. *Cell Reports* 2016; **16**(4)**:** 1126-1137.

54.   Zheng SC, Stein-O'Brien G, Augustin JJ, Slosberg J, Carosso GA, Winer B *et al.* Universal prediction of cell-cycle position using transfer learning. *Genome Biol* 2022; **23**(1)**:** 41.

55.   Codega P, Silva-Vargas V, Paul A, Maldonado-Soto AR, Deleo AM, Pastrana E *et al.* Prospective identification and purification of quiescent adult neural stem cells from their in vivo niche. *Neuron* 2014; **82**(3)**:** 545-559.

56.   Inta D, Cameron HA, Gass P. New neurons in the adult striatum: from rodents to humans. *Trends Neurosci* 2015; **38**(9)**:** 517-523.

57.   Iacono G, Massoni-Badosa R, Heyn H. Single-cell transcriptomics unveils gene regulatory network plasticity. *Genome Biol* 2019; **20**(1)**:** 110.

58.   Zou JX, Revenko AS, Li LB, Gemo AT, Chen H-W. ANCCA, an estrogen-regulated AAA+ ATPase coactivator for ERalpha;, is required for coregulator occupancy and chromatin modification. *Proceedings of the National Academy of Sciences* 2007; **104**(46)**:** 18067-18072.

59.   Murakami S, Li R, Nagari A, Chae M, Camacho CV, Kraus WL. Distinct Roles for BET Family Members in Estrogen Receptor α Enhancer Function and Gene Regulation in Breast Cancer Cells. *Mol Cancer Res* 2019; **17**(12)**:** 2356-2368.

60.   Browne JA, NandyMazumdar M, Paranjapye A, Leir SH, Harris A. The Bromodomain Containing 8 (BRD8) transcriptional network in human lung epithelial cells. *Mol Cell Endocrinol* 2021; **524:** 111169.

61.   Leung HW, Leung CON, Lau EY, Chung KPS, Mok EH, Lei MML *et al.* EPHB2 Activates β-Catenin to Enhance Cancer Stem Cell Properties and Drive Sorafenib Resistance in Hepatocellular Carcinoma. *Cancer Res* 2021; **81**(12)**:** 3229-3240.

62.   Shah M, Rennoll SA, Raup-Konsavage WM, Yochum GS. A dynamic exchange of TCF3 and TCF4 transcription factors controls MYC expression in colorectal cancer cells. *Cell Cycle* 2015; **14**(3)**:** 323-332.

63.   Pereira JD, Sansom SN, Smith J, Dobenecker MW, Tarakhovsky A, Livesey FJ. Ezh2, the histone methyltransferase of PRC2, regulates the balance between self-renewal and differentiation in the cerebral cortex. *Proc Natl Acad Sci U S A* 2010; **107**(36)**:** 15957-15962.

64.   Lee KE, Seo J, Shin J, Ji EH, Roh J, Kim JY *et al.* Positive feedback loop between Sox2 and Sox6 inhibits neuronal differentiation in the developing central nervous system. *Proc Natl Acad Sci U S A* 2014; **111**(7)**:** 2794-2799.

65.   Li L, Medina-Menendez C, Garcia-Corzo L, Cordoba-Beldad CM, Quiroga AC, Calleja Barca E *et al.* SoxD genes are required for adult neural stem cell activation. *Cell Rep* 2022; **38**(5)**:** 110313.

66.   Jeong WJ, Ro EJ, Choi KY. Interaction between Wnt/beta-catenin and RAS-ERK pathways and an anti-cancer strategy via degradations of beta-catenin and RAS by targeting the Wnt/beta-catenin pathway. *NPJ Precis Oncol* 2018; **2**(1)**:** 5.

67.   Improta-Brears T, Whorton AR, Codazzi F, York JD, Meyer T, McDonnell DP. Estrogen-induced activation of mitogen-activated protein kinase requires mobilization of intracellular calcium. *Proceedings of the National Academy of Sciences* 1999; **96**(8)**:** 4686-4691.

68.   Satoh Y, Kobayashi Y, Takeuchi A, Pagès G, Pouysségur J, Kazama T. Deletion of ERK1 and ERK2 in the CNS causes cortical abnormalities and neonatal lethality: Erk1 deficiency enhances the impairment of neurogenesis in Erk2-deficient mice. *J Neurosci* 2011; **31**(3)**:** 1149-1155.

69.   Vithayathil J, Pucilowska J, Goodnough LH, Atit RP, Landreth GE. Dentate Gyrus Development Requires ERK Activity to Maintain Progenitor Population and MAPK Pathway Feedback Regulation. *J Neurosci* 2015; **35**(17)**:** 6836-6848.

70. Shi Z, Geng Y, Liu J, Zhang H, Zhou L, Lin Q *et al.* Single-cell transcriptomics reveals gene signatures and alterations associated with aging in distinct neural stem/progenitor cell subpopulations. *Protein Cell* 2018; **9**(4)**:** 351-364.

71. Fujioka A, Fujioka T, Ishida Y, Maekawa T, Nakamura S. Differential effects of prenatal stress on the morphological maturation of hippocampal neurons. *Neuroscience* 2006; **141**(2)**:** 907-915.

72. Deryckere A, Stappers E, Dries R, Peyre E, van den Berghe V, Conidi A *et al.* Multifaceted actions of Zeb2 in postnatal neurogenesis from the ventricular-subventricular zone to the olfactory bulb. *Development* 2020; **147**(10).

73. Fardi M, Alivand M, Baradaran B, Farshdousti Hagh M, Solali S. The crucial role of ZEB2: From development to epithelial-to-mesenchymal transition and cancer complexity. *J Cell Physiol* 2019.

74. Dongre A, Weinberg RA. New insights into the mechanisms of epithelial–mesenchymal transition and implications for cancer. *Nature Reviews Molecular Cell Biology* 2019; **20**(2)**:** 69-84.

75. Birkhoff JC, Brouwer RWW, Kolovos P, Korporaal AL, Bermejo-Santos A, Boltsis I *et al.* Targeted chromatin conformation analysis identifies novel distal neural enhancers of ZEB2 in pluripotent stem cell differentiation. *Hum Mol Genet* 2020; **29**(15)**:** 2535-2550.

76. Miquelajauregui A, Van de Putte T, Polyakov A, Nityanandam A, Boppana S, Seuntjens E *et al.* Smad-interacting protein-1 (Zfhx1b) acts upstream of Wnt signaling in the mouse hippocampus and controls its formation. *Proc Natl Acad Sci U S A* 2007; **104**(31)**:** 12919-12924.

77. Morshead CM, Reynolds BA, Craig CG, McBurney MW, Staines WA, Morassutti D *et al.* Neural stem cells in the adult mammalian forebrain: a relatively quiescent subpopulation of subependymal cells. *Neuron* 1994; **13**(5)**:** 1071-1082.

78. Hardy D, Malvaut S, Breton-Provencher V, Saghatelyan A. The role of calretinin-expressing granule cells in olfactory bulb functions and odor behavior. *Sci Rep* 2018; **8**(1)**:** 9385.

79. Das S, Sadanandappa MK, Dervan A, Larkin A, Lee JA, Sudhakaran IP *et al.* Plasticity of local GABAergic interneurons drives olfactory habituation. *Proc Natl Acad Sci U S A* 2011; **108**(36)**:** E646-654.

80. Heinbockel T, Laaris N, Ennis M. Metabotropic glutamate receptors in the main olfactory bulb drive granule cell-mediated inhibition. *J Neurophysiol* 2007; **97**(1)**:** 858-870.

81. Nunes D, Kuner T. Disinhibition of olfactory bulb granule cells accelerates odour discrimination in mice. *Nature Communications* 2015; **6**(1)**:** 8950.

82. Alonso M, Viollet C, Gabellec MM, Meas-Yedid V, Olivo-Marin JC, Lledo PM. Olfactory discrimination learning increases the survival of adult-born neurons in the olfactory bulb. *J Neurosci* 2006; **26**(41)**:** 10508-10513.

83. Breton-Provencher V, Lemasson M, Peralta MR, 3rd, Saghatelyan A. Interneurons produced in adulthood are required for the normal functioning of the olfactory bulb network and for the execution of selected olfactory behaviors. *J Neurosci* 2009; **29**(48)**:** 15245-15257.

84. Moreno MM, Linster C, Escanilla O, Sacquet J, Didier A, Mandairon N. Olfactory perceptual learning requires adult neurogenesis. *Proc Natl Acad Sci U S A* 2009; **106**(42)**:** 17980-17985.

85. Whitman MC, Greer CA. Adult neurogenesis and the olfactory system. *Prog Neurobiol* 2009; **89**(2)**:** 162-175.

86. Yang M, Crawley JN. Simple behavioral assessment of mouse olfaction. *Curr Protoc Neurosci* 2009; **Chapter 8:** Unit 8 24.

87. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014; **15**(12)**:** 550.

88. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* 2008; **9:** 559.

89. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation (Camb)* 2021; **2**(3)**:** 100141.

90. Hu P, Fabyanic E, Kwon DY, Tang S, Zhou Z, Wu H. Dissecting Cell-Type Composition and Activity-Dependent Transcriptional State in Mammalian Brains by Massively Parallel Single-Nucleus RNA-Seq. *Mol Cell* 2017; **68**(5)**:** 1006-1015 e1007.

A

B

A

E14   P0   9-month-old

PNS   Foster   MAPK inhibitors
                                27days

B

C