Domain loss enabled evolution of novel functions in a gene superfamily

Ivan Koludarov[1,*,^], Tobias Senoner[1,2*], Timothy N. W. Jackson[3], Daniel Dashevsky[4], Michael Heinzinger[1], Steven D. Aird[5] & Burkhard Rost[1, 6]

TUM (Technical University of Munich) Department of Informatics, Bioinformatics & Computational Biology - i12, Boltzmannstr. 3, 85748 Garching/Munich, Germany

TUM Graduate School, Center of Doctoral Studies in Informatics and its Applications (CeDoSIA), Boltzmannstr. 11, 85748 Garching/Munich, Germany

Australian Venom Research Unit, Department of Biochemistry and Pharmacology, University of Melbourne, Australia

Australian National Insect Collection, Commonwealth Scientific & Industrial Research Organisation, Canberra, Australia

7744-23 Hotaka Ariake, Azumino-shi, Nagano-ken, Japan 399-8301

Institute for Advanced Study (TUM-IAS), Lichtenbergstr. 2a, 85748 Garching/Munich, Germany & TUM School of Life Sciences Weihenstephan (WZW), Alte Akademie 8, Freising, Germany

* Contributed equally

^ Corresponding author: koludarov@rostlab.org, www.rostlab.org, +49-289-17-811 (email rost: assistant@rostlab.org)

Abstract

Three-finger toxins (3FTxs) are a functionally diverse family of toxins, apparently unique to venoms of caenophidian snakes. Although the ancestral function of 3FTxs is antagonism of nicotinic acetylcholine receptors (nAChR), redundancy conferred by accumulation of duplicate genes has facilitated extensive neofunctionalization, such that derived members of the family interact with a range of targets. 3FTxs are members of the Ly6/uPAR family, but their non-toxin ancestor remains unknown. Combining traditional phylogenetic approaches, manual synteny analysis, and cutting-edge machine learning techniques (including *AlphaFold2* and *ProtT5*), we have reconstructed a detailed evolutionary history of 3FTxs. We identify their immediate ancestor as a non-secretory Ly6, unique to squamate reptiles, and propose that changes in molecular ecology resulting from loss of a membrane-anchoring domain (MaD) and changes in gene expression, paved the way for evolution of one of the most important families of snake toxins.

Key words: gene family evolution, evolution of function, comparative genomic analysis, protein natural language models, protein structure prediction.

35   Introduction

The low cost of accumulating sequence data has permitted molecular evolutionary studies to
focus on the central role of nucleic acid sequences in "storing and transmitting" biological
information. Furthermore, the linear nature (writable as 1D-strings of letters) of such
sequences makes them readily amenable to computational analyses that increase our
understanding of ways in which traits evolve and affect their functional roles in the life histories
of the organisms that possess them. However, as functional traits, including those of
individual molecules, are relational, we cannot understand their evolution by focusing on
sequences alone[1,2]. Gene-products act as three-dimensional (3D) protein structures therefore
one-dimensional (1D) gene/protein sequence data alone are insufficient to explain molecular
mechanisms or evolutionary history. Nucleic acids "store" information, deployment of which
is selective and context-specific[3]. A multi-level approach is therefore required to understand
phenotypic consequences of sequence variation upon which selection acts.

Venom systems are excellent to study the impact of gene sequence change upon protein
function because most proteinaceous toxins are encoded by single genes and adapted for
specific functions when injected into target organisms[4]. Few other systems offer such a clear
causal pathway from genetics to ecology. This tractability helped investigations of toxin-
derived molecules as drugs and their utilization as investigational ligands[5]. Here, we chronicle
origins and innovations of an ancient gene superfamily that includes one of the most
widespread and functionally diverse toxin families in snake venoms: three-finger toxins
(3FTxs) derived from a superfamily called lymphocyte antigen 6 (Ly6) or urokinase-type
plasminogen activator receptors (uPAR)[6]. Members of this superfamily are located on several
chromosomes in vertebrates and are also present in many groups of invertebrates, indicating
their ancient origin. Although 3FTxs appeared relatively recently in this superfamily, they are
its best characterized members, and as a result, the Ly6/uPAR superfamily is also sometimes
referred to as "toxin-like proteins" (TOLIP), or the snake-toxin-like superfamily[7].

Ly6/uPAR proteins possess a characteristic "LU" or "3FF" ("three-finger fold"; Fig. 1). The
group includes membrane-bound proteins such as multimeric uPARs, which have a C-
terminus that is post-translationally removed to attach a glycophosphatidylinositol (GPI)
anchor (MaD: the "membrane-anchoring domain/region"), and secretory proteins lacking this
region (for example secreted Ly-6/uPAR-related protein *SLURP* and 3FTxs)[6-8]. Many
Ly6/uPAR proteins function as lymphocyte antigens. Others bind and modulate nicotinic
acetylcholine receptors (nAChRs, Table 1)[7]. The latter function is mediated by the cysteine-
rich 3FF region, typically involving 60-74 residues. While this region has no enzymatic activity,
3FTxs have co-opted their affinity for the nAChR as the primordial toxic function[9]. In snakes,
these genes were recruited to serve as neurotoxins, leading to an explosive diversification of
toxin sequences and activities[7,10]. Plesiotypic forms of the 3FTx family possess 10 cysteines
in 5 disulfide bonds (Fig. 1). These forms exert "α-neurotoxicity" by antagonizing nAChRs,
with some forms exhibiting greater affinity for reptile or bird, rather than mammalian
receptors[8]. This primordial theme has been extensively varied within the venom systems of
colubroid snakes (particularly members of the family *Elapidae*), generating considerable
evolutionary and ecological impact[9,10].

The relation between plesiotypic toxins and derived elapid forms remains difficult to discern
despite substantial research on 3FTxs. Molecular phylogenies of 3FTxs are typically restricted
to either plesiotypic or derived forms, either to narrow the research question or because
Bayesian phylogenies including both types almost invariably collapse into extensive,
uninformative polytomies[11,12]. Ancestral relations within the Ly6/uPAR superfamily remain

unclear, although they have long been suspected to be orthologues of LYNX1 and its
homologues on Chromosome 8 in humans[8].
84

Fig. 1: Primary species and proteins investigated in this study. Panel A: Species and their
phylogenies on which this study is centered, including estimated divergence times in millions of years
ago; Panel B: Representative structural diversity in 3FTx/Ly6 family. 1: Structure of a typical membrane
bound Ly6 protein (human LYNX1), color highlights parts of the molecule that are encoded by different
exons: exon one in orange, exon two in light blue, exon three in pink. 2: The same structure with key
elements highlighted: signal peptide in maroon, membrane-anchoring domain in blue, cysteine bonds
common to the entire 3FTx/Ly6 family in yellow, cysteine bond that disappears in derived 3FTxs in

orange. 3: Structure of a derived 3FTx (alpha-bungaratoxin), highlighting changes in structure from a
plesiotypic form: cysteine bond new to long-chain 3FTxs in orange, extended loop in green, common
cysteine bonds in yellow, notice missing cysteine and membrane-anchoring domain (cf. 2). All
structures computed as a part of this study (see Materials and Methods and Supplementary Materials
SM).

While there has been an explosion of high-quality sequence data since the 1960s[13] many
sequences remain unannotated and inferring structure or activity from sequence has been
challenging. Recent advances in artificial intelligence (AI) promise to mitigate these
bottlenecks. As of spring, 2023, AlphaFold2[14] has been used to make accurate predictions of
3D protein structure for over 200 million proteins[15]. This wealth of data allows to infer similarity
between proteins based on their structures, such as FoldSeek[16] Independently, another
breakthrough built upon advances in natural language processing (NLP) by analogizing words
with amino acids and sentences with proteins. This simple adaption enabled the transfer of
NLP concepts to biology by learning some aspects of the "language" of life as written into
protein sequences through AI models dubbed protein Language Models (pLMs)[17-19].

In this study, we combined reliable 3D structure predictions from AlphaFold2[14] and
embeddings from the pLM ProtT5[18] with traditional bioinformatic analyses and extensive
manual microsyntenic analyses. Using public sequence data (including newly annotated and
re-annotated genes), we examined the Ly6 genomic region in various vertebrate genomes to
reconstruct the evolutionary history of 3FTxs and non-toxin family members. For simplicity,
we refer to genes in this region as "Ly6", while using "TOLIP" to refer to the larger parental
group of uPARs. This approach[20-23] enabled us to reconstruct the evolutionary history of the
Ly6 gene family with unprecedented detail and to catalogue emergence of novel structures
and functions that led from an ancestral Ly6 to the radiation of deadly snake venom 3FTxs.
117

118 **Table 1: Existing knowledge of human Ly6 genes, homologs of 3FTXs.** ◊

Protein name	Sketch of known information
GPIHBP1 Glycosylphosphatidylinositol anchored high density lipoprotein binding protein 1	Capillary endothelial cell membrane protein assisting lipoprotein lipase in processing dietary lipoprotein.
Ly6H Lymphocyte Antigen 6 Family Member H	Membrane-anchored; assumed to modulate nAChRs activity; seems to inhibit alpha-7/CHRNA7 signaling in hippocampal neurons.
Ly6L Lymphocyte Antigen 6 Family Member L	Membrane-anchored.
Ly6E Lymphocyte Antigen 6 Family Member E	Anchored cell surface protein regulating T-lymphocytes proliferation, differentiation, and activation.
Ly6D Lymphocyte Antigen 6 Family Member D	Predicted as membrane-anchored; predicted to be involved in lymphocyte differentiation.
LYNX1 Ly6/Neurotoxin 1	Membrane-anchored; modulates functional properties of nAChRs to prevent excessive excitation, and hence neurodegeneration.
SLURP2 Secreted Ly6/UPAR Related Protein 2	Secreted; binds and may modulate the functional properties of nAChRs; may regulate keratinocytes proliferation, differentiation and apoptosis.
LYPD2 Ly6/PLAUR Domain-Containing Protein 2	Predicted to be located in extracellular region and plasma membrane.

Protein name	Sketch of known information
SLURP1 Secreted Ly6/UPAR Related Protein 1	Secreted; found to be a marker of late differentiation of the skin; may be involved in regulation of intracellular Ca(2+) signaling in T cells.
Ly6K Lymphocyte Antigen 6 Family Member K	Predicted in cell surface, cytoplasm, and plasma membrane; predicted to be active in acrosomal vesicle and to be involved in binding of sperm to zona pellucida.
PSCA Prostate Stem Cell Antigen	Membrane-anchored; highly expressed in prostate, also expressed in bladder, placenta, colon, kidney, and stomach; possibly involved in regulation of cell proliferation; inhibits nicotine-induced signaling, *in vitro*.

120　◊　Based on UniProt and Entrez databases. In order of chromosomal position downstream from
121　　　TOP1MT gene.

122   **Results**

Genomic analysis reveals an evolutionary stable Ly6 cluster. To distinguish distantly
related TOLIP uPAR/PLAUR genes from Ly6s that are direct homologs of snake 3FTxs, we
surveyed genomes of 15 species of *Tetrapoda* (four-limbed vertebrates, see Table 2 in
Materials and Methods) for regions syntenic with snake 3FTxs (Fig. 2). We identified the
mitochondrial topoisomerase, TOP1MT, as the best marker gene for this region. In all
genomes surveyed, TOP1MT was present in a single copy and either flanked the 3FTx/Ly6
genomic region or was in the middle of it (SM 2). Another useful marker gene is THEM6
(Thioesterase Superfamily Member 6 – UniProt ID Q8WUY1 for a human form) present in
mammals, some reptiles (including snakes), and *Xenopus* (clawed frog). This gene is located
near SLURP1 homologs, implying that its stable synteny is rather ancient.

Fig. 2: Syntenic schema derived from combined phylogenetic and genomic analyses. <u>Panel A</u>
compares Ly6/3FTx clusters from Komodo dragon and humans. Gene names are from annotations.
Unless marked with Ψ, gene relationships are based on our phylogenetic analyses. <u>Panel B</u> provides
a syntenic map for representatives of tetrapod clades. Ly6 genes are colored according to the legend,
with Ly6 unique standing for genes who do not have direct orthologs in surveyed species. TOP1MT
are white, and other genes are depicted as hollow semi-circular, white arrowheads, with narrow
triangles representing orphan exons. Genes encoding forms predicted to be secreted are indicated
with asterisks (*). Human Ly6K is marked with (~*) because only some forms are secreted (see text for
details).
143
We located all Ly6/3FTx genes (both full-length and missing exons) using previously
established methods of BLASTing exons across a genomic region[20-23]. After this, we
described multiple genes, pseudogenes, and orphan exons in each species examined. In
most cases, the Ly6/3FTx cluster was located downstream from TOP1MT. It spanned several
megabase pairs (Mbp), contained 8-35 isoforms (10, on average, for non-venomous species)
from the Ly6/3FTx family and rarely included genes from any other protein family (Fig. 2, SM
2). These syntenic arrangements were extremely conserved among all species sampled, with
the notable exception of snakes (Fig. 3), and most genes were shared by all members within
each taxonomical class, i.e., Reptilia or Mammalia.

153
Fig. 3: **Snake 3FTxs.** <u>Panel A</u>: 3FTx genomic regions in snakes, top to bottom: *Protobothrops*
muscrusquamatus, *Naja naja*, *Bungarus multicinctus* and *Pseudonaja textilis*. Non-3FTx Ly6 in white,
genes unrelated to the Ly6/3FTx family indicated by white semi-circular arrowheads, short-chain 3FTx
genes in light blue, long-chain 3FTx genes in dark blue, plesiotypic 3FTx genes in bright green and
non-standard 3FTx genes in moss green. Exonic shapes above genes show non-canonical exonic
structure with duplicated Exon 3. In all cases, only a single copy of exon 3 is used for a mature
sequence, while another is pseudogenized. TOP1MT gene indicates continuous genomic region that

is homologous to Ly6 cluster region shown in Figure 2. Other chromosomal regions are indicated by
an X with numbers where the exact chromosome is known. Ends of bioinformatic scaffolds are
indicated with two diagonal parallel lines, where genomic regions continue to a line below is indicated
with black triangles. Orphan exons are indicated as follows: large triangle for exon 3, rounded small
rectangle for exon 2 and small triangle for exon 1. Panel B: phylogenetic relationships between cobra
genes and their respective genomic position, pseudogenized genes are marked with Greek letter psi.
167
The 3FTx cluster is dramatically expanded in Indian cobra (*Naja naja*) one of very few elapid
genomes sequenced to-date at the chromosomal level[23]. This pattern conformed to the
central role of this toxin family in the evolution of venoms in elapid snakes[9]. Another unique
feature of elapid genomes was the presence of two genes in the Indian cobra located outside
the TOP1MT neighborhood (on Chromosome 3). Those were single-copy Ly6/3FTx family
genes on Chromosomes 1 and 4. We managed to locate an ortholog for this gene only on
Chromosome 4 in the Eastern brown snake (*Pseudonaja textilis*) genome, whereas both were
found in the many-banded krait (*Bungarus multicinctus*). We found no related genes (or stray
exons) in homologous genomic regions of any other organism analyzed. Many 3FTx genes
have a 4-exon structure distinct from the typical 3-exon structure (Fig. 3). All additional exons
are duplicated and pseudogenized copies of exon 3, inserted before the original exon 3. We
did not find evidence of protein production involving these redundant exons. One of the genes
in Bearded dragons (*Pogona*) likely has a similar exonic structure (Povi_7 in our dataset, see
SM 2); however, we could only detect pieces (encoding 4-5 aa each) of the duplicated exon,
not its full, even if pseudogenized, sequence. A human Ly6K gene possesses a very similar
feature, with its third exon duplicated twice, and all three isoforms apparently functional.
Lastly, human LYPD2-SLURP1 has a known "read-through" 4-exon chimera (exons 1 and 2
from LYPD2 concatenated with exons 2 and 3 of SLURP1) that is very likely a uniquely
mammalian feature, given that in all other genomes surveyed those genes have a different
orientation and are facing each other. However, genomic annotations for both *Pogona* and
Varanus predict similar read-throughs involving LYPD2 and one of the unique reptilian Ly6
genes located upstream from LYPD2. Whether this is real or an annotation artefact due to
projection of human genomic annotation onto lizard genomes (in all the cases, the read-
through occurs right before THEM6 gene) remains an open question.
192
Protein 3D-structural analysis. We used the ColabFold[24] implementation of AlphaFold2[14] to
predict full sequence (where available) and mature protein 3D structures for all 1,427 Ly6
family proteins in our dataset (SM 1). Where experimental 3D structures were available, we
validated the AlphaFold2/ColabFold predictions against them (see SM 1 for comparison and
SM 6 for structures). All proteins in the dataset were observed and predicted in the same
canonical 3FTx-domain architecture. They differed in their compactness (density of inter-
residue contacts) and loop lengths (continuous regions with neither helix nor strand). Most
Ly6 genes (~86%) possessed a lengthy C-terminal "tail", in some cases predicted to adopt
an alpha helix. In humans that extension facilitates GPI-assisted membrane anchoring[6,7].
Following broad classification of 3FTx accepted in literature[10,11,12], we classified all toxins in
our dataset into 4 categories: plesiotypic (with ancestral Ly6 cysteine arrangement and no
loop extensions); short chain (8 cysteines), long chain (10 cysteines, of which one pair is novel,
loop extension present) and non-standard (that do not fit into any of the previous categories;
also called "non-canonical" in literature).
207
Phylogenetics (ExaBayes, iqTree, DALI). We removed signal peptides from sequences (to
avoid using potentially mis-predicted N-termini) and aligned cut sequences with MAFFT[25].
The resulting alignments seeded the Bayesian (ExaBayes[26]) and Maximum Likelihood

(iqTree[27]) phylogenetic trees. We also generated a phylogenetic tree based on DALI (Distance-
matrix ALIgnment[28]) comparisons of predicted 3D structures. While smaller protein clades
were consistent across the three trees, their interrelationships differed between the three trees
from ExaBayes, iqTree and DALI (Fig. 4, SM 3). The Bayesian ExaBayes phylogeny recovered
a single monophyletic 3FTx clade (albeit one that was plagued by extensive polytomy, but
still more robust than any previously published phylogeny (SM 3c). This clade was sister to
Ly6E-like sequences, including a clade of toxicoferan Ly6 to which we refer as "pre-3FTx"
(Fig. 2). The iqTree phylogeny looked largely similar, but suffered from low bootstrap values
for most nodes, except those that defined some of the smaller clades. In contrast, the DALI
phylogeny recovered the vast majority of 3FTxs in two clades, one with primarily cat-eyed
snake (*Boiga*) 3FTxs and other colubrid toxins, and the second including plesiotypic toxins
from other snake families and more derived elapid toxins. 3FTxs were even more polyphyletic
in the DALI than the iqTree topology, because several 3FTxs were orphaned on single
branches, sisters to a clade containing the vast majority of Ly6 sequences and the two main
3FTx clades. Some findings were consistent across phylogenies. Secretory Ly6 proteins
lacking the MaD were distributed among various clades, indicating that this domain has been
lost convergently on multiple occasions, and non-standard 3FTxs whose cysteine patterns
deviate from the stereotypical major toxin groups are found throughout the gene family
phylogeny and across taxa (see SM 3 for details). Both iqTree and ExaBayes phylogenies
(and DALI, to a lesser extent) revealed several Ly6 clades that were unique to reptiles. Those
clades do not have any names, unlike mammalian homologs of human genes, and as far as
we can tell we are the first to characterize them as groups. We labeled them "reptilian Ly6
groups" with numbers 1 to 7 (Fig. 4). For the rest of the study we rely on Bayesian phylogeny,
which seems to offer the closest representation of reality.

Fig. 4: Phylogenetic trees for the non-redundant dataset used in this study. Leaves are colored
according to the figure legend. Ly6 genes are indicated by a thick black arc and black arrows. <u>Panel</u>
<u>A</u>: DALI branch lengths have no meaning due to the nature of the analysis; <u>Panel B</u>: Maximum-
likelihood nodes with bootstrap higher than 85% are marked by red dots; <u>Panel C: Bayesian inference:</u>
Node support higher than 85% is indicated by a red dot. See Supplementary Materials 3 for full
resolution trees.
242
Embedding maps. We used the pLM ProtT5 to generate multidimensional (1024D) vector
representations (embeddings) from single protein sequences. Essentially, these vectors
resemble the simplicity of 20D vectors obtained by computing amino acid composition, i.e.
the fraction of each of the 20 amino acids in a protein. However, while amino acid composition
vectors ignore interactions between pairs of residues, pLM embeddings reflect such effects,
on average. We computed embedding vectors for all proteins in two sets: the core Ly6/3FTx
dataset (1,427 sequences), and the entire InterPro[29] TOLIP family (2466 non-identical
sequences on May 1st 2023, of which 163 were already part of our dataset). To facilitate
visual inspection, we projected the 1024D embeddings onto 3D through UMAP[30] (Fig. 5,
https://rostspace.onrender.com/ for an interactive plot). The projected embedding space
distribution for Ly6s from the TOP1MT region (including 3FTxs) was similar between the two
data sets (core Ly6/3FTx vs. TOLIP in InterPro). All Ly6 with a MaD occupied a compact area
of the projected sequence space. Secretory Ly6s and 3FTxs formed a separate, more diffuse
supercluster. Plesiotypic 3FTxs formed a distinct group along with Ly6s lacking the MaD
domain, e.g., SLURPs. Toxins of colubrid snakes were found along the length of this
elongated cluster while plesiotypic toxins of elapid snakes were located together at one end
(Fig. 5 and interactive plot). Near this end we found another long cluster that included all
short-chain derived 3FTxs. Once again, relations within this cluster appeared relevant:
cytotoxic 3FTxs were gathered at the end of one extended cluster whereas another small
subcluster contained most toxins from hydrophiine snakes. There were also three outliers in
this cluster: two sequences from *Helicops leopardinus* (leopard keelback) which have
convergently lost the same disulfide bond as elapid short-chain toxins and one sequence
from *Causus rhombeatus* (night adder) that appears to be a misattributed elapid sequence,
possibly due to cross-contamination between sequencing samples. Long-chain toxins
formed their own distinct cluster, which is far more compact than those of plesiotypic or short-
chain 3FTxs.
269

Fig. 5: Rendering of 3D UMAP projection embedding space for Ly6/3FTXs. Key structure-function
clusters are circled and named. Each protein in our set was represented by a 1024-dimensional vector
of the ProtT5 pLM embeddings[18]. These 1024D vectors were projected by UMAP [30] onto 2D (Panels A
through C) and 3D (Panel D). Panel A shows distribution of subgroups within the family; Panel B shows

predictions of secreted and membrane-bound proteins; Panels C and D are different representations
of the same information on smaller clades within the family, thus, only a single figure legend is given.
The resulting embedding map seeded hypotheses on the hypothetical evolutionary trajectory of this
family in protein space that were compared to existing sequence- or structure-based approaches.
Clustering, especially with the 3D projection, obscures many points and skews the actual distance.
The full detail of interactive plots available at: https://rostspace.onrender.com/ .

Discussion

Embeddings boost understanding of origins of three-finger toxins. Our results illustrate
why reconstructing the early evolution of 3FTxs has proven difficult[8]. The two traditional
phylogenetic methods compared, provided conflicting results. The Bayesian phylogeny
exhibited unresolved polytomies similar to those that have plagued previous efforts to
incorporate plesiotypic and derived toxin sequences into the same trees (Figs. 4 and SM 3c).
Despite the polytomous substructure of the 3FTx clade in the Bayesian phylogeny, our
synteny- and pLM-based methods are concordant with hypothesizing that the whole toxin
clade had a single origin, rather than multiple origins, as implied by the structure-based DALI
phylogeny.

Previous approaches relied mostly on statistical analysis of related proteins via multiple
sequence alignments (MSAs). In contrast, the main idea behind pLMs is to let an artificial
neural network (ANN) learn how to "read" a protein sequence, i.e., how to make an amino acid
sequence computer-readable. Similar to a child learning a new language, this is achieved by
training *transformers* (the underlying ANN architecture of the most successful LMs and pLMs)
on filling out cloze tests for LMs or residues for proteins. While repeating this task on billions
of sentences or protein sequences, the (p)LM learns to detect re-occurring patterns by
encoding them in its internal, trainable weights. This is achieved via the *attention mechanism*
[31] at the core of each transformer, which learns for each input token (a word or an amino acid)
a weighted average over all the other tokens in the same sentence or protein sequence. The
crucial advantage of training the network on cloze tests is that we can directly leverage
information from the gigantic amount of unlabeled data, i.e., we need only sequential data
without any other information on the language-specific grammar. The knowledge acquired by
pLMs during this process can later be transferred to any other task by providing a
sentence/sequence as input to the model and extracting its *hidden states* (internal activations
of the ANN). This is often dubbed *transfer learning*. Akin to a new experimental technique, the
numerical vectors, *embeddings*, extracted this way provide a novel perspective on the
relationship between proteins, orthogonal to established approaches such as homology-
based inference. Contrary to most existing approaches which rely on MSAs, pLMs only need
single protein sequences as input to provide protein comparisons and predictions[32-34]. These
embeddings form a sequence space that reveals previously obscured biological information[35],
an invaluable feature for studying gene evolution and the approach at the center of the present
study. Despite their success, the complexity of large transformers makes it hard to interpret
why a specific input sequence leads to a specific embedding. An AI subfield, *XAI* (explainable
artificial intelligence), is emerging to address such questions [36,37], hopefully at some future
point [38].

Our pLM ProtT5 embedding model differentiated the major molecular and functional forms
of 3FTxs that have been identified by decades of structure-function research. Impressively,
endophysiological Ly6, plesiotypic, short-chain, and long-chain toxins all form distinct
clusters, toxins of shared activity, such as cytotoxins, even group together within these
clusters (Fig. 5). Importantly, this grouping emerged intrinsically without requiring any
expertise other than inputting the 3FTx/Ly6 dataset or the entire InterPro[29] TOLIP family. Thus,
using embeddings from pLMs clearly advances predictions of molecular activity from
sequence data by analogy with prior knowledge and enable the annotation of uncharacterized
proteins extracted from the genomic data with greater confidence than is possible using only
phylogenies.

3FTxs evolved from a single-copy gene unique to toxicoferan reptiles. Identifying a
common, endophysiological ancestor of exophysiological 3FTxs has proven challenging,
although previous studies have suggested monomeric neuromodulatory Ly6s, such as LYNXs
and SLURPs, to be the most likely candidates[8]. Our embedding results put LYNXs in the main
cluster of Ly6 sequences while SLURPs and non-standard Ly6Ks clustered with plesiotypic
3FTxs as the closest human homologs. This appears intuitive, as secretory SLURPs lack the
MaD (membrane-anchoring domain) similar to 3FTxs; however, no Ly6 group present in
mammals was close to 3FTxs in our analyses. Instead, several unique reptilian groups of
genes were indicated as possible homologues to the ancestor of 3FTxs. These clades of
genes have no previous names assigned to them, so we labeled them (and other newly
discovered groups) "reptilian Ly6 groups" 1 to 7. Clustering methods applied to the
embedding space identified reptilian Ly6 group 3, and reptilian Ly6 group 4 as the sequences
closest in the embedding space to the 3FTxs. Interestingly, most members of this cluster
possess a signal peptide that begins with MKT, a character that is rare in the Ly6, but is by
far the most common start in 3FTxs. Given other lines of evidence, this could be the result of
molecular convergence. Ly6 with a MaD cluster together, separated from those without,
including these potential homologs to the ancestor of 3FTxs (groups 3, 4, 5 and 7), which are
in turn much closer to 3FTxs. Our phylogenetic and syntenic analyses strongly suggest that
MaDs have been lost repeatedly and convergently, which would imply that the embedding
space captures structural features of the protein sequences rather than their phylogeny. The
case for convergence is bolstered by the hierarchical clustering that places the candidates
closer to other non-toxicoferan Ly6 including secreted human forms than to the 3FTxs (SM
5). Our genomic and syntenic results offer yet-another possibility: while members of reptilian
Ly6 groups 3, 4 and 5 have been reconstructed from the genomes of snakes bearing 3FTx,
reptilian Ly6 group 7 is only known from genomes of non-snake toxicoferans where it occurs
in a similar microregion to where 3FTxs are found in snake genomes (Fig. 6). Signal peptides
of these reptilian group 7 genes begin with MK, but not MKT, and all possess a MaD. Our
Bayesian phylogeny revealed this group (reptilian Ly6 group 7) as a sister group to 3FTxs.
Based on the above, we labeled it "pre-3FTxs."
Our syntenic analyses indicate that all reptiles have a genomic cluster (of varying copy
number) of Ly6-like genes between TOP1MT and the more conserved SLURP1-like uPAR
genes. Given the stability and age of these syntenic groups, it is likely that each protein
encoded by these genes possesses a conserved endophysiological function, including, in
some cases, regulation of acetylcholinergic pathways. One of these copies in the cluster
became the "pre-3FTxs" in the *Toxicofera*. After losing the membrane-anchoring region
(convergently with other secretory Ly6s), and gaining increased affinity for nAChRs, this form
gave rise to snake venom 3FTxs (Fig. 7). This led to a dramatic expansion of the cluster
eventually resulting in genomes such as Indian cobra (*Naja naja*) with more than 30 3FTx
genes, many orphan exons and pseudogenes that testify to a long history of "birth-and-death"
evolution (Fig. 3, SM 2). Likewise, the taxonomic distribution of leaves and branches of
Ly6/3FTx protein phylogenies clearly indicates constant emergence of new forms and copies
in the genomic cluster.

Fig. 6: Synteny map of Ly6/3FTXs in tetrapods and an inferred evolutionary scenario. Ly6/3FTx family genes are colored according to the legend. Genes unrelated to the Ly6/3FTx family have been left out (except for TOP1MT, indicated by a white arrow with a T in it). Double diagonal bold lines signify breaks in bioinformatic scaffolds. Black triangular arrows show continuation of the chromosomal region, compressed for the sake of space. The black chromosome icon specifies chromosomes different from those that house the TOP1MT region. Dotted outlines show pseudogenes and orphan exons.

Genomic remnants discovered in our study are consistent with past research, which has repeatedly found that 3FTx genes are subject to high levels of diversifying selection and extremely high rates of molecular evolution[9,39,40]. Such rapid rates of evolution have been attributed partially to relaxation of selective constraints on individual members of multigene arrays possessing a degree of functional redundancy[40]. Accumulation of 3FTx genes in the Indian cobra (*N. naja*) and many-banded krait (*B. multicinctus*) genomes[23,41] provides a vivid example of this process. Unique syntenic arrangements of snake 3FTx and Ly6 genes also suggest that some initial "Ly6E-like" genes were lost to recombination with newly evolved 3FTxs. If so, this would indicate possible "reverse-recruitment"[42], or "moonlighting" (adaptation of separate function)[43], in which 3FTxs, derived for envenomation functions, acquired regulatory functions of their endophysiological ancestors. Although details of such molecular ecology are beyond the scope of this paper, expression of "3FTx-like" proteins in

diverse tissues [44] and various mutations in nAChRs that confer some resistance to 3FTxs [45]
suggest this possibility.

3FTxs evolved explosively in snakes following the loss of a MaD region. Over a period of
about 25-38 million years[46,47] Ly6, a previously stable cluster of genes, erupted into a massive
array of specialized proteins with a novel, exophysiological function: neurotoxic subjugation
of prey organisms. An initial small, but significant change in a membrane-bound protein in
lizards, visible both in terms of embeddings and in structure- and sequence-based trees, was
followed by a swap of its membrane-anchoring region for a signal sequence. This swap may
have resulted from partial recombination with secreted members of the Ly6 cluster located
farther away on the genome, i.e., SLURP1/LYPD2 or through a duplication of the exon that
encoded the C-terminus of the protein followed by a mutation that inserted the stop codon
before the MaD (a scenario that seems more likely given the previously described
observations in both human Ly6K and remnants of an anciently duplicated exon 3 in 3FTxs –
see Fig. 7). The novel secretory protein, perhaps constitutively expressed in oral glands of
early snakes [48,49] may have facilitated chemical subversion of neurotransmission in prey. If, as
some have conjectured[48,49], an incipient venom system has been one key adaptation behind
the explosive radiation of snakes, 3FTxs may have served as one of the first major toxin
families. Viperids, which do not express appreciable quantities of 3FTxs in their venoms,
except for Fea's viper (*Azemiops feae*)[50], have "fully functional", but dormant, 3FTx genes (Fig.
3). The reasons for this extreme conservation remain obscure but could include regulatory
spillover from molecular mechanisms responsible for conservation of Ly6 genes, or could
indicate an endophysiological role for 3FTxs in viperids, at least some of which possess 3FTx-
resistant nAChRs[45].
A recent analysis of the genome of the many-banded krait (*Bungarus multicinctus*)
concluded that 3FTxs may have originated via neofunctionalization of "Ly6E"[41]. Our results,
garnered utilizing a broader range of methods and genomic data, support the identification of
Ly6E as a close *mammalian* relative of 3FTxs. However, we identified a group of genes that
are quite likely direct descendants of transitional forms between Ly6E and 3FTxs. The initial
expansion of Ly6E-like genes was the result of multiplication, or mutation followed by
recombination throughout the cluster, of ancestral GPIHBP1-like genes, the only members of
the gene family present in the clawed frog (*Xenopus*) genome. PSCA and other *amniote*
(branch of tetrapods containing reptiles and mammals) genes (including SLURP1/LYPD2 and
Ly6H) are also likely derived from an ancestral GPIHBP1-like pool of genes. Seven of nine
frog genes display an alpha-helix membrane-anchoring loop, indicating the ancestrality of
this feature, which is shared by almost all family members except for those in the subfamilies
of SLURP1/LYPD2 and 3FTx genes. This distinction between non-secretory, membrane
bound proteins and secretory forms lacking the MaD also represents the primary disjunction,
or "leap" in protein configuration space (Fig. 5). The loss of the MaD is thus relatively common
in evolution of the Ly6 family, having occurred convergently on multiple occasions, including
in the origin of 3FTxs from "pre-3FTxs".

Fig. 7: Proposed mechanism of parallel evolution of membrane-anchored forms into secreted forms of Ly6 genes. Small triangles stand for exon 1, rounded rectangles for exon 2 and large triangles for exon 3. Exons that make up a gene are represented by circles underneath genomic schemas and are connected via a line. Newly evolved exons are colored yellow. A pseudogenized exon is indicated by a shattered triangle.

Snake toxin genes suggest convergent evolution. The evolutionary history of 3FTxs is marked by many unique and intriguing details. Nevertheless, its contours resemble scenarios previously described for snake venom serine proteases (SVSP) and viperid venom phospholipases A_2 (PLA$_2$)[1,21,22]. A more-or-less conserved group of physiologically important genes from one protein family cluster on a chromosome with a genomic architecture shared with representatives of extant clades of tetrapods. Among mammals and toxicoferan reptiles, a particular clade of genes is exapted for new functions, usually immune functions in mammals, and toxicity in snakes. However, the change always includes toxicoferan "lizards". In each of these cases, a single gene that mutated into a form distinct from the rest of the cluster has founded a whole sub-family of genes and functions, with its members being so numerous and clinically relevant that research efforts directed at them overshadow the older and supposedly more important parental clade.

As in the pattern described for PLAs$_2$[22] and SVSPs[21], the major "neofunctionalization" events in the Ly6 family follow an apparently "arbitrary" change in molecular ecology. The major structural innovation within the family concerns the loss of the MaD, i.e., a shift from a non-secretory to a secretory role. According to the classic neo-Darwinian view, such mutations are "random", or more "arbitrary". As the mutation effects are highly non-random and context-specific[1,20,51], they affect the context or molecular ecology, in which the secretory form is exposed to a novel milieu and the opportunity to engage novel interaction partners. The interactions of a secretory protein are distinct from those of a membrane-bound protein. Exposure to novel biochemical environments facilitates novel relations that originate in chance encounters between molecules and may subsequently be stabilized by selection[1]. "Recruitment" of an endophysiological protein, such as a secretory Ly6, as a venom toxin with an exophysiological target, follows a further arbitrary change in molecular ecology, likely facilitated by stochastic gene expression. Thus, the transition from membrane-bound "pre-3FTxs" to functional 3FTxs likely involved a two-stage process of transitioning between molecular ecologies, from non-secretory to secretory and from endophysiological to exophysiological. However, the possibility of 3FTxs moonlighting in endophysiological roles is intriguing, and further complicates linear conceptions of neofunctionalization. Investigation

470 of these scenarios will be further illuminated by additional high-quality caenophidian snake
471 genomes.

Conclusion

We benefitted from recent advances in machine learning (ML) and artificial intelligence (AI),
including reliable predictions of protein 3D structure by AlphaFold2[14] and embeddings from
protein Language Models (pLMs, here ProtT5[18]), to complement traditional phylogenetic and
genomic tools for analysis of the 3FTx/Ly6 family (Fig. 1, Table 1). We unveiled the complex
evolutionary history of a fascinating and diverse gene family (Fig. 6). Paired with manual
synteny analyses (Fig. 2, Fig. 6), this approach provides an unprecedented window on protein
evolution, enabling us to demonstrate that the major division within the Ly6/uPAR family is
not phylogenetic, but structural, resulting from loss of the membrane-anchoring
region/domain (MaD, Fig. 7), which has apparently occurred multiple times in evolution. Most
significantly, we could chart the evolutionary trajectory of 3FTxs from an ancestral Ly6 via an
intermediate, toxicoferan, membrane-bound "pre-3FTx" (Fig. 6). The functionally diverse
clade of snake venom 3FTxs emerges following another independent loss of the MaD. Thus,
contrary to prior hypotheses, snake venom 3FTxs cannot be said to have descended directly
from either LYNXs, Ly6Es or SLURPs.

Data and Code Availability

489 All data used in this study (including newly discovered sequences, embeddings and
490 computed structures) are given in Supplementary Materials, all necessary information of the
491 code used is given in Materials and Methods. No new code was created for this project.

Author Contributions

494 IK, SA, TNWJ and BR conceived the study. IK, TS, DD and MH performed the experiments.
495 All authors contributed to writing the manuscript.

Acknowledgements

Thanks primarily to Tim Karl for invaluable help with hardware and software and to Inga Weise
(TUM) for support with many other aspects of this work. Thanks to Matt Summerville and
Wolfgang Wuster who gave permission to use their photographs to create snake icons, as
well as Yung-Lun Lin (for *Bungarus*) and 王朝威 (for *Protobothrops*) whose images were used
under Creative Commons Attribution 4.0 International license. Also thanks to all those who
maintain public databases in particular Steven Burley (PDB, Rutgers), Alan Bridge (Swiss-
Prot, SIB, Lausanne), Alex Bateman (UniProt, EBI Hinxton) and their crews, and to all
experimentalists who enabled this analysis by making their data publicly available. IK is
funded by an Alexander von Humboldt Foundation Fellowship. We also thank the Bavarian
Ministry of Education through funding to the TUM and by a grant from the Alexander von
Humboldt foundation through the German Ministry for Research and Education (BMBF:
Bundesministerium für Bildung und Forschung); BMBF [031L0168 and program 'Software
Campus 2.0 (TUM) 2.0 '01IS17049]; Deutsche Forschungsgemeinschaft [DFG-GZ:
RO1320/4-1].

Materials and Methods

512 **Data genomes.** All data sets were downloaded from NCBI, with exception of *Bungarus,*
513
514 which is available at China National GeneBank DataBase (see accession numbers in Table 2).

515 **Table 2: Genomes used.**

Species	ID
Alligator sinensis	GCA_000455745.1
Anolis carolinensis	GCA_000090745.2
Bungarus multicinctus	CNP0002662
Chelonia mydas	GCA_015237465.2
Gallus gallus	GCA_000002315.5
Homo sapiens	GCA_000001405.29
Naja naja	GCA_009733165.1
Ornithorhynchus anatinus	GCA_004115215.4
Phascolarctos cinereus	GCA_002099425.1
Pogona vitticeps	GCA_900067755.1
Protobothrops mucrosquamatus	GCA_001527695.3
Pseudonaja textilis	GCA_900518735.1
Python bivittatus	GCA_000186305.2
Varanus komodoensis	GCA_004798865.1
Xenopus tropicalis	GCA_000004195.4

Genomic and synteny analysis. As in previous studies, including one on 3FTxs, we used
publicly available vertebrate genomes of good quality to establish locations and synteny of
the Ly6 cluster. We used genomes for which verified RNA-seq genomic annotations were
available as reference points and created an extensive map of genes that populate Ly6/3FTx
clusters in those genomes. These include TOP1MT, THEM6, cytP450, KCNMB2 and others
(SM 2). Thereby we could establish syntenic relationships of those regions in various
genomes. We then used those flanking genes as a database to BLAST[52] (NCBI-BLAST
v.2.7.1+ suite, blastn, e-value cutoff of 0.05, default restrictions on word count and gaps) the
genomes that were less well annotated. That gave us several hundred genomic scaffolds
potentially containing Ly6 genes. We used those scaffolds for a second round of BLAST
(tblastx, e-value cutoff of 0.01) against a database of sequences from well-annotated
mammalian and reptilian TOLIP genes. Positive hits were checked visually in Geneious[53], and
complete exons were manually annotated and later merged into coding sequences of newly
annotated genes if the exon order and count accorded with existing reliable Ly6 annotations.

We included all resulting genes that produced viable mature peptides, and then used these
for the phylogenetic analysis.
534
Dataset assembly. We extracted all complete gene coding regions from the genomic regions
surveyed, combined them with previously compiled 3FTx datasets[11,12,41,54] and then
supplemented them with all non-redundant reviewed 3FTx sequences from UniProt[55].
Duplicated, fragmented and incomplete sequences were removed. For broad comparison,
we used all sequences in the TOLIP protein family in InterPro[56]. We removed signal peptides
from all sequences, following SwissProt[55] annotations from humans and snakes as a guide
and SignalP6 for confirmation[57].
542
Phylogenetic analysis. We translated all viable genes located in the previous step into
proteins and aligned these with selected publicly available sequences of interest using the L-
INS-i method of MAFFT software v7.305[25] with 1000 iterations (--localpair --maxiterate 1000).
These parameters were used for all subsequent alignments.
We established a naming convention to differentiate between genomic sequences (first
two letters of both generic and specific epithets, followed by a number to differentiate
sequences from the same scaffold). We used sequences from previously published studies
and homologues from UniProt and SwissProt to provide outgroups and fill gaps in sequence
space keeping UniProt or SwissProt IDs. We visually inspected the alignments for apparent
errors (e.g., proper alignment of the cysteine backbone), and used to construct phylogenetic
trees in ExaBayes[26] (10 parallel runs of 4 chains each, and 33,000,000 generations – which
took almost 10 weeks to converge; SM 3c) and iqTree[27] (100 rounds of bootstrap; SM 3b).
555
Embedding space analysis. Over the last few years, algorithms in natural language
processing (NLP) have advanced substantially, yielding breakthroughs in tasks such as
automated translation or question answering. This was achieved in particular by training
Language Models (LMs) on large, but unlabeled text corpora[58,59]. Thanks to the sequential
nature of both natural language and protein sequences, these advances have been
transferred readily to protein sequences through so-called protein Language Models (pLMs[17]).
In both cases, masked or missing tokens (words in NLP, amino acids in computational
biology) are predicted from the remaining, uncorrupted sequence context (a sentence or a
protein sequence, respectively). As this objective does not require labeled data, it allows
mining the wealth unlabeled data[17-19,60], i.e., data without experimental annotations such as
exponentially growing protein sequence databases by relying solely on sequential patterns in
the input. Processing the information learned by such pLMs, e.g., by feeding protein
sequences as inputs to the network and constructing vectors from activation in the network's
last layers, yields a representation of protein sequences known as embeddings (for sketch:
Fig. 1 in Elnaggar et al.[18]). This allows transfer of features learned by the pLM to any predictive
task requiring numerical protein representations (transfer learning), which has already been
shown for various applications ranging from protein structure[61] to protein function[62]. Distance
in embedding space correlates with protein function and can be used as an orthogonal signal
for clustering proteins into functional families[62].
In this work, we generated embeddings for each protein sequence using the pLM ProtT5-
XL-UniRef50[18] (for simplicity referred to as *ProtT5*), which has been built in analogy to the
NLP model T5[59]. ProtT5 was trained solely on unlabeled protein sequences from BFD (Big
Fantastic Database; 2.5 billion sequences including meta-genomic sequences)[63] and
UniRef50. Ultimately this allowed ProtT5 to learn some of the constraints of protein
sequences. As ProtT5 was only trained on unlabeled protein sequences and no supervised
training or fine-tuning was performed, there is no risk of information leakage or overfitting to

a certain class or label. In order to transfer the knowledge or constraints that ProtT5 had
acquired to other tasks (transfer learning), we first created individual vector representations
for each residue in a protein. In order to derive fixed-length vector representations for single
proteins irrespective of protein length (per-protein embedding), we then averaged all residue
embeddings in a protein (Fig. 1 in Elnaggar et al.[18]). As a result, every protein was represented
as a 1024-dimensional embedding. Those high-dimensional representations were projected
to 3D using UMAP[30] (n_neighbors=25, min_dist=0.5, random_state=42, n_components=3)
and colored according to features of interest to allow visual analysis. Embeddings were
created using the bio_embeddings package[64] and 2D, 3D and interactive plots using
RostSpace (https://github.com/Rostlab/RostSpace).

Protein 3D structures. For fast protein structure prediction, 3D structures for all proteins in
our set were generated using default parameters of the ColabFold implementation (v1.3.0,
commit: 7ebcbe62e8d88400b0e75aa0878dce2ff3a6c71f) of AlphaFold 2 (AF2), i.e., no early-
stopping, no templates, and no amber-relaxation were used[24,65]. Input MSAs were generated
using the search-script provided by ColabFold with highest sensitivity. As AlphaFold2 is an
ensemble of five models each outputting its own 3D structure, only the best 3D structure for
each protein was used for further processing. For this filtering we used the structure of the
model with the highest predicted reliability, i.e., the highest predicted local distance difference
test - pLDDT.

Comparison between existing experimental 3D structures of Ly6s and 3FTxs and
predicted in this study. The dataset encompassed 59 available experimental 3D structures,
which were juxtaposed with the structures predicted by Alphafold2, employing Foldseek for
the comparison[16]. The resultant high LDDT (0.85 ± 0.08), TM-score (0.87 ± 0.08), and RMSD
(1.92 ± 0.85) provide robust evidence supporting the precision of the predicted 3D structures
across the entire dataset, please see results in SM 1.

Cluster analysis. Results of the embedding space analysis were used as coordinates for
clustering analyses carried out in R v3.6.2[66] using RStudio 2023.03.0[67]. The most basic
approach was a k-means clustering approach with k=7 implemented in the default stats
package[66]. We also used the DBSCAN algorithm with ε=0.55 and minimum points = 5 to
perform density-based clustering. Finally, we extracted a hierarchical density-based
dendrogram from the results of the OPTICS algorithm with Ξ=0.01. Density-based analyses
were implemented in the dbscan package[68].

3D structure-based trees. 3D structures predicted by AlphaFold2 were used as input for a
local version of DALI v5 (distance matrix alignment)[28] to generate trees based on structural
similarity between all proteins in our set. All parameters were left at default, but computation
was executed in parallel on 20 cores using the parameter, mpirun.
618

Protein property predictions. Beside protein structures, we used various other predicted
protein properties to enrich our analysis. More specifically, we used a recently published
pipeline that combines a variety of protein property predictors, all of which rely on ProtT5 to
encode protein sequences[69]. Here, we mainly focused on predicting whether a protein is
secreted using a method called LightAttention and on predicting membrane-anchoring
domains using the method, TMbed[70].

Supplementary Materials (SM)

1. Dataset in a spreadsheet format, including aligned sequences, grouping and localization
predictions
2. TOP1MT region gene map figure
3. Phylogenetic trees
4. CLANS analysis
5. DBSCAN clusters
6. Structures in .pdb format
7. TOLIP protein space clustering
8. Protein language models' embeddings

References

636
637
638
1   Jackson, T. N. W. & Koludarov, I. How the Toxin got its Toxicity. *Frontiers in*
Pharmacology **11** (2020). https://doi.org:10.3389/fphar.2020.574925
2   Guttinger, S. *A Process Ontology for Macromolecular Biology*. (2018).
3   Gardner, K. E., Allis, C. D. & Strahl, B. D. Operating on chromatin, a colorful
language where context matters. *J. Mol. Biol.* **409**, 36-46 (2011).
https://doi.org:10.1016/j.jmb.2011.01.040
4   Fry, B. G. *et al.* The toxicogenomic multiverse: convergent recruitment of proteins
into animal venoms. *Annu. Rev. Genomics Hum. Genet.* **10**, 483-511 (2009).
https://doi.org:10.1146/annurev.genom.9.081307.164356
5   Fry, B. G. *et al.* Seeing the Woods for the Trees: Understanding Venom Evolution as
a Guide for Biodiscovery in *Venomous Reptiles and Their Toxins: Evolution,*
Pathophysiology and Biodiscovery   1-36 (2015).
6   Loughner, C. L. *et al.* Organization, evolution and functions of the human and mouse
Ly6/uPAR family genes. *Hum. Genomics* **10**, 10 (2016).
https://doi.org:10.1186/s40246-016-0074-2
7   Kessler, P., Marchot, P., Silva, M. & Servent, D. The three-finger toxin fold: a
multifunctional structural scaffold able to modulate cholinergic functions. *J.*
Neurochem. **142 Suppl 2**, 7-18 (2017). https://doi.org:10.1111/jnc.13975
8   Utkin, Y. N., Sunagar, K., Jackson, T. N. W., Reeks, T. & Fry, B. G. Three-finger
toxins (3FTxs) in *Venomous Reptiles and Their Toxins: Evolution, Pathophysiology*
and Biodiscovery   (2015).
9   Sunagar, K. *et al.* Three-fingered RAVERs: Rapid Accumulation of Variations in
Exposed Residues of snake venom toxins. *Toxins* **5**, 2172-2208 (2013).
https://doi.org:10.3390/toxins5112172
10   Utkin, Y. N. Last decade update for three-finger toxins: Newly emerging structures
and biological activities. *World J. Biol. Chem.* **10**, 17-27 (2019).
https://doi.org:10.4331/wjbc.v10.i1.17
11   Xie, B. *et al.* Dynamic genetic differentiation drives the widespread structural and
functional convergent evolution of snake venom proteinaceous toxins. *BMC Biol.* **20**,
4 (2022). https://doi.org:10.1186/s12915-021-01208-9
12   Dashevsky, D. & Fry, B. G. Ancient Diversification of Three-Finger Toxins in Micrurus
Coral Snakes. *J. Mol. Evol.* **86**, 58-67 (2018). https://doi.org:10.1007/s00239-017-
9825-5
13   Mathé, E., Hays, J. L., Stover, D. G. & Chen, J. L. The Omics Revolution Continues:
The Maturation of High-Throughput Biological Data Sources. *Yearb. Med. Inform.* **27**,
211-222 (2018). https://doi.org:10.1055/s-0038-1667085
14   Jumper, J. *et al.* High accuracy protein structure prediction using deep learning in
Fourteenth Critical Assessment of Protein Structure Prediction (CASP14)   (2021).
15   Tunyasuvunakool, K. *et al.* Highly accurate protein structure prediction for the human
proteome. *Nature* (2021). https://doi.org:10.1038/s41586-021-03828-1
16   van Kempen, M. *et al.* Foldseek: fast and accurate protein structure search. *bioRxiv*
(2022). https://doi.org: https://doi.org/10.1101/2022.02.07.479398
17   Heinzinger, M. *et al.* Modeling aspects of the language of life through transfer-
learning protein sequences. *BMC Bioinformatics* **20**, 723 (2019).
https://doi.org:10.1186/s12859-019-3220-8

18   Elnaggar, A. *et al.* ProtTrans: Towards Cracking the Language of Lifes Code Through
Self-Supervised Deep Learning and High Performance Computing. *IEEE Trans*
Pattern Anal Mach Intell **PP** (2021). https://doi.org:10.1109/TPAMI.2021.3095381
19   Alley, E. C., Khimulya, G., Biswas, S., AlQuraishi, M. & Church, G. M. Unified rational
protein engineering with sequence-based deep representation learning. *Nature*
methods, 1-8 (2019).
20   Koludarov, I. *et al.* Bee core venom genes predominantly originated before aculeate
stingers evolved. *bioRxiv* (2022). https://doi.org:10.1101/2022.01.21.477203
21   Barua, A., Koludarov, I. & Mikheyev, A. S. Co-option of the same ancestral gene
family gave rise to mammalian and reptilian toxins. *BMC Biol.* **19**, 268 (2021).
https://doi.org:10.1186/s12915-021-01191-1
22   Koludarov, I., Jackson, T.N., Suranse, V., Pozzi, A., Sunagar, K. & Mikheyev A.S.
Reconstructing the evolutionary history of a functionally diverse gene family reveals
complexity at the genetic origins of novelty. *bioRxiv* (2019).
https://doi.org/10.1101/583344
23   Suryamohan, K. *et al.* The Indian cobra reference genome and transcriptome
enables comprehensive identification of venom toxins. *Nat. Genet.* **52**, 106-117
(2020). https://doi.org:10.1038/s41588-019-0559-8
24   Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat. Methods*
19, 679-682 (2022). https://doi.org:10.1038/s41592-022-01488-1
25   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772-780 (2013).
https://doi.org:10.1093/molbev/mst010
26   Aberer, A. J., Kobert, K. & Stamatakis, A. ExaBayes: massively parallel bayesian tree
inference for the whole-genome era. *Mol. Biol. Evol.* **31**, 2553-2556 (2014).
https://doi.org:10.1093/molbev/msu236
27   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530-1534 (2020).
https://doi.org:10.1093/molbev/msaa015
28   Holm, L. Using Dali for Protein Structure Comparison. *Methods Mol. Biol.* **2112**, 29-
42 (2020). https://doi.org:10.1007/978-1-0716-0270-6_3
29   Paysan-Lafosse, T. *et al.* InterPro in 2022. *Nucleic Acids Res* (2022).
https://doi.org:10.1093/nar/gkac993
30   Leland McInnes, J. H., James Melville. UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction. *arXiv [cs.CL]* (2018).
https://doi.org:10.48550/ARXIV.1802.03426
31   Vaswani, A. *et al.* Attention is all you need in *Advances in neural information*
processing systems.   5998-6008 (2017).
32   Heinzinger, M. *et al.* Contrastive learning on protein embeddings enlightens midnight
zone. *NAR Genomics and Bioinformatics* **4**, lqac043 (2022).
https://doi.org:https://doi.org/10.1093/nargab/lqac043
33   Littmann, M., Heinzinger, M., Dallago, C., Weissenow, K. & Rost, B. Protein
embeddings and deep learning predict binding residues for various ligand types.
Scientific Reports **11**, 23916 (2021). https://doi.org:10.1038/s41598-021-03431-4
34   Schütze, K., Heinzinger, M., Steinegger, M. & Rost, B. Nearest neighbor search on
embeddings rapidly identifies distant protein relations. *bioRxiv* (2022).
https://doi.org:doi: https://doi.org/10.1101/2022.09.04.506527
35   Detlefsen, N. S., Hauberg, S. & Boomsma, W. Learning meaningful representations
of protein sequences. *Nat Commun* **13**, 1914 (2022). https://doi.org:10.1038/s41467-
022-29443-w

36   Vig, J. *et al.* BERTology meets biology: interpreting attention in protein language
models. *arXiv preprint arXiv:2006.15222* (2020).
37   Arrieta, A. B. *et al.* Explainable Artificial Intelligence (XAI): Concepts, taxonomies,
opportunities and challenges toward responsible AI. *Information fusion* **58**, 82-115
(2020).
38   Neely, M., Schouten, S. F., Bleeker, M. & Lucic, A. A Song of (Dis) agreement:
Evaluating the Evaluation of Explainable Artificial Intelligence in Natural Language
Processing. *arXiv preprint arXiv:2205.04559* (2022).
39   Doley, R., Mackessy, S. P. & Kini, R. M. Role of accelerated segment switch in exons
to alter targeting (ASSET) in the molecular evolution of snake venom proteins. *BMC*
Evol. Biol. **9**, 146 (2009). https://doi.org:10.1186/1471-2148-9-146
40   Jackson, T. N. W. *et al.* Rapid Radiations and the Race to Redundancy: An
Investigation of the Evolution of Australian Elapid Snake Venoms. *Toxins* **8** (2016).
https://doi.org:10.3390/toxins8110309
41   Zhang, Z. Y. *et al.* The structural and functional divergence of a neglected three-
finger toxin subfamily in lethal elapids. *Cell Rep* **40**, 111079 (2022).
https://doi.org:10.1016/j.celrep.2022.111079
42   Casewell, N. R., Huttley, G. A. & Wüster, W. Dynamic evolution of venom proteins in
squamate reptiles. *Nat. Commun.* **3**, 1066 (2012).
https://doi.org:10.1038/ncomms2065
43   Jeffery, C. J. Moonlighting proteins: old proteins learning new tricks. *Trends Genet*
19, 415-417 (2003). https://doi.org:10.1016/S0168-9525(03)00167-7
44   Hargreaves, A. D., Swain, M. T., Logan, D. W. & Mulley, J. F. Testing the Toxicofera:
comparative transcriptomics casts doubt on the single, early evolution of the reptile
venom system. *Toxicon* **92**, 140-156 (2014).
https://doi.org:10.1016/j.toxicon.2014.10.004
45   Khan, M. A. *et al.* Widespread Evolution of Molecular Resistance to Snake Venom α-
Neurotoxins in Vertebrates. *Toxins* **12** (2020).
https://doi.org:10.3390/toxins12100638
46   Lee, M. S. Y., Sanders, K. L., King, B. & Palci, A. Diversification rates and phenotypic
evolution in venomous snakes (Elapidae). *R Soc Open Sci* **3**, 150277 (2016).
https://doi.org:10.1098/rsos.150277
47   Zaher, H. *et al.* Large-scale molecular phylogeny, morphology, divergence-time
estimation, and the fossil record of advanced caenophidian snakes (Squamata:
Serpentes). *PLoS One* **14**, e0216148 (2019).
https://doi.org:10.1371/journal.pone.0216148
48   Jackson, T. N. W. *et al.* Endless forms most beautiful: the evolution of ophidian oral
glands, including the venom system, and the use of appropriate terminology for
homologous structures. *Zoomorphology* **136**, 107-130 (2017).
https://doi.org:10.1007/s00435-016-0332-9
49   Fry, B. G. *et al.* Squeezers and leaf-cutters: differential diversification and
degeneration of the venom system in toxicoferan reptiles. *Mol. Cell. Proteomics* **12**,
1881-1899 (2013). https://doi.org:10.1074/mcp.M112.023143
50   Babenko, V. V. *et al.* Novel Bradykinin-Potentiating Peptides and Three-Finger
Toxins from Viper Venom: Combined NGS Venom Gland Transcriptomics and
Quantitative Venom Proteomics of the Azemiops feae Viper. *Biomedicines* **8** (2020).
https://doi.org:10.3390/biomedicines8080249

51   Jackson, T. N. W., Jouanne, H. & Vidal, N. Snake Venom in Context: Neglected
Clades and Concepts. *Frontiers in Ecology and Evolution* **7** (2019).
https://doi.org:10.3389/fevo.2019.00332
52   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local
alignment search tool. *J Mol Biol* **215**, 403-410 (1990).
https://doi.org:10.1016/S0022-2836(05)80360-2
53   Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software
platform for the organization and analysis of sequence data. *Bioinformatics* **28**,
1647-1649 (2012). https://doi.org:10.1093/bioinformatics/bts199
54   Chandna, R. *et al.* Drysdalin, an antagonist of nicotinic acetylcholine receptors
highlights the importance of functional rather than structural conservation of amino
acid residues. *FASEB bioAdvances* **1**, 115 (2019).
55   UniProt Consortium. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids*
Res **47**, D506-D515 (2019). https://doi.org:10.1093/nar/gky1049
56   Paysan-Lafosse, T. *et al.* InterPro in 2022. *Nucleic Acids Res* **51**, D418-D427 (2023).
https://doi.org:10.1093/nar/gkac993
57   Teufel, F. *et al.* SignalP 6.0 predicts all five types of signal peptides using protein
language models. *Nature biotechnology* **40**, 1023-1025 (2022).
58   Devlin, J., Chang, M.-W., Lee, K. & Toutanova, K. BERT: Pre-training of Deep
Bidirectional Transformers for Language Understanding. *arXiv [cs.CL]* (2018).
59   Raffel, C. *et al.* Exploring the Limits of Transfer Learning with a Unified Text-to-Text
Transformer. *arXiv [cs.LG]* (2019).
60   Bepler, T. & Berger, B. Learning the protein language: Evolution, structure, and
function. *Cell Syst* **12**, 654-669.e653 (2021).
https://doi.org:10.1016/j.cels.2021.05.017
61   Weissenow, K., Heinzinger, M. & Rost, B. Protein language-model embeddings for
fast, accurate, and alignment-free protein structure prediction. *Structure* **30**, 1169-
1177.e1164 (2022). https://doi.org:10.1016/j.str.2022.05.001
62   Littmann, M., Heinzinger, M., Dallago, C., Olenyi, T. & Rost, B. Embeddings from
deep learning transfer GO annotations beyond homology. *Sci. Rep.* **11**, 1160 (2021).
https://doi.org:10.1038/s41598-020-80786-0
63   Steinegger, M. & Söding, J. Clustering huge protein sequence sets in linear time.
Nature Communications **9** (2018). https://doi.org:10.1038/s41467-018-04964-5
64   Dallago, C. *et al.* Learned Embeddings from Deep Learning to Visualize and Predict
Protein Sets. *Curr Protoc* **1**, e113 (2021). https://doi.org:10.1002/cpz1.113
65   Skolnick, J., Gao, M., Zhou, H. & Singh, S. AlphaFold 2: Why It Works and Its
Implications for Understanding the Relationships of Protein Sequence, Structure,
and Function. *J. Chem. Inf. Model.* **61**, 4827-4831 (2021).
https://doi.org:10.1021/acs.jcim.1c01114
66   Team, R. D. C. A language and environment for statistical computing. *http://www. R-*
project. org (2009).
67   Team, R. RStudio: integrated development for R. *(No Title)* (2015).
68   Hahsler, M., Piekenbrock, M. & Doran, D. dbscan: Fast density-based clustering with
R. *Journal of Statistical Software* **91**, 1-30 (2019).
69   Ferruz, N. *et al.* From sequence to function through structure: deep learning for
protein design. *bioRxiv*, 2022.2008.2031.505981 (2022).
https://doi.org:10.1101/2022.08.31.505981
70   Bernhofer, M. & Rost, B. TMbed: transmembrane proteins predicted through
language model embeddings. *BMC bioinformatics* **23**, 1-19 (2022).