Title: Functional dissociations between prefrontal and parietal cortex during task switching: A combined fMRI and TMS study.

Abbreviated title: Frontoparietal task-switching dissociations after TMS

Authors: José A. Periáñez[1, *, †], Raquel Viejo-Sobera[2, †], Genny Lubrini[1, *], Juan Álvarez-Linera[3], Elisa Rodríguez Toscano[1, 4], María Dolores Moreno[4], Celso Arango[4], Diego Redolar-Ripoll[2, 7], Elena Muñoz Marrón[2, ††], Marcos Ríos-Lago[5, 6, ††]

[1] Dept. Experimental Psychology, Complutense University of Madrid, 28223, Spain.

[2] Cognitive NeuroLab, Universitat Oberta de Catalunya (UOC), Barcelona, 08018, Spain.

[3] Neuroradiology Department, Hospital Ruber Internacional. Madrid, 28034, Spain.

[4] Institute of Psychiatry and Mental Health. Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón Instituto de Investigación Sanitaria Gregorio Marañón (IiSGM), CIBERSAM, Madrid, 28007, Spain.

[5] Dept. Basic Psychology II, UNED, Madrid, 28040, Spain.

[6] Brain Damage Service, Beata Maria Ana Hospital, Madrid, 28007, Spain.

[7] Brain360 Neuroscience Institute, Barcelona, 08022, Spain.

* Corresponding authors:

Genny Lubrini, Ph.D.

Dept. Experimental Psychology, Complutense University of Madrid

Campus de Somosaguas, 28223, Madrid, Spain.

E-mail: glubrini@ucm.es

Phone: +34-91 394 3181 Fax: +34-91 394 3189

Jose A. Periáñez, Ph.D.

Dept. Experimental Psychology, Complutense University of Madrid

Campus de Somosaguas, 28223, Madrid, Spain.

E-mail: japerian@ucm.es

Phone: +34-91 394 3118 Fax: +34-91 394 3189

Submitting author:

Genny Lubrini, Ph.D.

Dept. Experimental Psychology, Complutense University of Madrid

Campus de Somosaguas, 28223, Madrid, Spain.

E-mail: glubrini@ucm.es

Phone: +34-91 394 3181 Fax: +34-91 394 3189

† José A. Periáñez and Raquel Viejo-Sobera contributed equally to the manuscript.

†† Elena Muñoz Marrón and Marcos Ríos-Lago contributed equally to the manuscript.

Conflict of interest. CA has been a consultant to or has received honoraria or grants from Acadia, Angelini, Boehringer, Gedeon Richter, Janssen Cilag, Lundbeck, Minerva, Otsuka, Pfizer, Roche, Sage, Servier, Shire, Schering Plough, Sumitomo Dainippon Pharma, Sunovion and Takeda. The remaining authors declare that they have no conflicts of interest.

Acknowledgements. This work was supported by the Spanish Ministry of Science and Innovation, Instituto de Salud Carlos III (ISCIII), CIBER-Consorcio Centro de Investigación Biomédica en Red- (CB/07/09/0023), financed by the European Union, ERDF Funds from the European Commission, "A way of making Europe", (PI1001061), and Fundación Alicia Koplowitz.

Abstract

Preparatory control in task-switching has been suggested to rely upon a set of distributed regions within a frontoparietal network, with frontal and parietal cortical areas cooperating to implement switch-specific preparation processes. Although recent causal evidences using transcranial magnetic stimulation (TMS) have generally supported this model, alternative evidences from both functional neuroimaging and neurophysiological studies have questioned the switch-specific role of both frontal and parietal cortices. The aim of the present study was to clarify the role of prefrontal and parietal areas supporting preparatory cognitive control in task-switching. Within this purpose, an fMRI study during task-switching performance was conducted to identify the specific brain areas involved in preparatory control during performance of a task-switching paradigm. Then, TMS was applied over the specific coordinates previously identified through fMRI, that is, the anterior portion of the inferior frontal junction (aIFJ) and the intraparietal sulcus (IPS). Results revealed that TMS over the aIFJ disrupted performance in both switch and repeat trails in terms of delayed responses as compared to Sham condition. In contrast, TMS over the IPS selectively interfered performance in switch trials. These findings support a multicomponent model of executive control with the aIFJ being involved in more general switch-unspecific process such as the episodic retrieval of goals, and the IPS being related to the implementation of switch-specific preparation mechanisms for activating stimulus-response mappings. The results help conciliating preceding evidences about the role of a frontoparietal network during task-switching, and support current models about a hierarchical organization within prefrontal cortex.

Key words: Executive control, Ventrolateral Prefrontal Cortex, fMRI, Task-switching, Transcranial Magnetic Stimulation.

Significance Statement

A combined fMRI and TMS study was conducted to clarify the brain areas involved in the executive control of attention during a cueing task-switching paradigm.

Functional dissociations were observed during TMS stimulation, with prefrontal and parietal areas playing different roles during task-switching preparation. While the anterior portion of the inferior prefrontal junction seemed to be involved in a general mechanism of memory retrieval for goal identification, the intraparietal sulcus seemed to be engaged in a switch-specific mechanism for the translation of abstract task goals into action rules.

The results help conciliating preceding evidences about the role of a frontoparietal network during task-switching, and support current models about a hierarchical organization within prefrontal cortex.

Introduction

Neuroanatomical models about executive control suggest interactions within a frontoparietal network (FPN) of brain regions cooperating for an adjusted and flexible control of human behaviour (Miller and Cohen, 2001; Koechlin and Summerfield, 2007; Petersen and Posner, 2012). General consensus exists among theories assuming that the prefrontal cortex will generate top-down signals to bias processing in posterior areas. Studies using task-switching paradigms in combination with neuroimaging, neurophysiological, and lesional techniques have contributed to clarify the role of different brain regions within this FPN to cognitive control (Sohn et al., 2000; Monchi et al., 2001; Periáñez et al., 2004; Kopp et al., 2006; Niendam et al., 2012; Muhle-Karbe et al., 2014; Worringer et al., 2019). Thus, task-switching requires to change frequently among a set of simple tasks, which is supposed to activate certain control operations that will ultimately allow a flexible adaptive behaviour (Wylie and Allport, 2000; Monsell, 2003; Logan and Bundesen, 2003).

Functional neuroimaging studies have identified switch-specific activation in prefrontal areas such as medial frontal gyrus, inferior frontal gyrus, and the inferior frontal junction (IFJ), and in parietal areas such as posterior parietal cortex and precuneus. For instance, the IFJ has been suggested to exert top-down preparatory control during switch trials for the updating of abstract task representations (Derrfuss, et al., 2005; Kim et al., 2012; Worringer et al., 2019). Within the parietal lobe, the intraparietal sulcus (IPS) would provide a more action-related task representations such as stimulus-response mappings (Andersen et al., 1997; Culham and Kanwisher, 2001; Worringer et al., 2019). The analysis of the time dynamics of the activation within this FPN by means of neurophysiological techniques have generally supported the idea that activation of prefrontal regions precedes the activation of posterior ones during the implementation of a switch in task (Stuss and Picton, 1978; Periáñez et al., 2004; Brass et al., 2005; Kopp et al., 2006).

Transcranial Magnetic Stimulation (TMS) has offered causal evidences about the contribution of different prefrontal (Rushworth et al., 2002; Muhle-Karbe et al., 2014) and parietal areas (Muhle-Karbe et al., 2014) to preparatory control in task-switching by lowering brain activity. Thus, Muhle-Karbe et al. (2014) found that the application of online inhibitory TMS over the IFJ increased reaction times in switch trials in an

experimental condition that involved the updating of the abstract task rules. In addition, TMS inhibition of the IPS increased the percentage of errors when switching involved remapping stimulus-response associations. The authors interpreted that prefrontal and parietal cortices would implement switch-specific preparation processes operating at two different levels of abstraction (i.e., task-goals and response-sets). However, alternative data have questioned the switch-specific preparatory role of prefrontal and parietal brain regions during task-switching (Bode and Haynes, 2009; Mansfield et al., 2012; Ruge et al., 2013). For instance, evidence about prefrontal activation during both switch and repeat trials suggests that prefrontal areas could implement more general switch-unspecific mechanisms (Braver et al., 2003; Barceló et al., 2008; Jamadar et al., 2010). Unfortunately, no other TMS study has explored further these alternative ideas, thus leading open the question about the role of the FPN in task-switching.

The aim of this work was to clarify the role of prefrontal and parietal areas supporting preparatory control during task-switching performance in a combined fMRI and TMS study. Performance of a well-known cueing task-switching paradigm using fMRI was analysed to identify brain areas associated to preparatory task-switching; then, reducing excitability of those areas using TMS allowed to analyse its disruptive behavioural effects on this task. Following preceding results (Muhle-Karbe et al., 2014), it was hypothesized that if both prefrontal and parietal cortices implement switch-specific preparation processes, then TMS over these areas should interfere with performance in switch, but not in repeat trials. Alternatively, if the activity of either prefrontal or parietal cortices during task-switching relates to a more general switch-unspecific process, then TMS over these regions would interfere performance in both switch and repeat trials.

Materials and methods

Participants

A study combining fMRI and TMS techniques in two separate sessions with two different groups of participants was conducted. fMRI activation data were used to determine the most suitable target areas for TMS stimulation. Twenty healthy university students (mean age 26.8±1.3 years; 12 female) participated in the fMRI study. Each participant underwent a screening interview excluding sensory-motor problems, history of

neurological, or psychiatric problems, or substance abuse. Twenty healthy university students (mean age 29.3±6.8 years; 13 female) participated in the TMS study. All of them met the TMS safety criteria (Rossi et al., 2009). They had normal, or corrected to normal, visual acuity, and none were taking any medication with effects over the nervous system, or had history of neurological or psychiatric disorder, or drug/alcohol abuse. All subjects signed informed consent for their participation. The study was approved by the Ethics Committees of both the Universitat Oberta de Catalunya, and the Gregorio Marañón University Hospital. All participants gave written informed consent to participate in the study following the Declaration of Helsinki.

Experimental Task

The experimental task-switching paradigm was inspired by the classical numerical judgment task by Allport et al. (1994). Individuals were instructed to respond to target stimuli centred on the screen (a number between 1 and 9, excluding 5) according to one of two possible task rules: Odd/Even or >5/<5 (see Figure 1a). Task rules were indicated on the basis of a trial-by-trial task-cueing procedure being cues presented between 1000-2000 milliseconds prior to the target number. This long cue-to-target interval was selected to favour cue-related task preparation and associated switch-related BOLD activation to occur (Ruge et al., 2013). The cues were two different symbols presented during 300 milliseconds duration that appeared centred on the screen: "x" or "+" for the odd/even and >5/<5 tasks, respectively. All the stimuli were presented in white colour over a black background in Arial font with a physical dimension of 1 x 1 inches. The target display remained on the screen for 300 milliseconds with a response time limit of 3000 milliseconds from the target onset. Participants were instructed to respond to target numbers according to whether the number was odd (left index finger) or even (right index finger) if the preceding cue was the "x" symbol, or whether the number was < 5 (left index finger) or >5 (right index finger) if the preceding cue was the "+" symbol. Responses were immediately followed by a 100 milliseconds feedback indicating right, wrong, too fast, or too slow performance. The interval between each response and the next cue adopted a random value between 1000-3000 milliseconds (see Figure 1). Both, speed and accuracy were stressed in the written instructions presented at the beginning of the task. The experimental task was controlled by Presentation software (www.neurobs.com).

A

B

Figure 1: Stimulus material and experimental conditions. (a) Schematic of trial sequences illustrating the two experimental conditions being analysed. Switch trials involved a change in task rule as compared with the immediately preceding trial while repeat trials involved using the same rule as before (either odd/even task or > 5/< 5 task. (b) Each trial consisted of a visual cue ("x" or "+" symbols) followed by a visual target display with one number between 1 and 9, excluding 5. Participants were instructed to respond to target numbers according to whether the number was odd (left index finger) or even (right index finger) if the preceding cue was the "x" symbol, or whether the number was < 5 (left index finger) or >5 (right index finger) if the preceding cue was the "+" symbol. Responses were immediately followed by feedback indicating right, wrong, too fast, or too slow performance.

Participants from both fMRI and TMS experiment complete blocks of 154 trials organized semi-randomly in 42 series consisting of an initial switch trial being followed by a varying number of repeat trials (between 2 and 4). The switch condition included the initial trial of a series where the current cue indicated to perform a different task than the one performed in the preceding trial (either odd/even or >5/<5). The repeat condition included the last trial of a series (either the 2nd, the 3rd, or the 4th trial after a switch trial) where the current cue indicated to respond according to the same rule as in the preceding trials. The experimental session lasted around 12 min. Before the experimental task participants practice the task until the experimenter was sure that they had understood the instructions. Reaction Times (RTs) were measured in both the first and the last correct trials of each series, corresponding with switch and repeat tasks conditions, respectively.

Behavioural analyses

The percentage of correct responses and RTs to correct trials were analysed in the fMRI study as a function of the Task factor (switch versus repeat) by means of Student *t* tests. In addition, in the TMS study, the percentage of correct responses and RTs to correct trials were each analysed by means of a 3 x 2 repeated measure ANOVA with the factors Area of stimulation (Prefrontal, Parietal, Sham) and Task (switch versus repeat). A significance level of 0.05 was set for all contrasts. A Bonferroni-corrected significance level of $p < 0.05$ was adopted for all tests of simple effects involving multiple comparisons. SPSS version 22.0 statistical software package was used for all the analyses.

fMRI scanning

The fMRI data were acquired on a 3.0T SignaHDx MR scanner (GE Healthcare, Waukesha, WI) using an eight-channel head coil (GE Coils, Cleveland, OH). Head motion was minimized with a vacuum pack system molded to fit each subject. Functional images were obtained with a gradient echoplanar sequence using blood oxygenation level–dependent (BOLD) contrast, each comprising a full volume of 39 contiguous axial slices (3 mm thickness, 0 mm spacing) covering the whole brain. Volumes were acquired continuously with a repetition time (TR) of 3 s [TE = 31 ms, flip angle = 90, field of view (FOV) = 21.7 cm]. A total of 420 scans were acquired for each participant in a single session (21 min run), with the first five volumes subsequently discarded to allow for T1 equilibration effects. High-resolution T1-weighted spoiled gradient recall (SPGR)

anatomical images were also obtained for each participant (184 1.8-mm-thick axial images, TR = 5.5, TE = 2.3, FOV = 24 cm, 256x256 matrix).

The Presentation software package (https://www.neurobs.com/) was used for stimuli presentation. Stimuli were presented to participants through optic-fiber-based glasses (MRVision 2000 ultra, Resonance Technology, Inc., Northridge, USA) connected to the stimulation computer. Responses were registered with Lumina LP400 response pads for fMRI.

fMRI data analysis

The data were analysed using a general linear model in SPM12 (Welcome Department of Imaging Neuroscience, London, UK, www.fil.ion.ucl.ac.uk/spm/) implemented in MATLAB R2018a (Mathworks, Inc., Sherborn, MA). Individual scans were i) spatially realigned and unwarped to compensate for head-movement; ii) corrected for differences in slice acquisition time (slice timing correction); iii) spatially normalized and iv) spatially smoothed to reduce noise and to compensate for anatomical inter-subject variability (Gaussian filter of 8 mm FWHM), using standard SPM methods. Population inference was made through a two-stage procedure. At the first level, we specified in a subject-specific analysis where the event-related activity for each voxel, condition and subject was modelled using a canonical haemodynamic response function plus temporal and dispersion derivatives. For each individual, the effect of switch and repeat conditions was determined (switch>repeat, and repeat>switch). Statistical parametric maps of the t-statistic (SPM{t}) were generated for each subject and the contrast images were stored. In a second level random-effects analysis, these images were then combined in a one-sample t-test model was used (a $p<0.001$ at the voxel level was applied in all cases). The surviving activated voxels were superimposed on high-resolution structural magnetic resonance (MR) scans of a standard brain [Montreal Neurological Institute (MNI)]. Anatomical identification was performed with reference to the Talairach Daemon Software (http://www.talairach.org/) and also via XjView8 (http://www.alivelearn.net/xjview/). In addition, the neuroanatomy atlases by Haines (2011), and Nolte and Angevine (2007) were consulted.

TMS Methods

Before their participation a high-resolution structural MRI was obtained from each participant to rule out any brain abnormalities, to locate the targets for TMS, and to guide the stimulation during de TMS sessions. The MRI was done on a 1.5T scanner (Siemens Magnetom Essenza). Sequences obtained were FSPGR-T1 3D (180 1-mm-thick sagittal slices, TR = 500 milliseconds, TE = 50 milliseconds, 256×256 matrix, FOV = 24 cm), SE T1, FSE DP-T2, FLAIR, and diffusion.

All participants underwent three separate sessions in which they received one of the three stimulation conditions: TMS over prefrontal areas, TMS over parietal areas or sham stimulation positioning the coil tilted 90º over vertex (control condition). After each of the stimulation conditions the participants performed the task. The order of the sessions was counterbalanced across participants. The stimulation was performed using a Magstim Super Rapid 2 stimulator (Magstim Company Ltd., Whitland, U.K.) with a 70 mm figure of eight-coil and the position of the coil location during stimulation was guided by a combination of Brainsight (Rogue Research, Montreal Canada) frameless stereotaxic system and Polaris (Northern Digital, Waterloo, ON, Canada) infrared tracking system.

At the beginning of the first session, and before stimulation, active motor threshold (aMT) was individually determined in order to set the appropriate output TMS intensity for each participant. To calculate the aMT, the coil was placed tangentially over the participant's right M1, with the handle positioned 45˚ backwards. The coil was repositioned until the hot-spot of the left hand first dorsal interosseous (FDI) was located. The aMT was defined as the minimum stimulator output that produced movement in 5 of 10 trials in the FDI when the muscle was gently contracted (approximately 20% of the maximum voluntary contraction). The mean aMT value for the group was 53.4±6.2% of the maximum stimulator output and the intensity applied was the 80% of aMT. After determining aMT, and before stimulation, participants performed practice trials of the experimental task.

Continuous Theta Burst Stimulation (cTBS) was used in order to induce transient decreases of local cortical excitability lasting beyond the stimulation patterns (Huang et al., 2009; Huang et al., 2011). The cTBS protocol consists on a total of 600 TMS pulses grouped in trains of three pulses at 50 Hz, with each train being repeated every 200 milliseconds (5 Hz). The stimulation lasted 40 s in total and was delivered at 80% of the

aMT of each participant (Huang et al., 2009; Rossi et al., 2009). After the cTBS, participants completed the task in a PC with a 17-inch monitor, the response buttons were letters M and Z of the keyboard. As part of the security protocol, in each session participants completed Mini Mental State Examination and Beck Depression Inventory before and after TMS to control for any changes in mood or cognitive functioning.

Targeted cortical areas were localized individually using the high-resolution MRI and the coordinates obtained from the previous fMRI study. MNI coordinates corresponding to maximum activation during the task performance were x=48, y=34, z=22 for prefrontal area, and x=42, Y=-54, z=58 for parietal area. Since these coordinates correspond to the anterior portion of the IFJ (aIFJ) and the IPS, coordinates were adjusted for individual anatomy to the closest gyrus area to the aIFJ and the IPS using Brainsight. For sham stimulation, the coil was positioned on vertex and tilted 90° to mimic the sound of the stimulation patterns used in the active conditions.

Results

fMRI Results

Accuracy

Subjects performed the task efficiently during the fMRI session and committed less than 17% errors. The average percentage of correct trials was 94.9%.

The Student *t* test on the percentage of correct responses during switch and repeat trials revealed the presence of marginally significant differences (t(19)=-1.8; p< 0.088) in line with a switch cost phenomenon where participants achieved less correct trials after switch cues as compared to repeat cues (94.5% vs. 96%, respectively).

Reaction Times

The Student *t* test on RTs to switch and repeat trials revealed the presence of significant differences (t(19)=3.71; p< 0.001) indicating behavioural switch cost with an increase in RTs during switch as compared to repeat trials (737.2±34.9 vs. 697.7±29.4, respectively).

fMRI activation results

Functional images were analysed by SPM12 using a general linear model applied at each voxel across the whole brain. We localized those brain areas that modulated their activity during the switch and repeat events. We focused on voxels for which the difference between the responses to the switch events and the repeat conditions was statistically significant. Results showed that cue-related brain activation in the switch > repeat contrast involved the right frontal, both left and right parietal lobes and in the cerebellum (see Table 1; Figure 2a). Target-related brain activation in the switch > repeat contrast involved a widespread number of brain areas within prefrontal, parietal, temporal, occipital lobes, and cerebellum (see Table 2, and Figure 2b).

Table 1: Locations and t-values of the main clusters of fMRI activity for the switch>repeat contrast during cues.

Lobe	Region	BA	H	cluster	x	y	z	t-value
Frontal	Inferior Frontal Junction anterior	45	R	539	48	34	22	5.59
	Inferior Frontal Junction posterior	44	R		50	10	24	5.37
	Superior Frontal Gyrus	6	R	27	14	12	54	4.36
Parietal	Precuneus	7	L	137	-10	-76	60	5.04
	Intraparietal Sulcus	40	L	15	-32	-80	48	4.38
	Precuneus	7	R	553	8	-74	48	4.97
	Intraparietal Sulcus	40	R		42	-54	58	4.95
Cerebellum	Cerebellar Tonsil posterior	*	L	161	-22	-64	-32	7.57
	Cerebellar Tonsil posterior	*	R	205	36	-56	-38	5.56

Figure 2: (a) fMRI group activation map showing activated brain when comparing cues in the switch>repeat contrast. Two main clusters of activations were identified at the anterior portion of the inferior frontal junction, and the intraparietal sulcus. (b) fMRI group activation map showing activated brain when comparing targets in the switch>repeat contrast. Activation was identified at a widespread network of brain areas within prefrontal, parietal, temporal, occipital lobes, and cerebellum. Images thresholded at p < 0.001. Neurological convention is followed (left side of the brain is shown on the left side of the figure).

Table 2: Locations and t-values of the main clusters of fMRI activity for the switch>repeat contrast during targets.

Lobe	Region	BA	H	cluster	x	y	z	t-value
Frontal	Superior Frontal Gyrus	8	L	1027	-20	26	50	6.36
	Superior Frontal Gyrus	10	L	138	-26	54	28	6.06
	Superior Frontal Gyrus	6	R	110	34	14	50	4.58
	Inferior Frontal Gyrus	*	R	17	60	16	4	4.28
	Inferior Frontal Gyrus	*	R	668	48	36	4	5.82
	Precentral Gyrus	44	L	78	-54	16	6	4.33
	Postcentral Gyrus	*	L	91	-46	-20	28	4.85
	Postcentral Gyrus	*	L	829	-20	-26	66	5.93
	Middle Frontal Gyrus	46	R	4	48	38	26	3.78
	Middle Frontal Gyrus	*	L	427	-32	40	6	6.38
	Anterior Cingulate	*	L	3	-8	34	2	3.74
Parietal	Inferior Parietal Lobule	40	L	2172	-36	-62	48	9.48
	Postcentral Gyrus	5	R	6285	30	-44	66	7.97

Lobe	Region	BA	H	cluster	x	y	z	t-value
	Precuneus	7	L	528	-10	-72	34	5.13
	Precuneus	7	R		2	-58	36	4.67
Temporal	Middle Temporal Gyrus	*	L	130	-56	-46	-2	5.35
	Inferior Temporal Gyrus	*	L	89	-58	-36	-18	4.79
	Middle Temporal Gyrus	21	R	252	66	-54	10	5.49
Occipital	Lingual Gyrus	*	R	26	16	-92	2	4.30
	Cuneus	19	R	49	8	-78	30	4.17
	Cuneus	18	R	11	20	-68	16	3.98
	Cuneus	*	L	9	-16	-96	10	3.77
	Cuneus	*	L	6	-4	-78	16	3.73
	Lingual Gyrus	*	L	52	-24	-54	0	6.18
Sub-lobar	Insula	*	L	1508	-36	2	4	6.24
	Insula	*	R	651	40	2	4	7.89
Cerebellum	Cerebellar Tonsil	*	R	1496	34	-50	-42	6.96
	Culmen	*	R		32	-50	-26	5.34
	Inferior Semi-Lunar Lobule	*	L	2622	-42	-64	-38	6.37
	Culmen	*	L		-26	-56	-24	5.76

TMS results

Accuracy

Subjects performed the task efficiently during the TMS session and typically committed less than 13% errors across each of the three experimental conditions. The average percentage of correct trials across conditions was 95.8%.

The 3 x 2 repeated measures ANOVA on the percentage of correct responses using Area (aIFJ, IPS, and Sham stimulation) and Task (switch and repeat) as the factors revealed the presence of a main Task effect ($F_{(1, 19)}=10.1$; $p< 0.005$), indicating a switch cost phenomenon with less correct trials after switch cues as compared to repeat cues (95,6% vs. 97,7%, respectively). Neither the effect of Area ($F_{(2, 38)}=1.18$; $p= 0.889$), nor the Area x Task interaction ($F_{(2, 38)}= 7.19$; $p= 0.494$) reached significance.

Reaction Times

A 3 x 2 repeated measures ANOVA with RTs as the dependent variable was performed to compare the effect of TMS stimulation in the three different brain areas (aIFJ, IPS, and Sham) as a function of Task (switch and repeat). A significant main effect of Area was found ($F_{(2, 38)}= 5.94$; $p< 0.006$) indicating that, overall, the stimulation of the aIFJ resulted in a significant increase of RTs across Task conditions as compared to Sham stimulation ($p< 0.020$), being the effect of IPS stimulation only marginal ($p= 0.078$), and with no differences in RTs between aIFJ and IPS conditions ($p = 0.514$). The presence of a main effect of Task ($F_{(1, 19)}=11.75$; $p< 0.003$) confirmed results from accuracy analyses, regarding the presence of a significant switch cost effect when switch and repeat trials were compared (693.7 ± 32.4 vs. 657.8 ± 26.7 milliseconds, respectively). More importantly was the presence of an Area x Task interaction ($F_{(2, 38)}= 4.25$; $p< 0.022$). Post-hoc analyses of this interaction revealed that while the effect of the aIFJ stimulation disrupted RTs in both switch ($p< 0.02$) and repeat ($p< 0.043$) conditions as compared to Sham stimulation, the effect of the IPS stimulation had a specific negative impact over switch trials ($p< 0.012$) but not in repeat trials ($p< 0.783$). No differences were found when RTs to switch trials were compared between the aIFJ and the IPS stimulation ($p= 0.921$), or when they were compared in repeat trials ($p= 0.267$; see Figure 3).

Figure 3: Reaction times during task switching performance in the TMS experiment. Mean reaction times in milliseconds (± S.E.M) for the task switch and the task repeat conditions, across the three conditions of TMS: aIFJ (stimulation of the anterior portion

of the Inferior Frontal Junction), IPS (stimulation of the intraparietal sulcus), or Sham stimulation.

Discussion

The aim of the present fMRI-TMS combined study was investigating the disruptive effect of TMS over specific prefrontal and parietal cortices within a FPN of brain regions previously related to preparatory task-switching. As expected, the results of the fMRI study revealed a behavioural switch cost with an increase in RTs during switch as compared to repeat trials. This behavioural cost was accompanied by an increased activation in the IFJ (anterior and posterior portions), and the IPS during switch compared to repeat cues (see Tables 1, and Figure 2a). These findings are coherent with a large body of evidence suggesting the involvement of a FPN in preparatory process during task-switching paradigms (Dove et al., 2000; Gruber et al., 2006; Jamadar et al., 2010; Ruge et al., 2013). Interestingly, the same switch versus repeat contrast did not reveal activation differences in these brain regions during target processing. In this regard, switch-specific activation during target processing included a widespread network of brain areas within prefrontal, parietal, temporal, occipital lobes, and cerebellum (see Table 2, and Figure 2b). These results led us to select the aIFJ, and the IPS as the brain regions for the TMS study, since they reached the higher activation in prefrontal and parietal areas, respectively (see Table 1).

TMS results revealed that reducing excitability in the aIFJ, and IPS had dissociable behavioural effects. TMS over the aIFJ disrupted performance in both switch and repeat trails in terms of delayed responses as compared to Sham condition. In contrast, TMS over the IPS selectively interfered performance in switch trials (see Figure 3). These results suggest that the IPS would be related to the implementation of a switch-specific process while the aIFJ would be involved in a more general switch-unspecific process of task preparation. Functional neuroimaging and neurophysiological evidences support the present results showing that preparatory task-switching would involve both general and specific processes associated to certain prefrontal and parietal areas, respectively. For instance, using fMRI during cued task-switching blocks against single-task blocks, Braver et al. (2003) found that, while the activity in inferior prefrontal areas was selectively associated with mixing costs (i.e., the extra time it takes to perform two tasks simultaneously), only IPS activity was related to the magnitude of the switch cost. In

agreement with the present results, the authors interpreted that prefrontal areas would support sustained components of cognitive control, such as the maintenance demands associated with keeping multiple task-sets at a relatively high level of activation, common to both switch and repeat trials. In contrast, parietal areas would support a more transient control process associated with task-switching, such as the updating of goals or the linking of task cues to their appropriate stimulus-response mapping. In another study integrating event related potentials and fMRI data, Jamadar et al. (2010) found that informative task-switch cues (switch or repeat) evoked larger amplitudes in an early cue-locked positive component as compared to cues providing no information about the next task (non-informative cues). Importantly, this amplitude modulation correlated with fMRI activity in lateral prefrontal cortex (cued informative vs. cue non-informative contrast), consistent with a general goal activation process. The authors also reported a cue-locked late positivity that was larger for informatively cued switch vs. repeat cues, whose amplitude correlated with fMRI activity in the posterior parietal cortex (cued switch vs. cue repeat contrast), compatible with a stimulus-response rule activation process. These findings would indicate that lateral prefrontal brain regions seem to relate to general preparation processes, instead of specific switching operations. Also, in another ERP study, Barceló et al. (2008) found enhanced fronto-centrally distributed cue-locked P3 activity in response to informative cues in a task-switching paradigm irrespective of switch or repeat demands, compared to analogous irrelevant cues in an oddball control task. However, only a late cue-locked P3 component in parietal areas was specifically modulated in response to switch versus repeat equiprobable cues. The authors suggested an association between the fronto-centrally P3 and a general control mechanism related to preparatory process in frontal areas, and between the posterior P3 and a switch-specific reconfiguration process for stimulus-response mapping in parietal areas. Taken together, the above-mentioned evidences and the present results support a multi-component view of anticipatory preparation in task-switching (Ruge et al., 2013). It could be that a general preparation process, such as the retrieval of cue information for goal identification, necessarily during both switch and repeat trials, will recruit the aIFJ. In this vein, areas in the vicinity of the aIFJ reported here have been widely related to different components of memory retrieval. For instance, Nyhus and Badre (2015), associate BA 45 (also referred as mid-ventrolateral prefrontal cortex) to post-retrieval selection needed both to solve competition among multiple memory representations and to align remembered information with current task goals. Also, the application of

inhibitory TMS over mid-ventrolateral prefrontal cortex selectively diminishes recall for episodic details in an old/new memory-discrimination task, and lead the authors to attribute a causal role of this area in goal-directed retrieval (Wais et al., 2018). On the other hand, a switch-specific preparation processes such as the translation of abstract task goals into specific action rules or stimulus-response maps for task implementation necessarily during switch trials only, seemed to recruit the IPS. The IPS has been largely related to response selection processes, or more precisely, to stimulus-response mapping. Worringer et al. (2019) interpreted that stronger IPS activation in switch versus repeat trials may reflect the controlled mapping of a given stimulus to the response that is adequate according to the current task-set. IPS projections to ventral premotor areas support the role of this areas in action planning and reorienting of motor attention (Binkofski et al., 1998; Rushworth et al., 2003). The specific area of parietal activation during task-switching may reflect variations based on specific task stimulus attributes, with numerical tasks like the one used there (i.e., >5/<5, and odd/even) activating IPS within the right hemisphere (Shi et al., 2014).

It has to be noted that the present findings differ from a previous TMS study using a task-switching paradigm (Muhle-Karbe et al., 2014). As mentioned above, these authors found that the application of inhibitory TMS over both the IFJ and IPS interfered performance over switch trials only, suggesting that both prefrontal and parietal regions would implement switch-specific preparation processes. On the contrary, the present results revealed that while inhibitory TMS over the aIFJ interference performance in both switch and repeat trials, TMS over IPS produced interference only in switch trials. Although apparently contradictory, a fine-grained anatomical analysis revealed that two different regions within the IFJ were stimulated in these two studies. Particularly, the prefrontal brain area stimulated by Muhle-Karbe et al. (2014) (MNI: -40, 4, 32, BA 44) was more posterior to the aIFJ (MNI: 48, 34, 22, BA 45) reported in the present study (note that both areas were activated in the present fMRI study). Such a distinction between anterior and posterior portions within the IFJ is coherent with current models suggesting the existence of a hierarchical rostro-caudal organization in lateral prefrontal cortex based in the abstractness of action representations (Koechlin and Summerfield, 2007; Badre and D'Esposito, 2009; O'Relly, 2010). On the one hand, the aIFJ would be involved in a general preparatory process such as the retrieval of episodic information for goal selection (Buckner, 2003). This operation would take place at a relatively high level of abstraction

any time a cue is presented, and irrespective of switching or repeating task. On the other hand, the pIFJ would be involved in a more specific preparatory process such as the updating of task-goals representations (Derrfuss et al., 2005; Brass et al., 2005). Following Muhle-Karbe et al. (2014) while the pIFJ (BA 44) would provide an initial representation of task goals that needs to be selected during the next switch trial at a more abstract level, the IPS might specify the resulting stimulus-response mappings, thereby providing a more action-related task representation (Brass and von Cramon, 2004; Brass et al., 2005). These interpretations fit well with the functional fractionation between episodic and contextual control as proposed by the hierarchical model about the architecture of cognitive control in prefrontal cortex (Koechlin & Summerfield, 2007).

In conclusion, the pattern of TMS observed supports a model of task-switching preparation that involves both general and specific reconfiguration processes. Particularly, the present results reveal for the first time that inhibitory stimulation in the aIFJ (BA 45) may have caused interference in the retrieval of episodic information for goal selection during task-switching. This interpretation is coherent with current data about the role of aIFJ in memory retrieval under high control demands (Buckner, 2003). In addition, our results confirm the purported role of the IPS in more specific mechanisms of goal setting and implementation (Binkofski et al., 1998; Rushworth et al., 2003). Further research needs to clarify the pinpointed dissociation between the anterior and posterior regions of IFJ. Combining fMRI and TMS data might be critical to provide a more complete picture about the complex interaction between brain areas within the frontoparietal network and cognitive control operations during task-switching.

References

Allport DA, Styles EA, Hsieh S (1994) Shifting intentional set: Exploring the dynamic control of tasks. In: Conscious and Nonconscious Information Processing: Attention and Performance (Umilta C Moscovitch M, ed), pp421–452. Cambridge, MA: Oxford University Press.

Andersen RA, Snyder LH, Bradley DC, Xing J (1997) Multimodal representation of space in the posterior parietal cortex and its use in planning movements. Annu Rev Neurosci 20:303-330.

Badre D, D'Esposito M (2009) Is the rostro-caudal axis of the frontal lobe hierarchical? Nat Rev Neurosci 10:659-669.

Barceló F, Periáñez JA, Nyhus E (2008) An information theoretical approach to task-switching: evidence from cognitive brain potentials in humans. Front Hum Neurosci 28:1-13.

Bode S, Haynes JD (2009) Decoding sequential stages of task preparation in the human brain. Neuroimage 45:606-613.

Brass M, Von Cramon DY (2004) Decomposing components of task preparation with functional magnetic resonance imaging. J Cogn Neurosci 16:609-620.

Brass M, Ullsperger M, Knoesche TR, Von Cramon DY, Phillips NA (2005) Who comes first? The role of the prefrontal and parietal cortex in cognitive control, J Cogn Neurosci 17:1367-1375.

Binkofski F, Dohle C, Posse S, Stephan KM, Hefter H, Seitz RJ, Freund HJ (1998) Human anterior intraparietal area subserves prehension: a combined lesion and functional MRI activation study. Neurology 50:1253-1259.

Braver TS, Reynolds JR, Donaldson DI (2003) Neural mechanisms of transient and sustained cognitive control during task switching. Neuron 39:713-726.

Buckner RL (2003) Functional-anatomic correlates of control processes in memory

J Neurosci 23:3999-4004.

Culham JC, Kanwisher NG (2001) Neuroimaging of cognitive functions in human parietal cortex. Curr Opin Neurobiol 11:157-163.

Derrfuss J, Brass M, Neumann J, von Cramon DY (2005) Involvement of the inferior frontal junction in cognitive control: meta-analyses of switching and Stroop studies. Hum Brain Mapp 25:22-34.

Dove A, Pollmann S, Schubert T, Wiggins CJ, von Cramon, DI (2000) Prefrontal cortex activation in task switching: an event-related fMRI study. Brain Res Cogn Brain Res 9:103-109.

Gruber O, Karch S, Schlueter EK, Falkai P, Goschke T (2006) Neural mechanisms of advance preparation in task switching. Neuroimage 31:887-895.

Haines DE (2011) Neuroanatomía. Atlas de estructuras, secciones y sistemas. Lippincott Williams &Wilkins.

Huang YZ, Sommer M, Thickbroom G, Hamada M, Pascual-Leonne A, Paulus W, Classen J, Peterchev AV, Zangen A, Ugawa Y (2009) Consensus: New methodologies for brain stimulation. Brain Stimul 2:2-13.

Huang YZ, Rothwell JC, Chen RS, Lu CS, Chuang WL (2011) The theoretical model of theta burst form of repetitive transcranial magnetic stimulation. Clin Neurophysiol 122:1011-1018.

Jamadar S, Hughes M, Fulham WR, Michie PT, Karayanidis F (2010) The spatial and temporal dynamics of anticipatory preparation and response inhibition in task-switching. Neuroimage 51:432-449.

Kim C, Cilles SE, Johnson NF, Gold B (2012) Domain general and domain preferential brain regions associated with different types of task switching: a meta-analysis. Hum Brain Mapp 33:130-142.

Koechlin E, Summerfield C (2007) An information theoretical approach to prefrontal executive function. Trends Cogn Sci 1:229-235.

Kopp B, Tabeling S, Moschner C, Wessel K (2006) Fractionating the neural mechanisms of cognitive control. J Cogn Neurosci 18:949-965.

Logan GD, Bundesen C (2003) Clever homunculus: Is there an endogenous act of control in the explicit task-cuing procedure? J Exp Psychol Hum Percept Perform 29:575–599.

Mansfield EL, Karayanidis F, Cohen MX (2012) Switch-related and general preparation processes in task-switching: evidence from multivariate pattern classification of EEG data. J Neurosci 32:18253-18258.

Miller EK, Cohen JD (2001) An integrative theory of prefrontal cortex function. Annu Rev Neurosci 24:167-202.

Monchi O, Petrides M, Petre V, Worsley K, Dagher A (2001) Wisconsin Card Sorting revisited: distinct neural circuits participating in different stages of the task identified by event-related functional magnetic resonance imaging. J Neurosci 21:7733-7741.

Monsell S (2003) Task switching. Trends Cogn Sci 7:134-140.

Muhle-Karbe PS, Andres M, Brass M (2014) Transcranial magnetic stimulation dissociates prefrontal and parietal contributions to task preparation. J Neurosci 4:12481-12489.

Niendam TA, Laird AR, Ray KL, Dean YM, Glahn DC, Carter CS (2012) Meta-analytic evidence for a superordinate cognitive control network subserving diverse executive functions. Cogn Affect Behav Neurosci 12:241–268.

Nyhus E, Badre D (2015) Memory retrieval and the functional organization of frontal cortex. In: The Wiley Handbook on the Cognitive Neuroscience of Memory (Addis DR, Barense M, Duarte A, eds), pp131–149. West Sussex, UK: John Wiley & Sons, Ltd.

Nolte N, Angevine JB (2007) The Human Brain in Photographs and Diagrams, 3rd Edition. Philadelphia, PA: Mosby Elsevier

O'Reilly RC (2010) The What and How of prefrontal cortical organization. Trends Neurosci 33:355-361.

Periáñez JA, Maestú F, Barceló F, Fernández A, Amo C, Ortiz Alonso T (2004) Spatiotemporal brain dynamics during preparatory set shifting: MEG evidence. Neuroimage 21:687-695.

Petersen SE, Posner MI (2012) The attention system of the human brain: 20 years after. Annu Rev Neurosci 35:73-89.

Rossi S, Hallett M, Rossini PM, Pascual-Leone A (2009) Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. Clin Neurophysiol 120:2008–2039.

Ruge H, Jamadar S, Zimmermann U, Karayanidis F (2013) The many faces of preparatory control in task switching: reviewing a decade of fMRI research. Hum Brain Mapp 34:12-35.

Rushworth MFS, Hadland KA, Paus T, Sipila PK (2002) Role of the human medial frontal cortex in task switching: a combined fMRI and TMS study. J Neurophysiol 87:2577-92.

Shi Y, Meindl T, Szameitat AJ, Müller HJ, Schubert T (2014) Task preparation and neural activation in stimulus-specific brain regions: an fMRI study with the cued task-switching paradigm. Brain Cogn 87:39-51.

Sohn MH, Ursu S, Anderson JR, Stenger VA, Carter CS (2000) The role of prefrontal cortex and posterior parietal cortex in task switching. Proc Natl Acad Sci U S A 97:13448-13453.

Stuss DT, Picton TW (1978) Neurophysiological correlates of human concept formation. Behav Biol 23:135-162.

Wais PE, Montgomery O, Stark CEL, Gazzaley A (2018) Evidence of a Causal Role for mid-Ventrolateral Prefrontal Cortex Based Functional Networks in Retrieving High-Fidelity Memory. Sci Rep 8:14877.

Worringer B, Langner R, Koch I, Eickhoff SB, Eickhoff CR, Binkofski FC (2019) Common and distinct neural correlates of dual-tasking and task-switching: a meta-

analytic review and a neuro-cognitive processing model of human multitasking. Brain Struct Funct 224:1845-1869.

Wylie G, Allport A (2000) Task switching and the measurement of "switch costs." Psychol Res 63:212–233.