Age but not menopausal status is linked to lower resting energy expenditure

Jari E. Karppinen[1], Petri Wiklund[2], Johanna K. Ihalainen[1], Hanna-Kaarina Juppi[1,3], Ville Isola[1], Matti Hyvärinen[1,3], Essi K. Ahokas[1], Urho M. Kujala[1], Jari Laukkanen[4,5], Juha J. Hulmi[1], Juha P. Ahtiainen[1], Sulin Cheng[1,6] & Eija K. Laakkonen[1,3]

[1]Faculty of Sport and Health Sciences, University of Jyväskylä, Finland

[2]Huawei Helsinki Research Center, Huawei Technologies Oy (Finland) Co. Ltd, Helsinki, Finland.

[3]Gerontology Research Center, University of Jyväskylä, Finland

[4]Central Finland Health Care District, Jyväskylä, Finland,

[5]Institute of Clinical Medicine, University of Eastern Finland, Kuopio, Finland

[6]Exercise Translational Medicine Centre, Shanghai Jiao Tong University, Shanghai, China.

Keywords: resting energy expenditure, menopause, estrogen, hormone replacement therapy

Corresponding author: Jari E. Karppinen, jari.e.j.karppinen@jyu.fi, Rautpohjankatu 8, 40700 Jyväskylä, Finland

Supported by the Academy of Finland grants 309504, 314181, 335249 (to EKL) and 135038 (to SC) and by the Ministry of Education and Culture of Finland grants OKM/54/627/2006, OKM/77/627/2008 (to SC) and OKM/10/626/2021 (to JKI). The manuscript preparation was supported by the Academy of Finland grant 330281 (to EKL). The contents of this article are solely the responsibility of the authors.

Author disclosures: The authors report no conflicts of interest.

ABSTRACT

Context

It remains uncertain whether aging before late adulthood and menopause are associated with fat-free mass and fat mass–adjusted resting energy expenditure (REE_{adj}).

Objectives

We investigated whether REE_{adj} differs between middle-aged and younger women and between middle-aged women with different menopausal statuses. We repeated the age group comparison between middle-aged mothers and their daughters to partially control for genotype. We also explored whether serum estradiol and follicle-stimulating hormone concentrations explain REE_{adj} in midlife.

Methods

We divided 120 women, including 16 mother–daughter pairs, into age groups; group I ($n = 26$) consisted of participants aged 17–21, group II ($n = 35$) of those aged 22–38 and group III ($n = 59$) of those aged 41–58 years. The women in group III were further categorized as pre- or perimenopausal ($n = 19$), postmenopausal ($n = 30$) or postmenopausal hormone therapy users ($n = 10$). REE was assessed using indirect calorimetry, body composition using dual-energy X-ray absorptiometry and hormones using immunoassays.

Results

The REE_{adj} of group I was 126 kcal/d (95% CI: 93–160) higher than that of group III, and the REE_{adj} of group II was 88 kcal/d (95% CI: 49–127) higher. Furthermore, daughters had a 100 kcal/d (95% CI: 63–138 kcal/d) higher REE_{adj} than their middle-aged mothers (all $P < 0.001$). In group III, REE_{adj} was not lower in postmenopausal women and did not vary by sex hormone concentrations.

Conclusions

We demonstrated that REE_{adj} declines with age in women before late adulthood, also when

controlling partially for genetic background, and that menopause may not contribute to this

decline.

Introduction

Energy expenditure is often assumed to begin declining in early to middle adulthood, but

Pontzer et al. (1) challenged this assumption by showing that fat-free mass (FFM) and fat

mass (FM)-adjusted total energy expenditure (TEE_{adj}) were stable between the ages of 20 and

63. However, they found that similarly adjusted resting energy expenditure (REE_{adj}) stabilizes

at adult levels at age 18 and declines from age 46 onwards, although the limited numbers of

middle-aged participants with a measured REE prevented the authors from making definitive

inferences about the onset of REE_{adj} decline, leading to a conclusion that age does not affect

energy expenditure in adults before the age of 60 (1). Nevertheless, previous studies are

consistent with an earlier turning point for REE_{adj} (2–4), and we therefore sought to assess

whether REE_{adj} declines before late adulthood. Our dataset also included mother–daughter

dyads, some of which had TEE measured with doubly labeled water, enabling us to partly

control the analyses for genetic background and to explore whether increasing age showed

similar associations with TEE_{adj} as it does with REE_{adj}.

Like aging, menopause is widely believed to reduce REE_{adj}, and the topic has

broad interest because many women gain FM during the menopausal transition (5,6) and

associate the change in body composition with slowing metabolism. During the menopausal

transition, ovarian follicular activity ceases, causing a striking shift in women's sex hormone

profile. The decline in systemic estradiol (E2) concentration in particular is thought to

decrease REE_{adj}, potentially via both central (7) and peripheral (8) mechanisms, while the

increase in follicle-stimulating hormone (FSH) secretion may also play a role (9,10).

Menopausal hormone therapy (MHT) can restore E2 and decrease FSH levels, which should

reverse the potential menopause-associated decline in REE_{adj}. However, whether menopause

truly decreases REE_{adj} is still uncertain because longitudinal studies following women over

the menopausal transition (11,12), cross-sectional studies comparing women with different

menopausal states (13–16) and MHT interventions (15,17–19) have been inconclusive.

Therefore, in addition to investigating whether REE_{adj} differs between young and middle-aged

women, we also assessed whether REE_{adj} differs between middle-aged women with different

menopause statuses. We restricted the menopause analysis to middle-aged participants to limit

the confounding effects of age. We also explored whether serum E2 and FSH concentrations

explain REE_{adj} in midlife.

Materials and Methods

Participants

The participants were 120 women who had taken part in one of four studies performed at the

Faculty of Sport and Health Sciences of the University of Jyväskylä (**Figure 1**). They were

required to be healthy and not taking medications that could affect metabolism, although

hormonal contraception and MHT use were allowed.

The Calex study (data collection 2008–2011) investigated whether lifestyle

factors influence muscle and adipose tissues (20). The current study used data from 17

middle-aged and 21 younger women with measured REE. This dataset included 16 mother–

daughter pairs; ten pairs also had their TEEs measured. The data has been partly used in an

earlier validation study (21). The EsmiRs study (Estrogen and microRNAs as Modulators of

Women's Metabolism, 2019–2020) examined resting and exercise metabolism in middle-aged

women (22), and we included all 42 participants with measured REEs from that study. The

Physique study (23) investigated the effects of competition weight loss in normal-weight

participants; here, we used the baseline data from 23 young women collected before their

weight loss. Finally, the Athletic Performance and Nutrition study (NO RED-S, 2021,

Ihalainen et al. unpublished) studied the health of winter sports athletes at the beginning of

their training and competition seasons; from that study, we used baseline data from 17 young

women, measured during the transition season when their training load was the lowest.

105   Studies were conducted according to the Declaration of Helsinki and approved

by the Ethics Committee of the Central Finland Health Care District (Calex; memo 22/8/2008

and 5/2009, EsmiRs; 9U/2018, Physique; 19U/2018) or the Ethics Committee of the

University of Jyväskylä (NO RED-S 514/13.00.04.00/2021). Participants gave informed

consent.

Figure 1

Age categorization

We used age as a continuous variable and categorized the participants into three age groups,

based on previous findings. REE_{adj} plateaued at age 18.0 (95% CI: 16.8–19.2) and started to

decline at age 46.5 (95% CI: 40.6–52.4), while TEE_{adj} plateaued at 20.5 (95% CI: 19.8–21.2)

and started to decline at 63.0 years (95% CI: 60.1–65.9) (1). We therefore assigned

participants aged 18~21 years to group I, 22~39 years to group II and 40~60 years to group

III.

Menopausal status

Group I and II participants were naturally menstruating women (group I, $n = 12$; group II, $n = 20$) or hormonal contraception users (group I, $n = 14$; group II, $n = 15$). We determined the menopausal status of group III women with the Stages of Reproductive Aging Workshop (STRAW) + 10 guidelines (24) using menstrual and serum FSH data: 11 were premenopausal (PRE), eight were perimenopausal (PERI), 30 were postmenopausal (POST) and ten were postmenopausal MHT users (MHT). We combined the PRE and PERI women into a PRE/PERI group to represent women with meaningful ovarian E2 production but performed a sensitivity analysis without the PERI women because E2 levels decline in perimenopause. All the MHT group women used E2 alone ($n = 1$) or with a progestogen ($n = 9$). The administration was oral in seven cases and transdermal in three. Details concerning menopausal status determination and the use of hormonal products are in **Supplemental Methods**.

Sex hormones

For sex hormone assessment, the serum was separated from fasting venous blood samples according to standard procedures and stored at −80°C. E2 concentrations were measured for all participants, except seven young women in the Calex study, and FSH concentrations were measured for group III participants using enzyme-amplified chemiluminescence immunoassays (IMMULITE 1000 or 2000 XPi, Siemens Medical Solution Diagnostics, Los Angeles, CA, USA). The analytical sensitivity for the E2 kit (Siemens, Llanberis, UK) is 0.055 nmol/l with an accurate reportable range of 0.073–7.342 nmol/l. The coefficient of variation in our lab using control samples has been 15%. We compared the used immunoassay method with liquid chromatography-mass spectrometry (HUSLAB, Helsinki

University Hospital, Helsinki, Finland) and found a good correlation in all test samples ($n =$

166, $r = 0.91$). However, when the comparison was restricted to samples with E2

concentrations less than 0.1 nmol/l, as determined by with liquid chromatography-mass

spectrometry, the correlation between methods was lower ($n = 76$, $r = 0.42$). The analytical

sensitivity of the FSH kit (Siemens, Llanberis, UK) is 0.1 IU/l, and the coefficient of variation

in our lab has been 5%.

Body composition

Body composition was assessed with dual-energy X-ray absorptiometry (DXA; DXA

Prodigy, GE Lunar Corp., Madison, WI, USA). We calculated the appendicular lean mass

index (ALMI) by scaling appendicular lean mass (kg) to height (m) squared.

Resting and total energy expenditure

REE was measured in all studies using the same Vmax Encore 92 metabolic cart

(Sensormedics, Yorba Linda, CA, USA) and ventilated hood in the same thermoneutral

laboratory; the cart was calibrated accordingly before each measurement. The REE

assessment details are in **Supplemental Methods**. Measurements were performed in the

morning after overnight fasting, with resting periods of 0–30 min and measurement periods of

15–30 min. We excluded at least the first 5 min of measurement data; for all participants, we

located a steady-state period of at least 5 min during which the coefficient of variation was

10% or less for $\dot{V}O_2$ and $\dot{V}CO_2$, and we calculated REE with the modified Weir equation (25).

We made REE comparable between different-sized participants using residuals—the

differences between measured and predicted values—from a linear regression model:

$$\text{REE (kcal/d)} = 413.4 + 20.2 \times \text{FFM (kg)} + 1.7 \times \text{FM (kg)}$$

Herein, we refer to residual REE as REE_{adj}. We also built three alternative explanatory

models (**Supplemental Table 1**): the first included age as a covariate with FFM and FM; to

the second we added ALMI; and in the third, the study data collection period.

To assess TEE in the Calex study, the overnight fasting participants gave a urine

sample and ingested a doubly labeled water dose of 1 g per kg of body mass (21). A second

urine sample was collected after 4–6 h, and a third 14 days later. The samples were analyzed

in triplicate using mass spectrometry (Metabolic Solutions Inc., Merrimack, NH, USA) at the

University of Alabama. TEE was calculated as in Schoeller et al. (26).

Statistical analyses

We performed the statistical analyses using R 4.2.1 (27). The analytic code is available

without restriction at https://doi.org/10.5281/zenodo.7684731. We report descriptive statistics

as means with standard deviations or as medians with first and third quartiles, but we did not

test group differences in order to preserve statistical power. We verified the model's

assumptions before accepting the results and used an alpha level of 0.05 for statistical

significance.

We estimated the association between age and REE_{adj} and compared the

measured REE and REE_{adj} between the age groups with linear mixed-effect models using the

nlme package (28), with family identification as a random effect. We also performed a

sensitivity analysis using FFM, FM and ALMI-adjusted REE residuals as the outcome. For

the mother–daughter pairs, we first compared the REE_{adj} and then TEE using the measured

TEE as the outcome and FFM and FM as covariates. We estimated intraclass correlation

coefficients using the *psych* package (29) with one-way random-effects models—intraclass

correlation compares within- and between-pair variations, thereby expressing how strongly

the mothers and daughters resemble each other.

193   Last, we used linear regression to compare the measured REE and REE$_{adj}$

194   between the menopause groups using the POST group as the reference. Given that body

195   composition parameters may explain REE differently among women of different ages, we

196   performed supporting analyses using measured REE as the outcome, menopause status or sex

197   hormone concentrations as the explanatory variable, and FFM, FM and age as covariates in

198   separate regression models.

Results

Participant characteristics

202   **Table 1** shows participant characteristics and energy expenditures across age and menopause

203   groups, and **Supplemental Table 2** shows the same in the mother–daughter pairs. Based on

204   the descriptive statistics, group II women had higher FFM and lower FM, and in group III,

205   sex hormone concentrations varied between the menopause groups, as expected.

206   *****Table 1*****

Age and energy expenditure

FFM and FM explained 47% of the REE variance, while the inclusion of age increased the

adjusted R^2 to 68%. **Figure 2** illustrates how age impacts REE estimation by presenting the

relationships between the predicted and measured REE values. Neither the ALMI nor

information on the study data collection period improved the explanatory value.

Supplemental Table 1 presents the full results.

Figure 2

Age was inversely associated with REE_{adj} (B = -3.9; 95% CI: -4.8 to -3.1; $P <$

0.001). Group I had 140 kcal/d (95% CI: 82–199) higher measured REE and 126 kcal/d (95%

CI: 93–160) higher REE_{adj} (**Figure 3A**) than group III, while group II had 187 kcal/d (95%

CI: 133–240) higher measured REE and 88 kcal/d (95% CI: 49–127) higher REE_{adj} ($P <$

0.001 for all). The group differences in the FFM, FM and ALMI-adjusted REE were slightly

smaller (**Supplemental Table 3**).

In the 16 mother–daughter pairs, the daughters had 100 kcal/d (95% CI: 63–138;

$P < 0.001$) higher REE_{adj} than their mothers (**Figure 3C**). In the ten pairs with REE and TEE

measurements, the daughters had 85 kcal/d (95% CI: 45–125; $P = 0.003$) higher REE_{adj} than

their mothers, but there was no significant difference in TEE_{adj} ($B = 26$ kcal/d; 95% CI: -128–

180; $P = 0.75$, **Figure 3D**). Intraclass correlation coefficients were 0.05 (95% CI: -0.43–0.52;

$P = 0.42$) for REE_{adj}, 0.83 (95% CI: 0.49–0.96; $P < 0.001$) for TEE and 0.92 (95% CI: 0.71–

0.98; $P < 0.001$) for TEE_{adj} (**Supplemental Figure 1**).

Figure 3

Resting energy expenditure in midlife

Compared with the POST group, the measured REE was not significantly different in either

the PRE/PERI ($B = 46$ kcal/d; 95% CI: -26–119; $P = 0.21$) or MHT groups ($B = -27$ kcal/d;

95% CI: -118–63; $P = 0.55$); neither was REE_{adj} (PRE/PERI; $B = 7$ kcal/d; 95% CI: -42–55; P

$= 0.78$; MHT: $B = -31$ kcal/d; 95% CI: -91–30; $P = 0.31$, **Figure 3B**). The exclusion of PERI

women did not alter the results (data not shown). In the specific models generated for the

middle-aged subsample that controlled for FFM, FM and age, the MHT group had a lower

REE than the POST group (**Table 2**). Furthermore, neither E2 nor FSH showed a statistically

significant association with REE.

Table 2

Discussion

Aging through adulthood and menopause are thought to slow basal metabolism, potentially

predisposing women to obesity. This study demonstrates that increasing age is associated with

a decline in REE_{adj} among young and middle-aged women, also after partly controlling for

genetic background. However, menopause did not contribute to the age-associated decline in

REE_{adj}. Furthermore, E2 or FSH concentrations were not related to REE_{adj} in middle-aged

women.

The decline in age-associated REE_{adj} aligns with previous cross-sectional studies

by Pontzer et al. (1), Geisler et al. (3) and Siervo et al. (4), in which the decline began in

women at 46.5, 35.2 and 47 years of age, respectively. The turning point for REE_{adj} therefore

may occur before 60 years of age (1), but whether the phenomenon represents an actual

slowing of cellular metabolism remains unclear. Low metabolic rate organs contribute more

to FFM as we age (30), but aging also causes tissue quality changes; for example, the brain

grey–white matter ratio (31) and skeletal muscle density (32) decline from young to middle

adulthood, meaning that each kilogram of the brain or muscle tissue has fewer metabolically

active cells as aging proceeds. Current body composition assessment methods, like DXA,

cannot detect such changes (30), so FFM adjustments that assume FFM composition and

quality are constant may overestimate the age-associated decline in $REE_{adj.}$ Indeed, Roubenoff

et al. (33) found no association between age and FFM-adjusted REE when assessing FFM

using total body potassium analysis, which directly estimates cell mass. Therefore, the direct

influence of age on REE_{adj} may be less impactful than initially observed, but it may enhance

REE estimation (as shown in **Figure 2B**) by enabling adjustments for age-related changes in

body composition within the model. If slowing cellular metabolism contributes to the

observed decline in REE_{adj}, it may result from altered systemic hormone and cytokine

stimulation (30), with intrinsic changes in hormone responsiveness, protein synthesis,

maintenance of membrane potentials and mitochondrial function (34).

267   We also compared the TEE_{adj} of middle-aged mothers and their daughters and

268   found that they were similar, despite differences in their REE_{adj}, indicating that TEE is highly

269   heritable (35). This also suggests that the possible age-associated decline in REE_{adj} may have

270   a negligible effect on TEE before late adulthood, as also reported by Pontzer et al. (1).

271   Our findings concur with previous research that menopausal status and sex

hormone levels do not robustly determine REE_{adj} during midlife. Although ovarian hormone

suppression studies in premenopausal women have shown some yet not fully convincing

evidence of a REE_{adj} decline (36,37), MHT interventions show no clear effect on REE in

postmenopause (15,17,18). In the present study, the MHT users had even lower REE than

postmenopausal women who did not use MHT after adjusting for FFM, FM and age.

However, given that the MHT group was the smallest in our study, this difference is unlikely

to be attributed to MHT use *per se*. Observational evidence also indicates that menopause has

a minimal impact on REE_{adj}. For instance, Lovejoy et al. (11) found that sleeping energy

expenditure and TEE_{adj} decreased in women transitioning from premenopause to

postmenopause during a four-year follow-up, but the changes were no different from

participants who remained premenopausal, which suggests that the decreases were related to

aging, not menopause. Furthermore, the menopausal transition was not associated with REE

in the longitudinal MONET study (12), and cross-sectional studies have shown no association

between menopausal status and measured REE (14,16), FFM-adjusted REE (13,15) or FFM-

adjusted TEE (38), while a study that did show a higher REE in MHT users than non-users

(14) failed to adjust the analyses for differing tissue masses. Finally, Pontzer et al. (1) found

no differences in REE_{adj} and TEE_{adj} trajectories between middle-aged women and men,

indicating that sex specific changes in energy expenditure are not observed during midlife.

The lack of a clear association between menopause and REE_{adj} is unexpected

because the cessation of reproductive functions and altered hormonal profile should decrease

basal metabolism. There are at least three potential explanations; first, with the limitations of

body composition assessment and indirect calorimetry methods (30), the energy expenditure

of female reproductive processes may be so small relative to other functions contributing to

REE that its loss is difficult to detect. The second explanation is that the effects of menopause

cannot be disentangled from the effects of aging, especially because aging progresses

differently between individuals, although this is insufficient to explain why MHT

interventions do not increase REE (15,17,19). The third, more speculative explanation is that

women reallocate energy during menopause from reproduction to other purposes; Hazda

hunter-gatherer women, for example, increase the time spent on gathering resources for their

offspring (39), potentially reallocating the freed energy to movement and maintenance of the

locomotor system (40). Women in industrialized nations live differently and may therefore

use the energy to build energy reserves and bodily defense mechanisms, reallocating the freed

energy inside the REE component (40). Increased FM, especially to the trunk, may further

promote metabolic deterioration (41), inflammation (42) and sympathetic nervous system

activity (43), thereby further elevating the REE (44–47). Such trade-offs could explain why

women's REE_{adj} does not decline and their cardiometabolic risk profile worsens after

menopause in industrialized societies (48).

Finally, it should be mentioned that a decrease in absolute REE following

skeletal muscle loss could also contribute to menopause-associated FM accumulation if

women do not match the drop by reducing energy intake. We previously reported that a peri-

to postmenopausal transition during a mean follow-up of 14 months resulted in a 0.2 kg lean

mass loss, likely from skeletal muscle (49); but, assuming that the mass-specific metabolic

rate of skeletal muscle is 13 kcal/d at rest (50), the loss would reduce REE by 2.6 kcal/d,

which cannot explain the 0.8 kg increase in FM (6), especially as the tissue changes happen

gradually.

The main limitations of this study are its cross-sectional and secondary nature.

We pooled existing studies to generate a sufficiently large sample and cannot entirely exclude

a clustering effect. Considering the impact of sex hormones, we did not control for the use of

hormonal products or menstrual cycle phases. Furthermore, serum E2 concentrations were

analyzed with immunoassays, whose accuracy is limited for the low E2 levels seen post-

menopause.

In conclusion, REE adjusted for DXA-measured FFM and FM declines in

women from young to middle adulthood, likely due to aging rather than menopause, but

whether falling cellular metabolic rates contribute is unclear. Current evidence does not

support the inference that menopause reduces REE. Longitudinal data from middle-aged

women with differing sex hormone trajectories are needed to reconcile whether menopause

truly affects REE in a meaningful way.

Acknowledgments

We thank the participants for their time and effort and the staff of the Faculty of Sport and

Health Sciences of the University of Jyväskylä.

The authors' responsibilities were as follows—JEK conducted EsmiRs study measurements,

gave guidance for the Physique and NO RED-S studies, analyzed data, and wrote the

manuscript. EKL led and designed the EsmiRs study with the help of JEK, H-KJ, MH, UMK,

and JL. PW conducted Calex study measurements and gave guidance for the EsmiRs study.

SC led and designed the Calex study with help from PW. JPA and JH led and designed the

Physique study with the help of VI, who also participated in data collection. JKI led,

designed, and conducted data in the NO RED-S study with the help of EKA. All authors

revised the manuscript and approved its final version.

Data availability

Restrictions apply to the availability of all data generated or analyzed during this study to

preserve patient confidentiality. The principal investigators of each study (Calex: CS and

EKL, EsmiRs [doi.10.17011/jyx/dataset/83491]: EKL, Physique: JPA, and NO RED-S: JKI)

will on request detail the restrictions and any conditions under which access to some data may

be provided.

REFERENCES

1. Pontzer H, Yamada Y, Sagayama H, Ainslie PN, Andersen LF, Anderson LJ, et al. Daily energy expenditure through the human life course. Science. 2021 Aug 13;373(6556):808–12.

2. Müller MJ, Bosy-Westphal A, Klaus S, Kreymann G, Lührmann PM, Neuhäuser-Berthold M, et al. World Health Organization equations have shortcomings for predicting resting energy expenditure in persons from a modern, affluent population: generation of a new reference standard from a retrospective analysis of a German database of resting energy expenditure. Am J Clin Nutr. 2004 Nov;80(5):1379–90.

3. Geisler C, Braun W, Pourhassan M, Schweitzer L, Glüer CC, Bosy-Westphal A, et al. Gender-Specific Associations in Age-Related Changes in Resting Energy Expenditure (REE) and MRI Measured Body Composition in Healthy Caucasians. J Gerontol A Biol Sci Med Sci. 2016 Jul;71(7):941–6.

4. Siervo M, Oggioni C, Lara J, Celis-Morales C, Mathers JC, Battezzati A, et al. Age-related changes in resting energy expenditure in normal weight, overweight and obese men and women. Maturitas. 2015 Apr;80(4):406–13.

5. Greendale GA, Sternfeld B, Huang M, Han W, Karvonen-Gutierrez C, Ruppert K, et al. Changes in body composition and weight during the menopause transition. JCI Insight. 2019 Mar 7;4(5).

6. Juppi HK, Sipilä S, Fachada V, Hyvärinen M, Cronin N, Aukee P, et al. Total and regional body adiposity increases during menopause-evidence from a follow-up study. Aging Cell. 2022 Jun;21(6):e13621.

7. Xu Y, López M. Central regulation of energy metabolism by estrogens. Mol Metab. 2018 Sep;15:104–15.

8. Klinge CM. Estrogenic control of mitochondrial function. Redox Biol. 2020 Apr;31:101435.

9. Liu P, Ji Y, Yuen T, Rendina-Ruedy E, DeMambro VE, Dhawan S, et al. Blocking FSH induces thermogenic adipose tissue and reduces body fat. Nature. 2017 Jun 1;546(7656):107–12.

10. Mattick LJ, Bea JW, Singh L, Hovey KM, Banack HR, Wactawski-Wende J, et al. Serum Follicle-Stimulating Hormone and 5-Year Change in Adiposity in Healthy Postmenopausal Women. J Clin Endocrinol Metab. 2022 Jul 14;107(8):e3455–62.

11. Lovejoy JC, Champagne CM, de Jonge L, Xie H, Smith SR. Increased visceral fat and decreased energy expenditure during the menopausal transition. Int J Obes (Lond). 2008 Jun;32(6):949–58.

12. Duval K, Prud'homme D, Rabasa-Lhoret R, Strychar I, Brochu M, Lavoie JM, et al. Effects of the menopausal transition on energy expenditure: a MONET Group Study. Eur J Clin Nutr. 2013 Apr;67(4):407–11.

13. Hodson L, Harnden K, Banerjee R, Real B, Marinou K, Karpe F, et al. Lower resting and
total energy expenditure in postmenopausal compared with premenopausal women
matched for abdominal obesity. J Nutr Sci. 2014 Feb 13;3:e3.

14. Monda V, Salerno M, Fiorenzo M, Villano I, Viggiano A, Sessa F, et al. Role of Sex
Hormones in the Control of Vegetative and Metabolic Functions of Middle-Aged
Women. Front Physiol. 2017;8:773.

15. Bessesen DH, Cox-York KA, Hernandez TL, Erickson CB, Wang H, Jackman MR, et al.
Postprandial triglycerides and adipose tissue storage of dietary fatty acids: impact of
menopause and estradiol. Obesity (Silver Spring). 2015 Jan;23(1):145–53.

16. Gould LM, Gordon AN, Cabre HE, Hoyle AT, Ryan ED, Hackney AC, et al. Metabolic
effects of menopause: a cross-sectional characterization of body composition and exercise
metabolism. Menopause. 2022 Feb 28;29(4):377–89.

17. Marlatt KL, Lovre D, Beyl RA, Tate CR, Hayes EK, Burant CF, et al. Effect of
conjugated estrogens and bazedoxifene on glucose, energy and lipid metabolism in obese
postmenopausal women. Eur J Endocrinol. 2020 Oct;183(4):439–52.

18. Anderson EJ, Lavoie HB, Strauss CC, Hubbard JL, Sharpless JL, Hall JE. Body
composition and energy balance: lack of effect of short-term hormone replacement in
postmenopausal women. Metabolism. 2001 Mar;50(3):265–9.

19. Lwin R, Darnell B, Oster R, Lawrence J, Foster J, Azziz R, et al. Effect of oral estrogen
on substrate utilization in postmenopausal women. Fertil Steril. 2008 Oct;90(4):1275–8.

20. Ojanen X, Cheng R, Törmäkangas T, Rappaport N, Wilmanski T, Wu N, et al. Towards
early risk biomarkers: serum metabolic signature in childhood predicts cardio-metabolic
risk in adulthood. EBioMedicine. 2021 Oct;72:103611.

21. Yu Z, Völgyi E, Wang R, Ember A, Wiklund P, Alén M, et al. Comparison of heart rate
monitoring with indirect calorimetry for energy expenditure evaluation. Journal of Sport
and Health Science. 2012 Dec 1;1(3):178–83.

22. Karppinen JE, Juppi HK, Hintikka J, Wiklund P, Haapala EA, Hyvärinen M, et al.
Associations of resting and peak fat oxidation with sex hormone profile and blood glucose
control in middle-aged women. Nutr Metab Cardiovasc Dis. 2022 Sep;32(9):2157–67.

23. Isola V, Hulmi JJ, Petäjä P, Helms ER, Karppinen JE, Ahtiainen JP. Weight loss induces
changes in adaptive thermogenesis in female and male physique athletes. Appl Physiol
Nutr Metab. 2023 Jan 16;

24. Harlow SD, Gass M, Hall JE, Lobo R, Maki P, Rebar RW, et al. Executive summary of
the Stages of Reproductive Aging Workshop + 10: addressing the unfinished agenda of
staging reproductive aging. Menopause. 2012 Apr;19(4):387–95.

25. Weir JB. New methods for calculating metabolic rate with special reference to protein
metabolism. 1949. Nutrition. 1990 Jun;6(3):213–21.

26. Schoeller DA, Ravussin E, Schutz Y, Acheson KJ, Baertschi P, Jéquier E. Energy
expenditure by doubly labeled water: validation in humans and proposed calculation. Am
J Physiol. 1986 May;250(5 Pt 2):R823-830.

27. R Core Team. R: A language and environment for statistical computing. 2022.

28. Pinheiro J, Bates D, DebRoy S, Deepayan S, Heisterkamp S, Van Willigen B, et al. nlme:
Linear and Nonlinear Mixed Effects Models [Internet]. 2022 [cited 2022 May 16].
Available from: https://CRAN.R-project.org/package=nlme

29. Revelle W. psych: Procedures for Psychological, Psychometric, and Personality Research
[Internet]. 2022 [cited 2022 May 16]. Available from: https://CRAN.R-
project.org/package=psych

30. Müller MJ, Geisler C, Hübers M, Pourhassan M, Braun W, Bosy-Westphal A.
Normalizing resting energy expenditure across the life course in humans: challenges and
hopes. Eur J Clin Nutr. 2018 May;72(5):628–37.

31. Giorgio A, Santelli L, Tomassini V, Bosnell R, Smith S, Stefano ND, et al. Age-related
changes in grey and white matter structure throughout adulthood. Neuroimage. 2010 Jul
7;51(3–2):943.

32. Lauretani F, Bandinelli S, Bartali B, Iorio AD, Giacomini V, Corsi AM, et al. Axonal
degeneration affects muscle density in older men and women. Neurobiology of Aging.
2006 Aug 1;27(8):1145–54.

33. Roubenoff R, Hughes VA, Dallal GE, Nelson ME, Morganti C, Kehayias JJ, et al. The
effect of gender and body composition method on the apparent decline in lean mass-
adjusted resting metabolic rate with age. J Gerontol A Biol Sci Med Sci. 2000
Dec;55(12):M757-760.

34. Heymsfield SB, Smith B, Dahle J, Kennedy S, Fearnbach N, Thomas DM, et al. Resting
Energy Expenditure: From Cellular to Whole-Body Level, a Mechanistic Historical
Perspective. Obesity (Silver Spring). 2021 Mar;29(3):500–11.

35. Joosen AMCP, Gielen M, Vlietinck R, Westerterp KR. Genetic analysis of physical
activity in twins. Am J Clin Nutr. 2005 Dec;82(6):1253–9.

36. Melanson EL, Gavin KM, Shea KL, Wolfe P, Wierman ME, Schwartz RS, et al.
Regulation of energy expenditure by estradiol in premenopausal women. J Appl Physiol
(1985). 2015 Nov 1;119(9):975–81.

37. Gavin KM, Melanson EL, Hildreth KL, Gibbons E, Bessesen DH, Kohrt WM. A
Randomized Controlled Trial of Ovarian Suppression in Premenopausal Women: No
Change in Free-Living Energy Expenditure. Obesity (Silver Spring). 2020
Nov;28(11):2125–33.

38. Tooze JA, Schoeller DA, Subar AF, Kipnis V, Schatzkin A, Troiano RP. Total daily
energy expenditure among middle-aged men and women: the OPEN Study. Am J Clin
Nutr. 2007 Aug;86(2):382–7.

39. Hawkes K, O'Connell JF, Blurton Jones NG. Hadza Women's Time Allocation, Offspring Provisioning, and the Evolution of Long Postmenopausal Life Spans. Current Anthropology. 1997;38(4):551–77.

40. Lieberman DE, Kistner TM, Richard D, Lee IM, Baggish AL. The active grandparent hypothesis: Physical activity and the evolution of extended human healthspans and lifespans. Proc Natl Acad Sci U S A. 2021 Dec 14;118(50):e2107621118.

41. Carey VJ, Walters EE, Colditz GA, Solomon CG, Willett WC, Rosner BA, et al. Body fat distribution and risk of non-insulin-dependent diabetes mellitus in women. The Nurses' Health Study. Am J Epidemiol. 1997 Apr 1;145(7):614–9.

42. Timpson NJ, Nordestgaard BG, Harbord RM, Zacho J, Frayling TM, Tybjærg-Hansen A, et al. C-reactive protein levels and body mass index: elucidating direction of causation through reciprocal Mendelian randomization. Int J Obes (Lond). 2011 Feb;35(2):300–8.

43. Scherrer U, Randin D, Tappy L, Vollenweider P, Jéquier E, Nicod P. Body fat and sympathetic nerve activity in healthy subjects. Circulation. 1994 Jun;89(6):2634–40.

44. Geisler C, Braun W, Pourhassan M, Schweitzer L, Glüer CC, Bosy-Westphal A, et al. Age-Dependent Changes in Resting Energy Expenditure (REE): Insights from Detailed Body Composition Analysis in Normal and Overweight Healthy Caucasians. Nutrients. 2016 Jun 1;8(6):E322.

45. Bosy-Westphal A, Wolf A, Bührens F, Hitze B, Czech N, Mönig H, et al. Familial influences and obesity-associated metabolic risk factors contribute to the variation in resting energy expenditure: the Kiel Obesity Prevention Study. Am J Clin Nutr. 2008 Jun;87(6):1695–701.

46. Calton EK, Pathak K, Soares MJ, Alfonso H, Keane KN, Newsholme P, et al. Vitamin D status and insulin sensitivity are novel predictors of resting metabolic rate: a cross-sectional analysis in Australian adults. Eur J Nutr. 2016 Sep;55(6):2075–80.

47. Soares MJ, Zhao Y, Calton EK, Pathak K. Triglycerides and systolic blood pressure negatively mediate the direct relationship of vitamin D status to resting energy expenditure: A cross sectional analysis. Diabetes Metab Syndr. 2022 Nov 9;16(12):102664.

48. Maas AHEM, Rosano G, Cifkova R, Chieffo A, van Dijken D, Hamoda H, et al. Cardiovascular health after menopause transition, pregnancy disorders, and other gynaecologic conditions: a consensus document from European cardiologists, gynaecologists, and endocrinologists. Eur Heart J. 2021 Mar 7;42(10):967–84.

49. Juppi HK, Sipilä S, Cronin NJ, Karvinen S, Karppinen JE, Tammelin TH, et al. Role of Menopausal Transition and Physical Activity in Loss of Lean and Muscle Mass: A Follow-Up Study in Middle-Aged Finnish Women. J Clin Med. 2020 May 23;9(5):E1588.

50. Elia M. Organ and Tissue Contribution to Metabolic Rate. In: Energy Metabolism: Tissue Determinants and Cellular Corollaries. New York: Raven Press; 1992. p. 61–79.

509 **TABLE 1.** Participant characteristics and energy expenditures according to age and menopause groups

	Age groups ($N = 120$)			Menopause groups ($n = 59$)		
	I	**II**	**III**	**PRE/PERI**	**MHT**	**POST**
Variable	**$n = 26$**	**$n = 35$**	**$n = 59$**	**$n = 19$**	**$n = 10$**	**$n = 30$**
Age, y	19.8 (1.1)	28.2 (4.3)	53.8 (3.5)	50.7 (4.3)	55.1 (1.7)	55.3 (1.9)
Sex hormone concentrations						
E2, nmol/l	0.10 (0.03–0.20) [#]	0.17 (0.06–0.41)	0.14 (0.08–0.33)	0.29 (0.18–0.55)	0.29 (0.17–0.38)	0.09 (0.06–0.12)
FSH, IU/l				8 (7–25)	38 (34–61)	80 (71–102)
Anthropometrics						
Height, cm	165.6 (5.6)	165.3 (5.8)	166.4 (5.0)	168.1 (4.6)	166.5 (5.6)	165.3 (4.8)
Body mass, kg	65.0 (10.5)	64.5 (7.9)	68.5 (9.1)	69.1 (9.0)	71.3 (12.3)	67.1 (7.9)
BMI, kg/m^2	23.7 (3.5)	23.6 (2.0)	24.7 (3.1)	24.5 (3.2)	25.7 (4.2)	24.5 (2.7)
Fat-free mass, kg	44.2 (5.0)	49.8 (5.3)	44.2 (4.2)	45.6 (4.1)	43.4 (4.2)	43.6 (4.1)
Fat mass, kg	20.9 (8.2)	14.7 (5.8)	24.2 (7.3)	23.6 (7.8)	27.9 (8.4)	23.4 (6.4)
Appendicular lean mass, kg	18.5 (2.6)	21.8 (2.7)	17.9 (1.9)	18.4 (2.0)	17.6 (1.9)	17.6 (1.9)
Appendicular lean mass index	6.7 (0.7)	8.0 (0.8)	6.5 (0.6)	6.5 (0.5)	6.4 (0.6)	6.4 (0.6)
Body fat percentage, %	31 (8)	23 (7)	36 (7)	35 (8)	40 (5)	36 (6)
Resting energy expenditure						
Measured, kcal/d	1,430 (110)	1,474 (146)	1,286 (124)	1,323 (105)	1,249 (143)	1,276 (128)
Predicted$_{FFM \& FM}$, kcal/d	1,339 (104)	1,442 (107)	1,345 (87)	1,372 (84)	1,335 (98)	1,332 (85)
Predicted$_{FFM, FM \& age}$, kcal/d	1,440 (103)	1,467 (94)	1,286 (92)	1,322 (86)	1,283 (115)	1,264 (84)
Total energy expenditure	**$n = 10$**		**$n = 10$**			
Measured, kcal/d	2,162 (310)		2,148 (236)			

Data as means (SD) or medians (IQR), [#] $n = 19$

E2, estradiol; FSH, follicle-stimulating hormone; MHT, menopausal hormone therapy; PRE/PERI, pre- or perimenopause; POST, postmenopause

511 **TABLE 2.** The associations between body composition, age, menopausal status, serum sex hormone concentrations, and resting energy
512 expenditure in midlife ($n = 59$)

	Mass and age		Menopause status		E2		FSH	
	B	*P*-value	*B*	*P*-value	*B*	*P*-value	*B*	*P*-value
Intercept	850.1	< 0.001	976.3	< 0.001	838.6	< 0.001	881.4	< 0.001
Fat-free mass, kg	17.2	< 0.001	17.2	< 0.001	17.2	< 0.001	17.4	< 0.001
Fat mass, kg	6.6	< 0.001	7.3	< 0.001	6.7	< 0.001	6.8	< 0.001
Age, y	-9.0	0.001	-11.2	< 0.001	-8.9	0.005	-10.1	0.001
Menopause status								
PRE/PERI			-39.7	0.11				
MHT			-58.4	0.025				
E2, nmol/l					5.2	0.89		
FSH, IU/l							0.2	0.43
R^2/adjusted R^2	0.70/0.68		0.73/0.71		0.70/0.68		0.70/0.68	

E2, estradiol; FSH, follicle-stimulating hormone; MHT, menopausal hormone therapy; PRE/PERI, pre- or perimenopause;
POST, postmenopause

514 **FIGURE 1**. Description of the study participants, outcomes, and statistical approaches.

521 **FIGURE 2.** The relationship between measured REE and predicted REE among the 120 participants. (A) REE predicted with fat-free mass
522 (FFM) and fat mass (FM), and (B) also with age.

FIGURE 3. The association of age and fat-free mass and fat mass–adjusted resting energy expenditure (REE$_{adj}$) and total energy expenditure (TEE$_{adj}$). (A) REE$_{adj}$ across age groups, (B) REE$_{adj}$ across menopause groups, (C) REE$_{adj}$ in 16 mother–daughter pairs, (D) TEE$_{adj}$ in 10 mother–daughter pairs. The boxplots show the mean and SD of each group, with whiskers representing $1.5 \times$ IQR.