Identification of a Non-Canonical Ciliate Nuclear Genetic Code Where UAA and UAG Code for Different Amino Acids

Authors:

Jamie McGowan[1], Estelle S. Kilias[2], Elisabet Alacid[2], James Lipscombe[1], Benjamin H. Jenkins[2], Karim Gharbi[1], Gemy G. Kaithakottil[1], Iain C. Macaulay[1], Seanna McTaggart[1], Sally D. Warring[1], Thomas A. Richards[2], Neil Hall[1,3], David Swarbreck[1]

Affiliations:

[1] Earlham Institute, Norwich Research Park, Norwich, NR4 7UZ, UK

[2] Department of Biology, University of Oxford, Oxford, OX1 3SZ, UK

[3] School of Biological Sciences, University of East Anglia, Norwich, UK

Correspondence:

thomas.richards@biology.ox.ac.uk, Neil.Hall@earlham.ac.uk,

David.Swarbreck@earlham.ac.uk

Keywords: Genetic code, stop codon reassignment, Ciliophora, Oligohymenophorea, suppressor tRNA, phylogenomics

Abstract

The genetic code is one of the most highly conserved features across life. Only a few lineages have deviated from the "universal" genetic code. Amongst the few variants of the genetic code reported to date, the codons UAA and UAG virtually always have the same translation, suggesting that their evolution is coupled. Here, we report the genome and transcriptome sequencing of a ciliate, belonging to the Oligohymenophorea class, where the translation of the UAA and UAG stop codons have changed to specify different amino acids. Genomic and transcriptomic analyses revealed that UAA has been reassigned to encode lysine, while UAG has been reassigned to encode glutamic acid. We identified multiple suppressor tRNA genes with anticodons complementary to the reassigned codons. We show that the retained UGA stop codon is enriched in the 3'UTR immediately downstream of the coding region of genes, suggesting that there is functional drive to maintain tandem stop codons. Using a phylogenomics approach, we reconstructed the ciliate phylogeny and mapped genetic code changes, highlighting the remarkable number of independent genetic code changes within the Ciliophora group of protists. According to our knowledge, this is the first report of a genetic code variant where UAA and UAG encode different amino acids.

Introduction

The genetic code is one of the most conserved features across life, emerging before

the last universal common ancestor (Knight et al. 2001). Virtually all organisms use the

canonical genetic code which has three stop codons (UAA, UAG, and UGA) and 61 sense

codons that specify one of 20 amino acids, including a translation start codon (AUG).

Variants of the genetic code, while rare, have been reported in several lineages of bacteria,

viruses, and eukaryotic organellar and nuclear genomes (Ivanova et al. 2014; Keeling 2016).

Ciliate nuclear genomes are a particular hotspot for genetic code variation. The phylum

Ciliophora is a large group of single-celled eukaryotes (protists) that diverged from other

Alveolates more than one billion years ago (Parfrey et al. 2011). Ciliates are highly unusual

in that they exhibit nuclear dimorphism whereby each cell has two types of nuclei, the

germline micronucleus (MIC) and the somatic macronucleus (MAC), each of which contains

its own distinct genome structure and function (Prescott 1994). The MIC genome functions as

the germline genome and is exchanged during sexual reproduction. MIC genomes are

typically diploid and are transcriptionally inactive during vegetative growth. The MIC

genome undergoes rearrangement and excision of non-coding sequences to serve as a

template to generate the transcriptionally active MAC genome (Arnaiz et al. 2012). MAC

genomes typically contain short, fragmented, gene-dense chromosomes that are present at

high ploidy levels (up to tens of thousands of copies) (Swart et al. 2013).

Known genetic code changes in ciliates involve reassignment of one or more stop

codons to specify for amino acids. Most reported ciliate genetic code changes involve

reassignment of both the UAA and UAG codons to specify glutamine as in *Tetrahymena*,

Paramecium, and *Oxytricha* (Lozupone et al. 2001), or glutamic acid in *Campanella*

umbellaria and *Carchesium polypinum* or tyrosine in *Mesodinium* species (Heaphy et al.

2016). Other known modifications include reassignment of the UGA stop codon to specify

tryptophan in *Blepharisma* (Lozupone et al. 2001), or cysteine in *Euplotes* (Meyer et al.

1991). The most extreme example of genetic code remodelling is found in *Condylostoma*

magnum where all three UAA, UAG, and UGA stop codons have been reassigned and can

specify either an amino acid (glutamine for UAA and UAG, and tryptophan for UGA) or

signal translation termination depending on their proximity to the mRNA 3' end (Swart et al.

2016; Heaphy et al. 2016). Not all ciliates use non-canonical genetic codes. For example,

Fabrea salina, *Litonotus pictus*, and *Stentor coeruleus* use the canonical genetic code

(Heaphy et al. 2016; Slabodnick et al. 2017).

Changing the meaning of codons from stop to sense requires modifications to the

translational apparatus. In eukaryotes, the eukaryotic release factor 1 (eRF1) protein

recognises the three standard stop codons in mRNA and triggers translation termination.

Studies have shown that mutations in the N-terminus of eRF1 can alter stop codon specificity

(Kryuchkova et al. 2013; Lekomtsev et al. 2007; Eliseev et al. 2011; Lozupone et al. 2001).

eRF1 specificity to recognise only the UGA codon has evolved independently via different

molecular mechanisms at least twice in ciliates with reassigned UAA and UAG codons

(Lekomtsev et al. 2007). Acquisition of tRNA genes with anticodons that recognise canonical

stop codons (suppressor tRNAs), via mutations, base modifications or RNA editing enables

translation of canonical stop codons into amino acids (Hanyu et al. 1986; Koonin &

Novozhilov 2009).

Tandem stop codons are additional stop codons located in the 3'-UTR within a few

positions downstream of a gene in the same reading-frame (Liang et al. 2005). They are

thought to act as "back-up" stop codons in the event of readthrough, minimising the extent of

erroneous protein elongation. For example, in yeast there is a statistical excess of stop codons

in the third in-frame codon position downstream of genes with a UAA stop codon (Liang et

al. 2005). Tandem stop codons have been shown to be overrepresented in ciliates that only

use UGA as a stop codon, compared to eukaryotes that use the canonical genetic code

(Fleming & Cavalcanti 2019). The level of overrepresentation is greater in highly expressed

genes (Adachi & Cavalcanti 2009). Tandem stop codons are thought to be particularly

important in ciliates where, following stop codon reassignment, readthrough events might

occur at a higher frequency due to mutations in eRF1 (Adachi & Cavalcanti 2009).

Several models have been proposed to describe genetic code changes. Under the

"codon capture" model, a codon that is rarely used (e.g., due to GC content) is gradually

eliminated from the genome followed by loss of the corresponding unused tRNA (Osawa et

al. 1992). Due to random genetic drift the codon could reappear and be captured by a

noncognate tRNA charged with a different amino acid, thus changing the genetic code. Such

a process would be essentially neutral, not resulting in mistranslated protein products as the

codon is eliminated from genes before the change in meaning occurs (Koonin & Novozhilov

2009). Alternatively, under the "ambiguous intermediate" model (Schultz & Yarus 1994),

reassignment of a codon takes place via an intermediate stage, where a codon is ambiguously

translated via competing tRNAs charged with different amino acids, or in the context of stop

codon reassignment, a suppressor tRNA competing with a release factor. This process would

be driven by selection and result in the elimination of the cognate tRNA if the new meaning

is advantageous. The "genome streamlining" model is more relevant to small genomes (e.g.,

organellar genomes or parasites) where there is pressure to minimise translational machinery

(Andersson & Kurland 1995). More recently the "tRNA loss driven codon reassignment"

mechanism was proposed to describe codon reassignments whereby tRNA loss, or alteration

of release factor specificity, results in an unassigned codon that can be captured by another

tRNA gene (Mühlhausen et al. 2016; Kollmar & Mühlhausen 2017).

In virtually all genetic code changes reported to date, the codons UAA and UAG have

the same meaning, i.e., they are either both used as canonical stop codons or are both

reassigned to the same amino acid (Kollmar & Mühlhausen 2017). This suggests that

evolutionary or mechanistic constraints couple the meaning of these two codons (Pánek et al.

2017). One such constraint is wobble binding of a suppressor tRNA gene with a UUA

anticodon, where uracil in the first anticodon position can bind to either adenine or guanine in

the third codon position of mRNA (Crick 1966). Thus, acquiring a suppressor tRNA gene

with a UUA anticodon could potentially change the meaning of both the UAA and UAG

codons. Wobble binding has been experimentally demonstrated in *Tetrahymena thermophila*,

where tRNA-Sup(UUA) was shown to suppress both the UAA and UAG codons, whereas

tRNA-Sup(CUA) suppressed only the UAG codon (Hanyu et al. 1986). The first report of

nuclear genetic code variants where UAA and UAG have different meanings were reported in

transcriptomics analyses where a Rhizarian species (Rhizaria sp. exLh) was shown to use

UAG to encode leucine and in a Fornicate (*Iotanema spirale*) where UAG has been

reassigned to glutamine (Pánek et al. 2017). However, in both cases, the UAA codon was

retained as a stop codon, thus avoiding the problem of genetic code ambiguity due to wobble

binding.

Here, we report the discovery of a novel variant of the genetic code in a ciliate

belonging to the Oligohymenophorea class, where the meaning of the UAA and UAG codons

have changed to specify different amino acids. Using G&T-Seq (Macaulay et al. 2015), we

performed parallel genome and transcriptome sequencing of small pools of ciliate cells.

Combining genome and transcriptome sequencing data from multiple independently

amplified samples enabled co-assembly of a highly complete macronuclear genome assembly

and annotation. Genomic and transcriptomic analysis revealed that the UAA codon has been

reassigned to specify lysine, while the meaning of the UAG codon has changed to specify

glutamic acid. We identified multiple suppressor tRNA genes of both types in the genome,

supporting the genetic code changes. We show that UGA codons are significantly enriched in

the 3'-UTR of genes suggesting that there is selective pressure to maintain tandem stop

codons, which may play a role in minimising erroneous protein elongation in the event of

translational readthrough. To our knowledge, this is the first report of a genetic code variant

where UAA and UAG specify different amino acids.

Results & Discussion

Genome Assembly of an Oligohymenophorean Ciliate

We isolated a ciliate species Oligohymenophorea sp. PL0344 from a freshwater pond at Oxford University Parks, Oxford, UK. Attempts to establish a stable long-term culture were unsuccessful so we applied low input single-cell based approaches to generate genomic and transcriptomic data. Small pools of cells (5 – 50 cells) were sorted into a microplate using fluorescence-activated cell sorting (FACS). Parallel genome and transcriptome sequencing was performed using G&T-Seq, which relies on whole genome amplification using multiple displacement amplification (MDA) and transcriptome analysis using a modified Smart-seq2 protocol (Macaulay et al. 2015).

A *de novo* genome assembly was generated by co-assembling reads from 10 samples (totalling approximately 6 Gb). Following removal of contaminant sequences, the resulting macronuclear genome assembly was 69.8 Mb in length, contained in 3671 scaffolds with an N50 of 59.6 Kb (**Table 1**). Approximately 89% of the corresponding RNA-Seq reads mapped to the genome assembly, indicating high completeness. GC content of the genome is low at 30.58% (**Table 1**), which is similar to previously sequenced ciliate genomes (Slabodnick et al. 2017). The mitochondrial genome was also recovered which is a linear molecule 35,635 bp in length with GC content of 25.33% and capped with repeats. The mitochondrial genome contains the small subunit (SSU) and large subunit (LSU) ribosomal RNA (rRNA) genes, 5 tRNA genes, 19 known protein-coding genes, and 13 open reading frames.

The nuclear encoded SSU rRNA gene sequence is 99.81% identical to an environmental sequence (AY821923) in the GenBank database, isolated from Orsay, France (Šlapeta et al. 2005). Maximum-likelihood phylogenetic analysis of the SSU rRNA gene placed it within a clade containing four unnamed ciliate species (AY821923, HQ219368, LR025746, HQ219418) and *Cinetochilum margaritaceum* (MW405094) with 100%

bootstrap support (**Supplementary Figure 1**). Thus, based on the SSU rRNA gene, *C.*

margaritaceum is the closest related named species. The SSU rRNA gene of *C.*

margaritaceum is 96.03% identical to that of Oligohymenophorea sp. PL0344. *C.*

margaritaceum belongs to the Loxocephalida order (Class Oligohymenophorea; Subclass

Scuticociliatia), which is considered a controversial order due to its non-monophyly

(Poláková et al. 2021; Gao et al. 2013). Our phylogenetic analysis places *C. margaritaceum*

as a divergent branch relative to other members of Loxocephalida (**Supplementary Figure**

1), which is congruent with previous analyses (Poláková et al. 2021; Gao et al. 2013),

suggesting taxonomic revision is required.

Oligohymenophorea sp. PL0344 Uses a Novel Genetic Code

Preliminary analysis of the genome sequence revealed that many coding regions

contained in-frame UAA and UAG codons. Consistent with codon reassignments in other

ciliate species, this suggested that the UAA and UAG stop codons have been reassigned to

code for amino acids. Surprisingly however, the meanings of these codons do not match any

known genetic code. An example gene (tubulin gamma chain protein), showing six in-frame

UAA codons and six in-frame UAG codons, translated and aligned to orthologous protein

sequences with representatives from across Eukaryota is displayed in **Figure 1**. Five in-frame

UAA codons correspond to highly conserved columns in the alignment where lysine is the

consensus amino acid (**Figure 1**). Four in-frame UAG codons correspond to highly

conserved columns in the alignment where glutamic acid is the consensus amino acid, and

another corresponds to a column with a mix of glutamic acid and aspartic acid (**Figure 1**).

We used two complementary tools to analyse the genetic code further. First, we used

the "genetic_code_examiner" utility from the PhyloFisher package (Tice et al. 2021), which

predicts the genetic code by comparing codon positions in query sequences to highly

conserved (> 70% conservation) positions in amino acid alignments from a database of 240

orthologous protein sequences. PhyloFisher identified 58 genes with 87 in-frame UAA

codons that correspond to highly conserved amino acid sites. Of these, 74 UAA codons

(85%) correspond to highly conserved lysine residues (**Figure 2A**). The second most

numerous match was to arginine, another positively charged amino acid, with 9 (10%) hits.

PhyloFisher identified 46 genes with 63 in-frame UAG codons that correspond to highly

conserved amino acid sites. Of these, 56 UAG codons (89%) correspond to highly conserved

glutamic acid residues (**Figure 2B**). The second most numerous match was to aspartic acid,

another negatively charged amino acid, with 4 (6%) of hits.

We also analysed the genetic code using Codetta (Shulgina & Eddy 2021). Codetta

predicts the genetic code by aligning profile hidden Markov models (HMMs) from the Pfam

database against a six-frame translation of the query genome assembly. The meaning of each

codon is inferred based on emission probabilities of the aligned HMM columns. From the

whole genome sequence, 14,633 UAA codons and 10,160 UAG codons had a Pfam position

aligned. Based on these alignments, Codetta also predicted that the UAA codon is translated

as lysine and UAG translated as glutamic acid, each with a log decoding probability of zero

(**Figure 2C**).

Thus, these results indicate that Oligohymenophorea sp. PL0344 uses a novel genetic

code where UAA is translated as lysine and UAG is translated as glutamic acid. This is the

first time this genetic code variant has been reported. Furthermore, according to our

knowledge, this is the first report of a genetic code variant where UAA and UAG have been

reassigned to specify different amino acids. Genetic code variants were previously reported

where UAG was reassigned to specify an amino acid (either leucine or glutamine) but UAA

was retained as a stop codon in both cases (Pánek et al. 2017). This is significant as it

suggests that the genetic code variant reported herein has overcome mechanistic constraints

linking the translation of these two codons.

Suppressor tRNA Genes

tRNA genes were annotated using tRNAscan (Chan et al. 2021), resulting in the

annotation of 320 tRNA genes, including 15 that are predicted to be pseudogenes. Amongst

the annotated tRNA genes are 23 putative suppressor tRNA genes. These are tRNA genes

with anticodons complementary to canonical stop codons (UAA, UAG, or UGA). The

annotated suppressor tRNA genes include 12 tRNA-Sup(UUA) genes and 10 tRNA-

Sup(CUA) genes. tRNAscan also predicted a single tRNA-Sup(UCA) gene, however this was

low scoring and was not predicted by ARAGORN (Laslett 2004), an alternative tool to

identify tRNA genes. tRNAscan also predicts the function of tRNAs. Many of the tRNAscan

isotype predictions were consistent with the predicted genetic code (i.e., UAA = lysine and

UAG = glutamic acid), however several putative tRNA genes had low-scoring or inconsistent

isotype predictions. To better characterise the suppressor tRNA genes, we compared their

sequences to the non-suppressor tRNA genes. Eight of the twelve predicted tRNA-Sup(UUA)

genes were most similar to tRNA-Lys genes with UUU or CUU anticodons (68.49% to

80.95% identical) (**Supplementary Table 1**), consistent with the genetic code prediction that

UAA has been reassigned to specify lysine. An example tRNA-Sup(UUA) predicted to

function as a lysine tRNA is shown in **Figure 3A**. All ten tRNA-Sup(CUA) genes were most

similar to tRNA-Glu genes with CUC or UUC anticodons (69.44% to 93.06% identical)

(**Supplementary Table 1**), consistent with the genetic code prediction that UAG has been

reassigned to specify glutamic acid. An example tRNA-Sup(CUA) predicted to function as a

glutamic acid tRNA is shown in **Figure 3B**. We also identified a tRNA gene for

selenocysteine, tRNA-SeC(UCA) (**Figure 3C**), suggesting that the UGA codon is used both

as a stop codon and to encode selenocysteine. Thus, all 64 codons can specify amino acids as

has been reported in other ciliate genomes (Eisen et al. 2006).

UAA and UAG codons differ only in the wobble position. According to wobble-

binding rules, uracil in the first tRNA anticodon position ("wobble position") (**Figure 3A**)

can bind to either adenine or guanine in the third codon position of mRNA (Crick 1966),

allowing tRNA with a UUA anticodon to recognise both UAA and UAG codons. It has been

experimentally demonstrated that *T. thermophila* tRNA-Sup(UUA) can recognise both UAA

and UAG codons (Hanyu et al. 1986). It has been suggested that wobble binding is a possible

explanation as to why UAA and UAG virtually always have the same meaning (Heaphy et al.

2016). Considering that Oligohymenophorea sp. PL0344 has tRNA-Sup(UUA) genes for

lysine and tRNA-Sup(CUA) genes for glutamic acid, this raises the question: are UAG

codons ambiguously translated as both glutamic acid and lysine? If not, how has it overcome

the mechanistic and evolutionary constraints that appear to couple the translation of these two

codons? Presumably, if wobble binding allows tRNA-Sup(UUA) to recognise the UAG

codon, it would be less efficient than tRNA-Sup(CUA) and outcompeted, possibly resulting

in some degree of stochastically translated protein products with glutamic acid residues

substituted by lysine at UAG codon positions. Attempts to establish a stable culture were

unsuccessful, and while we can overcome this problem to generate a genome assembly using

low-input sequencing methods designed for single-cell analysis, such low-input approaches

are not available for proteomics. Without proteomics data, it is not possible to determine if

UAG is ambiguously translated. Furthermore, from suppressor tRNA gene sequences alone,

it is not possible to determine if they incorporate modified nucleotides which could alter

codon-anticodon binding specificity.

Genome Annotation and Codon Usage Analysis

Genome annotation incorporating RNA-Seq data and protein alignments from other ciliates resulted in the annotation of 22,048 transcripts from 20,141 gene models (**Table 1**). BUSCO analysis estimates that the genome annotation is highly complete with 94.7% of BUSCO genes recovered as complete, which compares favourably to other high quality ciliate genomes (**Supplementary Table 2**). The median intron size of 57 bp (**Table 1**) is similar to previously sequenced ciliate genomes, such as *Tetrahymena thermophila* and *Oxytricha trifallax* (Eisen et al. 2006; Swart et al. 2013) but not as short as the extremely short introns (15 – 25 bp) found in *Stentor coeruleus* or *Paramecium tetraurelia* (Slabodnick et al. 2017; Aury et al. 2006). Lysine, leucine, and glutamic acid are the three most used amino acids at 10.67%, 9.06% and 8.62%, respectively (**Supplementary Table 3**). Codon usage is biased towards using codons with lower GC content. This bias is reduced in highly expressed genes which have higher GC content compared to all genes (38.51% versus 34.12%), similar to previous reports in *Paramecium* and *Tetrahymena* (Eisen et al. 2006; Salim et al. 2008). The reassigned UAA and UAG codons are the 7[th] and 13[th] most used codons at 3.25% and 2.10% respectively. Their usage is reduced in highly expressed genes to 1.87% (21[st] most used) and 0.98% (40[th] most used) respectively. Among lysine codons (UAA, AAA, and AAG), UAA usage is 30.42% across all genes, but reduced to 20.49% in highly expressed genes. For glutamic acid codons (UAG, GAA, and GAG), codon usage of UAG is 24.41% across all genes but is also reduced in highly expressed genes to 13.21%. 98.6% of genes have at least one UAA codon, which is reduced to 89.1% for highly expressed genes. For UAG, 96.4% of genes have at least one UAG codon compared to just 75.2% of highly expressed genes. Reduced codon usage in highly expressed genes could indicate translational inefficiency, or that selective pressure to retain canonical lysine and glutamic acid codons is higher in highly expressed genes.

We analysed tandem stop codons by counting UGA codons in the first 20 in-frame

codon positions downstream of genes. Our results show that UGA codons are significantly

overrepresented (chi-squared test, p-value < 0.05) in the first four in-frame codons

downstream of genes (**Figure 4**). 12.3% of genes have at least one UGA codon within the

first six in-frame codon positions downstream of genes, similar to the proportion reported for

T. thermophila (11.5%) where UAA and UAG have also been reassigned to encode amino

acids (Fleming & Cavalcanti 2019). For comparison, the reassigned UAA and UAG codons

are not overrepresented in this region. The frequency of UGA codons at these positions is

greater for highly expressed genes whereby 13.6% of highly expressed genes have at least

one UGA codon within the first six in-frame codon positions downstream of genes (**Figure**

4). These data add support that there is selective pressure for ciliates with reassigned UAA

and UAG codons to maintain tandem UGA stop codons at the beginning of the 3'-UTR. It is

tempting to speculate that these additional UGA stop codons play a role in minimising

deleterious consequences of readthrough events.

Phylogenomics Analysis of Genetic Code Changes in the Ciliophora

We carried out phylogenomics analyses to map genetic code changes onto the ciliate

phylogeny. A phylogenomic dataset consisting of genomic and transcriptomic data from 46

ciliate species and 9 outgroup species was constructed (**Supplementary Table 2**).

Phylogenomic reconstruction was performed on a concatenated alignment of 89 single-copy

BUSCO proteins (40,289 amino acid sites) using maximum-likelihood (IQ-TREE under

LG+F+I+R7 model) and Bayesian (PhyloBayes-MPI under CAT-GTR model) approaches.

We also conducted a partitioned analysis on the same dataset using IQ-TREE, with a

partitioning scheme which merged the 89 proteins into 14 partitions. The three resulting

phylogenies were largely in agreement with each other and with previously published

analyses, with full or high support from all three methods at most branches (**Figure 5**).

Oligohymenophorea sp. PL0344 was robustly placed within the Oligohymenophorea class in

a clade containing Hymenostomatida and *Pseudocohnilembus persalinus* with full support

from all three methods (**Figure 5**).

The position of *Paramecium* (order Peniculida) is unstable in our phylogenomic

analyses. Both the LG+F+I+R7 and partitioned phylogeny group *Paramecium* as sister to the

Peritrichia clade with 89% and 82% bootstrap support respectively (**Figure 5**). This is

congruent with some previous phylogenomic analyses which recover Peniculida as sister to

Peritrichia species (Feng et al. 2015; Jiang et al. 2019; Gentekaki et al. 2017). However, the

Bayesian phylogeny places *Paramecium* as sister to Hymenostomatida and Philasterida

(**Supplementary Figure 2**). This grouping has been recovered in some previous

phylogenomic analyses (Wang et al. 2021; Rotterová et al. 2020). The correct placement of

Peniculida is unclear based on the current datasets available. The *Paramecium* species

included in our analysis have a high proportion of missing data (**Figure 5**). We anticipate that

differences in topology may be influenced by varying levels of sensitivity to missing data in

the models used. *Mesodinium rubrum* is another problematic taxon which is thought to be

prone to long branch attraction (LBA) artefacts. Furthermore, existing *Mesodinium*

transcriptomes are contaminated with sequences from their prey (Lasek-Nesselquist &

Johnson 2019). Some previous phylogenomic and phylogenetic analyses place it as an early

branching ciliate (Lynn & Kolisko 2017; Chen et al. 2015), however these may have been

influenced by contamination (Lasek-Nesselquist & Johnson 2019). Here, we account for

contamination by removing any sequences from the *M. rubrum* transcriptome with best

BLAST hits outside of the Ciliophora (n = 3,574). Both the LG+F+I+R7 and partitioned

phylogeny group *M. rubrum* with *Litonotus pictus*, another member of the Litostomatea class,

with 85% and 93% bootstrap support respectively (**Figure 5**), while our Bayesian analysis

places it as a deep branching ciliate branching before *Protocruzia* (**Supplementary Figure**

2B). The grouping of *M. rubrum* with *L. pictus* agrees with a recent phylogenomics analysis

of *Mesodinium* species that accounts for LBA and contamination (Lasek-Nesselquist &

Johnson 2019).

Where genome or transcriptome assemblies were available, or raw sequencing reads

deposited in public databases, we validated the known genetic codes using Codetta and

PhyloFisher. All species had the expected genetic code except for *Plagiopyla frontata*.

Codetta and PhyloFisher both predicted that UAA and UAG are translated as glutamine in *P.*

frontata, which is not surprising given how many ciliate species use this genetic code (**Figure**

5). Interestingly however, both methods predict that UGA has also been reassigned to specify

tryptophan in *P. frontata*. From the PhyloFisher dataset of 240 query proteins, 3 (1.25%)

contain internal UGA codons that correspond to highly conserved tryptophan residues in

other species **(Supplementary Figure 3).** This suggests that *P. frontata* may use UGA both

as a stop codon and also rarely as a sense codon to specify tryptophan, similar to the

Condylostoma genetic code (**Figure 5**) (Swart et al. 2016; Heaphy et al. 2016).

We mapped genetic code reassignments onto the ciliate phylogeny, highlighting the

remarkable number of independent genetic code changes within the ciliates (**Figure 5**). Based

on our phylogeny, and assuming a non-canonical genetic code doesn't revert to the canonical

genetic code, the translation of UAR (UAA and UAG) codons to glutamine is the most

common genetic code variant and has independently evolved at least five times. From our

analysis, translation of UGA to tryptophan has independently evolved at least three times in

ciliate nuclear genomes. However, it has recently been reported that Karyorelict ciliates (not

included in this analysis) use a context-dependent genetic code similar to *Condylostoma*,

where UAR has been reassigned to glutamine and UGA specifies either tryptophan or stop

depending on context, indicating a fourth independent origin of UGA being translated as

tryptophan and a sixth independent origin of UAR being translated to glutamine in ciliates

(Seah et al. 2022). The translation of UGA to cysteine in *Euplotes*, UAR to tyrosine in

Mesodinium and UAR to glutamic acid in Peritrichia have all evolved once. The

Oligohymenophorea sp. PL0344 genetic code appears to be a relatively recent phenomenon

and is unique in that the two codons have different meanings. The Oligohymenophorea class

contains at least three different genetic code variants, and no sampled species which have

retained usage of UAA or UAG as a stop codon. Our phylogeny suggests that the stop codons

UAA and UAG were reassigned to glutamine in the ancestor of Oligohymenophorea (**Figure**

5). These codons were then reassigned to glutamic acid in the Peritrichs, or to lysine (UAA)

and glutamic acid (UAG) in Oligohymenophorea sp. PL0344.

It remains unclear why Ciliate genomes are such a hotspot for stop codon

reassignments. Our study shows that even within the Oligohymenophorea class, which is

relatively well sampled compared to other ciliate clades, there remain novel genetic code

reassignments to be discovered. Further sequencing of under-sampled ciliate lineages and

other microbial eukaryotes may reveal more variant genetic code changes and help to better

understand the evolution and mechanisms of genetic code changes.

Materials and methods

Sampling, Ciliate isolation, Culturing, and Cell-sorting

Surface water was collected from a margin of an artificial freshwater pond at Oxford University Parks (51°45'51.0"N 1°15'04.5"W), Oxford (UK) by directly submerging a 1L-autoclaved glass collection bottle. 200 mL of the water sample were concentrated using a 5-um filter into a final volume of 20 mL. Oligohymenophorea sp. PL0344 was identified using an inverted microscope (Olympus CKX41) and single cells were isolated manually using a glass micropipette by transferring them into successive drops of 0.2 μm pre-filtered and autoclaved environmental source water. When cells were free of any other contaminant, they were transferred into a 96 well-plate containing filtered and autoclaved environmental source water. In order to obtain a clonal culture, isolated cells were incubated during a week at 20°C with a 12h:12h light:dark photo-cycle with a photon flux of 32 μmoles·m^{-2}·s^{-1}. When ciliate cells divided and a dense culture was observed in the well, the mini-culture was scaled-up during a month by successively transferring the cells into larger volumes until a non-axenic dense ciliate culture of 20 mL of volume was established. Pools of ciliate cells (5 – 50 cells) were then sorted into a 384-well plate containing 5 μL of autoclaved source water using FACS (BD FACSMelody™ Cell Sorter, BD Biosciences). 10uL of RLT+ lysis buffer (Qiagen) was then added to each well and the plate was sealed and centrifuged (2000xg, 4ºC, 1min) to remove bubbles and to ensure that the lysis buffer was at the bottom of each well. The sorted plate was stored at -80ºC until processed.

G&T-Seq, Library Preparation, and Sequencing

Using a magnetic separator, Dynabeads™ MyOne™ Streptavidin C1 (Invitrogen) beads were washed according to the manufacturer's guidance and then incubated with 2 × Binding &Wash buffer (10mM Tris-HCl pH7.5, 1mM EDTA, 2M NaCl) and Biotynilated Oligo-dT

primer (IDT, 5'-/BiotinTEG/AAG CAG TGG TAT CAA CGC AGA GTA CTT TTT TTT

TTT TTT TTT TTT TTT TTT TTT TTT TTT TVN-3') at 100μM for 30 minutes at room temperature

on a rotator. The oligo-treated beads were washed four times in $1 \times$ Binding &Wash buffer

(5mM Tris-HCl pH7.5, 0.5mM EDTA, 1M NaCl) and then suspended in $1 \times$ SuperScript II

First Strand Buffer (Invitrogen) supplemented with SUPERase•In™ RNase Inhibitor

(Invitrogen) to a final concentration of 1U/μl. The lysate was thawed on ice. 10μl of prepared

oligo-dT beads was added to each well containing 12μl cell lysate. The lysate plate was

sealed and incubated on a ThermoMixer C (Eppendorf) at 21°C for 20 minutes shaking at

1000rpm. Using a Fluent 480 liquid handling robot (Tecan) and a Magnum FLX magnetic

separator (Alpaqua), the lysate supernatant was transferred to a new plate and the beads were

washed twice in a custom wash buffer (50mM Tris-HCl pH8.3, 75mM KCl, 3mM MgCl2,

10mM DTT, 0.5% Tween-20). The supernatant from the washes was added to the left-over

cell lysate - containing the genomic DNA - which was stored at -20°C overnight. The washed

beads were suspended in a reverse transcription mastermix of 1mM dNTPs, 0.01M MgCl2,1

$\times$ SuperScript II First Strand Buffer, 1M Betaine, 5.4M DTT, 1μM Template-Switching

Oligo (5'-AAGCAGTGGTATCAACGCAGAGTACrGrG+G-3', where "r" indicates a

ribonucleic acid base and "+" indicates a locked nucleic acid base, Qiagen) then incubated

using a ThermoMixer C with the following conditions: 42°C for 2 minutes at 200rpm, 42°C

for 60 minutes at 1500rpm, 50°C for 30 minutes at 1500rpm, 60°C for 10 minutes at

1500rpm. The cDNA was amplified using HiFi Hotstart Ready Mix (KAPA) and IS Primers

to a final concentration of 0.1μM (IDT, 5'-AAG CAG TGG TAT CAA AGA GT-3') with the

following thermocycling conditions: 98°C for 3 minutes, then 21 cycles of 98°C for 15

seconds, 67°C for 20 seconds, 72°C for 6 minutes and finally 72°C for 5 minutes. The cDNA

was then purified using $0.8 \times$ vols Ampure XP (Beckman Coulter) and 80% ethanol on the

Fluent 480 liquid handling robot and eluted in 10mM Tris-HCl. The remaining cell lysate

was thawed and subjected to a 0.6 × vols Ampure XP clean-up with 80% ethanol. The bead-

bound gDNA was isothermally amplified for 3 hours at 30°C then 10 minutes at 65°C using a

miniaturised (1/5 vols) Repli-g Single-Cell assay (Qiagen). The amplified gDNA was cleaned

up with 0.8 × vols Ampure XP and 80% ethanol, then eluted in 10mM Tris-HCl. The cDNA

and gDNA were quantified by fluorescence (Quant iT HS-DNA, Invitrogen) on an Infinite

Pro 200 plate reader (Tecan) then normalised to a final concentration of 0.2ng/µl in 10mM

Tris-HCl. Dual-indexed sequencing libraries (Nextera XT, Illumina) were prepared using

Mosquito and Dragonfly liquid handling instruments (SPT Labtech). The libraries were

pooled and cleaned up using 0.8× vols Ampure XP and 80% ethanol. The libraries were

eluted in 10mM Tris-HCl and assessed using a Bioanalyzer HS DNA assay (Agilent), HS

DNA Qubit assay (Invitrogen) and finally an Illumina Library Quantification Kit assay

(KAPA). Sequencing was conducted on a NovaSeq 6000 with a 300 cycle Reagent kit v1.5

(Illumina) to produce 150bp paired-end, dual-indexed reads.

Genome Assembly

Adapter and quality trimming were carried out using BBDuk ([https://jgi.doe.gov/data-](https://jgi.doe.gov/data-and-tools/bbtools)

[and-tools/bbtools](https://jgi.doe.gov/data-and-tools/bbtools)). Reads which mapped to a database of common lab contaminants (human

and mouse) were removed using BBMap. A co-assembly of genomic DNA reads from 10

samples was generated using SPAdes (v3.15.3) (Bankevich et al. 2012) with default settings

except -k 21, 33, 55, 77 and single-cell mode (--sc) was enabled. Contaminant contigs were

removed using a combination of metagenomic binning with MetaBAT2 (Kang et al. 2019)

based on tetra-nucleotide frequencies and taxonomic classification with CAT (v5.2) (Von

Meijenfeldt et al. 2019) and Tiara (v1.0.1) (Karlicki et al. 2022). Assembly statistics were

calculated using Quast (Gurevich et al. 2013). Genome completeness was assessed using

BUSCO (v4.1.2) (Manni et al. 2021) with the Alveolata_obd10 dataset.

Genome Annotation

The genetic code was predicted using Codetta (v2.0) (Shulgina & Eddy 2021) and also using the "Genetic Code Examiner" utility from the PhyloFisher package, with the included database of 240 orthologs (Tice et al. 2021).

Gene models were annotated via the Robust and Extendable eukaryotic Annotation Toolkit (REAT, https://github.com/EI-CoreBioinformatics/reat) and Minos (https://github.com/EI-CoreBioinformatics/minos) using a workflow incorporating repeat identification, RNA-Seq mapping / assembly, alignment of protein sequences from related species and evidence guided gene prediction with AUGUSTUS (Stanke et al., 2006).

A de novo repeat annotation was created using the RepeatModeller (Hubley et al. 2008) v1.0.11 -RepeatMasker v4.07 (Hubley et al. 2008) pipeline with defaults settings and the --gff output option enabled. To ensure high copy number 'bonafide' genes were excluded from repeat masking, the RepeatModeler library was hard masked using protein coding genes from 11 ciliate species (detailed below). The protein coding genes were first filtered to remove any genes with descriptions indicating "transposon" or "helicase". TransposonPSI (r08222010) http://transposonpsi.sourceforge.net was then run to remove any transposon hits by hard-masking them and using the filtered gene set to mask the RepeatModeler library. RepeatMasker v4.0.7 was run with the Repbase Alveolata library (RepBaseRepeatMaskerEdition-20170127.tar.gz) and additionally with the filtered RepeatModeler library. The interspersed repeats were combined and used as evidence in the gene build.

The REAT transcriptome workflow was run with RNA-Seq (total 77 million read pairs) from 28 samples. As transcriptome assembly is sensitive to depth of RNA-Seq coverage samples were combined into sets of 28, 10, 10 and 8 samples to ensure reasonable coverage but also allow alternative assemblies to be created. Illumina RNA-seq reads were

mapped to the genome with HISAT2 v2.2.1 (Kim et al., 2019) and high-confidence splice

junctions identified by Portcullis (Mapleson et al., 2018). The aligned reads were assembled

for each set of samples with StringTie2 v2.1.5 (Kovaka et al., 2019) and Scallop v0.10.5

(Shao and Kingsford, 2017). From the combined set of RNA-Seq assemblies a filtered set of

non-redundant gene-models were derived using Mikado (Venturini et al., 2018). The REAT

homology workflow was used to generate gene models based on alignment of proteins from

11 ciliate species (**Supplementary Table 2**). These together with the transcriptome derived

models were used to train the AUGUSTUS v3.4.0 gene predictor, with transcript and protein

alignments plus repeat annotation provided as hints in evidence guided gene prediction using

the REAT prediction workflow. Six alternative AUGUSTUS gene builds were generated

using different evidence inputs or weightings for the protein, transcriptome and repeat

annotation. The Minos pipeline was run to generate a consolidated set of gene models from

the transcriptome, homology, projected and AUGUSTUS predictions. The pipeline utilises

external metrics to assess how well supported each gene model is by available evidence,

based on these and intrinsic characteristics of the gene models a final set of models is

selected. For each gene model a confidence and biotype classification were determined based

on the type and extent of supporting data.

Annotation completeness was assessed using BUSCO (v4.1.2) (Manni et al. 2021)

with the Alveolata_obd10 dataset. tRNA genes were annotated using tRNAscan (v2.0.7)

(Chan et al. 2021). rRNA genes were annotated using barrnap (v0.9)

(https://github.com/tseemann/barrnap).

Tandem Stop Codon Analysis

To investigate if UGA stop codons are enriched in the 3'-UTR of genes, codon usage

of the first 20 in-frame codons downstream of each gene's stop codon was calculated.

Expected frequencies were determined by counting codons in all six reading frames in the 60

bp region downstream of each gene's stop codon. We also carried out this analysis for highly

expressed genes which we defined as the 10% of genes with the highest transcripts per

million (TPM) values, calculated using Kallisto (Bray et al. 2016). Statistical significance

was assessed using the chi-squared test.

Phylogenetic Analysis of SSU rRNA Genes

Small subunit ribosomal RNA sequences from related species were retrieved from

GenBank (**Supplementary Figure 1**). Sequences were aligned using MAFFT (v7.490) with

the G-INS-I algorithm (Katoh & Standley 2013). Maximum-likelihood phylogenetic analysis

was performed using IQ-TREE (v2.2.0) (Minh et al. 2020) under the GTR+F+R5 model,

which was the best fit model according to ModelFinder (Kalyaanamoorthy et al. 2017), with

100 non-parametric bootstrap replicates.

Phylogenomic Analyses

A phylogenomic dataset of 55 species was assembled including previously published

ciliate genomes and transcriptomes with outgroup species from the Alveolata, retrieved from

databases and published phylogenomics analyses (Irwin et al. 2021; Richter et al. 2022)

(**Supplementary Table 2**). *De novo* transcriptome assemblies were generated for two species

– *Campanella umbellaria* and *Carchesium polypinum*. RNA-Seq reads were retrieved from

the sequence read archive (SRR1768423 and SRR1768437) (Feng et al. 2015).

Transcriptome assemblies were generated using Trinity (Grabherr et al. 2011), redundancy

was reduced using CD-HIT-EST (Fu et al. 2012) with an identity cut-off of 98% and protein

coding transcripts were predicted using Transdecoder (Haas et al. 2013). Coding sequences

were translated into amino acids using the correct genetic code (UAR = E). The

transcriptome assembly of *Mesodinium rubrum* is contaminated with sequences from its prey.

We excluded any *M. rubrum* proteins with a best BLAST hit outside of the Ciliophora to

account for this contamination which resulted in the removal of 3,574 (22%) proteins.

BUSCO analysis using the Alveolata_obd10 dataset identified 89 proteins that are

present and single copy in at least 65% of species, i.e., at least 36 out of 55 species. Each

BUSCO family was individually aligned using MAFFT (v7.490) (Katoh & Standley 2013)

and then trimmed using trimAl (v1.4) with the "gappyout" parameter (Capella-Gutierrez et al.

2009). The trimmed alignments were concatenated together resulting in a supermatrix

alignment of 40,289 amino acid sites. Maximum-likelihood phylogenetic reconstruction was

performed using IQ-TREE (v2.2.0) (Minh et al. 2020) under the LG+F+I+R7 model, which

was the best fitting model according to ModelFinder (Kalyaanamoorthy et al. 2017), and 100

non-parametric bootstrap replicates were used to assess branch support. We also conducted a

partitioned analysis using IQ-TREE (Chernomor et. al 2016) with a partitioning scheme that

merged the 89 proteins into 14 partitions with model selection performed by ModelFinder,

with 100 non-parametric bootstrap replicates. Bayesian analyses were also performed on the

supermatrix alignment using PhyloBayes-MPI (v1.8c) (Lartillot et al. 2013) under the CAT-

GTR model. Constant sites (n = 3,299) were removed. Two independent Markov chain

Monte Carlo (MCMC) chains were run for approximately 12,000 generations. Convergence

was assessed using bpcomp and tracecomp with a burn-in of 20%.

Acknowledgements

This work was funded by Wellcome though the Darwin Tree of Life Discretionary Award

(218328) and supported by the Biotechnology and Biological Sciences Research Council

(BBSRC), part of UK Research and Innovation, through the Core Capability Grant

BB/CCG1720/1 at the Earlham Institute. Part of this work was delivered via the BBSRC

National Capability in Genomics and Single Cell Analysis (BBS/E/T/000PR9816) at Earlham

Institute by members of the Genomics Pipelines, Single☐Cell and Core Bioinformatics

Groups, the authors, note the specific contributions of Tom Barker, Vanda Knitlhoffer and

Chris Watkins. Additional work was delivered via the BBSRC National Capability in

e☐Infrastructure (BBS/E/T/000PR9814) at the Earlham Institute by members of the

e☐Infrastructure group. The authors would like to acknowledge the Scientific Computing

group, as well as support for the physical HPC infrastructure and data centre delivered via the

NBI Research Computing group. TAR is supported by a Royal Society University Research

Fellowship (URF/R/191005).

Data Availability

564 All sequencing data and the genome assembly of Oligohymenophorea sp. PL0344 have been

565 deposited to the European Nucleotide Archive under the study accession PRJEB58266.

566 Additional supporting data have been deposited on Zenodo (10.5281/zenodo.7373057).

Figure Legends

Figure 1: Genetic code change in Oligohymenophorea sp. PL0344. Example multiple sequence alignment of a tubulin gamma chain protein and orthologous sequences spanning Eukaryota. The alignment has been trimmed for visualisation purposes to remove poorly conserved regions and highlight internal UAA and UAG codons.

Figure 2: Genetic code prediction for Oligohymenophorea sp. PL0344. PhyloFisher genetic code prediction for the **(A)** UAA and **(B)** UAG codons using the PhyloFisher database of 240 orthologs. Only well conserved (>70%) amino acids are considered. Colours correspond to amino acid properties and match the multiple sequence alignment in Figure 1. **(C)** Codetta genetic code prediction. Log decoding probabilities for the UAA and UAG codons are shown for each of the 20 standard amino acids.

Figure 3: Example tRNA Genes. **(A)** Predicted secondary structure of an example tRNA-Sup(UUA) predicted to function as a lysine tRNA. The wobble position is highlighted. According to wobble-binding rules, uracil at this position can bind to either adenine or guanine in the third codon position of mRNA, allowing the suppressor tRNA to recognise both UAA and UAG stop codons. **(B)** Predicted secondary structure of an example tRNA-Sup(CUA) predicted to function as a glutamic acid tRNA. **(C)** Predicted secondary structure of the tRNA-SeC(UCA) for selenocysteine.

Figure 4: Enrichment of tandem stop codons. The proportion of codon positions occupied by UGA in the 20 in-frame codon positions immediately downstream of all genes and highly expressed genes. Positions where UGA is significantly overrepresented (chi-squared test, p-value < 0.05) are indicated with an asterisk.

Figure 5: **Phylogenomic analysis of genetic code changes in the Ciliophora.** Maximum-

likelihood phylogeny of 46 ciliate species and 9 outgroup species from the Alveolata, based

on a concatenated alignment of 89 BUSCO proteins (40,289 amino acid sites) under the

LG+F+I+R7 model using IQ-TREE. The values at branches represent statistical support from

100 non-parametric bootstraps with the LG+F+I+R7 model, 100 non-parametric bootstraps

from the IQ-TREE partitioned analysis, and Bayesian posterior probabilities determined

under the CAT-GTR model in PhyloBayes-MPI. Branches with full support from all three

approaches (i.e., 100/100/1) are indicated with solid black circles. Hyphens indicate branches

that weren't recovered. Stop codon reassignments are shown (*, STOP; Q, glutamine; W,

tryptophan; K, lysine; E, glutamic acid; Y, tyrosine; C, cysteine). The percentage of proteins

included in the concatenated alignment is shown in the bar plot, highlighting the amount of

missing data per species.

Supplementary Figure 1: Maximum-likelihood phylogeny of small subunit ribosomal RNA

genes under the GTR+F+R5 model using IQ-TREE with 100 non-parametric bootstraps.

Supplementary Figure 2: Bayesian phylogenomic analysis of 46 ciliate species and 9

outgroup Alveolata species, based on a concatenated alignment of 89 BUSCO proteins under

the CAT-GTR model using PhyloBayes-MPI.

Supplementary Figure 3: Example multiple sequence alignments of *Plagiopyla frontata*

genes with internal UGA codons identified by PhyloFisher with orthologous sequences

spanning Eukaryota. **A)** TM9SF1. **B)** PIK3C3. **C)** CRNL1.

Table Captions

Table 1: Genome assembly and annotation statistics

Supplementary Table 1: tRNA genes pairwise identities

Supplementary Table 2: Datasets used for phylogenomics and genome annotation

Supplementary Table 3: Amino acid and codon usage

Table 1. Genome Assembly and Annotation Statistics

Genome Assembly	
Total length	69,783,056 bp
Contigs	3671
N50	59,570 bp
GC content	30.58%
RNA-Seq mapping rate	89%
BUSCO completeness[1]	Complete: 84.8% Complete and single copy: 80.1% Complete and duplicated: 4.7% Fragmented: 3.5% Missing: 11.7%
Genome Annotation	
Number of genes	20,141
Number of transcripts	22,084
Number of monoexonic genes	7,080
Exons per transcript	3.76
GC content (CDS)	34.12%
% of genome covered by CDS	63.8%
BUSCO completeness[1]	Complete: 94.74% Complete and single copy: 87.72% Complete and duplicated: 7.02% Fragmented: 1.75% Missing: 3.51%

	Median	Mean
CDS size (bp)	1,506	2,014.12
Intron size (bp)	57	80.39
5'UTR size (bp)	65	96.74
3'UTR size (bp)	95	137.61
Intergenic distances	156	574.43

[1]BUSCO completeness assessed using V4 with the Alveolata_obd10 dataset

References

Adachi M, Cavalcanti ARO. 2009. Tandem stop codons in ciliates that reassign stop codons. J. Mol. Evol. 68:424–431.

Andersson SGE, Kurland CG. 1995. Genomic evolution drives the evolution of the translation system. Biochem. Cell Biol. 73:775–787.

Arnaiz O et al. 2012. The Paramecium Germline Genome Provides a Niche for Intragenic Parasitic DNA: Evolutionary Dynamics of Internal Eliminated Sequences. PLoS Genet. 8:e1002984.

Aury J-M et al. 2006. Global trends of whole-genome duplications revealed by the ciliate Paramecium tetraurelia. Nature. 444:171–178.

Bankevich A et al. 2012. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J. Comput. Biol. 19:455–477.

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq quantification. Nat. Biotechnol. 34:525–527.

Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 25:1972–1973.

Chan PP, Lin BY, Mak AJ, Lowe TM. 2021. tRNAscan-SE 2.0: improved detection and functional classification of transfer RNA genes. Nucleic Acids Res. 49:9077–9096.

Chen X et al. 2015. Phylogenomics of non-model ciliates based on transcriptomic analyses. Protein Cell. 6:373–385.

Chernomor O, von Haeseler A, Minh BQ. 2016. Terrace Aware Data Structure for Phylogenomic Inference from Supermatrices. Syst. Biol. 65:997–1008.

Crick FHC. 1966. Codon—anticodon pairing: The wobble hypothesis. J. Mol. Biol. 19:548–555.

Eisen JA et al. 2006. Macronuclear genome sequence of the ciliate Tetrahymena thermophila,

a model eukaryote. PLoS Biol. 4:e286.

Eliseev B, Kryuchkova P, Alkalaeva E, Frolova L. 2011. A single amino acid change of translation termination factor eRF1 switches between bipotent and omnipotent stop-codon specificity †. Nucleic Acids Res. 39:599–608.

Feng J-M et al. 2015. Phylogenomic analyses reveal subclass Scuticociliatia as the sister group of subclass Hymenostomatia within class Oligohymenophorea. Mol. Phylogenet. Evol. 90:104–111.

Fleming I, Cavalcanti ARO. 2019. Selection for tandem stop codons in ciliate species with reassigned stop codons Kapler, GM, editor. PLoS One. 14:e0225804.

Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-generation sequencing data. Bioinformatics. 28:3150–3152.

Gao F, Katz LA, Song W. 2013. Multigene-based analyses on evolutionary phylogeny of two controversial ciliate orders: Pleuronematida and Loxocephalida (Protista, Ciliophora, Oligohymenophorea). Mol. Phylogenet. Evol. 68:55–63.

Gentekaki E, Kolisko M, Gong Y, Lynn D. 2017. Phylogenomics solves a long-standing evolutionary puzzle in the ciliate world: The subclass Peritrichia is monophyletic. Mol. Phylogenet. Evol. 106:1–5.

Grabherr MG et al. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat. Biotechnol. 29:644–652.

Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. Bioinformatics. 29:1072–1075.

Haas BJ et al. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. Nat. Protoc. 8:1494–1512.

Hanyu N, Kuchino Y, Nishimura S, Beier H. 1986. Dramatic events in ciliate evolution: alteration of UAA and UAG termination codons to glutamine codons due to anticodon

mutations in two Tetrahymena tRNAs Gln. EMBO J. 5:1307–1311.

Heaphy SM, Mariotti M, Gladyshev VN, Atkins JF, Baranov P V. 2016. Novel Ciliate
Genetic Code Variants Including the Reassignment of All Three Stop Codons to Sense
Codons in Condylostoma magnum. Mol. Biol. Evol. 33:2885–2889.

Hubley, R. & Smit A.F.A. 2008. RepeatModeler at

http://www.repeatmasker.org/RepeatModeler/

Hubley, R., Green, P. & Smit A.F.A. 2008. RepeatMasker at http://repeatmasker.org

Irwin NAT et al. 2021. The Function and Evolution of Motile DNA Replication Systems in
Ciliates. Curr. Biol. 31:66-76.e6.

Ivanova NN et al. 2014. Stop codon reassignments in the wild. Science (80-.). 344:909–913.

Jiang C-Q et al. 2019. Insights into the origin and evolution of Peritrichia
(Oligohymenophorea, Ciliophora) based on analyses of morphology and phylogenomics.
Mol. Phylogenet. Evol. 132:25–35.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder:
fast model selection for accurate phylogenetic estimates. Nat. Methods. 14:587–589.

Kang DD et al. 2019. MetaBAT 2: an adaptive binning algorithm for robust and efficient
genome reconstruction from metagenome assemblies. PeerJ. 7:e7359.

Karlicki M, Antonowicz S, Karnkowska A. 2022. Tiara: deep learning-based classification
system for eukaryotic sequences Birol, I, editor. Bioinformatics. 38:344–350.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Keeling PJ. 2016. Genomics: Evolution of the Genetic Code. Curr. Biol. 26:R851–R853.

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. 2019. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. Nature Biotechnology, 37(8),
907–915.

Knight RD, Freeland SJ, Landweber LF. 2001. Rewiring the keyboard: evolvability of the genetic code. Nat. Rev. Genet. 2:49–58.

Kollmar M, Mühlhausen S. 2017. Nuclear codon reassignments in the genomics era and mechanisms behind their evolution. BioEssays. 39:1–12.

Koonin E V., Novozhilov AS. 2009. Origin and evolution of the genetic code: The universal enigma. IUBMB Life. 61:99–111.

Kovaka, S., Zimin, A. V., Pertea, G. M., Razaghi, R., Salzberg, S. L., & Pertea, M. 2019. Transcriptome assembly from long-read RNA-seq alignments with StringTie2. Genome Biology, 20(1), 1–13.

Kryuchkova P et al. 2013. Two-step model of stop codon recognition by eukaryotic release factor eRF1. Nucleic Acids Res. 41:4573–4586.

Lartillot N, Rodrigue N, Stubbs D, Richer J. 2013. PhyloBayes MPI: Phylogenetic Reconstruction with Infinite Mixtures of Profiles in a Parallel Environment. Syst. Biol. 62:611–615.

Lasek-Nesselquist E, Johnson MD. 2019. A Phylogenomic Approach to Clarifying the Relationship of Mesodinium within the Ciliophora: A Case Study in the Complexity of Mixed-Species Transcriptome Analyses Zufall, R, editor. Genome Biol. Evol. 11:3218–3232.

Laslett D. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. Nucleic Acids Res. 32:11–16.

Lekomtsev S et al. 2007. Different modes of stop codon restriction by the Stylonychia and Paramecium eRF1 translation termination factors. Proc. Natl. Acad. Sci. 104:10824–10829.

Liang H, Cavalcanti ARO, Landweber LF. 2005. Conservation of tandem stop codons in yeasts. Genome Biol. 6:R31.

Lozupone CA, Knight RD, Landweber LF. 2001. The molecular basis of nuclear genetic code change in ciliates. Curr. Biol. 11:65–74.

Lynn DH, Kolisko M. 2017. Molecules illuminate morphology: phylogenomics confirms convergent evolution among 'oligotrichous' ciliates. Int. J. Syst. Evol. Microbiol. 67:3676–3682.

Macaulay IC et al. 2015. G&T-seq: parallel sequencing of single-cell genomes and transcriptomes. Nat. Methods. 12:519–522.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes Kelley, J, editor. Mol. Biol. Evol. 38:4647–4654.

Mapleson, D., Venturini, L., Kaithakottil, G., & Swarbreck, D. 2018. Efficient and accurate detection of splice junctions from RNA-seq with Portcullis. GigaScience, 7(12).

Von Meijenfeldt FAB, Arkhipova K, Cambuy DD, Coutinho FH, Dutilh BE. 2019. Robust taxonomic classification of uncharted microbial sequences and bins with CAT and BAT. Genome Biol. 20:217.

Meyer F et al. 1991. UGA is translated as cysteine in pheromone 3 of Euplotes octocarinatus. Proc. Natl. Acad. Sci. 88:3758–3761.

Minh BQ et al. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era Teeling, E, editor. Mol. Biol. Evol. 37:1530–1534.

Mühlhausen S, Findeisen P, Plessmann U, Urlaub H, Kollmar M. 2016. A novel nuclear genetic code alteration in yeasts and the evolution of codon reassignment in eukaryotes. Genome Res. 26:945–955.

Osawa S, Jukes TH, Watanabe K, Muto A. 1992. Recent evidence for evolution of the genetic code. Microbiol. Rev. 56:229–264.

Pánek T et al. 2017. Nuclear genetic codes with a different meaning of the UAG and the UAA codon. BMC Biol. 15:8.

Parfrey LW, Lahr DJG, Knoll AH, Katz LA. 2011. Estimating the timing of early eukaryotic diversification with multigene molecular clocks. Proc. Natl. Acad. Sci. 108:13624–13629.

Poláková K, Čepička I, Bourland WA. 2021. Phylogenetic Position of Three Well-known Ciliates from the Controversial Order Loxocephalida Jankowski, 1980 (Scuticociliatia, Oligohymenophorea) and Urozona buetschlii (Schewiakoff, 1889) with Improved Morphological Descriptions. Protist. 172:125833.

Prescott DM. 1994. The DNA of ciliated protozoa. Microbiol. Rev. 58:233–67.

Richter DJ et al. 2022. EukProt: a database of genome-scale predicted proteins across the diversity of eukaryotes. bioRxiv. 2020.06.30.180687.

Rotterová J et al. 2020. Genomics of New Ciliate Lineages Provides Insight into the Evolution of Obligate Anaerobiosis. Curr. Biol. 30:2037-2050.e6.

Salim, H., Ring, K. and Cavalcanti, A., 2008. Patterns of Codon Usage in two Ciliates that Reassign the Genetic Code: Tetrahymena thermophila and Paramecium tetraurelia. Protist, 159(2), pp.283-298.

Schultz DW, Yarus M. 1994. Transfer RNA Mutation and the Malleability of the Genetic Code. J. Mol. Biol. 235:1377–1380.

Seah B, Singh A, Swart E. 2022. Karyorelict ciliates use an ambiguous genetic code with context-dependent stop/sense codons. Peer Community Journal 2.

Shao, M., & Kingsford, C. 2017. Accurate assembly of transcripts through phase-preserving graph decomposition. Nature Biotechnology, 35(12), 1167–1169.

Shulgina Y, Eddy SR. 2021. A computational screen for alternative genetic codes in over 250,000 genomes. Elife. 10:1–25.

Slabodnick MM et al. 2017. The Macronuclear Genome of Stentor coeruleus Reveals Tiny Introns in a Giant Cell. Curr. Biol. 27:569–575.

Šlapeta J, Moreira D, López-García P. 2005. The extent of protist diversity: insights from

molecular ecology of freshwater eukaryotes. Proc. R. Soc. B Biol. Sci. 272:2073–2081.

Stanke, M., & Morgenstern, B. 2005. AUGUSTUS: a web server for gene prediction in eukaryotes that allows user-defined constraints. Nucleic Acids Research, 33(Web Server), W465–W467.

Swart EC et al. 2013. The Oxytricha trifallax Macronuclear Genome: A Complex Eukaryotic Genome with 16,000 Tiny Chromosomes Eisen, JA, editor. PLoS Biol. 11:e1001473.

Swart EC, Serra V, Petroni G, Nowacki M. 2016. Genetic Codes with No Dedicated Stop Codon: Context-Dependent Translation Termination. Cell. 166:691–702.

Tice AK et al. 2021. PhyloFisher: A phylogenomic package for resolving eukaryotic relationships Hejnol, A, editor. PLOS Biol. 19:e3001365.

Venturini, L., Caim, S., Kaithakottil, G. G., Mapleson, D. L., & Swarbreck, D. 2018. Leveraging multiple transcriptome assembly methods for improved gene structure annotation. GigaScience, 7(8).

Wang C et al. 2021. Large-scale phylogenomic analysis provides new insights into the phylogeny of the class Oligohymenophorea (Protista, Ciliophora) with establishment of a new subclass Urocentria nov. subcl. Mol. Phylogenet. Evol. 159:107112.

	A	C	D	E	F	G	H	I	K	L	M	N	P	Q	R	S	T	V	W	Y	?
UAA	-7247.7	-12835.5	-8178.5	-5787.8	-14432.1	-10288.2	-6864.4	-12385.8	0.0	-10974.3	-10337.2	-6106.2	-11220.1	-4178.8	-3215.4	-6325.0	-6863.7	-10629.3	-15433.8	-11716.0	-3255.1
UAG	-5129.3	-9789.2	-3214.5	0.0	-10447.8	-7064.8	-5266.2	-9348.6	-4089.2	-8450.4	-8002.1	-4528.5	-7525.4	-3136.4	-4933.1	-4482.9	-5141.5	-8054.1	-11143.0	-8505.1	-2540.6

A. tRNA-Sup(UUA)   **B.** tRNA-Sup(CUA)   **C.** tRNA-SeC(UCA)

Wobble position

Tree scale: 0.1

Class

- Armophorea
- Colpodea
- Heterotrichea
- Litostomatea
- Muranotrichea
- Nassophorea
- Oligohymenophorea
- Plagiopylea
- Protocruziea
- Spirotrichea

Tree scale: 1

A) TM9SF1

B) PIK3C3

C) CRNL1