How sexual dimorphism in phenotypic plasticity may evolve

Anton S. Zadorin[1] and Olivier Rivoire[2]

[1]*Laboratory of Biophysics and Evolution, Chemistry Biology Innovation,*
ESPCI, Université PSL University, Paris, France.
[2]*Center for Interdisciplinary Research in Biology (CIRB),*
Collège de France, CNRS, INSERM, Université PSL, Paris, France.

Phenotypes are partly shaped by the environment, which can impact both short-term adaptation and long-term evolution. In dioecious species, the two sexes may exhibit different degrees of phenotypic plasticity and theoretical models indicate that such differences may confer an adaptive advantage when the population is subject to directional selection, either because of a systematically varying environment or of a load of deleterious mutations. The effect stems from the fundamental asymmetry between the two sexes: female fertility is more limited than male fertility. Whether this asymmetry is sufficient for sexual dimorphism in phenotypic plasticity to evolve is, however, not obvious. Here, we show that even in conditions where it provides an adaptive advantage, dimorphic phenotypic plasticity may be evolutionarily unstable due to sexual selection. This is the case, in particular, for panmictic populations where mating partners are formed at random. However, we show that the effects of sexual selection can be counteracted when mating occurs within groups of related individuals. Under this condition, sexual dimorphism in phenotypic plasticity can not only evolve but offset the twofold cost of males. We demonstrate these points with a simple mathematical model through a combination of analytical and numerical results.

I. INTRODUCTION

Phenotypic plasticity, the influence of the environment on the development of phenotypes, is a well-recognized component of evolutionary change [1] with the potential to constitute an essential adaptation to varying environments [2]. Sex differences in phenotypic plasticity are documented in several dioecious species [3, 4], but only recently has the question of the adaptive value of these sex differences started to be studied mathematically [4, 5]. Remarkably, these studies show that sexual dimorphism in phenotypic plasticity can enhance population growth and thus prevent the extinction of populations subject to directional selective pressures. They have left open, however, the question of the conditions under which such an adaptive dimorphism may evolve [4, 5]. A puzzling observation is indeed that sexual dimorphism can fail to evolve even in conditions where it is adaptive [5].

Understanding the conditions under which sexual dimorphism in phenotypic plasticity can evolve has at least two motivations. On one hand is the pressing issue of determining the role that phenotypic plasticity may play in the adaptation of extant species to current climate changes [6]. On the other hand is the long-standing question of the origin and prevalence of dioecious species given the "two-fold cost of males" that they incur compared to monoecious species: a monoecious species where each individual would be as fecund as the females of a dioecious species would indeed produce twice as many offsprings per generation and therefore have a major selective advantage [7]. This raises two questions: (1) Under which conditions may sexual dimorphism be an adaptation that offsets the two-fold cost of males? (2) Under which conditions may such a population-level adaptation be reachable through evolution?

In previous work [5], we started to examine these questions with a focus on the adaptive value and evolution of sexual dimorphism in developmental variances – the possibility that one sex displays higher developmental canalization than the other. We found that such a sexual dimorphism could in principle

be adaptive but only under restricted conditions that made it unlikely to be a generic cause of phenotypic differences between sexes, or a fortiori to explain why many species comprise two sexes. Further, we found that an optimal difference in developmental variances failed to evolve even in conditions where it was adaptive. Here we revisit these questions with a focus on sexual differences in phenotypic plasticity. Although the two problems can formally be mapped on each other [5], we show that sexual dimorphism in phenotypic plasticity can possibly confer a substantial adaptive advantage over a much broader range of conditions.

To demonstrate this result, we study a simple but generic model for the evolution of sexually dimorphic plasticity. This model extends previous models [2, 4, 5, 8, 9] with several crucial differences. First, it differs from models of quantitative genetics [2, 4, 8] by introducing an explicit dynamics for genetic changes, which allows us to readily analyze the evolution of plasticity through modifier genes [10]. Second, compared to our previous models [5, 9], it accounts for varying degrees of sex-specific limitations in fertility, as well as for the possibility that mating interactions are limited to groups of genetically related individuals. We show that mating patterns – the rules that govern the associations between males and females – are critical to both the adaptive value and the evolution of dimorphic plasticity. In particular, we identify simple mating patterns for which sexual dimorphism in phenotypic plasticity is evolvable with an adaptive value that offsets the two-fold cost of males.

II.   MODEL

A.   Model with fixed plasticity

We consider a population of individuals, each with a continuous one-dimensional genotype γ, a continuous one-dimensional phenotype ϕ and a sex $\bullet$, which can either be female ($\bullet = ♀$) or male ($\bullet = ♂$). At each discrete generation t, the environment x_t defines an optimal phenotype, which may be directionally varying as $x_t = c_E t$, where c_E therefore represents a rate of change. The phenotype ϕ of an individual dictates its chances of survival and reproduction. Following previous models [2, 4, 5, 8, 9], we assume that the probability for an individual with phenotype ϕ to survive and possibly reproduce in environment x_t is $S(\phi, x_t) = e^{-(\phi - x_t)^2/(2\sigma_S^2)}$ where σ_S^2 represents the stringency of selection. We assume that an individual whose parents have genotypes $\gamma^♀$ and $\gamma^♂$ inherits a mid-genotype

$$\gamma = \frac{\gamma^♀ + \gamma^♂}{2} - c_M + \mu, \qquad \mu \sim \mathcal{N}(\sigma_M^2) \tag{1}$$

where μ is a random variable normally distributed with zero mean and variance σ_M^2 and where c_M is a possible mutational bias towards deleterious mutations. This form of inheritance is justified if γ represents the contribution of many loci that are independently transmitted at random from each parent.

Following previous models [2, 4, 5, 9], we consider the phenotype ϕ of an individual of sex $\bullet$ to possibly depend both on its genotype γ and on its environment x_t through the relation

$$\phi = \lambda_\bullet \gamma + \kappa_\bullet x_t + \zeta, \qquad \zeta \sim \mathcal{N}(\sigma_D^2) \tag{2}$$

where σ_D^2 represents a developmental variance. We take $\lambda_\bullet = 1 - \kappa_\bullet$ to interpolate between $\kappa_\bullet = 0$ corresponding to no plasticity and $\kappa_\bullet = 1$ corresponding to perfect plasticity [5]. $\kappa_\bullet$ is interpreted as a reaction norm, which may here depend on the sex $\bullet$ of the individual. We previously examined the case of sex-specific developmental variances [5] but assume here that a common σ_D^2 applies to both sexes.

Finally, the sex of an individual is chosen at random with no bias, $\bullet = ♀$ or $♂$ with same probability $1/2$, which we write as $\bullet = \text{rand}([♀, ♂])$.

B.   Model with evolving plasticity

To study the evolution of sex-specific phenotypic plasticity we augment the genotype γ of an individual to include two modifier genes $y_{\female}, y_{\male}$ that respectively control the female and male reaction norms $\kappa_{\female}$ and $\kappa_{\male}$. The complete genotype of an individual is therefore of the form $\Gamma = (\gamma, y_{\female}, y_{\male})$. In this more general model, the phenotype of an individual consists of its continuous trait ϕ, its sex $\bullet$ and its reaction norm $\kappa_{\bullet}$. It has therefore the form $\Phi = (\phi, \bullet, \kappa_{\bullet})$. We take the relation between $y_{\bullet}$ and $\kappa_{\bullet}$ to be

$$\kappa_{\bullet} = \frac{\kappa^{*}}{1 + e^{y_{\bullet}}} \tag{3}$$

so as to map the continuous trait $y_{\bullet}$ into an interval $[0, \kappa^{*}]$ where $\kappa^{*} < 1$ represents a maximal reaction norm.

We assume that the modifier genes follow the same inheritance rules as γ, i.e., given parents with genotypes $\Gamma^{\female} = (\gamma^{\female}, y_{\female}^{\female}, y_{\male}^{\female})$ and $\Gamma^{\male} = (\gamma^{\male}, y_{\female}^{\male}, y_{\male}^{\male})$, we assume that an offspring inherits

$$y_{\bullet} = \frac{y_{\bullet}^{\female} + y_{\bullet}^{\male}}{2} + \nu, \qquad \nu \sim \mathcal{N}(\sigma_{V}^{2}) \tag{4}$$

with $\bullet = \female$ and $\male$. Here we introduce a mutational variance σ_{V}^{2} that may differ from σ_{M}^{2} but note that we could equivalently consider $\sigma_{V}^{2} = \sigma_{M}^{2}$ by redefining $\kappa_{\bullet}$ as $\kappa_{\bullet} = \kappa^{*}/(1 + e^{y_{\bullet} \sigma_{M}/\sigma_{V}})$. The present formulation is adopted to recover the case of a fixed plasticity as the limit where $\sigma_{V}^{2} = 0$.

The introduction of a maximal reaction norm $\kappa^{*} < 1$ is motivated by the fact that in absence of further constraints, $\kappa_{\bullet} = 1$ guarantees that $\phi = x_{t}$, i.e., perfect adaptation independently of the genotype γ. This extreme degree of plasticity is optimal but not biologically feasible and models usually integrate a fitness cost that penalizes large values of $\kappa_{\bullet}$ [2]. Our parameter κ^{*} is a simple representation of this cost, with values $\kappa \leq \kappa^{*}$ corresponding to no cost and values $\kappa > \kappa^{*}$ to infinite cost.

C.   Mating patterns

Following Eqs. (2)-(3), a newborn individual with genotype $\Gamma = (\gamma, y_{\female}, y_{\male})$ develops to have phenotype $\Phi = (\phi, \bullet, \kappa)$. This individual reaches maturation with probability $e^{-(\phi - x_{t})^{2}/(2\sigma_{S}^{2})}$ and otherwise dies. Next, we assume that each mature individual of sex $\bullet = \female$ or $\male$ has the capacity to participate in $q_{\bullet}$ mating events, each producing one individual of the next generation. This maximal capacity may, however, not be achieved if partners of the opposite sex are not available. If $M_{\female,t}$ and $M_{\male,t}$ are, respectively, the number of mature females and males at generation t, the number of newborn at generation $t + 1$ is therefore

$$N_{t+1} = \min(q_{\female} M_{\female,t}, q_{\male} M_{\male,t}). \tag{5}$$

This relationship accounts both for the case where the limitation is the number of females ($q_{\female} M_{\female,t} < q_{\male} M_{\male,t}$) or the number of males ($q_{\female} M_{\female,t} > q_{\male} M_{\male,t}$). By definition of females, $q_{\female} < q_{\male}$, with a limit case being when males have unlimited mating capacities ($q_{\male} = \infty$), in which case the limiting sex is necessarily females, with simply $N_{t+1} = q_{\female} M_{\female,t}$.

We also consider the possibility for the population to be divided into different groups with mating couples formed at random only within each group. We take these groups to have a maximal size K and assume that when the number of newborn individuals within a group exceeds K, this group splits into randomly formed sub-groups of size $K/2$. A limit case is here that of panmictic populations where couples are formed totally at random between available mature individuals of each sex, which corresponds to K being equal or larger than the population size.

In general, we expect a population following these rules to grow exponentially whenever $q_{\female}$ and $q_{\male}$ are sufficiently large, with a growth rate Λ such that $N_{t}/N_{0} \sim e^{\Lambda t}$ for large t. This growth rate can

Evolving variables	
$\bullet$	sex, which may be either female ($\bullet = \female$) or male ($\bullet = \male$)
ϕ	one-dimensional trait under direct selection
γ	genotype controlling the main trait ϕ
$\kappa_\bullet$	sex-specific reaction norm
$y_\bullet$	modifier gene controlling the reaction norm $\kappa_\bullet$
Environmental parameters	
x_t	environmental variable defining the optimal phenotype ϕ at generation t
$c_E = c$	rate of environmental change ($x_t = c_E t$)
σ_S^2	stringency of selection
Genetic parameters	
κ^*	maximal reaction norm
$c_M = c$	mean effect of deleterious mutations
σ_M^2	mutation rate for trait γ
σ_V^2	mutation rate for the modifier genes $y_\female$ and $y_\male$ ($\sigma_V^2 = 0$ for non-evolving reaction norms)
σ_D^2	developmental variance
Mating parameters	
$q_\bullet$	maximal number of matings to which an individual of sex $\bullet$ can participate
K	maximal size of the groups within which mating is restricted
Population parameters	
N	total population size
Λ	long-term population growth rate
$\chi_\bullet$	fraction of individuals of sex $\bullet$ reaching maturation at each generation

TABLE I: Summary of the main notations used in the text and figures.

be computed analytically for large panmictic populations (Methods). More generally, the dynamics can be studied numerically with an agent-based simulation. In practice, it is convenient in numerical simulations to impose a total population size N and resample individuals at each generation to maintain this size (Methods). For panmictic populations, we can then compare the analytically computed growth rate Λ to its numerical approximation $\Lambda_{T,N} = (1/T) \sum_{t=1}^{T} \ln(N_t/N)$ where N_t is the number of newborn individuals at generation t and T the total number of generations.

D. Parameters

As so far defined, the model has ten parameters, σ_S^2, σ_D^2, σ_M^2, σ_V^2, c_E, c_M, κ^*, $q_\female$, $q_\male$, K (see Table I), not counting the initial conditions, the total population size N and the total number of generations T that must also be set when performing numerical simulations. Our results, however, depend only on a subset of these parameters and we therefore set the others. For instance, by rescaling other variances we can always assume $\sigma_S^2 = 1$ [9]. When $q_\male = \infty$, $q_\female$ contributes only additively to the growth rate and as we are only interested in differences in growth rates, we can also fix $q_\female$. Additionally, we note that c_E and c_M play formally equivalent roles and can therefore be represented by a single parameter c which quantifies a directional selection independently of its environmental or genetic origin (Methods).

We first examine the model with non-evolving reaction norms ($\sigma_V^2 = 0$) in the limit of an infinite panmictic population ($N = K = \infty$). In this case, two additional parameters are the fixed reaction norms of each sex, $\kappa_\female$ and $\kappa_\male$. Next ,we consider the long-term evolution of large populations with $\sigma_V^2 > 0$, in which case $\kappa_\female$ and $\kappa_\male$ can differ between individuals of a same populations, thus leading us

FIG. 1: **A.** Long-term growth rate Λ of large unstructured populations with non-evolving reaction norms as a function of the male reaction norm $\kappa_\male$ for two extreme values of the female reaction norm $\kappa_\female$. The full lines are from analytical calculations and the dots from numerical simulations, showing very good agreement. **B.** Long-term growth rate Λ as a function of $\kappa_\male$ and $\kappa_\female$, showing a maximum for dimorphic plasticity when $\kappa_\male = 0$ and $\kappa_\female = \kappa^*$ (upper left corner). **C.** Values of the reaction norms that optimize the growth rate Λ as a function of the ratio $q_\male/q_\female$ where $q_\bullet$ controls the maximal number of mating events to which an individual of sex $\bullet$ can participate ($\bullet = \female$ or $\male$). The case of female demographic dominance corresponds to $q_\male/q_\female > 1$, in which case it is optimal for females to be plastic ($\hat\kappa_\female > 0$) and for males to be non-plastic ($\kappa_\male = 0$). For sufficiently large ratios $q_\male/q_\female$, it is even optimal for female reaction norms to be maximal ($\kappa_\female = \kappa^*$). The conclusions are symmetrically reversed if $q_\male/q_\female < 1$.

to analyze population-wise mean values $\bar\kappa_\female$ and $\bar\kappa_\male$.

Unless otherwise indicated, we take the following values of the parameters: $q_\male = \infty$, $q_\female = 4$, $c = 10^{-1}$, $\sigma_M^2 = 10^{-1}$, $\sigma_V^2 = 10^{-2}$, $\sigma_D^2 = 0$, $\kappa^* = 0.5$, and, when performing numerical simulations, $N = 10^3$ and $T = 10^4$ with initial conditions such that $\kappa_\female = \kappa_\male = \kappa^*/2$ for all individuals.

III. RESULTS

A. Optimal plasticity

For infinite panmictic populations ($N = K = \infty$) with non-evolving reaction norms ($\sigma_V^2 = 0$), the long-term growth rate Λ can be computed analytically (Methods), with results matching well with those of numerical simulations (Fig. 1A). These results indicate that the growth rate Λ is maximal when the two sex-specific reaction norms $\kappa_\female$ and $\kappa_\male$ take different values, namely $\hat\kappa_\female = \kappa^*$ and $\hat\kappa_\male = 0$ (Fig. 1B). This corresponds to maximally plastic females and minimally plastic males, consistent with previous findings showing that sexual dimorphism can optimize the long-term growth of populations subject to directional selection [4, 5]. Given that fertility is limited by females, this result may be interpreted as follows: maximizing female plasticity maximizes the number of offsprings at the next generation, while minimizing male plasticity maximizes the fitness value of the genotype that these offsprings inherit, as only males with well adapted genotypes reach maturity. In other words, males pay the price of natural selection to guarantee long-term adaptation while females buffer environmental effects to guarantee short-term growth. These results depend primarily on the assumption that fertility is limited by females ($q_\male \gg q_\female$) and the opposite conclusions are reached if $q_\male \ll q_\female$. When $q_\female < q_\male$ but $q_\male$ is comparable to $q_\female$, we also find that the optimal female reaction norm $\hat\kappa_\female$ takes a non-zero value that is independent of the maximal possible reaction norm κ^* (Fig. 1C).

FIG. 2: **A-C.** Examples of evolutionary trajectories, showing the mean values of the reaction norms $\kappa_{\female}$ (in pink) and $\kappa_{\male}$ (in blue) across 10 different populations as a function of the number t of generations, starting from populations with $\kappa_{\bullet} = \kappa^*/2$ for all individuals. In presence of small groups (A), dimorphic plasticity evolves where male reaction norms are minimal while in presence of large groups (C) monomorphic plasticity evolves where male reaction norms are maximal. In intermediate cases (B), male reaction norms may take arbitrary values. K indicates the maximal size of the groups and N the total population size. **D.** Mean value of the reaction norms $\kappa_{\female}$ and $\kappa_{\male}$ after $T = 10^4$ generations (averaged over 100 trajectories) showing how the evolution of dimorphic plasticity depends on the maximal group size K of the groups (displayed in log-scale). As in panels A-C, the simulations are performed under the hypothesis that fertility is only limited by females ($q_{\male} = \infty$). **E.** If instead of assuming that each male can participate in an arbitrary number of mating events we assume that they can participate in at most twice as many mating events as females ($q_{\male} = 2q_{\female}$), qualitatively similar results are obtained. **F.** If males and females are subject to the same limitations in fertility ($q_{\male} = q_{\female}$), however, the symmetry between the two sexes prevents any sexual dimorphism to evolve.

B.   Evolution of plasticity

While a dimorphic plasticity with $\kappa_{\female} = \kappa^*$ and $\kappa_{\male} = 0$ maximizes long-term population growth, we find that when reactions norms are subject to evolution ($\sigma_V^2 > 0$), they converge to the same value, namely $\bar{\kappa}_{\female} = \bar{\kappa}_{\male} = \kappa^*$ (Fig. 2C). Evolution thus leads to monomorphic plasticity that is sub-optimal for long-term growth. This may be interpreted as arising from a conflict between two levels of selection: long-term population growth benefits from low male plasticity but sexual selection among males promotes high male plasticity. The outcome of such a conflict can be reversed by group selection [11], as we verify here by considering mating to be structured into groups. We find that in presence of groups, evolution of the reaction norms can indeed lead to a dimorphic plasticity that maximizes population growth (Fig. 2A). This requires the size of the groups to be sufficiently small (Fig. 2D). This critical size K^* is found to scale with the total population size N as $N^{1/2}$ (Fig. S1), which indicates that for larger populations, proportionally larger groups can lead to the evolution of sexual dimorphism in plasticity. We also verify that K^* scales with σ_V^2 as $(\sigma_V^2)^\alpha$ (Fig. S1), as is typical for this type of models [12]. Group selection enables the evolution of dimorphic plasticity whenever fertility is controlled by females, even if male fertility is also limited, provided this limitation is to a lesser extent (Fig. 2E). When the two sexes have same fertility limits, however, no asymmetry is present and we verify that no dimorphism evolves (Fig. 2F). Finally, we note that the variance between groups that is necessary for group selection to operate arises here from the closer genetic relationships within than between groups, which can also be viewed as a form of kin selection [11].

FIG. 3: **A.** Gain in growth rate from sexual dimorphism in phenotypic plasticity, $\Delta\Lambda = \Lambda(\kappa_\female = \kappa^*, \kappa_\male = 0) - \Lambda(\kappa_\female = \kappa^*, \kappa_\male = \kappa^*)$, as a function of c, the rate at which the selective pressure is varying (in blue). The vertical dashed line marks the value c^* above which this difference exceeds the two fold cost of males, $\ln 2$ (in green). The fraction $\chi_\male$ of surviving males at each generation is indicated in red. The lines are analytical calculations and the dots results from numerical simulations. **B.** Fraction $\chi_\male$ of surviving males as a function of the maximal achievable reaction norm κ^* when considering $c = c^*$, shown here for three different values of the mutational variance σ_M^2. For instance, with $\sigma_M^2 = 0.1$ and $\kappa^* = 0.5$ (the values used in all other figures), $\chi_\male \simeq 0.5$, which indicates that half of the males are expected to reach maturation at any given generation, a survival rate that is easily sustained in finite-size populations.

C. Two-fold cost of males

Our results show that dioecious populations can benefit from sexual dimorphism in plasticity. This benefit is quantified by a supplement of growth rate given by $\Delta\Lambda = \Lambda(\kappa_\female = \kappa^*, \kappa_\male = 0) - \Lambda(\kappa_\female = \kappa^*, \kappa_\male = \kappa^*)$. Compared to monoecious populations, however, dioecious populations incur a two-fold cost per generation [7], which corresponds to a growth rate difference of $\ln 2$ (Methods). This cost is therefore offset if $\Delta\Lambda > \ln 2$. In general, $\Delta\Lambda$ increases with c, the extent of directional selection, and offsetting the two-fold cost of dioecy through dimorphic plasticity therefore requires c to exceed a threshold c^* (Fig. 3A). This is not, however, the only constraint: the survival rate of non-plastic individuals, which decreases with increasing c, must also be taken into account. With the optimal reaction norms $\kappa_\female = \kappa^*$ and $\kappa_\male = 0$, this corresponds to the fraction $\chi_\male$ of males reaching maturation becoming smaller and smaller as c becomes larger. Given a population size N, it is however necessary to have $\chi_\male N > 1$ or no male is expected to be available for mating with females [5]. We find that provided that κ^* is sufficiently large and σ_M^2 sufficiently low, the survival rate $\chi_\male$ of non-plastic males taken at $c = c^*$ is of the order $\chi_\male \sim 1/2$ (Fig. 3B), thus indicating that conditions where the two-fold cost of males can be overcome through dimorphic plasticity even in populations of moderate sizes.

D. Greater male variability

As the phenotypic variance (the variance of ϕ among mature individuals of a same generation) is a decreasing function of reaction norms, non-plastic males have higher phenotypic variance than plastic females (Methods). We assumed, however, that all members of a same generation experience the same environment x_t. A more realistic assumption is that they experience a common average environment x_t with fluctuations that are normally distributed with a finite variance σ_e^2 (fluctuations in the mutational bias, on the other hand, simply amount to a larger value of σ_M^2). Considering that each individual experience a particular environment $y_t = x_t + \epsilon$ with $\epsilon \sim \mathcal{N}(\sigma_e^2)$ changes both the probability to reach maturation, which becomes $S(\phi, y_t) = e^{-(\phi - y_t)^2/(2\sigma_S^2)}$, and the genotype-to-phenotype map, which becomes $\phi = (1 - \kappa_\bullet)\gamma + \kappa_\bullet y_t + \xi$ with $\xi \sim \mathcal{N}(\sigma_d^2)$, where σ_d^2 represents the contribution to the developmental variance that is independent of the environment. This more general model can,

however, be mapped to the previous model by considering a sex-specific effective developmental variance, $\sigma_{D,\bullet}^2 = \sigma_d^2 + (1 - \kappa_\bullet)^2 \sigma_e^2$. Our results are robust to this generalization, including the fact that dimorphic plasticity can evolve where males have a more limited phenotypic plasticity associated with a greater phenotypic variability (Fig. S2).

IV. DISCUSSION

We presented a mathematical model for the evolution of phenotypic plasticity in dioecious populations which led us to the following conclusions:

(1) Sexual dimorphism in phenotypic plasticity can be optimal for long-term population growth when selection is directional, either because of a directionally varying environment or a systematic mutational bias. The root of the asymmetry is the limited fertility of one of the two sexes, conventionally females. The optimum then corresponds to females being maximally plastic (with a high reaction norm) and to males being non-plastic (with no reaction norm).

(2) The evolution of plasticity in panmictic populations leads to maximal plasticity in the two sexes, i.e., a suboptimal long-term growth rate, which is interpreted as arising from sexual selection or, more precisely, from intra-male competition as our model assumes no female choice.

(3) Limiting mating to within groups of sufficiently small sizes can counter-act sexual selection and enable the evolution of sexual dimorphism in phenotypic plasticity.

(4) Sexual dimorphism in phenotypic plasticity can confer an evolutionary benefit of dioecy over monoecy despite the two-fold cost of males.

(5) As a smaller reaction norm implies larger phenotypic variance, and even more so when environmental differences are present within a generation, the sexual dimorphism in phenotypic plasticity that evolves implies a greater male variability.

The first point was demonstrated in previous works [4, 5] but the conditions under which dimorphic plasticity may evolve was not previously established. Here we show that limiting mating to within groups is essential to counter-act the adverse effect of intra-male competition and thus resolve a conflict between levels of selection in favor of the highest, population level. At variance with most other explanations of sexual dimorphism, sexual dimorphism in phenotypic plasticity thus arises in our model not because but in spite of sexual selection.

A greater male variability is commonly observed in many species, as for instance reviewed in [13]. Several alternative explanations have been proposed [14], including scenarios involving a stronger selection on males [15], consistent with our results where plasticity evolves to buffer selection on females. Other scenarios have also been proposed to explain how two sexes may provide an adaptive advantage that offsets the two-fold cost of males [16–23]. All these scenarios rely on the same fundamental asymmetry, differential mating success. As most previous models on the subject [19–21], we have only shown here that sexually dimorphic plasticity may provide a adaptive advantage for dioecy over monoecy but not examined the conditions under which dioecy may evolve from monoecy. While this question is beyond the scope of the present work, we note that the few available models demonstrating the evolution of dioecy [22, 23] assume an additional asymmetry between the two sexes, namely an asymmetry in the strength at which the two sexes are selected, with males being subject to a stronger selection than females. In our model, the only asymmetry is in differential reproductive investment – the defining distinction between males and females. We find that the two sexes evolve to experience differential selective pressure, but this is an emergent feature arising from the evolution of sexual dimorphism in phenotypic plasticity, not an a priori assumption. Our scenario thus most closely resembles the informal theory of Geodakyan [16] although he did not anticipate the antagonistic effect that sexual selection may have. This effect imposes a strong constraint: mating must occur within groups of sufficiently closely related individuals for dimorphic plasticity to evolve as a plastic male within a group of non-plastic males has always a higher chance of reproducing. More generally,

we note that the same mechanism by which sexual dimorphism evolves through group/kin selection in our model is at play in other models for the evolution of biological symmetry breaking [24]. This bolsters the hypothesis that the evolution of two sexes might instantiate a more general evolutionary phenomenon [25].

Acknowledgements – We thank Nobuto Takeuchi for stimulating discussions.

METHODS

A.   Dynamics for large unstructured populations

In the limit of large population sizes and in absence of group structure, the long-term growth rate Λ of the population can be obtained analytically by solving a recursion for the density $n_t(\Gamma)$ of newborn individuals at generation t with genotype $\Gamma = (\gamma, y_\female, y_\male)$, where $n_t(\Gamma) = N_t(\Gamma)/N_t$ if $N_t(\Gamma)$ is the mean number of individuals born at generation t with genotype Γ and $N_t = \int d\Gamma N_t(\Gamma)$ the mean total population size.

First, the mean number $M_{\bullet,t}(\Gamma)$ of individuals of sex $\bullet = \female$ or $\male$ that reach maturation is

$$M_{\bullet,t}(\Gamma) = \frac{1}{2} \int d\Phi \ S(\Phi, x_t) D_\bullet(\Phi|\Gamma, x_t) N_t(\Gamma) \tag{6}$$

where $D_\bullet(\Phi|\Gamma, x_t)$ is the probability that an individual of sex $\bullet$ and genotype Γ acquires a phenotype $\Phi = (\phi, y_\female, y_\male)$ and $S(\Phi, x_t)$ is the probability that it survives to reach maturation, given an environment x_t. The factor $1/2$ accounts for the fact that sexes are assigned randomly with same probability. This may be rewritten in terms of densities as

$$m_{\bullet,t}(\Gamma) = \frac{1}{\chi_{\bullet,t}} \int d\Phi \ S(\Phi, x_t) D_\bullet(\Phi|\Gamma, x_t) n_t(\Gamma) \tag{7}$$

where $m_{\bullet,t}(\Gamma) = M_{\bullet,t}(\Gamma)/M_{\bullet,t}$ with $M_{\bullet,t} = \int d\Gamma M_{\bullet,t}(\Gamma)$ is the density of mature individuals of sex $\bullet$ at generation t, and where $\chi_{\bullet,t}$ represents the fraction of individuals of sex $\bullet$ reaching maturation, which is given by

$$\chi_{\bullet,t} = \int d\Gamma d\Phi \ S(\Phi, x_t) D_\bullet(\Phi|\Gamma, x_t) n_t(\Gamma). \tag{8}$$

Second, random mating between individuals of the opposite sex lead to a density of newborn individuals with genotype Γ at generation $t+1$ given by

$$n_{t+1}(\Gamma) = \int d\Gamma^\female d\Gamma^\male \ H(\Gamma|\Gamma^\female, \Gamma^\male) m_{\male,t}(\Gamma^\male) m_{\female,t}(\Gamma^\female). \tag{9}$$

where $H(\Gamma|\Gamma^\female, \Gamma^\male)$ is the probability that a couple with genotypes $\Gamma^\female$ and $\Gamma^\male$ produces an individual of genotype Γ.

The total number of mating events at generation t is $\min(q_\female M_{\female,t}, q_\male M_{\male,t})$ where $M_{\bullet,t} = \int d\Gamma M_{\bullet,t}(\Gamma)$ is the mean total of individuals of sex $\bullet$ that reach maturation. Growth is thus limited by females if $q_\female \chi_{\female,t} < q_\male \chi_{\male,t}$ and by males otherwise. If for instance females are the limiting sex, the growth increase at generation t is

$$\ln \frac{N_{t+1}}{N_t} = \ln \frac{N_{t+1}}{M_{\female,t}} + \ln \frac{M_{\female,t}}{(N_t/2)} - \ln 2 = \ln q_\female + \ln \chi_{\female,t} - \ln 2 \tag{10}$$

and the same expression with $\male$ instead of $\female$ if males are the limiting sex.

Finally, for directionally varying selective pressures, $\chi_{\female,t}$ and $\chi_{\male,t}$ asymptotically reach constant values $\chi_{\female}$ and $\chi_{\male}$ so that the long-term population growth is given by

$$\Lambda = \min\left(\ln(q_{\female}\chi_{\female}), \ln(q_{\male}\chi_{\male})\right) - \ln 2. \tag{11}$$

For monoecious (hermaphroditic) populations, the same formulas apply but with $m_{\female,t}(\Gamma) = m_{\male,t}(\Gamma) = m_t(\Gamma)$ and without the term $\ln 2$ in Eq. (11). This term corresponds to the two-fold cost of males when fertility is limited by females and, more generally, to the two-fold cost of dioecy.

In our model, all the kernels are Gaussian, that is, if we denote Gaussian functions by $G_{\sigma^2}(z) = e^{-z^2/(2\sigma^2)}/(2\pi\sigma^2)^{1/2}$,

$$S(\Phi, x_t) = (2\pi\sigma_S^2)^{1/2} G_{\sigma_S^2}(\phi - x_t), \tag{12}$$

$$D_{\bullet}(\Phi|\Gamma, x_t) = G_{\sigma_D^2}(\phi - (1-\kappa)\gamma - \kappa x_t) G_0\left(\kappa - \frac{\kappa^*}{1 + e^{y_\bullet}}\right), \tag{13}$$

$$H(\Gamma|\Gamma^{\female}, \Gamma^{\male}) = G_{\sigma_M^2}\left(\gamma - \frac{\gamma^{\female} + \gamma^{\male}}{2}\right) \prod_{\bullet = \female, \male} G_{\sigma_V^2}\left(y_\bullet - \frac{y_\bullet^{\female} + y_\bullet^{\male}}{2}\right). \tag{14}$$

Here $G_0(z) = \delta(z)$ represents Dirac delta function.

B. Analytical calculations

We consider here the model with $\sigma_V^2 = 0$ and fixed values of $\kappa_\bullet$ for which the genotypes Γ and phenotypes Φ reduce to scalar variables γ and ϕ. We treat the general case where $\lambda_\bullet$ may differ from $1 - \kappa_\bullet$ so that

$$D_\bullet(\phi|\gamma, x_t) = G_{\sigma_{D,\bullet}^2}(\phi - \lambda_\bullet\gamma - \kappa_\bullet x_t). \tag{15}$$

For the sake of generality, we also allow for the possibility that σ_D^2, c_M and σ_M^2 are sex-dependent. We otherwise adhere to the definition of the model given in the main text. The derivation below follows the solution that we previously provided for a more general model [5].

As the densities $n_t(\gamma)$ and $m_{\bullet,t}(\gamma)$ are asymptotically normally distributed we make the following Gaussian Ansätze to solve Eqs. (7) and (9):

$$n_{\bullet,t}(\gamma) = G_{\varsigma_{\bullet,t}^2}(\gamma - u_{\bullet,t}), \quad m_{\bullet,t}(\gamma) = G_{\varrho_{\bullet,t}^2}(\gamma - v_{\bullet,t}). \tag{16}$$

This leads to

$$\chi_{\bullet,t} = (2\pi\sigma_S^2)^{1/2} G_{\sigma_S^2 + \sigma_{D,\bullet}^2 + \lambda_\bullet^2 \varsigma_{\bullet,t}^2}(\lambda_\bullet u_{\bullet,t} - (1-\kappa_\bullet)x_t) \tag{17}$$

$$v_{\bullet,t} = \alpha_{\bullet,t} u_{\bullet,t} + (1 - \alpha_{\bullet,t})\lambda_\bullet(1 - \kappa_\bullet)x_t \tag{18}$$

$$\varrho_{\bullet,t}^2 = \alpha_{\bullet,t}\varsigma_{\bullet,t}^2 \tag{19}$$

$$\alpha_{\bullet,t} = \frac{\sigma_S^2 + \sigma_{D,\bullet}^2}{\sigma_S^2 + \sigma_{D,\bullet}^2 + \lambda_\bullet^2 \varsigma_{\bullet,t}^2} \tag{20}$$

and

$$u_{\bullet,t+1} = (v_{\female,t} + v_{\male,t})/2 - c_{M,\bullet}, \tag{21}$$

$$\varsigma_{\bullet,t+1}^2 = (\varrho_{\female,t}^2 + \varrho_{\male,t}^2)/4 + \sigma_{M,\bullet}^2 \tag{22}$$

To accommodate for sex-specific genetic parameters, it is convenient to introduce

$$u_{\female,t} = u_{\bullet,t} + c_{M,\bullet} \tag{23}$$

$$\varsigma_{\female,t}^2 = \varsigma_{\bullet,t}^2 - \sigma_{M,\bullet}^2 \tag{24}$$

which are independent of $\bullet$ and follow the recursion

$$u_{\varphi,t+1} = \frac{1}{2}\left(\alpha_\varphi(u_{\varphi,t} - c_\varphi) + \alpha_\sigma(u_{\varphi,t} - c_\sigma)\right) + \left[(1 - \alpha_\varphi)\lambda_\varphi(1 - \kappa_\varphi) + (1 - \alpha_\sigma)\lambda_\sigma(1 - \kappa_\sigma)\right]x_t. \tag{25}$$

$\varsigma_{\varphi,t}^2$ reaches a fixed point ς_φ^2 when $t \to \infty$ that is solution of the cubic equation

$$\varsigma_\varphi^2 = \frac{1}{4}\left(\alpha_\varphi(\varsigma_\varphi^2 + \sigma_{M,\varphi}^2) + \alpha_\sigma(\varsigma_\varphi^2 + \sigma_{M,\sigma}^2)\right) \tag{26}$$

with

$$\alpha_\bullet = \frac{1}{1 + \lambda_\bullet^2\eta_\bullet(\varsigma_\varphi^2 + \sigma_{M,\bullet}^2)/\sigma_S^2}, \qquad \eta_\bullet = \frac{\sigma_S^2}{\sigma_S^2 + \sigma_{D,\bullet}^2}. \tag{27}$$

Finally, if we assume for instance that females are the limiting sex we have $\Lambda = \ln(q_\varphi\chi_\varphi/2)$ with

$$\ln\chi_\varphi = \lim_{t\to\infty}\ln\chi_{\varphi,t} \tag{28}$$

$$= \frac{1}{2}\ln\frac{\sigma_S^2}{\sigma_S^2 + \sigma_{D,\varphi}^2 + \lambda_\varphi^2\varsigma_\varphi^2} - \frac{1}{2}\frac{\lim_{t\to\infty}(\lambda_\varphi u_{\varphi,t} - (1 - \kappa_\varphi)x_t)^2}{\sigma_S^2 + \sigma_{D,\varphi}^2 + \lambda_\varphi^2\varsigma_\varphi^2}$$

$$= \frac{1}{2}\ln(\alpha_\varphi\eta_\varphi) - \frac{\alpha_\varphi\eta_\varphi\lambda_\varphi^2}{2\sigma_S^2}\lim_{t\to\infty}(u_{\varphi,t} - c_{M,\varphi} - (1 - \kappa_\varphi)/\lambda_\varphi x_t)^2$$

Here note that we can always assume $c_E = 0$ by redefining $c_{M,\varphi}$ as $c_{M,\varphi} + (1 - \kappa_\varphi)/\lambda_\varphi c_E$. We are therefore left with the evaluation of $\lim_{t\to\infty}(u_{\varphi,t} - c_{M,\varphi})^2$ where $u_{\varphi,t}$ is given by

$$u_{\varphi,t+1} = \alpha u_{\varphi,t} - \beta = \alpha^t u_{\varphi,0} - \beta\frac{1 - \alpha^t}{1 - \alpha} \tag{29}$$

with

$$\alpha = \frac{\alpha_\varphi + \alpha_\sigma}{2}, \qquad \beta = \frac{\alpha_\varphi c_{M,\varphi} + \alpha_\sigma c_{M,\sigma}}{2} \tag{30}$$

so that

$$\lim_{t\to\infty}(u_{\varphi,t} - c_{M,\varphi})^2 = \left(\frac{\beta}{1 - \alpha} + c_{M,\varphi}\right)^2 \tag{31}$$

Finally, considering $c_{M,\varphi} = c_{M,\sigma} = c_M$ and a non-zero c_E, we obtain Λ as given by Eq. (11) with

$$\ln\chi_\bullet = \frac{1}{2}\ln(\alpha_\bullet\eta_\bullet) - \frac{\alpha_\bullet\eta_\bullet(\lambda_\bullet c_M + (1 - \kappa_\bullet)c_E)^2}{2(1 - \alpha)^2\sigma_S^2} \tag{32}$$

C. Phenotypic variance

If $n_t(\gamma)$ is normally distributed, the phenotypes ϕ of mature individuals of sex $\bullet$ are also normally distributed with a distribution $m_{\bullet,t}(\phi)$ given up to a normalizing factor by

$$m_{\bullet,t}(\phi) \propto S(\phi, x_t)\int d\gamma D_\bullet(\phi|\gamma, x_t)n_{\bullet,t}(\gamma). \tag{33}$$

The phenotypic variance is therefore

$$\mathrm{Var}_\bullet(\phi) = \left(\sigma_S^{-2} + \left(\sigma_D^2 + (1 - \kappa_\bullet)^2\varsigma_{\bullet,t}^2\right)^{-1}\right)^{-1} \tag{34}$$

When the mutational variance σ_M^2 is independent of the sex, the genetic variances $\varsigma_{\bullet,t}^2$ reach a fixed point ς^2 that is also independent of the sex $\bullet$. In this case, the smaller the reaction norm, the larger the phenotypic variance. In presence of intra-generation environmental fluctuations, σ_D^2 is replaced by $\sigma_D^2 = \sigma_d^2 + (1 - \kappa_\bullet)^2\sigma_e^2$, which does not change this conclusion.

D. Numerical simulations

As in our previous works [5], numerical simulations are performed with a population of fixed size N. Genotypes Γ are initially identical for all members of the population and a phenotype Φ is assigned to each individual based on its genotype Γ and the environment x_t as described in the main text. Each individual is then selected with probability $S(\phi, x_t) = e^{-(\phi-x_t)^2/(2\sigma_S^2)}$. At generation t, this results in $M_{\female,t}$ mature females and $M_{\male,t}$ mature males. The total number of mating events is then $N_t = \min(q_{\female}M_{\female,t}, q_{\male}M_{\male,t})$. If for instance $q_{\female}M_{\female,t} < q_{\male}M_{\male,t}$, each female is involved in exactly $q_{\female}$ mating events and males are sampled at random without replacement in the set of size $q_{\male}M_{\male,t}$ where each male is represented $q_{\male}$ times or, if $q_{\male} = \infty$, sampled at random with replacement in the set of males. The same procedure is applied with the two sexes playing opposite roles if $q_{\female}M_{\female,t} > q_{\male}M_{\male,t}$. Each mating event leads to a newborn that inherits a genotype Γ based on the genotypes $\Gamma^{\female}$ and $\Gamma^{\male}$ of its parents following the rules given in the main text. The population at the next generation $t+1$ is then obtained by randomly sampling N of the N_t newborns. N may be larger than N_t, but we restrict the study to parameters where $N_t > 0$ at each generation t. After T generations, the long-term growth rate is estimated as $\Lambda_{N,T} = (1/T) \sum_{t=1}^{T} \ln(N_t/N)$.

[1] T D Price, A Qvarnström, and D E Irwin. The role of phenotypic plasticity in driving genetic evolution. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270(1523):1433–1440, 2003.

[2] L-M Chevin, R Lande, and G M Mace. Adaptation, Plasticity, and Extinction in a Changing Environment: Towards a Predictive Theory. *PLoS Biology*, 8(4):e1000357, 04 2010.

[3] R C Stillwell, W U Blanckenhorn, T Teder, G Davidowitz, and C W Fox. Sex differences in phenotypic plasticity affect variation in sexual size dimorphism in insects: from physiology to evolution. *Annual review of entomology*, 55(1):227 – 245, 00 2010.

[4] S Hangartner, C M. Sgrò, T Connallon, and I Booksmythe. Sexual dimorphism in phenotypic plasticity and persistence under environmental change: An extension of theory and meta?analysis of current data. *Ecology Letters*, 25(6):1550–1565, 2022.

[5] A S. Zadorin and O Rivoire. Sex as information processing: Optimality and evolution. *Physical Review E*, 103(6):062413, 2021.

[6] J Merilä and A P Hendry. Climate change, adaptation, and phenotypic plasticity: the problem and the evidence. *Evolutionary applications*, 7(1):1–14, 2014.

[7] J Maynard-Smith. *The evolution of sex*, Cambridge University Press Cambridge, 1978.

[8] B Charlesworth. Directional selection and the evolution of sex and recombination. *Genetical research*, 61(3):205 – 224, 06 1993.

[9] O Rivoire and S Leibler. A model for the generation and transmission of variations in evolution. *Proceedings of the National Academy of Sciences of the United States of America*, 111(19):E1940 – 9, 05 2014.

[10] S Karlin and J McGregor. Towards a theory of the evolution of modifier genes. *Theoretical Population Biology*, 5(1):59–103, 1974.

[11] D S Wilson. A theory of group selection. *Proceedings of the National Academy of Sciences*, 72(1):143–146, 1975.

[12] N Takeuchi, N Mitarai, and K Kaneko. A scaling law of multilevel evolution: how the balance between within- and among-collective evolution is determined. *Genetics*, 220(2):iyab182–, 2021.

[13] T P Hill. An Evolutionary Theory for the Variability Hypothesis. *arXiv*, 2017.

[14] S RK Zajitschek, F Zajitschek, R Bonduriansky, R C Brooks, W Cornwell, D S Falster, M Lagisz, J Mason, A M Senior, D WA Noble, and S Nakagawa. Sexual dimorphism in trait variability and its eco-evolutionary and statistical implications. *eLife*, 9:e63170, 2020.

[15] A. Pomiankowski and A P Møller. A resolution of the lek paradox. *Proceedings of the Royal Society of*

London. Series B: Biological Sciences, 260(1357):21–29, 1995.

[16] VA Geodakyan. Role of the sexes in the transmission and transformation of genetic information. *Problemy Peredachi Informatsii*, 1(1):105–112, 1965.

[17] R L Trivers. Sexual Selection and Resource-Accruing Abilities in Anolis garmani. *Evolution*, 30(2):253, 1976.

[18] A Kodric-Brown and J H. Brown. Anisogamy, sexual selection, and the evolution and maintenance of sex. *Evolutionary Ecology*, 1(2):95–105, 1987.

[19] J T Manning. Males and the advantage of sex. *Journal of Theoretical Biology*, 108(2):215–220, 1984.

[20] S Siller. Sexual selection and the maintenance of sex. *Nature*, 411(6838):689–692, 2001.

[21] A F Agrawal. Sexual selection and the maintenance of sexual reproduction. *Nature*, 411(6838):692–695, 2001.

[22] L Hadany and T Beker. Sexual selection and the evolution of obligatory sex. *BMC Evolutionary Biology*, 7(1):245, 2007.

[23] D Roze and S P Otto. Differential selection between the sexes and selection for sex. *Evolution*, 66(2):558–574, 2012.

[24] N Takeuchi and K Kaneko. The origin of the central dogma through conflicting multilevel selection. *Proceedings. Biological sciences*, 286(1912):20191359, 10 2019.

[25] VA Geodakian. Evolutionary theories of asymmetrization of organisms, brain and body. *Uspekhi fiziologicheskikh nauk*, 36(1):24–53, 2005.

FIG. S1: Scaling of the maximal group size below which dimorphic plasticity evolves – The calculations performed to obtain Figure 2D are repeated here with either different total population sizes N (A-B) or different mutational variances for the modifier genes σ_V^2 (C-D). **A.** As the total population size N increases, the maximal group size K^* below which dimorphic plasticity evolves increases. **B.** This maximal group size K^* scales as $K^* \sim N^{1/2}$. **C.** As the mutational variance σ_V^2 increases, the maximal group size K^* below which dimorphic plasticity evolves decreases. **B.** This maximal group size K^* scales as $K^* \sim \sigma_V^{-1/2}$. Only average male reactions norms are shown here as female reaction norms evolve to maximal values in all cases, and results are averaged over 10 trajectories instead of 100 trajectories in Figure 2D, which explains the larger error bars.

FIG. S2: Evolution of plasticity in presence of intra-generation environmental fluctuations – The parameters and calculations are as in Figure 2D, except that each individual at generation t experiences a local environment $y_t = x_t + \epsilon$ where ϵ is normally distributed with zero mean and variance σ_e^2. As a consequence, the probability to reach maturation is $S(\phi, y_t)$ instead of $S(\phi, x_t)$ and the phenotypes are related to genotypes by $\phi = (1 - \kappa_\bullet)\gamma + \kappa_\bullet y_t$ instead of $\phi = (1 - \kappa_\bullet)\gamma + \kappa_\bullet x_t$. Here we consider $c_M = 0$ and therefore $c = c_E$. The results for $\sigma_e^2 = 0$ correspond exactly to those of Figure 2D. For sufficiently small σ_e^2, the results are qualitatively unchanged: sexual dimorphism in plasticity evolves when mating occurs within groups of sufficiently small size K. Large values of σ_e^2, however, tend to disfavor the evolution of dimorphic plasticity.