Wings of Change: aPKC/FoxP–dependent plasticity in steering motor neurons underlies operant self–learning in *Drosophila*

Andreas Ehweiner[1], Carsten Duch[2] and Björn Brembs[1]*

1 Universität Regensburg, Institut für Zoologie - Neurogenetik, Regensburg, Ger-
many.

2 Johannes Gutenberg Universität Mainz, Institute of Developmental Biology and
Neurobiology (iDN), Mainz, Germany

* bjoern@brembs.net

9 Abstract

Background: Motor learning is central to human existence, such as learning to
speak or walk, sports moves, or rehabilitation after injury. Evidence suggests that
all forms of motor learning share an evolutionarily conserved molecular plasticity
pathway. Here, we present novel insights into the neural processes underlying
operant self-learning, a form of motor learning in the fruit fly *Drosophila*.

Methods: We operantly trained wild type and transgenic *Drosophila* fruit flies,
tethered at the torque meter, in a motor learning task that required them to initi-
ate and maintain turning maneuvers around their vertical body axis (yaw torque).
We combined this behavioral experiment with transgenic peptide expression,
CRISPR/Cas9-mediated, spatio-temporally controlled gene knock-out and confo-
cal microscopy.

Results: We find that expression of atypical protein kinase C (aPKC) in direct wing
steering motoneurons co-expressing the transcription factor *FoxP* is necessary for
this type of motor learning and that aPKC likely acts via non-canonical pathways.
We also found that it takes more than a week for CRISPR/Cas9-mediated knockout
of *FoxP* in adult animals to impair motor learning, suggesting that adult *FoxP* ex-
pression is required for operant self-learning.

Conclusions: Our experiments suggest that, for operant self-learning, a type of motor learning in *Drosophila*, co-expression of atypical protein kinase C (aPKC) and the transcription factor *FoxP* is necessary in direct wing steering motoneurons. Some of these neurons control the wing beat amplitude when generating optomotor responses, and we have discovered modulation of optomotor behavior after operant self-learning. We also discovered that aPKC likely acts via non-canonical pathways and that *FoxP* expression is also required in adult flies.

1. Introduction

Motor learning is an essential component of human behavior and ubiquitous throughout the animal kingdom. The process of learning a motor skill can be influenced by a number of factors, such as the amount of training, type of feedback, or the presence/absence of environmental cues. Regaining lost motor functions after brain or spinal cord injury is considered a crucial component of rehabilitation. Human language is acquired by a form of motor learning, and motor learning also appears to be a key invention that allowed the newly evolved animals in the Cambrian to become ambulatory (1–3). Vocal learning, such as learning to speak, is a form of motor learning (4) that involves the Forkhead Box transcription factor family P (FoxP) in vertebrates (5–15). Other, nonvocal forms of motor learning also involve FoxP genes in vertebrates (16–19) and invertebrates (20). Vocal learning also shares the involvement of protein kinase C (PKC) with other forms of motor learning (21–25), raising the possibility of a conserved motor learning

pathway extending beyond these two components.

While motor learning shares various features with other forms of learning, such as operant learning producing habits or skill-learning, it is debated how many common biological mechanisms these different concepts share (26–29). Understanding motor learning in a numerically smaller nervous system in a genetically tractable organism where one can not only study the motor learning process itself, but also its interactions with other forms of learning (30), may help inform these debates.

Here we provide further evidence about the specific manner in which FoxP and PKC are involved in a form of motor learning in the fruit fly *Drosophila*, operant self-learning at the torque meter. In this experiment, motor learning dissociates from other forms of learning such that genes involved in motor learning are not involved in other forms of learning and *vice versa* (31–33). At the torque meter, a fly is tethered between head and thorax such that it can move all other appendages. When beating its wings, the fly generates forces, some of which can be measured by the torque meter. Specifically, the torque meter measures torque around the vertical body axis, yaw torque (34). Even in the absence of any guiding cues, flies can learn to associate one torque domain (e.g.,

roughly corresponding to left or right, respectively, turning maneuvers) with a punishing heat beam (35). This experiment not only conceptually mimics other motor learning paradigms in that feedback is made immediately contingent on specific motor actions, but also via its dependence on *FoxP* and PKC genes (20,21). This form of motor learning has been termed operant self-learning to distinguish it from other forms of operant learning and to denote that the subject is learning about its own behavior, as opposed to some stimulus associated with the behavior (32).

For operant self-learning in *Drosophila*, it is not known in which neurons the *FoxP* gene is required and which PKC gene is involved. It is also unknown which pathway is engaged by PKC and whether FoxP expression is also required acutely in adult flies for this form of motor learning. In this work, we addressed all three research questions.

2. Methods

2.1. Strains and fly rearing

If not stated otherwise, flies (table 1) were raised on standard cornmeal/molasses medium (36) at 25°C and 60% humidity under a 12-hour light/dark cycle. For experiments requiring the expression of temperature-sensitive Gal80, animals were raised at 18°C. To set

up crosses for behavioral experiments, 20 females were placed together with five to eight males and were allowed to lay eggs for 24 h. They were flipped daily into fresh vials, to ensure appropriate larval density. In 30 years of research on learning and memory in tasks like ours, no difference was ever observed between male and female flies in terms of learning ability. Whenever genetically appropriate, we used female flies for practical reasons. The flies were prepared the day before the experiment, allowing them time to recover. Female flies (24 to 48 h old) were briefly immobilized under cold anesthesia. A thin triangular copper hook (0,05 mm diameter) was glued (3m Espe Sinfony, 3M Deutschland GmbH) between head and thorax, fixing both body parts to each other (37). Each animal was kept individually in a small moist chamber with a few grains of sugar. For $tub\text{-}Gal80^{ts}$ expression, animals were raised at 18°C and incubated at 30°C for two days. Experiments were always conducted at room temperature. For experiments using the gene-switch system, newly hatched flies were placed on *Drosophila* instant medium (351.204, Schlüter Biologie, Eutin-Neudorf, Germany) containing the steroid hormone RU486 (200 µg/ml, CAS No.: 84371-65-3, Sigma-Aldrich, St. Louis, MO) for two days.

The UAS-g-KIBRA line was generated in the laboratory of Michael Krahn, Universität

148 Münster. They cloned the following guide RNAs
149 from KIBRA Intron-2: GTACTTAC-
150 GACTGCTTCGAC and KIBRA Intron-4: GGGCAC-
151 CGTGCAGATCAGCA in pCDF6 and inserted
152 them in attP40.
153

Table 1: Fly strains used in this work

Genotype	use	Bloomington	Flybase
;;nSyb-GS	pan-neuronal driver	80699	FBti0201287
;;GMR65A06-GAL4	protocerebral bridge driver	39330	FBti0137511
;;GMR20H05-GAL4	central complex driver	47896	FBti0133817
;;ato-Gal4	driver for dorsal cluster neurons		
ELAV-Gal4;;	pan-neuronal driver		
ELAV-Gal4;Tub-Gal80ts;;	temperature-sensitive pan-neuronal driver		
;;FoxP-iB-Gal4/TM3	driver for FoxP-iB neurons		
;;FoxP-LexA	driver for all FoxP neurons		
;;UAS-g-aPKC	effector	85862	FBti0210993
;;UAS-g-BAZ	effector	84234	FBti0207133
;;UAS-g-PKC53e	effector	76612	FBti0194968
;UAS-g-KIBRA	effector		
;;UAS-Cas9;	effector		
;;UAS-PKCi	effector	4589	FBti0010565
;;UAS-t:gRNA(4xFoxP)	effector		
;UAS-Cas9;;	effector		
C380-Gal4;;	MN driver	80580	FBti0016294
;;D42-Gal4	MN driver	8816	FBti0002759
y[1] w[*]; Mi{Trojan GAL4.un} aPKC[MI10848-TG4.un]/SM6a	driver for aPKC cells	77814	FBti0196316
LexAop-mD8-RFP-UAS-mCD8-GFP;TM3/TM6b	effector		
Wild type Berlin	wild type strain		

2.2. Behavioral setups

Two different torque meters and setups had to be used for technical reasons, not by choice. Both torque meters were described previously: The 'Tang' meter (38) and the 'Götz' meter (34)(RRID:SCR_017276), named after the authors. With flies attached to the torque meter via a clamp, all devices measure the rotational force (torque) around the animal's vertical body axis. While the Götz meter is older than the Tang meter, it is technically more advanced because of its rotational compensation, and was included in a more modern version of the setup that was used in the later experiments, after the first data were collected using the Tang setup. Even later, the wild type experiments shown in Figure 4 were conducted with a third torque meter, still in the prototype phase, which combines the laser-based measurement of the Tang meter with induction-based compensation of the Götz meter. Documentation of this setup is in preparation, but the operating principles remain the same as for the two referenced devices. In all setups, the animal is surrounded by a cylindrical panorama (arena; diameter 58 mm Tang, 90 mm other setups), homogeneously illuminated from behind by either a projector (new setups: DLPLCR4500EVM, Texas Instruments) or a halogen lamp (Tang: OSRAM 100W/12V),

such that stationary flight in a controlled environment was achieved. An infrared laser (Stocker Yale Lasiris SNF series; 825 nm, 150 mW) was used as punishment in all setups. The laser was pointed from above onto the animal's head, pulsed (approximately 200 ms pulse width ~4 Hz) and the intensity adjusted empirically for maximal heat avoidance and learning. The experiment is fully computer controlled, using custom software (Tang: LabView, National Instruments, RRID:SCR_014325. New setups: DOI: 10.5281/zenodo.7102195).

2.3. Design of behavioral experiments

Before each self-learning experiment, the yaw torque range was adjusted using optomotor stimuli for each fly tethered to the torque meter. The optomotor response (OMR) is an innate, orienting behavior evoked by whole-field visual motion and is common to vertebrates and invertebrates. The OMR has algorithmic properties such that the direction of the whole-field coherent motion dictates the direction of the behavioral output (e.g., leftward visual stimuli lead to turning left, and rightward visual stimuli lead to turning right). For instance, when tethered Drosophila are surrounded by vertical black and white grating patterns rotating along the fly's azimuth (i.e, around the fly's vertical body axis), the fly will turn (i.e., produce yaw torque) in the direction

of perceived motion. Typical OMRs for tethered flies responding to horizontally rotating vertical stripes are depicted in Figure 4A1.

For the 'Tang' setup, arena rotation for the optomotor stimulus was operated by hand. The direction of the rotation was reversed after the fly reached its asymptotic optomotor torque. During optomotor presentations before the self-learning experiment, the torque was adjusted to be zero-symmetric. This was to facilitate unbiased torque preferences before training. Torque traces during OM presentations were not stored in experiments using the 'Tang' setup. Optomotor stimuli (15 vertical black stripes on a white background taking about 3.5s for a full rotation, i.e., a pattern wavelength of 24° at a pattern frequency of about 4.3 Hz) were presented for a duration of 30 s in each turning direction for flies in the new setups and recorded in the raw data files together with all other data from each experiment. Because of this difference, as experiments recorded with the Tang device did not record optomotor periods, periods in 'Tang' experiments are numbered from 1 to 9 (Table 2), while optomotor periods for the new setups were included, such that the periods in the new setups were numbered 1-17 (Table 3). Because we used four optomotor periods each before and after training in the new setup, periods 5

to 13 were the periods in which self-learning was studied in this setup.

The main self-learning experiment then consisted of nine periods of two minutes duration in both setups. The laser was permanently off during the first two periods, so that the fly could freely choose its direction of turning maneuvers without any feedback. In the following two training periods either the left or the right torque domain was associated with the punishing laser, without any hysteresis. The punished torque domain was alternated between experiments. The first two training periods were followed by one test period without punishment. Afterwards, the fly was trained again with the same side punished as before for another two 2-min. periods. Finally, no heat was applied in the final two test periods, allowing the fly to express its spontaneous yaw torque preference. The figures always show the preference in the first test period after the last training period, i.e., period 8 (performance index, PI8) in the Tang setup and period 12 (PI12) in the new setups (Tables 2, 3). When the axis labels in the figures differ with regard to PI8 or PI12, these differences only indicate which setup was used. In all cases, the same first test period after the last training period was used to test for learning, irrespective of setup.

Table 2: Experimental sequence "Tang Setup". All periods lasted 120s.

Period 1	Period 2	Period 3	Period 4	Period 5	Period 6	Period 7	Period 8	Period 9
Pretest	Pretest	Training	Training	Test	Training	Training	Test	Test
No heat	No heat	**Heat**	**Heat**	No heat	**Heat**	**Heat**	No heat	No heat

Table 3: Experimental sequence for new setups. OM - optomotor. Torque learning periods (5-13) lasted 120s, while OM periods lasted 30s.

OM before				Experiment									OM after			
Period 1	Period 2	Period 3	Period 4	Period 5	Period 6	Period 7	Period 8	Period 9	Period 10	Period 11	Period 12	Period 13	Period 14	Period 15	Period 16	Period 17
OM	OM	OM	OM	Pretest	Pretest	Training	Training	Test	Training	Training	Test	Test	OM	OM	OM	OM
No heat	No heat	No heat	No heat	No heat	No heat	**Heat**	**Heat**	No heat	Heat	Heat	No heat	No heat	No heat	No heat	No heat	No heat

2.4. Data handling and statistical analysis of behavioral experiments

2.4.1. Data selection

To ensure proper punishment by the laser, each fly was exposed to the laser after the experiment, to ensure it was adjusted correctly. If the fly survived the laser for 15 s or longer, the data were excluded from analysis. Data that did not show any or shifted OMRs, indicating either an unhealthy fly or an error with the measuring device, were also excluded. Data were also excluded if the fly had not ex-

perienced the laser at least once during training. Finally, flies with poor flight performance (constant stopping of flight) were also excluded from analysis. While these data were excluded from analysis, all complete traces are nevertheless included in the published data sets, such that the inclusion criteria can be independently tested.

2.4.2. Data availability and analysis

The preference of a fly for right or left torque domain was quantified as the performance index $PI = (ta − tb)/(ta + tb)$. During training periods, tb indicates the time the fly is exposed to the heat and ta the time without heat. During tests, ta and tb refer to the times when the fly chose the formerly (or subsequently) unpunished or punished situation, respectively. Thus, a PI of 1 indicates the fly spent the entire period in the situation not associated with heat, whereas a PI of −1 indicates that the fly spent the entire period in the situation associated with heat. Accordingly, a PI of zero indicates that the fly distributed the time evenly between heated and non-heated situations and a PI of 0.5 indicates that 90 of the 120 s in that period were spent in the unpunished situation.

Analogously, optomotor (Table 3, OM) behavior was quantified by computing an OM asymmetry index from the torque traces. Straight lines or double sigmoidal models were fitted to individual torque traces from OM periods, depending on the detected slope of the OMR. Each fit was generated separately for each turning direction. From the fitted lines/models, optomotor magnitude was derived as either the intercept (lines) or the asymptote (double sigmoidal model) for each turning direction. The magnitude of left-turning torque was subtracted from the magnitude of right-turning torque and divided by the sum of the two values. This optomotor asymmetry index becomes -1 for OMRs where clockwise ('right-turning') stimuli elicit no or left-turning torque (while counter-clockwise stimuli elicit left-turning torque). It becomes 1 for OMRs where counter-clockwise ('left-turning') stimuli elicit no or right-turning torque (while clockwise stimuli elicit right-turning torque). The OM asymmetry index becomes zero if the absolute magnitudes of OMRs in both directions are equal. In brief, a positive optomotor asymmetry index indicates shifts away from symmetrical torque towards right-turning torque and a negative index indicates shifts towards left-turning torque.

All behavioral data were analyzed using R (R Project for Statistical Computing)

(RRID:SCR_001905). The collection of R-scripts evaluating the time series data can be found at DOI 10.5281/zenodo.10041052 (39). The data model pertaining to the XML raw data files and the YAML data set files can be found at 10.5281/zenodo.7101734 (39). In brief, the XML data files contain both the meta-data for each single fly experiment as well as the time-series data covering the entire experiment. The single YAML file per dataset contains the experimental design, such as which data files belong to which experimental group, the type of statistics to be performed, significance levels used, experimenter comments and data inclusion/exclusion. The main R-Script reads the YAML dataset files and performs the appropriate computations for quality control, analysis and statistics.

Quality control is performed on each single-fly XML file and included in the published datasets. Each single-fly experiment XML file is thus accompanied by a single-fly HTML quality control report sheet containing plots of the raw time series data, as well as a number of evaluations necessary to assess the proper execution of the experiment and the quality of the resulting data.

Data analysis and statistics for each dataset are reported in an HTML dataset evaluation sheet. Thus, a complete dataset consists of one XML raw data file and one HTML quality

control sheet for each fly, plus a single YAML dataset file and one HTML dataset evaluation sheet.

The datasets were published using a custom Python script (DOI: 10.5281/zenodo.7101741)(39) that synchronizes the collected data on the local computer with the University of Regensburg publication server. The persistent identifiers for each dataset are listed in the figure legends.

2.4.3. Statistics

Motivated by ongoing efforts to improve statistical inference in science (e.g., (40–54), we chose to statistically evaluate PIs in two complementary ways, using both frequentist and Bayesian statistics (55–61). Following previous studies (30,31,62–65), individual PIs of the first test after the last training period were tested against zero to evaluate the ability of the manipulated flies to show a preference towards the unpunished torque domain. The rationale behind estimating a group of flies as either showing learning or not is to trade-off statistical power with a more nuanced measure of learning performance: comparing between experimental groups may yield more nuance, but also requires impractically large sample sizes for adequate (>80 %) statistical power. To further reduce the chance of statistical error, we used both Wilcoxon tests in a frequentist sce-

nario and computed the equivalent Bayes Factors for a Bayesian version. We set the alpha value for the Wilcoxon test to 0.5 % as suggested by (46), such that p-values below 0.005 and Bayes factors above 5 for the same group of flies would be considered compelling evidence that the flies were able to learn. Conversely, Bayes Factors below one together with p-values higher than 0.05 were considered evidence that the flies were not able to show self-learning. Finally, groups where the two statistics were in conflict or intermediate, were considered inconclusive. Thus, both the Bayesian and the frequentist criteria had to be met in order to claim that a genetic manipulation interfered with operant self-learning. These criteria were chosen to quantitatively distinguish between effective and ineffective manipulations without neglecting the uncertainty associated with all experimentation. At the same time, our evaluations were chosen to specifically identify large contributions to the learning processes that can be identified with sufficient statistical power (i.e., large effect sizes). All statistical results are published with the raw data and the code used to compute them is openly available.

2.5. Gene expression analysis in flight steering motor neuron terminals

2.5.1. Dissection

The axon terminals on flight steering muscles were tested immunocytochemically for the expression of *FoxP* and aPKC in flies expressing LexAop RFP under the control of FoxP-Lex and UAS td-GFP under the control of aPKC-Trojan GAL4 (LexAop RFP, UAS td-GFP / aPKC-Trojan-GAL4; FoxP-LexA / +). In addition, synapses at neuromuscular junctions were labeled with the active zone marker *bruchpilot* (*brp*, (66)). Animals were dissected in normal saline along the dorsal midline and bent open with minute pins inserted through the dorsalmost edge of the dorsal longitudinal flight muscles (DLMs). The heart, gut, fat tissue, and other connective tissue were removed to expose the ventral nerve cord (VNC) and the musculature. Next, on both sides the DLMs and dorsoventral flight muscles (DVMs) were carefully removed layer-by-layer to expose the direct flight steering muscles, which are located close to the lateral cuticle of the thorax. Preparations were rinsed 5-10 times in saline to remove debris from the dissection procedure, and specimens were fixed for 1 h in 4 % paraformaldehyde in 0.1 M PBS buffer at room temperature (22°C). After fixation specimen paraformaldehyde was exchanged with 0.1 M PBS buffer.

2.5.2. Immunohistochemistry of motor terminals

Following fixation, specimens were washed 6 × 20 min in 0.1 M PBS buffer at room temperature. Next, preparations were washed 3 × 1 h in 0.1 M PBS-Tx (0.3 %) buffer at room temperature and incubated with primary antibodies (mouse α-brp, 1:500 (Hybridoma bank, NC82)); chicken α-GFP (1:1000, Life Technologies A10262), and rabbit α-mCherry, (1:500, PAS-34974)) in 0.1 M PBS-Tx (0.3 %) buffer at 4° C for 24 to 36 h. Following primary antibody incubation, preparations were washed 6 × 1 h in 0.1 M PBS buffer at room temperature. Next animals were incubated in secondary antibodies: donkey α-mouse Alexa 647 (JacksonImmunoResearch 715-605-150), donkey α-chicken Alexa 488 (Dianova 703-545-155), and donkey α-rabbit Alexa 568 (Invitrogen A10042) in 0.1 M PBS-Tx (0.15%) buffer at 4° C for 24 to 36 h. Following secondary antibody incubation, preparations were washed 3 × 1 h in 0.1 M PBS buffer at room temperature, dehydrated in an ascending ethanol series (50, 70, 90, and 2 × 100 % EtOH, 15 min each), and cleared for 5 min in methyl salicylate. Finally, preparations were mounted in methyl salicylate in between two coverslips that were glued onto both sides of a round hole (10 mm diameter) drilled into custom-made metal slides of 188 µm thickness (67). Briefly, one cover slip was fixated with superglue underneath the hole, the space was

filled with methyl salicylate, the preparation transferred into the mounting media, and another coverslip was carefully placed on top of the hole, so that no air remained in the methyl salicylate filled hole. The top coverslip was fixed to the metal slide using transparent nail polish (DM Markt, Mainz, Germany). After 20 min to let the nail polish dry the preparation was transferred to the microscope. In total 11 animals were subjected to immunocytochemistry on direct flight muscles. In 9 of these 11 preparations, we were able to identify only subsets of the direct wing steering muscles under investigation. These subsets yielded identical results with respect to aPKC and FoxP expression as the two preparations with complete sets of direct muscles as shown in Figure 3.

2.5.3. Confocal laser scanning microscopy

Preparations were scanned using a Leica (Leica Microsystems, Germany) SP8 confocal laser scanning microscope (CLSM) under either a 20x oil (NA = 0.75) or a 40x oil (NA = 1.3) lens at an image format of 1024 × 1024 pixels. Z-step size was 2 µm for the 20x lens (resulting in voxel dimensions of 0.57 × 0.57 × 2 µm) and 1 µm for the 40x lens (resulting in voxel dimensions of 0.28 × 0.28 × 1 µm). Alexa 488 was excited with an argon laser at 488 nm and detected with a photomultiplier between 495 and 530 nm wavelength. Alexa 488 was excited

with a solid state laser at 561 nm and detected with a photomultiplier between 570 and 610 nm wavelength. Alexa 647 was excited with a red helium neon laser at 633 nm and detected with a photomultiplier between 640 and 670 nm wavelength. All image stacks were stored as .lei files and further analyzed using Las X software (Leica Microsystems, Germany). Selected fields of view were used for maximum intensity projection views, which were exported as 24 bit three color tiff images and further processed with [Corel Draw 11](#).

2.5.4. Data availability

Confocal image stacks can be found at: 10.5281/zenodo.10606166 (68).

3. Results

3.1. Self-learning requires a*PKC* in *FoxP* motor neurons

Colomb and Brembs (2016) discovered that blocking all protein kinase C (PKC) isoforms in neurons using the inhibitory peptide PKCi abolished operant self-learning. We replicated their results by pan-neuronal expression of PKCi. In one group, PKCi was expressed both during development and in the adult flies. In the other group, PKCi expression was restricted to adulthood only, by using the temperature-sensitive Gal4 inhibitor *tub-Gal80^{ts}* (see Methods for details). Temporally

unrestricted pan-neural expression did not impair self-learning, whereas expression restricted to neurons in adult flies abolished self-learning (Figure 1A). This result may seem surprising, but compensation for experimental manipulations of PKC activity through development has been reported numerous times in the literature (69–73) and our data reproduced our previously published, identical experiments with PKCi (21), demonstrating the PKCi construct is still performing as expected. The same publication also reported that PKCi expression in MNs was sufficient to impair operant self learning. The expression of the driver lines used in Colomb and Brembs overlap with the expression of FoxP in the MNs of the ventral nerve cord (74), so we tested if driving PKCi in neurons expressing the isoform B of FoxP was sufficient to impair operant self-learning. Corroborating the notion that MNs are crucial for operant self-learning, expressing PKCi only in cells expressing isoform B of the FoxP gene also abolished self-learning, even without temporal control, suggesting that PKC activity is required in FoxP-positive MNs of the ventral nerve cord.

Figure 1: For operant self-learning, *aPKC* is required in motor neurons.

Performance indices for the first period after training (PI8/PI12) are plotted. Each colored dot represents a single fly experiment. Red dots denote experiments where the fly was punished on its left turning torque domain, blue dots denote flies that were punished on their right turning domain. Box plots denote medians, quartiles and non-outlier range. Statistical analyses test for differences of PIs against zero.

A. Inhibiting all protein kinase C isoforms with the inhibitory peptide PKCi. Constitutive, pan-neuronal expression PKCi (left, yellow), leads to high PIs and a large Bayes Factor, indicating this manipulation left self-learning intact. Expressing PKCi either in FoxP-isoform B positive neurons without temporal control (middle, green), or in all neurons but restricted to adulthood using *tub-Gal80^{ts}* (right, blue, see Methods for details) yields low PIs, high p-values and low Bayes factors, indicating self-learning was impaired. Data: 10.5283/epub.52958 (75)

B. Pan-neuronal knock-out of two different PKC genes with CRISPR/Cas9 in adulthood (using the GeneSwitch system and feeding RU486 to adult flies, see Methods for details) suggests aPKC is necessary for operant self-learning. Knocking out atypical PKC (yellow, left) yields moderate PIs, p-values and Bayes factors, indicating some effect on operant self-learning, while the high PIs, low p-values and high Bayes factor of the group where PKC53E was knocked out (right, green) indicate their self-learning was intact. Data: 10.5283/epub.52957 (76)

C. Knocking out aPKC in motor neurons or *FoxP*-neurons impairs operant self-learning. Expressing the CRISPR/Cas9 components either in *FoxP* isoform B-positive neurons (green, middle) or in motor neurons (blue, right) leads to low PIs, high p-values and low Bayes Factors, indicating their self-learning is strongly impaired. Control flies with only the CRISPR/Cas9 genetic elements but no driver, showed high PIs, a low p-value and a high Bayes Factor, indicating their self-learning was intact. Data: 10.5283/epub.52944 (77)

D. Bayesian statistics for the three datasets.

To help determine which PKC may be involved in this mechanism, we screened RNA-Seq databases for PKC genes expressed in MNs. Restricting candidates to those where gRNA lines were available for CRISPR/Cas9-mediated gene knockout yielded only two genes: the atypical PKC (aPKC) and the diacylglycerol-activated PKC53E. Knocking out each gene pan-neuronally in adult flies and testing the manipulated animals for operant self-learning showed excellent learning performance in PKC53E-manipulated flies (Figure 1B), clearly ruling out PKC53E as the gene involved in operant self-learning. These results also confirmed prior experiments with PKC53E mutant flies (21), ruling out any PKC53E involvement. In contrast to the PKC53E manipulated flies, the performance of the aPKC-manipulated flies was inconclusive: their preference scores were somewhat lower, not reaching our criteria for significant learning, but at the same time too high to be confident in the result (Figure 1B). To

minimize the possibility of a false-positive result, we replicated the aPKC experiments with different driver lines.

As PKC activity is required in MNs (21), we limited the aPKC knockout to these neurons which abolished self-learning (Figure 1C). To test the hypothesis articulated above that aPKC activity is required in *FoxP* neurons, we also knocked out aPKC in *FoxP* neurons, which also abolished self-learning. (Figure 1C). Using two different driver lines also controls for driver-specific effects and potential expression outside of motor neurons. Both driver lines support operant self-learning in principle (article in preparation, data at DOI: 10.5283/epub.52962), with C380-Gal4 also already in the peer-reviewed literature (21). To our knowledge, C380 and FoxP-iB expression overlaps only in MNs. Thus, three manipulations of aPKC showed effects on self-learning, while two manipulations of PKC53E both failed to show an effect on self-learning.

3.2. *aPKC* and *FoxP* are co-expressed in identified flight steering MNs

Thus, the behavioral data presented above support the hypothesis that the plasticity mediating operant self-learning takes place in neurons that co-express both *FoxP* and *aPKC*. Whole-mount confocal microscopy of fly central nervous systems with *aPKC*-Gal4 and *Foxp*-

LexA expression suggested that neurons expressing both *aPKC* and *FoxP* exist only in the ventral nerve cord (VNC, Figure 2A, B). We identified co-expressing neurons in all neuromers of the VNC, with the ventral location of the mesothoracic *aPKC/FoxP* neurons suggesting potential wing MNs (Figure 2C)(78). To ascertain the identity of this ventral cluster of co-expressing neurons, we marked MNs with the driver line D42-Gal4 and used FoxP-LexA to stain all *FoxP* neurons. With these labels, we identified a ventral sub-population of putative wing MNs expressing *FoxP*, which matched the location of the *aPKC/FoxP* neurons identified before (Figure 2D). The overlap (or lack thereof) may not be clearly visible in these 2D renderings presented in this text, which is why we made the 3D image stacks available for closer scrutiny (DOI: 10.5281/zenodo.10047941)(79). Both the C380-Gal4 driver used in Figure 1 and the D42-Gal4 driver used here, label not only MNs but also other neurons, in particular cholinergic neurons. However, the lines overlap in MNs in the VNC. This is the main reason why we have both historically and also in this work used both drivers (c380-Gal4 and D42-Gal4) interchangeably. *FoxP* is also expressed in these MNs and knocking out aPKC in these neurons also impaired operant self-learning. None of these lines of evidence is sufficient to conclude that steering MNs are the site of plasticity for

operant self-learning. Taken together, they justify testing the hypothesis that wing steering MNs expressing both aPKC and FoxP may be important for operant self-learning. These suggestive results, together with the recently published draft VNC connectome (80), motivated us to analyze the direct wing steering muscles for innervation by MNs with both aPKC and *FoxP* expression, instead of further quantifying the VNC dataset. The expected outcome of this analysis was to obtain a more high-quality and higher resolution picture of aPKC/FoxP expression in direct steering muscle MNs.

Figure 2: FoxP and aPKC are co-expressed in MNs.

Confocal stacks of whole mount preparations of central nervous systems; A-C: green - aPKC-Gal4>CD8::GFP, red - FoxP-LexA>CD8::RFP; D: green - D42-Gal4>CD8::GFP, red - FoxP-LexA>CD8::RFP. Confocal image stacks available at: 10.5281/zenodo.10047941.

A. Adult brain (top) with ventral nerve cord (VNC, bottom) attached. No co-expressing cells can be observed in the brain, whereas such neurons(yellow) are readily observable in all neuromers of the VNC (arrowheads).

B. VNC with *aPKC/Foxp* co-expression (yellow) both in cell bodies and fiber tracts in nerves (arrowheads).

C. C1: Dorsal view of motor neuron reconstruction (modified from (78). C2: Confocal image stack of dorsal view of the mesothoracic neuromer with putative wing MNs expressing both *aPKC* (green) and *FoxP* (red) marked.

D. D1: Lateral view of motor neuron reconstruction (modified from (78). D2: Confocal image stack of mesothoracic neuromer with all MNs (green) and FoxP neurons (red) marked. This lateral view supports the hypothesis that the ventral cluster of *aPKC/FoxP* neurons comprises wing MNs.

Each direct steering muscle is innervated by one single identified MN (81). Thus, identifying the muscles reveals the identity of the corresponding MNs. To test for the expression of aPKC and *FoxP* in MNs that innervate direct wing muscles, we expressed UAS-GFP under the control of aPKC-Gal4 (Figure 3, second column) and RFP under the control of FoxP-lexA (Figure 3, third column). The signal of each fluorescent reporter was further enhanced by immunocytochemistry, and the active zone marker *bruchpilot* (*brp*; (66) was used to label the neuromuscular synapses in flight MN axon terminals (Figure 3, fourth column). In total, *Drosophila* is equipped with 12 flight steering muscles each of which likely contributes to a distinct function in flight control (82). The activation patterns of four of these muscles have

been analyzed in response to optomotor stimulation during flight (first and second basalar muscles b1 and b2, as well as first pterale I muscle (i1) and third pterale II muscle (iii1) (82–84). A schematic of these 4 direct flight muscles plus five adjacent ones that were analyzed in this study illustrates their spatial arrangement, shapes, and depicts which ones are innervated by steering MNs that express both aPKC (green) and FoxP (magenta), only one of both, or neither of them (grey; Figure 3A1).

Representative maximum intensities projection views at 20x magnification allow us to visualize all nine flight steering muscles investigated in one field of view (Figs. 3A2-A4). Except the pleurosternal muscle 1 (ps1) and the posterior notal wing process muscles (hg), all other steering muscles are innervated by aPKC-positive MNs (Figs. 3A1, A2, green). In contrast, six of the nine steering muscles are innervated by *FoxP*-positive MNs, including the basalars b1 and b3, the pterale i1, iii3, and iii4, as well as the pleurosternal muscle ps1 (Figs. 3A1, A3, magenta). A subset of five flight steering muscles is innervated by MNs that express both genes required for operant self-learning. These are b1, b3, i1, iii3, and iii4 (Figure 3A).

Figure 3. A subset of direct wing muscles is innervated by FoxP/aPKC co-expressing MNs. Representative projection view of MN terminals on the direct flight steering muscles in animals with GFP label in aPKC expressing cells (aPKC-Gal4>CD8::GFP, green), RFP expression in FoxP expressing cells (FoxP-LexA>CD8::RFP, magenta), and immunolabeling for the presynaptic active zone marker *bruchpilot* (*brp*, cyan) reveal which direct flight steering MNs express either aPKC, FoxP, or both, or none of them but only *brp* in presynaptic active zones. **(A1)** depicts the orientation, shape, and abbreviated names of direct flight steering muscles and summarizes which ones are innervated by aPKC expressing MNs (green), by FoxP-expressing MNs (magenta), or by MNs without FoxP and aPKC expression (grey). **(A2)** Projection view of direct flight muscles and their innervation with GFP expression under the control of aPKC-GAL4 (green) at 20x magnification. **(A3)** Same preparation, image stack, and field of view but with RFP expression under the control of FoxP-lexA (magenta). **(A4)** Same preparation, image stack, and field of view with *brp* immunolabel (cyan) in presynaptic active zones of flight steering MNs. **(B1-B4).** Same preparation but with selective enlargement of the three basalare muscles (b1-b3), with all three labels in **(B1)**, GFP label in aPKC-expressing cells (green, **B2**), RFP label in FoxP expressing cells (magenta, **B3**), and Brp label in presynaptic active zones (cyan, **B4**). Muscles b1 and b3 are innervated by steering MNs with aPKC and FoxP expression, but b2 is devoid of FoxP-expressing innervation. **(C1-C4)** Same preparation but with selective enlargement of second basalare (b2) and the adjacent pterale 1 (i1) and pterale II (iii3) muscles with all three labels **(C1)**, GFP label in aPKC-expressing cells (green, **C2**), RFP label in FoxP-expressing cells (magenta, **C3**), and *brp* label in presynaptic active zones (cyan, **C4**). Only i1 is faintly labeled for terminals with aPKC FoxP expression. **(D1-D4)** Same preparation but with selective enlargement of the pterale II muscles iii3 and iii4, the adjacent pleurosternal muscle (ps1), and the posterior notal wing process muscles (hg) with all three labels **(D1)**, GFP label in aPKC-expressing cells (green, **D2**), RFP label in FoxP-expressing cells (magenta, **D3**), and Brp label in presynaptic active zones (cyan, **C4**). The pterale II muscles iii3 and iii4 are innervated by terminals with aPKC and FoxP expression. Image stacks available at: 10.5281/zenodo.10606166

Selective enlargements at 40x magnification were used to better visualize the axon terminals on those steering muscles of particular interest, with figure 3 panels B1-B4 focussing on the three muscles that insert at the basalar sclerite, and are thus named basalar 1-3 (b1-b3). Prominent labels of both aPKC (green) and *FoxP* (magenta) are present in the MNs to steering muscles b1 and b3, but the motor axon innervating b2 is devoid of *FoxP* signals (Figure 3, B1-B4).

Muscle b1 has been reported to fire once every wingbeat (83) during straight flight, during the transition from up- to downstroke. Optomotor stimulation causes phase shifts of

b1 firing in the wing beat cycle, which in turn cause changes in wing beat amplitude both in the fruit fly *Drosophila* (83) as well as in the blowfly, *Calliphora* (85). The activity of b3 during flight has not been recorded electrophysiologically, but the muscle exhibits very similar morphological properties compared to b1. Both b1 and b3 are orientated similarly relative to the body axis and the wing hinge, and both are innervated by MNs with particularly large diameter axons (Figs. 3B1-B3) and particularly large active zones (Figure 3B4), suggesting similar functional roles. Steering muscle b2 is also innervated by large diameter axons (Figure 3C1) with large presynaptic active zones (Figure 3C4). Although some aPKC reporter label is detected in the MN to b2 (Figs. 3C1, C2, C4), labeling intensity is considerably fainter than that in b1 and b3 MNs (Figs. 3B1-B3). Fainter reporter labeling indicates weaker aPKC-Gal4 expression. The b2 MN is devoid of the *FoxP* reporter label (Figure 3C3). Although b2 has been reported to respond to optomotor stimulation, its phasic bursting responses correlate with rapid changes in wing beat amplitude as observed during body saccades (83).

Figure 3 panels C1-C4 move the field of view posteriorly and show the pterale 1 muscle i1 and the pterale 2 muscle iii1. Muscle i1 is innervated by a MN with faint aPKC (Figure 3C2) and faint *FoxP* (Figure 3C3) label. i1 has been

reported to respond to optomotor stimulation, but its specific role in optomotor control remains largely unknown (83). Steering muscle iii1 shows some faint aPKC signal (Figure 3C2) but is devoid of any *FoxP* label (Figure 3C3) and does not participate in optomotor flight control (83).

The roles in flight control of the remaining four steering muscles (ps1, iii3, iii4, and hg; Figs. 3D1-D4) in *Drosophila* are not fully understood. However, iii3 and iii4 are innervated by MNs with aPKC (Figure 3D2) and *FoxP* (Figure 3D3) expression. Axon terminals on ps1 are *FoxP* positive (Figure 3D3) but lack any aPKC label (Figure 3D2). Active zones on steering muscle hg are visible through the *brp* label (Figure 3D4) but the motor axon on hg is devoid of both aPKC (Figure 3D2) and *FoxP* (Figure 3D3).

3.3.   Self-learning breaks optomotor symmetry

Because it is not known at which torque meter reading the fly actually generates zero angular momentum, before each torque learning experiment, it is crucial to set the average of the asymptotic left and right optomotor response (OMR) magnitudes to zero (Figure 4A1, see Methods for details). This is to avoid an initial bias in the torque preference before training and with the assumption that the mid-point between the maximal OMRs roughly corresponds to flying straight (i.e, zero angular momentum). During the ensuing conditioning

procedure, flies show spontaneous torque fluctuations that can reach or even exceed those elicited by optomotor stimulation in magnitude. Within each torque domain (i.e., 'left' or 'right', respectively), the temporal patterns (i.e., slow or fast) of torque fluctuations and their relative direction (i.e., more or less torque) are irrelevant for the heat stimulus as long as the zero point is not crossed: heat remains either on or off until the fly switches torque domains. Until this research, there was no reason to assume any relation between elicited OMRs and spontaneous torque fluctuations, neither conceptually nor anatomically. Now, however, the described role of the steering MNs co-expressing aPKC and FoxP (Figure 3) in large torque fluctuations elicited by optomotor stimuli prompted the hypothesis that there may be a neuroanatomical connection between spontaneous torque fluctuations and elicited OMRs after all: the motor neurons that innervate the steering muscles may be involved in both elicited and spontaneous torque fluctuations. If these neurons were indeed common to elicited OMRs and spontaneous torque fluctuations, we should observe a change in the OMRs after operant yaw torque learning.

To test this hypothesis, we analyzed the OMRs after training of a cohort of wild type

Berlin flies (Figure 4B; control flies from a sep-
arate research project). We found that the as-
ymptotic magnitude of the OMR on the pun-
ished side was reduced after operant training,
while the OMR on the unpunished side re-
mained unaltered compared to before training
(Figure 4B1). Quantifying this observation re-
vealed a significant difference in optomotor
asymmetry after training (Figure 4B2) but not
before training (Figure 4A2) between the two
experimental groups.

Before Training

After Training

909   **Figure 4. Associating one torque domain with heat changes optomotor behavior.**

910   A.   Measurements before training
911   **A1** Averaged optomotor traces of flies punished either on the 'left'(yellow) or on the 'right' (green)
912   torque domain. Both groups show similar response magnitudes in either direction of the optomo-
913   tor stimulus. Errors are standard deviations.
914   **A2** Optomotor asymmetry indices for flies punished either on the 'left'(yellow) or on the 'right'
915   (green) torque domain. The values for both groups spread around zero. Positive values indicate a
916   shift towards positive (right-turning) torque. Both frequentist and Bayesian analyses are displayed
917   to the right of the plots and indicate no difference between the groups.
918   **A3** Regression analysis between torque preference and optomotor asymmetry. Optomotor values
919   were adjusted such that positive values indicate a shift towards the unpunished torque domain. No
920   significant correlation was observed. Left-pointing arrowheads (yellow) denote flies punished on
921   left-turning torque and right-pointing arrowheads (green) denote flies punished on right-turning
922   torque.
923   **A4** Comparison of optomotor asymmetry (left, blue) and torque preference(right, red) indices. Here
924   again, optomotor values were adjusted such that positive values indicate a shift towards the un-
925   punished torque domain. Both measures vary around the zero point and Wilcoxon tests against
926   zero are not significant (p-values above each plot). Left- and right-pointing arrowheads denote pun-
927   ishment directions as before.

928   B.   Measurements after training
929   **B1** Averaged optomotor traces of flies punished either on the 'left' (yellow) or on the 'right' (green)
930   torque domain. A reduction in the OMR magnitude can be observed on the punished, but not on
931   the unpunished side. Errors are standard deviations.
932   **B2** Optomotor asymmetry indices for flies punished either on the 'left' (yellow) or on the 'right'
933   (green) torque domain. Positive values indicate a shift towards positive (right-turning) torque. The
934   values for each group have now shifted towards the unpunished side compared to the values be-
935   fore training. Both frequentist and Bayesian analyses are displayed to the right of the plots and
936   indicate a significant difference between groups.
937   **B3** Regression analysis between torque preference and optomotor asymmetry. Optomotor values
938   were adjusted such that positive values indicate a shift towards the unpunished torque domain. A
939   significantly positive correlation was observed, such that higher torque preferences entailed larger
940   optomotor asymmetry. Left-pointing arrowheads (yellow) denote flies punished on left-turning
941   torque and right-pointing arrowheads (green) denote flies punished on right-turning torque.
942   **B4** Comparison of optomotor asymmetry (left, blue) and torque preference (right, red) indices.
943   Here again, optomotor values were adjusted such that positive values indicate a shift towards the
944   unpunished torque domain. Both measures are shifted towards more positive values and Wilcoxon
945   tests against zero are now significant for both variables (p-values above each plot). Left- and right-
946   pointing arrowheads denote punishment directions as before.

947   Data available at: 10.5283/epub.54804 (86)

948

949

950

An important question regarding the functional significance of this change in optomotor asymmetry is whether the amount of torque preference after training reflects the amount of optomotor asymmetry. There was no significant correlation between torque preference and optomotor asymmetry before training (Figure 4A3; as expected as the OMR was adjusted to be as symmetrical as practically possible). Once the flies have completed the training phase, torque preference becomes a significant predictor of optomotor asymmetry (Figure 4B3). These results corroborate our hypothesis that the wing MNs identified above, specifically those that have been shown to be involved in generating OMRs (82,83), constitute a site of the plasticity mediating operant self-learning in *Drosophila*. We also tested whether it would be sufficient to test for optomotor asymmetry after training as a proxy measure for torque preference. Before training, as expected, both measures showed similar values ranging around the zero point (Figure 4A4). After training, both measures did deviate from zero towards the unpunished direction, however, the effect was noticeably larger for torque preference than for optomotor asymmetry (Figure 4B4). Together with the correlation explaining about a third of the variance (Figure 4B3), these data may indicate that

the MNs are an important, but not the only site of plasticity in this form of learning.

3.4. Spatial genome editing: *FoxP* likely not required in the brain

Optomotor analysis (Figure 4) suggests there may be additional sites of plasticity besides wing steering MNs. *FoxP*-positive neurons in the brain are straightforward potential candidates for such additional sites. Therefore, we performed spatial CRISPR/Cas9-based genome editing without restricting the manipulation to the adult stage. At the time of these experiments, no driver lines with *FoxP*-overlapping expression in the Saddle and Vest regions were available. Therefore, we tested driver lines expressing in the protocerebral bridge (PCB, Figure 5A), the PCB and adjacent central complex neuropils (see Materials and Methods for details, Figure 5B) and the dorsal cluster neurons (Figure 5C). Overlap of Gal4 expression patterns with FoxP expression was verified using the FoxP-LexA line (74,87) and FoxP-knockout efficiency was quantified previously (74). No self-learning impairment was observed following *FoxP* knockout in these brain areas, suggesting that *FoxP* expression is not necessary in these areas for operant self-learning. The results from these experiments, performed before aPKC was discovered as the likely gene necessary for operant self-learning, appear to corroborate the results of the

1012	aPKC/FoxP co-expression analysis in the brain:
1013	if both aPKC and FoxP are required for operant
1014	self-learning and there are no neurons in the
1015	brain that co-express both aPKC and FoxP,
1016	then knocking out either of them in the brain
1017	should not have any effects on operant self-
1018	learning.

1019

Figure 5: For operant self-learning, *FoxP* is likely not required in the brain.

Performance indices for the first period after training (PI8/12) are plotted. Each colored dot represents a single fly experiment. Red dots denote experiments where the fly was punished on its left turning torque domain, blue dots denote flies that were punished on their right turning domain. Box plots denote medians, quartiles and non-outlier range. Statistical analyses test for differences of PIs against zero (above plots and D).

- A. Flies with *FoxP* knocked out in the protocerebral bridge (left, yellow) as well as the gRNA (middle, green) and Cas9 (right, blue) control flies showed high PIs, large Bayes factors (D) and small p-values, indicating their self-learning was intact. Data: 10.5283/epub.52956 (88)
- B. Flies with *FoxP* knocked out in the protocerebral bridge and additional components of the central complex (left. yellow) as well as the gRNA (middle, green) and Cas9 (right, blue) control flies showed high PIs, large Bayes factors (D) and small p-values, indicating their self-learning was intact. Data: 10.5283/epub.52951 (89)
- C. Flies with *FoxP* knocked out in the dorsal cluster neurons (left, yellow) as well as the gRNA (middle, green) and Cas9 (right, blue) control flies showed high PIs, large Bayes factors (D) and small p-values, indicating their self-learning was intact. Data: 10.5283/epub.52946 (89,90)
- D. Bayesian statistics for the three datasets.

We discovered that aPKC is required in MNs for operant self-learning (see above). As *FoxP* is also expressed in MNs, we knocked *FoxP* out in MNs using two different driver lines, C380-Gal4 and D42-Gal4. However, CRISPR/Cas9-mediated *FoxP* knockout in MNs disrupted flight-performance of manipulated flies to an extent that precluded any torque learning experiments. Our results above (Figure 3) will now allow us to select more specific driver lines, expressing only in the identified wing steering MNs (91), which may yield *FoxP* knock-out flies (or any other manipulation) with sufficient flight performance.

3.5. Temporal genome editing: Self-learning impaired only after a 2-week *FoxP* knockout

FoxP was shown to be important for normal development (74,92). Strong motor impairments have been reported after a developmental *FoxP* knockout, for instance rendering the animals unable to fly (see above). Since the ability to fly is a basic requirement for torque learning, we used a CRISPR/Cas9-based approach to pan-neuronally knock out *FoxP* in adult flies. No effect was observed on self-learning two days after the start of the knock-out induction (Figure 6 A), but waiting for 14 days after the pan-neural knock-out yielded a significant self-learning impairment (Figure 6B). As a test seven days after induction was also without effect (DOI: 10.5283/epub.52965),

self-learning remains functional for at least 7-
14 days after cessation of *FoxP* gene transcrip-
tion in all neurons.

Figure 6: Knockout of *FoxP* in adult *Drosophila* shows learning impairments only after 14 days.
Plotted are performance indices for the first period after training (PI12). Each colored dot represents a single fly experiment. Red dots denote experiments where the fly was punished on its left turning torque domain, blue dots denote flies that were punished on their right turning domain. Box plots denote medians, quartiles and non-outlier range. Statistical analyses test for differences of PIs against zero.

A. Self-learning two days after *FoxP* knockout induction with RU486. Experimental animals (left, yellow) as well as gRNA (middle, green) and Cas9 (left, blue) control animals showed all high PIs as well as large Bayes Factors above 1000 and p-values below 0.005, indicating that all groups showed unimpaired self-learning. Data: 10.5283/epub.52963 (93)

B. Self-learning 14 days after *FoxP* knockout induction with RU486 and 12 days after cessation of RU486 administration. Experimental animals (left, yellow) showed low PIs as well as a Bayes Factor of less than one together with a large p-value, whereas both the genetic control animals without RU486 treatment (middle, green) and the pooled RU486-treated gRNA and Cas9 controls (right, blue) showed high PIs, large Bayes Factors and p-values smaller than 0.005, indicating an impairment in self-learning only in the experimental group. Data: 10.5283/epub.52964 (94)

3.6. *aPKC* acts via non-canonical pathways in self-learning

Having established that *aPKC* is required in *FoxP*-positive MNs, we sought to identify further components of the *aPKC*-dependent plasticity underlying operant self-learning in *Drosophila*. It is not uncommon for plasticity mechanisms in the adult animal to recruit genes with a function during neuronal development (95–97). With PKCs being notorious for being able to compensate for genetic manipulations (69–73), one reliable countermeasure has proven to shorten the time period between manipulation and testing sufficiently to ensure compensation has no time to take place (21,31). Following this tried-and-tested approach, our manipulations of prominent PKC interaction partners were thus restricted to adult neurons. One such prominent interaction partner of *aPKC* during *Drosophila* nervous system development is *bazooka* (*baz*), a crucial component of the highly conserved PAR complex (98,99). However, knocking out *baz* in all adult neurons did not disrupt operant self-learning (Figure 7A), suggesting that the Par complex signaling pathway is not involved in operant self-learning. A second prominent aPKC interaction partner is the kidney and brain protein (KIBRA), which acts in the conserved Hippo pathway (100,101) and also proposed to be involved in

 learning/memory (102–106). Knocking out KI-
BRA in all adult neurons also did not disrupt
operant self-learning (Figure 7B).

A

	bf	error
nsybGS>g:Baz;Cas9(+RU)	29900	5.62e-12
nsybGS>g:Baz;Cas9(-RU)	69.9	1.13e-07

B

	bf	error
nsybGS>g:Kibra;Cas9(+RU)	389	2.11e-10
nsybGS>g:Kibra;Cas9(-RU)	757	1.11e-10

Figure 7: Neither *bazooka* nor KIBRA are involved in operant self-learning.

Plotted are performance indices for the first period after training (PI12). Each colored dot represents a single fly experiment. Red dots denote experiments where the fly was punished on its left turning torque domain, blue dots denote flies that were punished on their right turning domain. Box plots denote medians, quartiles and non-outlier range. Statistical analyses test for differences of PIs against zero.

A. Knocking out *bazooka* using CRISPR/Cas9-mediated genome editing in adult animals had no effect on operant self-learning. Both flies that were fed the steroid hormone RU486 (left, yellow) and the genetically identical flies without the hormone (right, green), showed high PIs, low p-values and high Bayes Factors, indicating a preference for the unpunished torque domain. Data: 10.5283/epub.52947 (107)

B. Knocking out KIBRA using CRISPR/Cas9-mediated genome editing in adult animals had no effect on operant self-learning. Both flies that were fed the steroid hormone RU486 (left, yellow) and the genetically identical flies without the hormone (right, green), showed high PIs, low p-values and high Bayes Factors, indicating a preference for the unpunished torque domain. Data: 10.5283/epub.53685 (108)

4. Discussion

Operant self-learning in *Drosophila* is a form of motor learning that appears to be conserved among bilaterians. The transcription factor *FoxP* is involved in various forms of motor learning in chordates (5–9,11,13,16–18), as is PKC (22–24,109–112). PKC is also involved in motor learning in the feeding behavior of the lophotrochozoan *Aplysia (113)* and both are involved in motor learning in the ecdysozoan *Drosophila* (this work and (20,21,31). This wealth of evidence supports what has been called 'deep homology' for motor learning in bilaterians (12).

In this work, we present new insights into the neurobiological mechanisms underlying this form of motor learning.

4.1. Self-learning requires *aPKC* in *FoxP* wing steering motor neurons

While previous evidence suggested that some PKC activity was required in some MNs for operant self-learning (21), it was not clear which PKC gene was involved and in which MNs. We now hypothesize that *aPKC* activity is required in aPKC/*FoxP*-positive direct wing steering MNs in the ventral nerve cord (VNC). Our analysis of these direct wing muscles, responsible for, e.g. wing beat amplitude (a major contributing factor to yaw torque) (114), re-

vealed *aPKC/FoxP* co-expressing MNs innervating a specific subset of these muscles (Figure 3), corroborating our hypothesis.

4.2. Motor neurons involved in generating yaw torque co-express aPKC and *FoxP*

Specifically, the basalar muscle b1 which is known to regulate wing beat amplitude during optomotor stimulation through phase shifts of its action potential within the wing beat cycle (83) is innervated by a MN with very strong aPKC and *FoxP* expression in all 11 animals analyzed. Similarly, b3 is also innervated by a MN with very strong aPKC and *FoxP* expression and the b3 muscle is reported to increase activity when the ipsilateral wing decreases its amplitude (115). The b1 and the b3 muscles thus both share similar sizes and morphologies, are innervated by MNs with particularly thick processes and presynaptic active zones, insert at opposite sides of the basalar sclerite and act in a push-pull fashion in controlling wing beat amplitude (115), a major contributor to yaw torque. A recent comprehensive re-analysis of all steering muscles has confirmed the antagonist role b1 and b3 play in generating yaw torque (116). Although the MN innervating the third basalar muscle, b2, also expresses aPKC along the axon, we did not detect aPKC signal in axon terminals or active zones, and it does not show *FoxP* label, thus

ruling out a function of b2 in aPKC/FoxP mediated operant self-learning. In contrast to b1 and b3, b2 firing is not linked to the wingbeat cycle; instead, it fires in bursts during turning maneuvers (82). Corroborating the conclusion that it is not involved in mediating operant self-learning, b2 is silent during straight flight, but it is likely involved in regulating body saccades (Heide, Götz, 1996), which are by themselves not relevant for controlling the heat in our experiments. For the basalar muscles the emerging picture is that b1 and b3 are the top candidates for self-learning because they likely regulate wingbeat amplitude and thus also yaw torque on a wingbeat cycle by cycle basis (116). Strikingly, the MNs innervating these two muscles show high levels of aPKC and FoxP expression. Although much less is known about the pterale muscles, and neither iii3 nor iii4 (both with aPKC and FoxP co-expression) have been recorded electrophysiologically during flight, a similar picture begins to emerge. The MN to the left muscle i1 is known to be active during right turns (83) and co-expresses aPKC and FoxP, whereas iii1 reportedly (83) does not participate in OMRs and is innervated by a MN with weak aPKC and no *FoxP* label. Also these roles of steering muscles have recently been confirmed (116). In summary, flight steering muscles that are known to regulate yaw torque

in response to horizontal optomotor stimulation receive aPKC and *FoxP* positive motor innervation, while those not involved in yaw torque production are not innervated by such doubly labeled MNs. Because endogenous yaw torque fluctuations, such as those rewarded/punished in our operant experiments also have to use these MNs, it becomes clear that OMRs and operant yaw torque learning share these steering motor neurons as a common component. Future experiments with (likely soon to be available) more specific driver lines will be able to test the effect of aPKC/FoxP manipulations in specific subsets of steering MNs.

4.3. Operant self-learning modifies optomotor responses

From this realization, the question arises if there are other shared neural components between OMRs and operant learning in addition to the doubly labeled steering MNs. Evidence suggests that the steering commands for OMRs are communicated directly from visual areas in the brain via (both identified and yet to be identified) descending neurons with direct synaptic connections onto the steering MNs in the VNC (117,118). This suggests that in the ventral nerve cord, the steering motor neurons are the only neurons that OMRs and operant self-learning share. This evidence entails that any additional overlap would need to be

localized in the brain, where no neurons expressing both aPKC and FoxP. seem to exist (Fig. 2). The remote hypothetical possibility remains that OMR plasticity after operant learning may be caused by non-aPKC/Foxp-dependent plasticity mechanisms in the same visual parts of the brain where also the steering commands for OMRs are computed. Future research will address this possibility. The currently available evidence thus points towards the steering MNs we have identified as the only set of neurons where a modification due to operant learning could have an effect on OMRs. The fact that we have observed precisely such an effect (Fig. 4), only leaves the doubly labeled steering neurons (Fig. 3) as the sites for the aPKC/FoxP mediated plasticity mechanisms underlying operant yaw torque learning.

Then again, only about a third of the variance in the torque preference after training can be explained by this optomotor asymmetry. This result explains the observations that there are flies with a strong conditioned torque preference but with an optomotor asymmetry in the opposite direction, as well as flies with a weak conditioned torque preference and large optomotor asymmetry (Figure 4B3). Clearly, plasticity in steering MNs appears to be important, but it does not reflect the entirety of the learning processes. On the other hand, it may be that the strong OM stimuli we

use here overshadow potentially larger asymmetries in OMRs. We will test this in future experiments using weaker OM stimuli after operant self-learning.

Plasticity in steering movements such as OMRs have been observed before, such as in classic "inversion goggles" experiments where the coupling between the fly's movements and the environment was reversed (119) or in a more recent experiment revealing adaptation processes (120). It has long been recognized that insects with asymmetrical wing damage need to adjust the neural commands for generating torque to compensate for the changed physical torque (e.g., (117,118,121–125)). Plasticity in wing steering motor neurons provides a potential mechanism for such adjustments.

4.4. Motoneuron plasticity mediates operant self-learning

Taken together, these results converge on the hypothesis that plasticity in MNs that innervate the direct muscles involved in generating yaw torque, but not other steering MNs, mediates an important aspect of operant self-learning in *Drosophila*. The importance of MN plasticity is emphasized by *FoxP*-dependent plasticity apparently not being required in the brain (Figure 5). This MN plasticity could either be implemented by (a) postsynaptic plasticity

of the input synapses to these MNs, postsynaptic because the MNs but not the interneurons express the proteins required for self-learning, or (b) on the level of the intrinsic excitability of flight steering motoneurons, or (c) on the level of the output synapses to the respective steering muscles. However, the mechanism and subcellular localization of self-learning in MNs remains to be determined.

A second interesting, yet unstudied population of *aPKC/FoxP* co-expressing neurons resides in the abdominal neuromer of the VNC. As flies use their abdomen analogously to a rudder during turns in flight (126,127), involvement of these neurons seems plausible in addition to wing MNs.

While few studies in insects have shown MN plasticity, the *Aplysia* sensorimotor synapse is a classical model for research on plasticity mechanisms. There is a rich literature on MN plasticity in this preparation, some of which reports PKC-dependent mechanisms (104,128–138). Also in mammals (including humans) MN plasticity in the spinal cord is a readily observable phenomenon in nonclinical and clinical settings (139–148). The discovery of aPKC-dependent plasticity in *Drosophila* MNs expands this body of literature to a genetically tractable organism and inasmuch as clinical practice relies on MN plasticity, may even help

instruct the development of clinical applications.

Until our work, PKC activity had only been shown to be important for memory consolidation/maintenance in world-learning experiments in flies, but not for learning/acquisition (149,150). Also in *Drosophila* at the torque meter, PKCs are dispensable for world-learning (31). The literature on PKCs in learning and memory in other animals is complex and multifaceted. In some preparations, PKC isoforms are required during memory maintenance, in some also during acquisition and in others different isoforms distinguish between acquisition and consolidation (69,70,102,104,129,134,136,151–159). As the manipulations in our experiments lasted throughout training together with the tests immediately following training and we did not test for long-term memory, we can only ascertain that aPKC is required in a very narrow time window of minutes around training. Future research will address whether aPKC must be present during training, test, or both.

Notably, there is one other preparation where PKC activity is involved and which is also conceptually most closely resembling the one we used here, operant reward learning in *Aplysia* feeding behavior. However, in this preparation, the calcium-dependent Apl-I PKC not the atypical Apl-III PKC appears to be mediating

the plasticity (113). Given the degeneracy between the different PKC genes and the fact that they can not only compensate for long-term PKCi-mediated inhibition (this work and (21), but also for each other (69), more research is needed to elucidate how these different mechanisms of plasticity evolved and are related to each other.

4.5. Self-learning may be mediated by a non-canonical *aPKC/FoxP* pathway

The observation that neither *bazooka* (*baz*) nor the kidney and brain gene (KIBRA), two prominent interaction partners of aPKC (98–104,106,160,161), showed an effect on operant self-learning when they were knocked out in the adult nervous system (Figure 7), raises questions about the effectiveness of the CRISPR/Cas9 method in these cases. In particular, one may question the approach of temporally limiting the manipulation to adult neurons. Although this had been effective with previous PKC manipulations using an inhibitory peptide, PKCi (21,31), it only yielded an intermediate effect with a CRISPR-mediated knockout targeting aPKC (Figure 1B). In contrast, temporally uncontrolled knockout of aPKC in MNs of the VNC (Figure 1C) proved surprisingly effective, given expectations from prior experience.

Nonetheless, we are confident that both manipulations successfully knocked out each of these genes in a way sufficient to induce a phenotype, if either were necessary for self-learning. First, we found that the *baz* knockout, rather than impairing self-learning, actually *increases* learning performance (manuscript in preparation, thesis: (87)), suggesting the PAR complex may be sequestering *aPKC* and thereby limiting its availability for self-learning. Thus, these data suggest that *baz* is indeed not directly involved in mediating the aPKC activity contributing to plasticity in steering MNs; instead, it may be binding aPKC in the PAR complex, preventing it from playing its role in MN plasticity (162). Second, our KIBRA knock-out had a severe effect on flight performance when elicited during development, suggesting that also this manipulation was, in principle, effective. That being said, without clear evidence that the *baz* and KIBRA proteins are completely absent, these results remain suggestive rather than conclusive. In addition to protein-level analysis of CRISPR efficacy (as we have performed in CRISPR-mediated FoxP knock-out (74)), future experiments will use FoxP-iB and c380/D42 drivers to drive baz and KIBRA knock-outs. The forthcoming full description of the KIBRA gRNA line by the Krahn laboratory will also help bolster or refute the results we have obtained.

Interestingly, at least during development, *baz* and KIBRA have been reported to have opposite effects on the function of *aPKC*, with the PAR complex (*baz*) and the Hippo pathway (KIBRA) mutually inhibiting each other (163). While *baz* is thought to mediate *aPKC* activity, KIBRA is thought to exert negative regulatory effects on *aPKC*. Thus, knocking out each one of them should have revealed a decrement in self-learning in at least one of them, if the processes during development were recapitulated during self-learning. On the other hand, experiments in which KIBRA has been shown to be involved in learning/memory have suggested a positive rather than a negative regulatory role (104,154), albeit with an emphasis on long-term memory rather than learning (164). Whichever way *aPKC* may be interacting with components of these canonical pathways, the literature predicts that at least one of our manipulations should have revealed a decrement in operant self-learning. The fact that this prediction was falsified may suggest that *aPKC* exerts its function in a non-canonical manner in operant self-learning plasticity. We are currently pursuing research into the possibility that *bazooka* may be a negative regulator of aPKC activity (162) during operant self-learning.

4.6. Persistent *FoxP* effect on operant self-learning in adults

As *FoxP* mutants are impaired in operant self-learning (20), two hypotheses about the role of this prominent transcription factor arise. First, *FoxP* may be directly involved in the learning process via some unknown, cytosolic function. A transcription factor function appears unlikely, because of the short duration of our experiments. Second, *FoxP* may exert its effects as a developmental regulator, being crucial for the development of the circuits mediating operant self-learning. As the developmental role of FoxP genes is well documented (7,14,74,92,165,166) and there are few domains in the gene that would lend themselves to a hypothetical cytosolic function, the latter hypothesis appeared more plausible. The result that adult knockout of *FoxP* had no immediate effect on operant self-learning (Figure 6A) seemed to corroborate this hypothesis. In contrast, supporting a continued role of FoxP genes in motor learning also after development are data from songbirds where *FoxP2* gene expression is not only regulated by singing (15,167–169), but where normal *FoxP2* expression is necessary in adults to maintain learned song (8,170). To also test the second hypothesis, we aged the flies after the *FoxP* knockout and tested them 7 and 14-days later. Our results (Fig. 6) suggest a role for adult *FoxP*

expression in maintaining operant self-learning capabilities after development, analogous to the role of FoxP2 in songbird vocal learning (8,170). There are three possible explanations for this result: For one, the genes regulated by *FoxP (171)* may continue to exert their functions for this amount of time also without the FoxP protein present. Second, the half-life of the FoxP protein may be long enough. Third, a combination of these two explanations. Further research is needed to distinguish between these options.

Data availability

Underlying data, figures

1A: 10.5283/epub.52958

1B: 10.5283/epub.52957

1C: 10.5283/epub.52944

2: 10.5281/zenodo.10047941

3: 10.5281/zenodo.10606166

4: 10.5283/epub.54804

5A: 10.5283/epub.52956

5B: 10.5283/epub.52951

5C: 10.5283/epub.52946

6A: 10.5283/epub.52963

6B: 10.5283/epub.52964

7A: 10.5283/epub.52947

1524   7B: 10.5283/epub.53685

1525   License: CC0

1526

Software availability

Latest source code available from: https://github.com/brembslab/DTSevalua-
tions

Software controlling experiments and collecting data:
https://www.doi.org/10.5281/zenodo.7102195

Software evaluating data:

https://www.doi.org/10.5281/zenodo.10041052

Data model:

10.5281/zenodo.10041052

Python script keeping local and public datasets synchronized:

10.5281/zenodo.7101741

License: GPL-3.0

1539

Acknowledgements: We thank Thomas Kopp in the electronics workshop of the
University of Regensburg, without whose tireless support this work could not have
been accomplished. We are indebted to Michael Krahn for very helpful feedback
on PKC interaction partners and for sending us his KIBRA gRNA flies before publi-
cation. We are grateful to several anonymous peer-reviewers as well as the named
reviewers of this submission (Efthimios M C Skoulakis, Katrin Vogt and Wayne Sos-
sin) who have helped improve the manuscript substantially. We thank the DFG for
financial support: DFG BR 1892/17-1. This work is dedicated to the late Jochen
Pflüger, at whose memorial CD and BB agreed to collaborate on motor neurons.

5.   References

1.   Arendt D. Elementary nervous systems. Philos Trans R Soc Lond B Biol Sci. 2021
Mar 29;376(1821):20200347.

2.   Jékely G. The chemical brain hypothesis for the origin of nervous systems. Philos
Trans R Soc Lond B Biol Sci. 2021 Mar 29;376(1821):20190761.

3.   Keijzer F, van Duijn M, Lyon P. What nervous systems do: early evolution, input–out-
put, and the skin brain thesis. Adapt Behav. 2013 Apr;21(2):67–85.

4.   Walsh B, Smith A, Weber-Fox C. Short-term plasticity in children's speech motor sys-
tems. Dev Psychobiol. 2006 Dec;48(8):660–74.

5.   Lai CS, Fisher SE, Hurst JA, Vargha-Khadem F, Monaco AP. A forkhead-domain gene
is mutated in a severe speech and language disorder. Nature. 2001 Oct
4;413(6855):519–23.

6.   Bolhuis JJ, Okanoya K, Scharff C. Twitter evolution: converging mechanisms in bird-
song and human speech. Nat Rev Neurosci. 2010 Nov;11(11):747–59.

7.   Co M, Anderson AG, Konopka G. FOXP transcription factors in vertebrate brain de-
velopment, function, and disorders. Wiley Interdiscip Rev Dev Biol. 2020
Sep;9(5):e375.

8.   Day NF, Hobbs TG, Heston JB, White SA. Beyond Critical Period Learning: Striatal
FoxP2 Affects the Active Maintenance of Learned Vocalizations in Adulthood.
eNeuro [Internet]. 2019 Mar;6(2). Available from:
http://dx.doi.org/10.1523/ENEURO.0071-19.2019

9.   Fisher SE, Scharff C. FOXP2 as a molecular window into speech and language.
Trends Genet. 2009 Apr;25(4):166–77.

10. Zhang J, Webb DM, Podlaha O. Accelerated protein evolution and origins of human-
specific features: Foxp2 as an example. Genetics. 2002 Dec;162(4):1825–35.

11. Haesler S, Rochefort C, Georgi B, Licznerski P, Osten P, Scharff C. Incomplete and in-
accurate vocal imitation after knockdown of FoxP2 in songbird basal ganglia nu-
cleus Area X. PLoS Biol. 2007 Dec;5(12):e321.

12. Scharff C, Petri J. Evo-devo, deep homology and FoxP2: implications for the evolu-
tion of speech and language. Philos Trans R Soc Lond B Biol Sci. 2011 Jul
27;366(1574):2124–40.

13. French CA, Jin X, Campbell TG, Gerfen E, Groszer M, Fisher SE, et al. An aetiological
Foxp2 mutation causes aberrant striatal activity and alters plasticity during skill
learning. Mol Psychiatry. 2012 Nov;17(11):1077–85.

14. Takahashi H, Takahashi K, Liu FC. FOXP genes, neural development, speech and lan-
guage disorders. Adv Exp Med Biol. 2009;665:117–29.

15. Teramitsu I, Poopatanapong A, Torrisi S, White SA. Striatal FoxP2 is actively regu-
lated during songbird sensorimotor learning. PLoS One. 2010 Jan 6;5(1):e8548.

16. French CA, Vinueza Veloz MF, Zhou K, Peter S, Fisher SE, Costa RM, et al. Differential
effects of Foxp2 disruption in distinct motor circuits. Mol Psychiatry. 2019
Mar;24(3):447–62.

17. Groszer M, Keays DA, Deacon RMJ, de Bono JP, Prasad-Mulcare S, Gaub S, et al. Im-
paired synaptic plasticity and motor learning in mice with a point mutation impli-
cated in human speech deficits. Curr Biol. 2008 Mar 11;18(5):354–62.

18. Schreiweis C, Bornschein U, Burguière E, Kerimoglu C, Schreiter S, Dannemann M,
et al. Humanized Foxp2 accelerates learning by enhancing transitions from declara-
tive to procedural performance. Proc Natl Acad Sci U S A. 2014 Sep
30;111(39):14253–8.

19. Lozano R, Gbekie C, Siper PM, Srivastava S, Saland JM, Sethuram S, et al. FOXP1 syn-
drome: a review of the literature and practice parameters for medical assessment
and monitoring. J Neurodev Disord. 2021 Apr 23;13(1):18.

20. Mendoza E, Colomb J, Rybak J, Pflüger HJ, Zars T, Scharff C, et al. Drosophila FoxP
mutants are deficient in operant self-learning. PLoS One. 2014 Jun 25;9(6):e100648.

21. Colomb J, Brembs B. PKC in motorneurons underlies self-learning, a form of motor
learning in Drosophila. PeerJ. 2016 Apr 25;4:e1971.

22. Sakaguchi H, Yamaguchi A. Early song-deprivation affects the expression of protein
kinase C in the song control nuclei of the zebra finch during a sensitive period of
song learning. Neuroreport. 1997 Aug 18;8(12):2645–50.

23. Metzger F, Kapfhammer JP. Protein kinase C: its role in activity-dependent Purkinje
cell dendritic development and plasticity. Cerebellum. 2003;2(3):206–14.

24. Gallimore AR, Kim T, Tanaka-Yamamoto K, De Schutter E. Switching On Depression
and Potentiation in the Cerebellum. Cell Rep. 2018 Jan 16;22(3):722–33.

25. Rochefort C, Arabo A, André M, Poucet B, Save E, Rondi-Reig L. Cerebellum shapes
hippocampal spatial code. Science. 2011 Oct 21;334(6054):385–9.

26. Fresco N, Tzelgov J, Shmuelof L. How can caching explain automaticity? Psychon Bull
Rev [Internet]. 2022 Oct 12; Available from: http://dx.doi.org/10.3758/s13423-022-
02191-0

27. Krakauer JW, Hadjiosif AM, Xu J, Wong AL, Haith AM. Motor Learning. Compr Physiol.
2019 Mar 14;9(2):613–63.

28. Salmon DP, Butters N. Neurobiology of skill and habit learning. Curr Opin Neuro-
biol. 1995 Apr;5(2):184–90.

29. Du Y, Krakauer JW, Haith AM. The relationship between habits and motor skills in
humans. Trends Cogn Sci. 2022 May;26(5):371–87.

30. Brembs B. Mushroom bodies regulate habit formation in Drosophila. Curr Biol.
2009 Aug 25;19(16):1351–5.

31. Brembs B, Plendl W. Double dissociation of PKC and AC manipulations on operant
and classical learning in Drosophila. Curr Biol. 2008 Aug 5;18(15):1168–71.

32. Colomb J, Brembs B. The biology of psychology: "Simple" conditioning? Commun In-
tegr Biol. 2010 Mar;3(2):142–5.

33. Brembs B. Spontaneous decisions and operant conditioning in fruit flies. Behav Pro-
cesses. 2011 May;87(1):157–64.

34. Götz KG. [Optomoter studies of the visual system of several eye mutants of the fruit
fly Drosophila]. Kybernetik. 1964 Jun;2(2):77–92.

35. Wolf R, Heisenberg M. Basic organization of operant behavior as revealed in Dro-
sophila flight orientation. J Comp Physiol A. 1991 Dec;169(6):699–705.

36. Guo A, Li L, Xia SZ, Feng CH, Wolf R, Heisenberg M. Conditioned visual flight orienta-
tion in Drosophila: dependence on age, practice, and diet. Learn Mem. 1996 Jul-
Aug;3(1):49–59.

37. Brembs B. Operant learning of Drosophila at the torque meter. J Vis Exp [Internet].
2008 Jun 16;(16). Available from: http://dx.doi.org/10.3791/731

38. Tang S, Juusola M. Intrinsic activity in the fly brain gates visual information during
behavioral choices. PLoS One. 2010 Dec 30;5(12):e14455.

39. Brembs B, SimonBnm, AnEhw, Eriksson A, von der Linde M. brembslab/DTSevalua-
tions: Second public release [Internet]. Zenodo; 2023. Available from: https://ze-
nodo.org/doi/10.5281/zenodo.10041052

40. Ioannidis JPA. Why most published research findings are false. PLoS Med. 2005
Aug;2(8):e124.

41. Berrar D, Dubitzky W. Should significance testing be abandoned in machine learn-
ing? Int J Data Sci Anal. 2019 Jun;7(4):247–57.

42. Berner D, Amrhein V. Why and how we should join the shift from significance test-
ing to estimation. J Evol Biol. 2022 Jun;35(6):777–87.

43. Maier M, Lakens D. Justify your alpha: A primer on two practical approaches. Adv
Methods Pract Psychol Sci. 2022 Apr;5(2):251524592210803.

44. Wulff JN, Taylor L. EXPRESS: How and why alpha should depend on sample size: A
Bayesian-frequentist compromise for significance testing. Strateg Organ [Internet].
2023 Nov 6; Available from: http://jour-
nals.sagepub.com/doi/10.1177/14761270231214429

45. Miller J, Ulrich R. The quest for an optimal alpha. PLoS One. 2019 Jan
2;14(1):e0208631.

46. Benjamin DJ, Berger JO, Johannesson M, Nosek BA, Wagenmakers EJ, Berk R, et al.
Redefine statistical significance. Nat Hum Behav. 2018 Jan;2(1):6–10.

47. Trafimow D, Amrhein V, Areshenkoff CN, Barrera-Causil CJ, Beh EJ, Bilgiç YK, et al.

Manipulating the Alpha Level Cannot Cure Significance Testing. Front Psychol. 2018
May 15;9:699.

48. Wasserstein RL, Schirm AL, Lazar NA. Moving to a world beyond "*p* < 0.05." Am Stat.
2019 Mar 29;73(sup1):1–19.

49. Amrhein V, Greenland S, McShane B. Scientists rise up against statistical signifi-
cance. Nature. 2019 Mar;567(7748):305–7.

50. Ioannidis JPA. Retiring statistical significance would give bias a free pass. Nature.
2019 Mar;567(7749):461.

51. Lakens D, Adolfi FG, Albers CJ, Anvari F, Apps MAJ, Argamon SE, et al. Justify your al-
pha. Nat Hum Behav. 2018 Feb 26;2(3):168–71.

52. Gigerenzer G. Statistical rituals: The replication delusion and how we got there. Adv
Methods Pract Psychol Sci. 2018 Jun;1(2):198–218.

53. Wasserstein RL, Lazar NA. The ASA statement on p-values: Context, process, and
purpose. Am Stat. 2016 Apr 2;70(2):129–33.

54. Colquhoun D. An investigation of the false discovery rate and the misinterpretation
of p-values. R Soc Open Sci. 2014 Nov;1(3):140216.

55. Wagenmakers EJ, Lee M, Lodewyckx T, Iverson GJ. Bayesian Versus Frequentist In-
ference. In: Bayesian Evaluation of Informative Hypotheses. New York, NY: Springer
New York; 2008. p. 181–207.

56. Wagenmakers EJ, Love J, Marsman M, Jamil T, Ly A, Verhagen J, et al. Bayesian infer-
ence for psychology. Part II: Example applications with JASP. Psychon Bull Rev. 2018
Feb;25(1):58–76.

57. Wagenmakers EJ, Marsman M, Jamil T, Ly A, Verhagen J, Love J, et al. Bayesian infer-
ence for psychology. Part I: Theoretical advantages and practical ramifications. Psy-
chon Bull Rev. 2018 Feb;25(1):35–57.

58. Wagenmakers EJ, Morey RD, Lee MD. Bayesian benefits for the pragmatic re-
searcher. Curr Dir Psychol Sci. 2016 Jun;25(3):169–76.

59. Malone HE, Coyne I. Complementing the P-value from null-hypothesis significance
testing with a Bayes factor from null-hypothesis Bayesian testing. Nurse Res [Inter-
net]. 2020 Nov 4; Available from: http://dx.doi.org/10.7748/nr.2020.e1756

60. Wei Z, Yang A, Rocha L, Miranda MF, Nathoo FS. A Review of Bayesian Hypothesis
Testing and Its Practical Implementations. Entropy [Internet]. 2022 Jan 21;24(2).
Available from: http://dx.doi.org/10.3390/e24020161

61. Tendeiro JN, Kiers HAL. A review of issues about null hypothesis Bayesian testing.
Psychol Methods. 2019 Dec;24(6):774–95.

62. Brembs B, Wiener J. Context and occasion setting in Drosophila visual learning.
Learn Mem. 2006 Sep-Oct;13(5):618–28.

63. Liu L, Wolf R, Ernst R, Heisenberg M. Context generalization in Drosophila visual
learning requires the mushroom bodies. Nature. 1999 Aug 19;400(6746):753–6.

64. Liu G, Seiler H, Wen A, Zars T, Ito K, Wolf R, et al. Distinct memory traces for two vis-
ual features in the Drosophila brain. Nature. 2006 Feb 2;439(7076):551–6.

65. Brembs B, Hempel de Ibarra N. Different parameters support generalization and
discrimination learning in Drosophila at the flight simulator. Learn Mem. 2006 Sep-
Oct;13(5):629–37.

66. Kittel RJ, Wichmann C, Rasse TM, Fouquet W, Schmidt M, Schmid A, et al. Bruchpilot
promotes active zone assembly, Ca2+ channel clustering, and vesicle release. Sci-
ence. 2006 May 19;312(5776):1051–4.

67. Vonhoff F, Kuehn C, Blumenstock S, Sanyal S, Duch C. Temporal coherency between
receptor expression, neural activity and AP-1-dependent transcription regulates
Drosophila motoneuron dendrite development. Development. 2013 Feb
1;140(3):606–16.

68. Duch C. Original CLSM image stack for figure 3 for [Internet]. Zenodo; 2024. Availa-
ble from: https://zenodo.org/doi/10.5281/zenodo.10606166

69. Tsokas P, Hsieh C, Yao Y, Lesburguères E, Wallace EJC, Tcherepanov A, et al. Com-
pensation for PKMζ in long-term potentiation and spatial long-term memory in mu-
tant mice. Elife [Internet]. 2016 May 17;5. Available from:
http://dx.doi.org/10.7554/eLife.14846

70. Sacktor TC, Hell JW. The genetics of PKMζ and memory maintenance. Sci Signal [In-
ternet]. 2017 Nov 14;10(505). Available from:
http://dx.doi.org/10.1126/scisignal.aao2327

71. Teegala LR, Elshoweikh Y, Gudneppanavar R, Thodeti S, Pokhrel S, Southard E, et al.
Protein Kinase C α and β compensate for each other to promote stem cell factor-
mediated KIT phosphorylation, mast cell viability and proliferation. FASEB J. 2022
May;36(5):e22273.

72. Hiroki S, Iino Y. The redundancy and diversity between two novel PKC isotypes that
regulate learning in. Proc Natl Acad Sci U S A [Internet]. 2022 Jan 18;119(3). Available
from: http://dx.doi.org/10.1073/pnas.2106974119

73. Trzesniewski J, Altmann S, Jäger L, Kapfhammer JP. Reduced Purkinje cell size is
compatible with near normal morphology and function of the cerebellar cortex in a
mouse model of spinocerebellar ataxia. Exp Neurol. 2019 Jan;311:205–12.

74. Palazzo O, Rass M, Brembs B. Identification of circuits involved in locomotion and
object fixation in. Open Biol. 2020 Dec;10(12):200295.

75. Ehweiner A. PKCi expression in all or FoxPiB postive neurons [Internet]. Universität Regensburg; 2022. Available from: https://epub.uni-regensburg.de/id/eprint/52958

76. Ehweiner A. Knockout of aPKC or PKC53e [Internet]. Universität Regensburg; 2022. Available from: https://epub.uni-regensburg.de/id/eprint/52957

77. Ehweiner A. aPKC knockout [Internet]. Universität Regensburg; 2022. Available from: https://epub.uni-regensburg.de/id/eprint/52944

78. Phelps JS, Hildebrand DGC, Graham BJ, Kuan AT, Thomas LA, Nguyen TM, et al. Reconstruction of motor control circuits in adult Drosophila using automated transmission electron microscopy. Cell. 2021 Feb 4;184(3):759–74.e18.

79. Ehweiner A. Confocal image stack of aPKC/FoxP co-staining [Internet]. Zenodo; 2023. Available from: https://zenodo.org/doi/10.5281/zenodo.10047941

80. Azevedo A, Lesser E, Mark B, Phelps J, Elabbady L, Kuroda S, et al. Tools for comprehensive reconstruction and analysis of *Drosophila* motor circuits [Internet]. bioRxiv. 2022. Available from: http://biorxiv.org/lookup/doi/10.1101/2022.12.15.520299

81. Trimarchi JR, Schneiderman AM. The motor neurons innervating the direct flight muscles of Drosophila melanogaster are morphologically specialized. J Comp Neurol. 1994 Feb 15;340(3):427–43.

82. Whitehead SC, Leone S, Lindsay T, Meiselman MR, Cowan NJ, Dickinson MH, et al. Neuromuscular embodiment of feedback control elements in flight. Sci Adv. 2022 Dec 14;8(50):eabo7461.

83. Heide G, Götz KG. Optomotor control of course and altitude in Drosophila melanogaster is correlated with distinct activities of at least three pairs of flight steering muscles. J Exp Biol. 1996 Aug;199(Pt 8):1711–26.

84. Dickinson MH, Tu MS. The function of dipteran flight muscle. Comp Biochem Physiol A Comp Physiol. 1997 Mar;116(3):223–38.

85. Tu MS, Dickinson MH. The control of wing kinematics by two steering muscles of the blowfly (Calliphora vicina). J Comp Physiol A. 1996 Jun;178(6):813–30.

86. Brembs B. Wild Type Berlin Torque Learning, left+right [Internet]. Universität Regensburg; 2023. Available from: https://epub.uni-regensburg.de/id/eprint/54804

87. Ehweiner A. The neuronal basis of operant self-learning in Drosophila melanogaster [Internet]. Universität Regensburg; 2022. Available from: https://epub.uni-regensburg.de/id/eprint/53425

88. Ehweiner A. GMR65A06-Gal4 x Cas9gFoxP [Internet]. Universität Regensburg; 2022. Available from: https://epub.uni-regensburg.de/id/eprint/52956

89. Ehweiner A. GMR20H05-Gal4 x Cas9gFoxP [Internet]. Universität Regensburg; 2022. Available from: https://epub.uni-regensburg.de/id/eprint/52951

90. Ehweiner A. Ato x Cas9gFoxP [Internet]. Universität Regensburg; 2022. Available
from: https://epub.uni-regensburg.de/id/eprint/52946

91. Ehrhardt E, Whitehead SC, Namiki S, Minegishi R, Siwanowicz I, Feng K, et al. Single-
cell type analysis of wing premotor circuits in the ventral nerve cord of. bioRxiv [In-
ternet]. 2023 Jun 1; Available from: http://dx.doi.org/10.1101/2023.05.31.542897

92. Castells-Nobau A, Eidhof I, Fenckova M, Brenman-Suttner DB, Scheffer-de Gooyert
JM, Christine S, et al. Conserved regulation of neurodevelopmental processes and
behavior by FoxP in Drosophila. PLoS One. 2019 Feb 12;14(2):e0211652.

93. Ehweiner A. gen switch (nSyb) adult Foxp knockout [Internet]. Universität Regens-
burg; 2022. Available from: https://epub.uni-regensburg.de/id/eprint/52963

94. Ehweiner A. nsyb-GS adult Foxp knockout, test after 14 days [Internet]. Universität
Regensburg; 2022. Available from: https://epub.uni-regensburg.de/id/eprint/52964

95. Leslie JH, Nedivi E. Activity-regulated genes as mediators of neural circuit plasticity.
Prog Neurobiol. 2011 Aug;94(3):223–37.

96. Mayford M, Kandel ER. Genetic approaches to memory storage. Trends Genet. 1999
Nov;15(11):463–70.

97. Greenspan RJ. Flies, genes, learning, and memory. Neuron. 1995 Oct;15(4):747–50.

98. Pichaud F. PAR-Complex and Crumbs Function During Photoreceptor Morphogene-
sis and Retinal Degeneration. Front Cell Neurosci. 2018 Mar 29;12:90.

99. Thompson BJ. Par-3 family proteins in cell polarity & adhesion. FEBS J. 2022
Feb;289(3):596–613.

100.Yoshihama Y, Chida K, Ohno S. The KIBRA-aPKC connection: A potential regulator of
membrane trafficking and cell polarity. Commun Integr Biol. 2012 Mar 1;5(2):146–
51.

101.Yoshihama Y, Sasaki K, Horikoshi Y, Suzuki A, Ohtsuka T, Hakuno F, et al. KIBRA sup-
presses apical exocytosis through inhibition of aPKC kinase activity in epithelial
cells. Curr Biol. 2011 Apr 26;21(8):705–11.

102.Sacktor TC. Memory maintenance by PKMζ--an evolutionary perspective. Mol Brain.
2012 Sep 18;5:31.

103.Heitz FD, Farinelli M, Mohanna S, Kahn M, Duning K, Frey MC, et al. The memory
gene KIBRA is a bidirectional regulator of synaptic and structural plasticity in the
adult brain. Neurobiol Learn Mem. 2016 Nov;135:100–14.

104.Ferguson L, Hu J, Cai D, Chen S, Dunn TW, Pearce K, et al. Isoform Specificity of
PKMs during Long-Term Facilitation in Is Mediated through Stabilization by KIBRA. J
Neurosci. 2019 Oct 30;39(44):8632–44.

105. Ji Z, Li H, Yang Z, Huang X, Ke X, Ma S, et al. Kibra Modulates Learning and Memory
via Binding to Dendrin. Cell Rep. 2019 Feb 19;26(8):2064–77.e7.

106. Zlomuzica A, Preusser F, Roberts S, Woud ML, Lester KJ, Dere E, et al. The role of KI-
BRA in reconstructive episodic memory. Mol Med. 2018 Mar 15;24(1):7.

107. Ehweiner A. Baz knockout in all neurons [Internet]. Universität Regensburg; 2022.
Available from: https://epub.uni-regensburg.de/id/eprint/52947

108. Ehweiner A. gKibra adult 2g [Internet]. Universität Regensburg; 2023. Available
from: https://epub.uni-regensburg.de/id/eprint/53685

109. Watanabe A, Li R, Kimura T, Sakaguchi H. Lesions of an avian forebrain nucleus pre-
vent changes in protein kinase C levels associated with deafening-induced vocal
plasticity in adult songbirds. Eur J Neurosci. 2006 May;23(9):2447–57.

110. Sakaguchi H. Effect of social factors on the development of PKC expression in song-
bird brain. Neuroreport. 2004 Dec 22;15(18):2819–23.

111. Yoshida Y, Yamada T, Sakaguchi H. Activation of protein kinase C by the error signal
from a basal ganglia-forebrain circuit in the zebra finch song control nuclei. Neu-
roreport. 2003 Mar 24;14(4):645–9.

112. Watanabe A, Kimura T, Sakaguchi H. Expression of protein kinase C in song control
nuclei of deafened adult male Bengalese finches. Neuroreport. 2002 Jan
21;13(1):127–32.

113. Lorenzetti FD, Baxter DA, Byrne JH. Molecular mechanisms underlying a cellular an-
alog of operant reward learning. Neuron. 2008 Sep 11;59(5):815–28.

114. Dickinson MH. The initiation and control of rapid flight maneuvers in fruit flies. In-
tegr Comp Biol. 2005 Apr;45(2):274–81.

115. Lindsay T, Sustar A, Dickinson M. The Function and Organization of the Motor Sys-
tem Controlling Flight Maneuvers in Flies. Curr Biol. 2017 Feb 6;27(3):345–58.

116. Melis JM, Siwanowicz I, Dickinson MH. Machine learning reveals the control mechan-
ics of an insect wing hinge. Nature. 2024 Apr;628(8009):795–803.

117. Namiki S, Ros IG, Morrow C, Rowell WJ, Card GM, Korff W, et al. A population of de-
scending neurons that regulates the flight motor of Drosophila. Curr Biol. 2022 Mar
14;32(5):1189–96.e6.

118. Palmer EH, Omoto JJ, Dickinson MH. The role of a population of descending neurons
in the optomotor response in flying*Drosophila* [Internet]. bioRxiv. 2022. Available
from: http://biorxiv.org/lookup/doi/10.1101/2022.12.05.519224

119. Wolf R, Heisenberg M. Visual orientation in motion-blind flies is an operant behav-
iour. Nature. 1986 Sep;323(6084):154–6.

120. Mano O, Choi M, Tanaka R, Creamer MS, Matos NCB, Shomar JW, et al. Long-time-
scale anti-directional rotation in optomotor behavior. Elife [Internet]. 2023 Sep
26;12. Available from: http://dx.doi.org/10.7554/eLife.86076

121. Heisenberg M, Wolf R. The sensory-motor link in motion-dependent flight control of
flies. Rev Oculomot Res. 1993;5:265–83.

122. Kassner Z, Dafni E, Ribak G. Kinematic compensation for wing loss in flying damsel-
flies. J Insect Physiol. 2016 Feb;85:1–9.

123. Vance JT, Roberts SP. The effects of artificial wing wear on the flight capacity of the
honey bee Apis mellifera. J Insect Physiol. 2014 Jun;65:27–36.

124. Haas CA, Cartar RV. Robust flight performance of bumble bees with artificially in-
duced wing wear. Can J Zool. 2008 Jul;86(7):668–75.

125. Fernández MJ, Springthorpe D, Hedrick TL. Neuromuscular and biomechanical com-
pensation for wing asymmetry in insect hovering flight. J Exp Biol. 2012 Oct
15;215(Pt 20):3631–8.

126. Zanker JM. How does lateral abdomen deflection contribute to flight control ofDro-
sophila melanogaster? J Comp Physiol A Neuroethol Sens Neural Behav Physiol.
1988 Sep;162(5):581–8.

127. Götz KG, Hengstenberg B, Biesinger R. Optomotor control of wing beat and body
posture in drosophila. Biol Cybern. 1979 Nov;35(2):101–12.

128. Alexandrescu A, Carew TJ. Postsynaptic effects of cysteine-rich neurotrophic factor
in the induction of activity-dependent long-term facilitation in. Learn Mem. 2020
Apr;27(4):124–9.

129. Hu J, Adler K, Farah CA, Hastings MH, Sossin WS, Schacher S. Cell-Specific PKM
Isoforms Contribute to the Maintenance of Different Forms of Persistent Long-Term
Synaptic Plasticity. J Neurosci. 2017 Mar 8;37(10):2746–63.

130. Hu JY, Levine A, Sung YJ, Schacher S. cJun and CREB2 in the postsynaptic neuron
contribute to persistent long-term facilitation at a behaviorally relevant synapse. J
Neurosci. 2015 Jan 7;35(1):386–95.

131. Choi YB, Kadakkuzha BM, Liu XA, Akhmedov K, Kandel ER, Puthanveettil SV. Hun-
tingtin is critical both pre- and postsynaptically for long-term learning-related syn-
aptic plasticity in Aplysia. PLoS One. 2014 Jul 23;9(7):e103004.

132. Jin I, Kandel ER, Hawkins RD. Whereas short-term facilitation is presynaptic, inter-
mediate-term facilitation involves both presynaptic and postsynaptic protein ki-
nases and protein synthesis. Learn Mem. 2011 Feb;18(2):96–102.

133. Reissner KJ, Pu L, Schaffhausen JH, Boyle HD, Smith IF, Parker I, et al. A novel
postsynaptic mechanism for heterosynaptic sharing of short-term plasticity. J Neu-
rosci. 2010 Jun 30;30(26):8797–806.

134.Villareal G, Li Q, Cai D, Fink AE, Lim T, Bougie JK, et al. Role of protein kinase C in the
induction and maintenance of serotonin-dependent enhancement of the glutamate
response in isolated siphon motor neurons of Aplysia californica. J Neurosci. 2009
Apr 22;29(16):5100–7.

135.Glanzman DL. New tricks for an old slug: the critical role of postsynaptic mecha-
nisms in learning and memory in Aplysia. Prog Brain Res. 2008;169:277–92.

136.Bougie JK, Cai D, Hastings M, Farah CA, Chen S, Fan X, et al. Serotonin-induced cleav-
age of the atypical protein kinase C Apl III in Aplysia. J Neurosci. 2012 Oct
17;32(42):14630–40.

137.Cai D, Chen S, Glanzman DL. Postsynaptic regulation of long-term facilitation in Ap-
lysia. Curr Biol. 2008 Jun 24;18(12):920–5.

138.Fulton D, Condro MC, Pearce K, Glanzman DL. The potential role of postsynaptic
phospholipase C activity in synaptic facilitation and behavioral sensitization in Ap-
lysia. J Neurophysiol. 2008 Jul;100(1):108–16.

139.Wolpaw JR, Lee CL. Memory traces in primate spinal cord produced by operant con-
ditioning of H-reflex. J Neurophysiol. 1989 Mar;61(3):563–72.

140.Wolpaw JR, Carp JS, Lee CL. Memory traces in spinal cord produced by H-reflex con-
ditioning: effects of post-tetanic potentiation. Neurosci Lett. 1989 Aug
14;103(1):113–9.

141.Wang Y, Pillai S, Wolpaw JR, Chen XY. Motor learning changes GABAergic terminals
on spinal motoneurons in normal rats. Eur J Neurosci. 2006 Jan;23(1):141–50.

142.Carp JS, Wolpaw JR. Motoneuron plasticity underlying operantly conditioned de-
crease in primate H-reflex. J Neurophysiol. 1994 Jul;72(1):431–42.

143.Carp JS, Chen XY, Sheikh H, Wolpaw JR. Operant conditioning of rat H-reflex affects
motoneuron axonal conduction velocity. Exp Brain Res. 2001 Jan;136(2):269–73.

144.Kaneko N, Sasaki A, Yokoyama H, Masugi Y, Nakazawa K. Effects of action observa-
tion and motor imagery of walking on the corticospinal and spinal motoneuron ex-
citability and motor imagery ability in healthy participants. PLoS One. 2022 Apr
18;17(4):e0266000.

145.Simonyan K, Avetisyan L, Isoyan A, Chavushyan V. Plasticity in Motoneurons Follow-
ing Spinal Cord Injury in Fructose-induced Diabetic Rats. J Mol Neurosci. 2022
Apr;72(4):888–99.

146.Rochat C, Bernard-Marissal N, Källstig E, Pradervand S, Perrin FE, Aebischer P, et al.
Astrocyte-targeting RNA interference against mutated superoxide dismutase 1 in-
duces motoneuron plasticity and protects fast-fatigable motor units in a mouse
model of amyotrophic lateral sclerosis. Glia. 2022 May;70(5):842–57.

147.Wolpaw JR. Harnessing neuroplasticity for clinical applications. Brain. 2012

Apr;135(Pt 4):e215; author reply e216.

148. Eftekhar A, Norton JJS, McDonough CM, Wolpaw JR. Retraining Reflexes: Clinical Translation of Spinal Reflex Operant Conditioning. Neurotherapeutics. 2018 Jul;15(3):669–83.

149. Kane NS, Robichon A, Dickinson JA, Greenspan RJ. Learning without performance in PKC-deficient Drosophila. Neuron. 1997 Feb;18(2):307–14.

150. Drier EA, Tello MK, Cowan M, Wu P, Blace N, Sacktor TC, et al. Memory enhancement and formation by atypical PKM activity in Drosophila melanogaster. Nat Neurosci. 2002 Apr;5(4):316–24.

151. Michel M, Green CL, Lyons LC. PKA and PKC are required for long-term but not short-term in vivo operant memory in Aplysia. Learn Mem. 2011 Jan;18(1):19–23.

152. Cai D, Pearce K, Chen S, Glanzman DL. Protein kinase M maintains long-term sensitization and long-term facilitation in aplysia. J Neurosci. 2011 Apr 27;31(17):6421–31.

153. Lyons LC, Gardner JS, Lentsch CT, Gandour CE, Krishnan HC, Noakes EJ. Differential role of calpain-dependent protein cleavage in intermediate and long-term operant memory in Aplysia. Neurobiol Learn Mem. 2017 Jan;137:134–41.

154. Hu J, Ferguson L, Adler K, Farah CA, Hastings MH, Sossin WS, et al. Selective Erasure of Distinct Forms of Long-Term Synaptic Plasticity Underlying Different Forms of Memory in the Same Postsynaptic Neuron. Curr Biol. 2017 Jul 10;27(13):1888–99.e4.

155. Farley J, Schuman E. Protein kinase C inhibitors prevent induction and continued expression of cell memory in Hermissenda type B photoreceptors. Proc Natl Acad Sci U S A. 1991 Mar 1;88(5):2016–20.

156. Etcheberrigaray R, Matzel LD, Lederhendler II, Alkon DL. Classical conditioning and protein kinase C activation regulate the same single potassium channel in Hermissenda crassicornis photoreceptors. Proc Natl Acad Sci U S A. 1992 Aug 1;89(15):7184–8.

157. Muzzio IA, Talk AC, Matzel LD. Incremental redistribution of protein kinase C underlies the acquisition curve during in vitro associative conditioning in Hermissenda. Behav Neurosci. 1997;111(4):739–53.

158. Blackwell KT. Subcellular, cellular, and circuit mechanisms underlying classical conditioning in Hermissenda crassicornis. Anat Rec B New Anat. 2006 Jan;289(1):25–37.

159. Gao PP, Goodman JH, Sacktor TC, Francis JT. Persistent Increases of PKMζ in Sensorimotor Cortex Maintain Procedural Long-Term Memory Storage. iScience. 2018 Jul 27;5:90–8.

160. Soriano EV, Ivanova ME, Fletcher G, Riou P, Knowles PP, Barnouin K, et al. aPKC Inhibition by Par3 CR3 Flanking Regions Controls Substrate Access and Underpins Apical-Junctional Polarization. Dev Cell. 2016 Aug 22;38(4):384–98.

161. Wieschaus E, Noell E. Specificity of embryonic lethal mutations in Drosophila ana-
lyzed in germ line clones. Rouxs Arch Dev Biol. 1986 Jan;195(1):63–73.

162. Ataman B, Budnik V, Thomas U. Scaffolding proteins at the Drosophila neuromuscu-
lar junction. Int Rev Neurobiol. 2006;75:181–216.

163. Zhang L, Wei X. The Roles of Par3, Par6, and aPKC Polarity Proteins in Normal Neu-
rodevelopment and in Neurodegenerative and Neuropsychiatric Disorders. J Neuro-
sci. 2022 Jun 15;42(24):4774–93.

164. Zha C, Sossin WS. The molecular diversity of plasticity mechanisms underlying
memory: An evolutionary perspective. J Neurochem. 2022 Dec;163(6):444–60.

165. Rocca DL, Wilkinson KA, Henley JM. SUMOylation of FOXP1 regulates transcriptional
repression via CtBP1 to drive dendritic morphogenesis. Sci Rep. 2017 Apr
13;7(1):877.

166. Bowers JM, Perez-Pouchoulen M, Roby CR, Ryan TE, McCarthy MM. Androgen mod-
ulation of Foxp1 and Foxp2 in the developing rat brain: impact on sex specific vocal-
ization. Endocrinology. 2014 Dec;155(12):4881–94.

167. Teramitsu I, White SA. FoxP2 regulation during undirected singing in adult song-
birds. J Neurosci. 2006 Jul 12;26(28):7390–4.

168. Thompson CK, Schwabe F, Schoof A, Mendoza E, Gampe J, Rochefort C, et al. Young
and intense: FoxP2 immunoreactivity in Area X varies with age, song stereotypy,
and singing in male zebra finches. Front Neural Circuits. 2013 Feb 28;7:24.

169. Chen Q, Heston JB, Burkett ZD, White SA. Expression analysis of the speech-related
genes FoxP1 and FoxP2 and their relation to singing behavior in two songbird spe-
cies. J Exp Biol. 2013 Oct 1;216(Pt 19):3682–92.

170. Xiao L, Merullo DP, Koch TMI, Cao M, Co M, Kulkarni A, et al. Expression of FoxP2 in
the basal ganglia regulates vocal motor sequences in the adult songbird. Nat Com-
mun. 2021 May 11;12(1):2617.

171. Kuo HY, Chen SY, Huang RC, Takahashi H, Lee YH, Pang HY, et al. Speech- and lan-
guage-linked FOXP2 mutation targets protein motors in striatal neurons. Brain [In-
ternet]. 2023 May 4; Available from: http://dx.doi.org/10.1093/brain/awad090

1979