1

UBXN1 maintains ER proteostasis and represses UPR activation by modulating translation

independently of the p97 ATPase

Brittany A. Ahlstedt[1], Rakesh Ganji[1], Sirisha Mukkavalli[1*], Joao A. Paulo[2], Steve P. Gygi[2], and

Malavika Raman[1#]

[1] Department of Developmental Molecular and Chemical Biology, Tufts University School of

Medicine, Boston MA.

[2] Department of Cell Biology Harvard Medical School, Boston MA.

* Present address Dana Farber Cancer Research Institute, Boston MA.

Address correspondence to Malavika Raman, malavika.raman@tufts.edu

Abstract

Endoplasmic reticulum (ER) protein homeostasis (proteostasis) is essential to facilitate proper folding and maturation of proteins in the secretory pathway. Loss of ER proteostasis due to cell stress or mutations in ER proteins can lead to the accumulation of misfolded or aberrant proteins in the ER and triggers the unfolded protein response (UPR). In this study we find that the p97 adaptor UBXN1 is an important negative regulator of the UPR. Loss of UBXN1 significantly sensitizes cells to ER stress and activates canonical UPR signaling pathways. This in turn leads to widespread upregulation of the ER stress transcriptional program. Using comparative, quantitative proteomics we show that deletion of UBXN1 results in a significant enrichment of proteins belonging to ER-quality control processes including those involved in protein folding and import. Notably, we find that loss of UBXN1 does not perturb p97-dependent ER associated degradation (ERAD). Our studies indicate that loss of UBXN1 increases translation in both resting and ER-stressed cells. Surprisingly, this process is independent of p97 function. Taken together, our studies have identified a new role for UBXN1 in repressing translation and maintaining ER proteostasis in a p97 independent manner.

Introduction

The ER is a major site for protein maturation and oversees the folding and secretion of one-third of the cellular proteome. A tight equilibrium exists between the ER protein folding capacity and protein load to maintain ER proteostasis. Alterations in this equilibrium due to various physiological conditions such as disruption of ER-calcium homeostasis, mutations that impair protein folding, or cellular aging can result in protein misfolding[1]. ER stress is elicited by protein misfolding and aggregation and can become pathological if not alleviated. Indeed, protein misfolding and aggregation are hallmarks of many age-associated neurodegenerative disorders. Numerous studies indicate that in neurodegenerative disorders such as Parkinson's Disease (PD) and amyotrophic lateral sclerosis (ALS), age associated protein aggregation in the ER is causally linked to cell death[2-5]. Protein aggregates can inappropriately interact with ER chaperones, disrupting ER morphology and vesicular trafficking to the Golgi apparatus[6-8]. In the context of these age-related disorders, it is known that a decline in the efficiency and capacity of protein quality control systems diminishes ER-protein homeostasis. However, it is currently unclear how decline in specific protein quality control mechanisms impacts cell health. Recent studies suggest that aging cells rewire chaperone subnetworks and find that proteasome and autophagy capacities are altered[9-11]. Thus, a thorough understanding of protein quality control mechanisms is warranted to comprehend how their decline triggers disease onset.

Distinct ER-quality control mechanisms sense ER stress. These include the unfolded protein response (UPR) and ER-associated degradation (ERAD) which function to improve the folding capacity of the ER, and clear misfolded proteins from the ER through the ubiquitin-proteasome system respectively. The UPR is an adaptive ER-quality control pathway comprised of three, parallel, ER-resident transmembrane stress sensors that monitor the misfolded protein burden within the ER. These sensors include activating transcription factor 6 (ATF6), inositol requiring enzyme 1 (IRE1), and protein kinase R (PKR)-like ER kinase (PERK). Signaling pathways downstream of each sensor ultimately activates a distinct transcription factor that

induces the expression of a suite of UPR-specific target genes, encoding chaperones and protein

degradation machinery[12]. The prevailing theory suggests that the ER sensors are held in an

inactive state when their luminal domains are bound to the abundant ER chaperone, binding

immunoglobulin protein (BiP) (also known as glucose-regulated protein 78 kDa, GRP78)[13]. When

misfolded proteins accumulate, BiP is titrated away from the sensors, triggering their

activation[14,15]. Under these conditions, ATF6 traffics from the ER to the Golgi apparatus where

site-1 and site-2 proteases release the cytosolic domain of ATF6 to generate an active

transcription factor. In contrast, PERK dimerizes upon activation and phosphorylates eukaryotic

initiation factor 2α (eIF2α), diminishing the translation initiation of most proteins[16]. In response to

eIF2α phosphorylation, the translation of activating transcription factor 4 (ATF4) is selectively

induced in response to reinitiation of ribosomes at the upstream *atf4* coding region. Lastly, IRE1

forms oligomers and mediates the unconventional mRNA splicing of an intron from the x-box

binding protein-1 (*xbp1*) mRNA to generate the active transcription factor xbp1s[17,18]. Additionally,

IRE1 degrades ER-targeted mRNAs in a process termed regulated IRE1-dependent decay

(RIDD)[19]. The rapid degradation of mRNAs through RIDD functions to alleviate ER stress by

preventing the continued influx of new polypeptides into the ER[20,21]. If these mechanisms fail to

alleviate ER stress, PERK and IRE1 signaling will induce apoptosis through expression of the

pro-apoptotic transcription factor C/EBP homologous protein (CHOP) and JNK signaling

respectively[22-24].

The UPR works in tandem with ERAD which purges terminally misfolded clients from the

ER by the ubiquitin-proteasome system. During ERAD, misfolded ER proteins are recognized and

handed off to ER membrane resident E3 ligases to initiate ubiquitin-dependent degradation by

cytosolic proteasomes. The AAA+ ATPase p97 (also known as valosin containing protein, VCP)

is a critical intermediate in this process that recognizes ubiquitylated proteins and retro-

translocates them from the ER to the cytosol prior to proteasomal degradation. The p97 homo-

hexamer utilizes ATP hydrolysis to unfold substrates[25-27] and participates in an array of cellular

pathways from the cell cycle to autophagy[28,29]. p97 interacts with a variety of dedicated adaptor

proteins that facilitate substrate specificity and targeting of p97. The largest family of p97 adaptors

contain a ubiquitin regulatory X (UBX) domain that enables binding to p97[30]. Many of the UBX

domain containing adaptors also harbor a ubiquitin associated domain (UBA) that associates with

polyubiquitin chains on client proteins. There are ~40 adaptors, though the functions and

substrates of most remain poorly characterized. Studies have shown that p97 mutation can

disrupt p97-adaptor binding which has been causally linked to multisystem proteinopathy 1 (MSP-

1), a degenerative disorder wherein individuals present with inclusion body myopathy, Paget's

disease of the bone and/or frontotemporal dementia[31-35]. More recently, p97 mutation has been

linked to the development of familial ALS (fALS), PD, and Charcot-Marie-Tooth disease type IIB

(CMTIIB)[36-39]. A full understanding of adaptor function is required to better understand how their

dysfunction disrupts protein quality control pathways to contribute to disease pathogenesis.

In a previous study, we reported that the p97 adaptor UBX domain protein 1 (UBXN1)

associated with the multifunctional BCL2-associated athanogene 6 (BAG6) chaperone. BAG6

facilitates the insertion of tail anchored proteins into the ER[40-45]. When clients fail to insert (due to

mutations, saturation of insertion machinery at the ER, or ER stress), BAG6 recruits the E3 ligase

RNF126 for ubiquitylation and degradation of the substrate[44]. We showed that UBXN1 was

important for the recruitment of p97 to the BAG6 triage complex to facilitate extraction and

degradation of non-inserted substrates. In addition, BAG6 has also been reported to function in

ERAD in concert with p97 by acting as a chaperone 'holdase' for unfolded p97 clients[46-48].

However, our studies indicated that UBXN1 is not a ERAD adaptor for p97[40]. Here, we have

identified a novel role for UBXN1, as a regulator or ER proteostasis. In this study, we demonstrate

that cells depleted of UBXN1 have elevated UPR activation both in unperturbed and ER-stressed

cells. Using quantitative proteomics, we find a significant increase in the abundance of ER

proteins in UBXN1 deleted cells. Increased steady state protein levels are not due to the

participation of UBXN1 in p97-dependent ERAD. We further demonstrate that loss of UBXN1

increases protein translation in a p97-independent manner and this may be the cause of ER

stress. Our studies have identified a new regulator of protein translation and ER proteostasis.

Results

UBXN1 depletion induces activation of the unfolded protein response.

Based on our prior study on p97-UBXN1 regulation of BAG6-associated protein triage, we

presumed that one consequence of UBXN1 loss may be imbalance in the ER proteome and

subsequent ER stress. Indeed, loss of p97 or BAG6 function can cause the accumulation of

misfolded proteins in the ER via multiple mechanisms to activate the UPR. Silencing BAG6 with

siRNA or inhibiting p97 with the ATP-competitive inhibitor CB-5083 caused robust activation of

the UPR as expected by immunoblotting for the ER chaperone BiP and transcription factor ATF4

(Supplementary Figure 1 A-E). To assess if UBXN1 was also involved in ER stress responses,

we systematically assessed UPR activation in UBXN1 knock-out (KO) HeLa Flp-In T-REx (HFT)

cell lines generated by CRISPR-Cas9 gene editing. Wildtype and UBXN1 KO cells were treated

with the reducing agent dithiothreitol (DTT), and lysates were probed for the ER chaperone BiP,

the transcription factor ATF4, and phosphorylation of eIF2α (Figure 1A and B). In wildtype cells,

a time-dependent increase in these proteins was observed, however a significant increase was

apparent at earlier time points in UBXN1 KO cells. Notably, these markers were already

upregulated in untreated UBXN1 KO cells (Figure 1A, compare lane 5 and 1). We next monitored

the levels of cleaved ATF6 in wildtype and UBXN1 KO cells treated with DTT. Cells were co-

treated with the proteasome inhibitor Bortezomib (BTZ) to prevent turnover of and facilitate

visualization of the N-terminus of ATF6[49]. Compared to control cells, there was a modest but

significant increase in cleaved ATF6 in UBXN1 KO cells in response to ER stress (Figure 1C and

D). Under ER stress conditions, IRE1α is activated by dynamic clustering in the ER

membrane[50,51]. IRE1α monomers cluster into higher order oligomers that can be imaged and

quantified by immunofluorescence. We generated a stable cell line using a previously published

doxycycline-inducible GFP-tagged IRE1α reporter to observe IRE1α clustering[51]. Utilizing this cell

line, IRE1 clustering into oligomers was quantified by quantifying the number of GFP puncta.

UBXN1 was depleted with two independent siRNAs and cells were imaged for GFP-foci (Figure

1E). We found that 40% of cells depleted of UBXN1 contained GFP-IRE1α foci compared to the

10% observed in cells transfected with a control siRNA (Figure 1F). Furthermore, there were

significantly more GFP-foci per cell in UBXN1-depleted cells compared to control (Figure 1F).

Additionally, we measured IRE1 activation via downstream *xbp1s* expression in UBXN1 KO cells

by quantitative real-time PCR using a primer set that spans the splice site removed by IRE1α[52].

We detected 19-fold greater *xbp1s* expression in UBXN1 KO cells compared to control in

response to stress, while total *xbp1* levels remained unchanged (Figure 1G). A similar phenotype

was observed in UBXN1 KO HEK-293T cells, (Supplementary Figure 1F). Notably, the repression

of ER stress by UBXN1 is not due to a role for UBXN1 as a p97 adaptor for ERAD[40,53]. Using a

panel of GFP-tagged ERAD reporters we have previously shown that p97 and its adaptor UBXD8

but not UBXN1 is required for the turnover of multiple ERAD substrates[40].

Our group recently reported that the p97-UBXN1 complex was required for recognizing

ubiquitylated protein aggregates and consolidating them in a perinuclear structure known as the

aggresome[54]. ERAD substrates can also be recruited to these aggregates that can associate with

the ER membrane to cause ER stress. To determine whether UPR activation caused by UBXN1

loss was due to the accumulation of ubiquitylated aggregates we treated wildtype and UBXN1 KO

cells with ER stressors tunicamycin (Tu), which inhibits the first step in N-linked glycosylation, or

thapsigargin (Tg), which inhibits the endoplasmic reticulum Ca^{2+} ATPase. Cells were also treated

with Bortezomib to induce aggresomes as a positive control. We visualized ubiquitylated proteins

by microscopy and immunoblotting. Bortezomib-treated cells contained a single ubiquitin-positive

aggresome that was disrupted in UBXN1 KO cells as we previously reported[54]. However, wildtype

and UBXN1 KO cells treated with Tu or Tg had no ubiquitin positive aggregates or overt

accumulation of ubiquitylated proteins (Supplementary Figure 1G and H). Thus, UPR activation

upon UBXN1 deletion is not due to mishandling of aggregates destined for the aggresome. Taken

together, our findings implicate UBXN1 as a novel regulator of the UPR independent of ERAD.

Loss of UBXN1 activates an ER stress-related transcriptional program leading to cell

death.

The UPR ultimately terminates in a transcriptional response downstream of ATF4, ATF6-

N, and xbp1s, therefore, we analyzed UPR target gene expression using quantitative real-time

PCR of 84 UPR-specific target genes (Supplementary Table 1). Gene expression analysis in

wildtype and UBXN1 KO cells identified a global upregulation (88%, 74 of 84 targets) of UPR-

specific target genes in UBXN1 KO cells with 40 genes (47%) experiencing a $\log_2$ fold change $\geq$

0.5 (Figure 2). While this appears to be a modest change, previous microarray analysis has

demonstrated that only a handful (22 out of 1868 analyzed) of UPR related genes have 2-fold

induction in HeLa cells stressed with tunicamycin[55]. Of the genes that demonstrated a $\log_2$ fold

change $\geq$ 0.5, there are known targets of each of the UPR sensors, including PERK (*ddit3/chop*),

ATF6 (calreticulin, *(calr)* and *herpud1*), and IRE1 (*sec62* and *edem3*), among others[12,56].

Hierarchical clustering analysis demonstrates that the gene expression pattern observed in

UBXN1 KO cells more closely resembles wildtype cells stressed with DTT than untreated wildtype

cells based on similar $\log_2$ fold change values (Figure 2). Additionally, we observed that the

transcriptional response in UBXN1 KO cells treated with DTT is comparable to wildtype cells

treated with DTT (Figure 2). We attribute this to a ceiling effect that is often seen with UPR target

gene expression upon ER stress[57,58].

UPR signaling during acute ER stress is a protective mechanism, whereas chronic ER

stress and UPR activation primes cells toward apoptosis[59-62]. We assessed recovery from ER

stress in wildtype and UBXN1 KO cells. Cells were treated with thapsigargin for a time-course

spanning 0.5 to 3 hours and released into drug-free media (Figure 3A). Viable cells remaining

after three days were stained with crystal violet and quantified. We observed significantly reduced

growth of UBXN1 KO cells relative to wildtype cells suggesting that elevated UPR signaling

primes UBXN1 KO cells for apoptosis (Figure 3A and B). In agreement with this finding, we

observed that UBXN1 KO cells have elevated levels of the apoptosis inducing transcription factor

C/EBP homologous protein (CHOP) compared to wildtype cells in response to DTT (Figure 3C

and D). Thus, UBXN1 loss is detrimental to cell viability in the face of ER stress.

Loss of UBXN1 leads to a perturbed ER proteome

To understand how UBXN1 loss causes ER stress, we first asked if UBXN1 was

associated with the ER. We isolated ER-derived microsomes by biochemical fractionation from

HEK293T cells and subjected the microsomes to proteinase K digestion in the presence and

absence of detergent (Triton X-100) (Figure 3E). As previously reported, a significant fraction of

p97 is peripherally associated with the ER to regulate ERAD[63,64] and this localization is sensitive

to Proteinase K (Figure 3E). Similarly, we found that UBXN1 was also associated with the ER

membrane and was readily digested in Proteinase K treated samples (Figure 3E). We next asked

if UBXN1 associated with the SEC61 translocon to regulate import of proteins into the ER,

however in co-immunoprecipitation studies we were unable to discern an interaction between

UBXN1 and the translocon components SEC61α or β, whereas UBXN1 interacted with p97 as

expected (Supplementary Figure 2A and B).

We decided to take an unbiased approach to assess how UBXN1 deletion impacted ER

protein homeostasis. We employed quantitative tandem mass tag (TMT) proteomics. Peptides

derived from duplicate wildtype and UBXN1 KO cells were labelled with four-plex TMT labels,

combined and peptide abundance was quantified by liquid chromatography and mass

spectrometry (Figure 4A). We quantified a total of 6,673 proteins from this study of which 53 were

significantly enriched in UBXN1 KO cells ($\log_2$ fold change KO:WT ≥ 1.0 and p ≤ 0.5) (Figure 4B,

Supplementary Table 2). Strikingly, gene ontology analysis identified significant enrichment of ER

proteins involved in protein folding, ER-quality control, and the response to ER stress (Figure 4B,

C, Supplementary Figure 3A, B). These include proteins involved in N-linked glycosylation

(glycosyltransferase enzymes ALG1, 3, and 5), ERAD (HRD1 E3 ligase adaptor SEL1L and the

ER-anchored protein escortase TMUB1[65]), and ER import (components of the heterotrimeric

SEC61 complex: SEC61A1, SEC62, and SEC63 and the ER membrane complex, EMC) (Figure

4C and Supplementary Figure 3A). Notably, when we categorized proteins based on their

organelle enrichment, we found that ER proteins in particular were increased in abundance

relative to all other organelles (Figure 4D). In contrast, mitochondrial proteins were depleted in

UBXN1 KO cells (Figure 4D). Thus, UBXN1 localizes to the ER and loss of UBXN1 alters protein

abundance within the ER.

Depletion of UBXN1 increases the expression and aggregation of ER-destined proteins

without preventing their degradation.

To further validate our proteomic studies, we examined the expression of significantly

enriched proteins by immunoblotting lysates from wildtype and UBXN1 depleted cells. In

agreement with the proteomic data, we observed a significant increase in the abundance of the

translocon associated protein α (TRAPα) and alkaline phosphatase placenta-like 2 (ALPP2)

(Figure 5A, B, and G). Interestingly, we found that this phenotype extends beyond our proteomics

hits; we detected a significant increase in the abundance of ER-targeted proteins α-galactosidase

(AGAL)[66] and Prion protein (PrP)[44] in cells transfected with the corresponding epitope tagged

constructs and depleted of UBXN1 (Figure 5C, 5G, Supplementary Figure 4A-C). The elevated

levels of these proteins are not due to increased transcription as *prp* transcript levels were

unchanged between wildtype and UBXN1 KO cells (Supplementary Figure 4C). We asked if the

elevated levels of AGAL and PrP represented ER imported forms or was due to inefficient

cytosolic degradation in the absence of UBXN1. We first monitored the glycosylation status of

FLAG-AGAL and HA-PrP. AGAL glycosylation was verified by EndoH treatment of cell lysates

that resulted in a faster migrating band (Supplementary Figure 4D). HA-PrP glycosylation could

be discerned by three distinctly migrating bands representing the unmodified, ER-resident, and

fully glycosylated post-ER forms (Supplementary Figure 4A). While UBXN1 depletion resulted in

increased steady state levels of these proteins, it did not impact the glycosylation of these proteins

indicating that they were targeted appropriately to the ER (Figure 5C, Supplementary Figure 4A,

B).

Next, we monitored the degradation of HA-PrP and FLAG-AGAL in wildtype and UBXN1

depleted cells. HA-PrP has a weak signal sequence and is mislocalized and degraded prior to

import in response to ER stress[44,46,67]. This can be visualized by the accumulation of a non-

glycosylated cytosolic from in immunoblots (Supplementary Fig 4A). Cells expressing HA-PrP

were treated with thapsigargin and the proteasome inhibitor Bortezomib to induce mislocalization

and prevent degradation respectively[67]. UBXN1 depletion did not induce greater mislocalization

and degradation of HA-PrP relative to total protein (Supplementary Figure 4A compare lane 8 to

4). Furthermore, UBXN1 depletion did not prevent the turnover of FLAG-AGAL or HA-PrP

monitored by cycloheximide chase studies (Supplementary Figure 4E-G). UBXN1 depleted cells

had significantly higher levels of AGAL compared to control cells to begin with (Figure 5C, 5G,

Supplementary Figure 4E), but AGAL was efficiently degraded in both control and UBXN1

depleted cells (Supplementary Figure 4E and F). We note that at earlier timepoints, there was a

delay in AGAL degradation in UBXN1-depleted cells and we attribute this to the overall increase

in protein abundance of AGAL when UBXN1 is depleted (Figure 5C, 5G). We also assessed the

turnover of TRAPα and ALPP2 but these proteins had long half-lives and did not appreciably turn

over after 9 hours of cycloheximide treatment. Nevertheless, increased protein abundance was

obvious (Supplementary Figure 4H).

We asked if increased levels of these protein in the ER impacted their solubility. We

monitored AGAL solubility in control and UBXN1-depleted cells treated with DTT and fractionated

into soluble and insoluble fractions. In control cells, AGAL was soluble and was not found in the

insoluble fraction even upon DTT treatment (Supplementary Figure 4I compare lanes 1 to 3 to 7).

Strikingly, in cells depleted of UBXN1 and treated with DTT, AGAL shifted completely into the

insoluble fraction (Supplementary Figure 4I compare lanes 2 to 4 and 8).

We were next interested if increased protein abundance upon UBXN1 depletion was

unique to ER proteins as suggested by the proteomics (Figure 4D). Our proteomics study

identified several mitochondrial proteins depleted in UBXN1 KO cells, but no change in other

organellar proteins (Figure 4D). Therefore, we investigated by immunoblot if loss of UBXN1

impacted cytosolic, nuclear, or mitochondrial proteins. While we observed no change in the

abundance of proteins in the cytosol or nucleus (Supplementary Figure 4J-M), we observed a

significant decrease in the expression of several mitochondrial proteins when UBXN1 was

depleted (for example, the mitochondrial outer membrane translocon component TOMM20 and

the complex III subunit CYC1) (Figure 5D-G, Figure 4D). Previous studies have demonstrated

that the physical interaction between the ER and mitochondria at contact sites facilitates the

transmission of ER stress to the mitochondria[68,69]. To prevent mitochondrial dysfunction

transmitted by ER stress, translational attenuation through eIF2α phosphorylation mediates the

protective degradation of the TIMM17A subunit of the TIMM23 import complex to prevent protein

import into mitochondria[70]. Whether other mitochondrial proteins are similarly downregulated is

unknown, but our studies suggest that loss of UBXN1 differentially regulates mitochondrial protein

levels in response to ER stress.

Taken together, loss of UBXN1 increases the abundance of ER proteins and leads to their

aggregation upon ER stress.

UBXN1 represses protein translation

Our findings thus far suggested that loss of UBXN1 increased ER protein abundance in a

degradation independent manner. We therefore asked if UBXN1 directly impacted protein

synthesis. We utilized puromycin incorporation into newly synthesized proteins as a proxy for

translation[71]. Puromycin is structurally similar to aminoacyl-tRNAs and can occupy the acceptor

(A) site on a ribosome to be incorporated into the nascent polypeptide[72,73]. Puromycilated proteins

are pre-maturely released from the ribosome and can be visualized by immunoblotting with a

puromycin antibody[74]. Wildtype and UBXN1 KO cells were briefly labelled (~30 min) with

puromycin, and the levels of puromycilated proteins were quantified. We observed a significant

increase in puromycilated proteins in UBXN1 KO cells suggestive of increased translation (Figure

6A, B and Supplementary Figure 5A). We additionally treated cells with DTT or thapsigargin to

induce ER stress and shut down protein translation through eIF2α phosphorylation. DTT,

thapsigargin, or cycloheximide treatment resulted in a complete shutdown of protein synthesis in

wildtype cells as evidenced by the loss in puromycin signal (Figure 6A, compare lanes 7 and 5 to

3, Supplementary Figure 5B compare lanes 3 to 1). However, although significantly reduced in

UBXN1 KO cells, translation could still be observed (Figure 6A, B, Supplementary Figure 5A, B).

To assess the role of p97 in this process, we performed these studies in cells depleted of p97.

Surprisingly, we found that loss of p97 did not impact translation (Figure 6C). In fact, in p97-

depleted and thapsigargin treated cells, there was an even more robust termination of translation

compared to control cells (Figure 6C and D).

To determine the specificity of the phenotype, we next asked if we could rescue the

perturbed translation phenotype by re-expressing wildtype Myc-UBXN1 in UBXN1 KO cells. As

seen in Figure 6E, transfection of Myc-UBXN1 (but not Myc-GFP control) restored translation to

wildtype levels (Figure 6E, compare lanes 3 and 4 to lane 1). However, expressing UBXN1

containing mutations in the UBX domain (Phe^{265} Pro^{266} Arg^{267} truncated to Ala-Gly) that we have

shown previously to disrupt interaction with p97[54], also rescued the increase in protein synthesis

(Figure 6E and F, compare lanes 5 to 1). However, a UBXN1 UBA domain mutant (Met^{13} and

Phe[15] to Ala) that fails to interact with ubiquitin[40,54], did not rescue the increase in protein synthesis

to levels observed in wildtype levels (Figure 6E and F compare lanes 6 to 1).

To further confirm the increased translation phenotype, we isolated polysomes to directly

analyze the amount of actively translating ribosomes present on mRNA. Inhibiting protein

synthesis with cycloheximide halts ribosomes on mRNA such that they can be isolated by sucrose

gradient centrifugation and quantified. We found that UBXN1 KO cells have an increase in the

levels of actively translating polysomes compared to wildtype cells (Figure 6G). Strikingly, while

DTT treatment in wildtype cells halted protein synthesis and collapses polysomes to 80S

monosomes, translation persists in UBXN1 KO cells (Figure 6G). In agreement with our

puromycin incorporation data, depletion of p97 decreased polysomes in untreated cells that was

further depleted upon DTT treatment (Supplementary Figure 5C and D). Taken together UBXN1

represses translation in a p97-independent but ubiquitin-dependent manner (Figure 7).

<u>Discussion</u>

Cellular stress can provoke an imbalance between the protein folding machinery and ER-

protein load. Numerous studies have illuminated how protein misfolding and consequent ER

stress is a major disease mechanism that exacerbates the pathogenesis of a variety of disease

states. Failure to adapt to ER stress has been linked to inflammation and UPR-induced apoptosis

in cardiovascular disease, liver fibrosis, and diabetes mellitus[75-78]. Thus, it is critical to understand

the protein quality control mechanisms that function to prevent ER stress and maintain ER

proteostasis to prevent disease.

In the current study, we have identified a novel role for the p97 adaptor protein UBXN1 in

ER-quality control as a repressor of the UPR (Figure 7). We found that cells devoid of UBXN1

have significant activation of pathways downstream of PERK, IRE1, and ATF6 (Figure 1) resulting

in a transcriptional program to reset ER proteostasis (Figure 2). Our proteomic studies indicate

that UBXN1 KO cells have a perturbed ER-proteome with an enrichment of proteins involved in

protein translocation into the ER, protein folding, ER-quality control, and the response to ER

stress (Figure 4). The increased abundance of these ER proteins is not due to defective

degradation via ERAD (Supplementary Figure 4). Furthermore, increased protein abundance in

the ER triggers aggregation as observed with AGAL. We propose that the increased influx of

proteins into the ER in UBXN1 KO cells causes UPR activation.

A recent study reported that re-initiation of protein synthesis before the restoration of ER

proteostasis can induce cell death in response to ER stress[60]. This study found that forced CHOP

and ATF4 expression after ER stress induces the expression of several aminoacyl-tRNA

synthetases and ribosomal proteins leading to increased translation. Increased protein synthesis

elevated reactive oxygen species (ROS) to trigger ER-stress induced apoptosis[60,61]. Therefore,

the ability to attenuate protein synthesis is paramount to cell viability. Indeed, MEFs harboring an

S51A mutation in eIF2α which prevents phosphorylation of eIF2α by PERK were significantly

sensitized to ER stress, whereas ATF4 null cells, which more slowly restored protein synthesis,

were resistant to ER-stress induced cell death[60]. We find robust elevation in the levels of ATF4

(Figure 1) and CHOP (Figure 3) in UBXN1 null cells and these cells are more sensitive to ER

stress than their wildtype counterparts (Figure 3).

Finally, we show that UBXN1 is a translation repressor and loss of UBXN1 increases

protein translation under basal conditions and in response to ER stress. Whether this occurs

through increased ATF4 expression is under investigation. Our study does not address whether

ER proteins are specifically repressed by UBXN1 or if it represses translation globally. However,

our analysis of the proteomic results and validation studies suggest that ER proteins in particular

are more abundant in UBXN1 deleted cells (Figure 4 and 5). How UBXN1 represses translation

is currently under investigation.

Surprisingly, we observed that the role of UBXN1 in repressing protein translation is

independent of p97 function but reliant on its ability to bind ubiquitin (Figure 6). p97 adaptors are

often thought to function exclusively in complex with p97, however, some studies suggest that

UBXN1 may function in the absence of p97. For example, after infection by RNA viruses, RIG-I-

like receptors in the cytoplasm bind to mitochondrial-antiviral signaling protein (MAVS) to rapidly

induce the expression of type I interferons (IFNs)[79]. UBXN1 is recruited to MAVS where it sterically

inhibits the binding of downstream effector molecules that contribute to IFNs production[80]. This

function was found to be p97 independent as p97 depletion with siRNA, or mutation of the UBX

domain did not impact suppression of these pathways. Only the UBA domain was shown to be

necessary and sufficient to inhibit antiviral signaling[80,81]. Whether other adaptors have functions

independent of the p97 ATPase remains to be elucidated.

Emerging studies indicate the presence of "pre-emptive" ER-quality control systems that

preclude protein misfolding in the ER by preventing the insertion of aberrant proteins. These

pathways act upstream of UPR and ERAD. The BAG6-RNF126 mediated degradation of

mislocalized ER proteins is one such pre-emptive quality control mechanism[40,44,46,67]. Recently,

Ufmylation of the ribosomal protein uL24 (Rpl26) has been reported to enable the degradation of

stalled polypeptides in the translocon[82,83]. It is likely that multiple measures exist to maintain an

optimal ER proteome prior to UPR and ERAD activation but they remain to be fully elucidated.

We propose that UBXN1-mediated translation repression is one such measure that warrants

further investigation.

FIGURE LEGENDS

Figure 1. UBXN1 depletion induces activation of the unfolded protein response.

(A) Immunoblot of HFT wildtype and UBXN1 KO cells treated with 1.5 mM dithiothreitol (DTT) for

the indicated timepoints (0-8 hours) to visualize downstream PERK targets. **(B)** Ratio (UBXN1

KO/wildtype) of the band intensity quantifications of BiP, ATF4, and peIF2α at the 8-hour timepoint

corresponding to Figure 1A. (*n* = three biologically independent samples). We only provide the 8-

hour quantification as the signal is not present in wildtype cells in earlier timepoints for accurate

quantification. **(C)** Immunoblot of HFT wildtype and UBXN1 KO cells co-treated with 1.5 mM DTT

and 1 µM Bortezomib (BTZ) to visualize full-length and cleaved ATF6. **(D)** ATF6 activation was

measured by band intensity quantification and calculation of the percentage of cleaved ATF6 to

total ATF6. The ratio of the percentage of ATF6 activation in UBXN1 KO cells to wildtype is

reported. (*n* = four biologically independent samples). **(E)** Representative immunofluorescent

image of IRE1α clustering in a stable, doxycycline inducible HFT cell line expressing GFP-tagged

IRE1α. UBXN1 was depleted by siRNA transfection and cells were treated with 2.5 µM

tunicamycin (Tu) for 4 hours to induce IRE1α clustering (Scale bar: 10 µm). **(F)** Quantification

corresponding to Figure 1E reporting the percent of cells with GFP-IRE1α foci as well as the

number of GFP-IRE1α foci per cell (*n* = three biologically independent samples). **(G)** Transcript

levels of *xbp1s* and total *xbp1* in HFT wildtype and UBXN1 KO cells quantified by real-time PCR.

Cells were treated with 10 nM thapsigargin (Tg) for 4 hours as indicated. (*n* = three biologically

independent samples). Data are means ± SEM (*, **, ***, **** where *P* < 0.05, 0.01, 0.001, and

0.0001, respectively.) Unpaired two-tailed *t* test **(B, D)** or One-way ANOVA with Tukey's multiple

comparisons test **(F, G)**.

Figure 2. Loss of UBXN1 activates an ER stress transcriptional program.

Heat map displays the hierarchical clustering of the $\log_2$ fold change of 84 UPR target genes

determined by the Human Unfolded Protein Response real-time PCR profiler array. The range is

from a $\log_2$ fold change of -2 (blue) to a $\log_2$ fold change of 4 (red). The untreated wildtype column

is displayed as an average of two replicates with the $\log_2$ fold change set to zero for this column.

Additional columns represent two replicates of untreated UBXN1 KO cells, two replicates of

UBXN1 KO cells treated with 1.5 mM DTT, and two replicates of HFT wildtype cells treated with

1.5 mM DTT for 4 hours respectively. Section 1 and 2 represent UPR target genes that experience

increased expression in the untreated UBXN1 KO cells as well as UBXN1 KO and wildtype cells

treated with DTT. Section 3 represents a set of UPR target genes that experience decreased

expression in all samples compared to untreated wildtype samples. Genes are color coded by the

upstream UPR stress sensor that induces their expression: PERK (green); IRE1 (red); ATF6

(blue); PERK and IRE1 (purple); ATF6 and IRE1 (orange). (n = two biologically independent

samples).

Figure 3. UBXN1 localizes to the ER and preserves cell viability.

(A) Cell viability of wildtype or UBXN1 KO cells was determined by crystal violet staining. Cells

were treated with 200 nM thapsigargin (Tg) for the indicated timepoints and released into drug

free media. Colonies were allowed to grow out for three days and live cells were stained with

crystal violet. **(B)** Percent viability calculated from the remining cells in Figure 3A. (n = four

biologically independent samples) **(C)** Immunoblot of HFT wildtype and UBXN1 KO cells treated

with 1.5 mM dithiothreitol (DTT) for the indicated timepoints (0-8 hours) to visualize CHOP

expression. **(D)** Band intensity quantifications of CHOP expression at the 8-hour timepoint

corresponding to Figure 3C. (n = three biologically independent samples)**.** We only provide the 8-

hour quantification as the signal is not present in wildtype cells in earlier timepoints for accurate

quantification. **(E)** Subcellular fractions enriched in ER-derived microsomes were isolated from

HEK-293T cells by biochemical fractionation. Protease protection assay was used to localize

UBXN1 to the ER periphery by immunoblot. Calnexin and Sec61β are ER specific markers, β-

actin is a cytosolic marker, and TOMM20 is a mitochondrial marker. (PNS: post-nuclear

supernatant sample, cyto: cytosolic sample). Data are means ± SEM (*, **, **** where $P < 0.05$,

0.01, and 0.0001, respectively.) Unpaired two-tailed t test **(B and D)**

Figure 4. Loss of UBXN1 leads to a perturbed ER proteome.

(A) Schematic workflow of quantitative tandem mass tag (TMT) proteomics in HFT wildtype and

CRISPR-Cas9 UBXN1 knock-out (KO) cells. **(B)** Volcano plot of the ($-\log_{10}$-transformed P value

versus the $\log_2$-transformed ratio of wildtype/UBXN1 KO) proteins identified by TMT proteomics

in wildtype and UBXN1 KO cells. A negative $\log_2$ fold change indicates proteins with increased

abundance in UBXN1 KO cells. Proteins with a $\log_2$ fold change $\geq |1|$ are considered significant.

ER membrane and luminal proteins are outlined by a red and blue circle respectively. Proteins

involved in translocation into the ER and glycosylation are colored in green and blue respectively

(n = 2 biologically independent samples for each genotype). **(C)** Metascape gene ontology

network of the proteins identified to be enriched in UBXN1 KO cells grouped by functional ontology

category. The size of the node represents the number of genes in that gene ontology cluster and

the thickness of the circle outline corresponds to the $-\log_{10}$-transformed P value of each term.

The different circle colors correspond to a specific gene ontology category. **(D)** Proteins identified

in the quantitative proteomics study categorized by organellar compartment. Proteins associated

with the ER experience a significant increase in abundance in UBXN1 KO cells where

mitochondrial proteins are significantly decreased. Data are means ± SEM (**** where $P <$

0.0001). One-way ANOVA with Brown-Forsythe and Welch ANOVA tests of -logFC **(D)**.

Figure 5. Depletion of UBXN1 increases the expression of ER-destined proteins.

(A and B) Representative immunoblots of proteins identified to be enriched in the proteomics

study. The ER and secretory proteins TRAPα and ALPP2 were measured in control cells or cells

depleted of UBXN1 for 48 hours with siRNA. Cells were either untreated or treated with 10 mM

DTT for 4 hours. **(C)** Immunoblot of a stable HFT cell line expressing doxycycline inducible FLAG-

tagged α-galactosidase (AGAL). UBXN1 was depleted with siRNA for 24 hours before AGAL

induction by 2 μg/ml doxycycline for 24 hours. Cells were untreated or treated with 10 mM DTT

for 4 hours. **(D-F)** Representative immunoblots of the mitochondrial resident proteins Aconitase-

2 (ACO2), TOMM20, and CYC1 measured in control or UBXN1 depleted cells (48 hours) either

untreated or treated with 10 mM DTT for 4 hours. **(G)** Band intensity quantifications of protein

expression corresponding to Figure 5A-F. (n = three biologically independent samples). Data are

means ± SEM (*, **, ***, **** where *P* < 0.05, 0.01, 0.001, and 0.0001, respectively.) One-way

ANOVA with Tukey's multiple comparisons test **(G)**.

Figure 6. UBXN1 represses protein translation.

(A) Representative immunoblot of puromycin incorporation into HFT wildtype and UBXN1 KO

cells. Cells were pulsed with 1 μM puromycin or 1 μM puromycin in combination with 10 μg/mL

cycloheximide (CHX) for 30 minutes. When applicable, cells were pretreated with 1.5 mM DTT

for 1 hour before 1 μM puromycin pulse for 30 minutes. The level of puromycin incorporation

reflects the rate of protein synthesis. **(B)** Puromycin, DTT, and CHX plots correspond to the

intensity of each lane in the immunoblot in Figure 6A. The intensity of the traces for each wildtype

and UBXN1 KO sample was determined by the plot profile feature in Fiji. The x-axis represents

the distance along the lane. **(C)** Representative immunoblot of 1 μM puromycin incorporation into

control cells or cells depleted of p97 with siRNA for 48 hours. Cells were either untreated or pre-

treated with 1 μM thapsigargin for 30 minutes. **(D)** Quantification of the whole lane corresponding

to each lane in the immunoblot in Figure 6C (*n* = four biologically independent samples). **(E)** Myc-

tagged GFP, wildtype, UBX domain mutant, or UBA domain mutant UBXN1 was expressed in

cells for 48 hours before 1 μM puromycin pulse for 30 minutes. **(F)** Quantification of each lane

intensity from Figure 6E. (*n* = three biologically independent samples) **(G)** Polysome profile traces

of HEK-293T wildtype and UBXN1 KO cells either untreated or treated with 2 mM DTT for 60

minutes. The 40S, 60S, and 80S ribosomal subunits are labeled as well as actively translating

polysomes. Traces below correspond to the polysomes seen in the main trace. Data are means

± SEM (**** where *P* < 0.0001.) One-way ANOVA with Tukey's multiple comparisons test **(D, F).**

Figure 7. UBXN1 represses protein translation to prevent ER stress and UPR activation.

UBXN1 plays an important role in ER-quality control to maintain ER proteostasis. Loss of UBXN1

increases protein synthesis. Increased abundance of ER proteins leads to protein misfolding and

activation of the UPR.

Supplementary Figure 1. BAG6 depletion and pharmacological inhibition of p97 induces

activation of the unfolded protein response.

(A) Immunoblot of BiP and ATF4 expression levels in HFT wildtype cells treated with 1 μM

thapsigargin (Tg), 5 μM of the ATP-competitive p97 inhibitor CB-5083, or both for the indicated

timepoints (0-6 hours). **(B-C)** Band intensity quantifications of BiP **(B)** and ATF4 **(C)**

corresponding to Supplementary Figure 1A reporting the fold change compared to the untreated

condition (*n* = three biologically independent samples)**. (D)** Immunoblot of HFT wildtype cells

depleted of BAG6 with siRNA for 48 hours before treatment with 1.5 mM DTT for the indicated

timepoints (0-8 hours). **(E)** Band intensity quantifications of BiP and ATF4 corresponding to

Supplementary Figure 1D. at the 8-hour timepoint. (*n* = four biologically independent

experiments). **(F)** Transcript levels of *xbp1s* and total *xbp1* in HEK-293T wildtype and UBXN1 KO

cells quantified by real-time PCR. Cells were treated with 10 nM thapsigargin (Tg) for 4 hours as

indicated (*n* = three biologically independent samples). **(G)** Immunofluorescent staining of

ubiquitin (FK2) and nuclei (Hoechst dye) in HFT wildtype and UBXN1 KO cells. Cells were treated

with 1 μM of the proteasome inhibitor bortezomib (BTZ), Tunicamycin (Tu), or Thapsigargin (Tg)

for 8 hours or 1 μM BTZ, 500 nM Tu, or 500 nM Tg for 18 hours. **(H)** Immunoblot assessing total

ubiquitin levels corresponding to Supplementary Figure 1G. Data are means ± SEM (*, **, ***, ****

where *P* < 0.05, 0.01, 0.001, and 0.0001, respectively.) One-way ANOVA with Dunnetts's multiple

comparisons test **(B and C).** Unpaired two-tailed *t* test **(E).** One-way ANOVA with Tukey's multiple

comparisons test **(F)**.

Supplementary Figure 2. UBXN1 does not interact with the Sec61 translocon.

(A) Representative immunoblot for the immunoprecipitation of endogenous UBXN1 from HEK-293T lysates. 10 and 20 µg of whole cell lysate was used for the input. Sec61β was used as a marker for the Sec61 translocon. **(B)** UBXN1 and Sec61β were separately immunoprecipitated from ER-microsomes isolated from HEK-293T cells. 10 and 20 µg of whole cell lysate was used for the input.

Supplementary Figure 3. Loss of UBXN1 leads to a perturbed ER proteome.

(A) Interaction network of the gene ontology clusters from the wildtype/UBXN1 KO quantitative proteomics study with a significant increase in abundance in UBXN1 KO cells. Proteins found in each node are labelled and the color of each circle corresponds to the $-\log_{10}$-transformed P value. **(B)** Bubble plot corresponding to significantly enriched gene ontology groups in the wildtype (navy) and UBXN1 KO (teal). The size of the circle corresponds to the number of genes identified in each term.

Supplementary Figure 4. Depletion of UBXN1 increases the expression of ER-destined proteins.

(A) Immunoblot of HFT wildtype cells expressing HA-tagged prion protein (PrP). PrP was expressed in HeLa cells for before siRNA mediated depletion of UBXN1. Cells were treated with 500 nM thapsigargin, 1 µM bortezomib (BTZ), or both for 16 hours. The percent mislocalization was calculated from the ratio of the mislocalized (cytosolic) form to all forms of PrP for that sample. **(B)** Total PrP protein expression was calculated by the sum of all forms of PrP for that sample. The untreated condition was quantified (lanes 1 and 5) (n = three biologically independent experiments). **(C)** Transcript levels of HA-PrP in cells depleted of UBXN1 quantified by real-time PCR. Control is cDNA generated from cells not expressing HA-PrP. (n = three biologically independent experiments). **(D)** Immunoblot of a stable HFT cell line expressing doxycycline

inducible FLAG-tagged AGAL. Glycans were digested with EndoH for 4 hours to generate the

non-glycosylated form. **(E)** Immunoblot examining the turnover of FLAG-AGAL in siRNA control

and UBXN1 depleted cells (48 hours). Cells were treated with 10 μg/mL cycloheximide (CHX) for

the indicated timepoints (0-9 hours). **(F)** Band intensity quantification of each time point of the

CHX time course corresponding to Supplementary Figure 5E. The zero hour time point was set

to 100 % for each siRNA and the % of AGAL remaining for each time point was quantified as a

ratio from the zero hour timepoint for that siRNA. ($n \geq$ three biologically independent samples. **(G)**

Immunoblot examining the turnover of HA-PrP in siRNA control and UBXN1 depleted cells. Cells

were treated with 10 μg/mL cycloheximide (CHX) for the indicated timepoints (0-8 hours). **(H)**

Immunoblot examining the turnover of the TMT-proteomics hits (ALPP2 and TRAPα) that were

identified to be significantly enriched in cells depleted of UBXN1. Control and UBXN1 depleted

cells were treated with 10 μg/mL cycloheximide (CHX) for the indicated timepoints (0-9 hours). **(I)**

Immunoblot of soluble and insoluble fractions of FLAG-tagged AGAL in control cells or cells

depleted of UBXN1 with siRNA. Cells were either untreated or treated with 10 mM DTT for 4

hours. (In this experiment only, we quantified and ran 2.5x less siUBXN1 protein so we could

accurately visualize and compare siControl to siUBXN1. The AGAL expression in siUBXN1 cells

often bleaches out what can be observed with siControl.) **(J-L)** Immunoblots of samples from

Figure 5A. examining the protein expression of additional organellar proteins including the nuclear

protein MCM3 and the cytosolic proteins G3BP1 and HPS90. **(M)** Band intensity quantifications

of protein expression corresponding to Supplementary Figure 5J-L. (n = three biologically

independent samples). Data are means ± SEM (*, **, and *** where P < 0.05, 0.01, and 0.001

respectively.) Unpaired two-tailed t test **(B and F)**, One-way ANOVA with Tukey's multiple

comparisons test **(C and M)**.

Supplementary Figure 5. UBXN1 suppresses protein synthesis.

(A) Densitometry quantifications of the entire lanes corresponding to Figure 6A ($n \geq$ three

biologically independent experiments). **(B)** Immunoblot of HFT wildtype and UBXN1 KO cells

pulsed with 1 μM puromycin for 30 minutes after pre-treatment with 1.5 μM thapsigargin for 30

minutes. **(C)** Polysome profile traces of HEK-293T control cells or cells depleted of p97 with siRNA

for 72 hours. Cells were either untreated or treated with 2 mM DTT for 60 minutes. The 40S, 60S,

and 80S ribosomal subunits are labeled as well as actively translating polysomes. Traces on the

right correspond to the polysomes seen in the traces on the left. **(D)** Immunoblot validating the

siRNA knockdown corresponding to Supplementary Figure 6C. Data are means ± SEM (*** and

**** where $P < 0.001$ and 0.0001, respectively.) One-way ANOVA with Tukey's multiple

comparisons test **(A)**.

MATERIALS AND METHODS

Cell culture, transfections, immunoblot, and immunoprecipitation

HeLa Flp-IN T-REX (HFT) and HEK-293T cells were cultured in Dulbecco's modified Eagle's

medium (DMEM) supplemented with 10 % fetal bovine serum (FBS) and 100 U/mL penicillin and

streptomycin. Cells were grown in a humidified, 5 % CO_2 atmosphere at 37 $^{\circ}$C. For siRNA

transfections, cells were either forward or reverse transfected with 20 nM siControl, siUBXN1,

siBAG6, or sip97 using Lipofectamine RNAiMax (Invitrogen) in 6-well plates. After 8 or 24 hours

depending on the study, cells were split into 12-well plates for treatment. 48 hours post

transfection, cells were treated with the indicated drugs for the indicated timepoints before harvest

for immunoblot or fixation for immunofluorescence. For DNA transfections, 1 μg HA-prion protein

(PrP), 0.5 μg Myc-tagged wildtype UBXN1, 0.5 μg of either Myc-tagged UBX mutant (Phe^{265} Pro^{266}

Arg^{267} truncated to Ala-Gly), or Myc-tagged UBA mutant (Met^{13} and Phe^{15} to Ala), and 50 ng Myc-

tagged GFP constructs were forward transfected into HFT cells seeded in a 6-well plate using

Lipofectamine 3000 (Invitrogen). The cells were harvested 48 or 72 hours post transfection. To

establish an HFT cell line stably expressing doxycycline inducible FLAG-AGAL, 1 μg FLAG-AGAL

construct and 4.5 μg pOG44 was transfected into HFTs seeded onto a 6-well plate using

Lipofectamine 2000 (Invitrogen). After 48 hours the transfected cells were selected with

hygromycin (200 μg/mL) for about 10 days.

Cell pellets were lysed in either radioimmunoprecipitation assay (RIPA) buffer (25 mM

Tris-Cl [pH 7.6], 150 mM NaCl, 1 % NP-40, 1 % sodium deoxycholate, 0.1 % SDS, HALT protease

inhibitors (Pierce PI-78425), sodium vanadate, sodium fluoride) or in 1 % SDS lysis buffer (1 %

SDS, 50 mM Tris-Cl, 150 mM NaCl, 0.5 % NP-40, HALT protease inhibitors, sodium vanadate,

sodium fluoride). FLAG-AGAL and HA-PrP cell pellets were resuspended in RIPA buffer and

incubated on ice for 20 minutes with a 10 second vortex pulse every 5 minutes. Lysates were

then centrifuged at 14,000 rpm for 15 minutes at 4 °C. For UPR induction studies in Figure 1, cell

pellets were resuspended in SDS lysis buffer, vortexed briefly, boiled for 10 minutes at 95 °C, and

briefly vortexed. Lysates were incubated at 65 °C for an additional 5 minutes before centrifugation

at 14,000 rpm at room temperature. The supernatant was collected, and protein concentration

was estimated utilizing the DCA protein assay kit (Biorad). Protein expression was analyzed by

SDS-PAGE and immunoblot.

For immunoprecipitations, cell pellets were lysed in mammalian cell lysis buffer (50 mM

Tris-HCl [pH 7.6], 150 mM NaCl, 1 % Nonidet P-40, HALT protease inhibitors (Pierce), and 1 mM

DTT). Protein G agarose beads (Thermo Scientific 20398) was pre-washed with MCLB before

addition of the cell lysate and antibody (UBXN1, Sec61β, or rabbit IgG control). Samples were

incubated with rotation overnight at 4 °C. Beads was collected by centrifugation at 3000 rpm for

1 minute and was washed four times with MCLB buffer. Beads were resuspended in 50 μl SDS

sample buffer, briefly boiled, and processed for SDS-PAGE and immunoblotting.

Antibodies, siRNA, and reagents

The rabbit UBXN1 (16135-1-AP 1:7000 dilution), p97 (10736-1-AP), BAG6 (26417-1-AP), ATF6

(24169-1-AP), Sec61β (51020-2-AP 1:3000 dilution), Aconitase 2 (11134-1-AP), calnexin (10427-

2-AP), CYC1 (10242-1-AP), and G3BP1 (13057-2-AP) antibodies were obtained from Proteintech

Inc; rabbit BiP (3177 dilution), peIF2α (3398 1:500 dilution), and MCM3 (4012S) antibodies were

obtained from Cell Signaling Technology; mouse PCNA (sc-56 1:3000 dilution), ATF4 (sc-390063

1:500 dilution), Tom20 (sc-17764), ubiquitin (P4D1; sc8017), β-actin (sc-69879 1:3000 dilution),

GAPDH (sc-47724 1:3000 dilution), c-myc (sc-40 1:3000 dilution), Sec61α (sc-393182 1:500

dilution), and TRAPα (sc-373916) were from Santa Cruz Biotechnology; mouse ubiquitin FK2 (04-

263 1:100 dilution) for immunofluorescence and Anti-Puromycin, clone 12D10 (MABE343 1:3000

dilution) was from EMD Millipore; mouse FLAG-M2 antibody (F3165 1:1000 dilution) was from

Sigma Aldrich. The rabbit anti-Prion protein PrP antibody (ab52604) was purchased from Abcam.

The rabbit ALPP/ALPP2 was purchased from St. John's Laboratory (STJ96740). The mouse

HSP90AB1 was purchased from Origene (TA500494). All antibodies were used at a dilution of

1:1000 unless otherwise specified. Secondary antibodies, HRP conjugated anti-rabbit (W4011)

and anti-mouse (W4021) were from Promega and used at a dilution of 1:10000. Thapsigargin

(5860051MG), Tunicamycin (ICN15002801), Proteinase K (FEREO0491), and Hoechst (51-17

1:10000 dilution) for immunofluorescence were purchased from Fisher Scientific. Dithiothreitol

(DTT25) and puromycin (P-600-100) were purchased from GoldBio. Cyclohexmide (97064-724)

and Triton X-100 (97063-864) was purchased from VWR. CB-5083 were purchased from Cayman

Chemical Company (19311). Bortezomib (S1013) was purchased from Selleckchem. EndoH

(P0702S) was purchased from New England Biolabs. Crystal violet was purchased from Sigma-

Aldrich (C6158-50G). Doxycycline was purchased from EMD Millipore (324385-1GM). Wild-type

PrP (HA) was generously gifted by Ramanujan Hegde (MRC, UK). FLAG-tagged AGAL were

generously gifted by Malaiyalam Mariappan[66]. siRNAs against UBXN1-2 (D-008652-02), UBXN1-

3 (D-008652-03), and UBXN1-4 (D-008652-04) were purchased from Dharmacon. siRNA against

BAG6 (s15467) was purchased from Ambion (Themo Fisher Scientific). siControl (SIC001) was

from Millipore Sigma. siRNA against p97 was purchase from Invitrogen (HSS111263).

Generation of CRISPR gene knockout cell lines

UBXN1 knockout cells were generated in both HFT and HEK-293T cells using the CRISPR-Cas9

gene editing method. The guide sequence 5′-GCCGTCCCAGGATATGTCCAA-3′ was cloned into

the pX459 vector carrying hSpCas9 as previously published[40]. 500 ng of the construct was

transiently transfected into HFT and HEK-293T cells seeded in a 6-well plate with Lipofectamine

3000. 36 hours post-transfection, the cells were cultured with 1 μg/mL puromycin for an additional

24 hours. The surviving cells were counted and serially diluted to a concentration of 1 cell per well

in a 96-well plate. Protein expression of UBXN1 was examined by immunoblot.

Crystal violet cell viability assay

Wildtype and UBXN1 KO cells were plated at a density of 8000 cells/well in a 12-well dish. The

next day, cells were treated with 200 nM thapsigargin for the indicated timepoints. After treatment,

wells were washed twice with PBS before re-addition of fresh media. Every day for three days,

cells were washed once with PBS and media was replaced. On the third day, cells were washed

twice with PBS before incubation with 0.5 % crystal violet staining solution for 20 minutes at room

temperature on an orbital shaker. After staining, wells were washed thrice with PBS to remove

unbound stain. 1 % SDS-solution was added to each well to solubilize bound stain and the plate

was incubated for one hour at room temperature on an orbital shaker until color was uniform in

solution. The absorbance was read at 570 nm on a multiwell plate reader (SpectraMax iD3). The

average absorbance of wells without cells was used to background subtract.

Real-time PCR

For all real-time PCR experiments, total RNA was isolated using the *Quick*-RNA Miniprep Kit

(Zymo Research cat. no. R1055). The purified RNA was quantified by NanoDrop and 1 µg of RNA

for each sample was used to generate cDNA using the iScript cDNA synthesis kit (Biorad cat. no.

1708890). Real-time PCR was performed with PowerUp SYBR Green Master Mix (Applied

Biosystems cat. no. A25741) on an Applied Biosystems StepOnePlus real-time PCR system. Data

analyses utilized the $2^{-\Delta\Delta Ct}$ method and GAPDH was used as a housekeeping gene to normalize

transcript expression across samples. The XBP1s primers were previously published[52].

XBP1s forward: 5′ TGCTGAGTCCGCAGCAGGTG 3′

XBP1s reverse: 5′ GCTGGCAGGCTCTGGGGAAG 3′

Total XBP1 forward: 5′ AAACAGAGTAGCTCAGACTGC 3′

Total XBP1 reverse: 5′ TCCTTCTGGGTAGACCTCTGGGAG 3′

HA-PrP forward: 5' CAGCCTCATGGTGGTGGCTG 3'

HA-PrP Reverse: 5' GTCGGTCTCGGTGAAGTTCTCC 3'

RT² Profiler PCR array

Wildtype or UBXN1 KO HeLa Flp-in T-REX cells were grown in 10cm tissue culture dishes and

either left untreated or treated with 1.5 mM DTT for 4 hours. Post-harvest, RNA was extracted

from each sample with the *Quick*-RNA MiniPrep Kit (Zymo Research cat. no. R1055) and cDNA

was generated with the RT² First Strand Kit (Qiagen cat no. 330401). Each Human Unfolded

Protein Response RT² Profiler PCR Array (cat no. 330231 PAHS-089ZA) plate consists of 84

UPR specific target genes as well as a panel of housekeeping genes and PCR efficiency controls.

Each sample was distributed to an individual plate for two biological replicates each. The

GeneGlobe Data Analysis Center was used to determine the fold change between samples. The

data were normalized by automatic selection from the housekeeping gene panel utilizing the

geometric mean. The untreated wildtype HFT samples were defined as the "control" sample for

the data set and used to calculate fold changes. Heatmap visualizations for PCR array data were

performed using the RStudio software (v2022.07.1 Build 554), including ggplot2 (v 3.3.5), circlize

(v0.4.14), ComplexHeatmap version 2.10.0, and InteractiveComplexHeatmap package to export

heatmap generated by ComplexHeatmap into an interactive Shiny app[84].

Immunofluorescence and microscopy

To assess aggresome formation, cells were grown on coverslips (12mm) in a 24-well plate treated

with 1 µM bortezomib (Btz), Tunicamycin (Tu), or Thapsigargin (Tg) for 8 hours or 1 µM Btz, 500

nM Tu, or 500nM Tg for 18 hours. Cells were washed briefly in phosphate-buffered saline (PBS)

and fixed with ice-cold methanol for 10 min at −20°C. Cells were washed in PBS then blocked in

2 % bovine serum albumin (BSA) with 0.3 % Triton X-100 in PBS for 1 hour. The coverslips were

incubated with the indicated antibodies in a humidified chamber overnight at 4 °C. Coverslips

were washed and incubated with the appropriate Alexa Fluor-conjugated secondary antibodies

(Molecular Probes) for 1 hour in the dark at room temperature. Cells were washed with PBS, and

nuclei were stained with Hoechst 33342 dye and mounted onto slides. All images were collected

using a Nikon A1R scan head with a spectral detector and resonant scanners on a Ti-E motorized

inverted microscope equipped with a 60× Plan Apo 1.4 NA objective lens. The indicated

fluorophores were excited with either a 405-nm, 488-nm or 594-nm laser line. Images were

analyzed by using FIJI (https://imagej.net/Fiji).

IRE1α-GFP foci imaging and quantification

Stable HFT wildtype cells expressing doxycycline inducible IRE1α-3xFLAG-6xHis-GFP (construct

a generous gift from Peter Walter, UCSF) were seeded onto a 6-well plate and reverse transfected

with an siControl or siUBXN1[51]. Next day, cells were trypsinized and seeded onto coverslips in a

12-well plate. 48 hours after siRNA transection, IRE1α-GFP was induced with 4 µg/mL

doxycycline for an additional 24 hours. Next day, cells were either left untreated or treated with

2.5 µM tunicamycin for 4 hours. Coverslips were briefly washed with phosphate-buffered saline

(PBS) and fixed with ice-cold methanol for 10 min at -20 $^{\circ}$C. Coverslips were washed with PBS

and nuclei were stained with Hoechst 33342 dye and mounted on slides. Images were collected

using either a 405-nm or 488-nm laser. One hundred cells were counted by creating a cell mask

and utilizing the "Find Maxima" function in FIJI (https://imagej.net/Fiji).

Subcellular fractionation of ER-derived microsomes

HEK-293T cells were seeded into four 150 mm sterile tissue culture dishes. Cells were scraped

off the plate into PBS, centrifuged at 600 x g for 5 minutes at 4 $^{\circ}$C, and resuspended in ice-cold

homogenization buffer (225 mM Mannitol, 75 mM Sucrose, 30 mM Tris [pH 7.4]). Cells were

homogenized with a Dounce homogenizer on ice. Every 150 passes of the homogenizer, cells

were centrifuged for 5 minutes at 600 x g, to remove non-lysed cells and nuclei. The integrity of

the homogenized cells was examined under a light microscope until 80-90 % cell lysis was

attained. The supernatant was collected as the post-nuclear supernatant (PNS) fraction. The

clarified lysate was centrifuged at 7000 x g for 10 min at 4 $^{\circ}$C, isolating the supernatant including

ER-microsomes. The supernatant was collected, centrifuged at 20,000 x g for 30 mins at 4 $^{\circ}$C to

clear lysosomal and plasma membrane fractions. The resulting supernatant was further

centrifuged at high speed, 100,000 x g for 1 hour in a swinging-bucket rotor SW-41 and Beckman

Coulter ultracentrifuge. This results in the isolation of pure ER-microsomes (pellet) and cytosolic

fraction (supernatant). PNS, cytosolic, and ER-microsomes were solubilized using 0.5 % (v/v)

SDS and protein concentrations were determined by DC protein assay. An equivalent amount of

ER-microsomes were used in the protease protection assay. Of four aliquots, one was left

untreated, one was incubated with 1 % (v/v) Triton X-100, and the other two were incubated with

10 µg/mL proteinase K with or without 1 % (v/v) Triton X-100. All samples are incubated at 37 $^{\circ}$C

for 10 minutes before reaction termination with 2 mM phenylmethylsulfonyl fluoride (PMSF). 5X-

Laemmli sample buffer was added to each sample before SDS-PAGE and immunoblot.

Puromycilation translation assay

Protein synthesis was measured in HFT wildtype and UBXN1 KO cells utilizing the non-

radioactive SUnSET assay[74]. HFT wildtype or UBXN1 KO cells were seeded onto a 6-well plate

at 400 k cells /well to achieve 70-80 % confluence at the time of puromycin addition. Cells were

incubated with 1 µM puromycin with or without 100 µg/ml cycloheximide for 30 minutes in

complete medium. Samples treated with DTT or thapsigargin were pre-incubated with 1.5 mM

DTT for 1 hour or 1 µM thapsigargin for 30 minutes before co-treatment of 1 µM puromycin with

1.5 mM DTT for 30 minutes in complete medium. At the end of the incubation period, cells were

washed with ice-cold PBS and the plate was immediately placed on ice. Ice-cold RIPA buffer was

added to the wells and cells were gently scraped into the buffer using a cell scraper. Lysates were

placed on ice and vortexed for 10 seconds every 5 minutes for 30 minutes before centrifugation

at 12,000 rpm for 10 minutes at 4 °C. Protein concentration was estimated using the DC assay

(Bio-Rad). An equivalent amount of protein for each sample were resolved by SDS-PAGE and

probed with the anti-puromycin antibody, clone 12D10.

Polysome profiling

HEK-293T wildtype and UBXN1 KO cells were plated in a 150 mm dish and allowed to grow for

two days before polysome isolation. For p97 depletion experiments, HEK-293T cells were plated

in a 150 mm dish and forward transfected next day with siRNA against p97 or an siControl. 24

hours later the plate was split into two 150 mm dishes and the cells were grown for two additional

days. The day before profiling, sucrose gradients were prepared with fresh 15 % and 60 %

sucrose solutions using 1X low-salt buffer (5 mM Tris-HCl [pH 7.5], 2.5 mM $MgCl_2$, 1.5 mM KCl,

1 mM PMSF). To generate a continuous gradient, 5 mL of 60 % sucrose solution was placed at

the bottom of an ultra-clear Beckman Coulter centrifuge tube. 5 mL of 15 % sucrose solution was

layered on top and the gradient was inverted horizontally and left on its side at room temperature

for 3 hours. After 3 hours, the tube was brought back to its vertical position and stored at 4 °C

overnight. When cells reached about 70-80 % confluency, cells were either pre-treated with 2 mM

DTT for 60 minutes or left untreated. After treatment, cells were incubated with 100 ug/mL

cycloheximide at 37 °C for 15-20 minutes. Cells were then washed twice with ice cold PBS

supplemented with 100 ug/mL cycloheximide and 1 mM PMSF, scraped off the plate, and pelleted

at 200 x g for 5 minutes at 4 °C. Cells were washed once with ice cold PBS, pelleted, and

resuspended in 480 uL 1X low salt buffer supplemented with 100 ug/mL cycloheximide, 2 uM

DTT, 1 mM PMSF, and 1 mM HALT protease inhibitor cocktail. 60 units of RNaseOUT

Recombinant RNase Inhibitor (Invitrogen, 10777019) was added to each sample and vortexed

for 5 seconds. Triton X-100 and sodium deoxycholate was added to each sample to a final

concentration of 0.5 % and samples were vortexed for an additional 5 seconds. Samples were

kept on ice for 5 minutes before centrifugation at 16,000 x g for 7 minutes at 4 °C. Samples were

quantified at 260 nm in UV compatible plates on a multiwell plate reader. 20 units of each sample

was loaded onto the continuous sucrose gradient and spun at 24,200 rpm for 3 hours 30 minutes

using an SW-41 swing bucket rotor in a Beckman Coulter ultracentrifuge. After the spin, the

samples were run on a continuous UA-6 detector, density gradient fractionator system with peak

detection software. Values determined by the fraction collector were graphed on GraphPad Prism.

TMT-based proteomics

Sample preparation, digestion, and tandem-mass tag (TMT) labeling

The TMT-based proteomics was performed exactly as previously described[85]. Briefly, 100 μg

protein of each sample was obtained by cell lysis in lysis buffer (8 M Urea, 200 mM N-(2-

Hydroxyethyl)piperazine-N′-(3-propanesulfonic acid) (EPPS) pH 8.5) followed by reduction using

5 mM tris(2-carboxyethyl)phosphine (TCEP), alkylation with 14 mM iodoacetamide and quenched

using 5 mM dithiothreitol. The reduced and alkylated protein was precipitated using methanol and

chloroform. The protein mixture was digested with LysC (Wako) overnight followed by Trypsin

(Pierce) digestion for 6 hours at 37°C. The trypsin was inactivated with 30% (v/v) acetonitrile. The

digested peptides were labelled with 0.2 mg per reaction of 4-plex TMT reagents (ThermoFisher

scientific) (126, 127N, 129N, and 129C) at room temperature for 1 hour. The reaction was

quenched using 0.5% (v/v) Hydroxylamine for 15 min. A 2.5 μL aliquot from the labeling reaction

was tested for labeling efficiency. TMT-labeled peptides from each sample were pooled together

at a 1:1 ratio. The pooled peptide mix was dried under vacuum and resuspended in 5% formic

acid for 15 min. The resuspended peptide sample was further purified using C18 solid-phase

extraction (SPE) (Sep-Pak, Waters).

Off-line basic pH reverse-phase (BPRP) fractionation.

We fractionated the pooled, labeled peptide sample using BPRP HPLC[86]. We used an Agilent

1200 pump equipped with a degasser and a detector (set at 220 and 280 nm wavelength).

Peptides were subjected to a 60-min linear gradient from 5% to 35% acetonitrile in 10 mM

ammonium bicarbonate pH 8 at a flow rate of 0.25 mL/min over an Agilent 300Extend C18 column

(3.5 μm particles, 2.1 mm ID and 250 mm in length). The peptide mixture was fractionated into a

total of 96 fractions, which were consolidated into 24 super-fractions[87]. Samples were

subsequently acidified with 1% formic acid and vacuum centrifuged to near dryness. Each

consolidated fraction was desalted via StageTip, dried again via vacuum centrifugation, and

reconstituted in 5% acetonitrile, 5% formic acid for LC-MS/MS processing.

Liquid chromatography and tandem mass spectrometry.

Mass spectrometric data were collected on an Orbitrap Fusion mass spectrometer coupled to a

Proxeon NanoLC-1000 UHPLC. The 100 μm capillary column was packed with 35 cm of Accucore

150 resin (2.6 μm, 150Å; ThermoFisher Scientific). The scan sequence began with an MS1

spectrum (Orbitrap analysis, resolution 120,000, 350−1400 Th, automatic gain control (AGC)

target 5 x10^5, maximum injection time 100 ms). Data were acquired for 150 minutes per fraction.

SPS-MS3 analysis was used to reduce ion interference[88,89]. MS2 analysis consisted of collision-

induced dissociation (CID), quadrupole ion trap analysis, automatic gain control (AGC) 1 x10^4,

NCE (normalized collision energy) 35, q-value 0.25, maximum injection time 60 ms), isolation

window at 0.7 Th. Following acquisition of each MS2 spectrum, we collected an MS3 spectrum in

which multiple MS2 fragment ions were captured in the MS3 precursor population using isolation

waveforms with multiple frequency notches. MS3 precursors were fragmented by HCD and

analyzed using the Orbitrap (NCE 65, AGC 2.0 x10^5, isolation window 1.3 Th, maximum injection

time 150 ms, resolution was 50,000).

Data analysis.

Spectra were converted to mzXML via MSconvert[90]. Database searching included all entries from

the Human UniProt Database (downloaded: April 2016). The database was concatenated with

one composed of all protein sequences for that database in the reversed order. Searches were

performed using a 50-ppm precursor ion tolerance for total protein level profiling. The product ion

tolerance was set to 0.9 Da. These wide mass tolerance windows were chosen to maximize

sensitivity in conjunction with Comet searches and linear discriminant analysis[91,92]. TMT tags on

lysine residues and peptide N-termini (+229.163 Da for TMT) and carbamidomethylation of

cysteine residues (+57.021 Da) were set as static modifications, while oxidation of methionine

residues (+15.995 Da) was set as a variable modification. Peptide-spectrum matches (PSMs)

were adjusted to a 1% false discovery rate (FDR)[93,94]. PSM filtering was performed using a linear

discriminant analysis, as described previously[92] and then assembled further to a final protein-

level FDR of 1% [94]. Proteins were quantified by summing reporter ion counts across all matching

PSMs, also as described previously[95]. Reporter ion intensities were adjusted to correct for the

isotopic impurities of the different TMT reagents according to manufacturer specifications. The

signal-to-noise (S/N) measurements of peptides assigned to each protein were summed and

these values were normalized so that the sum of the signal for all proteins in each channel was

equivalent to account for equal protein loading. Finally, each protein abundance measurement

was scaled, such that the summed signal-to-noise for that protein across all channels equaled

100, thereby generating a relative abundance (RA) measurement.

Downstream data analyses for TMT datasets were carried out using the R statistical package

(v4.0.3) and Bioconductor (v3.12; BiocManager 1.30.10). TMT channel intensities were quantile

normalized and then the data were log-transformed. The log transformed data were analyzed with

limma-based R package where p-values were FDR adjusted using an empirical Bayesian

statistical. Differentially expressed proteins were determined using a $\log_2$ (fold change (WT *vs*

UBXN1 KO)) threshold of > +/- 0.65.

Gene ontology (GO) functional enrichment analyses of proteomics data.

The differentially expressed proteins were further annotated and GO functional enrichment

analysis was performed using Metascape online tool (http://metascape.org)[96]. The GO cluster

network and protein-protein interaction network generated by metascape and the STRING

database (https://string-db.org/), respectively, were imported into Cytoscape software (v3.8.2) to

add required attributes (fold changes, p-values, gene number, and conditions) and prepared for

the visualization. Other proteomic data visualizations were performed using the RStudio software

(v1.4.1103), including hrbrthemes (v0.8.0), viridis (v0.6.1), dplyr (v.1.0.7), and ggplot2 (v 3.3.5).

Quantification and statistical analysis

Immunoblots were quantified using densitometry on FIJI and protein levels were normalized to
the loading control for that experiment (GAPDH or β-actin). Fold changes, SEM, and statistical
significance was calculated using GraphPad Prism (version 9.4). Statistical tests performed are
indicated in figure legends and include unpaired two-tailed Student's *t*-test or one-way ANOVA
with post-hoc analysis indicated in figure legends.

Acknowledgements
We thank Peter Juo and Michelle Johnson and Rakesh Ganji for critical reading of the
manuscript. We are grateful to Peter Walter (University of California San Francisco and Altos
Labs) for the GFP-IRE1 construct, and Mals Mariappan (Yale University) for the FLAG-AGAL
construct. This work is supported by the NIH grants GM127557, NS123631 and a Research
Scholar Grant RSG-19-022-01 from the American Cancer Society to M.R. This work was funded
in part by NIH/NIGMS grant GM67945 (S.P.G.) and R01 GM132129 (J.A.P.).

Respective Contributions
B.A and M.R conceived the studies. R.G performed TMT proteomics and data analysis, S.M
performed aggresome studies. J.P and S.P.G assisted with proteomic studies. B.A and M.R wrote
the manuscript.

Competing Interests
The authors declare no conflicts of interest.

Request for reagents
Please contact the corresponding author, M.R for reagent requests.

Data availability

All raw proteomic data will be made available on public servers upon acceptance of the manuscript. Any other data is available from the corresponding author upon request.

<u>REFERENCES</u>

Kaufman, R. J. Orchestrating the unfolded protein response in health and disease. *J Clin Invest* **110**, 1389-1398, doi:10.1172/JCI16886 (2002).

Bellucci, A. *et al.* Induction of the unfolded protein response by alpha-synuclein in experimental models of Parkinson's disease. *J Neurochem* **116**, 588-605, doi:10.1111/j.1471-4159.2010.07143.x (2011).

Duennwald, M. L. & Lindquist, S. Impaired ERAD and ER stress are early and specific events in polyglutamine toxicity. *Genes Dev* **22**, 3308-3319, doi:10.1101/gad.1673408 (2008).

Atkin, J. D. *et al.* Endoplasmic reticulum stress and induction of the unfolded protein response in human sporadic amyotrophic lateral sclerosis. *Neurobiol Dis* **30**, 400-407, doi:10.1016/j.nbd.2008.02.009 (2008).

Hoozemans, J. J. *et al.* The unfolded protein response is activated in pretangle neurons in Alzheimer's disease hippocampus. *Am J Pathol* **174**, 1241-1251, doi:10.2353/ajpath.2009.080814 (2009).

Nishitoh, H. *et al.* ALS-linked mutant SOD1 induces ER stress- and ASK1-dependent motor neuron death by targeting Derlin-1. *Genes Dev* **22**, 1451-1464, doi:10.1101/gad.1640108 (2008).

Cooper, A. A. *et al.* Alpha-synuclein blocks ER-Golgi traffic and Rab1 rescues neuron loss in Parkinson's models. *Science* **313**, 324-328, doi:10.1126/science.1129462 (2006).

Yang, H. *et al.* Huntingtin interacts with the cue domain of gp78 and inhibits gp78 binding to ubiquitin and p97/VCP. *PLoS One* **5**, e8905, doi:10.1371/journal.pone.0008905 (2010).

9   Vonk, W. I. M. *et al.* Differentiation Drives Widespread Rewiring of the Neural Stem Cell

Chaperone Network. *Mol Cell* **78**, 329-345 e329, doi:10.1016/j.molcel.2020.03.009

(2020).

10   Brehme, M. *et al.* A chaperome subnetwork safeguards proteostasis in aging and

neurodegenerative disease. *Cell Rep* **9**, 1135-1150, doi:10.1016/j.celrep.2014.09.042

(2014).

11   Rousseau, A. & Bertolotti, A. An evolutionarily conserved pathway controls proteasome

homeostasis. *Nature* **536**, 184-189, doi:10.1038/nature18943 (2016).

12   Shoulders, M. D. *et al.* Stress-independent activation of XBP1s and/or ATF6 reveals three

functionally diverse ER proteostasis environments. *Cell Rep* **3**, 1279-1292,

doi:10.1016/j.celrep.2013.03.024 (2013).

13   Bertolotti, A., Zhang, Y., Hendershot, L. M., Harding, H. P. & Ron, D. Dynamic interaction

of BiP and ER stress transducers in the unfolded-protein response. *Nat Cell Biol* **2**, 326-

332, doi:10.1038/35014014 (2000).

14   Kopp, M. C., Larburu, N., Durairaj, V., Adams, C. J. & Ali, M. M. U. UPR proteins IRE1

and PERK switch BiP from chaperone to ER stress sensor. *Nat Struct Mol Biol* **26**, 1053-

1062, doi:10.1038/s41594-019-0324-9 (2019).

15   Pincus, D. *et al.* BiP binding to the ER-stress sensor Ire1 tunes the homeostatic behavior

of the unfolded protein response. *PLoS Biol* **8**, e1000415,

doi:10.1371/journal.pbio.1000415 (2010).

16   Harding, H. P., Zhang, Y., Bertolotti, A., Zeng, H. & Ron, D. Perk is essential for

translational regulation and cell survival during the unfolded protein response. *Mol Cell* **5**,

897-904, doi:10.1016/s1097-2765(00)80330-5 (2000).

17   Calfon, M. *et al.* IRE1 couples endoplasmic reticulum load to secretory capacity by

processing the XBP-1 mRNA. *Nature* **415**, 92-96, doi:10.1038/415092a (2002).

18   Yoshida, H., Matsui, T., Yamamoto, A., Okada, T. & Mori, K. XBP1 mRNA is induced by

968   ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription

969   factor. *Cell* **107**, 881-891, doi:10.1016/s0092-8674(01)00611-0 (2001).

970   19   Hollien, J. *et al.* Regulated Ire1-dependent decay of messenger RNAs in mammalian cells.

971   *J Cell Biol* **186**, 323-331, doi:10.1083/jcb.200903014 (2009).

972   20   Lipson, K. L., Ghosh, R. & Urano, F. The role of IRE1alpha in the degradation of insulin

973   mRNA in pancreatic beta-cells. *PLoS One* **3**, e1648, doi:10.1371/journal.pone.0001648

974   (2008).

975   21   Oikawa, D., Tokuda, M. & Iwawaki, T. Site-specific cleavage of CD59 mRNA by

976   endoplasmic reticulum-localized ribonuclease, IRE1. *Biochem Biophys Res Commun* **360**,

977   122-127, doi:10.1016/j.bbrc.2007.06.020 (2007).

978   22   Ghosh, A. P., Klocke, B. J., Ballestas, M. E. & Roth, K. A. CHOP potentially co-operates

979   with FOXO3a in neuronal cells to regulate PUMA and BIM expression in response to ER

980   stress. *PLoS One* **7**, e39586, doi:10.1371/journal.pone.0039586 (2012).

981   23   Urano, F. *et al.* Coupling of stress in the ER to activation of JNK protein kinases by

982   transmembrane protein kinase IRE1. *Science* **287**, 664-666,

983   doi:10.1126/science.287.5453.664 (2000).

984   24   Kato, H. *et al.* mTORC1 serves ER stress-triggered apoptosis via selective activation of

985   the IRE1-JNK pathway. *Cell Death Differ* **19**, 310-320, doi:10.1038/cdd.2011.98 (2012).

986   25   Twomey, E. C. *et al.* Substrate processing by the Cdc48 ATPase complex is initiated by

987   ubiquitin unfolding. *Science* **365**, doi:10.1126/science.aax1033 (2019).

988   26   Pan, M. *et al.* Mechanistic insight into substrate processing and allosteric inhibition of

989   human p97. *Nat Struct Mol Biol* **28**, 614-625, doi:10.1038/s41594-021-00617-2 (2021).

990   27   Bodnar, N. O. & Rapoport, T. A. Molecular Mechanism of Substrate Processing by the

991   Cdc48 ATPase Complex. *Cell* **169**, 722-735 e729, doi:10.1016/j.cell.2017.04.020 (2017).

28   Meyer, H. & Weihl, C. C. The VCP/p97 system at a glance: connecting cellular function to

disease pathogenesis. *J Cell Sci* **127**, 3877-3883, doi:10.1242/jcs.093831 (2014).

29   Ye, Y., Meyer, H. H. & Rapoport, T. A. Function of the p97-Ufd1-Npl4 complex in

retrotranslocation from the ER to the cytosol: dual recognition of nonubiquitinated

polypeptide segments and polyubiquitin chains. *J Cell Biol* **162**, 71-84,

doi:10.1083/jcb.200302169 (2003).

30   Raman, M. *et al.* Systematic proteomics of the VCP-UBXD adaptor network identifies a

role for UBXN10 in regulating ciliogenesis. *Nat Cell Biol* **17**, 1356-1369,

doi:10.1038/ncb3238 (2015).

31   Buchberger, A., Schindelin, H. & Hanzelmann, P. Control of p97 function by cofactor

binding. *FEBS Lett* **589**, 2578-2589, doi:10.1016/j.febslet.2015.08.028 (2015).

32   Zhang, X. *et al.* Altered cofactor regulation with disease-associated p97/VCP mutations.

Proc Natl Acad Sci U S A **112**, E1705-1714, doi:10.1073/pnas.1418820112 (2015).

33   Weihl, C. C., Dalal, S., Pestronk, A. & Hanson, P. I. Inclusion body myopathy-associated

mutations in p97/VCP impair endoplasmic reticulum-associated degradation. *Hum Mol*

Genet **15**, 189-199, doi:10.1093/hmg/ddi426 (2006).

34   Watts, G. D. *et al.* Inclusion body myopathy associated with Paget disease of bone and

frontotemporal dementia is caused by mutant valosin-containing protein. *Nat Genet* **36**,

377-381, doi:10.1038/ng1332 (2004).

35   Schroder, R. *et al.* Mutant valosin-containing protein causes a novel type of frontotemporal

dementia. *Ann Neurol* **57**, 457-461, doi:10.1002/ana.20407 (2005).

36   Tang, W. K. & Xia, D. Mutations in the Human AAA(+) Chaperone p97 and Related

Diseases. *Front Mol Biosci* **3**, 79, doi:10.3389/fmolb.2016.00079 (2016).

37   Gonzalez, M. A. *et al.* A novel mutation in VCP causes Charcot-Marie-Tooth Type 2

disease. *Brain* **137**, 2897-2902, doi:10.1093/brain/awu224 (2014).

38   Gonzalez-Perez, P. *et al.* Novel mutation in VCP gene causes atypical amyotrophic lateral

sclerosis. *Neurology* **79**, 2201-2208, doi:10.1212/WNL.0b013e318275963b (2012).

39   Chan, N. *et al.* Valosin-containing protein mutation and Parkinson's disease. *Parkinsonism*

Relat Disord **18**, 107-109, doi:10.1016/j.parkreldis.2011.07.006 (2012).

40   Ganji, R., Mukkavalli, S., Somanji, F. & Raman, M. The VCP-UBXN1 Complex Mediates

Triage of Ubiquitylated Cytosolic Proteins Bound to the BAG6 Complex. *Mol Cell Biol* **38**,

doi:10.1128/MCB.00154-18 (2018).

41   Hegde, R. S. & Keenan, R. J. Tail-anchored membrane protein insertion into the

endoplasmic reticulum. *Nat Rev Mol Cell Biol* **12**, 787-798, doi:10.1038/nrm3226 (2011).

42   Hegde, R. S. & Zavodszky, E. Recognition and Degradation of Mislocalized Proteins in

Health and Disease. *Cold Spring Harb Perspect Biol* **11**,

doi:10.1101/cshperspect.a033902 (2019).

43   Guna, A. & Hegde, R. S. Transmembrane Domain Recognition during Membrane Protein

Biogenesis and Quality Control. *Curr Biol* **28**, R498-R511, doi:10.1016/j.cub.2018.02.004

(2018).

44   Rodrigo-Brenni, M. C., Gutierrez, E. & Hegde, R. S. Cytosolic quality control of

mislocalized proteins requires RNF126 recruitment to Bag6. *Mol Cell* **55**, 227-237,

doi:10.1016/j.molcel.2014.05.025 (2014).

45   Mariappan, M. *et al.* A ribosome-associating factor chaperones tail-anchored membrane

proteins. *Nature* **466**, 1120-1124, doi:10.1038/nature09296 (2010).

46   Wang, Q. *et al.* A ubiquitin ligase-associated chaperone holdase maintains polypeptides

in soluble states for proteasome degradation. *Mol Cell* **42**, 758-770,

doi:10.1016/j.molcel.2011.05.010 (2011).

47   Ernst, R. *et al.* Enzymatic blockade of the ubiquitin-proteasome pathway. *PLoS Biol* **8**,

e1000605, doi:10.1371/journal.pbio.1000605 (2011).

48   Claessen, J. H. & Ploegh, H. L. BAT3 guides misfolded glycoproteins out of the

endoplasmic reticulum. *PLoS One* **6**, e28542, doi:10.1371/journal.pone.0028542 (2011).

49   Hong, M., Li, M., Mao, C. & Lee, A. S. Endoplasmic reticulum stress triggers an acute

proteasome-dependent degradation of ATF6. *J Cell Biochem* **92**, 723-732,

doi:10.1002/jcb.20118 (2004).

50   Belyy, V., Tran, N. H. & Walter, P. Quantitative microscopy reveals dynamics and fate of

clustered IRE1alpha. *Proc Natl Acad Sci U S A* **117**, 1533-1542,

doi:10.1073/pnas.1915311117 (2020).

51   Li, H., Korennykh, A. V., Behrman, S. L. & Walter, P. Mammalian endoplasmic reticulum

stress sensor IRE1 signals by dynamic clustering. *Proc Natl Acad Sci U S A* **107**, 16113-

16118, doi:10.1073/pnas.1010580107 (2010).

52   van Schadewijk, A., van't Wout, E. F., Stolk, J. & Hiemstra, P. S. A quantitative method

for detection of spliced X-box binding protein-1 (XBP1) mRNA as a measure of

endoplasmic reticulum (ER) stress. *Cell Stress Chaperones* **17**, 275-279,

doi:10.1007/s12192-011-0306-2 (2012).

53   LaLonde, D. P. & Bretscher, A. The UBX protein SAKS1 negatively regulates endoplasmic

reticulum-associated degradation and p97-dependent degradation. *J Biol Chem* **286**,

4892-4901, doi:10.1074/jbc.M110.158030 (2011).

54   Mukkavalli, S., Klickstein, J. A., Ortiz, B., Juo, P. & Raman, M. The p97-UBXN1 complex

regulates aggresome formation. *J Cell Sci* **134**, doi:10.1242/jcs.254201 (2021).

55   Okada, T., Yoshida, H., Akazawa, R., Negishi, M. & Mori, K. Distinct roles of activating

transcription factor 6 (ATF6) and double-stranded RNA-activated protein kinase-like

endoplasmic reticulum kinase (PERK) in transcription during the mammalian unfolded

protein response. *Biochem J* **366**, 585-594, doi:10.1042/BJ20020391 (2002).

56   Gonen, N., Sabath, N., Burge, C. B. & Shalgi, R. Widespread PERK-dependent repression
of ER targets in response to ER stress. *Sci Rep* **9**, 4330, doi:10.1038/s41598-019-38705-
5 (2019).

57   Shinjo, S., Mizotani, Y., Tashiro, E. & Imoto, M. Comparative analysis of the expression
patterns of UPR-target genes caused by UPR-inducing compounds. *Biosci Biotechnol*
Biochem **77**, 729-735, doi:10.1271/bbb.120812 (2013).

58   Travers, K. J. *et al.* Functional and genomic analyses reveal an essential coordination
between the unfolded protein response and ER-associated degradation. *Cell* **101**, 249-
258, doi:10.1016/s0092-8674(00)80835-1 (2000).

59   Harding, H. P. *et al.* An integrated stress response regulates amino acid metabolism and
resistance to oxidative stress. *Mol Cell* **11**, 619-633, doi:10.1016/s1097-2765(03)00105-9
(2003).

60   Han, J. *et al.* ER-stress-induced transcriptional regulation increases protein synthesis
leading to cell death. *Nat Cell Biol* **15**, 481-490, doi:10.1038/ncb2738 (2013).

61   Guan, B. J. *et al.* Translational control during endoplasmic reticulum stress beyond
phosphorylation of the translation initiation factor eIF2alpha. *J Biol Chem* **289**, 12593-
12611, doi:10.1074/jbc.M113.543215 (2014).

62   Krokowski, D. *et al.* A self-defeating anabolic program leads to beta-cell apoptosis in
endoplasmic reticulum stress-induced diabetes via regulation of amino acid flux. *J Biol*
Chem **288**, 17202-17213, doi:10.1074/jbc.M113.466920 (2013).

63   Ye, Y. *et al.* Recruitment of the p97 ATPase and ubiquitin ligases to the site of
retrotranslocation at the endoplasmic reticulum membrane. *Proc Natl Acad Sci U S A* **102**,
14132-14138, doi:10.1073/pnas.0505006102 (2005).

64   Ye, Y., Shibata, Y., Yun, C., Ron, D. & Rapoport, T. A. A membrane protein complex
mediates retro-translocation from the ER lumen into the cytosol. *Nature* **429**, 841-847,
doi:10.1038/nature02656 (2004).

65   Wang, L. *et al.* TMUB1 is an endoplasmic reticulum-resident escortase that promotes the

p97-mediated extraction of membrane proteins for degradation. *Mol Cell* **82**, 3453-3467

e3414, doi:10.1016/j.molcel.2022.07.006 (2022).

66   Sun, S., Li, X. & Mariappan, M. Signal sequences encode information for protein folding

in the endoplasmic reticulum. *J Cell Biol* **222**, doi:10.1083/jcb.202203070 (2023).

67   Kang, S. W. *et al.* Substrate-specific translocational attenuation during ER stress defines

a pre-emptive quality control pathway. *Cell* **127**, 999-1013, doi:10.1016/j.cell.2006.10.032

(2006).

68   Hori, O. *et al.* Transmission of cell stress from endoplasmic reticulum to mitochondria:

enhanced   expression   of   Lon   protease.   *J   Cell   Biol*   **157**,   1151-1160,

doi:10.1083/jcb.200108103 (2002).

69   Bravo, R. *et al.* Increased ER-mitochondrial coupling promotes mitochondrial respiration

and bioenergetics during early phases of ER stress. *J Cell Sci* **124**, 2143-2152,

doi:10.1242/jcs.080762 (2011).

70   Rainbolt, T. K., Atanassova, N., Genereux, J. C. & Wiseman, R. L. Stress-regulated

translational attenuation adapts mitochondrial protein import through Tim17A degradation.

Cell Metab **18**, 908-919, doi:10.1016/j.cmet.2013.11.006 (2013).

71   Schmidt, E. K., Clavarino, G., Ceppi, M. & Pierre, P. SUnSET, a nonradioactive method

to monitor protein synthesis. *Nat Methods* **6**, 275-277, doi:10.1038/nmeth.1314 (2009).

72   Yarmolinsky, M. B. & Haba, G. L. Inhibition by Puromycin of Amino Acid Incorporation into

Protein. *Proc Natl Acad Sci U S A* **45**, 1721-1729, doi:10.1073/pnas.45.12.1721 (1959).

73   Nathans, D. Puromycin Inhibition of Protein Synthesis: Incorporation of Puromycin into

Peptide Chains. *Proc Natl Acad Sci U S A* **51**, 585-592, doi:10.1073/pnas.51.4.585 (1964).

74   Ravi, V., Jain, A., Mishra, S. & Sundaresan, N. R. Measuring Protein Synthesis in Cultured

Cells and Mouse Tissues Using the Non-radioactive SUnSET Assay. *Curr Protoc Mol Biol*

133, e127, doi:10.1002/cpmb.127 (2020).

75   Kim, J. Y. *et al.* ER Stress Drives Lipogenesis and Steatohepatitis via Caspase-2

Activation of S1P. *Cell* **175**, 133-145 e115, doi:10.1016/j.cell.2018.08.020 (2018).

76   Okada, K. *et al.* Prolonged endoplasmic reticulum stress in hypertrophic and failing heart

after aortic constriction: possible contribution of endoplasmic reticulum stress to cardiac

myocyte apoptosis. *Circulation* **110**, 705-712, doi:10.1161/01.CIR.0000137836.95625.D4

(2004).

77   Ljubkovic, M. *et al.* Disturbed Fatty Acid Oxidation, Endoplasmic Reticulum Stress, and

Apoptosis in Left Ventricle of Patients With Type 2 Diabetes. *Diabetes* **68**, 1924-1933,

doi:10.2337/db19-0423 (2019).

78   Ramalingam, L., Sopontammarak, B., Menikdiwela, K. R. & Moustaid-Moussa, N.

Endoplasmic Reticulum (ER) Stress in Part Mediates Effects of Angiotensin II in

Pancreatic Beta Cells. *Diabetes Metab Syndr Obes* **13**, 2843-2853,

doi:10.2147/DMSO.S257797 (2020).

79   Kawai, T. *et al.* IPS-1, an adaptor triggering RIG-I- and Mda5-mediated type I interferon

induction. *Nat Immunol* **6**, 981-988, doi:10.1038/ni1243 (2005).

80   Wang, P. *et al.* UBXN1 interferes with Rig-I-like receptor-mediated antiviral immune

response by targeting MAVS. *Cell Rep* **3**, 1057-1070, doi:10.1016/j.celrep.2013.02.027

(2013).

81   Hu, Y. *et al.* Multiple UBXN family members inhibit retrovirus and lentivirus production and

canonical NFkappaBeta signaling by stabilizing IkappaBalpha. *PLoS Pathog* **13**,

e1006187, doi:10.1371/journal.ppat.1006187 (2017).

82   Walczak, C. P. *et al.* Ribosomal protein RPL26 is the principal target of UFMylation. *Proc*

Natl Acad Sci U S A **116**, 1299-1308, doi:10.1073/pnas.1816202116 (2019).

83   Wang, L. *et al.* UFMylation of RPL26 links translocation-associated quality control to

endoplasmic reticulum protein homeostasis. *Cell Res* **30**, 5-20, doi:10.1038/s41422-019-

0236-6 (2020).

84   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in

multidimensional   genomic   data.   *Bioinformatics*   **32**,   2847-2849,

doi:10.1093/bioinformatics/btw313 (2016).

85   Ganji, R. *et al.* The p97-UBXD8 complex modulates ER-Mitochondria contact sites by

modulating membrane lipid saturation and composition. *bioRxiv*, 2021.2012.2008.471763,

doi:10.1101/2021.12.08.471763 (2021).

86   Wang, Y. *et al.* Reversed-phase chromatography with multiple fraction concatenation

strategy for proteome profiling of human MCF10A cells. *Proteomics* **11**, 2019-2026,

doi:10.1002/pmic.201000722 (2011).

87   Paulo, J. A. *et al.* Quantitative mass spectrometry-based multiplexing compares the

abundance of 5000 S. cerevisiae proteins across 10 carbon sources. *J Proteomics* **148**,

85-93, doi:10.1016/j.jprot.2016.07.005 (2016).

88   Gygi, J. P. *et al.* Web-Based Search Tool for Visualizing Instrument Performance Using

the Triple Knockout (TKO) Proteome Standard. *J Proteome Res* **18**, 687-693,

doi:10.1021/acs.jproteome.8b00737 (2019).

89   Paulo, J. A., O'Connell, J. D. & Gygi, S. P. A Triple Knockout (TKO) Proteomics Standard

for Diagnosing Ion Interference in Isobaric Labeling Experiments. *J Am Soc Mass*

Spectrom **27**, 1620-1625, doi:10.1007/s13361-016-1434-9 (2016).

90   Chambers, M. C. *et al.* A cross-platform toolkit for mass spectrometry and proteomics. *Nat*

Biotechnol **30**, 918-920, doi:10.1038/nbt.2377 (2012).

91   Beausoleil, S. A., Villen, J., Gerber, S. A., Rush, J. & Gygi, S. P. A probability-based

approach for high-throughput protein phosphorylation analysis and site localization.

Nature biotechnology **24**, 1285-1292, doi:10.1038/nbt1240 (2006).

92   Huttlin, E. L. *et al.* A tissue-specific atlas of mouse protein phosphorylation and

expression. *Cell* **143**, 1174-1189, doi:10.1016/j.cell.2010.12.001 (2010).

93   Elias, J. E. & Gygi, S. P. Target-decoy search strategy for mass spectrometry-based

1170   proteomics. *Methods Mol Biol* **604**, 55-71, doi:10.1007/978-1-60761-444-9_5 (2010).

1171   94   Elias, J. E. & Gygi, S. P. Target-decoy search strategy for increased confidence in large-

1172   scale protein identifications by mass spectrometry. *Nat Methods* **4**, 207-214,

1173   doi:10.1038/nmeth1019 (2007).

1174   95   McAlister, G. C. *et al.* Increasing the Multiplexing Capacity of TMTs Using Reporter Ion

1175   Isotopologues with Isobaric Masses. *Analytical Chemistry* **84**, 7469-7478,

1176   doi:10.1021/ac301572t (2012).

1177   96   Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of

1178   systems-level datasets. *Nat Commun* **10**, 1523, doi:10.1038/s41467-019-09234-6 (2019).

1179

Figure 1

Figure 2

Figure 3

A. Tg (hrs)

Fix and stain

Release

0 0.5 1.5 3

Wildtype
Tg (hours)
0   0.5

UBXN1 KO
Tg (hours)
0   0.5

1.5   3
Tg (hours)

1.5   3
Tg (hours)

B.

Colony outgrowth

% Viability

**

*

****

WT
UBXN1 KO

0   1   2   3
Tg (hours)

C.

Wildtype   UBXN1 KO
0 2 4 8   0 2 4 8   DTT (hours)

CHOP ─   ─25 kDa
UBXN1 ─
                                    ─37 kDa
GAPDH ─   ─37 kDa

1 2 3 4   5 6 7 8

D.

CHOP 8 hours

Fold Change

*

WT UBXN1 KO

E.

ER-microsomes
PNS Cyto -   +   -   +   Proteinase K
           -   -   +   +   TX-100

UBXN1 ─   ─37 kDa
p97 ─   ─100 kDa
Calnexin full length ─   ─100 kDa
Calnexin ER lumen ─   ─50 kDa
Sec61β ─   ─15 kDa
β-actin ─   ─50 kDa
TOMM20 ─   ─20 kDa

1   2   3   4   5   6

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary Figure 1

A.

	Input		IP:UBXN1
	10µg	20µg	IgG

UBXN1 — 37 kDa

Sec61β — 15 kDa

p97 — 100 kDa

B.

UBXN1
Sec61α — 37 kDa

Sec61β — 15 kDa

Supplementary Figure 3

A.

B.

Supplementary Figure 4

Supplementary Figure 5

A.

B.

C.

D.