1
A phylogenetics-based nomenclature system for steroid receptors in teleost fishes

9
10
11
12

Andrew P. Hoadley[1†], Kathleen M. Munley[1†], and Beau A. Alward[1,2*]

14
15
16

[1]*Department of Psychology, University of Houston, Houston, TX, USA, and [2]Department of Biology and Biochemistry, University of Houston, Houston, TX, USA*

19
20

[†]These authors share first authorship

*Corresponding Author:
Beau A. Alward
Department of Psychology and Department of Biology and Biochemistry
University of Houston
4349 Martin Luther King Boulevard
Houston, TX 77204
Telephone: 713-743-4138
E-mail: balward@uh.edu

ABSTRACT (250/250 words)

Teleost fishes have emerged as tractable models for studying the neuroendocrine

regulation of social behavior via molecular genetic techniques, such as CRISPR/Cas9

gene editing. Moreover, teleosts provide an opportunity to investigate the evolution of

steroid receptors and their functions, as species within this lineage possess novel steroid

receptor paralogs that resulted from a teleost-specific whole genome duplication.

Although teleost fishes have grown in popularity as models for behavioral

neuroendocrinology, there is not a consistent nomenclature system for steroid receptors

and their genes, which may impede a clear understanding of steroid receptor paralogs

and their functions. Here, we used a phylogenetic approach to assess the relatedness of

protein sequences encoding steroid receptor paralogs in 18 species from 12 different

orders of the Infraclass Teleostei. While most similarly named sequences grouped based

on the established phylogeny of the teleost fish lineage, our analysis revealed several

inconsistencies in the nomenclature of steroid receptor paralogs, particularly for

sequences encoding estrogen receptor beta (ERβ) and estrogen-related receptor gamma

(ERγ). Based on our results, we propose a nomenclature system for teleosts in which

Greek symbols refer to proteins and numbers refer to genes encoding different subtypes

of steroid receptors within the five major groups of this nuclear receptor subfamily.

Collectively, our results bridge a critical gap by providing a cohesive naming system for

steroid receptors in teleost fishes, which will serve to improve communication, promote

collaboration, and enhance our understanding of the evolution and function of steroid

receptors across vertebrates.

Keywords: evolution, fishes, genome duplication, paralogs, steroid hormones, steroid

receptors

HIGHLIGHTS (3-5 bullet points, 85 characters or less each)

- Teleosts have emerged as models for studying neuroendocrine regulation of behavior.
- Teleost fishes possess additional steroid receptor paralogs due to the TS-WGD.
- Teleosts lack consistent nomenclature system for steroid receptors and their genes.
- Phylogenetic analyses of protein sequences revealed discrepancies for ERβ and ERγ.
- We propose a novel, cohesive naming system for steroid receptors in teleost fishes.

1. INTRODUCTION

Over the past few decades, teleost fishes have emerged as important model systems in numerous biological fields, including molecular biology, genetics, evolution, neuroendocrinology, and behavior (reviewed in Glasauer and Neuhass, 2014; Sato and Nishida, 2010; Volff, 2005). Teleost fishes (Class: Actinopterygii, Infraclass: Teleostei) are the largest and most diverse group of ray-finned fishes that comprise >95% of all extant fish species and roughly half of all extant vertebrate species (Nelson, 1994; reviewed in Volff, 2005). Consequently, teleosts show a remarkable level of diversity in morphology, ecology, physiology, and behavior, enabling researchers to address evolutionary questions related to various facets of biology. In recent years, the popularity of teleosts in biological research has surged, particularly for their tractability in molecular genetic approaches. The development of high-throughput and rapid DNA sequencing methods has enabled researchers to sequence the whole genomes of several fish species from distinct orders within the Infraclass Teleostei, including channel catfish (*Ictalurus punctatus*; Liu et al., 2016), zebrafish (*Danio rerio*; Howe et al., 2013), Atlantic salmon (*Salmo salar*; Lien et al., 2016), spotted green pufferfish (*Tetraodon nigroviridis*; Jaillon et al., 2004), three-spined stickleback (*Gasterosteus aculeatus*; Nath et al., 2021; Peichel et al. 2017, 2020), Nile tilapia (*Oreochromis niloticus*; Conte et al., 2017), and medaka (*Oryzias latipes*; Kasahara et al., 2007). The availability of whole genome sequences, in addition to the prevalence of external fertilization as a mode of reproduction among many teleost species, makes techniques such as CRISPR/Cas9 relatively straightforward to implement (reviewed in Barman et al., 2017; Lu et al., 2021; Zhu and Ge, 2018). Collectively, these advantages make teleost fishes excellent models for examining the molecular regulation of physiological mechanisms and behavior via gene editing.

Genomic and phylogenetic evidence strongly supports the occurrence of a single whole-genome duplication (WGD) event approximately 350 mya in the common ancestor of all teleosts (Brunet et al., 2006), which took place after the emergence of the non-teleost ray-finned fishes (e.g., gar and bowfin), but before the divergence of the superorder Osteoglossomorpha (Hoegg et al., 2004; Fig. 1). This event is often referred to as the teleost-specific WGD [TS-WGD, also known as 3R WGD or third round WGD;

reviewed in Glasauer and Neuhauss (2014)]. After a WGD event, redundant genes can

face one or more of several fates. Frequently, relaxed selective pressure on a duplicated

gene leads to the buildup of deleterious mutations, which results in non-functionalization

(Ohno, 1970). In some cases, however, one or more gene paralogs may be under positive

selection due to the accumulation of mutations conferring novel functions, a process

known as neo-functionalization (Force et al., 1999; Ohno, 1970). Alternatively, duplicate

genes may undergo sub-functionalization, in which both paralogs are maintained due to

the complementary partitioning of ancestral functions between two daughter genes, such

that their complimentary actions comprise the total function of their single ancestral gene

(Force et al., 1999).

Among the genes that have been duplicated following the TS-WGD are steroid

receptors, a family of ligand-dependent nuclear receptors that function as signal

transducers and transcription factors (Carson-Jurica et al., 1990). Steroid receptors

modulate suites of physiological mechanisms, such as homeostatic functions, growth and

development, cellular signaling, sexual differentiation and reproduction, as well as

behavior (Szego et al. 2003; Whirledge and Cidlowski, 2019). Steroid receptors consist

of five major groups: 1) the androgen receptors (ARs), 2) estrogen receptors (ERs), and

3) progesterone receptors (PRs), which mediate the actions of androgens (e.g.,

testosterone, dihydrotestosterone, and 11-ketotestosterone), estrogens (e.g., estradiol),

and progestins (e.g., progesterone) on the development and maintenance of the

reproductive, musculoskeletal, cardiovascular, immune, and central nervous systems

(reviewed in Brinton et al., 2008; Chen et al., 2022; Davey and Grossmann, 2016; Mani

et al. 1997), and 4) the glucocorticoid receptors (GRs) and 5) mineralocorticoid receptors

(MRs), which facilitate the actions of glucocorticoids (e.g., cortisol) and mineralocorticoids

(e.g., aldosterone) on metabolism, growth and development, cardiovascular function, ion

transport, and salt balance (reviewed in Pascual-Le Tallec and Lombés, 2005;

Timmermans et al., 2019). Prior work suggests that extant steroid receptors evolved from

an ancestral estrogen receptor that had estrogen-like functionality and preceded the

origin of bilaterally symmetric animals, followed by an ancestral progesterone receptor

(Thornton, 2001; Thornton et al., 2003). Specifically, extant steroid receptors evolved

novel functions from these ancestral receptors following two large-scale genome

expansions, one before and one after the advent of jawed vertebrates (the 1R-WGD and
2R-WGD, respectively). Because the ancestral steroid receptor had estrogen-like
functionality, these findings support a model of ligand exploitation for extant steroid
receptors, in which the terminal ligand in a biosynthetic pathway (in this case, estrogens)
is the first for which a receptor evolves, thereby indirectly selecting for the synthesis of
intermediates within this pathway and enabling duplicated receptors to evolve affinity for
these substances (Baker, 2004; reviewed in Eick and Thornton, 2011). Taken together,
these studies suggest that ERs and PRs evolved from more ancient lineages, whereas
ARs, GRs, and MRs are relatively recent in origin.

The novel steroid receptor paralogs resulting from the TS-GWD present both
opportunities and challenges for researchers. For instance, novel steroid receptor
paralogs provide opportunities to dissect their functions to discover fundamental roles of
steroid hormones across species (reviewed in Alward et al., 2023). Some challenges
include difficulty in manipulating the functions of these genes (e.g., two steroid receptors
bind the same steroid ligand in many teleosts, whereas in other vertebrates, only one
gene product does). Another practical challenge is there is not a cohesive nomenclature
system for the steroid receptors. This challenge at first may seem inconsequential, but a
lack of a logical nomenclature system for novel teleost steroid receptor genes may hinder
a clear understanding of their functions. With the advent of CRISPR/Cas9 gene editing
technologies, numerous non-traditional laboratory species, including teleost fishes, are
emerging as genetically tractable for a variety of research questions (reviewed in Alward
et al., 2023; Juntti, 2019), including those related to steroid hormone function (Alward et
al., 2023, 2020; Crowder et al., 2018; Faught and Vijayan, 2019, 2018; Muto et al., 2013;
Yin et al., 2017; Yong et al., 2017; Ziv et al., 2013). Therefore, it is important for logical,
consistent nomenclature to be used for steroid receptors and their genes across teleost
fishes.

The goal of the current study was to use a quantitative approach to propose a
nomenclature system for steroid receptors in teleosts. To achieve this, we used
phylogenetics to determine the relatedness of protein sequences encoding teleost steroid
receptor paralogs from the five major groups of this nuclear receptor subfamily in
vertebrates: ARs, ERs, PRs, GRs, and MRs. This study focused solely on steroid

receptors, as the nomenclature issue across teleosts is especially apparent for this class
of molecules. We predicted that steroid receptor protein sequences would generally group
based on the established phylogeny of the teleost fish lineage, but that there would be a
few instances in which protein sequences of certain steroid receptor paralogs show
greater similarity to a different group of related paralogs than for that which the sequence
is currently named. Together, our findings bridge a critical gap among researchers who
study teleost fishes by providing a consistent nomenclature system for steroid receptors,
clarifying scientific communication among and between fields and promoting the
progression of teleost fish research during rapid technological advancements in genetic
technology. We hope that our approach may act as a model to benefit other researchers
to establish a consistent nomenclature for other species and/or other molecular systems
with an inconsistent nomenclature system.

2. MATERIALS AND METHODS

2.1. Species selection

To assess the relatedness of steroid receptor protein sequences among teleost
fishes, we chose 18 species representing 12 different orders from the Infraclass Teleostei
[Order Cypriniformes (carps, minnows, and loaches), Order Cyprinodontimformes
(toothcarps), Order Characiformes (tetras and piranhas), Order Siluriformes (catfish),
Order Salmoniformes (salmonids), Order Gadiformes (cods), Order Beloniformes
(medakas, ricefishes, and needlefishes), Order Cichliformes (cichlids and convict
blennies), Order Perciformes (perches, darters, basses, and groupers), Order
Gasterosteiformes (sticklebacks, pipefishes, and seahorses), Order Pleuronectiformes
(flatfishes), Order Tetraodontiformes (puffers and filefishes), and family Pomacentridae
in the subseries Ovalentaria (damselfishes and clownfishes)]. Specifically, we selected
the following species for phylogenetic analysis: *Anguilla japonica* (Japanese eel), *Danio*
rerio (zebrafish), *Carassius auratus* (goldfish), *Fundulus heteroclitus* (Atlantic killifish),
Poecilia mexicana (shortfin molly), *Gambusia affinis* (western mosquitofish), *Astyanax*
mexicanus (Mexican tetra), *Ictalurus punctatus* (channel catfish), *Oncorhynchus mykiss*
(rainbow trout), *Gadus morhua* (Atlantic cod), *Oryzias latipes* (medaka), *Astatotilapia*
(*Haplochromis) burtoni* (Burton's mouthbrooder), *Oreochromis niloticus* (Nile tilapia),

Amphiprion percula (orange clownfish), *Dicentrarchus labrax* (European seabass),
Gasterosteus aculeatus (three-spined stickleback), *Paralichthys olivaceus* (olive
flounder), and *Tetraodon nigroviridis* (spotted green pufferfish). These species were
chosen based on the availability of protein sequences for steroid receptors, many of which
were acquired via whole-genome sequencing projects, and to represent the diversity of
fishes within the teleost lineage.

In addition, we included 1 non-teleost ray-finned fish [*Lepisosteus oculatus*
(spotted gar), Infraclass: Holostei], which diverged from teleost fishes before the TS-
WGD; and 2 cartilaginous fishes [*Leucoraja erinacea* (little skate), Order: Rajiformes;
Callorhinchus milli (Australian ghostshark), Order: Chimaeriformes] and 1 jawless fish
[*Petromyzon marinus* (sea lamprey), Order: Petromyzontiformes], which are ancestral to
ray-finned fishes. These species were selected to examine potential differences between
steroid receptor protein sequences from more ancestral fish species (*P. marinus*, *L.
erinacea*, and *C. milli*) and from a species that diverged from teleosts prior to the TS-
WGD (*L. oculatus*) to steroid receptor paralogs within the teleost lineage. Finally, we
chose 4 additional species that represent traditional animal models across biological
fields [*Xenopus laevis* (African clawed frog), *Gallus gallus* (red junglefowl), *Mus musculus*
(house mouse), and *Homo sapiens* (humans)] as outgroups for our phylogenetic
analyses. A phylogeny showing established relationships between the teleost fishes, non-
teleost fishes, and outgroup species included in our phylogenetic analyses are shown in
Figure 1.

2.2. Protein sequence collection and criteria

Protein sequences for androgen, estrogen, progesterone, glucocorticoid, and
mineralocorticoid receptors, including paralogs and isoforms resulting from the TS-WGD,
were acquired from the National Center for Biotechnology Information (NCBI) Protein and
UniProt databases. This search yielded a total of 211 protein sequences across 26
species (36 androgen receptor sequences, 95 estrogen receptor sequences, 27
progesterone receptor sequences, 30 glucocorticoid receptor sequences, and 23
mineralocorticoid receptor sequences). Sequences for the following steroid receptors,
paralogs, and isoforms were included in our analysis: ARα, ARβ, ARa, ARb, ERα, ERβ,

ERβ1, ERβ2, estrogen-related receptor gamma (ERγ), estrogen-related receptor gamma

1 (ERγ1), estrogen-related receptor gamma 2 (ERγ2), PR, PR1, PR2, GR, GR1, GR2,

GR2a, GR2b, and MR. Sequences that were described as alternative splice variants or

as partial sequences were excluded from analysis. Protein sequences were downloaded

from NCBI Protein and UniProt in FASTA format for inclusion in phylogenetic analysis

(see **Section 2.3**). A complete list of protein sequences that were used for phylogenetic

analysis, along with their sources, accession numbers, and references, is available in the

Supplementary Material (Tables S1-S5).

2.3. Phylogenetic analyses

For each major group of steroid receptors (excluding ERγ, which was run

separately from the other ER subtypes because it is highly conserved among teleosts),

available or inferred full-length amino acid sequences were used to generate a molecular

phylogeny. Construction of multiple alignments of receptor protein sequences with the

neighbor-joining method were performed using the MUSCLE Alignment version 3.8.425

(Edgar, 2004; Saitou and Nei, 1987). To estimate molecular phylogenies, trees were

constructed using the Jukes-Cantor genetic distance model (Jukes and Cantor, 1969).

Trees were rooted using the most basal organism for which we had an available sequence

[*P. marinus* (sea lamprey), *L. erinacea* (little skate), *C. milii* (Australian ghostshark), or *H.*

sapiens (human)]. Bootstrap probabilities for the neighbor-joining analyses were

calculated using bootstrap resampling with 1000 replicates (Felsenstein, 1985). All

analyses were performed using Geneious Prime version 2022.2.2.

2.4 Approach for steroid receptor nomenclature

Our approach to determining a nomenclature system for protein and gene

sequences encoding steroid receptors was based on clustering within our phylogenies.

We hypothesized that collections of closely related genes across the species would form

large clusters that generally correspond to the current understanding of the proteins

present in teleosts in which the genes and/or proteins have been identified. For example,

we predicted that there would be two distinct clusters for the ARs because two distinct

paralogs that have been discovered for these genes in several species of teleosts (Lorin

et al., 2015; Ogino et al., 2016).

Protein names were assigned for each cluster based on majority rule, in which the

name proposed for a given cluster reflected the current protein name that was listed for

most teleost species included in our analyses. Once protein names were assigned, we

proposed names for their corresponding genes (see Table 1 for a basic consensus of all

steroid receptor proteins and genes and Supplementary Material, Tables S1-S5 for a full

list of proposed protein and gene names for each individual species). We propose a new

system for naming novel paralogous steroid receptor genes using a combination of the

existing nomenclature system in teleosts as well as the model of steroid receptor gene

nomenclature in mice (*M. musculus*).

3. RESULTS

Our phylogenetic approach led to clear classifications of a variety of steroid

receptor protein sequences across species. The proposed names for the steroid receptor

proteins are shown in each figure for the receptors in distinct, broad clusters delineated

by differently colored, transparent rectangles. The corresponding gene name for each

protein is provided in Table 1. Based on clusters in the phylogenies, we propose names

across defined clusters for each steroid receptor.

3.1. Androgen receptors

Our analysis of AR strongly resolves the split between the teleost specific ARα and

ARβ paralogs (Fig. 2). Overall, protein sequences for AR, ARα, and ARβ clustered into

two well-supported clades (90% bootstrap value) that generally agreed with established

phylogenetic relationships in the teleost lineage (Fig. 1). The first clade primarily

contained ARα sequences ("Androgen Receptor Alpha" group), whereas the second

clade mostly consisted of ARβ sequences ("Androgen Receptor Beta" group; Fig. 2).

These results agree with previous work showing that many species of teleosts possess

two distinct AR paralogs (Lorin et al., 2015; Ogino et al., 2016). Like the current naming

system for AR genes in mice (Juntti et al., 2010), we propose referring to the genes for

an AR protein with an *ar* base*.* Thus, genes encoding ARα will be called *ar1*, and genes
encoding ARβ will be called *ar2* (Table 1).

The AR sequences for the tetrapod species clustered with *L. erinacea* (little skate)
at the base of the AR phylogenetic tree (100% bootstrap value; Fig. 2). Most ARα
sequences clustered into the "Androgen Receptor Alpha" group (100% bootstrap value)*,*
while the majority of ARβ sequences grouped into the "Androgen Receptor Beta" group
(94% bootstrap value). Importantly, we identified several discrepancies in how ARs are
currently named in the NCBI Protein and UniProt databases. Specifically, sequences
labeled "androgen receptor" for *G. morhua* (Atlantic cod) and *A. burtoni* (Burton's
mouthbrooder) clustered with the ARα sequences in the "Androgen Receptor Alpha"
group (≥ 96% bootstrap values), whereas similarly named sequences for *T. nigroviridis*
(spotted green pufferfish), *D. rerio* (zebrafish), *C. auratus* (goldfish), *A. mexicanus*
(Mexican tetra), *D. labrax* (European seabass), *A. percula* (orange clownfish), *F.*
heteroclitus (Atlantic killifish), and *G. affinis* (western mosquitofish) clustered with ARβ
sequences in the "Androgen Receptor Beta" group (>90% bootstrap values).
Interestingly, the ARα and ARβ sequences for *O. mykiss* (rainbow trout) and *A. japonica*
(Japanese eel) clustered in the "Androgen Receptor Beta" group, suggesting that the ARα
sequences in these species are more similar to ARβ sequences in other teleost species
than to other ARα sequences.

3.2. Estrogen receptors

Our phylogenetic analysis for ERs revealed several inconsistencies in the
nomenclature of estrogen receptor paralogs in teleost fishes based on protein sequence
similarities, particularly for ERβ and ERγ (Fig. 3). Protein sequences for ER, ERα, and
ERβ clustered into two well-supported clades (100% bootstrap values). The first clade
largely consisted of ERα sequences ("Estrogen Receptor Alpha" group), whereas the
second clade primarily contained ERβ sequences ("Estrogen Receptor Beta" group).
Furthermore, there were two additional clusters within the "Estrogen Receptor Beta"
group that were supported by our analysis (70% bootstrap value): 1) the "Estrogen
Receptor Beta 2" group, which was mostly comprised of ERβ2 sequences (93% bootstrap
value), and 2) the "Estrogen Receptor Beta 1" group, which mainly included ERβ1

sequences (60% bootstrap value; Fig. 3). These results are consistent with the

expression of two distinct ERβ genes, *esr2a* and *esr2b*, in medaka (*O. latipes*; Nishiike et

al., 2021) and Burton's mouthbrooder (*A. burtoni*; Maruska et al., 2020). We propose

calling the genes for ERα, ERβ1, and ERβ2, and ERγ *esr1, esr2a, esr2b,* and *esr3*,

respectively (Table 1).

The ERα sequence for *P. marinus* (sea lamprey) clustered by itself at the base of

the ERα and ERβ phylogenetic tree (100% bootstrap value; Fig. 3). All other ERα

sequences clustered into a single clade ("Estrogen Receptor Alpha," 100% bootstrap

value), including those of *X. laevis* (African clawed frog; "estrogen receptor alpha 1" and

"estrogen receptor alpha 2") and the two cartilaginous fish species [*L. erinacea* (little

skate) and *C. milli* (Australian ghostshark)], and generally agreed with established

phylogenetic relationships within teleost fishes (Fig. 1). Notably, all sequences labeled

"estrogen receptor" in NCBI Protein and UniProt, including those of *D. rerio* (zebrafish),

C. auratus (goldfish), *A. mexicanus* (Mexican tetra), *O. mykiss* (rainbow trout), *O. latipes*

(medaka), and *A. percula* (orange clownfish), as well as "estrogen receptor 1" for *A.*

japonica (Japanese eel) and the "type I estrogen receptor" for *O. niloticus* (Nile tilapia),

clustered with the ERα sequences in the "Estrogen Receptor Alpha" group (≥ 99%

bootstrap values; Fig. 3).

Conversely, we found some discrepancies in nomenclature based on protein

sequence alignment for the "Estrogen Receptor Beta" clade (Fig. 3). The tetrapod

outgroup sequences for ERβ ["estrogen receptor beta 1" and "estrogen receptor beta 2"

for *X. laevis* (African clawed frog)] clustered with *L. erinacea* (little skate), *C. milli*

(Australian ghostshark), and *L. oculatus* (spotted gar), species that diverged from teleosts

prior to the TS-WGD, and these sequences were basal to all teleost ERβ sequences that

were included in our analysis (100% bootstrap value). The majority of ERβ2 sequences,

which are labeled "estrogen receptor beta 2" and "estrogen receptor 2b," clustered into a

single well-supported clade ("Estrogen Receptor Beta 2," 93% bootstrap value) that was

consistent with known phylogenetic relationships in the teleost lineage (Fig. 1).

Importantly, there was one sequence within the "Estrogen Receptor Beta 2" group that

was unexpected: the "estrogen receptor beta 1" sequence for *D. rerio* (zebrafish, 100%

bootstrap value). The sequences "estrogen receptor type beta" for *I. punctatus* (channel

catfish) and "estrogen receptor BetaB" for *F. heteroclitus* (Atlantic killifish) also clustered

with the ERβ2 sequences in the "Estrogen Receptor β2" group (≥ 96% bootstrap values).

Similarly, most ERβ1 sequences, which are labeled "estrogen receptor beta 1" or

"estrogen receptor 2a," clustered into a single clade (60% bootstrap value) that typically

followed known phylogenetic relationships in teleost fishes (Fig. 1). Moreover, all

sequences labeled "estrogen receptor beta" in NCBI Protein and UniProt, including those

of *C. auratus* (goldfish), *O. mykiss* (rainbow trout), *O. latipes* (medaka), and *P. olivaceus*

(olive flounder), as well as "estrogen receptor BetaA" for *F. heteroclitus* (Atlantic killifish)

and "type II estrogen receptor" for *O. niloticus* (Nile tilapia), clustered with the ERβ

sequences in the "Estrogen Receptor Beta 1" group (≥ 80% bootstrap values; Fig. 3).

Of the ER paralogs examined in this study, ERγ showed the highest level of protein

sequence similarity across teleost species (Fig. 4). Because the ERγ protein sequence is

highly conserved among teleosts relative to the protein sequences of other ER paralogs,

we performed our ERγ phylogenetic analysis separately from that of ERα and ERβ.

Interestingly, "estrogen-related receptor gamma" for *P. marinus* (sea lamprey) and

"estrogen-related receptor beta" for *A. percula* (orange clownfish) clustered in a single

group that was located at the base of the ERγ phylogenetic tree (100% bootstrap values).

Overall, ERγ sequences clustered into two clades, one that primarily consisted of ERγb

sequences ("Estrogen-Related Receptor Gamma B" group; 92% bootstrap value) and

one that was mostly comprised of ERγa sequences ("Estrogen-Related Receptor Gamma

A" group; 98% bootstrap value). Notably, there were more discrepancies between the

relatedness of ERγ protein sequences and established phylogenetic relationships in

teleost fishes compared to the phylogenetic tree generated from ERα and ERβ

sequences (Figs. 1 and 3). The "Estrogen-Related Receptor Gamma B" group included

sequences labeled "estrogen-related receptor gamma b" with a few exceptions:

"estrogen-related receptor gamma type 2" for *O. latipes* (medaka), "estrogen-related

receptor alpha" for *T. nigroviridis* (spotted green pufferfish) and *A. percula* (orange

clownfish), and "estrogen-related receptor gamma-like" for *C. auratus* (goldfish; 100%

bootstrap values). Conversely, the "Estrogen-Related Receptor Gamma A" group

consisted of sequences labeled "estrogen-related receptor gamma a" as well as most

sequences labeled "estrogen-related receptor gamma" in NCBI Protein or UniProt,

including those of *C. milli* (Australian ghostshark), *L. oculatus* (spotted gar), *D. rerio*

(zebrafish), *I. punctatus* (channel catfish), *O. mykiss* (rainbow trout), *F. heteroclitus*

(Atlantic killifish), *G. morhua* (Atlantic cod), *P. mexicana* (shortfin molly), *O. niloticus* (Nile

tilapia), and *A. burtoni* (Burton's mouthbrooder; ≥ 98% bootstrap values). In addition, the

"estrogen-receptor gamma type 1" sequence from *O. latipes* (medaka) clustered with the

ERγa sequences in the "Estrogen-Related Receptor Gamma A" group (76% bootstrap

value; Fig. 4). To our knowledge, it is unknown whether ERγa and Erγb are encoded by

distinct genes in the species in which distinct ERγ proteins have been discovered.

3.3. Progesterone receptors

Our analysis of PR suggests that there was no widespread retention of orthologous

PR genes following the TS-WGD, consistent with previous studies (Ren et al. 2019; Fig.

5). The tetrapod sequences for PR clustered as a sister group to all teleost sequences

(100% bootstrap value), and all teleost PR sequences clustered into a single well-

supported clade (100% bootstrap value) that mostly agreed with established phylogenetic

relationships in this lineage (Fig. 1). Importantly, our search yielded two PR sequences in

C. auratus (goldfish; PR1 and PR2), which clustered next to each other (86% bootstrap

value; Fig. 5). No additional PR sequences have been discovered in closely-related

teleost species (e.g., zebrafish; Tang et al., 2016), suggesting that these sequences might

represent a gene duplication that is specific to the goldfish lineage. The current name for

the PR gene is *pgr*, and this nomenclature was consistent across all teleost species

examined in our study*.*

3.4. Glucocorticoid receptors

Our analysis revealed several inconsistencies in the nomenclature of GR (Fig. 6).

Protein sequences for GR, GR1, and GR2 clustered into two clades (78% bootstrap

value) that were consistent with known phylogenetic relationships in teleosts (Fig. 1). The

first clade mostly consisted of GR2 sequences ("Glucocorticoid Receptor 2" group), while

the second clade primarily contained GR1 sequences ("Glucocorticoid Receptor 1" group;

Fig. 6), suggesting that two glucocorticoid receptors persisted in the teleost lineage

following the TS-WGD. To remain consistent with the naming systems for the other

steriod receptor proteins, we propose calling "glucocorticoid receptor 1" GRα and

"glucocorticoid receptor 2" GRβ with gene names of *nr3c1a* and *nr3c1b* (Table 1).

The tetrapod outgroup sequences for GR clustered with that of *L. oculatus* (spotted

gar) at the base of the GR phylogenetic tree (100% bootstrap value; Fig. 6). Interestingly,

most GR sequences used in our analysis were labeled "glucocorticoid receptor," yet these

sequences still grouped into two separate clades. Specifically, sequences labeled

"glucocorticoid receptor" for *D. rerio* (zebrafish), *T. nigroviridis* (spotted green pufferfish),

F. heteroclitus (Atlantic killifish), *O. latipes* (medaka), *G. aculeatus* (three-spined

stickleback), and *A. burtoni* (Burton's mouthbrooder) clustered in the "Glucocrticoid

Receptor 2" group (100% bootstrap value), whereas sequences labeled "glucocorticoid

receptor" for *A. japonica* (Japanese eel), *A. mexicanus* (Mexican tetra), *O. mykiss*

(rainbow trout), *A. percula* (orange clownfish), *O. niloticus* (Nile tilapia), *D. labrax*

(European seabass), and *P. olivaceus* (olive flounder), in addition to "glucocorticoid

receptor-like" for *G. morhua* (Atlantic cod) and *P. mexicana* (shortfin molly), clustered in

the "Glucocorticoid Receptor 1" group (93% bootstrap value). A few species in our

analysis had two different GR sequences, including *I. punctatus* (channel catfish;

"glucocorticoid receptor 1" and "glucocorticoid receptor 2"), *A. burtoni* [Burton's

mouthbrooder; "glucocorticoid receptor" and "glucocorticoid receptor (2)"] and *O. mykiss*

(rainbow trout; "glucocorticoid receptor" and "glucocorticoid receptor 2"). Although these

sequences generally clustered with other similarly named sequences, the "glucocorticoid

receptor (2)" sequence for *A. burtoni* clustered with sequences in the "Glucocorticoid

Receptor 1" group, suggesting that this sequence should be called GRα. Because

duplicate GR sequences were discovered in highly derived clades (e.g., *A. burtoni*), our

results suggest that GR paralogs may exist, but remain undescribed in some teleost

species.

3.5. Mineralocorticoid receptors

Our analysis of MR indicates no widespread retention of orthologous MR genes

following the TS-WGD (Fig. 7). The tetrapod outgroup sequences for MR clustered with

P. marinus (sea lamprey) and were basal to all teleost MR sequences (100% bootstrap

value). MR protein sequences from teleost fishes clustered into a single well-supported

clade (100% bootstrap value) that was consistent with known phylogenetic relationships

in this lineage (Fig. 1). While one of the sequences included in our analysis was labeled

"mineralocorticoid receptor-like" [from *P. mexicana* (shortfin molly)], this sequence

confidently clustered with other MRs in our analysis (100% bootstrap value), suggesting

that this protein sequence encodes MR. The current name for the MR gene is *nr3c2*, and

this nomenclature was consistent in all teleost species included in our study*.*

4. DISCUSSION

As the most specious clade in the animal kingdom, teleost fishes exhibit profound

variation in life-history traits, especially those related to the actions of steroid hormones,

such as reproductive and physiological plasticity (reviewed in Godwin, 2010). As such,

steroid production and signaling mechanisms have been studied extensively in many

teleost fishes, presenting both opportunities and challenges in making comparative

insights in the field of behavioral neuroendocrinology. Because the genome editing

revolution has enhanced research progress and genetic tractability in diverse non-

traditional species like teleosts (reviewed in Alward et al., 2023; Juntti, 2019), it is

important that basic tools, such as the names of widely-studied proteins and genes, are

consistently and logically applied. To facilitate both ongoing and future collaboration,

researchers in this diverse field must endeavor to work from the same set of facts. As an

initial step in that direction, we used a phylogenetic approach to propose a consistent

naming scheme for steroid receptors and their genes.

In our analysis, we identified several instances of supporting evidence for novel

steroid receptors, including PRs in goldfish, which possess two distinct PRs encoded by

distinct genes, and ERγ, for which a few species had two copies.   Future work

investigating whether other species possess extra PR genes and whether the different

ERγ proteins are encoded by distinct genes is warranted. Several intriguing observations

were also made for the ARs. For example, in the Japanese eel (*A. japonica*), both

identified AR sequences clustered with ARβ sequences in the "Androgen Receptor Beta"

group, suggesting this species has two ARβ receptors. It is unclear whether these two

receptors are encoded by distinct genes, but unlike in other teleosts, no significant

differences in the androgen-binding capacity of the two eel receptors has been detected

(Peñaranda et al., 2014). These results suggest that the eel ARα and ARβ known in the

literature (Ikeuchi et al., 1999; Todo et al., 1999) are either splice variants or paralogs of

the teleost ARβ. It is intriguing to speculate whether these sequences represent evidence

of early loss of function (e.g., non-functionalization) of the ARα receptor in this lineage.

The AR sequences in rainbow trout *(O. mykiss)* also clustered with ARβ sequences in the

"Androgen Receptor Beta" group, which is consistent with prior work suggesting that

these paralogs resulted from a lineage specific WGD occurred in the Salmoniformes

(Allendorf et al., 1984; Takeo and Yamashita, 1999). The framework we propose can

provide researchers with novel hypotheses to test, enabling them to disentangle AR

evolution and selection in teleost fishes and vertebrates more broadly.

Most of the name changes we propose are straightforward, aligning

inconsistencies between species with what is otherwise a consensus within the field.

Proposing any universal naming system for certain proteins and genes, however, remains

a challenge. For example, some teleosts, such as *D. rerio,* only have one described AR

protein and one AR gene. Traditionally, animals with a single AR use the name AR (*ar*).

Our analysis shows, however, that the AR protein sequence in *D. rerio* clusters with ARβ

sequences. Thus, this sequence should be referred to as ARβ (*ar2*) based on our

proposed nomenclature system in teleosts, which takes into account that the vast majority

of teleost species possess two ARs (ARα and ARβ). We propose that for such cases,

including widely studied models like zebrafish and goldfish, a phylogenetic approach

should be used to unambiguously identify that the gene in question belongs to the ARβ

(*ar2*) lineage of receptors. When, as is the case of this example, an extensive body of

literature exists that retains the ambiguous naming convention, we suggest that explicit

reference is made to the protein or gene of interest's evolutionary context. By rendering

synonymous the original name AR or *ar* and the more nuanced ARβ or *ar2*, we believe

future research will facilitative collaboration as well as yield comparative insights within

the diverse teleost clade.

5. CONCLUSIONS

Within the most popular genetic model systems, such as house mice, there is

broad consensus on names of some of the most widely-studied genes that are relevant

to numerous subfields of biology. This is clearly the case for steroid receptors, in which

the names of proteins and genes encoding steroid receptors are consistent across the

literature. For teleost fishes, where studies of steroid signaling functions are extensive

and broadly studied, the names used for these proteins and genes have been

inconsistently assigned. Our study addresses this challenge by providing a cohesive

nomenclature system for steroid receptors in teleosts, an essential step for enhancing

communication and promoting collaborative efforts. More broadly, we hope our analysis

and recommendations will propagate a shift towards consistency in the names used for

these widely studied proteins and genes, enabling us to better understand the evolution

and function of steroid receptors across taxa.

DECLARATION OF COMPETING INTEREST

The authors declare no competing or financial interests.

DATA AVAILABILITY

A complete list of sources and accession numbers for the protein sequences used

in our phylogenetic analyses are available in the Supplementary Material (Tables S1-S5).

ACKNOWLEDEGMENTS

We thank members of the Alward Lab for their intellectual contributions to the ideas

discussed in this paper. This research was funded by NIGMS grant R35GM142799 (to

B.A.A.), University of Houston – National Research University Fund grant R0503962 (to

B.A.A.), and a Beckman Young Investigator Award (to B.A.A.).

AUTHOR CONTRIBUTIONS

A.P.H. conceived of the original idea for the manuscript. B.A.A., A.P.H., and

K.M.M. conceived of the approach to analyses and final framing of the ideas presented

in the manuscript. A.P.H. and K.M.M. chose the species to include in the analyses. A.P.H.

conducted the analyses, generated the phylogenies, and made the figures. K.M.M. did

the protein sequence collection, made the table in the main text, and made the

supplementary material. B.A.A., A.P.H., and K.M.M. edited final versions of the

supplementary tables. A.P.H. and K.M.M. wrote the methods and the results sections;
B.A.A., A.P.H., and K.M.M. wrote the first draft of the introduction; and B.A.A. wrote the
first draft of the discussion and conclusion. K.M.M. wrote the abstract, highlights, and
references. K.M.M. edited significant portions of all sections of subsequent drafts. All
authors reviewed, edited, and approved the final versions of the manuscript and
supplementary files.
572

REFERENCES

Alward, B.A., Hoadley, A.P., Jackson, L.R., Lopez, M.S., 2023. Genetic dissection of steroid-hormone modulated social behavior: Novel paralogous genes are a boon for discovery. Horm Behav 147, 105295. https://doi.org/10.1016/j.yhbeh.2022.105295

Alward, B.A., Laud, B., Skalnik, C.J., York, R.A., Juntii, S., Fernald, R.D., 2020. Modular Genetic Control of Social Status in a cichlid fish. PNAS 1–30.

Baker, M.E., 2004. Evolution of Steroid Receptors, in: Martini, L. (Ed.) Encyclopedia of Endocrine Diseases. Academic Press, pp. 312-319.

Barman, H.K., Rasal, K.D., Chakrapani, V., Ninawe, A.S., Vengayil, D.T., Asrafuzzaman, S., Sundaray, J.K., Jayasankar, P., 2017. Gene editing tools: state-of-the-art and the road ahead for the model and non-model fishes. Transgenic Res. 26, 577-589.

Betancur-R, R., Wiley, E.O., Arratia, G., Acero, A., Bailly, N., Miya, M., Lecointre, G., Ortí, G., 2017. Phylogenetic classification of bony fishes. BMC Evol Biol 17, 162. https://doi.org/10.1186/s12862-017-0958-3.

Brinton, R.D., Thompson, R.F., Foy, M.R., Baudry, M., Wang, J., Finch, C.E., Morgan, T.E., Pike, C.J., Mack, W.J., Stanczyk, F.Z., Nilsen, J., 2008. Progesterone receptors: Form and function in brain. Front. Neuroendocrinol. 29(2), 313-339.

Brunet, F.G., Roest Crollius, H., Paris, M., Aury, J.-M., Gibert, P., Jaillon, O., Laudet, V., Robinson-Rechavi, M., 2006. Gene loss and evolutionary rates following whole-genome duplication in teleost fishes. Mol Biol Evol 23, 1808–1816. https://doi.org/10.1093/molbev/msl049

Carson-Jurica, M.A., Schrader, W.T., O'Malley, B.W., 1990. Steroid receptor family: Structure and functions. Endocr Rev 11, 201–220. https://doi.org/10.1210/edrv-11-2-201

Chen, P., Li, B., Ou-Yang, L., 2022. Role of estrogen receptors in health and disease. Front. Endocrinol. 13, 839005.

Conte, M.A., Gammerdinger, W.J., Bartie, K.L., Penman, D.J., Kocher, T.D., 2017. A high quality assembly of the Nile Tilapia (*Oreochromis niloticus*) genome reveals the structure of two sex determination regions. BMC Genomics 18(1), 341.

Crowder, C.M., Lassiter, C.S., Gorelick, D.A., 2018. Nuclear Androgen Receptor Regulates Testes Organization and Oocyte Maturation in Zebrafish. Endocrinology 159, 980–993. https://doi.org/10.1210/en.2017-00617

Davey, R.A., Grossmann, M., 2016. Androgen receptor structure, function and biology: From bench to bedside. Clin. Biochem. Rev. 37(1), 3-15.

Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res 32, 1792–1797.
https://doi.org/10.1093/nar/gkh340

Eick, G.N., Thornton, J.W., 2011. Evolution of steroid receptors from an estrogen-
sensitive ancestral receptor. Mol. Cell. Endocrinol. 334(1-2), 31-38.

Faught, E., Vijayan, M.M., 2019. Loss of the glucocorticoid receptor in zebrafish improves
muscle glucose availability and increases growth. Am J Physiol Endocrinol Metab
316, E1093–E1104. https://doi.org/10.1152/ajpendo.00045.2019

Faught, E., Vijayan, M.M., 2018. The mineralocorticoid receptor is essential for stress
axis regulation in zebrafish larvae. Sci Rep 8, 1–11.
https://doi.org/10.1038/s41598-018-36681-w

Force, A., Lynch, M., Pickett, F.B., Amores, A., Yan, Y., Postlethwait, J., 1999.
Preservation of Duplicate Genes by Complementary, Degenerative Mutations.
Genetics 151, 1531–1545. https://doi.org/10.1093/genetics/151.4.1531

Glasauer, S.M.K., Neuhauss, S.C.F., 2014. Whole-genome duplication in teleost fishes
and its evolutionary consequences. Mol. Genet. Genomics 289(6), 1045-1060.

Godwin, J. (2010). Neuroendocrinology of sexual plasticity in teleost fishes. Frontiers in
Neuroendo. 31, 203-216.

Hoegg, S., Brinkmann, H., Taylor, J.S., Meyer, A., 2004. Phylogenetic timing of the fish-
specific genome duplication correlates with the diversification of teleost fish. J
Mol Evol 59, 190–203. https://doi.org/10.1007/s00239-004-2613-z

Howe, K., Clark, M.D., Torroja, C.F., Torrance, J., Berthelot, C., Muffato, M., Collins, J.E.,
Humphray, S., McLaren, K., Matthews, L., McLaren, S., Sealy, I., Caccamo, M.,
Churcher, C., Scott, C., Barrett, J.C., Koch, R., Rauch, G.-J., White, S., Chow, W.,
Kilian, B., Quintais, L.T., Guerra-Assunção, J.A., Zhou, Y., Gu, Y., Yen, J., Vogel,
J.-H., Eyre, T., Redmond, S., Banerjee, R., Chi, J., Fu, B., Langley, E., Maguire,
S.F., Laird, G.K., Lloyd, D., Kenyon, E., Donaldson, S., Sehra, H., Almeida-King,
J., Loveland, J., Trevanion, S., Jones, M., Quail, M., Willey, D., Hunt, A., Burton,
J., Sims, S., McLay, K., Plumb, B., Davis, J., Clee, C., Oliver, K., Clark, R., Riddle,
C., Elliott, D., Threadgold, G., Harden, G., Ware, D., Begum, S., Mortimore, B.,
Kerry, G., Heath, P., Phillimore, B., Tracey, A., Corby, N., Dunn, M., Johnson, C.,
Wood, J., Clark, S., Pelan, S., Griffiths, G., Smith, M., Glithero, R., Howden, P.,
Barker, N., Lloyd, C., Stevens, C., Harley, J., Holt, K., Panagiotidis, G., Lovell, J.,
Beasley, H., Henderson, C., Gordon, D., Auger, K., Wright, D., Collins, J., Raisen,
C., Dyer, L., Leung, K., Robertson, L., Ambridge, K., Leongamornlert, D., McGuire,
S., Gilderthorp, R., Griffiths, C., Manthravadi, D., Nichol, S., Barker, G., Whitehead,
S., Kay, M., Brown, J., Murnane, C., Gray, E., Humphries, M., Sycamore, N.,
Barker, D., Saunders, D., Wallis, J., Babbage, A., Hammond, S., Mashreghi-
Mohammadi, M., Barr, L., Martin, S., Wray, P., Ellington, A., Matthews, N.,
Ellwood, M., Woodmansey, R., Clark, G., Cooper, J.D., Tromans, A., Grafham, D.,

Skuce, C., Pandian, R., Andrews, R., Harrison, E., Kimberley, A., Garnett, J.,
Fosker, N., Hall, R., Garner, P., Kelly, D., Bird, C., Palmer, S., Gehring, I., Berger,
A., Dooley, C.M., Ersan-Ürün, Z., Eser, C., Geiger, H., Geisler, M., Karotki, L., Kirn,
A., Konantz, J., Konantz, M., Oberländer, M., Rudolph-Geiger, S., Teucke, M.,
Lanz, C., Raddatz, G., Osoegawa, K., Zhu, B., Rapp, A., Widaa, S., Langford, C.,
Yang, F., Schuster, S.C., Carter, N.P., Harrow, J., Ning, Z., Herrero, J., Searle,
S.M.J., Enright, A., Geisler, R., Plasterk, R.H.A., Lee, C., Westerfield, M., de Jong,
P.J., Zon, L.I., Postlethwait, J.H., Nüsslein-Volhard, C., Hubbard, T.J.P., Crollius,
H.R., Rogers, J., Stemple, D.L., 2013. The zebrafish reference genome sequence
and its relationship to the human genome. Nature 496(7446), 498-503.

Hughes, L.C., Ortí, G., Huang, Y., Sun, Y., Baldwin, C.C., Thompson, A.W., Arcila, D.,
Betancur-R., R., Li, C., Becker, L., Bellora, N., Zhao, X., Li, X., Wang, M., Fang,
C., Xie, B., Zhou, Z., Huang, H., Chen, S., Venkatesh, B., Shi, Q., 2018.
Comprehensive phylogeny of ray-finned fishes (Actinopterygii) based on
transcriptomic and genomic data. Proc. Natl. Acad. Sci. U.S.A. 115, 6249–6254.
https://doi.org/10.1073/pnas.1719358115

Jaillon, O., Aury, J.-M., Brunet, F., Petit, J.-L., Stange-Thomann, N., Mauceli, E.,
Bouneau, L., Fischer, C., Ozouf-Costaz, C., Bernot, A., Nicaud, S., Jaffe, D.,
Fisher, S., Lutfalla, G., Dossat, C., Segurens, B., Dasilva, C., Salanoubat, M.,
Levy, M., Boudet, N., Castellano, S., Anthouard, V., Jubin, C., Castelli, V., Katinka,
M., Vacherie, B., Biémont, C., Skalli, Z., Cattolico, L., Poulain, J., De Berardinis,
V., Cruaud, C., Duprat, S., Brottier, P., Coutanceau, J.-P., Gouzy, J., Parra, G.,
Lardier, G., Chapple, C., McKernan, K.J., McEwan, P., Bosak, S., Kellis, M., Volff,
J.-N., Guigó, R., Zody, M.C., Mesirov, J., Lindblad-Toh, K., Birren, B., Nusbaum,
C., Kahn, D., Robinson-Rechavi, M., Laudet, V., Schachter, V., Quétier, F., Saurin,
W., Scarpelli, C., Wincker, P., Lander, E.S., Weissenbach, J., Crollius, H.R., 2004.
Genome duplication in the teleost fish *Tetraodon nigroviridis* reveals the early
vertebrate proto-karyotype. Nature 431(7011), 946-957.

Jukes, T.H., Cantor, C.R., 1969. Evolution of Protein Molecules, in: Mammalian Protein
Metabolism. Elsevier, pp. 21–132. https://doi.org/10.1016/B978-1-4832-3211-
9.50009-7

Juntti, S., 2019. The future of gene-guided neuroscience research in non-traditional
model organisms. Brain Behav Evol. https://doi.org/10.1159/000500072

Juntti, S.A., Tollkuhn, J., Wu, M.V., Fraser, E.J., Soderborg, T., Tan, S., Honda, S.-I.,
Harada, N., Shah, N.M., 2010. The androgen receptor governs the execution, but
not programming, of male sexual and territorial behaviors. Neuron 66, 260–72.
https://doi.org/10.1016/j.neuron.2010.03.024

Kasahara, M., Naruse, K., Sasaki, S., Nakatani, Y., Qu, W., Ahsan, B., Yamada, T.,
Nagayasu, Y., Doi, K., Kasai, Y., Jindo, T., Kobayashi, D., Shimada, A., Toyoda,
A., Kuroki, Y., Fujiyama, A., Sasaki, T., Shimizu, A., Asakawa, S., N., S.,
Hashimoto, S., Yang, J., Lee, Y., Matsushima, K., Sugano, S., Sakaizumi, M.,

Narita, T., Ohishi, K., Haga, S., Ohta, F., Nomoto, H., Nogata, K., Morishita, T.,
Endo, T., Shin-I, T., Takeda, H., Morishita, S., Kohara, Y., 2007. The medaka draft
genome and insights into vertebrate genome evolution. Nature 447(7145), 714-
719.

Lien, S., Koop, B.F., Sandve, S.R., Miller, J.R., Kent, M.P., Nome, T., Hvidsten, T.R.,
Leong, J.S., Minkley, D.R., Zimin, A., Grammes, F., Grove, H., Gjuvsland, A.,
Walenz, B., Hermansen, R.A., von Schalburg, K., Rondeau, E.B., Di Genova, A.,
Samy, J.K.A., Olav Vik, J., Vigeland, M.D., Caler, L., Grimholt, U., Jentoft, S., Inge
Våge, D., de Jong, P., Moen, T., Baranski, M., Palti, Y., Smith, D.R., Yorke, J.A.,
Nederbragt, A.J., Tooming-Klunderud, A., Jakobsen, K.S., Jiang, X., Fan, D., Hu,
Y., Liberles, D.A., Vidal, R., Iturra, P., Jones, S.J.M., Jonassen, I., Maass, A.,
Omholt, S.W., Davidson, W.S., 2016. The Atlantic salmon genome provides
insights into rediploidization. Nature 533(7602), 200-205.

Liu, Z., Liu, S., Yao, J., Bao, L., Zhang, J., Li, Y., Jiang, C., Sun, L., Wang, R., Zhang, Y.,
Zhou, T., Zeng, Q., Fu, Q., Gao, S., Li, N., Koren, S., Jiang, Y., Zimin, A., Xu, P.,
Phillippy, A.M., Geng, X., Song, L., Sun, F., Li, C., Wang, X., Chen, A., Jin, Y.,
Yuan, Z., Yang, Y., Tan, S., Peatman, E., Lu, J., Qin, Z., Dunham, R., Li, Z.,
Sonstegard, T., Feng, J., Danzmann, R.G., Schroeder, S., Scheffler, B., Duke,
M.V., Ballard, L., Kucuktas, H., Kaltenboeck, L., Liu, H., Armbruster, J., Xie, Y.,
Kirby, M.L., Tian, Y., Flanagan, M.E., Mu, W., Waldbieser, G.C., 2016. The
channel catfish genome sequence provides insights into the evolution of scale
formation in teleosts. Nat. Commun. 7, 11757.

Lorin, T., Salzburger, W., Böhne, A., 2015. Evolutionary fate of the androgen receptor-
Signaling pathway in ray-finned fishes with a special focus on cichlids. G3: Genes,
Genomes, Genetics 5, 2275–2283. https://doi.org/10.1534/g3.115.020685

Lu, J., Fang, W., Huang, J., Li, S., 2021. The application of genome editing technology in
fish. Mar. Life Sci. Technol. 3, 326-346.

Mani, S.K., Blaustein, J.D., O'Malley, B.W., 1997. Progesterone receptor function from a
behavioral perspective. Horm. Behav. 31(3), 244-255.

Maruska, K.P., Butler, J.M., Anselmo, C., Tandukar, G., 2020. Distribution of aromatase
in the brain of the African cichlid fish Astatotilapia burtoni: Aromatase expression,
but not estrogen receptors, varies with female reproductive-state. Journal of
Comparative Neurology 528, 2499–2522. https://doi.org/10.1002/cne.24908

Muto, A., Taylor, M.R., Suzawa, M., Korenbrot, J.I., Baier, H., 2013. Glucocorticoid
receptor activity regulates light adaptation in the zebrafish retina. Front Neural
Circuits 7, 1–8. https://doi.org/10.3389/fncir.2013.00145

Nath, S., Shaw, D.E., White, M.A., 2021. Improved contiguity of the threespine stickleback
genome using long-read sequencing. G3 (Bethesda) 11(2), jkab007.

Nelson, J.S., 1994. Fishes of the World, 3 ed. John Wiley and Sons, New York.

Nishiike, Y., Miyazoe, D., Togawa, R., Yokoyama, K., Nakasone, K., Miyata, M., Kikuchi, Y., Kamei, Y., Todo, T., Ishikawa-Fujiwara, T., Ohno, K., Usami, T., Nagahama, Y., Okubo, K., 2021. Estrogen receptor 2b is the major determinant of sex-typical mating behavior and sexual preference in medaka. Current Biology 31, 1699-1710.e6. https://doi.org/10.1016/j.cub.2021.01.089

Ogino, Y., Kuraku, S., Ishibashi, H., Miyakawa, H., Sumiya, E., Miyagawa, S., Matsubara, H., Yamada, G., Baker, M.E., Iguchi, T., 2016. Neofunctionalization of androgen receptor by Gain-of-Function mutations in teleost fish lineage. Mol Biol Evol 33, 228–244. https://doi.org/10.1093/molbev/msv218

Ohno, S., 1970. The Enormous Diversity in Genome Sizes of Fish as a Reflection of Nature's Extensive Experiments with Gene Duplication. Transactions of the American Fisheries Society 99, 120–130. https://doi.org/10.1577/1548-8659(1970)99<120:TEDIGS>2.0.CO;2

Pascual-Le Tallec, L., Lombès, M., 2005. The mineralocorticoid receptor: A journey exploring its diversity and specificity of action. Mol. Endocrinol. 19(9), 2211-2221.

Peichel, C.L., McCann, S.R., Ross, J.A., Naftaly, A.F.S., Urton, J.R., Cech, J.N., Grimwood, J., Schmutz, J., Myers, R.M., Kingsley, D.M., White, M.A., 2020. Assembly of the threespine stickleback Y chromosome reveals convergent signatures of sex chromosome evolution. Genome Biol. 21(1), 177.

Peichel, C.L., Sullivan, S.T., Liachko, I., White, M.A., 2017. Improvement of the threespine stickleback genome using a Hi-C-based proximity-guided assembly. J. Hered. 108(6), 693-700.

Saitou, N., Nei, M., 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. Mol Biol Evol 4, 406–425. https://doi.org/10.1093/oxfordjournals.molbev.a040454

Sato, Y., Nishida, M., 2010. Teleost fish with specific genome duplication as unique models of vertebrate evolution. Environ. Biol. Fishes 88, 169-188.

Szego, C.M., Nemere, I., Pietras, R.J., 2003. Membrane Steroid Receptors, in: Henry, H.L., Norman, A.W. (Eds.), Encyclopedia of Hormones. Academic Press, pp. 657-671.

Thornton, J.W., 2001. Evolution of vertebrate steroid receptors from an ancestral estrogen receptor by ligand exploitation and serial genome expansions. Proc. Natl. Acad. Sci. USA 98(10), 5671-5676.

Thornton, J.W., Need, E., Crews, D., 2003. Resurrecting the ancestral steroid receptor: ancient origin of estrogen signaling. Science 301(5640), 1714-1717.

Timmermans, S., Souffriau, J., Libert, C., 2019. A general introduction to glucocorticoid biology. Front. Immunol. 10, 1545.

Volff, J.-N., 2005. Genome evolution and biodiversity in teleost fish. Heredity 94(3), 280-
294.

Whirledge, S., Cidlowski, J.A., 2019. Chapter 5 - Steroid Hormone Action, in: Strauss III,
J.F., Barbieri, R.L. (Eds.), Yen & Jaffe's Reproductive Endocrinology: Physiology,
Pathophysiology, and Clinical Management. Elsevier, pp. 115-131.e114.

Yin, Y., Tang, H., Liu, Y., Chen, Y., Li, G., Liu, X., Lin, H., 2017. Targeted disruption of
aromatase reveals dual functions of cyp19a1a during sex differentiation in
zebrafish. Endocrinology 158, 3030–3041. https://doi.org/10.1210/en.2016-1865

Yong, L., Thet, Z., Zhu, Y., 2017. Genetic editing of the androgen receptor contributes to
impaired male courtship behavior in zebrafish. Journal of Experimental Biology
220, 3017–3021. https://doi.org/10.1242/jeb.161596

Ziv, L., Muto, A., Schoonheim, P.J., Meijsing, S.H., Strasser, D., Ingraham, H.A., Schaaf,
M.J.M., Yamamoto, K.R., Baier, H., 2013. An affective disorder in zebrafish with
mutation of the glucocorticoid receptor. Mol Psychiatry 18, 681–691.
https://doi.org/10.1038/mp.2012.64

Zhu, B., Ge, W., 2018. Genome editing in fishes and their applications. Gen. Comp.
Endocrinol. 257, 3-12.

798 **TABLES**

800 **Table 1.** Proposed nomenclature for proteins and genes encoding steroid receptors in

801 teleost fishes.

Protein	Gene	Figure
AR		
ARα	*ar1*	Fig. 2
ARβ	*ar2*	Fig. 2
ER		
ERα	*esr1*	Fig. 3
ERβ1	*esr2a*	Fig. 3
ERβ2	*esr2b*	Fig. 3
ERγ	*esr3*	Fig. 4
ERγ1	*esr3*	Fig. 4
ERγ2	*esr3*	Fig. 4
PR	*pgr*	
PRα	*pgr1*	Fig. 5
PRβ	*pgr2*	Fig. 5
GR	*nr3c1*	
GRα	*nr3c1a*	Fig. 6
GRβ	*nr3c1b*	Fig. 6
MR	*nr3c2*	Fig. 7

803
Proposed protein and gene names for androgen receptors, estrogen receptors, progesterone receptors,

glucocorticoid receptors, and mineralocorticoid receptors in teleost fishes based on phylogenetic analysis.

The location of figures containing phylogenies associated with each steroid receptor subtype are also

shown. *Abbreviations: AR, androgen receptor; ar1, androgen receptor 1 gene; ARα, androgen receptor*

alpha protein; ar2, androgen receptor 2 gene; ARβ, androgen receptor beta protein; ER, estrogen

receptor; ERα, estrogen receptor alpha protein; ERβ, estrogen receptor beta protein; ERβ1, estrogen

receptor beta 1 protein; ERβ2, estrogen receptor beta 2 protein; ERγ, estrogen-related receptor gamma

protein; ERγ1, estrogen-related receptor gamma 1 protein; ERγ2, estrogen-related receptor gamma 2

protein; esr1, estrogen receptor 1 gene; esr2a, estrogen receptor 2a gene; esr2b, estrogen receptor 2b

gene; esr3, estrogen receptor 3 gene; GR, glucocorticoid receptor protein; GRα, glucocorticoid receptor

alpha protein; GRβ, glucocorticoid receptor beta protein; MR, mineralocorticoid receptor; nr3c1, nuclear

receptor subfamily 3 group C member 1 gene; nr3c1a, nuclear receptor subfamily 3 group C member 1a

gene; nr3c1b, nuclear receptor subfamily 3 group C member 1b gene; nr3c2, nuclear receptor subfamily

3 group C member 2 gene; pgr, progesterone receptor gene; pgr1, progesterone receptor 1 gene; pgr2,

progesterone receptor 2 gene; PR, progesterone receptor; PRα, progesterone receptor alpha protein;

PRβ, progesterone receptor beta protein.

FIGURE LEGENDS

Figure 1: Phylogeny identifying the evolutionary relationships between taxa considered

in this paper [after Betancur-R et al. (2017) and Hughes et al. (2018)]. Gray and orange

circles indicate genome duplication events. *Abbreviations: 2R, second-round; 3R, third-*

round; WGD, whole-genome duplication.

Figure 2: Consensus phylogenetic tree of androgen receptors. Numbers shown at each

branch node indicate bootstrap values (%), and the scale bar indicates phylogenetic

distance (0.1 substitutions/site). The orange circle represents the approximate position of

the teleost-specific whole-genome duplication event (TS-WGD).

Figure 3: Consensus phylogenetic tree of estrogen receptors alpha (α) and beta (β).

Numbers shown at each branch node indicate bootstrap values (%), and the scale bar

indicates phylogenetic distance (0.1 substitutions/site). Orange circles represent the

approximate position of the teleost-specific whole-genome duplication event (TS-WGD),

which impacted the independent duplication of ERβ, but not ERα.

Figure 4: Consensus phylogenetic tree of estrogen-related receptor gamma (ERγ).

Numbers shown at each branch node indicate bootstrap value (%), and the scale bar

indicates phylogenetic distance (0.1 substitutions/site). The orange circle represents the

approximate position of the teleost-specific whole-genome duplication event (TS-WGD).

Figure 5: Consensus phylogenetic tree of progesterone receptor. Numbers shown at

each branch node indicate bootstrap value (%), and the scale bar indicates phylogenetic

distance (0.1 substitutions/site). The orange circle represents the approximate position of

the teleost-specific whole-genome duplication event (TS-WGD).

Figure 6: Consensus phylogenetic tree of glucocorticoid receptors. Numbers shown at
each branch node indicate bootstrap value (%), and the scale bar indicates phylogenetic
distance (0.1 substitutions/site). The orange circle represents the approximate position of
the teleost-specific whole-genome duplication event (TS-WGD).
Figure 7: Consensus phylogenetic tree of mineralocorticoid receptor. Numbers shown at
each branch node indicate bootstrap value (%), and the scale bar indicates phylogenetic
distance (0.1 substitutions/site). The orange circle represents the approximate position of
the teleost-specific whole-genome duplication event (TS-WGD).

879 **FIGURES**

880 **Figure 1.**

Figure 2.

0.1 substitutions / site

906   **Figure 3.**

915 **Figure 4.**

0.1 substitutions / site

933   **Figure 5.**

Figure 6.

975 **Figure 7.**