RNA binding protein PRRC2B mediates translation of specific proteins and regulates cell cycle progression

Feng Jiang[1,2], Omar M. Hedaya[1,2], EngSoon Khor[1], Jiangbin Wu[1], Matthew Auguste[3], Peng Yao[1,2,4,5]*

[1] Aab Cardiovascular Research Institute, Department of Medicine, University of Rochester School of Medicine & Dentistry, Rochester, New York 14642

[2] Department of Biochemistry & Biophysics, University of Rochester School of Medicine & Dentistry, Rochester, New York 14642

[3] Undergraduate Program in Biology and Medicine, Department of Biological Sciences: Molecular Genetics, University of Rochester, Rochester, New York 14642

[4] The Center for RNA Biology, University of Rochester School of Medicine & Dentistry, Rochester, New York 14642

[5] The Center for Biomedical Informatics, University of Rochester School of Medicine & Dentistry, Rochester, New York 14642

Correspondence to Peng Yao, Ph.D., Aab Cardiovascular Research Institute, Department of Medicine, University of Rochester School of Medicine & Dentistry, Rochester, New York 14642.

E-mail: peng_yao@urmc.rochester.edu

Keywords: PAR-CLIP, PRRC2B, eIF4G2, mRNA translation, cell proliferation

Short title: PRRC2B regulates mRNA translation and cell proliferation

Abstract

Accumulating evidence suggests that posttranscriptional control of gene expression, including RNA splicing, transport, modification, translation, and degradation, primarily relies on RNA binding proteins (RBPs). However, the functions of many RBPs remain understudied. Here, we characterized the function of a novel RBP, Proline-Rich Coiled-coil 2B (PRRC2B). Through photoactivatable ribonucleoside-enhanced crosslinking and immunoprecipitation and sequencing (PAR-CLIP-seq), we identified transcriptome-wide CU- or GA-rich PRRC2B binding sites around the translation initiation codon on a specific cohort of mRNAs in HEK293T cells. These mRNAs, including oncogenes and cell cycle regulators such as *CCND2* (cyclin D2), exhibited decreased translation upon PRRC2B knockdown as revealed by polysome-associated RNA-seq, resulting in reduced G1/S phase transition and cell proliferation. Antisense oligonucleotides blocking PRRC2B interactions with *CCND2* mRNA decreased its translation, thus inhibiting G1/S transition and cell proliferation. Mechanistically, PRRC2B interactome analysis revealed RNA-independent interactions with eukaryotic translation initiation factors 3 (eIF3) and 4G2 (eIF4G2). The interaction with translation initiation factors is essential for PRRC2B function since the eIF3/eIF4G2-interacting defective mutant, unlike wild-type PRRC2B, failed to rescue the translation deficiency or cell proliferation inhibition caused by PRRC2B knockdown. Altogether, our findings reveal that PRRC2B is essential for efficient translation of specific proteins required for cell cycle progression and cell proliferation.

Introduction

RNA-binding proteins (RBPs) are vital regulators affecting the fate of nearly all classes of RNA molecules (1,2). Through one or multiple RNA-binding domains (3,4), RBPs interact selectively or globally with RNAs throughout their lifespan with temporal, spatial, and functional dynamics (5). Once recruited to RNAs, RBPs participates in posttranscriptional regulation of gene expression, including RNA splicing, transport, modification, translation, and degradation (6,7). Accumulating evidence has demonstrated the essentiality of RBP-directed posttranscriptional regulations at cellular and organismal levels (7) with alterations in the abundance or functionality of RBPs linked with various human disorders (8,9). Thus, exploring the role of RBPs in posttranscriptional regulation of gene expression contributes significantly to our understanding of basic biology and human diseases.

mRNA translation is one of the posttranscriptional processes affected by RBPs extensively (10). As the most energy-consuming and precisely regulated process in cells, translation is tightly controlled through cis-acting RNA elements such as terminal oligopyrimidine (TOP) motifs and CA-rich elements (11-13) and through trans-acting factors such as mRNA binding proteins (mRBPs) (2,10,14). The latter affects all steps in global or transcript-specific translation (10). Numerous mRBPs and (m)RNA-binding domains have been identified through mass-spectrometry-based "mRNA interactome capture" methods where probes, mostly deoxythymidine oligonucleotides (oligo(dT)), were used to capture poly(A)-bearing mRNAs together with their interacting mRBPs after ultraviolet (UV) light crosslinking (15-17). These methods identify mRBPs physically bound to sequence- or structure-specific elements in mRNAs that modulate gene expression at the posttranscriptional level (6). However, despite continuous attempts at functional characterization of individual mRBPs (5), the majority of the newly identified mRBPs remain unstudied.

One of the functionally unannotated mRBPs that may regulate translation is P̲ro-r̲ich C̲oiled-coil 2B (PRRC2B). PRRC2B is identified as an mRNA-binding protein based on its enrichment with the oligo(dT) captured mRNAs across multiple different cell types (15,18) and its putative arginine-glycine (RG)-rich domains documented to interact with

RNA (19,20). Previous studies suggested that PRRC2B is part of the eukaryotic initiation factor 4G2 (eIF4G2)-mediated translation initiation complex. This is based on its co-immunoprecipitation with eIF4G2 and eukaryotic translation initiation factor 3 (eIF3) in human breast cancer cells and mouse embryonic stem cells (21-23). However, the exact function and mRNA targets of PRRC2B in translational regulation remain unknown.

Although understudied at the mechanistic level, PRRC2B has been linked with various human cancers. PRRC2B is highly expressed in fast-proliferating tumor cells such as Wilms' tumor, and its high expression is tightly associated with poor overall survival of patients (24,25). In addition, fusion genes of PRRC2B with MGMT, DEK, and ALK have been reported in multiple different tumors (26-30). Therefore, understanding the function of PRRC2B may shed light on translational regulation with relevance to tumor cell proliferation and human diseases.

In this study, we perform a comprehensive characterization of PRRC2B in translational regulation of gene expression in human embryonic kidney (HEK293T) cells. We demonstrate that PRRC2B and a 750 amino acid-long fragment of PRRC2B bind to GA- or CU-rich elements around the translation initiation codon of the main open reading frame on a subset of mRNAs, promoting their translation via interacting with translation initiation factors. Loss of PRRC2B binding leads to decreased translation efficiency of PRRC2B-bound mRNAs encoding oncogenes and cell cycle regulators such as cyclin D2, resulting in reduced cell cycle progression. Altogether, our study revealed the function of PRRC2B in translational regulation of specific proteins affecting cell proliferation.

Results

PRRC2B directly interacts with a subset of mRNAs

We first applied photoactivatable ribonucleoside-enhanced crosslinking and immunoprecipitation (PAR-CLIP) (31) to identify transcriptome-wide PRRC2B binding sites in HEK293T cells. During PAR-CLIP, RBPs were crosslinked to their bound RNAs by 4-thiouridine (4sU) and UV exposure, enriched by immunoprecipitation (IP) after RNase T1 digestion, and visualized on Bis-Tris gel by radioactive labeling of bound RNAs (**Figure 1A, S1A**). Following this method, we enriched FLAG-tagged full-length and three truncated variants of PRRC2B (P1, P2, and P3) (**Figure 1B**). While the extent of enrichment is comparable (**Figure S1B**), only full-length and 750 amino acid-long RG-rich motif containing P2 variant of PRRC2B showed significant RNA signal at the position of the protein with corresponding size (**Figure 1C**). Crosslinked RNAs (mostly 19-35 nt) were then purified from PRRC2B and P2, visualized on polyacrylamide gel (**Figure 1D**), and subjected to library construction and deep sequencing. Following an established analysis pipeline with in-house modifications, we identified a total of 1062 binding sites for full-length PRRC2B (**Table S1**) and 749 binding sites for P2 (**Table S2**) from two biological replicates that highly correlate with each other (**Figure S1C, S1D**). Among PRRC2B and P2 CLIP sites, 462 overlapped binding sites were found (**Table S1, S2**), accounting for ~44% of the full-length binding sites and ~62% of the P2 binding sites (**Figure 1E**). The majority of the overlapped binding sites were on mRNAs (326, ~71%), with a small portion on long noncoding RNA (lncRNA) (62, ~13%), miRNA (25, ~5%), snoRNA (3, ~1%), and other RNA species (2, < 1%) (**Figure 1F**). Similar distributions were also observed in full-length PRRC2B and P2 binding sites when analyzed separately (**Figure S1E**). Further examination of the binding sites on mRNAs revealed that 5' untranslated region (5' UTR) possessed the highest density (binding site per 1000 nt), followed by the coding sequence (CDS), 3' untranslated region (3' UTR), and intron (**Figure 1G, S1F**). The enrichment in mature mRNAs (5' UTR, CDS, 3' UTR) versus introns was further supported by the fact that the majority of PRRC2B are localized in the cytosol where mature mRNAs (5' UTR, CDS, 3' UTR) are dominant (**Figure S1G**). Notably, the highest peak of binding sites was found at the start codon region of the main open

reading frames spanning 5' UTR and CDS (**Figure 1H, S1H**). Subsequent scanning of consensus sequences in the overlapped binding sites across intact mRNAs and in the sub-regions (5' UTR, CDS, 3' UTR) on mRNAs by MEME (32) showed motifs that are either GA- or CU-rich (**Figure 1I, S1I**), which hardly accrue within the same binding site. Together, these results suggest that PRRC2B directly binds to a select cohort of mRNAs at the region near translation initiation sites.

Loss of PRRC2B decreases translation of its bound mRNAs

The enrichment of PRRC2B binding sites around the translation initiation sites suggested its potential involvement in translational regulation. Accordingly, we examined changes in translation efficiency at the transcriptomic level upon PRRC2B knockdown by polysome profiling and RNA sequencing (**Figure 2A**) (33,34). A doxycycline (Dox)-inducible short hairpin RNA (shRNA) targeting PRRC2B (shPRRC2B) was used to knockdown PRRC2B with a non-targeting shRNA as a control (shCtrl) (**Figure 2B, 2C)**. RNA-seq was performed on polysome-associated mRNAs (associated to > 2 ribosomes, considered actively translated) and total mRNAs from cells 24 hours after Dox induction of PRRC2B knockdown to calculate translation efficiency (TE; ratio of FPKM value of polysome-associated RNA to that of total RNA) for each mRNA in PRRC2B knockdown (shPRRC2B) and control (shCtrl) cells. In the sequencing results, a total of 25537 mRNAs were detected of which 1455 mRNAs exhibited increased TE (Log_2 ($\text{TE}_{shPRRC2B}/\text{TE}_{shCtrl}$) > 0.5; adjusted P value < 0.05; TE-up) upon PRRC2B knockdown while 3198 mRNAs exhibited decreased TE (Log_2 ($\text{TE}_{shPRRC2B}/\text{TE}_{shCtrl}$) < -0.5; adjusted P value < 0.05; TE-down) (**Figure 2D, Table S3**). Gene Ontology (GO) analysis reveals enrichment of the TE-up genes in "pattern specification process" and "mitochondrial transport" while enrichment of TE-down genes in 'cytosolic transport' and 'regulation of mRNA processing' (**Figure S2A, S2B, Table S4**). To further elucidate the TE changes caused directly by loss of PRRC2B binding, we focused on mRNAs with the overlapped binding sites (PRRC2B-bound mRNAs) (**Table S5**). We observed significantly changed TE in 96 out of total 221 PRRC2B-bound mRNAs, among which only 17 mRNAs exhibited increased TE (Log_2 ($\text{TE}_{shPRRC2B}/\text{TE}_{shCtrl}$) > 0.5; adjusted P value < 0.05; TE-up) while 79 mRNAs showed decreased TE (Log_2 ($\text{TE}_{shPRRC2B}/\text{TE}_{shCtrl}$) < -0.5; adjusted P value < 0.05; TE-down)

(**Figure 2E**). The TE-up PRRC2B-bound mRNAs were functionally enriched in "cytoplasmic translation" and "ribosome assembly" while TE-down PRRC2B-bound mRNAs were enriched in "stem cell population maintenance" and "maintenance of cell number" (**Figure S2C, S2D, Table S6**).

Since dysregulated TE in unbound mRNAs were detected, we suspected that secondary effects might have been triggered by the primary changes due to PRRC2B knockdown. To distinguish the primary and secondary changes, we examined the translational changes of some PRRC2B-bound and unbound mRNAs at earlier time points after Dox induction by performing reverse transcription-quantitative PCR (RT-qPCR) on RNAs extracted from polysome profiling fractions (**Figure 2A**). All tested PRRC2B-bound mRNAs were first validated to interact with the endogenous PRRC2B in cells by RNA-binding protein immunoprecipitation (RIP) and RT-qPCR (**Figure S2E**). All tested TE-down PRRC2B-bound mRNAs showed shifts towards the light polysome or pre-polysome fractions (with smaller number of ribosomes on mRNAs) from heavier polysomes fractions (with larger number of ribosomes on mRNAs) at 6 or 12 hours after induction of PRRC2B knockdown, suggesting reduced translation efficiency (**Figure 2F, S2F**). Similar shifts were not observed in the TE-unchanged PRRC2B-bound mRNA (*ACTB*) or in PRRC2B-unbound mRNAs, including unbound TE-unchanged mRNAs (*GAPDH*, *GATA4*), unbound TE-down mRNAs (*PTBP1*, *HSPA4*), and unbound TE-up mRNA (*RPL37A*) (**Figure 2G, S2G**). This indicates that translation of unbound mRNAs changed at later stages (24 hours of DOX-induced PRRC2B knockdown) while translation of PRRC2B-bound mRNAs was reduced at earlier stages (6-12 hours of DOX-induced PRRC2B knockdown). Therefore, translational changes of PRRC2B-unbound mRNAs are due to secondary effects. The reduction of translation efficiency in some of the TE-down PRRC2B-bound mRNAs (*CCND2*, *CRKL*, *YWHAZ*) was further confirmed by decreased protein abundance upon PRRC2B knockdown (**Figure 2H**). Notably, the validated TE-down mRNAs possess PRRC2B binding sites in 5' UTR (*CRKL*, *CCND2*), 3' UTR (*YWHAZ*, *CTC1*, *ATAD5*), or CDS (*SERBP1*, *BPTF*, *PPP2R1A*) (**Figure 2I, S2H**), suggesting that the location of PRRC2B binding site may not affect PRRC2B function in translational control. Unlike TE-down PRRC2B-bound mRNAs, none of the tested TE-up PRRC2B-bound mRNAs showed constant shifts towards heavier polysomes (**Figure S2I**).

Together, these results suggest that loss of PRRC2B primarily decreases translation of its bound mRNAs.

PRRC2B binding to *CCND2* mRNA facilitates CCND2 protein expression

To further elucidate the role of PRRC2B binding on mRNAs, we selected cyclin D2 (*CCND2*) mRNA, which has a CU-rich PRRC2B binding site in 5' UTR (**Figure 3A**), as an example to examine the relationship between PRRC2B binding and protein expression. Antisense oligonucleotides (ASO) base-paring with the PRRC2B binding site on *CCND2* mRNA (ASO1) were used to specifically block PRRC2B binding by forming double-stranded RNA at the PRRC2B binding site (**Figure 3A**). ASO1 significantly inhibited the binding of PRRC2B, while ASO base-paring with the region adjacent to binding sites (ASO2) (**Figure 3A**) and ASO without targets in human cells (Ctrl ASO) had minor effects (**Figure 3B**). While none of the ASOs tested caused a significant alteration in *CCND2* mRNA abundance (**Figure 3C**), ASO1 significantly decreased CCND2 protein expression in wild-type cells (**Figure 3D**). Notably, decreased CCND2 protein or mRNA expression was not observed in PRRC2B knockdown cells treated with ASO1 (**Figure S3A, S3B**), confirming the requirement of PRRC2B for the mechanism of action of ASO1.

We next investigated the importance of the CU-rich elements in PRRC2B binding to *CCND2* mRNA by dual-luciferase reporter assays. Wild-type (WT-FLuc) and mutated 5' UTR of *CCND2* were cloned as the 5' UTR of the open reading frame of a firefly luciferase reporter (**Figure 3E**). These mutants include 5' UTR of *CCND2* mRNA with (i) PRRC2B binding sites replaced by an adjacent sequence showing less PRRC2B binding (Del-FLuc), (ii) all the "C" in PRRC2B binding sites mutated to "G" (C2G-FLuc), (iii) all the "U" in PRRC2B binding sites mutated to "A" (T2A-FLuc), and (iv) all the "C & U" in PRRC2B binding sites mutated to "G & A" (CT2GA-FLuc) (**Figure 3E, S3C**). As expected, the mRNA of WT-FLuc bound strongly with PRRC2B while the mRNAs of Del-FLuc, C2G-FLuc, and T2A-FLuc showed significantly decreased binding affinity (**Figure 3F**). Intriguingly, the mRNA of CT2GA-FLuc only showed a moderate decrease in PRRC2B binding (**Figure 3F**). Subsequent dual-luciferase reporter assays showed decreased luciferase activity of Del-FLuc, C2G-FLuc, and T2A-FLuc and a modest change in luciferase activity of CT2GA-FLuc in wild-type cells, which suggests that PRRC2B binding

to the reporter mRNA facilitates luciferase expression (**Figure 3G**). These effects are further confirmed to occur at the translational level since comparable abundance of the reporter mRNAs was observed despite the sequence differences (**Figure 3H**). Notably, similar trends of luciferase expression were not observed in PRRC2B knockdown cells (**Figure S3D).** Together, these results suggest that PRRC2B binding to the target mRNA, *CCND2*, promotes protein expression without changing the mRNA abundance.

PRRC2B knockdown inhibits cell proliferation

We next examined the biological consequence of PRRC2B binding on target mRNAs by focusing on cellular changes upon PRRC2B knockdown. As "maintenance of cell number" is the top functionally enriched GO term of TE-down PRRC2B-bound mRNAs (**Figure S2D, Table S6**), changes in cell proliferation were first examined. Decreased cell proliferation and increased doubling time were observed from 1 to 7 days after induction of PRRC2B knockdown (**Figure 4A, 4B**). This change could be partially explained by the observed decrease in the G1/S phase transition (**Figure 4C, 4D**). Of note, similar changes were not observed in cells without induction of PRRC2B knockdown (**Figure S4A-D**). The decreased G1/S phase transition could result from the reduced expression of cyclin D2, which has been established to facilitate the G1/S phase transition (35,36). Accordingly, decreased G1/S phase transition was observed when PRRC2B binding to *CCND2* mRNA was blocked by ASO1 (**Figure 4E, 4F**). Together, these results suggest that PRRC2B affects cell proliferation through regulating the translation of cell cycle regulators such as cyclin D2.

PRRC2B interacts with eukaryotic initiation factors eIF4G2 and eIF3

To explore mechanistically how PRRC2B regulates the translation of PRRC2B-bound mRNAs, we performed IP followed by liquid chromatography-mass spectrometry analysis to capture the interactome of endogenous PRRC2B (**Figure 5A**). Ninety-five high-confidence interactions were identified (SAINT (Significance Analysis of INTeractome) probability > 0.95) (37), of which translation initiation factors eIF4G2 and eIF3 subunits are predominant (**Figure 5B, Table S7**). The 3' UTR-binding protein FXR1 (38) known to co-IP with eIF4G2 (23) was also detected (**Figure 5B, Table S7**). These highly confident interactions were validated by western blot following IP of either endogenous or FLAG-

tagged PRRC2B (**Figure 5C, S5A**) and were further confirmed to be RNA-independent since they were retained following RNase T1 digestion of RNA to less than 35-nt long (**Figure S5B, S5C**). In contrast, no interaction of PRRC2B with eIF4G1 or cap-binding protein eIF4E was observed even though they are highly abundant and drive canonical cap-dependent translation initiation (**Figure 5B, 5C, S5A, S5C**). This suggests that PRRC2B may participate specifically in the eIF4G2-mediated translation initiation (22). This notion is further supported by co-presence of PRRC2B, eIF4G2, eIF3, and FXR1 in the 40S and 60S fractions where translation initiation is happening (**Figure 5D**). Surprisingly, a significant amount of PRRC2B was also observed in 80S and disome fractions (**Figure 5D**). Subsequent IP experiments showed that P2 and P3 variants of PRRC2B contribute to its interaction with FXR1 while P1 is primarily responsible for interactions with eIF4G2 and eIF3 (**Figure S5D**). A more detailed IP-based mapping of truncated P1 showed that the N-terminal 1-150 amino acids of PRRC2B contains eIF3 binding domain while 1-450 amino acids region interacts with eIF4G2 (**Figure S5E, S5F**). Sequences in these N-terminal regions are highly conserved among PRRC2B orthologues among vertebrates (from zebrafish to human), highlighting its functional importance (**Extended Data 1**). Together, these results suggest that PRRC2B interacts with translation initiation factors on mRNAs being translated.

PRRC2B functions in translational regulation by interacting with translation initiation factors eIF3 and eIF4G2

Subsequently, we examined whether PRRC2B functions through interacting with translation initiation factors eIF3 and eIF4G2. As eIF4G2 is a noncanonical translation initiation factor, we first examined its role in translating three PRRC2B-bound mRNAs (*CCND2*, *CRKL*, *YWHAZ*). We observed that all three proteins showed decreased expression upon siRNA-mediated eIF4G2 knockdown, suggesting that eIF4G2 is involved in translation of PRRC2B-bound mRNAs (**Figure S6A**). This notion is further supported by the fact that increased expression of the three proteins were not observed in eIF4G2 knockdown cells but in wild-type cells upon overexpressing PRRC2B (**Figure 6A**). We then asked whether interacting with eIF4G2 and eIF3 is essential for PRRC2B function. We constructed a truncated PRRC2B variant with no eIF4G2 or eIF3 binding ability (del450) (**Figure 6B, 6C**) and tested whether del450 could facilitate the translation

of PRRC2B-bound mRNAs. Our rescue experiments showed that del450 failed to fully rescue the protein expression and translational changes on *CCND2*, *CRKL*, and *YWHAZ* mRNAs caused by PRRC2B knockdown, while the full length PRRC2B rescued the protein expression successfully (**Figure 6D, S6B**), suggesting del450 cannot facilitate translation of PRRC2B-bound mRNAs. Similar changes were also observed when rescuing inhibitions in cell proliferation and G1/S phase transition by wild-type PRRC2B or del450 (**Figure 6E-G, S6C**). Together, these findings demonstrate that interaction with translation initiation factors eIf4G2 and eIF3 is essential for the function of PRRC2B in translational regulation.

Discussion

In this study, we demonstrate the role of PRRC2B, a novel RNA-binding protein, in translational regulation of PRRC2B-bound mRNAs. We discover that PRRC2B binds to CU- or GA-rich sequences near the translation initiation site, facilitating translation of a group of proteins related to cell proliferation. This effect can be inhibited by antisense oligonucleotide targeting PRRC2B binding sites in PRRC2B-bound mRNAs such as *CCND2* mRNA. Mechanistically, the function of PRRC2B primarily depends on its interaction with eIF4G2 and eIF3, suggesting a positive regulatory role of PRRC2B in eIF4G2-mediated translation. Altogether, this study reveals a novel translation regulatory pathway that maintains efficient cell proliferation and can be targeted to diminish cell cycle progression.

Using PAR-CLIP (31), we demonstrated the binding of PRRC2B to mature mRNAs. Unlike many promiscuous RNA-binding proteins, PRRC2B binds to both CU- and GA-rich RNA elements. Although CU- and GA-rich RNA sequences tend to be complementary, we assume that PRRC2B binds to single-stranded RNA since these elements seldom appear adjacent to each other, and therefore are unlikely to base pair to form double-stranded RNA. Our mutagenesis assays demonstrated that mutating a CU-rich binding site to GA-rich has minor effect on PRRC2B binding and function. From this, we conclude

that PRRC2B binds to both motifs with a similar affinity. Although CU- and GA-rich motifs are found throughout mature mRNAs (in 5' UTR, CDS, and 3' UTR), PRRC2B predominately binds the motif sequence near the translation initiation sites where it interacts with translation initiation factors eIF4G2 and eIF3. Despite the fact that eIF4G2 has been accounted for transcript selectivity (23,39), interacting with it hardly affects PRRC2B binding since the PAR-CLIP for P2, the 751-1500 amino acid-containing fragment of PRRC2B not interacting with eIF4G2 or eIF3, exhibited similar binding profile and target sequences as the full-length PRRC2B. Based on this, we conclude that P2 region of PRRC2B dictates at least some degree of transcript selectivity in an eIF4G2- and eIF3-independent manner.

eIF4G2 has been evident to drive noncanonical eIF4G1- and eIF4E-independent translation initiation (22,23,40) together with eIF3D, which replaces eIF4E in cap-binding (40), and FXR1 which binds to mRNA 3' UTR and facilitates mRNA circularization (38). From the PRRC2B interactome (**Figure 5, S5**), we conclude that PRRC2B selectively participates in eIF4G2-mediated translation initiation but not the canonical eIF4G1-mediated translation initiation. This discovery echoes previous studies on eIF4G2 (22,23) and is essential for understanding PRRC2B function (**Figure 6**). We speculate that PRRC2B may act as a scaffold recruiting other factors since it interacts with mRNA, eIF4G2, eIF3, and FXR1 through distinct regions with little dependency on each other (**Figure1, S5**). Besides, considering that PRRC2B dictates transcript selectivity, it may navigate eIF4G2 to specific mRNAs. However, given the mRNA abundance of *eIF4G2* is roughly three times of *PRRC2B* in HEK293T cells (RPKM value detected by our RNA-seq), PRRC2B likely only participate in a fraction of eIF4G2-mediated translation. In addition, PRRC2B does not completely co-sediment with translation initiation complexes in 40S to 80S fractions (**Figure 5D**), suggesting that it may have other functions beyond translation initiation (e.g., initiation-to-elongation transition). To establish the role of PRRC2B in eIF4G2-mediated translation initiation, future studies involving *in vitro* translation are required. However, obtaining recombinant PRRC2B will be significantly challenging given that full length PRRC2B is 243 KDa and highly disordered as predicted by AlphaFold (41).

Although the mechanism is not fully defined, PRRC2B facilitates eIF4G2-mediated translation of its bound mRNAs (**Figure 2, 3**). These mRNAs include multiple oncogenes and cell cycle regulators such as *CCND2* (35,36) and *CRKL* (42-44), agreeing with the link between PRRC2B and various types of human cancer (24-30). Our results suggest that PRRC2B promotes cell proliferation by facilitating translation of *CCND2* mRNA, which encodes the D-type cyclin facilitating G1/S transition in cancer and stem cells (35,45). Blocking PRRC2B binding to *CCND2* mRNA by ASOs significantly decreases the CCND2 expression in cells expressing PRRC2B (**Figure 3**), thereby inhibiting cell proliferation. This discovery inspired us about inhibiting expression of oncogene and cell cycle proteins by ASOs exclusively in cells highly expressing PRRC2B. Because PRRC2B is highly expressed in cancer cells such as large B cell lymphoma (DLBC), thymus cancer (THYM) (25), and Wilms' tumor (24), these ASOs can be applied for potential anti-cancer treatment, which deserves future investigation in biological or disease-relevant models.

While PRRC2B is evident to facilitate translation of its bound mRNAs, we still found targets to have unchanged or increased translation efficiency upon PRRC2B knockdown. The unaltered TE could possibly indicate that translation of PRRC2B-bound mRNAs may not entirely depend on PRRC2B-mediated translation, as there could be other mechanisms sufficient to carry out the translation when PRRC2B is absent. The increased TE is probably due to secondary or compensatory effects since these mRNAs, similar as the TE-changed unbound mRNAs, exhibited no translational changes at earlier time points after induction of PRRC2B knockdown (**Figure S2I**). These secondary or compensatory effects could result from the primary changes in TE of PRRC2B-bound mRNAs or the altered cell cycle progression upon PRRC2B knockdown (**Figure 4**) since translation varies dramatically in different phases of cell cycle (46).

The PRRC2 proteins have been repeatedly linked with mRNA binding and posttranscriptional regulation (21-23,47-50). A recent study suggests that PRRC2 proteins facilitate leaky scanning of upstream open reading frames (uORF) (47). However, our results showed both mRNAs with and without uORFs are bound and regulated by PRRC2B, suggesting that the effect of PRRC2B is not limited to uORFs but determined by the presence of GA- and CU-rich binding sites. Besides, previous studies showed

PRRC2A interacts with N6-methyladenosine (m^6A) around the stop codon to stabilize mRNA (48), while PRRC2C facilities efficient formation of stress granules (49,50). Although PRRC2B shares sequence similarities with PRRC2A and PRRC2C (**Extended Data 2**), the reported function of PRRC2A and PRRC2C were not recapacitated in PRRC2B (**Figure 1H, S1H, 2I, S2H**). On the other hand, PRRC2B loss-of-function mutations have been linked with human congenital heart defects (51) and mouse cardiovascular developmental defects (52), which is not observed in a PRRC2C loss-of-function mouse model (52). Therefore, the functional diversity and redundancy of the three members of PRRC2 protein family warrants future elucidation.

Material and Methods

Molecular cloning

DNA fragments were obtained and amplified by PCR from Genomic DNA or cDNA extracted from HEK293T cells by Trizol (ThermoFisher) according to manufacturer's instructions. Amplified DNA fragments were separated on 1% agarose gel, purified, digested by restriction endonucleases together with the backbone plasmid at 37°C overnight, and ligated to the cleaved backbone by T4 DNA ligase (NEB). The ligated products were transformed to *E.coli* competent cells (DH5α). Miniprep was performed on single colonies to purify the plasmids. Plasmid sequences were then confirmed by Sanger sequencing.

Cell culture and transfections

Human HEK293T cells were propagated in Dulbecco's modified Eagle's medium (DMEM; Corning) supplemented with 10% fetal bovine serum. Cells were transfected with plasmids using Lipofectamine 3000 (Invitrogen) as described by the manufacturer.

PRRC2B knockdown

shRNA cassettes were cloned into 5′-XhoI and 3′-EcoRI sites of tetracycline-inducible lentiviral pTRIPZ vector driving the expression of a TurboRFP fluorescent reporter (GE Dharmacon technology). The shRNA cassette sequences were as follows:

PRRC2B shRNA: 5′- ATCGTTTGAGGAACCAGCC -3′

Non-silencing sequence control shRNA: 5′-AATTCTCCGAACGTGTCACGT-3′

Stable single clone cell lines were generated using puromycin selection. shRNA expression was induced by 1–2 μg/ml Doxycycline and indicated as red fluorescent protein expression.

Luciferase reporter construction

The 5' UTR of *CCND2* mRNA was cloned into the pGL3-TK-5UTR-BsmBI-Luciferase reporter plasmid purchased from Addgene (https://www.addgene.org/114670/). The DNA fragment was amplified from cDNA prepared from HEK293T using primers with the extra 5' end corresponding to the BsmbI cut sites in the plasmid: 5′-AACGTCTCCACAC-3′ for

the forward primer and 5′-AACGTCTCTCTTCCAT-3′ for the reverse primer. The DNA fragment and backbone were then cleaved via the restriction enzyme BsmbI for 1 hour at 55°C and then ligated using T4 DNA ligase at room temperature, followed by selection on X-Gal coated plates. White colonies were cultured and verified using Sanger sequencing. To construct mutant 5' UTRs, Q5 high fidelity polymerase was used to introduce the desired mutations. 0.5 μl of the PCR reaction is then incubated with a mixture of T4 DNA ligase, T4 PNK, and DpnI in T4 DNA ligase buffer for 1 hour (37°C, 15 min; 25°C,30 min; 37°C, 15 min). 5 μl of this reaction is transformed into competent cells and then plated on ampicillin agarose plates.

PAR-CLIP

FLAG-tagged full-length and fragments of PRRC2B were cloned into 5′-KpnI and 3′-NotI sites in pcDNA3.1 plasmid for overexpression in HEK293T cells. PAR-CLIP was performed essentially following the protocol previously described (31) with some modifications. Briefly, cells were treated with 800 mM 4sU 16 hours before crosslinking by 312 nm ultraviolet exposure using Stratagene Stratalinker UV 1800. Crosslinked cells were lysed in a native lysis buffer with 50 mM HEPES, pH 7.5, 150 mM KCl, 2 mM EDTA, 1 mM NaF, 0.1% (v/v) Nonidet P-40, 1 mM DTT, 25 μl/ml protease inhibitor cocktail for mammalian tissues (Sigma-Aldrich), and 1U/μl RNase T1. Immunoprecipitation was performed using anti-DYKDDDDK magnetic agarose (ThermoFisher) followed by on-bead RNA digestion with 5U/μl RNase T1, on-bead dephosphorylation with calf intestinal alkaline phosphatase (CIP), and on-bead phosphorylation with T4 PNK and ^{32}[P]-ATP. Radio-labeled RNA-protein complexes were extracted from beads by boiling in 70 μl of SDS loading buffer (Bio-Rad), separated on a 4-12% Bis-Tris gel, and imaged with phosphorimager. Bands corresponding to the protein size were cut off from the gel and subjected to proteinase K digestion and RNA purification by phenol-chloroform. The purified RNAs were run on 15% denaturing polyacrylamide gel and purified for deep sequencing. Sequencing libraries were constructed using the NEBNext® Small RNA Library Prep Set following the manufacturer's instructions. Sequencing was performed on Illumina HiSeq2500 sequencer with 75 bp single-end sequencing.

PAR-CLIP data analysis

We applied the CLIP Tool Kit (CTK) to perform all steps of the analysis from raw reads to binding sites and target genes as previously described (53). Briefly, raw reads were filtered based on the mean quality score with CTK built-in function and processed by cutadapt/b1 (54) to remove adapters. Processed reads with the identical sequences were labeled as PCR duplicates and removed from downstream analysis. Mapping was performed using bowtie/1.2.1.1/b1 (55) with the --best –strata option against the Homo sapiens reference genome (GRCh38, hg38) downloaded from the UCSC genome browser (http://genome.ucsc.edu). One mismatch for each alignment was allowed to tolerate the T to C mutation introduced by 4sU crosslinking. Clustering and peak calling were performed on aligned reads with CTK built-in functions using the 'valley seeking' algorithm (53). Peak significance was first assigned based on whether the observed peak height is more than expected by chance using background models and scan statistics, then further evaluated based on the presence of T to C mutations. Significant peaks around T to C mutations were then treated as binding sites. Sequences flanking the T-to-C mutation sites (-20 to +20 nt) in the overlapped binding sites between full-length PRRC2B and P2 on mRNA exons were submitted to MEME (32) for de novo motif discovery with settings to find the longest possible motif enriched against a randomized set of sequences of the same length. Motifs with E-values < 0.05 were reported.

RNA purification and RT-qPCR

Cells were lysed by 1000 μl of Trizol (Qiagen) and mixed with 200 μl chloroform. The mixture was spun down at 16,000 g for 10 min. RNA was precipitated from the aqueous layer by adding two volumes of isopropanol and spinning down at 16,000 g for 10 min. The pellet was washed twice with 70% ethanol, left to dry, and resuspended in nuclease-free water. For quantitation of mRNA, cDNAs were prepared using iScript master mix RT Kit (Biorad) and subsequently qPCR-amplified using SYBR Primer Assay kits (Biorad). Notably, when a primer set was first used, the identity of the resulting PCR product was confirmed by cloning and Sanger sequencing. Once confirmed, melting curves were used in each subsequent PCR to verify that each primer set reproducibly and specifically generates the same PCR product.

Polysome profiling and RNA extraction

1×10^9 cells were incubated with cycloheximide (100 µg/ml; Sigma) for 10 min and harvested using a native lysis buffer with 100 mM KCl, 5 mM MgCl₂, 10 mM HEPES, pH 7.0, 0.5% Nonidet P-40, 1 mM DTT, 100 U/ml RNasin RNase inhibitor (Promega), 2 mM vanadyl ribonucleoside complexes solution (Sigma-Aldrich (Fluka BioChemika)), 25 µl/ml protease inhibitor cocktail for mammalian tissues (Sigma-Aldrich), cycloheximide (100 µg/ml). The lysate was spun down at 1,500 g for 5 min to pellet the nuclei. The supernatant was loaded onto a 10-50% sucrose gradient and spun in an ultracentrifuge at 150,000 g for 2 hours. The gradients were then transferred to a fractionator coupled to an ultraviolet absorbance detector that outputs an electronic trace across the gradient. Using a 60% sucrose chase solution, the gradient was pumped into the fractionator and divided equally into 9 or 10 fractions. For subsequent RNA extraction, 400 µl of each fraction was mixed with an equal amount of chloroform: phenol: isoamyl alcohol and 0.1 x of 3M sodium acetate (pH 5.2), and then spun down at 16,000 g for 10 min. The upper aqueous layer was mixed with 3 volumes of 100% ethanol and 0.1 volume of 3M sodium acetate (pH 5.2) and incubated at -20°C overnight. The solution was then spun at maximum speed for 30 min to pellet the RNA, which was washed twice with 70% ethanol and resuspended in nuclease-free water. For RNA deep sequencing, RNA purified from fractions #6-10 was combined as polysome-associated RNA.

RNA sequencing and data processing

Total and polysome-associated RNA samples were subjected to DNA removal and polyA enrichment before library construction by NGS Library Prep. Paired-end sequencing was conducted at Novogene using NovaSeq6000 S4 with a depth of 20M reads/sample. Reads were demultiplexed using bcl2fastq version 2.19.0. Quality filtering and adapter removal were performed using Trimmomatic version 0.36 (56) with the following parameters: "ILLUMINACLIP:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:5:25 MINLEN:32 HEADCROP:10". Processed/cleaned reads were then mapped to the Homo sapien reference genome (GRCh38, hg38) with Hisat version 2.1.0 (57) with the default settings. The subread-2.0.1(58) package (featureCounts) was used to derive gene counts based on utilizing the gencode.v38 gene annotations (59) with the following parameters: "-T 10 -g gene_name -B -C -p --ignoreDup --fracOverlap 0.1". For normalizing raw counts

and identifying significantly changed genes, DESeq2 (60) was used within an R-4.1.2 (61) environment. Gene ontology analyses were performed on significantly changed genes using clusterProfiler package version 4.0 (62).

Co-immunoprecipitation

Cells from one 80% confluent 10-cm culture dish were harvested using a native lysis buffer (300 μL) with 50 mM HEPES, pH 7.5, 150 mM KCl, 2 mM EDTA, 1 mM NaF, 0.1 % (v/v) Nonidet P-40, 1 mM DTT, and 25 μl/ ml protease inhibitor cocktail (Sigma-Aldrich) for mammalian tissues. RNasin RNase inhibitor (Promega) or RNase T1 was added for RNA-dependent and independent interactome capture. Immunoprecipitation (IP) of endogenous PRRC2B was performed using 1 μg (dilution factor 1:125) rabbit anti-human endogenous PRRC2B antibody (Invitrogen) followed by Magnetic dynabead protein A (Invitrogen) pull-down. IPs of FLAG-tagged PRRC2B and PRRC2B fragments were performed using anti-DYKDDDDK Magnetic agarose (ThermoFisher Scientific). Beads were mixed with 70 μl of SDS loading buffer (Bio-Rad), and boiled for 5 min to elute proteins for western blot and mass spectrometry analysis.

Mass spectrometry analysis

Mass spectrometry (MS) was performed by the Mass Spectrometry Resource Lab of University of Rochester Medical Center. Briefly, the IP elution was loaded onto a 4-12% gradient SDS-PAGE gel, run for 10 min, and stained with SimplyBlue SafeStain (Invitrogen). The stained regions were then excised, cut into 1mm cubes, de-stained, reduced and alkylated with DTT and IAA (Sigma), dehydrated with acetonitrile, and incubate with trypsin (Promega) at 37°C overnight. Peptides were then extracted by 0.1% TFA and 50% acetonitrile, dried down in a CentriVap concentrator (Labconco), and injected onto a homemade 30 cm C18 column with 1.8 μm beads (Sepax) on an Easy nLC-1000 HPLC (ThermoFisher Scientific) connected to a Q Exactive Plus mass spectrometer (ThermoFisher Scientific) for MS. Raw data from MS experiments were mapped against SwissProt human database using the SEQUEST search engine within the Proteome Discoverer software platform, version 2.2 (ThermoFisher Scientific). The Minora node was used to determine relative protein abundance using the default settings. Percolator was used as the FDR calculator and peptides with q-value > 0.1 were filtered

out. For calculating SAINT probability, biological replicates and two IPs using rabbit pre-immune IgG (negative control) were included. Calculation was performed by the CRAPome web tool (63) according to the user guide.

Western blotting

Cells were lysed in RIPA buffer (Thermo Fisher Scientific), and total cell proteins were separated in a 10% denaturing polyacrylamide gel, transferred to polyvinylidene difluoride membranes (PVDF; Amersham Biosciences), probed using primary antibodies, and incubated with either a mouse or rabbit secondary antibody conjugated with horseradish peroxidase (GE Biosciences). For quantitative western blotting, protein abundance was quantified by Pierce BCA Protein Assay kit (Thermo, UL294765) according to the manufacturer's suggestions. The same protein mass was loaded, and β-actin was used as an internal control, if applicable.

RNA-binding protein immunoprecipitation (RIP)

Cells from one 80% confluent 10-cm culture dish were harvested in native lysis buffer (300 μL) with 50 mM HEPES, pH 7.5, 150 mM KCl, 2 mM EDTA, 1 mM NaF, 0.1 % (v/v) Nonidet P-40, 1 mM DTT, and RNasin RNase inhibitor (Promega). Immunoprecipitation of endogenous PRRC2B was performed using 1 μg (dilution factor 1:125) rabbit anti-human PRRC2B antibody (Invitrogen) followed by Magnetic dynabead protein A (Invitrogen) pull-down. Bound RNA was extracted from agarose following the same Trizol (Qiagen) extraction protocol, as mentioned previously.

Dual-luciferase reporter assay

Control or PRRC2B knockdown HEK293T cells were grown in 96-well plates until 80% confluency. The cells were then transfected with an equal amount of experimental FLuc reporter plasmid and a control renilla luciferase (RLuc) plasmid for 18 hours. The cells were then incubated with Dual-Glo luciferase substrate (Promega) according to the manufacturer's recommendations. The final readings of the FLuc were normalized to RLuc to obtain the relative luminescence reading.

Cellular phenotype

For cell proliferation, cells were seeded at 500/well in 96-well plates as biological triplicates. MTT assay was performed using MTT Cell Proliferation Kit I (Sigma-Aldrich) according to the manufacturer's recommendations every 24 hours until day 7. Cell-doubling time was calculated for cells in the log phase of the growth curve. For the cell cycle, flow cytometry was performed with Propidium iodide (PI) stained cells according to the standardized protocol (64). Biological triplicates were performed for flow cytometry. G0/G1, S, and G2/M peaks were autodetected and quantified by FlowJo, and percentages of total cells detected by flow cytometry were reported.

Statistics

All quantitative data were presented as mean ± SD and analyzed using Excel (Microsoft Office). Statistical analyses were performed using the two-tailed Student's *t* test with $P < 0.05$ considered significant.

Figure 1. Characterization of PRRC2B-mRNA interaction through PAR-CLIP of full-length PRRC2B and its RNA-binding fragment. (A) A sketch of the workflow of PAR-CLIP. **(B)** Schematic showing the fragmentation of PRRC2B protein. The putative RNA-binding arginine-glycine (RG) rich sequences were covered in one fragment, P2. **(C)** An autoradiographic image showing the migration of [32][P]-radiolabeled crosslinked RNA-protein complexes enriched in immunoprecipitation of FLAG-tagged full-length PRRC2B

and fragments (P1, P2, P3) on a 4-12% Bis-Tris gel. (**D**) Autoradiogram of ^{32}[P]-radiolabeled size marker (19 and 35 nt) and RNase T1-digested RNA fragments crosslinked with full-length PRRC2B and P2 on 15% polyacrylamide gel. Results are representative of three and four independent experiments in (C, D). (**E**) A Venn diagram showing the 462 overlapped peaks (binding sites) on RNAs enriched in full-length PRRC2B and P2 immunoprecipitation. Peaks having at least 1 nt overlapping were reported as overlapped peaks. (**F**) Distribution of the 462 overlapped binding sites on different RNA species and intergenic regions. Binding sites on exons and introns were included. (**G**) A histogram showing the density of binding sites (number of binding sites per 1000 nt) on 5' untranslated region (UTR), coding sequence (CDS), 3' UTR, and intron regions of mRNA. Only the 326 overlapped binding sites on mRNAs were considered. (**H**) Distribution of the 326 overlapped binding sites on mature mRNAs. Only the longest transcript was included for each gene. All transcripts were scaled to the same length with the start and stop codons of the main open reading frames aligned. Kernel density estimation (KDE) was plotted as Y-axis. (**I**) The longest significant consensus motifs that MEME can identify from the sequences flanking the T-to-C mutation sites (-20 to +20 nt) in the 326 overlapped binding sites on mRNAs. Significance was tested against randomized sequences according to the MEME user guide. E-value represents an estimate of the expected number of motifs found in a similarly sized set of random sequences (32). E-value < 0.05 was considered statistically significant.

Figure 2. PRRC2B bound mRNAs exhibit decreased translation efficiency upon PRRC2B knockdown. (**A**) Schemes of the polysome profiling assay. Cell lysates were fractionated into 10 fractions to extract RNA for RT-qPCR. RNA sequencing was performed on the combined polysome fractions 6-10 after polysome profiling. (**B**) Schematic of the Doxycycline-inducible shRNA-mediated PRRC2B knockdown by the "Tet-on" system. TRE, tetracycline response element in Tet operator fused promoter; rtTA, reverse tetracycline-controlled trans-activator; DOX, Doxycycline; shPRRC2B, shRNA targeting the coding region of *PRRC2B* mRNA. (**C**) A western blot result showing the Doxycycline-inducible knockdown of PRRC2B. shCtrl, a HEK293T cell line expressing a non-silencing control shRNA upon DOX induction; shPRRC2B, a HEK293T cell line expressing shPRRC2B upon DOX induction. (**D**) A scatterplot showing the $\log_2$ transformed translation efficiency (TE) values of control cells (shCtrl $\log_2$ (TE)) and PRRC2B knockdown cells (shPRRC2B $\log_2$ (TE)) for all mRNAs detected in polysome-seq (sequencing of pooled polysome-associated RNA) 24 hours after DOX induction. (**E**) A scatterplot showing the $\log_2$ transformed translation efficiency values of control cells (shCtrl $\log_2$ (TE)) and $\log_2$ transformed translation efficiency values of PRRC2B knockdown cells (shPRRC2B $\log_2$ (TE)) for all PRRC2B bound mRNAs (mRNAs with overlapped binding sites identified in Figure 1E) 24 hours after DOX induction. (**F, G**) mRNA abundance distribution of exemplary TE-down PRRC2B-bound mRNAs (*CRKL*, *CCND2*, *YWHAZ*), TE-unchanged PRRC2B-unbound mRNAs (*GAPDH*, *GATA4*), and TE-unchanged PRRC2B-bound mRNAs (*ACTB*) in different polysome profiling fractions (1-10) at 0, 6, 12 hours after induction of PRRC2B knockdown. The distribution was measured by RT-qPCR and represented as a percentage of total RNA. (**H**) Western blot results of the protein abundance of TE-down PRRC2B-bound mRNAs (*CRKL*, *YWHAZ*, *CCND2*) at 0, 6, and 12 hours after induction of PRRC2B knockdown. (**I**) Gene body coverages of TE-down PRRC2B-bound mRNAs (*CRKL*, *YWHAZ*, *CCND2*) in PAR-CLIP of full-length PRRC2B, total RNA-seq of control cells, and total RNA-seq of PRRC2B knockdown cells. Biological duplicates were performed for all RT-qPCR in (F, G), and representative data was shown. Western blot results are representative of >5 independent experiments in (C) and two replicates in (H). β-actin was used as an internal control in (C, H).

Figure 3. PRRC2B binding to the 5' UTR of *CCND2* mRNA increases protein expression. (**A**) Schematic of the design of an antisense oligonucleotide (ASO) base-pairing with the PRRC2B binding site on the 5' UTR of *CCND2* mRNA (ASO1) and the design of an ASO base-pairing with an adjacent region (ASO2). (**B**) Fold enrichment (RIP versus input) of *CCND2* mRNA in RNA-binding protein immunoprecipitation (RIP) of endogenous PRRC2B in cells treated with control ASO having no targets in human cells (Ctrl ASO), ASO1, and ASO2. (**C**) Relative mRNA abundance (normalized by 18S rRNA) of *CCND2* mRNA detected by RT-qPCR in cells treated with control ASO (Ctrl ASO), ASO1, and ASO2. (**D**) Western blot analysis of the protein abundance of CCND2 in cells treated with Ctrl ASO, ASO1, and ASO2. β-actin is used as an internal control. Western blot results are representative of 3 independent experiments. (**E**) Schematic of the construction of firefly luciferase reporters with wild-type 5' UTR of *CCND2* mRNA (WT-FLuc), 5' UTR of *CCND2* mRNA with PRRC2B binding sites replaced by an adjacent sequence showing less PRRC2B binding (Del-FLuc), 5' UTR of *CCND2* mRNA with all the "C" in PRRC2B binding sites mutated to "G" (C2G-FLuc), *CCND2* 5' UTR with all the "U" in PRRC2B binding sites mutated to "A" (T2A-FLuc), and *CCND2* 5' UTR with all the

"C & U" in PRRC2B binding sites mutated to "G & A" (CT2GA-FLuc). (**F**) Fold enrichment (RIP versus input) of reporter mRNAs in RIP of endogenous PRRC2B. (**G**) Relative luciferase activity of all reporters. (**H**) Relative mRNA abundance (normalized by 18S rRNA) of reporter mRNAs detected by RT-qPCR. For all the histograms in this figure, biological triplicates were performed. Data were expressed as mean ± SD, and the Student *t* test was used to calculate the significance. NS, not significant; * *P* < 0.05; ** *P* < 0.01; *** *P* < 0.001.

Figure 4. PRRC2B knockdown reduces cell proliferation and G1/S transition. (**A**) Cell proliferation curves from day 1 to day 7 of control cells (shCtrl) and PRRC2B knockdown cells (shPRRC2B) were measured by MTT assay. (**B**) Doubling time was calculated by fitting the cell proliferation data (A) to exponential functions. (**C**) Representative flow cytometry images of control cells (left, shCtrl) and PRRC2B knockdown cells (right, shPRRC2B). The experiments were performed 24 hours after induction of PRRC2B knockdown. (**D**) The quantification of relative cell number across different cell cycle stages in (C). (**E**) Representative flow cytometry images of cells treated

with control ASO (left) and ASO1 reducing the binding of PRRC2B to *CCND2* mRNA (right). (**F**) The quantification of relative cell number across different cell cycle stages in (E). For all flow cytometry results, biological triplicates were performed, G0/G1, S, and G2/M peaks were autodetected and quantified by FlowJo, and percentages of total cells detected by flow cytometry were reported. Biological triplicates were performed for all experiments. Data were expressed as mean $\pm$ SD in A, B, D, and F, and the Student *t* test was used to calculate the significance. NS, not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Figure 5. PRRC2B interacts with translation initiation factors eIF4G2 and eIF3. (**A**) Schematic of PRRC2B interactome capture workflow with or without RNase T1 digestion in HEK293T cells. (**B**) SAINT probabilities and the Log$_2$ transformed fold enrichments (Log$_2$FC) of all PRRC2B interacting partners identified in the PRRC2B interactome capture assay without RNase T1 treatment. Biological duplicates were performed. SAINT probability > 0.95 is considered high confidence. (**C**) Western blot analysis of the co-immunoprecipitation of endogenous PRRC2B together with translation initiation factors

(eIF4G2 and eIF3) and FXR1 without RNase T1 treatment. (**D**) Western bot detection of the distribution of PRRC2B and its interacting partners in different polysome profiling fractions. Protein was purified from an equal volume of each fraction, and an equal volume of the purified protein was loaded onto a 10% SDS-PAGE gel. Western blot results are representative of >2 independent experiments in (C, D).

Figure 6. Interacting with eIF4G2/eIF3 is essential for PRRC2B-mediated translation of target mRNAs. (**A**) Western blot measurement of changes in the abundance of proteins encoded by PRRC2B-bound mRNAs (*CCND2*, *YWHAZ*, *CRKL*) in control cells (shCtrl), PRRC2B knockdown cells (shPRRC2B), PRRC2B-overexpressing cells (PRRC2B OE), and PRRC2B-overexpressing cells with eIF4G2 knockdown (PRRC2B OE siEIF4G2). (**B**) Schematic of the construction of FLAG-tagged truncated PRRC2B with eIF4G2/eIF3 binding region removed (del450). (**C**) Western blot detection of the Co-IP of FLAG-tagged full-length and truncated PRRC2B (del450) together with translation initiation factors (eIF4G2 and eIF3) upon RNase T1 treatment (5U/μl). (**D**) Western blot

analysis of the protein abundance changes of CCND2, YWHAZ, and CRKL in control cells overexpressing FLAG tag (shCtrl KD + FLAG OE), PRRC2B knockdown cells overexpressing FLAG tag (shPRRC2B + FLAG OE), PRRC2B knockdown cells overexpressing FLAG-tagged PRRC2B (shPRRC2B + FL-FLAG OE), and PRRC2B knockdown cells overexpressing FLAG-tagged truncated PRRC2B (shPRRC2B + del450 OE). (**E**) Cell proliferation curves from day 1 to day 7 of cells under various conditions. (**F**) Doubling time was calculated by fitting the cell proliferation data (E) to exponential functions. (**G**) The quantification of relative cell number distribution across different cell cycle stages under various conditions. (**H**) Schematic model showing PRRC2B-bearing translation initiation complexes on mRNA. Biological triplicates were performed for all the experiments, and representative images were shown in (A, C, D). Flow cytometry results were analyzed and quantified by FlowJo. β-actin is used as an internal control for western blot in (A, D). Western blot results are representative of 3 independent experiments in (A, C, D). PRRC2B knockdown experiments were performed 24 hours after induction of gene knockdown in (A, D, G). Data were expressed as mean $\pm$ SD in (E, F, G), and the Student t test was used to calculate the significance. NS, not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Data availability

Data supporting the findings of this manuscript are available from the corresponding author upon request. The sequencing data raw files and processed files were deposited at GEO database and are accessible under accession numbers GSE220057, GSE220058, and GSE220059.

Acknowledgments

We are grateful for the consulting contributions from Elizabeth Grayhack, Joshua Munger, Yitao Yu, and Chen Yan. We appreciate Jared Hollinger and Huan Liu for their technical support and Emily Bonanno for her polishing the manuscript. This work was supported in part by the National Institutes of Health (R01 HL132899 and R01 HL147954 to P.Y.), NIH T32 Fellowship (T32 GM068411 to O.H.), American Heart Association Career Development Award (848985 to J.W.), and start-up funds from Aab Cardiovascular Research Institute of University of Rochester Medical Center (to P.Y.).

Author contributions

PY obtained the funding and launched the study. PY, JW, and FJ conceived the ideas. PY and FJ designed the experiments, analyzed the data, and wrote the manuscript. FJ carried out the experimental work and bioinformatic analysis. OH, ESK, and MA provided conceptual feedback or technical assistance. All the authors discussed the results and had the opportunity to comment on the manuscript.

Conflict of interest

None of the authors have any financial conflict of interest related to the research described in this manuscript.

Reference

1. ZJ, L. (2012) *RNA binding proteins*. Austin, Tex.: Landes Bioscience.
2. Hentze, M.W., Castello, A., Schwarzl, T. and Preiss, T. (2018) A brave new world of RNA-binding proteins. *Nat Rev Mol Cell Biol*, **19**, 327-341.
3. Lunde, B.M., Moore, C. and Varani, G. (2007) RNA-binding proteins: modular design for efficient function. *Nat Rev Mol Cell Biol*, **8**, 479-490.
4. Allain*., A.C.a.F.H.-T. (2011) *From structure to function of RNA binding domains In RNA Binding Proteins*, Zdravko L, ed. Landes Bioscience Springer Science+Business Media.
5. Van Nostrand, E.L., Freese, P., Pratt, G.A., Wang, X., Wei, X., Xiao, R., Blue, S.M., Chen, J.-Y., Cody, N.A.L., Dominguez, D. *et al.* (2020) A large-scale binding and functional map of human RNA-binding proteins. *Nature*, **583**, 711-719.
6. Glisovic, T., Bachorik, J.L., Yong, J. and Dreyfuss, G. (2008) RNA-binding proteins and post-transcriptional gene regulation. *FEBS Lett*, **582**, 1977-1986.
7. Mitchell, S.F. and Parker, R. (2014) Principles and Properties of Eukaryotic mRNPs. *Molecular Cell*, **54**, 547-558.
8. Gerstberger, S., Hafner, M. and Tuschl, T. (2014) A census of human RNA-binding proteins. *Nat Rev Genet*, **15**, 829-845.
9. Gebauer, F., Schwarzl, T., Valcárcel, J. and Hentze, M.W. (2021) RNA-binding proteins in human genetic disease. *Nature Reviews Genetics*, **22**, 185-198.
10. Harvey, R.F., Smith, T.S., Mulroney, T., Queiroz, R.M.L., Pizzinga, M., Dezi, V., Villenueva, E., Ramakrishna, M., Lilley, K.S. and Willis, A.E. (2018) Trans-acting translational regulatory RNA binding proteins. *WIREs RNA*, **9**, e1465.
11. Cockman, E., Anderson, P. and Ivanov, P. (2020) TOP mRNPs: Molecular Mechanisms and Principles of Regulation. *Biomolecules*, **10**.
12. Leppek, K., Das, R. and Barna, M. (2018) Functional 5' UTR mRNA structures in eukaryotic translation regulation and how to find them. *Nat Rev Mol Cell Biol*, **19**, 158-174.
13. Venkata Subbaiah, K.C., Hedaya, O., Wu, J., Jiang, F. and Yao, P. (2019) Mammalian RNA switches: Molecular rheostats in gene regulation, disease, and medicine. *Comput Struct Biotechnol J*, **17**, 1326-1338.
14. Gebauer, F. and Hentze, M.W. (2004) Molecular mechanisms of translational control. *Nat Rev Mol Cell Biol*, **5**, 827-835.
15. Castello, A., Fischer, B., Eichelbaum, K., Horos, R., Beckmann, Benedikt M., Strein, C., Davey, Norman E., Humphreys, David T., Preiss, T., Steinmetz, Lars M. *et al.* (2012) Insights into RNA Biology from an Atlas of Mammalian mRNA-Binding Proteins. *Cell*, **149**, 1393-1406.
16. Kastelic, N. and Landthaler, M. (2017) mRNA interactome capture in mammalian cells. *Methods*, **126**, 38-43.
17. Baltz, A.G., Munschauer, M., Schwanhausser, B., Vasile, A., Murakawa, Y., Schueler, M., Youngs, N., Penfold-Brown, D., Drew, K., Milek, M. *et al.* (2012) The mRNA-bound proteome and its global occupancy profile on protein-coding transcripts. *Mol Cell*, **46**, 674-690.
18. Castello, A., Horos, R., Strein, C., Fischer, B., Eichelbaum, K., Steinmetz, L.M., Krijgsveld, J. and Hentze, M.W. (2016) Comprehensive Identification of RNA-Binding Proteins by RNA Interactome Capture. *Methods Mol Biol*, **1358**, 131-139.

19. Chowdhury, M.N. and Jin, H. (2022) The RGG motif proteins: Interactions, functions, and regulations. *WIREs RNA*, **n/a**, e1748.
20. Thandapani, P., O'Connor, T.R., Bailey, Timothy L. and Richard, S. (2013) Defining the RGG/RG Motif. *Molecular Cell*, **50**, 613-623.
21. Stark, C., Breitkreutz, B.J., Reguly, T., Boucher, L., Breitkreutz, A. and Tyers, M. (2006) BioGRID: a general repository for interaction datasets. *Nucleic Acids Res*, **34**, D535-539.
22. de la Parra, C., Ernlund, A., Alard, A., Ruggles, K., Ueberheide, B. and Schneider, R.J. (2018) A widespread alternate form of cap-dependent mRNA translation initiation. *Nature Communications*, **9**, 3068.
23. Sugiyama, H., Takahashi, K., Yamamoto, T., Iwasaki, M., Narita, M., Nakamura, M., Rand, T.A., Nakagawa, M., Watanabe, A. and Yamanaka, S. (2017) Nat1 promotes translation of specific proteins that induce differentiation of mouse embryonic stem cells. *Proceedings of the National Academy of Sciences*, **114**, 340-345.
24. He, S., Yang, L., Xiao, Z., Tang, K. and Xu, D. (2021) Identification of key carcinogenic genes in Wilms' tumor. *Genes Genet Syst*, **96**, 141-149.
25. Tang, Z., Li, C., Kang, B., Gao, G., Li, C. and Zhang, Z. (2017) GEPIA: a web server for cancer and normal gene expression profiling and interactive analyses. *Nucleic Acids Research*, **45**, W98-W102.
26. Kewitz, S., Stiefel, M., Kramm, C.M. and Staege, M.S. (2014) Impact of O6-methylguanine-DNA methyltransferase (MGMT) promoter methylation and MGMT expression on dacarbazine resistance of Hodgkin's lymphoma cells. *Leuk Res*, **38**, 138-143.
27. Panagopoulos, I., Gorunova, L., Torkildsen, S., Tjonnfjord, G.E., Micci, F. and Heim, S. (2017) DEK-NUP214-Fusion Identified by RNA-Sequencing of an Acute Myeloid Leukemia with t(9;12)(q34;q15). *Cancer Genomics Proteomics*, **14**, 437-443.
28. Tsurubuchi, T., Nakano, Y., Hirato, J., Yoshida, A., Muroi, A., Sakamoto, N., Alexander, Z., Matsuda, M., Ishikawa, E., Kohno, T. *et al.* (2019) Subependymal giant cell astrocytoma harboring a PRRC2B-ALK fusion: A case report. *Pediatr Blood Cancer*, **66**, e27995.
29. Gupta, A., Liu, H., Schieffer, K.M., Koo, S.C., Cottrell, C.E., Mardis, E.R., Roberts, R.D. and Yeager, N.D. (2021) Targeted Therapy in a Young Adult With a Novel Epithelioid Tumor Driven by a PRRC2B-ALK Fusion. *J Natl Compr Canc Netw*, **19**, 1116-1121.
30. Wang, Z., Geng, Y., Yuan, L.Y., Wang, M.M., Ye, C.Y., Sun, L., Dai, W.P. and Zang, Y.S. (2022) Durable Clinical Response to ALK Tyrosine Kinase Inhibitors in Epithelioid Inflammatory Myofibroblastic Sarcoma Harboring PRRC2B-ALK Rearrangement: A Case Report. *Front Oncol*, **12**, 761558.
31. Danan, C., Manickavel, S. and Hafner, M. (2016) PAR-CLIP: A Method for Transcriptome-Wide Identification of RNA Binding Protein Interaction Sites. *Methods Mol Biol*, **1358**, 153-173.
32. Bailey, T.L., Johnson, J., Grant, C.E. and Noble, W.S. (2015) The MEME Suite. *Nucleic Acids Research*, **43**, W39-W49.

33. Chassé, H., Boulben, S., Costache, V., Cormier, P. and Morales, J. (2017) Analysis of translation using polysome profiling. *Nucleic Acids Research*, **45**, e15-e15.

34. Wu, J., Subbaiah, K.C.V., Xie, L.H., Jiang, F., Khor, E.S., Mickelsen, D., Myers, J.R., Tang, W.H.W. and Yao, P. (2020) Glutamyl-Prolyl-tRNA Synthetase Regulates Proline-Rich Pro-Fibrotic Protein Synthesis During Cardiac Fibrosis. *Circ Res*, **127**, 827-846.

35. Koyama-Nasu, R., Nasu-Nishimura, Y., Todo, T., Ino, Y., Saito, N., Aburatani, H., Funato, K., Echizen, K., Sugano, H., Haruta, R. *et al.* (2013) The critical role of cyclin D2 in cell cycle progression and tumorigenicity of glioblastoma stem cells. *Oncogene*, **32**, 3840-3845.

36. Ando, K., Ajchenbaum-Cymbalista, F. and Griffin, J.D. (1993) Regulation of G1/S transition by cyclins D2 and D3 in hematopoietic cells. *Proc Natl Acad Sci U S A*, **90**, 9571-9575.

37. Choi, H., Liu, G., Mellacheruvu, D., Tyers, M., Gingras, A.C. and Nesvizhskii, A.I. (2012) Analyzing protein-protein interactions from affinity purification-mass spectrometry data with SAINT. *Curr Protoc Bioinformatics*, **Chapter 8**, Unit8.15.

38. George, J., Li, Y., Kadamberi, I.P., Parashar, D., Tsaih, S.W., Gupta, P., Geethadevi, A., Chen, C., Ghosh, C., Sun, Y. *et al.* (2021) RNA-binding protein FXR1 drives cMYC translation by recruiting eIF4F complex to the translation start site. *Cell Rep*, **37**, 109934.

39. Weber, R., Kleemann, L., Hirschberg, I., Chung, M.-Y., Valkov, E. and Igreja, C. (2021) DAP5 enables translation re-initiation on structured messenger RNAs. *bioRxiv*, 2021.2001.2021.427569.

40. Lee, A.S., Kranzusch, P.J., Doudna, J.A. and Cate, J.H. (2016) eIF3d is an mRNA cap-binding protein that is required for specialized translation initiation. *Nature*, **536**, 96-99.

41. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A. *et al.* (2021) Highly accurate protein structure prediction with AlphaFold. *Nature*, **596**, 583-589.

42. Fu, L., Dong, Q., Xie, C., Wang, Y. and Li, Q. (2015) CRKL protein overexpression enhances cell proliferation and invasion in pancreatic cancer. *Tumour Biol*, **36**, 1015-1022.

43. Kim, Y.H., Kwei, K.A., Girard, L., Salari, K., Kao, J., Pacyna-Gengelbach, M., Wang, P., Hernandez-Boussard, T., Gazdar, A.F., Petersen, I. *et al.* (2010) Genomic and functional analysis identifies CRKL as an oncogene amplified in lung cancer. *Oncogene*, **29**, 1421-1430.

44. Cai, L., Wang, H. and Yang, Q. (2017) CRKL overexpression promotes cell proliferation and inhibits apoptosis in endometrial carcinoma. *Oncol Lett*, **13**, 51-56.

45. Sherr, C.J. and Roberts, J.M. (2004) Living with or without cyclins and cyclin-dependent kinases. *Genes Dev*, **18**, 2699-2711.

46. Stonyte, V., Boye, E. and Grallert, B. (2018) Regulation of global translation during the cell cycle. *Journal of Cell Science*, **131**.

47. Bohlen, J., Roiuk, M. and Teleman, A.A. (2022) PRRC2 proteins regulate translation initiation by promoting leaky scanning. *bioRxiv*, 2022.2011.2011.516176.

48. Wu, R., Li, A., Sun, B., Sun, J.-G., Zhang, J., Zhang, T., Chen, Y., Xiao, Y., Gao, Y., Zhang, Q. *et al.* (2019) A novel m6A reader Prrc2a controls oligodendroglial specification and myelination. *Cell Research*, **29**, 23-41.

49. Youn, J.Y., Dunham, W.H., Hong, S.J., Knight, J.D.R., Bashkurov, M., Chen, G.I., Bagci, H., Rathod, B., MacLeod, G., Eng, S.W.M. *et al.* (2018) High-Density Proximity Mapping Reveals the Subcellular Organization of mRNA-Associated Granules and Bodies. *Mol Cell*, **69**, 517-532 e511.

50. Yang, P., Mathieu, C., Kolaitis, R.M., Zhang, P., Messing, J., Yurtsever, U., Yang, Z., Wu, J., Li, Y., Pan, Q. *et al.* (2020) G3BP1 Is a Tunable Switch that Triggers Phase Separation to Assemble Stress Granules. *Cell*, **181**, 325-345 e328.

51. Jin, S.C., Homsy, J., Zaidi, S., Lu, Q., Morton, S., DePalma, S.R., Zeng, X., Qi, H., Chang, W., Sierant, M.C. *et al.* (2017) Contribution of rare inherited and de novo variants in 2,871 congenital heart disease probands. *Nat Genet*, **49**, 1593-1601.

52. Dickinson, M.E., Flenniken, A.M., Ji, X., Teboul, L., Wong, M.D., White, J.K., Meehan, T.F., Weninger, W.J., Westerberg, H., Adissu, H. *et al.* (2016) High-throughput discovery of novel developmental phenotypes. *Nature*, **537**, 508-514.

53. Shah, A., Qian, Y., Weyn-Vanhentenryck, S.M. and Zhang, C. (2017) CLIP Tool Kit (CTK): a flexible and robust pipeline to analyze CLIP sequencing data. *Bioinformatics*, **33**, 566-567.

54. Martin, M. (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal; Vol 17, No 1: Next Generation Sequencing Data Analysis*.

55. Langmead, B., Trapnell, C., Pop, M. and Salzberg, S.L. (2009) Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biology*, **10**, R25.

56. Bolger, A.M., Lohse, M. and Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, **30**, 2114-2120.

57. Kim, D., Paggi, J.M., Park, C., Bennett, C. and Salzberg, S.L. (2019) Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol*, **37**, 907-915.

58. Liao, Y., Smyth, G.K. and Shi, W. (2014) featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, **30**, 923-930.

59. Frankish, A., Diekhans, M., Jungreis, I., Lagarde, J., Loveland, J.E., Mudge, J.M., Sisu, C., Wright, J.C., Armstrong, J., Barnes, I. *et al.* (2021) GENCODE 2021. *Nucleic Acids Res*, **49**, D916-d923.

60. Love, M.I., Huber, W. and Anders, S. (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, **15**, 550.

61. Team, R.C.R. (2020) A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. . *Vienna, Austria. URL https://www.R-project.org/. (2020).*

62.   Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L. *et al.* (2021) clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation (Camb)*, **2**, 100141.

63.   Mellacheruvu, D., Wright, Z., Couzens, A.L., Lambert, J.P., St-Denis, N.A., Li, T., Miteva, Y.V., Hauri, S., Sardiu, M.E., Low, T.Y. *et al.* (2013) The CRAPome: a contaminant repository for affinity purification-mass spectrometry data. *Nat Methods*, **10**, 730-736.

64.   Krishan, A. (1975) Rapid flow cytofluorometric analysis of mammalian cell cycle by propidium iodide staining. *J Cell Biol*, **66**, 188-193.

Supplementary Information for

RNA binding protein PRRC2B mediates translation of specific proteins and regulates cell cycle progression

Feng Jiang, Omar M. Hedaya, EngSoon Khor, Jiangbin Wu, Matthew Auguste, Peng Yao

Correspondence to Peng Yao

E-mail: peng_yao@urmc.rochester.edu

This document file includes:

Figures S1 to S6

Tables S1 to S7

Extended Data 1 and 2

SI Materials and Methods

Supplemental figures

Figure S1. PAR-CLIP analysis of PRRC2B-RNA interaction. (**A**) An autoradiographic image showing the migration of ³²[P]-radiolabeled crosslinked RNA-protein complexes (with 4-thiouridine, 4sU) and non-crosslinked RNAs (without 4sU) enriched in the

immunoprecipitation step of PAR-CLIP on a 4-12% Bis-Tris gel. (**B**) Western blot analysis shows the enrichment of comparable protein amounts of FLAG-tagged full-length PRRC2B and fragments (P1, P2, P3) by anti-FLAG antibody-conjugated magnetic beads in PAR-CLIP. 2% of the total volume of enriched proteins was loaded on a 10% SDS-PAGE gel. (**C**) Distribution of the length of sequenced reads after trimming and quality filtering of the two biological replicates in full-length PRRC2B (left) and P2 PAR-CLIP (right). (**D**) Strong correlations (Pearson's correlation) between the two biological replicates in full-length PRRC2B (left) and P2 PAR-CLIP (right), as indicated by scatterplots. The coordinates of each dot represent the FPKM (fragments per kilobase of exon per million mapped fragments) values of each gene in the two biological replicates. (**E**) Distribution of the binding sites identified in full-length PRRC2B (left) and P2 PAR-CLIP (right) on different RNA species and intergenic regions. Binding sites on exons and introns were included. (**F**) Histograms showing the density of binding sites (number of binding sites per 1000 nt) on 5' UTR, CDS, 3' UTR, and intron regions of mRNA in full-length PRRC2B (left) and P2 PAR-CLIP (right). Only the binding sites on mRNAs were considered. (**G**) Subcellular localization of endogenous and FLAG-tagged PRRC2B in HEK293T cells detected by immunofluorescent staining. (**H**) Distribution of binding sites identified in full-length PRRC2B (left) and P2 PAR-CLIP (right) on mature mRNAs. Only the longest transcript was included for each gene. All transcripts were scaled to the same length with start and stop codons of the main open reading frames aligned. Kernel density estimation (KDE) was plotted as Y-axis. (**I**) The longest significant consensus motifs that MEME can identify from the sequences flanking the T-to-C mutation sites (-20 to +20 nt) in the overlapped binding sites on 5' UTR, CDS, and 3' UTR. Significance was tested against randomized sequences according to the MEME user guide. E-value represents an estimate of the expected number of motifs found in a similarly sized set of random sequences (32). E-value < 0.05 was considered statistically significant.

Figure S2. PRRC2B-bound mRNAs exhibit decreased translation efficiencies upon PRRC2B knockdown. (**A-D**) Top enriched gene ontology (GO) biological processes in all TE-up mRNAs, TE-down mRNAs, TE-up PRRC2B-bound mRNAs, and TE-down PRRC2B-bound mRNAs. GO terms were filtered by adjusted *P* < 0.05 and simplified by the R package 'ClusterProfiler' to remove redundant terms. (**E**) RNA-binding protein immunoprecipitation (RIP) of endogenous PRRC2B followed by RT-qPCR for PRRC2B-bound and unbound mRNAs. Input mRNA was used as a normalizer. Results are presented as mean $\pm$ SD. (**F, G, I**) mRNA abundance distribution (detected by RT-qPCR and represented as a percentage of total RNA) of exemplary TE-down PRRC2B-bound mRNAs (*CTC1*, *ATAD5*, *SERBP1*, *BPTF*, *PPP2R1A*), TE-down unbound mRNAs (*PTBP1*, *HSPA4*), TE-up unbound mRNAs (*RPL37A*), and TE-up bound mRNAs (*RPL35*, *RPLP1*, *RPL38*) in different polysome profiling fractions at 0, 6, 12 hours after induction of PRRC2B knockdown. Biological duplicates were performed for RT-qPCR experiments included in (F, G, I), and representative data were shown. (**H**) Gene body coverages (exhibited as FPKM values) of exemplary TE-down PRRC2B-bound mRNAs (*CTC1*, *ATAD5*, *SERBP1*, *BPTF*, *PPP2R1A*) in PAR-CLIP of full-length PRRC2B, total RNA-seq of control cells, and total RNA-seq of PRRC2B knockdown cells.

Figure S3. PRRC2B knockdown cells are insensitive to ASOs and express the mutant luciferase reporters differently from wild-type cells. (**A**) Western blot analysis of the protein abundance of CCND2 in PRRC2B knockdown cells treated with Ctrl ASO, ASO1, and ASO2. (**B**) RT-qPCR measurement of *CCND2* mRNA in PRRC2B knockdown cells treated with Ctrl ASO, ASO1, and ASO2. (**C**) A snapshot showing the sequences of the PRRC2B binding region on WT-FLuc and corresponding mutated sequences in Del-FLuc, C2G-FLuc, T2A-FLuc, and CT2GA-FLuc. (**D**) Relative luciferase activity of wildtype and four mutant reporters in PRRC2B knockdown cells. For all assays involving PRRC2B knockdown, experiments were performed 24 hours after DOX induction. Western blots were representative of >2 biological replicates. Data were expressed as mean ± SD in (B, D), and the Student *t* test was used to calculate the significance. NS, not significant; * *P* < 0.05; ** *P* < 0.01.

Figure S4. Cell proliferation or G1/S transition does not change without induction of PRRC2B knockdown. (**A**) Cell proliferation curves from day 1 to day 7 of control cells and PRRC2B knockdown cells without induction of PRRC2B knockdown (no DOX treatment) measured by MTT assay. (**B**) Doubling time was calculated by fitting the cell proliferation data (A) to exponential functions. (**C**) Representative flow cytometry images of control (left) and PRRC2B knockdown cells (right) without DOX induction. (**D**) The quantification of relative cell number across different cell cycle stages in (C). For all the line plots and histograms, biological triplicates were performed. For all flow cytometry results, biological triplicates were performed, G0/G1, S, and G2/M peaks were autodetected and quantified by FlowJo, and percentages of total cells detected by flow cytometry were reported. Data were expressed as mean ± SD, and the Student *t* test was used to calculate the significance. NS, not significant.

Figure S5. Co-immunoprecipitation of PRRC2B with translation initiation factors.
(**A**) Western blot detection of the co-immunoprecipitation (Co-IP) of FLAG-tagged

PRRC2B together with translation initiation factors (eIF4G2 and eIF3) and FXR1 without RNase T1 treatment. (**B**) SYBR gold staining shows the fragmentation of RNA in total cell lysates by 1 U/μl, 5 U/μl, and 10 U/μl RNase T1. (**C**) Western blot detection of the Co-IP of endogenous PRRC2B together with translation initiation factors and FXR1 upon 5 U/μl RNase T1 treatment. (**D**) Western blot detection of the Co-IP of FLAG-tagged PRRC2B fragments (P1, P2, P3) together with translation initiation factors and FXR1 upon 5 U/μl RNase T1 treatment. (**E**) Upper: Schematic of the fragmentation of the P1 region of PRRC2B protein. Truncated P1 (T1-T5) was made by removing a series of 150 amino acids from the C-terminus of P1. Lower: Western blot detection of the Co-IP of FLAG-tagged truncated P1 fragments (T1-T5) together with translation initiation factors upon 5U/μl RNase T1 treatment. (**F**) Schematic of the regions of PRRC2B protein interacting with eIF3, eIF4G2, FXR1, and RNA. Western blot results are representative of >2 independent experiments in (A, C, D, E).

Figure S6. eIF4G2/eIF3-PRRC2B interaction is essential for translation of PRRC2B mRNA targets. (**A**) Western blot detection of the abundance changes of proteins (CCND2, YWHAZ, CRKL) in cells transfected with control siRNA with no target in human cells (siCtrl) and cells transfected with siRNA targeting CDS of *EIF4G2* mRNA (siEIF4G2). (**B**) mRNA abundance distribution (detected by RT-qPCR and represented as a percentage of total RNA) of PRRC2B-bound mRNAs (*CCND2*, *YWHAZ*, *CRKL*) in different polysome profiling fractions from cells under different conditions. (**C**) Representative flow cytometry images of cells under different conditions. For all flow cytometry results, biological triplicates were performed. β-actin is used as an internal control for western blot. For all assays involving PRRC2B knockdown, experiments were performed 24 hours after induction of PRRC2B knockdown. Western blot results are representative of >2 biological replicates in (A).

Extended Data 1.

Sequence alignment of PRRC2B protein in five representative species was shown. PRRC2B gene is conserved in 460 species, including all kinds of mammals, reptiles, amphibians, and fish (e.g., zebrafish and shark) based on sequence information from NCBI.

```
zebrafish   MSDRLGQITKSKDGKSKYSSLSLFDKYKGKSIETQKTTVVARHGLQSLGKVAAARRMPPP    60
frog        MSDRLGQITKGKDGKSKYSSLNLFDKYKGKSIEAVRTTVIPRHGLQSLGKVAAARRMPPP    60
chicken     MSDRLGQITKGKDGKSKYSTLSLFDKYKGKSIEAIRTTVIPRHGLQSLGKVAAARRMPPP    60
human       MSDRLGQITKGKDGKSKYSTLSLFDKYKGKSVDAIRSSVIPRHGLQSLGKVAAARRMPPP    60
mouse       MSDRLGQITQGKDGKSKYSTLSLFDKYKGRSVGAVRSSVIPRHGLQSLGKVATARRMPPP    60
            *********:.*********:*.********:*: : :::*: *********** :******

zebrafish   AHLPSLKSESKGNDPNVIIVPKDGTGWANKQDQPDPKSSVASSAQLPEPQPPLALQKSVS   120
frog        ANLPSLKSENKGNDPNIFIVPKDGTGWANKQEQQDQKSSSVTAPPQQESLPPQGLQKSVS   120
chicken     ANLPSLKSENKGNDPNIVIVPKDGTGWANKQDQSDQKNSSATAAQLQESLLQQGLQKSVS   120
human       ANLPSLKSENKGNDPNIVIVPKDGTGWANKQDQQDPKSSSATASQPPESLPQPGLQKSVS   120
mouse       ANLPSLKSENKGNDPNIVIVPKDGTGWANKQDQQDPKSSSVTASQPPESQPQPGLQKSVS   120
            *:*******.******:.**************:* * *.* .::     *      .******

zebrafish   NLQKPMPITSPEST--STGGPKQWAQLNGKAVDQD-GLKVSSRLQPFSHEDFPTLKAAGE   177
frog        NLQKPTPSINQENINLVPGGPKSWAQLNGKPAGQEGGSRGSNRLLSFSPEEFPTLKAAGE   180
chicken     NLQKPTQSIAQESTNSVPGGPKSWAQLNGKPAGLEGGSRASSRLLSFSPEEFPTLKAAGE   180
human       NLQKPTQSISQENTNSVPGGPKSWAQLNGKPVGHEGGLRGSSRLLSFSPEEFPTLKAAGG   180
mouse       NLQKPTQSISQENTNSVPGGPKSWAQLSGKPVGHEGGLRGSSRLLSFSPEEFPTLKAAGG   180
            *****     *.    ****.****.** .. : * : *.**   ** *:********

zebrafish   QDKVGKERSVFDPSYGPGPSLRPQ--------NVSSWREGGGRNLQPPSLSSAPPGDTDT   229
frog        QDKAGKEKSVLDPSYGPGPSLRPQSKPLALLPDVTSWRDGGGRSIASSLSPTASPTEPGS   240
chicken     QDKVGKEKGALDLSYGPGPSLRPQ--------NVTSWREGGGRNITSATSLTASPAELGS   232
human       QDKAGKEKGVLDLSYGPGPSLRPQ--------NVTSWREGGGRHIISATSLTSTPTELGS   232
mouse       QDKAGKEKGALDLSYGPGPSLRPQ--------NVTSWREGGGRNIISAASLSASPTELGS   232
            ***.***:..:* ***********          :*:***:**** :    :: * : .:

zebrafish   KSSGSVEAPLPPPPSTSTSTSSSASPAASSSSTAAAAAPSSEVKEPSLRPAPPLTRRAAP   289
frog        KTAV-------------------------PGEGPPSTSLANDTKELSLRPAQPTRKGPAQ   275
chicken     KTSS-------------------------TGDGAPSSVSASDPKEPSLRPAQPVRKGASQ   267
human       RNSS-------------------------TGDGAPSSACTSDSKDPSLRPAQPVRKGASQ   267
mouse       RNAS-------------------------GADGAPSLACTSDSKEPSLRPAQPSRRGASQ   267
            :.:                          ..   :   :.: *: ***** *  :  :

zebrafish   PALQYQHHTTTTYHDMLPAFMCPKETREAPGSSDGPAAVVA--PVRFDSRPIFRPPFPAP   347
frog        --FMGNTYHPPTYHDMLPAFMCSQHPPEPSGTLDRTSFPTLNSQSRLEPRVPFRQ-----   328
chicken     --FMGNVYQPPTYHDMLPAFMCPQQPSETPAPLDRGSFPVP--QLRLEPRVPFRQ-----   318
human       --FMGNVYHPPTYHDMLPAFMCSPKSSENQGTVERGSFPLP--QLRLEPRVPFRQ-----   318
mouse       --FMGHGYQPPTYHDMLPAFMCSPQSSENQTTVERSSFPLP--QLRLEPRVPFRQ-----   318
            :  :  : :  **********   .  *     :  :       *:: *   **

zebrafish   EPVNGDVRREPRFPRAPPRPSSRPIRRPGDRAPRPAIINPDDLKDLDELDNDCEDGWAGL   407
frog        FQMNDNEGRESRFPIAPPRS-ARPPRPPAERPPRATIINAEDLKELDDLDNDAEDGWAGL   387
chicken     YQMNDQDGKENRLGLSR--P-ARPLRQLVERAPRPTIINAENLKGLDELDNDADDGWAGI   375
human       FQMNDQDGKENRLGLSR--P-LRPLRQLVERAPRPTIINAENLKGLDDLDADADDGWAGL   375
mouse       FQMNDQDGKE-RPGVAR--P-VRPLRQLVERAPRPTIINAENLKGLDDLDTDADDGWAGL   374
            :*.:  :* *   :       ** *   :* ** :*** ::** **:** *.:*****:
```

```
zebrafish   HEEVDYSEKLKFSDDEEDHAP--REKSKIWDDWDHHHHHHHLQGQHSSQSSGDGAFPQEP     465
frog        HDEVDYSEKLKFSEDEEEEDGAAKDGRSKWNGWDGKRQRQ---VSFNSTDSVEGKHPADE     444
chicken     HDEVDYSEKLKFSEDEEEE-SLKDGRQKWNSWDPRRQRQ---LSMSSADSADVKHTLEE     431
human       HEEVDYSEKLKFSDDEEEE-VVKDGRPKWNSWDPRRQRQ---LSMSSADSADAKRTREE     431
mouse       HEEVDYSEKLKFSDDEDED-VVKDGRSKWNNWDPRRQRA---LSLSSADSTDAKRTQEE     430
            *:***********:**::.     ::    *:.** ::::     . .* .* :      :

zebrafish   ED---------EPYLR-QQE--PPASRKTSTRFPAEPQQK-----ASVNSESAEEQDEPQR     509
frog        GKVWNDTLSHSRPARRTPEPAPQGPRKPNTWAAVADHQKPPAPAAVLRQPSI---EEKDE     501
chicken     GKNWSDSVGLSRSVRKAQDSQQ-PPRKINGWNSTSEYQKPS-QGSALRQQSL---EDKEE     486
human       GKDWAEAVGASRVVRKAPDPQP-PPRKLHGWAPGPDYQKSS-MGSMFRQQSI---EDKED     486
mouse       GKDWSGTAGGSRVIRKVPEPQP-PSRKLHSWASGPDYQKPT-MGSMFRQHSA---EDKED     485
             .          .     * :        **          **       : ... *     :: :

zebrafish   SAPPRGKFVSADV-SAVERARRRREEEERRAREERLAACAEKLKKLDEKFGKTEKPARSG     568
frog        KLPQRKNFVQSEISEAVERARKRREEEERRNREERLAACAAKLKQLDQKNKQALKPSADS     561
chicken     KVPLRQKFVHSEISEAVERARKRREEEERRAREERLAACAAKLKQLDQKCKLAQKSGE--     544
human       KPPPRQKFIQSEMSEAVERARKRREEEERRAREERLAACAAKLKQLDQKCKQARKAGE--     544
mouse       KPPPRQKFIQSEMSEAVERARKRREEEERRAREERLAACAAKLKQLDQKCRQAQKANE--     543
            . * * :*: ::: .******:******** ******** ***:**:*     : *

zebrafish   ----EADGKELAQSPARRSAKPQQDGWQYGSKEV----T---------DTPAESASSQDYK     612
frog        SRQLDNKENEDPRSPVSDRSNPQENAHTFRREYS----QESPSEYPEEEPSGT-------     610
chicken     --TQKHTENEDLRPPNTEKSAVQENGHVFRRATPEFHTQDVSGYLEEETPAPAAP-----     597
human       --ARKQAEKEVPWSPSAEKASPQENGPAVHKGSPEFPAQETPTTFPEEAPTVSPAV-AQS     601
mouse       --TPKPVEKEVPRSPGIEKVSPPENGPVVRKGSPEFPVQEAPTMFLEETPATSPTV-AQS     600
                      .   :*     *       ::.                    : *: :

zebrafish   DEGCNFHNNDDEDDDGGESTSPVPEYSRHQKPVPPRFQKQQ-----------QQQEQVYKL     662
frog        -----LKAETSSEEEYAEPTSPAQDFSKHQKLIPPRFQRQQ-----------QQQEQLYKM     655
chicken     ---------QSSSEDELREAPSPAQEFSKYQKSLPPRFQRQQ-----------QQEQLYKM     638
human       NSS---------EEEAREAGSPAQEF-KYQKSLPPRFQRQQQQQ---------QQQEQLYKM     643
mouse       NSSSSSSSSSSSIEEEVRESGSPAQEFSKYQKSLPPRFQRQQQQQQQQQQQQQQQQEQLYKM     660
                       ::: *  **. :: ::** :*****:**        ****:**:

zebrafish   ASWQQSAH--PAQPSSAHPQRGFYPPH--VLGFDPRWMMMPPYMDPRLAQGCSPVDYYPP     718
frog        QQWQQQQQQAYVASTHSNPPRTFYSPHPQMLGFDPRWMMMPSYMDPRMTQGRAPVDFYPS     715
chicken     QHWQQQQV----YPPPSHSRTFYPPHPQMLGFDPRWMMMPSYLDPRMAQSRTPVDFYPS     694
human       QHWQ--PV----YPPPSHPQRTFYPHHPQMLGFDPRWMMMPSYMDPRITPTRTPVDFYPS     697
mouse       QHWQ--PV----YPPPSHPQRTFYPHHPQMLGFDPRWMMMPSYMDPRITPTRTPVDFYPS     714
             **            ::  *  **  *  :*********** *:***::    :***:**

zebrafish   GVHSSGLVKPVIQPDHLNSPGST--SDEGCHPSMHQERRAPSTEPYQVWNQDSYPSARSF     776
frog        SIHPPGVMKHMIQQDSMSGSGSCHSDEQNCQSE----RRPQSAEPMSGWGQESYVQLHSK     771
chicken     ALHPSGIMKPMIQQDSIGGN-SCRSEDQNCQAG-QVERKAAPMDPVPVWGQESYASLQSK     752
human       ALHPSGLMKPMMPQESLNGT-GCRSEDQNCVPP-LQERKVTPIDSPPVWSPEGYMALQSK     755
mouse       ALHPSGLMKPMMPQESLSGT-GCRSEDQNCVPS-LQERKVTALDPAPVWSPEGYMALQNK     772
            .:*  *::* ::  : :..  .    .::.*     *:     :     *. :.*    :.

zebrafish   TPPYQ--RTHENGER-----------APADDRSDRSTSRHDSYEEHRHERTDSPAEETTH     823
frog        ACSLPQQKQAETAAV-GSYSRNESSYSSAREKSDGSPTHPEPLEREEEYLS-VSYEKKST     829
chicken     GYSLSHQKQSDNMTMEGLHARNGSSYSSSPGRPENLNTQRDLFEERGEEYL-NAFDKKTQ     811
human       GYPLPHPKSSDTLAM-DMRVRNESSFSASLGRAGGVSAQRDLFEERGEEYL-SAFDKKAQ     813
mouse       GYSLPHPKSADTLAM-GMHVRNERSYCASPGRPGGISAQRDLFEERGEEFL-SAFDKKAQ     830
               : :.        . : :     :: :  *.. .       :::.:
```

```
zebrafish   --------------------------QGFRQGR---VSDGSQREPMMLSRT---------      845
frog        GSFSSCISPQRRGQDALYQHLESGIESGVNRGRQMAARDSSSRGSEFIKTDKKPQFNGWG      889
chicken     GDFDSCLSSQRLGQDLLFQHQESVQETCPSSNRHANL-RCSPLEPDFIQAEKKPEYNGWD      870
human       ADFDSCISSQRIGQELLFPPQENVQDAGAPGGHTQNL-RCSPLEPDFVPDEKKPECGSWD      872
mouse       ADFDSCISSQRIGQELLFPPQENVQEVGAPGGCTPNL-RCSPLEPDFVPAEKKPEYGSWD      889
                                        .             *         ::

zebrafish   -------HDDAQHQKEEPYDSKEKSYEADFWRRDVRKE----SSGQTQWSDHGSSSSSSIS      895
frog        YAPHQKSSETASRAEEDLSKEDLSDELEHWKKEDGGSSQDG-TGETCWRSESANSN--SQ      946
chicken     ISHHQKPSEPAAEITEEVPRNEQSFAADPWKKEGASNTKQPAEETAEWAPENR-ST--GV      927
human       VSHQPETADTAHGVERETPREGTAFNISSWDKNGSPNKQP--SSEPEWTPEPR-SS--SS      927
mouse       VGHQPKAADTANGVELEAPRQEPSFHVSSWEKEGSPKKQP--NPEPEWTPEPR-SS--SG      944
                    :         :  :  . * ::  .          *  .   *.

zebrafish   QPSETSGRTLRRTGPIKKPVLKPLKVEDKENEKPKAEPEEKPV--PYRLEKEV--ITNVY      951
frog        QQSEHLGR-SRRSGPIKKPVLKALKIEDKELEKNKA--EIKEPVKPFKE---KI-LPKTE      999
chicken     QQQEQIGR-TRRSGPIKKPVLKALKVEEKEKEMEKVKLEGEDTSRHL-KE--KVTVQKVE      983
human       QHPEQTGR-TRRSGPIKKPVLKALKVEDKEKELEKIKQELGEESTRLAKEKEQSPTAEKD      986
mouse       QHQEQPGR-TRRSGPIKKPVLKALKVEEKEKALERGRQGLREESSQRAPEKEP--VGRAE     1001
            *    *  **   **:********* **:*:**      :                  .

zebrafish   DLKKDAPHVSNRHSSSVEDKQADALKMEKT--------SALSEELHKESCWEKSQSDCSD     1003
frog        CVVKTEPVVSSPPPSMIEVKTPPPPSPVQE---------TDKFPQEKAERSWETKPLR--E     1049
chicken     NESDDSAALLNSTRYLLDDKGSSQASLAREAEKSQEEEEEEEEDKPERTWENKVSR--E     1041
human       EDEENDASLANSSTTTLEDKGPGHATFGREATKF-------EEEEKPDKAWEARPPR--E     1037
mouse       EDEENNPALANA-SSALEDKAASRAGFAHEASKL-------DEDEKADKTWESRPSR--E     1051
                      .     : .     :: *          :          : : :  **          :

zebrafish   SRDPTAPRRNNWIFIDEEQAFAGA-RGRGRGRGFREFNSRGGGARGGRSDTNRG--AYNN     1060
frog        SSSTSLQKRSDWIFIDEEQAFGGRTQGRGRGRGFKDFGFRNRGVSG---------------     1095
chicken     SSELPPTKRNNWIFIDEEQAFGGRGQGRGRGRGFREFTFRGRG------------TVVSS     1089
human       SSDVPPMKRNNWIFIDEEQAFGVRGQARGRGRGFREFTFRG-RPAGGNGSGLCGGGVLGA     1096
mouse       ASDIPPTKRNNWIFIDEEQAFGGRGQARSRGRGFREFTFRGGRPAGSSTSGLCGTGVLGS     1111
            :  .      :*.:**********.   :.*.*****::*   *.

zebrafish   NNNNTGAQRLTRGRGNRVFR-GDDLQRGKPRRRNVSETHSETSEYEEQPKRPRQKVSENG     1119
frog        --TY-SGQKVSRGRGVREYNPPEDLRGKASRRRIASETHSEGSEYEELPKRRRQRGTENG     1152
chicken     RGVYN-NQRSSRGRGLREFNQPEDFPRGKPRRRIASETHSEGSEYEELPKRRRQRGSENS     1148
human       RSIYCSSQRSGRGRGLREFARPEDCPRAKPRRRVASETHSEGSEYEELPKRRRQRGSENG     1156
mouse       RGMYSSGQRSNRGRGLRDFPPPEDCPRAKPRRRIASETHSEGSEYEELPKRRRQRGSEHS     1171
                 *:    **** * :      :*        *** .****** ***** *** **: :*:.

zebrafish   GEASGE-------VKKADRESWRSNKVYTDDQSEK----PKSRVFGRSFPPRLNA-AYNR     1167
frog        NDPAVQEREAEDLRKGDFQDSWRSNRNYSDDQNNTDSKSRAPRAFGRSLPPRLSN-SYGR     1211
chicken     NDGSVLDREDSDLKKGDFKESWRSNKIYSDDHTSLDPKMRAPRAFGRSLPPRLSNSGYGR     1208
human       NEGSLLEREESTLKKGDCRDSWRSNKGCSEDHSGLDAKSRGPRAFGRALPPRLSNCGYGR     1216
mouse       HEGMLTERDEG------ALKDSWRSNRTYTEDQGGIDTRSRGSRTCGRALPPRLSNCSYGR     1226
              :                ::*****: ::*:       *. **::****.  .*.*

zebrafish   GFTGTRDISTWRGRGTQF-----GS-------SSSQENGYNFVADSYSKRTEPLKYPAKF     1215
frog        RPLSTKESSHWQAKSGGSSWQEYGA--PAEAYGSRHHADRDYAH-DY-------RYADSF     1261
chicken     RGFMGKEPTQWQGRSGGAGWQEYSHTSPSDGFGSRQQSDRDYIQDSY-------KHVDSF     1261
human       RTFVSKESPHWQSKSPGSSWQEYGP---SDTCGSRRPTDRDYVPDSY-------RHPDAF     1266
mouse       RTFVAKEPPHWQSRSPGSSWQEYGS---SDPCGPRRGTDRDYIPDSY-------RQSDTF     1276
```

```
                ::    *:.:.            .          .  :  .::  .*        :     *
zebrafish  TGTFAENGVEDRDGGYY------VDDNPDNRPLRRRRPPRQDKPPRFRRLRQERDGAGAW   1269
frog       PSRGFDESHGDERRSFFQEEYS-DRDSLEKRSFGRRRPPRQDKPPRFRRLRQERDPLGQW   1320
chicken    SSRVFDESHLDDKRHFFQEDYLADQENIENRPFRRRRPPRQDKPPRFRRLRQERELVGQW   1321
human      GGRGFEDSRAEDKRSFFQDEHVADSENAENRPFRRRRPPRQDKPPRFRRLRQERESLGLW   1326
mouse      GSRLFEDSRTEDKRSFFQDDHGADSENAENRPFRRRRPPRQDKPPRFRRLRQERESLGLW   1336
              .   ::.  :    ::          :. ::* : ********************:  * *

zebrafish  SNDDYV-NGEFPNQWPSRAKSAEEHWQNHYPGGRS---QSQEMTAQSQGEDWETGSDNSD   1325
frog       SGEETVGTGNSSEHWQSRPRVPLNDKSG--PVHRRSPEMSHHN-SDHVTEDWETASESSD   1377
chicken    NPEEG-GPNLLPSQWPGRPKLSTTEKSS--ISGRRSPELSYQNSSDHANEEWETASESSD   1378
human      GPEEE--PHLLAGQWPGRPKLCSGDKSG--TVGRRSPELSYQNSSDHANEEWETASESSD   1382
mouse      GPEEE--SHLLASQWPGRSKLCPGDKSG--PGGHRSPELSYQNSSDHANEEWETASESSD   1392
             . ::        :* .* :    . ..      :      * .  ::    *:***.*:.**

zebrafish  FGDWREKRGPHGD-PL---TDAALGDPGSEKRELSKRSFSSQRPLD-RQNRKNDVSVMEN   1380
frog       FSEKRPG---------DMDGDGSQSGSGLSEKRELSKRSFSSQRPLMDRQSRKVEPSGYDE   1429
chicken    FSERRERRDGVS-ESEGQLESGLGSGSLGEKRELAKRSFSSQRPLVDRQNRKAEPAGYAE   1437
human      FSERRERREGPGSEPDSQVDGGLSGASLGEKKELAKRSFSSQRPVVDRQSRKLEPGGFGE   1442
mouse      FSERRERREGLVAEPEAQGDGGLSGSSLGEKKELAKRSFSSQRPLADRQSRKLEPGGFGE   1452
           *.: *             .. .   .**:**:**********:  **.** : .   :

zebrafish  SK-MGRPADAGSRNDSWQNGVASGSKRSPEESVSGL-------SSASVYGVEQNEDRS--   1430
frog       KPSRVSGSS--SRNDFQKSSGPLKSSRCSDDSYG-------AESSHHRYAMDRSMHCDSG   1480
chicken    QSVRTVGA--ASRYESQQNGTLIKSKRSPEE--GGGLGNTSGGSSHSIYSLDRASLANSE   1493
human      KPVRPGGGDTSPRYESQQNGTPLKVKRSPDEALPGGLSGCSSGSGHSPYALERAAHASAD   1502
mouse      KPVRPGGGEPSPRCESQQSGTPLKVKRSPDEALPGGLG------SHSPYALERTTHASSD   1506
             .      .   * :  :..    .*. ::       .   *.::: .

zebrafish  TPEPTGKIIEKELKPRN-MKADMTEPLSQYDLNTYSIESDSG---------ASVPSPEVF   1480
frog       E---AGKKQERDSRTPGLKSSDKNEVIGQFDLKYGDSMLEEE-------SEV---------   1522
chicken    SAEGPGKKSEKEPKSAVQRASEKGETLSQFELNYGSTIIDNRVSSTAEENEVGSMAGEGF   1553
human      LPEASSKKAEKEAKLAAPRAGEQGEAMKQFDLNYGSAIIENCGSSPGEESEVGSMVGEGF   1562
mouse      GPETPSKKSEREVSLPTQRASEQEEARKQFDLGYGNALIDNCASSPGEENEASSVVGEGF   1566
               .*   *::         .: *    *::*    .   :.                .

zebrafish  QDSLSKKQRRPQEDERRRKEQGPPSSVKSRTIASKMPPRFAKKQGSMSMEQPEETLTANN   1540
frog       -GDIGNKPRRVLEKDRRKKDQVVQVPAKGSSIQSRIPPRFAKKQNGMCLDQADVTA----   1577
chicken    IEVLTKKQRRLLEEERRKKEQAAQAPAKARVLQSRIPPRFAKKQNSLCLEQSDVTVSGNS   1613
human      IEVLTKKQRRLLEEERRKKEQAVQVPVKGRGLSSRIPPRFAKKQNNLCLEQGDVTVPGSS   1622
mouse      IEVLTKKQRRLLEEERRKKEQAAQVPVKGRGLSSRIPPRFAKKQNGLCLEQ-DVTVPGSS   1625
             : :* **  *.:**:*:*     .*.  : *::********..:.::* : *

zebrafish  LGTEIWETNSTALTVQSSGGDSWTKQVSYTGSEP-------------------------   1574
frog       --KEIWESSGQGISVQ-PGTDTWSKPVSSFSTES--SSTEGFKGSQGDSGIDLSAESRES   1632
chicken    LGTEIWESNSPALSVQSPGSDSWSKPVNTFNGTESSSTEQGFKGSQGDSGIDLSAESRES   1673
human      LGTEIWESSSQALPVQAPANDSWRKAVTAFSSTETGSAEQGFKSSQGDSGVDLSAESRES   1682
mouse      LGTEIWENSSQALPVQGAASDSWRTAVTAFSSTEPGT-SEGFKSSQGDSGVDLSAESRES   1684
            .****... .: **  . *:* . *.  .

zebrafish  ----------------------------------------NSEDSDAGPEQSKEHKPGPI   1594
frog       SATSSQRSSPYGTMKPEEMNGTGVAEPKPDCPKEQGQKQSEKKDSDPGSGLNKEHKPGPI   1692
chicken    SATSSQRSSPYGTLKPEEMNGAGLVDPKPDCQKEQVQKQSDKKDSDQGSGQNKEHKPGPI   1733
human      SATSSQRSSPYGTLKPEEMSGPGLAEPKADSHKEQAPKPSEQKDSEQGSGQSKEHRPGPI   1742
```

```
mouse       SATSSQRSSPYGTLKPEEISGPGLAESKADSHKDQAQKQAEHKDSEQGSAQSKEHRPGPI        1744
                              : :**: *    .***:****

zebrafish   GNERSLKNRKGSEGLERLEGS-ITPVNGVDIHVENVIPVPPIEFGVNAKDSDFSLPAGSA        1653
frog        GNERSLKNRKGSEGTERLEGG-VPPVNGVEIHVDSVLPVPPIEFGVSAKDADYALPPGTA        1751
chicken     GNERSLKNRKGSEGTERLEGN-IPPVNGVEIHVDSVLPVPPIEFGVNPKDSDFSLPPGSA        1792
human       GNERSLKNRKGSEGAERLQGAVVPPVNGVEIHVDSVLPVPPIEFGVSPKDSDFSLPPGSA        1802
mouse       GNERSLKNRKGSEGAERLPGAVVPPVNGVEIHVDSVLPVPPIEFGVSPKDSDFSLPPGSV        1804
            ************** *** *  : *****:***:.*:*********. **:*::** *:.

zebrafish   AVPVSNPVSKLQDALAGNAGLTQAIPMLRRDH-LQPAITLNPISFPSADLTLKMESARKA        1712
frog        TVQAANTVTKLQDALASKAGLTQSIPILRRDHHMQQGMSLN-MSYPTADLTLKMESARKA        1810
chicken     SGTAANPVTKLQDALASNAGLTQSIPILRRDHHLQRCIGLNPMSFPTAELTLKMESARKA        1852
human       SGPTGSPVVKLQDALASNAGLTQSIPILRRDHHIQRAIGLSPMSFPTADLTLKMESARKA        1862
mouse       SGPVGNPVAKLQDVLASNAGLTQSIPILRRDHHMQRAIGLSPMSFPTADLTLKMESARKA        1864
            :  ... * ****.**.:*****:**:***** :*  : *. :*:*:*:**********

zebrafish   WENSQAVPEQGSPGS-GSSAQPVCSVGSS-SVS---YSSFSMPPMPVASVAPSVSMQGSH        1767
frog        WENSPSLPEQNSPAGPGSGIQPPSSVGASTGVNYSSFGGVSMPPMPVASVAPSASMPGNH        1870
chicken     WENSPSLPEQNSPGGAGSGIQPPSSVGASNGVSYSSFGGVSMPPMPVASVAPSASIPGNH        1912
human       WENSPSLPEQSSPGGAGSGIQPPSSVGASSGVNYSSFGGVSMPPMPVASVAPSASMPGSH        1922
mouse       WENSPSLPEQSSPGGAGSGIQPPSSVGASNGVNYSSFGGVSMPPMPVASVAPSASIPGSH        1924
            **** ::***.**.. **. ** .***:* .*.   :...*************.*: *.*

zebrafish   IPPLYLDGHVFPSQPRLVPPL-TQQPSYQQ-ATPQQIPISLHTSLQAQAQLGLRGALPVS        1825
frog        IPPLYLDGHVFTNQPRLVQQTIPQQQGYQQAAAAQQIPISLHTSLQAQAQLGLRGGLPVS        1930
chicken     IPPLYLDGHVFASQPRLVPQTIPQQQSYQQAAAAQQIPISLHTSLQAQAQLGLRGGLPVS        1972
human       LPPLYLDGHVFASQPRLVPQTIPQQQSYQQAAAAQQIPISLHTSLQAQAQLGLRGGLPVS        1982
mouse       LPPLYLDGHVFASQPRLVPQTIPQQQSYQQAATAQQIPISLHTSLQAQAQLGLRGGLPVS        1984
            :********** .*****      ** .*** *: ********************.****

zebrafish   QSQEIFSSIPPFRSQVYMHPNLSQASPMVLSGGAALKGPYSAFPGLQPSDLVKSQSGSHY        1885
frog        QSQEIYSSMQPFRSQVYMHPSLSQPSAMVLTSGTGLKPPYNPFPGMQTLEMVKTQPTSPY        1990
chicken     QSQEMYSSIQPFRSQVYMHPSLSQPSTMVLTGGTALKPPYSAFPGMQPLEVVKTQSGSPY        2032
human       QSQEIFSSLQPFRSQVYMHPSLSPPSTMILSGGTALKPPYSAFPGMQPLEMVKPQSGSPY        2042
mouse       QSQEIFSSLQPFRSQVYMHPSLSPPSTMILSGGTALKPPYSAFPGIQPLEMVKPQSGSPY        2044
            ****::**: **********.**  * *:*:.*:.** **. ***:*  ::** *   * *

zebrafish   QPINGSTPLVYD----QPTGMSTSQLMDSQLIQVTMPMPGSQL---RYGSAQQHLILPQS        1938
frog        QPLNGSQQLVYESQLNQA-----SQMMDSQLTQLTMPMPGSQLQMPRYSSGQQTMILPQS        2045
chicken     QPLNGSQTLVYEGQINQAAGMGASQMMDSQLTQLTMPVPGSQLPLPRYGSGQQPLILPQS        2092
human       QPMSGNQALVYEGQLSQAAGLGASQMLDSQLPQLTM-------PLPRYGSGQQPLILPQS        2095
mouse       QPMSGNQALVYEGQLGQAAGLGTSQMLDSQLPQLTM-------PLPRYGSGQQPLILPQS        2097
            **:.*. ***:     *    **::**** *:**        **.*.** :*****

zebrafish   IQLQQNQNLSVGAPRRMMPPGSQP-VLSGSREVSQMDMKGFQFSDKPSHSPGI----PSG        1993
frog        IQLPQGQNLPVGAPRRMQP------SMLTSRESSQMEMKSFHFTDGKQNMPTAMQ--AQH        2097
chicken     IQLPQGQNLSVGAPRRILPPGSQPSVLAASRESSQMEMKGFHFSDGKQNMSSGGSVPSPH        2152
human       IQLPPGQSLSVGAPRRIPPPGSQPPVLNTSREPSQMEMKGFHFADSKQNVPSGGPVPSPQ        2155
mouse       IQLPPGQSLSVGAPRRVPPPGSQPPVLNTSRESAPMELKGFHFADSKQNVPTGGSAPSPQ        2157
            ***  .*.* ****** *      :  *** : *::*.*:*:*  .:

zebrafish   SYR--------------PGSASPSGKASGAP---AVASLAGHYTQQVSGPQGSMVMHMRPP        2037
frog        SYR--------------PSSASPSGKSPGPGPSANLGSVQGHYPQQVKHRSDDSKGGLR--        2142
chicken     AYRIYSMNVVDSSISRPSSASPNGKPS--GPAVSMGSVQGHYVQQAKQRVDENKANLG--        2208
```

```
human       TYR-------------PSSASPSGKPS--GSAVNMGSVQGHYVQQAKQRVDEK-PSLG--        2197
mouse       AYR-------------PSSASPSGKPS--GSAVNMGSVQGHYVQQAK-RVDEK-PGLG--        2198
            :**             *.****.**         :.*: *** **..    ..     :

zebrafish   SSGPFPTPIQRPIMQVNKPVIIRSPPYPGPNTGPAHTHAHTHNPEGPGKGPEDGLKECVA        2097
frog        ------------------------------------------------------SPKVPE        2148
chicken     ------------------------------------------------------AVKLQE        2214
human       ------------------------------------------------------AVKLQE        2203
mouse       ------------------------------------------------------TVKLQE        2204
                                                                    :

zebrafish   SAAPVKASASRSGAIKPQPLKLEEGKA        2124
frog        QPSASPMKPARTGAIKPQAVKVEESKA        2175
chicken     TASTSQMKPVRTGAIKPQAVKVEESKA        2241
human       APSAA-SQMKRTGAIKPRAVKVEESKA        2229
mouse       ASSAT-SQMKRTGAIKPRAVKVEGSKA        2230
            :     .   *:*****: :*:* .**
```

Extended Data 2. Sequence alignment of PRRC2A, PRRC2B, and PRRC2C proteins in humans was shown.

```
PRRC2A[Homo_sapiens(human)]   MSDRSGPTAKGKDGK-KYSSLNLFDTYKGKSLEIQKPAVAPRHGLQSLGKVAIARRMPPP   59
PRRC2B[Homo_sapiens(human)]   MSDRLGQITKGKDGKSKYSTLSLFDKYKGKSVDAIRSSVIPRHGLQSLGKVAAARRMPPP   60
PRRC2C[Homo_sapiens(human)]   MSEKSGQSTKAKDGK-KYATLSLFNTYKGKSLETQKT--TARHGLQSLGKVGISRRMPPP   57
                              **::  *   :*.**** **::*.**:.*****::   :      **********. :******

PRRC2A[Homo_sapiens(human)]   ANLPSLKAENKGNDPNVSLVPKDGTGWASKQEQSDPKSSDASTAQPPESQPLPASQTPAS   119
PRRC2B[Homo_sapiens(human)]   ANLPSLKSENKGNDPNIVIVPKDGTGWANKQDQQDPKSSSATASQPPESLPQPGLQKSVS   120
PRRC2C[Homo_sapiens(human)]   ANLPSLKAENKGNDPNVNIVPKDGTGWASKQEQHEEEKTP----EVPPAQPKPGVA----   109
                              *******:*********: :*********.**:* : :.:       : * : * *.

PRRC2A[Homo_sapiens(human)]   NQPKR-PPAAPENTPLVPSGVKSWAQASVTHGAHGDGGRASSLLSRFSREEFPTLQAAGD   178
PRRC2B[Homo_sapiens(human)]   NLQKPTQSISQENTNSVPGGPKSWAQLNGKPVGHEGGLRGSSRLLSFSPEEFPTLKAAGG   180
PRRC2C[Homo_sapiens(human)]   ------------APPEVAPAPKSWASN--KQGGQGDGIQVN----SQFQQEFPSLQAAGD   151
                                          *  . ****.    .  .: .* : .        :***:*:***.

PRRC2A[Homo_sapiens(human)]   QDKAAKERESAEQSSGPGPSLRPQNSTTWRDGGGRGPDELEGP----------DSKLHHG   228
PRRC2B[Homo_sapiens(human)]   QDKAGKEKGVLDLSYGPGPSLRPQNVTSWREGGGRHIISATSLSTSPTELGSRNSSTGDG   240
PRRC2C[Homo_sapiens(human)]   QEKK--EKETNDDNYGPGPSLRPPNVACWRDGGKAAGSPSSSDQ-DEKLPGQDESTAGTS   208
                              *:*    *:   :  . ******** * : **:**       .            :*.   .

PRRC2A[Homo_sapiens(human)]   HD--------PRGGLQPS-----------G-PPQFPPYRGMMPPFMYPPYLPFPPPYGPQ   268
PRRC2B[Homo_sapiens(human)]   APSSACTSDSKDPSLRPAQPVRKGASQFMGNVYHPPTYHDMLPAFMCSPKSSENQGTVER   300
PRRC2C[Homo_sapiens(human)]   EQNDILKVVEKRIACGPPQAKLNGQ-----QAALASQYRAMMPPYMFQQYPRMTYPP-LH   262
                                          .   *                *: *:* :*            :

PRRC2A[Homo_sapiens(human)]   GPYRYPT--------------PDGPSRFPRVAGPRGSGPPMRLVEPVGRPSILKEDNLKE   314
PRRC2B[Homo_sapiens(human)]   GSFPLPQLRLEPRVPFRQFQMNDQDGKENRLGLSRPLRPLRQLVERAPRPTIINAENLKG   360
PRRC2C[Homo_sapiens(human)]   GPMRFPPS-----------------LSETNKGLRGRGPPPSWASEPERPSILSASELKE   304
                              *     *                          *   *     ..   **:*:. .:**

PRRC2A[Homo_sapiens(human)]   ---FDQLDQENDDGWAGAHEEVDYTEKLKFSDEEDGRDSDEEGAEGHRDSQSASGEER-P   370
PRRC2B[Homo_sapiens(human)]   ---LDDLDADADDGWAGLHEEVDYSEKLKFSDDEEEEEVVKDGRPKWNSWDPRRQRQLSM   417
PRRC2C[Homo_sapiens(human)]   LDKFDNLDAEADEGWAGAQMEVDYTEQLNFSDDDEQGSNSPKENNSEDQGSKASENNENK   364
                                 :*:** : *:**** : ****:*:*:***::: .    .        . .   .:

PRRC2A[Homo_sapiens(human)]   PEADGKKGN----------SPNSEPPTPKTAWAETSRPPETEPGPPAPKPPLPPPHRGPA   420
PRRC2B[Homo_sapiens(human)]   SSADS--------------ADAKRTREEGKDWAEAVGASRVVRKAPDPQP-----PPRKL   458
PRRC2C[Homo_sapiens(human)]   KETDEVSNTKSSSQIPAQPSVAKVPYGKGPSFNQERGTSSH----------LPP-PPKLL   413
                               .:*               :  .        : :

PRRC2A[Homo_sapiens(human)]   GNWGPPGDYPDR--GG-PPCKPPAPEDEDEAWRQRRKQSSSEISLAVERARRRREEEERR   477
PRRC2B[Homo_sapiens(human)]   HGWAPGPDYQKS--SMGSMFRQQSIEDKEDKPPPRQKFIQSEMSEAVERARKRREEEERR   516
PRRC2C[Homo_sapiens(human)]   AQQHPPPDRQAVPGRPGPFPSKQQVADEDEIWKQRRR-QQSEISAAVERARKRREEEERR   472
                               *   *                *:::    *:: .**:* ******:********

PRRC2A[Homo_sapiens(human)]   MQEERRAACAEKLKRLDEKFGAPDKRLKAEPAAPPAAPSTPAPPPAVPKELPAPPAPP--   535
PRRC2B[Homo_sapiens(human)]   AREERLAACAAKLKQLDQKCKQARKAGEARKQAEKEVPWSPSAEKASPQEN----G----   568
PRRC2C[Homo_sapiens(human)]   MEEQRKAACAEKLKRLDEKLGILEKQPSPEEIRERERKERERKELEKEQEQEREKERE    532
                               .*:* **** ***:**:*       *  . .                :*
```

```
PRRC2A[Homo_sapiens(human)]    --PA---------------SAPTP---ETEPEE----------PAQAPPAQS--------    557
PRRC2B[Homo_sapiens(human)]    --PA-----------------VH---KGSP----------------------------    576
PRRC2C[Homo_sapiens(human)]    KDRERQQEKEKELEKEQEKQREMEKERKQEKEKELERQKEKEKELQKMKEQEKECELEKE    592
                                                                        :  .

PRRC2A[Homo_sapiens(human)]    ------TPTPGVAAAPTLVSGGGSTSSTSSGSFEASPVEPQLPSKEGPEPPEEVPPPTTP    611
PRRC2B[Homo_sapiens(human)]    ----------------------------------------EFPAQETPTTFPEEAPTVSP    596
PRRC2C[Homo_sapiens(human)]    REKLEEKIEPREPNLEPMVEKQES---ENSC---NKEEEPVFTRQDSNRSEKE----ATP    642
                                                                    : ::       *     .:*

PRRC2A[Homo_sapiens(human)]    PVPKVEPKGDGIGPTRQPPSQGLGYPKYQKSLPPRFQRQQQEQLLKQQQ-QHQWQQHQQG    670
PRRC2B[Homo_sapiens(human)]    AVAQSNSSEE----EAREAGSPAQEFKYQKSLPPRFQRQQQQQQQEQLYKMQHWQPVY--    650
PRRC2C[Homo_sapiens(human)]    VVH---ETEPESGSQPRPAVLSGYFKQFQKSLPPRFQRQQEQM------KQQQWQQQQQQ    693
                               *       .       :       ::************::        ::**

PRRC2A[Homo_sapiens(human)]    SAPPTPVPPSPPQPVTLGAVPAPQAPPPPPKALYPGALGRPPPMPPMNFDPRWMMIPPYV    730
PRRC2B[Homo_sapiens(human)]    --P----PP-----------------SHPQRTFYPH------HPQMLGFDPRWMMMPSYM    681
PRRC2C[Homo_sapiens(human)]    GVLPQTVPSQPS---------SSTVPPPPHRPLYQPMQPHPQHLASMGFDPRWLMMQSYM    744
                                      *                   * : :*           :.*****:*:  *:

PRRC2A[Homo_sapiens(human)]    DPRLLQGRPPLDFYPPG---VHPSGLVPRERSD--SGGSS--SEPFDRHAPAMLRERGTP    783
PRRC2B[Homo_sapiens(human)]    DPRITPTRTPVDFYPSA---LHPSGLMKPMMPQESLNGTG--CRSEDQNCVPPLQERKVT    736
PRRC2C[Homo_sapiens(human)]    DPRMMSGRPAMDIPPIHPGMIPPKPLMRRDQ----MEGSPNSSESF-EHIARSARDHAIS    799
                               ***:    *   :*: *      : *. *:       *:    ..    .:     :::

PRRC2A[Homo_sapiens(human)]    PVDPKLAWVGDVFTATPAEPRPLTSPLRQAADEDDKGMRSETPP----------------    827
PRRC2B[Homo_sapiens(human)]    PIDSPPVWSPEGYMALQSKGYPLPHPKSSDTLAMDMRVRNESSFSASLGRAGGVS-----    791
PRRC2C[Homo_sapiens(human)]    LSEPRMLWGSDPYPHAEPQQA--TTPKAT---EEPEDVRSEAALDQEQITAAYSVEHNQL    854
                                :      *  : :       :        *           :*.*:

PRRC2A[Homo_sapiens(human)]    ---------VPPPPPYLASYP------------------------------------GFP    842
PRRC2B[Homo_sapiens(human)]    ---AQRDLFEERGEEYLSAFDKKAQAD----------FDSCISSQRIGQELLFPPQENVQ    838
PRRC2C[Homo_sapiens(human)]    EAHPKADFIRESSEAQVQKFLSRSVEDVRPHHTDANNQSACFEAP--DQKTLSAPQEER-    911
                                        :  :

PRRC2A[Homo_sapiens(human)]    ENGAPGPPISR---------FPLEEPGPRPLPWPPGSDEVAKIQTPPPKKEPPKEETAQ-    892
PRRC2B[Homo_sapiens(human)]    DAGAPGGHTQNLRCSPLEPDFVPDEKKPECGSWDVSHQPETADTAHGVERETPREGTA--    896
PRRC2C[Homo_sapiens(human)]    ---------ISAVESQPSR---------KRSVSHGSNHT-----QKPDEQRSEPSAGIPKV    949
                                       .              .         .          ::. *

PRRC2A[Homo_sapiens(human)]    ----------LTGPEAG-RKPARGVGSGGQGPPPPRRESRTETRWGPRPGSSRRGIPPEE    941
PRRC2B[Homo_sapiens(human)]    ----------------------FNISS-WDKNGSPNKQPSSEPEWTPEPRSSSSQHPE--    931
PRRC2C[Homo_sapiens(human)]    TSRCIDSKEPIERPEEKPKKEGFIRSSEG-PKPEKVYKSKSETRWGPRPSSNRREEV---   1005
                                         .*              :  :* .* *.* *.

PRRC2A[Homo_sapiens(human)]    PGAPPRRAGPIKKPPPPTKVEELPPKP-----------------LEQG---DETPKPPKP    981
PRRC2B[Homo_sapiens(human)]    QTGRTRRSGPIKKPVLKALKVEDKEKELE---K-------IK--QELG---EESTRLAKE    976
PRRC2C[Homo_sapiens(human)]    NDRPVRRSGPIKKPVLRDMKEEREQRKEKEGEKAEKVTEKVVVKPEKTEKKDLPPPPPPP   1065
                                   **:******        *   :                *       :

PRRC2A[Homo_sapiens(human)]    D----------------------------------------------------PLKITK    988
PRRC2B[Homo_sapiens(human)]    KEQSP------------TAEKDEDE-----------ENDASLANSSTTTLEDKGPGHATF   1013
PRRC2C[Homo_sapiens(human)]    QPPAPIQPQSVPPPIQPEAEKFPSTETATLAQKPSQDTEKPLEPVSTVQVE---PAVKTV   1122
                               .                                               *   *
```

```
PRRC2A[Homo_sapiens(human)]    --GKL-GG-----------P-K----ETPPNGNLSP------APRLRRDYS-------YER      1017
PRRC2B[Homo_sapiens(human)]    --GRE-ATKFEEEEKPDKAWE----ARPPRESSDV------PPMKRNNWIFIDEEQAFGV      1060
PRRC2C[Homo_sapiens(human)]    NQQTMAAPVVKEEKQPEKVISKDLVIERPRPDSRPAVKKESTLPPRTYWKEARERD---W      1179
                                     .                 .       *. .            *   :

PRRC2A[Homo_sapiens(human)]    VGPTSCRGRGRGEYFARGRGFRGT----YGGRGRGARSREFRSYREFRGDDGRGGGTG--      1071
PRRC2B[Homo_sapiens(human)]    RGQARGRGRGFREFTFRGRPAGGNGSGLCGGGVLGARSIYCSSQ-----RSGRGRGLREF      1115
PRRC2C[Homo_sapiens(human)]    FPDQGYRGRGRGEYYSRGRSYRGS----YGGRGRGGRGHT-RDYPQYRDNKPR-------      1227
                                      ****  *:  ***    *.       **   *.*.    .         . *

PRRC2A[Homo_sapiens(human)]    ---GPNHPPAPRGRTASETRSEGSEYEEIPKRRRQRGSETGSETHESDLAPSDKEAPTPK      1128
PRRC2B[Homo_sapiens(human)]    ARPEDCPRAKPRRRVASETHSEGSEYEELPKRRRQRGSENGNEGSLLEREEST-----LK      1170
PRRC2C[Homo_sapiens(human)]    --AEHIPSGPLRQREESETRSESSDFEVVPKRRRQRGSETDTDSEIHESASD---KDSLS      1282
                                        * *   ***:**.*::* :**********...:    :    .       .

PRRC2A[Homo_sapiens(human)]    EGTLTQVPLAPPPPGAPP---------SPAPA----------RFTARGGRVFTPRGVPSRR      1170
PRRC2B[Homo_sapiens(human)]    KGDCRDSWR--SNKGCSE--------DHSGL----------DAKSRGPRAFGRALPPRLS      1210
PRRC2C[Homo_sapiens(human)]    KGKLPKREERPENKKPVKPHSSFKPDNHVRIDNRLLEKPYVRDDDKAKPGFLPKGEPTRR      1342
                                :*    .                    .                  :.   *      *

PRRC2A[Homo_sapiens(human)]    --G----RGGGRP---P----------PQVCPGWSPPAKSLAPKKPPTGPLPPSKEPLKEK      1212
PRRC2B[Homo_sapiens(human)]    NCG----YGRRTF---V---------SKESPHWQSKSPGSSWQEYGPSDTCGSRRPTDRD      1254
PRRC2C[Homo_sapiens(human)]    GRGGTFRRGGRDPGGRPSRPSTLRRPAYRDNQWNPRQSEVPKPE------DGEPPRRHEQ      1396
                                *       *                     *.           :         .   ...

PRRC2A[Homo_sapiens(human)]    LIPGPLSP-VARGGSN--------------GGSNVGMEDGERPRRRHGRAQQQDKPPRFR      1258
PRRC2B[Homo_sapiens(human)]    YVPDSYRHPDAFGGRGFEDSRAEDKRSFFQDEHVADSENAENRPFRRRRPPRQDKPPRFR      1314
PRRC2C[Homo_sapiens(human)]    FIPIA----------------ADKRPPKF--ERKFDPARERPRRQRPTRPPRQDKPPRFR      1438
                                :*                    ..          . *    *    *  :********

PRRC2A[Homo_sapiens(human)]    RLKQERENAARG-SEGKPSLTLPASAPGPEEALTTVTVAPAPRRAAAKSPDLSNQ-NSDQ      1316
PRRC2B[Homo_sapiens(human)]    RLRQERESLGLWGPEEEPHL-LAGQWPGRPKLC----SGDKSGTVGRRSPELSYQNSSDH      1369
PRRC2C[Homo_sapiens(human)]    RLREREAASKSNEV-------VAVPTNGTVNNVAQEPVNTLGDISGNKTPDLSNQNSSDQ      1491
                                **::..         :       *  :            . ::*:** * .**:

PRRC2A[Homo_sapiens(human)]    ANEEWETASESSDFTSERRGDKEAPPPVLLTPKAVGTPGGGGGGAVPGISAMSRGDLSQR      1376
PRRC2B[Homo_sapiens(human)]    ANEEWETASESSDFSERRERRE-GPGSE---------PDSQVDG---GL----SGASLGE      1412
PRRC2C[Homo_sapiens(human)]    ANEEWETASESSDFNERRERDEKKNADLNA-Q----------------TVVKVGENVLPP      1534
                                **************...*.  :

PRRC2A[Homo_sapiens(human)]    AKDLSKRSFSSQRPGMERQNRRPGPGGKA-GSSGSS----------SGGGGG-----GPG      1420
PRRC2B[Homo_sapiens(human)]    KKELAKRSFSSQRPVVDRQSRKLEPGGFG-EKPVR-----------PGGGDT-----SPR      1455
PRRC2C[Homo_sapiens(human)]    KREIAKRSFSSQRPVD-RQNRRGNNGPPKSGRNFSGPRNERRSGPPSKSGKRGPFDDQPA      1593
                                :::::*********   **.*:   *                     .*        *

PRRC2A[Homo_sapiens(human)]    GRTGP----------------GRGDKRSWPSPKNRSRPPEERPPGLPLPPPPPSSSAVFR      1464
PRRC2B[Homo_sapiens(human)]    Y-----------------------ESQQNGTPLKVKRSPDEALPGGLSGCSSGSGHSPYA      1492
PRRC2C[Homo_sapiens(human)]    GTTGVDLINGSSAHHQEGVPNGTGQKNSKDSTGKK---REDPKPGPKKPKEKVDALSQFD      1650
                                                 :... : :     :: **       .. : :

PRRC2A[Homo_sapiens(human)]    LDQVIH--------------------SNPAGIQQALAQLSSRQ------------------      1487
PRRC2B[Homo_sapiens(human)]    LERAAH--------------------ASA-DLPEASSKKAEKE------------------      1514
PRRC2C[Homo_sapiens(human)]    LNNYASVVIIDDHPEVTVIEDPQSNLNDDGFTEVVSKKQQKRLQDEERRKKEEQVIQVWN      1710
                                *:.                       .  .: :. ::  .:.
```

```
PRRC2A[Homo_sapiens(human)]    -------G----SVTAPGGHPRHKPGLPQAP-------------Q-------------GPS    1511
PRRC2B[Homo_sapiens(human)]    -------A----KLAAPR-----------------------A-------------GEQ    1525
PRRC2C[Homo_sapiens(human)]    KKNANEKGRSQTSKLPPRFAKKQATGIQQAQSSASVPPLASAPLPPSTSASVPASTSAPL    1770
                                     .     .    *                                    .

PRRC2A[Homo_sapiens(human)]    PR-------------PPTRY----------------EPQRVNSGLSS-------------    1529
PRRC2B[Homo_sapiens(human)]    GE-------------AMKQF----------------D---LNY----------------    1536
PRRC2C[Homo_sapiens(human)]    PATLTPVPASTSAPVPASTLAPVLASTSAPVPASPLAPVSASASVSASVPASTSAAAITS    1830
                                            .                            .

PRRC2A[Homo_sapiens(human)]    --DP-HFEEPGPMVRGVGGTPRDSAGVS---------PFPPKR-----------------    1560
PRRC2B[Homo_sapiens(human)]    ----------GSAIIENCGSSPGEESEVGSMVGEGFIEVLTKK-----------------    1569
PRRC2C[Homo_sapiens(human)]    SSAPASAPAPTPILASVS-TPA---SV-TILASASIPILASALASTSAPTPAPAASSPAA    1885
                                        ::  .  . :*       *           .  .

PRRC2A[Homo_sapiens(human)]    -----------------------RERPP----------------RKPELLQEESLP-PPH    1580
PRRC2B[Homo_sapiens(human)]    -----------------------QRRLL----------------EEERRKKEQAVQVPVK    1590
PRRC2C[Homo_sapiens(human)]    PVITAPTIPASAPTASVPLAPASASAPAPAPTPVSAPNPAPPAPAQTQAQTHKPVQNPLQ    1945
                                                                   :.    .: :   * :

PRRC2A[Homo_sapiens(human)]    SSGFLGSKPE-------GPGPQAE-------S--RDTGTEALTPHIWNRLHTA-------    1617
PRRC2B[Homo_sapiens(human)]    GRGLSSRIPPRFAKKQNNLCLEQG-------D--VTVPGSSLGTEIWESSSQA-------    1634
PRRC2C[Homo_sapiens(human)]    TTSQSSKQPPPSIRLPSAQTPNGTDYVASGKSIQTPQSHGTLTAELWDNKVAPPAVLNDI    2005
                                 .    *           :        .       :*   .:*:

PRRC2A[Homo_sapiens(human)]    ----TSRKSYRPSSMEPWMEPLSPFED------VAGTEMSQSDSGVDLSGDSQVSSGPCS    1667
PRRC2B[Homo_sapiens(human)]    ----LP---VQAPANDSWRKAVTAFSSTETGSAEQGFKSSQGDSGVDLSAESRESSATSS    1687
PRRC2C[Homo_sapiens(human)]    SKKLGPISPPQPPSVSAWNKPLTSFGSAPSSEGAKN----GQESGLEIGTDTIQFGAPAS    2061
                                    :  : . * : :: * .      .        :**:::. ::   .. .*

PRRC2A[Homo_sapiens(human)]    QRSSPDGGLKGAAEGPPKRPGGSSPLNAVPCEGPPGSEPPRRPPPAPHDGDRKELPREQP    1727
PRRC2B[Homo_sapiens(human)]    QRSSPYGTLKPEEMSGPGL---AEPKAD--------SHKEQAPKPSEQKDSEQGSGQSKE    1736
PRRC2C[Homo_sapiens(human)]    NGNENEVVPVLSEKSADKIPEPKEQRQ----------KQPRAGPIKAQKLPDLSPVENKE    2111
                               : ..         .            .          . :      :.       ..:

PRRC2A[Homo_sapiens(human)]    LPPGPIGTERSQRTDRGTEPGP----------------------------IRPSH    1754
PRRC2B[Homo_sapiens(human)]    HRPGPIGNERSLKNRKGSEGAERLQG-------AVVP-----------PVNGVEIHVDSV    1778
PRRC2C[Homo_sapiens(human)]    HKPGPIGKERSLKNRKVKDAQQVEPEGQEKPSPATVRSTDPVTTKETKAVSEMSTEIGTM    2171
                                 *****.*** :. : .:                                      :

PRRC2A[Homo_sapiens(human)]    RPGPPVQFGTSDKDSDLRLVVG--DSLKAEKEL----------TASVTEAIPVSRDWELL    1802
PRRC2B[Homo_sapiens(human)]    LPVPPIEFGVSPKDSDFSLPPG--SASGPTGSPVVKLQDALASNAGLTQSIPILRRDHHI    1836
PRRC2C[Homo_sapiens(human)]    ISVSSAEYGTNAKESVTDYTTPSSSLPNTVATNNTKMEDTLVNNVPLPNTLPLPKRE-TI    2230
                                :: *.. *:*         .                .. : :::*: :     :

PRRC2A[Homo_sapiens(human)]    PSA-------AASA------EPQSKNLDSGHCVPEPSSS---GQRLYPEVFYG------S    1840
PRRC2B[Homo_sapiens(human)]    QRAIGLSPMSFPTADLTLKMESARKAWENSPSLPEQSSPGGAGSGIQPPSSVG------A    1890
PRRC2C[Homo_sapiens(human)]    QQSSSLTSVPPTTFSLTFKMESARKAWENSPNVREKGSPVTS---TAPPIATGVSSSASG    2287
                                :      :       *   *  :..  : * .*        *      *        .

PRRC2A[Homo_sapiens(human)]    -AGPSSSQISGGAMDSQ----LHP-----------------------------------    1859
PRRC2B[Homo_sapiens(human)]    SSGVNYSSFGGVSMPPMPVASVAPSASMPGSHL--------------------------    1923
PRRC2C[Homo_sapiens(human)]    PSTANYNSFSSASMPQIPVASVTPTASLSGAGTYTTSSLSTKSTTTSDPPNICKVKPQQL    2347
                                :   . ..:.. :*         : *
```

```
PRRC2A[Homo_sapiens(human)]    ------------------------------------------------------------    1859
PRRC2B[Homo_sapiens(human)]    ----------------------------------------PPLYLDG-HVFA-SQPRLVP    1941
PRRC2C[Homo_sapiens(human)]    QTSSLPSASHFSQLSCMPSLIAQQQQNPQVYVSQSAAAQIPAFYMDTSHLFNTQHARLAP    2407

PRRC2A[Homo_sapiens(human)]    -------------------------------------NSGGFRPG----------T-PSLHPY    1874
PRRC2B[Homo_sapiens(human)]    QTIPQQQSYQ----QAAAAQQIPISLHTSLQA--QAQLGLRGGLPVSQSQEIF-SSLQPF    1994
PRRC2C[Homo_sapiens(human)]    PSLAQQQGFQPGLSQPTSVQQIPIPIYAPLQGQHQAQLSLGAGPAVSQAQELFSSSLQPY    2467
                                                        .  .:  *              **:*:

PRRC2A[Homo_sapiens(human)]    RSQPLYLPPGPAPPSA--LLSGLALKGQFLDFSTMQATELGKLPAGGVLYPPP-----SF    1927
PRRC2B[Homo_sapiens(human)]    RSQV-YMHPSLSPPSTMILSGGTALKPPYSAFPGMQPLEMVKPQSGS-------------    2040
PRRC2C[Homo_sapiens(human)]    RSQPAFMQSSLSQPSV--VLSGTAI----HNFPTVQHQELAKAQSGLAFQQTSNTQPIPI    2521
                               ***   ::  . : **.  : .* *:      *  :*  *: *  :*

PRRC2A[Homo_sapiens(human)]    LYS------PAFCPSPLPDTSLLQVRQDLPSPSDFYSTPLQPG-GQSGFLPSGAPAQ---    1977
PRRC2B[Homo_sapiens(human)]    ----------------------------------------PYQPMSGNQALVYEGQLSQAAG    2062
PRRC2C[Homo_sapiens(human)]    LYEHQLGQASGLGGSQLIDTHLLQARANLTQASNLYSGQVQQP-GQTNFYNTAQS-----    2575
                                                                     *   *:  :    .

PRRC2A[Homo_sapiens(human)]    QMLLPMVDSQLPVVN-----FGSLPPAPPPAPPPLSLLPVGPAL----------QPPSL--    2021
PRRC2B[Homo_sapiens(human)]    LGASQMLDSQLPQLTMPLPRYGSGQQ--PLILPQSIQLPPGQSL---------SVGAP--    2109
PRRC2C[Homo_sapiens(human)]    PSALQQVTVPLPASQLSLPNFGSTGQ--PLIALPQTLQPPLQHTTPQAQAQSLSRPAQVS    2633
                                    :   **            :**      *          *          .  :

PRRC2A[Homo_sapiens(human)]    ---AVRPPPAPATRVLPSPARPFPASLGRAELHPVELKPFQDYQK-LSSNLGGPGSSR--    2075
PRRC2B[Homo_sapiens(human)]    ---RRIPPPGSQPPVLNTS----------REPSQMEMKGFHFADSKQNVPSGGPVPSPQT    2156
PRRC2C[Homo_sapiens(human)]    QPFRGLIPAGTQHSMIA----------TTGKMSEMELKAFGSG---IDIKPGTPPIAGRS    2680
                                  *  .    ::              :  :*:* *       .   * *  :

PRRC2A[Homo_sapiens(human)]    TPPTGRSFSGL-----NSRLKATPSTYSGVFRTQRVDLYQQASPPDALRWIPK-------    2123
PRRC2B[Homo_sapiens(human)]    YRPSSASPSGK-----PSGSAVNMGSVQGHYVQQAKQ----------------------    2188
PRRC2C[Homo_sapiens(human)]    TTPTSSPFRATSTSPNSQSSKMNSIVYQKQFQSAPATVRM--TQPFPTQFAPQAKQRAEV    2738
                                *:.        .           .      .      .  :

PRRC2A[Homo_sapiens(human)]    -------------------------------PWERTGPPPREGPSRRA----EEPGSR-G    2147
PRRC2B[Homo_sapiens(human)]    --------------------------------RVDEKPSLGAVKLQ----EAPSAASQ    2210
PRRC2C[Homo_sapiens(human)]    LQSTQRFFSEQQQSKQIGGGKAQKVDSDSSKPPETLTDPPGVCQEKVEEKPPPAPSIATK    2798
                                                                *       :          *.

PRRC2A[Homo_sapiens(human)]    DKEPGLPPPR---------    2157
PRRC2B[Homo_sapiens(human)]    MKRTGAIKPRAVKVEESKA    2229
PRRC2C[Homo_sapiens(human)]    PVRTGPIKPQAIKTEETKS    2817
                                 . *   *:
```

Supplemental Material and Methods

Reagents, antibodies, plasmids, and siRNAs

Cloning primers (F, forward; R, reverse)	Product	Sequence (5' - 3')
human_PRRC2B_F	FLAG-tagged full-length PRRC2B, P1, T1, T2, T3, T4	CGGGGTACCGCCACCATGTCCGATCGTTTGGGGCA
human_PRRC2B_750_R	FLAG-tagged P1	ATAAGAATGCGGCCGCCACATGTAGCCCTCTGGGCTC
human_PRRC2B_150_R	FLAG-tagged T1	ATAAGAATGCGGCCGCCACTTTCCATTCAGCTGTGCCCAT
human_PRRC2B_300_R	FLAG-tagged T2	ATAAGAATGCGGCCGCCATCGTTCCACTGTACCCTGGTT
human_PRRC2B_450_R	FLAG-tagged T3	ATAAGAATGCGGCCGCCAGTCTGGCGCCTTTCGGAC
human_PRRC2B_600_R	FLAG-tagged T4	ATAAGAATGCGGCCGCCACTGTGCCACTGCTGGGGA
human_PRRC2B_750_F	FLAG-tagged P2	CGGGGTACCGCCACCATGGCACTGCAGAGCAAGGGC
human_PRRC2B_1500_R	FLAG-tagged P2	ATAAGAATGCGGCCGCCAACTGGCATGGGCTGCCCGC
human_PRRC2B_1500_F	FLAG-tagged P3	CGGGGTACCGCCACCATGGCTGACCTTCCCGAAGCCTCC
human_PRRC2B_2230_R	FLAG-tagged P3, full-length	ATAAGAATGCGGCCGCCAGGCCTTACTCTCCTCCACTTTGACAG
CCND2_UTR5_F	WT-FLuc	AAGCTAATACGACTCACTATAGGCAGCCCCGAGGCTCTG
CCND2_UTR5_R	WT-FLuc	CCTTTGGCTAAATAGGGGGTTTT
CCND2_UTR5_mutant_R	Del-FLuc, T2A-FLuc, C2G-FLuc, CT2GA-FLuc	GCCTCGGGGCTGCCTC
CCND2_UTR5_T2A_F	T2A-FLuc	ACAGCACGCCCACCACCCAAACCACGCCACCCAACAGCACCACCAACACACACAGCCCACACCACACCCCCG AAAACCCCCTATTTAGCCAAAGG

CCND2_UTR5_C2G_F	C2G-FLuc	TGTGTGGGGGAGGAGGGA ATGGTGGGGTGGGTTGTGG TGGAGGTTGTGTGTGTGGG GTGAGGTGTGGGGGG AAAACCCCCTATTTAGCCAA AGG
CCND2_UTR5_Del_F	Del-FLuc	AAGGAGGTCAGGGGAACG CTCTCCCCTCCCCTTCCAA AAAACAAAAACAGAAAAACC TTTTTCCAGGCCGGG AAAACCCCCTATTTAGCCAA AGG
CCND2_UTR5_CT2GA_ F	CT2GA-Fluc	AGAGGAGGGGGAGGAGGG AAAGGAGGGGGAGGGAAGA GGAGGAGGAAGAGAGAGA GGGGAGAGGAGAGGGGGG AAAACCCCCTATTTAGCCAA AGG
RT-qPCR primers (F, forward; R, reverse)	**Targeted transcript (RefSeq)**	**Sequence (5' - 3')**
CCND2_F	NM_001759.4	CTCTGCTGAGCGGTACTAA AC
CCND2_R	NM_001759.4	CTCCCTTCAACTATCATCCC ATAC
CRKL_F	NM_005207.4	CACACGGAAAGCATGGAAA TAG
CRKL_R	NM_005207.4	AAACTGCAGGTAGAGGAGT TG
YWHAZ_F	NM_145690.3	AGCAGAGAGCAAAGTCTTC TATT
YWHAZ_R	NM_145690.3	GACTGATCGACAATCCCTTT CT
GAPDH_F	NM_002046.7	GACCACTTTGTCAAGCTCAT TTC
GAPDH_R	NM_002046.7	CTCTCTTCCTCTTGTGCTCT TG
GATA4_F	NM_002052.5	TTCCAGCAACTCCAGCAAC G
GATA4_R	NM_002052.5	GCTGCTGTGCCCGTAGTGA G

ACTB_F	NM_001101.5	GGATCAGCAAGCAGGAGTATG
ACTB_R	NM_001101.5	AGAAAGGGTGTAACGCAACTAA
CTC1_F	NM_025099.6	CTCTTCTGGGTGTGTGCTTATT
CTC1_R	NM_025099.6	CTGATGATGGCCTGGCTTATAG
ATAD5_F	NM_024857.5	AGCTGATCCTGTCCCTAGTT
ATAD5_R	NM_024857.5	TAGGAAGTCACGCTTTGCTTTA
SERBP1_F	NM_015640.4	GTGACTGAGGAAACACCTGAA
SERBP1_R	NM_015640.4	AGCCTTCCACTCATCCAAAG
BPTF_F	NM_182641.4	TTATCCGGAGGAGATGGAAGA
BPTF_R	NM_182641.4	GAGTGCTGCTGTAGGTACTATG
PPP2R1A_F	NM_014225.6	ATGGGTCTCTCTCCCATCTT
PPP2R1A_R	NM_014225.6	GTTCACACAGTCCAGGTTAGAG
RPL35_F	NM_007209.4	GGAGAACCTGAAGACCAAGAAG
RPL35_R	NM_007209.4	CTCAGCCAGCTGTGCTTTAT
RPLP1_F	NM_001003.3	GCTGTTGGTCTTGTCCATAGT
RPLP1_R	NM_001003.3	GCCTGTGTCATGGCAGAATA
RPL38_F	NM_000999.4	GGGAGGCCATGATGGAAATA
RPL38_R	NM_000999.4	GCGAGTGGTAGATGTGCTATAC

PTBP1_F	NM_002819.5	GACGGCATTGTCCCAGATATAG
PTBP1_R	NM_002819.5	GGTCCGTTAGTGACACAAGTAG
HSPA4_F	NM_002154.4	CATTGCAGTGTGCCATCTTATC
HSPA4_R	NM_002154.4	CACAGTCACTTGACCCTTCTT
RPL37A_F	NM_000998.5	ATTACTTGAGGCCAGGAGATTG
RPL37A_R	NM_000998.5	CCTCCAAAGTAGCTGGGATTAC

ASO	Effect	Sequence (5' - 3')
ctrl ASO	control ASO with no target in eukaryotic cells	CmCmAmGmGmAmUmUmCmAmAmCmCmUmAmoCm
ASO1	ASO targeting PRRC2B-binding sites on 5' UTR of *CCND2* mRNA	UmUmAmGmGmAmGmCmGmGmAmGmGmGmAmAmGmAm
ASO2	ASO targeting an adjacent region of PRRC2B-binding sites on 5' UTR of *CCND2* mRNA	CmCmUmCmCmUmUmCmCmUmUmUmGmGmCmUm

Note: "m" indicates a 2'-O-methyl modification.

Antibodies	Cat. #	Dilution
Mouse anti-β-actin	MA5-11869; Invitrogen	IB (1: 2000)
Mouse anti-DAP5 (EIF4G2)	135999; Santa Cruz	IB (1: 2000)
Donkey anti-rabbit IgG	NA9340V; Roche	IP (1: 100)
Rabbit anti-eIF3G Polyclonal antibody	11165-1-AP; Proteintech	IB (1:1000)
Mouse monoclonal anti-eIF3D	sc-271515; Santa Cruz	IB (1:500)
Mouse monoclonal anti-eIF4G1	sc-133155; Santa Cruz	IB (1:500)

Mouse monoclonal anti-eIF4E	sc-9976; Santa Cruz	IB (1:500)
Mouse monoclonal anti-FXR1	sc-137418; Santa Cruz	IB (1:500)
Rabbit polyclonal anti-PRRC2B	PA5-66677; Thermo Fisher Scientific	IB (1:1000); IP (1:125)
Mouse polyclonal anti-RPLP0	sc-293260; Santa Cruz	IB (1:1000)
Rabbit polyclonal anti-RPS6	14823-1-AP; Proteintech	IB (1:1000)
Pierce™ Anti-DYKDDDDK Magnetic Agarose	A36797; Thermo Fisher Scientific	IP (1:50)
Mouse monoclonal anti-CRKL	sc-365092; Santa Cruz	IB (1:500)
Rabbit polyclonal anti-YWHAZ	A7639; Abclonal	IB (1:1000)
Rabbit polyclonal anti-cyclin D2	A1773; Abclonal	IB (1:1000)
Cell lines	**Cat. #**	
HEK293T	CRL-3216; ATCC	
Chemicals & reagents	**Cat. #**	
Ethanol	V1016; Decon Labs	
Chloroform	C2432-500ML; Millipore-Sigma	
RIPA Lysis and Extraction Buffer	89900; Thermo Fisher Scientific	
[gamma-32P]-ATP (250 mCi)	NEG502A250UC; PerkinElmer	
HEPES-KOH	BB-2076-K; Boston BioProducts	
Potassium chloride	P3911-25G; Millipore-Sigma	

MgCl$_2$	M8266-100G; Millipore-Sigma	
Sodium Acetate 3M, pH 5.2	41920020; Bio-world	
Dithiothreitol	D0632-10G; Millipore-Sigma	
IGEPAL® CA-630	I8896-50ML; Millipore-Sigma	
Vanadyl ribonucleoside complexes	94740-250MG; Millipore-Sigma	
RNasin® Ribonuclease Inhibitor	N2611; Promega	
iScript™ cDNA Synthesis Kit	1708890; Bio-Rad	
cOmplete™ Protease Inhibitor Cocktail	11697498001; Millipore-Sigma	
iTaq Universal SYBR Green Supermix	1725124, Bio-Rad	
Picrosirius red stain	ab150681; Abcam	
TrueCut Cas9	A36496; Thermo Fisher Scientific	
Trypsin-EDTA (0.25%), phenol red	25200056; Thermo Fisher Scientific	
6-well cell culture plates	140675; Thermo Fisher	
10-cm cell culture plates	353002; Corning	
OPTI-MEM	31985062; Thermo Fisher Scientific	
Immobilon®-P PVDF Membrane	IPVH00010; Millipore-Sigma	
Lipofectamine 3000	L3000015; Thermo Fisher Scientific	
Superscript III	18080093; Thermo Fisher Scientific	

Dual-Glo Luciferase Assay System	E2920; Promega	
Bsmb1 restriction enzyme	R0739S; NEB	
Kpn1 restriction enzyme	R3142S; NEB	
Not1 restriction enzyme	R3189S; NEB	
Quick CIP	M0525S; NEB	
RNase T1 (1000 U/µL)	EN0542; Thermo Fisher Scientific	
T4 DNA Ligase	M0202S; NEB	
T4 PNK	M0201S; NEB	
Dpn1 restriction enzyme	R0176S; NEB	
Proteinase K	3115879001; Millipore Sigma	
GlycoBlue™ Coprecipitant (15 mg/mL)	AM9515; Thermo Fisher Scientific	
Acid-Phenol: Chloroform, pH 4.5 (with IAA, 125:24:1)	AM9722; Thermo Fisher Scientific	
RNA Gel Loading Dye (2X)	R0641; Thermo Fisher Scientific	
NuPage transfer buffer 20X	NP00061; Thermo Fisher Scientific	
UltraPure Phenol:Chloroform: Isoamyl Alcohol (25:24:1, v/v)	15593031; Thermo Fisher	
Trizol	15596026; Thermo Fisher	
TRIPZ Inducible Lentiviral shRNA targeting PRRC2B	V3THS_374329; Dharmacon	

4-Thiouridine	T4509; Sigma-Aldrich	
Cycloheximide	C7698; Sigma-Aldrich	
siRNA smartpool (eIF4G2)	L-011263-00-0005; Dharmaco	
Plasmids	**Cat. #**	
pGL3-TK-5UTR-BsmBI-Luciferase	114670; Addgene	
pRL Renilla Luciferase Control Reporter	E2231; Promega	
pCDNA3.1(+)-3xFLAG	Homemade	