TITLE

An important role for triglyceride in regulating spermatogenesis

AUTHORS

Charlotte F. Chao[1&], Yanina-Yasmin Pesch[1&], Huaxu Yu[2], Chenjingyi Wang[2], Maria

Aristizabal[3], Tao Huan[2], Guy Tanentzapf[1], Elizabeth J. Rideout[1*]

AFFILIATIONS

[1] Department of Cellular and Physiological Sciences, Life Sciences Institute, The

University of British Columbia, Vancouver, BC, Canada V6T 1Z3

[2] Department of Chemistry, The University of British Columbia, Vancouver, BC, Canada

V6T 1Z1

[3] Department of Biology, Queen's University, Kingston, ON, Canada K7L 3N6

* Corresponding author

[&] Equally contributing authors

[*] Correspondence: elizabeth.rideout@ubc.ca

ABSTRACT

Drosophila is a powerful model to study how lipids affect spermatogenesis. Yet, the

contribution of neutral lipids, a major lipid group which resides in organelles called lipid

droplets (LD), to normal sperm development is largely unknown. Emerging evidence

suggests that LD are present in the testis and that loss of neutral lipid- and LD-

associated genes causes subfertility; however, key regulators of testis neutral lipids and

LD remain unclear. Here, we show that LD are present in early-stage somatic and

germline cells within the *Drosophila* testis. We identified a role for triglyceride lipase

brummer (*bmm*) in regulating testis LD, and found that whole-body loss of *bmm* leads to

defects in sperm development. Importantly, these represent cell-autonomous roles for

bmm in regulating testis LD and spermatogenesis. Because lipidomic analysis of *bmm*

mutants revealed excess triglyceride accumulation, and spermatogenic defects in *bmm*

mutants were rescued by genetically blocking triglyceride synthesis, our data suggest

that *bmm*-mediated regulation of triglyceride influences sperm development. This

identifies triglyceride as an important neutral lipid that contributes to *Drosophila* sperm

development, and reveals a key role for *bmm* in regulating testis triglyceride levels

during spermatogenesis.

KEYWORDS

Triglyceride, lipid droplet, spermatogenesis, testis, *brummer*, adipose triglyceride lipase

(ATGL), lipidomics.

INTRODUCTION

Lipids play an essential role in regulating spermatogenesis across animals [1–4].

Studies in *Drosophila* have illuminated key roles for multiple lipid species in regulating

sperm development [5–7]. For example, phosphatidylinositol and its phosphorylated

derivatives participate in diverse aspects of *Drosophila* spermatogenesis including

meiotic cytokinesis [1,8–11], somatic cell differentiation [12], germline and somatic cell

polarity maintenance [13–16], and germline stem cell (GSC) maintenance and

proliferation [17]. Membrane lipids also influence sperm development [18,19], whereas

fatty acids play a role in processes such as meiotic cytokinesis [20] and sperm

individualization [21,22]. While these studies suggest key roles for membrane lipids and

fatty acids during *Drosophila* spermatogenesis, some of which are conserved in

mammals [23–25], much less is known about how neutral lipids contribute to

spermatogenesis.

Neutral lipids are a major lipid group that includes triglyceride and cholesterol

ester, and reside within specialized organelles called lipid droplets (LD) [26]. LD are

found in diverse cell types (*e.g.* adipocytes, muscle, liver, glia, neurons) [27,28,26], and

play key roles in maintaining cellular lipid homeostasis. In nongonadal cell types, correct

regulation of LD contributes to cellular energy production [29–31], sequestration and

redistribution of lipid precursors [32–36], and regulation of lipid toxicity [37–39]. The

importance of LD to normal cellular function in nongonadal cell types is shown by the

fact that dysregulation of LD causes defects in cell differentiation, survival, and energy

production [26,37,40,41]. In the testis, much less is known about the regulation and

function of neutral lipids and LD, and how this regulation affects sperm development.

Multiple lines of evidence suggest a potential role for neutral lipids and LD during

spermatogenesis. First, genes that encode proteins associated with neutral lipid

metabolism and LD are expressed in the testis across multiple species [42–44].

Second, testis LD have been identified in mammals and flies under both normal

physiological conditions [27,44–48] and after mitochondrial stress [49]. Third, loss of

genes associated with neutral lipid metabolism and LD cause subfertility phenotypes in

both flies and mammals [27,50–52]. While studies suggest that mammalian testis LD

contribute to steroidogenesis [53,54], the spatial, temporal, and cell-type specific

requirements for neutral lipids and LD in the testis have not been explored in detail in

any animal. It remains similarly unclear which genes are responsible for regulating

neutral lipids and LD during spermatogenesis.

To address these knowledge gaps, we used *Drosophila* to investigate the

regulation and function of neutral lipids and LD during sperm development. Our detailed

analysis of spermatogenesis under normal physiological conditions revealed the

presence of LD in early-stage somatic and germline cells in the testis. We identified

triglyceride lipase *brummer* (*bmm*) as a regulator of testis LD, and showed that this

represents a cell-autonomous role for *bmm*. Importantly, we found that the *bmm*-

mediated regulation of testis LD was significant for spermatogenesis, as both whole-

body and cell-autonomous loss of *bmm* caused defects in sperm development. Given

that our lipidomic analysis revealed an excess accumulation of triglyceride in animals

lacking *bmm*, and that genetically blocking triglyceride synthesis rescued many

spermatogenic defects associated with *bmm* loss, our data suggests that *bmm*-

mediated regulation of triglyceride is important for normal *Drosophila* sperm

development. This reveals previously unrecognized roles for neutral lipids such as

triglyceride in regulating spermatogenesis, and for *bmm* in regulating sperm

development under normal physiological conditions. Together, these findings advance

knowledge of the regulation and function of neutral lipids during spermatogenesis.

RESULTS

Lipid droplets are present in early-stage somatic and germline cells

We previously reported the presence of small circular punctae (<1 μm) corresponding to

LD near the apical tip of the testis [27]. We confirm these results in w^{1118} males using

neutral lipid stain BODIPY (4,4-Difluoro-1,3,5,7,8-Pentamethyl-4-Bora-3a,4a-Diaza-*s*-

Indacene) (Figure 1A). Importantly, we reproduced this spatial distribution of LD in two

independent genetic backgrounds and at two additional ages (Figure 1B,1C). In all

cases, LD were in a testis region that contains stem cells and early-stage somatic and

germline cells (Figure 1A-A', arrows), and in the hub, an organizing center and stem cell

niche in the *Drosophila* testis (Figure 1A''-A''', arrows) [55]. LD were largely absent from

the testis region occupied by spermatocytes (Figure 1A and A', arrowheads). While LD

may contain multiple neutral lipid species[56], cholesterol-binding fluorescent polyene

antibiotic filipin III did not detect cholesterol within testis LD (Figure S1A), suggesting

triglyceride is the main neutral lipid in *Drosophila* testis LD.

Drosophila spermatogenesis requires the codevelopment and differentiation of

two cell lineages, the germline and the somatic cells [57]. To identify LD in each lineage,

we used the GAL4/UAS system to overexpress a transgene in which GFP is fused to

the LD-targeting motif of motor protein Klarsicht [58] (*UAS-GFP-LD*). We targeted *UAS-*

GFP-LD to somatic cells with *Traffic jam* (*Tj*)-*GAL4* and to the germline using *nanos*

(*nos*)-*GAL4*; LD were visualized using neutral lipid dye LipidTox. We found LD in the

somatic cells of 0-day-old males (Figure 1D), and showed that the majority of somatic

LD were <30 µm from the hub (Figure 1E). Because the somatic LD distribution

coincided with a marker for somatic stem cells and their immediate daughter cells (Zinc

finger homeodomain 1, Zfh-1) (Figure 1F; two-sample Kolmogorov-Smirnov test) [59],

but not with a marker for late somatic cells (Eyes absent, Eya) [12,60], our data

suggests LD are present in early somatic cells. In the germline, GFP punctae

corresponding to LD were found near the apical tip of the testis in 0-day-old males

(Figure 1G,H). We found that the disappearance of germline LD coincided with peak

expression of a GFP reporter that reflects the expression of Bag-of-marbles (Bam)

protein in the testis (Bam-GFP) [61] (Figure 1I,1J). Because peak Bam expression

signals the last round of transient amplifying mitotic cell cycle prior to the germline's

transition into the meiotic cell cycle [62–64], our data suggests that germline LD, like

somatic LD, are present at early stages of germline development.

brummer plays a cell-autonomous role in regulating testis lipid droplets

Adipose triglyceride lipase (*ATGL*) is a critical regulator of neutral lipid metabolism and

LD [65–74]. Loss of *ATGL* in many cell types triggers LD accumulation, and *ATGL*

overexpression decreases LD number [30,67,68,71,73,73,75,76]. Given that the

Drosophila ATGL homolog *brummer* (*bmm*) regulates testis LD induced by

mitochondrial stress [49], we explored whether *bmm* regulates testis LD under normal

physiological conditions. We first examined *bmm* expression in the testis by isolating

this organ from flies in which a *bmm* promoter fragment drives *GFP* expression (*bmm-*

GFP). Indeed, *bmm-GFP* accurately reproduces changes to *bmm* mRNA levels [77].

GFP expression was present in the germline of *bmm-GFP* testes, and we found

germline GFP levels were higher in spermatocytes than at earlier stages of sperm

development (Figure 2A,2B; one-way ANOVA with Tukey multiple comparison test).

Supporting this, our analysis of a publicly available single-cell RNA sequencing data set

of the male reproductive organ [78] suggested a similar trend in *bmm* mRNA levels

between different stages of germline (Figure S2A,S2B) and somatic cell (Figure

S2C,S2D) development. Importantly, germline GFP levels were negatively correlated

with testis LD in *bmm-GFP* flies (Figure 2A,2C), suggesting regions with higher *bmm*

expression had fewer LD.

To test whether *bmm* regulates testis LD, we compared LD in testes from 0-day-

old males carrying a loss-of-function mutation in *bmm* (*bmm^1*) to control male testes

(*bmm^{rev}*)[67]. *bmm^1* testes had significantly more LD across all LD sizes compared with

control males (Figure 2D–2G; Welch two-sample t-test with Bonferroni correction), and

showed a significantly expanded LD distribution (Figure 2D–2F,2H; two-sample

Kolmogorov-Smirnov test). This suggests *bmm* normally restricts LD to the region near

the apical tip of the testis, a role we confirm in both somatic and germline lineages

(Figure S2E–S2H). Importantly, after inducing homozygous *bmm^1* or *bmm^{rev}* clones in

the testes using *FLP-FRT* system[79], we found *bmm^1* spermatocyte clones had

significantly more LD at 3 days post-clone induction (Figure 2I; Welch two-sample t-

test), a stage at which LD were absent from *bmm^{rev}* clones. This indicates a previously

unrecognized cell-autonomous role for *bmm* in regulating testis LD, a role we were

unable to assess in somatic cells as we recovered no *bmm^1* somatic cell clones.

***brummer* plays a cell-autonomous role in regulating germline development**

To determine the physiological significance of *bmm*-mediated regulation of testis LD, we

investigated testis and sperm development in males without *bmm* function. In 0-day-old

bmm^1 males reared at 25°C, testis size was significantly smaller than in age-matched

bmm^{rev} controls (Figure S3A; Welch two-sample t-test), and the number of spermatid

bundles was significantly lower (Figure S3B; Kruskal-Wallis rank sum test). Defects in

testis size and sperm development were also observed in 14-day-old *bmm^1* males

(Figure S3C,S3D Welch two-sample t-test). When the animals were reared at 29°C, a

temperature that exacerbates spermatogenesis defects associated with changes in lipid

metabolism [21], *bmm^1* phenotypes were more pronounced (Figure 3A-3C). This

suggests loss of *bmm* affects testis development and spermatogenesis. Because similar

phenotypes are observed in male mice without ATGL [52], and supplementing the diet

of *bmm^1* males with medium-chain triglycerides (MCT) partially rescues the testis and

spermatogenic defects we observed in flies (Figure S3E,S3F; one-way ANOVA with

Tukey multiple comparison test), as it does in mice [52,80], our data suggests flies are a

good model to study how *bmm*/*ATGL* influences sperm development.

To explore spermatogenesis in *bmm^1* animals, we used germline-specific marker

Vasa to visualize the germline in the testes of *bmm^1* and *bmm^{rev}* males (Figure 3D,3E)

[81]. We observed a significant increase in the number of germline stem cells (GSC)

(Figure 3F; Kruskal-Wallis rank sum test) and higher variability in GSC number in *bmm^1*

males (p=5.7×10^{-12} by F-test). Given that GSC number is affected by hub size and GSC

proliferation [82,83], we monitored both parameters in bmm^1 and bmm^{rev} controls. While

hub size in bmm^1 testes was significantly larger than in testes from bmm^{rev} controls

(Figure S3G,S3H; Welch two-sample t-test), the number of phosphohistone H3-positive

GSC, which indicates proliferating GSC, was unchanged in bmm^1 animals (Figure S3I;

Kruskal-Wallis rank sum test). While this indicates a larger hub may partly explain

bmm's effect on GSC number, bmm also plays a cell-autonomous role in regulating

GSC, as we recovered a higher proportion of bmm^1 clones in the GSC pool compared

with bmm^{rev} clones at 14 days after clone induction (Figure 3G; Welch two-sample t-

test).

Beyond GSC, we uncovered additional spermatogenesis defects in bmm^1 testes.

Peak Bam-GFP expression in testes from 0-day-old bmm^1 and bmm^{rev} males showed

that GFP-positive cysts with were significantly further away from the hub in bmm^1 testes

(Figure 3H,S3J; Welch two-sample t-test). Indeed, 15/18 bmm^1 testes contained Vasa-

positive cysts with large nuclei in the distal half of the testis (Figure 3I, arrowheads), a

phenotype not present in bmm^{rev} testes (0/8) (p=0.0005 by Pearson's Chi-square test).

Because these phenotypes are also seen in testes with differentiation defects [13,84],

we recorded the stage of sperm development reached by the germline in bmm^1 testes.

Most bmm^1 testes contained post-meiotic cells in males raised at 25°C (Figure S3K);

however, germline development did not progress past the spermatocyte stage in most

bmm^1 testes from animals raised at 29°C (Figure S3K). Testes from bmm^1 males reared

at 25°C also had a smaller Boule-positive area (Figure 3J,S3L; Welch two-sample t-

test), and fewer individualization complexes and waste bags (Figure S3M,S3N; Kruskal-

Wallis rank sum test). Together, these data indicate loss of *bmm* delays germline

development. Because we recovered fewer *bmm^1* spermatocyte and spermatid clones

14 days after clone induction (Figure 3K,3L; Kruskal-Wallis rank sum test), this effect on

germline development represents a cell-autonomous role for *bmm*.

brummer-dependent regulation of testis triglyceride levels affects

spermatogenesis

ATGL catalyzes the first and rate-limiting step of triglyceride hydrolysis [73,85,86]. Loss

of this enzyme or its homologs leads to excess triglyceride accumulation

[27,30,67,73,75] and shifts in multiple lipid classes [66,87–89]. To determine how loss of

bmm affects spermatogenesis, we carried out mass spectrometry (MS)-based

untargeted lipidomic profiling of *bmm^1* and *bmm^rev* males. Hierarchical clustering of lipid

species suggests that *bmm^1* and *bmm^rev* males show distinct lipidomic profiles (Figure

4A). Overall, we detected 2464 and 1144 lipid features with high quantitative confidence

in positive and negative ion modes, respectively. By matching experimental *m/z*,

isotopic ratio, and tandem MS spectra to lipid libraries, we confirmed 293 unique lipid

species (Supplemental table 1). We found 107 lipids had a significant change in

abundance between *bmm^1* and *bmm^rev* males ($p_{adj}<0.05$): 85 species were upregulated

in *bmm^1* males and 22 lipid species were downregulated. Among differentially regulated

species from different lipid classes, triglyceride had the largest residual above expected

proportion ($p=5.00\times10^{-4}$ by Pearson's Chi-squared test). This suggests triglyceride is the

lipid class most affected by loss of *bmm* (Figure 4B,4C).

In *bmm¹* males, most triglyceride species (55/97) were significantly higher.

Because we observed a positive correlation between the fold increase in triglyceride

abundance with both the number of double bonds (p=7.52×10⁻⁸ by Kendall's rank

correlation test; Figure S4A) and the number of carbons (p=2.77×10⁻¹⁰ by Kendall's rank

correlation test; Figure S4B), our data align well with *bmm/ATGL's* known role in

regulating triglyceride levels[67,68,73] and its substrate preference of long-chain

polyunsaturated fatty acids[85]. While we also detected changes in species such as

fatty acids, acylcarnitine, and membrane lipids (Figure S4C–S4H), in line with recent

Drosophila lipidomic data[90,91], the striking accumulation of triglyceride in *bmm¹* males

suggested that excess testis triglyceride in *bmm¹* males may contribute to their

spermatogenic defects. To test this, we examined spermatogenesis in *bmm¹* males

carrying loss-of-function mutations in *midway* (*mdy*). *mdy* is the *Drosophila* homolog of

diacylglycerol O-acyltransferase 1 (*DGAT1*), and whole-body loss of *mdy* reduces

whole-body triglyceride levels[92–94]. Importantly, testes isolated from males lacking

both *bmm* and *mdy* (genotype *mdy*^{QX25/k03902}*;bmm¹*) had fewer LD than testes dissected

from *bmm¹* males (Figures 4D,S4I; one-way ANOVA with Tukey multiple comparison

test).

We found that testes isolated from *mdy*^{QX25/k03902}*;bmm¹* males were significantly

larger and had more spermatid bundles than testes from *bmm¹* males (Figure 4E–G;

one-way ANOVA with Tukey multiple comparison test). The elevated number of GSC in

bmm¹ male testes was similarly rescued in *mdy*^{QX25/k03902}*;bmm¹* males (Figure 4H; one-

way ANOVA with Tukey multiple comparison test). These data suggest that defective

spermatogenesis in *bmm¹* males can be partly attributed to excess triglyceride

accumulation. Notably, at least some of these defects are cell-autonomous: RNAi-

mediated knockdown of *mdy* in the germline of *bmm*[1] males partially rescued the

defects in testis size (Figure 4I; Kruskal-Wallis rank sum test with Dunn's multiple

comparison test) and GSC variance (Figure S4J; *p*=4.5 x 10^{-5} and 8.2 x 10^{-3} by F-test

from the GAL4- and UAS-only crosses, respectively). *bmm*-mediated regulation of testis

triglyceride therefore plays a previously unrecognized role in regulating sperm

development.

DISCUSSION

In this study, we used *Drosophila* to gain insight into how the neutral lipids, a major lipid

class, contribute to sperm development. We describe the distribution of LD under

normal physiological conditions in the *Drosophila* testis, and show that LD are present

at the early stages of development in both somatic and germline cells. While many

factors are known to regulate LD in nongonadal cell types, we reveal a cell-autonomous

role for triglyceride lipase *bmm* in regulating testis LD during spermatogenesis. Indeed,

our data indicates loss of *bmm* delays germline differentiation leading to an

accumulation of early-stage germ cells. These defects in germline differentiation can be

partially explained by the excess accumulation of triglyceride in flies lacking *bmm*, as

genetically blocking triglyceride synthesis rescues multiple spermatogenic defects in

bmm mutants. Together, our data reveals previously unrecognized roles for LD and

triglycerides during spermatogenesis, and for *bmm* as an important regulator of testis

LD and germline development under normal physiological conditions.

One key outcome of our study was increased knowledge of LD regulation and

function in the testis. Despite rapidly expanding knowledge of LD in cell types such as

adipocytes or skeletal muscle, less is known about how LD influence spermatogenesis

under normal physiological conditions. In mammals, testis LD contain cholesterol and

play a role in promoting steroidogenesis [95,96]. In flies, we show that LD are present in

the testis, and that excess accumulation of these LD affects sperm development. In

nongonadal cell types, triglycerides provide a rich source of fatty acids for cellular ATP

production, lipid building blocks to support membrane homeostasis and growth, and

metabolites that can act as signaling molecules [26]. Because ATP production, lipid

precursors, and lipid signaling all play roles in supporting normal sperm development

[97,98], future studies will need to determine how each of these processes is affected

when excess triglyceride accumulates in testis LD. This will provide critical insight into

how triglyceride stored within testis LD contributes to overall cellular lipid metabolism

during spermatogenesis. Because of the parallel spermatogenic defects we observed in

bmm mutants and *ATGL*-deficient mice, we expect that these mechanisms will also

operate in other species.

A more comprehensive understanding of neutral lipid metabolism during sperm

development will also emerge from studies on the upstream signaling networks that

regulate testis LD and triglyceride. Given that we show an important and cell-

autonomous role for *bmm* in regulating testis LD and triglyceride, future studies will

need to identify factors that regulate *bmm* in the testis. Based on public single-cell

RNAseq data and the *bmm-GFP* reporter strain, our data suggest *bmm* mRNA levels

are differentially regulated between early and later stages of sperm development.

Candidates for mediating this regulation include the insulin/insulin-like growth factor

signaling pathway (IIS), Target of rapamycin (TOR) pathway, and nuclear factor

κB/Relish pathway (NFκB), as all of these pathways influence *bmm* mRNA levels in

nongonadal cell types [99–105]. Beyond mRNA levels, Bmm protein levels and post-

translational modifications may also be differentially regulating during spermatogenesis.

For example, studies show that the proteins encoded by *bmm* homologs in other

animals are regulated by phosphorylation [106], mediated by kinases such as

adenosine monophosphate-activated protein kinase (AMPK) and protein kinase A

(PKA) [107–109]. Importantly, many of these pathways, including IIS, TOR, AMPK,

NFκB and possibly PKA influence *Drosophila* sperm development [110–115]. Identifying

the signaling networks that influence *bmm* regulation during sperm development will

therefore lead to a deeper understanding of how testis LD and triglyceride are

coordinated with physiological factors to promote normal spermatogenesis. Because

pathways such as IIS and AMPK, and others, regulate sperm development in other

species [116–118], these insights may reveal conserved mechanisms that govern the

regulation of cellular neutral lipid metabolism during sperm development.

ACKNOWLEDGEMENTS

We thank Dr. Ronald Kühnlein for *bmm*[1] and *bmm*[rev] lines [67], Dr. Michael Welte for

UAS-GFP-LD[58], and Dr. Kaeko Kamei for *bmm-GFP* [77]. We used stocks from the

Bloomington *Drosophila* Stock Center (NIH P40OD018537) and Vienna *Drosophila*

Resource Center (VDRC). We acknowledge critical resources and information provided

by FlyBase [119] (supported by the National Human Genome Research Institute at the

U.S. National Institutes of Health (U41 HG000739) and the British Medical Research

Council (MR/N030117/1)). This work was supported by the Life Sciences Institutes

Imaging Core, supported by the UBC GREx Biological Resilience Initiative. Funding for

this study was provided by grants to EJR from the Canadian Institutes for Health

Research (PJT-153072), Michael Smith Foundation for Health Research (16876), and

the Canadian Foundation for Innovation (JELF-34879). GT was supported by a grant

from the Natural Sciences and Engineering Research Council (NSERC; 2018-04648),

TH/HY/CW were supported by NSERC (2020-04895), MA was supported by the

Jacob's foundation. We would like to acknowledge that our research takes place on the

traditional, ancestral, and unceded territory of the Musqueam people; a privilege for

which we are grateful.

AUTHOR CONTRIBUTIONS

Conceptualization, C.C. and E.J.R.; Methodology, C.C. and Y.Y.P.; Software, C.C.;

Investigation, C.C., H.Y., and Y.Y.P.; Lipidomics, M.A., H.Y., C.W., T.H.; Writing –

Original Draft, C.C. and E.J.R.; Writing – Review and Editing, C.C., E.J.R., and Y.Y.P.;

Supervision, E.J.R., G.T., and T.H.; Project administration, E.J.R.; Funding Acquisition,

E.J.R., G.T., and T.H.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1 – Lipid droplets are present in early-stage somatic and germ cells. (A)

Testis lipid droplets (LD) in w^{1118} animals visualized with neutral lipid dye BODIPY.

(A,A') Scale bar=50 µm; (A",A"') scale bar=15 µm. Asterisk indicates hub in all images.

Arrows point to LD; arrowheads point to spermatocytes in A,B. (B) Testis LD visualized

with BODIPY in newly-eclosed males from two wild-type genotypes. Scale bars: main

image=50 µm; inset image=10 µm. (C) Testis LD from w^{1118} animals at different times

post-eclosion. Scale bars=50 µm. (D) Testis LD visualized with LipidTox Red in animals

with somatic cell overexpression of GFP-LD (*Tj-GAL4>UAS-GFP-LD*). GFP- and

LipidTox Red-positive punctae are somatic LD (D–D" arrows); LipidTox punctae without

GFP indicate germline LD (D–D" arrowheads). Scale bars=10 µm. (E) Histogram

showing the spatial distribution of somatic cell LD; error bars represent standard error of

the mean (SEM). (F) Cumulative frequency distributions of somatic LD (blue line, data

reproduced from E), zfh-1-positive somatic cells (zfh-1$^+$ cells, orange line), and Eya-

positive somatic cells (Eya$^+$ cells, grey line). (G) Testis LD visualized with LipidTox Red

in males with germline overexpression of GFP-LD (*nos-GAL4>UAS-GFP-LD*). GFP- and

LipidTox Red-positive punctae indicate germline LD (arrows); LipidTox punctae without

GFP indicate non-germline LD (arrowheads). Scale bars=10 µm. (H) Histogram

representing the spatial distribution of LD within the germline; error bars represent SEM.

(I) Histogram representing the spatial distribution of LD and GFP fluorescence (green

line) (arbitrary units, a.u.) in a representative testis of a *bam-GFP* animal (panel J). (J)

Testis LD in a *bam-GFP* animal; arrows point to LD and arrowheads point to

spermatocytes. Scale bar=50 µm. See also Supplemental Figure 1.

Figure 2 – *bmm* regulates testis lipid droplets in a cell-autonomous manner. (A)

Testis lipid droplets (LD) indicated by LipidTox Red in *bmm-GFP* animals. Arrows point

to LD in all images. Arrowheads point to spermatocytes. Scale bars=50 µm. Asterisks

indicate the hub in all images. (B) Quantification of nuclear GFP intensity in testes

isolated from *bmm-GFP* animals (n=3). Germline stem cell (GSC), spermatogonia (SG),

spermatocyte (SC). (C) Spatial distribution of LD (grey histogram) and GFP expression

(green line) in testes from *bmm-GFP* animals as a function of distance from the hub

(n=3). (D,E) LD near the apical region of the testis in *bmm^{rev}* (D) or *bmm^1* (E) animals.

(F) LD further away from the apical tip in *bmm^1* animals. (D–F) Scale bars=50 µm. (G)

Histogram representing testis LD size distribution in *bmm^{rev}* (grey) and *bmm^1* (orange).

(H) Apical tip of the testes is at the left of the graph; individual dots represent a single

LD and its relative position to the hub marked by an asterisk. Cumulative frequency

distribution of the distance between LDs and the apical tip of the testes are drawn as

solid lines. (I) Number of testis LD in *bmm^{rev}* (grey) or *bmm^1* (orange) in FLP-FRT

clones 3 days post-clone induction; dots represent measurements from a single clone.

The number of cells in each cyst (CC) counted is indicated. See also Supplemental

Figure 2.

Figure 3 – A cell-autonomous role for *bmm* in regulating spermatogenesis. Testes

isolated from *bmm^rev* (A) and *bmm^1* (A') animals raised at 29°C stained with phalloidin.

Scale bars=100 μm. (B) Testis size in *bmm^1* and *bmm^rev* animals raised at 29°C. (C)

Spermatid bundle number in *bmm^1* and *bmm^rev* testes from animals reared at 29°C.

(D,E) Representative images of *bmm^rev* (D) or *bmm^1* (E) testes stained with DAPI and

anti-Vasa antibody. Arrows indicate germline stem cells (GSC). Scale bar=50 μm. The

hub is marked by an asterisk in all images. (F) GSC number in *bmm^1* and *bmm^rev*

testes. (G) Proportion of GSCs that were either *bmm^1* or *bmm^rev* clones at 3 and 14

days post-clone induction. (H) Representative images of *bmm^rev* (H) and *bmm^1* (H')

testes carrying *bam-GFP*; data quantified in Figure S3J. Arrows indicate regions with

high Bam-GFP. Scale bars=50 μm. (I) Representative images of *bmm^rev* (I) or *bmm^1*

(I',I'') testes stained with anti-Vasa antibody. Arrows indicate Vasa-positive cysts in

bmm^1 testis. Panel I'' is magnified from the boxed region in I'. (I,I') Scale bars=100 μm;

(I'') scale bar=50 μm. (J) Maximum projection of *bmm^rev* (J) or *bmm^1* (J') testes stained

with anti-Boule antibody (green) and DAPI (blue). Scale bars=100 μm. Number of *bmm^1*

and *bmm^rev* spermatocyte clones (K) or post-meiotic clones (L) at 3 and 14 days post-

clone induction. See also Supplemental Figure 3.

Figure 4 – Loss of *bmm* disrupts triglyceride homeostasis and leads to

spermatogenic defects. (A) Hierarchical clustering of lipid species detected in *bmm^rev*

and *bmm¹* animals. (B) Histograms showing the proportion of species in each lipid class

with different levels between *bmm¹* and *bmm^rev*. Numbers on histograms indicate the

number of species with differences in abundance. (C) Volcano plot showing fold change

in abundance of triglycerides (green; 97 species) and non-triglyceride lipids (grey; 186

species) in our dataset. (D) Arrows indicate testis LD stained with LipidTox Red in

bmm^rev (D), *bmm¹* (D'), or *mdy^{QX25/k03902}*; *bmm¹* (D'') animals. (E) Whole testes isolated

from *bmm^rev* (E), *bmm¹* (E'), or *mdy^{QX25/k03902}*;*bmm¹* (E'') animals stained with anti-Vasa

antibody (red) and DAPI (blue). Arrowheads indicate spermatid bundles. Scale

bars=100 µm. (F) Testis size in *bmm^rev*, *bmm¹*, and *mdy^{QX25/k03902}*;*bmm¹* animals.

Spermatid bundles (G) and number of germline stem cells (H) in *bmm^rev*, *bmm¹*, and

mdy^{QX25/k03902};*bmm¹* animals. (I) Testis size in animals with germline-specific *mdy*

knockdown (*nos-GAL4>mdy RNAi*; *bmm¹*) compared with controls (*nos-GAL4>+*; *bmm¹*

and +>*mdy RNAi*; *bmm¹*). See also Supplemental Figure 4.

MATERIALS AND METHODS

Materials and Resource availability. *Drosophila* strains and their source are listed in a Key Resources table. Further information and requests for resources and reagents should be directed to, and will be fulfilled by, lead contact Dr. Elizabeth J. Rideout (elizabeth.rideout@ubc.ca).

Data and Code availability. All raw data and results of statistical tests reported in this paper are located in Supplementary files 1-4. This paper does not report original code. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Fly husbandry. Fly stocks were maintained at room temperature in 12:12 hour light:dark cycle. Unless otherwise indicated, all flies were raised at 25°C with a density of 50 larvae per 10 mL fly media. Because this project examines sperm development, we used male flies in all experiments. Fly media contained 20.5 g sucrose (SU10, Snow Cap), 70.9 g Dextrose (SUG8, Snow Cap), 48.5 g cornmeal (AO18006, Snow Cap), 30.3 g baker's yeast (NB10, Snow Cap), 4.55 g agar (DR-820-25F, SciMart), 0.5 g calcium chloride dihydrate (CCL302.1, BioShop Canada), 0.5 g magnesium sulfate heptahydrate (MAG511.1, BioShop Canada), 4.9 mL propionic acids (P1386, Sigma-Aldrich), and 488 μL phosphoric acid (P5811, Sigma-Aldrich) per 1L of media. For diets with medium- or long-chain triglyceride, 4 g of coconut oil (medium chain triglyceride) or olive oil (long chain triglyceride) was added per 100 mL of media described above prior

to cooling. Males were collected and dissected within 24 hours of eclosion unless

otherwise indicated. Fixations were performed at room temperature with 4%

paraformaldehyde (CA11021-168, VWR) in PBS for 20 minutes on a rotating platform

followed by washing in PBS twice before staining. Fly strains used in our study are

listed in a Key Resources table.

Testis cell stage classification and measurements. Cells at an early stage of

development (stem cells and early-stage somatic and germline cells) were located in

the apical region of the testis, and were identified by their small and dense nuclei[120].

GSC were defined as Vasa-positive cells in direct contact with the hub; proliferating

GSC were identified as Vasa-positive cells in direct contact with the hub that were also

phospho-H3 positive. Cells in the testis region occupied by primary spermatocytes were

identified by their large cell size and decondensed chromosome staining occupying

three nuclear domains[120]. Spermatid bundles were identified by their condensed and

needle-shaped nuclei, which roughly corresponds to nuclei with protamine-based

chromatin[121]. Testis size was measured by quantifying the length of a line drawn

down the middle of a testis image; starting from the apical tip of the testis and ending

where the testis meets the seminal vesicle.

FLP-FRT clone induction. Adult males were collected at 3-5 days post-eclosion and

heat-shocked three times at 37°C with a 10 min rest period at room temperature

between heat shocks. After heat-shock, the flies were incubated at room temperature

until dissection.

Immunohistochemistry. Fixed samples were rinsed three times with blocking solution

containing 0.2% bovine serum albumin (A4503, Sigma-Aldrich), 0.3% Triton-X in PBS,

then blocked for 1 hr on a rotating platform at room temperature. During the incubation,

the blocking solution was changed every 15 minutes. After blocking, the sample were

resuspended in blocking solution with the appropriate concentration of primary antibody

(see Key Resources table), and incubated overnight at 4°C. Samples were rinsed three

times with blocking solution after removing primary antibody, and blocked for one hour

on a rotating platform in blocking solution. Secondary antibody was applied in blocking

solution and left on the rotating platform at room temperature for 40 min. The sample

was rinsed with blocking solution three more times, and washed four times for 15 min

per wash in blocking solution. Testis samples were resuspended in Vectashield

mounting media with DAPI (H-1200-10, Vector Laboratory) or SlowFade Diamond

mounting media (S36972, Thermo Fisher Scientific) prior to mounting.

Lipid droplet staining. Fixed testes were briefly permeabilized with 0.1% Triton-X in

PBS for 5 min prior to applying phalloidin. For BODIPY staining, samples were

suspended in PBS containing 10 µg/mL DAPI (2879083-5mg, PeproTech), 1:500

BODIPY 495/503 (Thermo Fisher Scientific D3922), and 1:1000 phalloidin iFluor647

(ab176759, Abcam) or 1:40 phalloidin TexasRed (T7471, Thermo Fisher Scientific). For

staining with LipidTox Red, samples were suspended in PBS containing 10 µg/mL DAPI

(2879083-5mg, PeproTech), 1:200 LipidTox Red (H34476, Thermo Fisher Scientific),

and 1:1000 phalloidin iFluor647 (ab176759, Abcam). For staining free sterols, samples

were prepared as for BODIPY staining with 50 µg/mL filipin in place of BODIPY for 30

min. Samples were incubated on a rotating platform for 40 minutes at room

temperature. After incubation, samples were washed twice with PBS, then resuspended

in SlowFade Diamond mounting media (Thermo Fisher Scientific S36972) prior to

mounting.

Image acquisition and processing. All images were acquired on a Leica SP5 confocal

microscope system with 20X or 40X objectives and quantified with Fiji image analysis

software[122].

Drosophila lipidomics. *Drosophila* extracts were prepared following the previously

reported protocol[123]. Briefly, 10 *Drosophila* males (~10 mg) were weighed, 300 µL of

ice-cold methanol/water mixture (9:1, v:v) was added to these males, and the samples

were homogenized with glass beads using a bead beater (mini-beadbeater-16,

BioSpec, Bartlesville, Ok, USA). Sample weight was used for sample normalization. Fly

lysate was kept at -20°C for 4 hours for protein precipitation. Then, 900 µL of methyl

tert-butyl ether was added and the solution was shaken for 5 min to extract lipids. To

induce phase separation 285 µL of water was added, followed by centrifugation. The

upper layer was separated, dried, and reconstituted in isopropanol/acetonitrile (1:1, v:v)

for liquid chromatography-mass spectrometry (LC-MS) analysis. The volume of

reconstitution solution was proportional to sample weight for normalization. Quality

control (QC) samples were prepared by pooling 20 µL aliquot from each sample. The

method blank sample was prepared using an identical workflow but without adding

Drosophila.

Drosophila extracts were analyzed on an UHR-QqTOF (Ultra-High Resolution

Qq-Time-Of-Flight) mass spectrometry Impact II (Bruker Daltonics, Bremen, Germany)

interfaced with an Agilent 1290 Infinity II LC Systems (Agilent Technologies, Santa

Clara, CA, USA). LC separation was performed using a Waters reversed-phase (RP)

UPLC Acquity BEH C18 Column (1.7 μm, 1.0 mm ×100 mm, 130 Å) (Milford, MA, USA)

maintained at 30°C. For positive ion mode, the mobile phase A was 60% acetonitrile in

water and the mobile phase B was 90% isopropanol in acetonitrile, both containing 5

mM ammonium formate (pH = 4.8, adjusted by formic acid). For negative ion mode, the

mobile phase A was 60% acetonitrile in water and the mobile phase B was 90%

isopropanol in acetonitrile, both containing 5 mM NH_4FA (pH = 9.8, adjusted by

ammonium hydroxide). The LC gradient for positive and negative ion modes was set as

follows: 0 min, 5% B; 8 min, 40% B; 14 min, 70% B; 20 min, 95% B; 23 min, 95% B; 24

min, 5% B; 33 min, 5% B. The flow rate was 0.1 mL/min. The injection volume was

optimized to 2 μL in positive mode and 5 μL in negative mode using QC sample. The

ESI source conditions were set as follows: dry gas temperature, 220 °C; dry gas flow, 7

L/min; nebulizer gas pressure, 1.6 bar; capillary voltage, 4500 V for positive mode and

3000 V for negative mode. The MS1 analysis was conducted using following

parameters: mass range, 70-1000 m/z; spectrum type: centroid, calculated using

maximum intensity; absolute intensity threshold: 250. Data-dependent MS/MS analysis

parameters: collision energy: 16-30 eV; cycle time, 3 s; spectra rate: 4 Hz when

intensity < 10^4 and 12 Hz when intensity > 10^5, linearly increased from 10^4 to 10^5.

External calibration was applied using sodium formate to ensure the m/z accuracy

before sample analysis.

The raw LC-MS data were processed using MS-DIAL (ver. 4.38)[124]. The

detailed MS-DIAL parameters are: MS1 tolerance, 0.01 Da; MS/MS tolerance, 0.05;

mass slice width, 0.05 Da; smoothing method, linear weighted moving average;

smoothing level, 3 scans; minimum peak width, 5 scans. Lipid features with high

quantitative confidence were selected by the following criteria: retention time was within

the gradient elution time (< 23 min); average intensity in QC samples is larger than 5-

fold of the intensity in method blank sample. Lipid identification was performed by

matching experimental precursor m/z, isotopic ratio and MS/MS spectrum against the

LipidBlast libraries embedded in MS-DIAL. To improve the quantification accuracy, the

measured MS signal intensities were corrected using serial diluted QC samples

following the reported workflow[125].

Quantification and statistical analysis. All microscopy images were quantified using

Fiji software[122]. For lipid droplet counts, a single optical slice through the middle of

the testis containing the hub was used with the exception of FLP-FRT experiment where

all lipid droplets within a GFP-negative cyst were counted (Figure 2I). All statistical

analyses were done using R (obtained from https://cran.r-project.org). With exception of

data concerning spatial distribution, and lipidomic data, Shapiro-Wilk test (via

shapiro.test in base R) was used to assess normality of distribution prior to testing for

significance. Kruskal-Wallis rank sum test (from the R package *coin*) and Dunn's test

(from the R package *dunn.test*) were used in place of Welch two-sample t-test and

Tukey's multiple comparison test when the assumption of normality was not met. For

testing differences in variance between two populations, F-test (via *var.test* in base R)

was used. For testing differences in spatial distribution, two-sample Kolmogorov-

Smirnov test (via *ks.test* in base R) was used. All *p*-values are indicated in figures;

extremely small *p*-values are listed as $p < 2.2 \times 10^{-16}$.

 RESOURCE TABLE

REAGENT or RESOURCE	SOURCE	RESOURCE #
Antibodies		
Anti-Vasa (1:200)	Gift from Dr. R. Lehman, MIT	
Anti-Eya (1:50)	Developmental Studies Hybridoma Bank (DSHB)	eya10H6
Anti-zfh1 (1:1000)	Gift from Dr. J. Skeath, WUSTL	
Anti-boule (1:1000)	Gift from Dr. S. Wasserman, UCSD	
Anti-phospho-histone H3 (1:1000)	Millipore Sigma	05-1354
Experimental models: *Drosophila melanogaster*		
w^{1118}	Bloomington *Drosophila* stock center	3605
CantonS	Bloomington *Drosophila* stock center	64349
OregonR	Bloomington *Drosophila* stock center	25211
bmm^1	Gift from Dr. R. Kühnlein	
bmm^{rev}	Gift from Dr. R. Kühnlein	
mdy[Qx25], cn[1], bw[1]/CyO, I(2)DTS513[1]	Bloomington *Drosophila* stock center	5095
y[1],w[67c23];P{lacW}Cse1[k03802],mdy[k03902]/CyO	Bloomington *Drosophila* stock center	10536
w[1118];P{GD1749}v6367 (UAS-mdy-RNAi)	Vienna *Drosophila* resource center	6367

REAGENT or RESOURCE	SOURCE	RESOURCE #
nos-GAL4::VP16	Bloomington Drosophila stock center	7303
Tj-GAL4	Gift from Dr. D. Godt, University of Toronto	
c587-GAL4	Bloomington Drosophila stock center	67747
Bam-GFP	[40]	
bmm-GFP	Gift from Dr. K. Kamei[57]	
GFP-LD	Gift from Dr. M. Welte[36]	
P{neoFRT}82B, bmm[1]	This study	
P{neoFRT}82B, bmm[rev]	This study	
bam-GFP, bmm[1]	This study	
bam-GFP, bmm[rev]	This study	
Software and algorithms		
Fiji	https://imagej.net/software/fiji/	
R	https://cran.r-project.org	

SUPPLEMENTAL TABLE AND FILES

Supplemental table 1 – Table showing identified lipid species from untargeted lipidomic analysis.

Supplementary file 1 – Raw data and statistical outputs from Figure 1.

Supplementary file 2 – Raw data and statistical outputs from Figure 2 and Supplemental figure 2.

Supplementary file 3 – Raw data and statistical outputs from Figure 3 and

Supplemental figure 3.

Supplementary file 4 – Raw data and statistical outputs from Figure 4 and

Supplemental figure 4.

SUPPLEMENTAL FIGURE LEGENDS

Supplemental Figure 1 related to Figure 1 – Cholesterol is absent from testis lipid

droplets. (A) Testes stained with BODIPY (A) to detect neutral lipids and Filipin III (A')

to detect free cholesterol. Scale bars=50 μm.

Supplemental Figure 2 related to Figure 2 – *bmm* **mRNA levels during**

spermatogenesis in germline and somatic lineages. (A) Pseudotime trajectory of

germline (black line) based on single-cell RNA sequencing data[62]. Individual cells are

labeled according to the annotation within the data set. (B) Rolling average of

normalized *bmm* transcript counts in the germline along the trajectory shown in panel A

are plotted as a black line on the upper panel. Composition of cell types mapped on to

the trajectory at each time point is shown at the lower half of panel B. (C) Pseudotime

trajectory of the somatic cells (black line) based on publicly available single-cell RNA

sequencing data[62]. Individual cells are labeled according to the annotation within the

data set. (D) Rolling average of normalized *bmm* transcript counts in somatic cells

plotted as a black line along the trajectory shown in C (upper panel). Composition of cell

types mapped on to the trajectory at each time point (lower panel). (E-H)

Representative images of *bmm^rev^* (E and F) and *bmm^1^* (G and H) testes with somatic

over-expression of *GFP-LD* (*Tj-GAL4>UAS-GFP-LD*). Panel F and H contain magnified

images of the area indicated by the boxes in panel E and G, respectively. In *bmm^rev*

testes, LD were restricted to a region near the apical tip (E) of the testis in both somatic

(F–F''' arrows) and germline cells (F–F''' arrowheads). In *bmm^1* testes, LD were present

in both somatic (G–H arrows) and germline cells (G–H arrowheads), near the apical tip

of the testis in a region corresponding to early-stage germ cells and in the region

corresponding to spermatocytes. (E,G) Scale bars=50 µm; (F,H) scale bars=20 µm.

Supplemental Figure 3 related to Figure 3 – Additional characterization of testis

development and spermatogenesis in animals lacking *bmm*. (A) Testis size was

smaller in *bmm^1* mutant animals compared with *bmm^rev* controls at <24 hr post-eclosion

when raised at 25°C (A; Welch two-sample t-test). (B) The number of spermatid bundles

was significantly lower in *bmm^1* mutant animals compared with *bmm^rev* controls at <24

hr post-eclosion when raised at 25°C (Kruskal-Wallis rank sum test). (C) Testis size was

significantly smaller in *bmm^1* mutant males compared with *bmm^rev* control males at 14-

days post-eclosion (Welch two-sample t-test). (D) While the median number of

spermatid bundles was not significantly different between *bmm^1* mutant males and

bmm^rev control males at 14 days post-eclosion (Welch two-sample t-test), 8/27 *bmm^1*

testis had no spermatid bundles, a phenotype absent in age-matched *bmm^rev* males

(0/22) (p=0.0163, Pearson's Chi-squared test), suggesting a subtle defect is present.

(E) Food supplemented with 4% medium chain triglyceride (MCT), but not long chain

triglyceride (LCT), significantly increased testis length in *bmm^1* animals but had no effect

on this phenotype in *bmm^rev* control animals (one-way ANOVA with Tukey multiple

comparison test). (F) Food supplemented with 4% medium chain triglyceride

significantly increased the number of spermatid bundles in *bmm¹* testes but had no

effect on this phenotype in *bmm^rev* control animals (one-way ANOVA with Tukey

multiple comparison test). (G) Representative images of *bmm^rev* (G–G') or *bmm¹* (G''–

G''') testes stained for Fas3 (G and G'') and Vas (G' and G'''). Scale bars=25 μm. (H)

Quantification of hub area in *bmm^rev* or *bmm¹* testes showed a significantly larger hub

size in *bmm¹* testes (Welch two-sample t-test). (I) The number of germline stem cells

(GSC) undergoing mitosis (phospho-histone H3⁺ GSC/total GSC) was not significantly

different between *bmm¹* and *bmm^rev* testes (Kruskal-Wallis rank sum test). (J) The

distance between the hub and the first Bam-GFP positive cyst (Figure 3H) was

significantly higher in *bmm¹* testes than in *bmm^rev* testes (Welch two-sample t-test). (K)

All *bmm^rev* testes and most *bmm¹* testes contained spermatids when raised at 25°C;

however, the most advanced stage of spermatogenesis observed in the majority of

bmm¹ testes isolated from animals reared at 29°C was the spermatocyte stage. (L)

Testes isolated from *bmm¹* animals showed a significantly smaller Boule-positive area

than control testes (Welch two-sample t-test). (M) Testes isolated from *bmm¹* animals

contain fewer individualization complexes than *bmm^rev* control testes (Kruskal-Wallis

rank sum test). (N) Fewer waste bags were present in testes isolated from *bmm¹*

animals compared with *bmm^rev* control testes (Kruskal-Wallis rank sum test).

Supplemental Figure 4 related to Figure 4 – Lipidomic analysis of animals lacking

***bmm*.** (A) Higher fold-changes of triglycerides in *bmm¹* animals were associated with

less saturation in the acyl-groups (Kendall's rank correlation test). (B) Higher fold-

changes of triglycerides in *bmm^1* animals were associated with higher number of

carbons in the acyl-groups (Kendall's rank correlation test). Each dot represents a single

triglyceride species for panel B and C. (C) Volcano plot of identified lipids;

monoglycerides shown in blue and diglycerides shown in orange. Many monoglycerides

and diglycerides show increase in fold-change in *bmm^1* males. (D) The number of

carbon and the degree of saturation of monoglycerides (MAG) and diglycerides (DAG)

with significant changes in abundance between *bmm^1* and *bmm^{rev}* males. (E) Volcano

plot of identified lipids; fatty acids shown in magenta and acyl-carnitine shown in green.

Many fatty acids show an increase in fold-change while many acyl-carnitines show a

decrease in fold-change in *bmm^1* males. (F) The number of carbon and the degree of

saturation of fatty acids (FA) and acyl-carnitines (ACar) with significant changes in

abundance between *bmm^1* and *bmm^{rev}* males. (G) Volcano plot of identified lipids;

membrane lipids shown in yellow. (H) The number of carbon and the degree of

saturation of membrane lipids with significant changes in abundance between *bmm^1*

and *bmm^{rev}* males. For panel G and H, PC: phosphatidylcholine; PE:

phosphatidylethanolamine; PI: phosphatidylinositol; LPC: lysophosphatidylcholine; LPE:

lysophosphatidylethanolamine; SM: sphingomyelin; PG: phosphatidylglycerol. (I) Loss

of *mdy* function rescued the elevated number of LD in *bmm^1* testes to control levels

(one-way ANOVA with Tukey multiple comparison test). (J) Germline-specific loss of

mdy in *bmm^1* animals did not reduce GSC numbers, but the variance in GSC number

was significantly rescued (*nos-GAL4>+*; *bmm^1* vs *nos-GAL4>mdy RNAi*; *bmm^1*: *p*=4.5 ×

10^{-5}; *+>mdy RNAi*; *bmm^1* vs *nos-GAL4>mdy RNAi*; *bmm^1*: *p*=0.0082 by F-test).

655
656

REFERENCES

1. Brill JA, Yildirim S, Fabian L. Phosphoinositide signaling in sperm development. Semin Cell Dev Biol. 2016;59: 2–9. doi:10.1016/j.semcdb.2016.06.010

2. Keber R, Rozman D, Horvat S. Sterols in spermatogenesis and sperm maturation. J Lipid Res. 2013;54: 20–33. doi:10.1194/jlr.R032326

3. Rato L, Alves MG, Socorro S, Duarte AI, Cavaco JE, Oliveira PF. Metabolic regulation is important for spermatogenesis. Nat Rev Urol. 2012;9: 330–338. doi:10.1038/nrurol.2012.77

4. Wang C, Huang X. Lipid metabolism and Drosophila sperm development. Sci China Life Sci. 2012;55: 35–40. doi:10.1007/s11427-012-4274-2

5. de Cuevas M, Matunis EL. The stem cell niche: lessons from the Drosophila testis. Dev Camb Engl. 2011;138: 2861–2869. doi:10.1242/dev.056242

6. Demarco RS, Eikenes ÅH, Haglund K, Jones DL. Investigating Spermatogenesis in Drosophila melanogaster. Methods San Diego Calif. 2014;68: 218–227. doi:10.1016/j.ymeth.2014.04.020

7. Fabian L, Brill JA. Drosophila spermiogenesis: Big things come from little packages. Spermatogenesis. 2012;2: 197–212. doi:10.4161/spmg.21798

8. Brill JA, Hime GR, Scharer-Schuksz M, Fuller MT. A phospholipid kinase regulates actin organization and intercellular bridge formation during germline cytokinesis. Development. 2000;127: 3855.

9. Giansanti MG, Bonaccorsi S, Kurek R, Farkas RM, Dimitri P, Fuller MT, et al. The Class I PITP Giotto Is Required for Drosophila Cytokinesis. Curr Biol. 2006;16: 195–201. doi:10.1016/j.cub.2005.12.011

10. Wong R, Hadjiyanni I, Wei H-C, Polevoy G, McBride R, Sem K-P, et al. PIP2 Hydrolysis and Calcium Release Are Required for Cytokinesis in Drosophila Spermatocytes. Curr Biol. 2005;15: 1401–1406. doi:10.1016/j.cub.2005.06.060

11. Wong R, Fabian L, Forer A, Brill JA. Phospholipase C and myosin light chain kinase inhibition define a common step in actin regulation during cytokinesis. BMC Cell Biol. 2007;8: 15. doi:10.1186/1471-2121-8-15

12. Amoyel M, Hillion K-H, Margolis SR, Bach EA. Somatic stem cell differentiation is regulated by PI3K/Tor signaling in response to local cues. Dev Camb Engl. 2016;143: 3914. doi:10.1242/dev.139782

13. Fairchild MJ, Islam F, Tanentzapf G. Identification of genetic networks that act in the somatic cells of the testis to mediate the developmental program of

spermatogenesis. PLOS Genet. 2017;13: e1007026.
doi:10.1371/journal.pgen.1007026

14. Inaba M, Venkei ZG, Yamashita YM. The polarity protein Baz forms a platform for
the centrosome orientation during asymmetric stem cell division in the Drosophila
male germline. eLife. 2015;4: e04960. doi:10.7554/eLife.04960

15. Krahn MP, Klopfenstein DR, Fischer N, Wodarz A. Membrane Targeting of
Bazooka/PAR-3 Is Mediated by Direct Binding to Phosphoinositide Lipids. Curr
Biol. 2010;20: 636–642. doi:10.1016/j.cub.2010.01.065

16. Papagiannouli F, Berry CW, Fuller MT. The Dlg Module and Clathrin-Mediated
Endocytosis Regulate EGFR Signaling and Cyst Cell-Germline Coordination in
the Drosophila Testis. Stem Cell Rep. 2019;12: 1024–1040.
doi:10.1016/j.stemcr.2019.03.008

17. Ueishi S, Shimizu H, Inoue YH. Male Germline Stem Cell Division and
Spermatocyte Growth Require Insulin Signaling in Drosophila. Cell Struct Funct.
2009;34: 61–69. doi:10.1247/csf.08042

18. Phan VH, Herr DR, Panton D, Fyrst H, Saba JD, Harris GL. Disruption of
Sphingolipid Metabolism Elicits Apoptosis-Associated Reproductive Defects in
Drosophila. Dev Biol. 2007;309: 329–341. doi:10.1016/j.ydbio.2007.07.021

19. Steinhauer J, Gijón MA, Riekhof WR, Voelker DR, Murphy RC, Treisman JE.
Drosophila Lysophospholipid Acyltransferases Are Specifically Required for Germ
Cell Development. Mol Biol Cell. 2009;20: 5224–5235. doi:10.1091/mbc.e09-05-
0382

20. Szafer-Glusman E, Giansanti MG, Nishihama R, Bolival B, Pringle J, Gatti M, et al.
A Role for Very-Long-Chain Fatty Acids in Furrow Ingression during Cytokinesis in
Drosophila Spermatocytes. Curr Biol. 2008;18: 1426–1431.
doi:10.1016/j.cub.2008.08.061

21. Ben-David G, Miller E, Steinhauer J. Drosophila spermatid individualization is
sensitive to temperature and fatty acid metabolism. Spermatogenesis. 2015;5.
doi:10.1080/21565562.2015.1006089

22. Jung A, Hollmann M, Schafer MA. The fatty acid elongase NOA is necessary for
viability and has a somatic role in Drosophila sperm development. J Cell Sci.
2007;120: 2924–2934. doi:10.1242/jcs.006551

23. Rabionet M, Bayerle A, Jennemann R, Heid H, Fuchser J, Marsching C, et al. Male
meiotic cytokinesis requires ceramide synthase 3-dependent sphingolipids with
unique membrane anchors. Hum Mol Genet. 2015;24: 4792–4808.
doi:10.1093/hmg/ddv204

24. Santiago Valtierra FX, Peñalva DA, Luquez JM, Furland NE, Vásquez C, Reyes
JG, et al. Elovl4 and Fa2h expression during rat spermatogenesis: a link to the
very-long-chain PUFAs typical of germ cell sphingolipids. J Lipid Res. 2018;59:
1175–1189. doi:10.1194/jlr.M081885

25. Zadravec D, Tvrdik P, Guillou H, Haslam R, Kobayashi T, Napier JA, et al. ELOVL2
controls the level of n-6 28:5 and 30:5 fatty acids in testis, a prerequisite for male
fertility and sperm maturation in mice. J Lipid Res. 2011;52: 245–255.
doi:10.1194/jlr.M011346

26. Walther TC, Farese RV. Lipid Droplets and Cellular Lipid Metabolism. Annu Rev
Biochem. 2012;81: 687–714. doi:10.1146/annurev-biochem-061009-102430

27. Wat LW, Chao C, Bartlett R, Buchanan JL, Millington JW, Chih HJ, et al. A role for
triglyceride lipase brummer in the regulation of sex differences in Drosophila fat
storage and breakdown. Hassan BA, editor. PLOS Biol. 2020;18: e3000595.
doi:10.1371/journal.pbio.3000595

28. Thiele C, Spandl J. Cell biology of lipid droplets. Curr Opin Cell Biol. 2008;20: 378–
385. doi:10.1016/j.ceb.2008.05.009

29. Bosma M. Lipid droplet dynamics in skeletal muscle. Exp Cell Res. 2016;340: 180–
186. doi:10.1016/j.yexcr.2015.10.023

30. Grönke S, Müller G, Hirsch J, Fellert S, Andreou A, Haase T, et al. Dual Lipolytic
Control of Body Fat Storage and Mobilization in Drosophila. PLOS Biol. 2007;5:
e137. doi:10.1371/journal.pbio.0050137

31. Rambold AS, Cohen S, Lippincott-Schwartz J. Fatty Acid Trafficking in Starved
Cells: Regulation by Lipid Droplet Lipolysis, Autophagy, and Mitochondrial Fusion
Dynamics. Dev Cell. 2015;32: 678–692. doi:10.1016/j.devcel.2015.01.029

32. Dichlberger A, Schlager S, Maaninka K, Schneider WJ, Kovanen PT. Adipose
triglyceride lipase regulates eicosanoid production in activated human mast cells.
J Lipid Res. 2014;55: 2471–2478. doi:10.1194/jlr.M048553

33. Mitsche MA, Hobbs HH, Cohen JC. Patatin-like phospholipase domain–containing
protein 3 promotes transfers of essential fatty acids from triglycerides to
phospholipids in hepatic lipid droplets. J Biol Chem. 2018;293: 6958–6968.
doi:10.1074/jbc.RA118.002333

34. Rajakumari S, Rajasekharan R, Daum G. Triacylglycerol lipolysis is linked to
sphingolipid and phospholipid metabolism of the yeast Saccharomyces
cerevisiae☆. Biochim Biophys Acta BBA - Mol Cell Biol Lipids. 2010;1801: 1314–
1322. doi:10.1016/j.bbalip.2010.08.004

35. Schlager S, Goeritzer M, Jandl K, Frei R, Vujic N, Kolb D, et al. Adipose triglyceride
lipase acts on neutrophil lipid droplets to regulate substrate availability for lipid

mediator synthesis. J Leukoc Biol. 2015;98: 837–850. doi:10.1189/jlb.3A0515-
206R

36. Zanghellini J, Natter K, Jungreuthmayer C, Thalhammer A, Kurat CF, Gogg-
Fassolter G, et al. Quantitative modeling of triacylglycerol homeostasis in yeast -
metabolic requirement for lipolysis to promote membrane lipid synthesis and
cellular growth: Triacylglycerol mobilization in yeast. FEBS J. 2008;275: 5552–
5563. doi:10.1111/j.1742-4658.2008.06681.x

37. Bailey AP, Koster G, Guillermier C, Hirst EMA, MacRae JI, Lechene CP, et al.
Antioxidant Role for Lipid Droplets in a Stem Cell Niche of Drosophila. Cell.
2015;163: 340–353. doi:10.1016/j.cell.2015.09.020

38. Liu L, MacKenzie KR, Putluri N, Maletić-Savatić M, Bellen HJ. The Glia-Neuron
Lactate Shuttle and Elevated ROS Promote Lipid Synthesis in Neurons and Lipid
Droplet Accumulation in Glia via APOE/D. Cell Metab. 2017;26: 719-737.e6.
doi:10.1016/j.cmet.2017.08.024

39. Nguyen TB, Louie SM, Daniele JR, Tran Q, Dillin A, Zoncu R, et al. DGAT1-
Dependent Lipid Droplet Biogenesis Protects Mitochondrial Function during
Starvation-Induced Autophagy. Dev Cell. 2017;42: 9-21.e5.
doi:10.1016/j.devcel.2017.06.003

40. Henne M. And three's a party: lysosomes, lipid droplets, and the ER in lipid
trafficking and cell homeostasis. Curr Opin Cell Biol. 2019;59: 40–49.
doi:10.1016/j.ceb.2019.02.011

41. Welte MA. Expanding Roles for Lipid Droplets. Curr Biol. 2015;25: R470–R481.
doi:10.1016/j.cub.2015.04.004

42. Casado ME, Pastor O, Mariscal P, Canfrán-Duque A, Martínez-Botas J, Kraemer
FB, et al. Hormone-sensitive lipase deficiency disturbs the fatty acid composition
of mouse testis. Prostaglandins Leukot Essent Fatty Acids. 2013;88: 227–233.
doi:10.1016/j.plefa.2012.12.005

43. El-Shehawi AM, El-Shazly S, Ahmed M, Alkafafy M, Sayed S, Farouk S, et al.
Transcriptome Analysis of Testis from HFD-Induced Obese Rats (Rattus
norvigicus) Indicated Predisposition for Male Infertility. Int J Mol Sci. 2020;21:
6493. doi:10.3390/ijms21186493

44. Wang W, Wei S, Li L, Su X, Du C, Li F, et al. Proteomic analysis of murine testes
lipid droplets. Sci Rep. 2015;5: 12070. doi:10.1038/srep12070

45. Bajpai M, Gupta G, Setty BS. Changes in carbohydrate metabolism of testicular
germ cells during meiosis in the rat. Eur J Endocrinol. 1998;138: 322–327.

46. Kerr JB, Kretser DMD. CYCLIC VARIATIONS IN SERTOLI CELL LIPID CONTENT
THROUGHOUT THE SPERMATOGENIC CYCLE IN THE RAT. Reproduction.
1975;43: 1–8. doi:10.1530/jrf.0.0430001

47. Mori H, Christensen AK. Morphometric analysis of Leydig cells in the normal rat
testis. J Cell Biol. 1980;84: 340–354. doi:10.1083/jcb.84.2.340

48. Paniagua R, Rodríguez MC, Nistal M, Fraile B, Amat P. Changes in the lipid
inclusion/Sertoli cell cytoplasm area ratio during the cycle of the human
seminiferous epithelium. Reproduction. 1987;80: 335–341.
doi:10.1530/jrf.0.0800335

49. Demarco RS, Uyemura BS, D'Alterio C, Jones DL. Mitochondrial fusion regulates
lipid homeostasis and stem cell maintenance in the Drosophila testis. Nat Cell
Biol. 2019;21: 710–720. doi:10.1038/s41556-019-0332-3

50. Chen M, Wang H, Li X, Li N, Xu G, Meng Q. PLIN1 deficiency affects testicular
gene expression at the meiotic stage in the first wave of spermatogenesis. Gene.
2014;543: 212–219. doi:10.1016/j.gene.2014.04.021

51. Hermo L, Chung S, Gregory M, Smith CE, Wang SP, El-Alfy M, et al. Alterations in
the testis of hormone sensitive lipase-deficient mice is associated with decreased
sperm counts, sperm motility, and fertility. Mol Reprod Dev. 2008;75: 565–577.
doi:10.1002/mrd.20800

52. Masaki H, Kim N, Nakamura H, Kumasawa K, Kamata E, Hirano K, et al. Long-
chain fatty acid triglyceride (TG) metabolism disorder impairs male fertility: a study
using adipose triglyceride lipase deficient mice. MHR Basic Sci Reprod Med.
2017;23: 452–460. doi:10.1093/molehr/gax031

53. Freeman DA, Ascoli M. Studies on the source of cholesterol used for steroid
biosynthesis in cultured Leydig tumor cells. J Biol Chem. 1982;257: 14231–14238.

54. Wang F, Chen Z, Ren X, Tian Y, Wang F, Liu C, et al. Hormone-sensitive lipase
deficiency alters gene expression and cholesterol content of mouse testis.
Reproduction. 2017;153: 175–185. doi:10.1530/REP-16-0484

55. de Cuevas M, Matunis EL. The stem cell niche: lessons from the Drosophila testis.
Development. 2011;138: 2861–2869. doi:10.1242/dev.056242

56. Walther TC, Chung J, Jr RVF. Lipid Droplet Biogenesis. Annu Rev Cell Dev Biol.
2017;33: 491–510. doi:10.1146/annurev-cellbio-100616-060608

57. Boyle M, DiNardo S. Specification, migration and assembly of the somatic cells of
the Drosophila gonad. Development. 1995;121: 1815–1825.
doi:10.1242/dev.121.6.1815

58. Yu YV, Li Z, Rizzo NP, Einstein J, Welte MA. Targeting the motor regulator Klar to
lipid droplets. BMC Cell Biol. 2011;12: 9. doi:10.1186/1471-2121-12-9

59. Leatherman JL, DiNardo S. Zfh-1 Controls Somatic Stem Cell Self-Renewal in the
Drosophila Testis and Nonautonomously Influences Germline Stem Cell Self-
Renewal. Cell Stem Cell. 2008;3: 44–54. doi:10.1016/j.stem.2008.05.001

60. Fabrizio JJ, Boyle M, DiNardo S. A somatic role for eyes absent (eya) and sine
oculis (so) in drosophila spermatocyte development. Dev Biol. 2003;258: 117–
128. doi:10.1016/S0012-1606(03)00127-1

61. Chen D, McKearin DM. A discrete transcriptional silencer in the bam gene
determines asymmetric division of the Drosophila germline stem cell.
Development. 2003;130: 1159–1170. doi:10.1242/dev.00325

62. Insco ML, Leon A, Tam CH, McKearin DM, Fuller MT. Accumulation of a
differentiation regulator specifies transit amplifying division number in an adult
stem cell lineage. Proc Natl Acad Sci U S A. 2009;106: 22311–22316.
doi:10.1073/pnas.0912454106

63. Chen D, Wu C, Zhao S, Geng Q, Gao Y, Li X, et al. Three RNA Binding Proteins
Form a Complex to Promote Differentiation of Germline Stem Cell Lineage in
Drosophila. PLoS Genet. 2014;10. doi:10.1371/journal.pgen.1004797

64. Gonczy P, Matunis E, DiNardo S. bag-of-marbles and benign gonial cell neoplasm
act in the germline to restrict proliferation during Drosophila spermatogenesis.
Development. 1997;124: 4361–4371.

65. Athenstaedt K, Daum G. YMR313c/TGL3 Encodes a Novel Triacylglycerol Lipase
Located in Lipid Particles of Saccharomyces cerevisiae*. J Biol Chem. 2003;278:
23317–23323. doi:10.1074/jbc.M302577200

66. Chitraju C, Trötzmüller M, Hartler J, Wolinski H, Thallinger GG, Haemmerle G, et
al. The impact of genetic stress by ATGL deficiency on the lipidome of lipid
droplets from murine hepatocytes. J Lipid Res. 2013;54: 2185–2194.
doi:10.1194/jlr.M037952

67. Grönke S, Mildner A, Fellert S, Tennagels N, Petry S, Müller G, et al. Brummer
lipase is an evolutionary conserved fat storage regulator in Drosophila. Cell
Metab. 2005;1: 323–330. doi:10.1016/j.cmet.2005.04.003

68. Haemmerle G, Lass A, Zimmermann R, Gorkiewicz G, Meyer C, Rozman J, et al.
Defective Lipolysis and Altered Energy Metabolism in Mice Lacking Adipose
Triglyceride Lipase. Science. 2006;312: 734–737. doi:10.1126/science.1123965

69. Haemmerle G, Moustafa T, Woelkart G, Büttner S, Schmidt A, van de Weijer T, et
al. ATGL-mediated fat catabolism regulates cardiac mitochondrial function via
PPAR-α and PGC-1. Nat Med. 2011;17: 1076–1085. doi:10.1038/nm.2439

70. Huijsman E, van de Par C, Economou C, van der Poel C, Lynch GS, Schoiswohl G,
et al. Adipose triacylglycerol lipase deletion alters whole body energy metabolism
and impairs exercise performance in mice. Am J Physiol-Endocrinol Metab.
2009;297: E505–E513. doi:10.1152/ajpendo.00190.2009

71. Korbelius M, Vujic N, Sachdev V, Obrowsky S, Rainer S, Gottschalk B, et al.
ATGL/CGI-58-Dependent Hydrolysis of a Lipid Storage Pool in Murine
Enterocytes. Cell Rep. 2019;28: 1923-1934.e4. doi:10.1016/j.celrep.2019.07.030

72. Kurat CF, Natter K, Petschnigg J, Wolinski H, Scheuringer K, Scholz H, et al.
Obese Yeast: Triglyceride Lipolysis Is Functionally Conserved from Mammals to
Yeast *. J Biol Chem. 2006;281: 491–500. doi:10.1074/jbc.M508414200

73. Zimmermann R, Strauss JG, Haemmerle G, Schoiswohl G, Birner-Gruenberger R,
Riederer M, et al. Fat Mobilization in Adipose Tissue Is Promoted by Adipose
Triglyceride Lipase. Science. 2004;306: 1383–1386.
doi:10.1126/science.1100747

74. Attané C, Peyot M-L, Lussier R, Poursharifi P, Zhao S, Zhang D, et al. A beta cell
ATGL-lipolysis/adipose tissue axis controls energy homeostasis and body weight
via insulin secretion in mice. Diabetologia. 2016;59: 2654–2663.
doi:10.1007/s00125-016-4105-2

75. Lee JH, Kong J, Jang JY, Han JS, Ji Y, Lee J, et al. Lipid Droplet Protein LID-1
Mediates ATGL-1-Dependent Lipolysis during Fasting in Caenorhabditis elegans.
Mol Cell Biol. 2014;34: 4165–4176. doi:10.1128/MCB.00722-14

76. Tuohetahuntila M, Molenaar MR, Spee B, Brouwers JF, Houweling M, Vaandrager
AB, et al. ATGL and DGAT1 are involved in the turnover of newly synthesized
triacylglycerols in hepatic stellate cells. J Lipid Res. 2016;57: 1162–1174.
doi:10.1194/jlr.M066415

77. Men TT, Thanh DNV, Yamaguchi M, Suzuki T, Hattori G, Arii M, et al. A *Drosophila*
Model for Screening Antiobesity Agents. BioMed Res Int. 2016;2016: 1–10.
doi:10.1155/2016/6293163

78. Li H, Janssens J, Waegeneer MD, Kolluru SS, Davie K, Gardeux V, et al. Fly Cell
Atlas: a single-cell transcriptomic atlas of the adult fruit fly. bioRxiv; 2021. p.
2021.07.04.451050. doi:10.1101/2021.07.04.451050

79. Xu T, Rubin GM. Analysis of genetic mosaics in developing and adult Drosophila
tissues. Development. 1993;117: 1223–1237.

80. Kim N, Nakamura H, Masaki H, Kumasawa K, Hirano K, Kimura T. Effect of lipid
metabolism on male fertility. Biochem Biophys Res Commun. 2017;485: 686–692.
doi:10.1016/j.bbrc.2017.02.103

81. Lasko PF, Ashburner M. The product of the Drosophila gene vasa is very similar to
eukaryotic initiation factor-4A. Nature. 1988;335: 611–617. doi:10.1038/335611a0

82. Resende LPF, Boyle M, Tran D, Fellner T, Jones DL. Headcase Promotes Cell
Survival and Niche Maintenance in the Drosophila Testis. Bökel C, editor. PLoS
ONE. 2013;8: e68026. doi:10.1371/journal.pone.0068026

83. Kiger AA, White-Cooper H, Fuller MT. Somatic support cells restrict germline stem
cell self-renewal and promote differentiation. Nature. 2000;407: 750–754.
doi:10.1038/35037606

84. Lin TY, Viswanathan S, Wood C, Wilson PG, Wolf N, Fuller MT. Coordinate
developmental control of the meiotic cell cycle and spermatid differentiation in
Drosophila males. Development. 1996;122: 1331–1341.

85. Eichmann TO, Kumari M, Haas JT, Farese RV, Zimmermann R, Lass A, et al.
Studies on the substrate and stereo/regioselectivity of adipose triglyceride lipase,
hormone-sensitive lipase, and diacylglycerol-O-acyltransferases. J Biol Chem.
2012;287: 41446–41457. doi:10.1074/jbc.M112.400416

86. Schweiger M, Schreiber R, Haemmerle G, Lass A, Fledelius C, Jacobsen P, et al.
Adipose Triglyceride Lipase and Hormone-sensitive Lipase Are the Major
Enzymes in Adipose Tissue Triacylglycerol Catabolism *. J Biol Chem. 2006;281:
40236–40241. doi:10.1074/jbc.M608048200

87. Missaglia S, Maggi L, Mora M, Gibertini S, Blasevich F, Agostoni P, et al. Late
onset of neutral lipid storage disease due to novel PNPLA2 mutations causing
total loss of lipase activity in a patient with myopathy and slight cardiac
involvement. Neuromuscul Disord. 2017;27: 481–486.
doi:10.1016/j.nmd.2017.01.011

88. Williams ML, Coleman RA, Placezk D, Grunfeld C. Neutral lipid storage disease: a
possible functional defect in phospholipid-linked triacylglycerol metabolism.
Biochim Biophys Acta BBA - Mol Basis Dis. 1991;1096: 162–169.
doi:10.1016/0925-4439(91)90055-E

89. Yang L, Liang J, Lam SM, Yavuz A, Shui G, Ding M, et al. Neuronal lipolysis
participates in PUFA-mediated neural function and neurodegeneration. EMBO
Rep. 2020;21. doi:10.15252/embr.202050214

90. Nazario-Yepiz NO, Fernández Sobaberas J, Lyman R, Campbell MR, Shankar V,
Anholt RRH, et al. Physiological and metabolomic consequences of reduced
expression of the Drosophila brummer triglyceride Lipase. Melkani GC, editor.
PLOS ONE. 2021;16: e0255198. doi:10.1371/journal.pone.0255198

91. Giedt MS, Thomalla JM, Johnson MR, Lai ZW, Tootle TL, Welte MA. Adipose
triglyceride lipase promotes prostaglandin-dependent actin remodeling by

regulating substrate release from lipid droplets. Cell Biology; 2021 Aug.
doi:10.1101/2021.08.02.454724

92. Beller M, Bulankina AV, Hsiao H-H, Urlaub H, Jäckle H, Kühnlein RP. PERILIPIN-
Dependent Control of Lipid Droplet Structure and Fat Storage in Drosophila. Cell
Metab. 2010;12: 521–532. doi:10.1016/j.cmet.2010.10.001

93. Buszczak M, Lu X, Segraves WA, Chang TY, Cooley L. Mutations in the midway
Gene Disrupt a Drosophila Acyl Coenzyme A: Diacylglycerol Acyltransferase.
Genetics. 2002;160: 1511–1518.

94. Martínez BA, Hoyle RG, Yeudall S, Granade ME, Harris TE, Castle JD, et al. Innate
immune signaling in Drosophila shifts anabolic lipid metabolism from triglyceride
storage to phospholipid synthesis to support immune function. PLOS Genet.
2020;16: e1009192. doi:10.1371/journal.pgen.1009192

95. Freeman DA, Ascoli M. Studies on the source of cholesterol used for steroid
biosynthesis in cultured Leydig tumor cells. J Biol Chem. 1982;257: 14231–14238.

96. Wang F, Chen Z, Ren X, Tian Y, Wang F, Liu C, et al. Hormone-sensitive lipase
deficiency alters gene expression and cholesterol content of mouse testis.
Reproduction. 2017;153: 175–185. doi:10.1530/REP-16-0484

97. Walther TC, Farese RV. Lipid droplets and cellular lipid metabolism. Annu Rev
Biochem. 2012;81: 687–714. doi:10.1146/annurev-biochem-061009-102430

98. Olzmann JA, Carvalho P. Dynamics and functions of lipid droplets. Nat Rev Mol
Cell Biol. 2019;20: 137–155. doi:10.1038/s41580-018-0085-z

99. Birse RT, Choi J, Reardon K, Rodriguez J, Graham S, Diop S, et al. High-fat-diet-
induced obesity and heart dysfunction are regulated by the TOR pathway in
Drosophila. Cell Metab. 2010;12: 533–44. doi:10.1016/j.cmet.2010.09.014

100. Molaei M, Vandehoef C, Karpac J. NF-κB Shapes Metabolic Adaptation by
Attenuating Foxo-Mediated Lipolysis in Drosophila. Dev Cell. 2019;49: 802-
810.e6. doi:10.1016/j.devcel.2019.04.009

101. Alic N, Andrews TD, Giannakou ME, Papatheodorou I, Slack C, Hoddinott MP, et
al. Genome-wide dFOXO targets and topology of the transcriptomic response to
stress and insulin signalling. Mol Syst Biol. 2011;7: 502. doi:10.1038/msb.2011.36

102. Junger MA, Rintelen F, Stocker H, Wasserman JD, Vegh M, Radimerski T, et al.
The Drosophila forkhead transcription factor FOXO mediates the reduction in cell
number associated with reduced insulin signaling. J Biol. 2003;2: 20.
doi:10.1186/1475-4924-2-20

103. Zinke I, Schutz CS, Katzenberger JD, Bauer M, Pankratz MJ. Nutrient control of
gene expression in Drosophila: microarray analysis of starvation and sugar-
dependent response. EMBO J. 2002;21: 6162–73.

104. Puig O, Tjian R. Transcriptional feedback control of insulin receptor by
dFOXO/FOXO1. Genes Dev. 2005;19: 2435–2446. doi:10.1101/gad.1340505

105. Kang P, Chang K, Liu Y, Bouska M, Birnbaum A, Karashchuk G, et al. Drosophila
Kruppel homolog 1 represses lipolysis through interaction with dFOXO. Sci Rep.
2017;7: 16369. doi:10.1038/s41598-017-16638-1

106. Bartz R, Zehmer JK, Zhu M, Chen Y, Serrero G, Zhao Y, et al. Dynamic activity of
lipid droplets: protein phosphorylation and GTP-mediated protein translocation. J
Proteome Res. 2007;6: 3256–3265. doi:10.1021/pr070158j

107. Pagnon J, Matzaris M, Stark R, Meex RCR, Macaulay SL, Brown W, et al.
Identification and functional characterization of protein kinase A phosphorylation
sites in the major lipolytic protein, adipose triglyceride lipase. Endocrinology.
2012;153: 4278–4289. doi:10.1210/en.2012-1127

108. Narbonne P, Roy R. Caenorhabditis elegans dauers need LKB1/AMPK to ration
lipid reserves and ensure long-term survival. Nature. 2009;457: 210–214.
doi:10.1038/nature07536

109. Ahmadian M, Abbott MJ, Tang T, Hudak CSS, Kim Y, Bruss M, et al.
Desnutrin/ATGL is regulated by AMPK and is required for a brown adipose
phenotype. Cell Metab. 2011;13: 739–748. doi:10.1016/j.cmet.2011.05.002

110. Amoyel M, Hillion K-H, Margolis SR, Bach EA. Somatic stem cell differentiation is
regulated by PI3K/Tor signaling in response to local cues. Dev Camb Engl.
2016;143: 3914–3925. doi:10.1242/dev.139782

111. Amoyel M, Simons BD, Bach EA. Neutral competition of stem cells is skewed by
proliferative changes downstream of Hh and Hpo. EMBO J. 2014;33: 2295–2313.
doi:10.15252/embj.201387500

112. Ueishi S, Shimizu H, H. Inoue Y. Male Germline Stem Cell Division and
Spermatocyte Growth Require Insulin Signaling in Drosophila. Cell Struct Funct.
2009;34: 61–69. doi:10.1247/csf.08042

113. Hof-Michel S, Cigoja L, Huhn S, Bökel C. Innate immune signalling drives loser
cell elimination during stem cell competition in the *Drosophila* testis.
Developmental Biology; 2020 Mar. doi:10.1101/2020.03.05.979161

114. Couderc J-L, Richard G, Vachias C, Mirouse V. Drosophila LKB1 is required for
the assembly of the polarized actin structure that allows spermatid
individualization. White-Cooper H, editor. PLOS ONE. 2017;12: e0182279.
doi:10.1371/journal.pone.0182279

115. Steinhauer J, Statman B, Fagan JK, Borck J, Surabhi S, Yarikipati P, et al.
Combover interacts with the axonemal component Rsp3 and is required for
Drosophila sperm individualization. Dev Camb Engl. 2019;146: dev179275.
doi:10.1242/dev.179275

116. Tartarin P, Guibert E, Touré A, Ouiste C, Leclerc J, Sanz N, et al. Inactivation of
AMPKα1 induces asthenozoospermia and alters spermatozoa morphology.
Endocrinology. 2012;153: 3468–3481. doi:10.1210/en.2011-1911

117. Martin-Hidalgo D, Hurtado de Llera A, Calle-Guisado V, Gonzalez-Fernandez L,
Garcia-Marin L, Bragado MJ. AMPK Function in Mammalian Spermatozoa. Int J
Mol Sci. 2018;19: 3293. doi:10.3390/ijms19113293

118. Pitetti J-L, Calvel P, Zimmermann C, Conne B, Papaioannou MD, Aubry F, et al.
An Essential Role for Insulin and IGF1 Receptors in Regulating Sertoli Cell
Proliferation, Testis Size, and FSH Action in Mice. Mol Endocrinol. 2013;27: 814–
827. doi:10.1210/me.2012-1258

119. Thurmond J, Goodman JL, Strelets VB, Attrill H, Gramates LS, Marygold SJ, et al.
FlyBase 2.0: the next generation. Nucleic Acids Res. 2018;47: D759–D765.
doi:10.1093/nar/gky1003

120. White-Cooper H. Spermatogenesis. In: Henderson DS, editor. Drosophila
Cytogenetics Protocols. Totowa, NJ: Humana Press; 2004. pp. 45–75.
doi:10.1385/1-59259-665-7:45

121. Fabian L, Brill JA. Drosophila spermiogenesis. Spermatogenesis. 2012;2: 197–
212. doi:10.4161/spmg.21798

122. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al.
Fiji: an open-source platform for biological-image analysis. Nat Methods. 2012;9:
676–682. doi:10.1038/nmeth.2019

123. Yu H, Villanueva N, Bittar T, Arsenault E, Labonté B, Huan T. Parallel
metabolomics and lipidomics enables the comprehensive study of mouse brain
regional metabolite and lipid patterns. Anal Chim Acta. 2020;1136: 168–177.
doi:10.1016/j.aca.2020.09.051

124. Tsugawa H, Cajka T, Kind T, Ma Y, Higgins B, Ikeda K, et al. MS-DIAL: data-
independent MS/MS deconvolution for comprehensive metabolome analysis. Nat
Methods. 2015;12: 523–526. doi:10.1038/nmeth.3393

125. Yu H, Huan T. Patterned Signal Ratio Biases in Mass Spectrometry-Based
Quantitative Metabolomics. Anal Chem. 2021;93: 2254–2262.
doi:10.1021/acs.analchem.0c04113

1051

1052
1053

Figure 1 – Lipid droplets are present in early-stage somatic and germline cells.

Figure 2 – *brummer* regulates testis lipid droplets in a cell-autonomous manner.

Figure 3 – A cell-autonomous role of *brummer* in regulating spermatogenesis.

Figure 4 – Loss of *brummer* disrupts triglyceride homeostasis and leads to spermatogenic defects.

Supplemental Figure 1 – Cholesterol is absent from testis lipid droplets.

Supplemental Figure 2 – *brummer* regulates testis lipid droplets in a cell-autonomous manner.

Supplemental Figure 3 – Additional characterization of testis development and spermatogenesis in animals lacking *brummer*.

Supplemental Figure 4 – Lipidomic analysis of animals lacking *brummer*.