TITLE

An important role for triglyceride in regulating spermatogenesis

AUTHORS

Charlotte F. Chao[1&], Yanina-Yasmin Pesch[1&], Huaxu Yu[2], Chenjingyi Wang[2], Maria Aristizabal[3], Tao Huan[2], Guy Tanentzapf[1], Elizabeth J. Rideout[1*]

AFFILIATIONS

[1] Department of Cellular and Physiological Sciences, Life Sciences Institute, The University of British Columbia, Vancouver, BC, Canada V6T 1Z3

[2] Department of Chemistry, The University of British Columbia, Vancouver, BC, Canada V6T 1Z1

[3] Department of Biology, Queen's University, Kingston, ON, Canada K7L 3N6

* Corresponding author

[&] Equally contributing authors

[*] Correspondence: elizabeth.rideout@ubc.ca

SUMMARY

Drosophila is a powerful model to study how lipids affect sperm development[1–3]. Studies in flies showed that phosphatidylinositol and its phosphorylated derivatives influence spermatogenesis[4–14], and uncovered roles for membrane lipids and fatty acids in this process[15–19]. Less is known about how neutral lipids, another major lipid group, affect sperm development. Neutral lipids include triglycerides and cholesterol esters, and reside within specialized organelles called lipid droplets (LD)[20]. Emerging evidence shows testis LD are present in normal and stressful contexts[21–26]. Genes associated with neutral lipid metabolism and LD also show testis expression[27–29], and dysregulation of these genes causes subfertility[30–32]. Yet, many questions remain about the spatial, temporal, and cell type-specific requirements for neutral lipids and LD during spermatogenesis. Here, we show that LD are present in early-stage somatic and germline cells within the *Drosophila* testis. We identified a role for triglyceride lipase *brummer* (*bmm*) in regulating testis LD, and show that whole-body and cell-autonomous loss of *bmm* leads to defects in sperm development. Because lipidomic analysis of *bmm* mutants revealed excess triglyceride accumulation, and spermatogenic defects in *bmm* mutants were rescued by genetically blocking triglyceride synthesis, our data suggest that *bmm*-mediated regulation of triglyceride influences sperm development. This identifies triglyceride as an important neutral lipid that contributes to *Drosophila* sperm development, and reveals a key role for *bmm* in regulating testis triglyceride levels and spermatogenesis.

39 **KEYWORDS**

40 Triglyceride, lipid droplet, spermatogenesis, testis, *brummer*, adipose triglyceride lipase

41 (ATGL), lipidomics.

RESULTS AND DISCUSSION

Lipid droplets are present in early-stage somatic and germline cells

We previously reported the presence of small circular punctae (<1 μm) corresponding to LD near the apical tip of the testis[21]. We confirm these results in w^{1118} males using neutral lipid stain BODIPY (4,4-Difluoro-1,3,5,7,8-Pentamethyl-4-Bora-3a,4a-Diaza-*s*-Indacene) (Figure 1A,S1A). Importantly, we reproduced this spatial distribution of LD in two independent genetic backgrounds and at two additional ages (Figure 1B,1C). In all cases, LD were in a testis region that contains stem cells and early-stage somatic and germline cells (Figure 1A-A', arrows), and in the hub, an organizing center and stem cell niche in the *Drosophila* testis (Figure 1A''-A''', arrows)[33]. LD were largely absent from the testis region occupied by spermatocytes (Figure 1A and A', arrowheads). While LD may contain multiple neutral lipid species[34], cholesterol-binding fluorescent polyene antibiotic filipin III did not detect cholesterol within testis LD (Figure S1A'), suggesting triglyceride is the main neutral lipid in *Drosophila* testis LD.

Drosophila spermatogenesis requires the codevelopment and differentiation of two cell lineages, the germline and the somatic cells[35]. To identify LD in each lineage, we used the GAL4/UAS system to overexpress a transgene in which GFP is fused to the LD-targeting motif of motor protein Klarsicht[36] (*UAS-GFP-LD*). We targeted *UAS-GFP-LD* to somatic cells with *Traffic jam* (*Tj*)-*GAL4* and to the germline using *nanos (nos)-GAL4*; LD were visualized using neutral lipid dye LipidTox. We found LD in the somatic cells of 0-day-old males (Figure 1D), and showed that the majority of somatic LD were <30 µm from the hub (Figure 1E). Because the somatic LD distribution coincided with a marker for somatic stem cells and their immediate daughter cells (Zinc

finger homeodomain 1, Zfh-1) (Figure 1F; two-sample Kolmogorov-Smirnov test)[37], but

not with a marker for late somatic cells (Eyes absent, Eya)[38,39], our data suggests LD

are present in early somatic cells. In the germline, GFP punctae corresponding to LD

were found near the apical tip of the testis in 0-day-old males (Figure 1G,H). We found

that the disappearance of germline LD coincided with peak expression of a GFP

reporter that reflects the expression of Bag-of-marbles (Bam) protein in the testis (Bam-

GFP)[40] (Figure 1I,1J). Because peak Bam expression signals the last round of transient

amplifying mitotic cell cycle prior to the germline's transition into the meiotic cell cycle[41–

43], our data suggests that germline LD, like somatic LD, are present at early stages of

germline development.

***brummer* plays a cell-autonomous role in regulating testis lipid droplets**

Adipose triglyceride lipase (*ATGL*) is a critical regulator of neutral lipid metabolism and

LD[44–53]. Loss of *ATGL* in many cell types triggers LD accumulation, and *ATGL*

overexpression decreases LD number[46,47,50,52,52,54–56]. Given that the *Drosophila ATGL*

homolog *brummer* (*bmm*) regulates testis LD induced by mitochondrial stress[26], we

explored whether *bmm* regulates testis LD under normal physiological conditions. We

first examined *bmm* expression in the testis by isolating this organ from flies in which a

bmm promoter fragment drives *GFP* expression (*bmm-GFP*). Indeed, *bmm-GFP*

accurately reproduces changes to *bmm* mRNA levels[57]. GFP expression was present in

the germline of *bmm-GFP* testes, and germline GFP levels were higher in

spermatocytes than at earlier stages of sperm development (Figure 2A,2B; one-way

ANOVA with Tukey multiple comparison test). Supporting this, our analysis of a publicly

available single-cell RNA sequencing data set of the male reproductive organ[58]

suggested a similar trend in *bmm* mRNA levels between different stages of germline

(Figure S2A,S2B) and somatic cell (Figure S2C,S2D) development. Importantly,

germline GFP levels were negatively correlated with testis LD in *bmm-GFP* flies (Figure

2A,2C), suggesting regions with higher *bmm* expression had fewer LD.

To test whether *bmm* regulates testis LD, we compared LD in testes from 0-day-

old males carrying a loss-of-function mutation in *bmm* (*bmm^1*) to control male testes

(*bmm^{rev}*)[46]. *bmm^1* testes had significantly more LD across all LD sizes compared with

control males (Figure 2D–2G; Welch two-sample t-test with Bonferroni correction), and

showed a significantly expanded LD distribution (Figure 2D–2F,2H; two-sample

Kolmogorov-Smirnov test). This suggests *bmm* normally restricts LD to the region near

the apical tip of the testis, a role we confirm in both somatic and germline lineages

(Figure S2E–S2H). Importantly, after inducing homozygous *bmm^1* or *bmm^{rev}* clones in

the testes using *FLP-FRT* system[59], we found *bmm^1* spermatocyte clones had

significantly more LD at 3 days post-clone induction (Figure 2I; Welch two-sample t-

test), a stage at which LD were absent from *bmm^{rev}* clones. This indicates a previously

unrecognized cell-autonomous role for *bmm* in regulating testis LD, a role we were

unable to assess in somatic cells as we recovered no *bmm^1* somatic cell clones.

***brummer* plays a cell-autonomous role in regulating germline development**

To determine the physiological significance of *bmm*-mediated regulation of testis LD, we

investigated testis and sperm development in males without *bmm* function. In 0-day-old

bmm^1 males reared at 25°C, testis size was significantly smaller than in age-matched

bmm^rev controls (Figure S3A; Welch two-sample t-test), and the number of spermatid

bundles was significantly lower (Figure S3B; Kruskal-Wallis rank sum test). Defects in

testis size and sperm development were also observed in 14-day-old *bmm^1* males

(Figure S3C,S3D Welch two-sample t-test). When the animals were reared at 29°C, a

temperature that exacerbates spermatogenesis defects associated with changes in lipid

metabolism[18], *bmm^1* phenotypes were more pronounced (Figure 3A-3C). This suggests

loss of *bmm* affects testis development and spermatogenesis. Because similar

phenotypes are observed in male mice without ATGL[32], and supplementing the diet of

bmm^1 males with medium-chain triglycerides (MCT) partially rescues the testis and

spermatogenic defects we observed in flies (Figure S3E,S3F; one-way ANOVA with

Tukey multiple comparison test), as it does in mice[32,60], our data suggests flies are a

good model to study how *bmm/ATGL* influences sperm development.

To explore spermatogenesis in *bmm^1* animals, we used a germline-specific

marker Vasa to visualize the germline in the testes of *bmm^1* and *bmm^rev* males (Figure

3D,3E)[61]. We observed a significant increase in the number of germline stem cells

(GSC) (Figure 3F; Kruskal-Wallis rank sum test) and higher variability in GSC number in

bmm^1 males (p=5.7×10^{-12} by F-test). Given that GSC number is affected by hub size

and GSC proliferation[62,63], we monitored both parameters in *bmm^1* and *bmm^rev* controls.

While hub size in *bmm^1* testes was significantly larger than in testes from *bmm^rev*

controls (Figure S3G,S3H; Welch two-sample t-test), the number of phosphohistone H3-

positive GSC, which indicates proliferating GSC, was unchanged in *bmm^1* animals

(Figure S3I; Kruskal-Wallis rank sum test). While this indicates a larger hub may partly

explain *bmm's* effect on GSC number, *bmm* also plays a cell-autonomous role in

regulating GSC, as we recovered a higher proportion of bmm^1 clones in the GSC pool

compared with bmm^{rev} clones at 14 days after clone induction (Figure 3G; Welch two-

sample t-test).

Beyond GSC, we uncovered additional spermatogenesis defects in bmm^1 testes.

Peak Bam-GFP expression in testes from 0-day-old bmm^1 and bmm^{rev} males showed

that GFP-positive cysts with were significantly further away from the hub in bmm^1 testes

(Figure 3H,S3J; Welch two-sample t-test). Indeed, 15/18 bmm^1 testes contained Vasa-

positive cysts with large nuclei in the distal half of the testis (Figure 3I, arrowheads), a

phenotype not present in bmm^{rev} testes (0/8) (p=0.0005 by Pearson's Chi-square test).

Because these phenotypes are also seen in testes with differentiation defects[10,64], we

recorded the stage of sperm development reached by the germline in bmm^1 testes.

Most bmm^1 testes contained post-meiotic cells in males raised at 25°C (Figure S3K);

however, germline development did not progress past the spermatocyte stage in most

bmm^1 testes from animals raised at 29°C (Figure S3K). Testes from bmm^1 males reared

at 25°C also had a smaller Boule-positive area (Figure 3J,S3L; Welch two-sample t-

test), and fewer individualization complexes and waste bags (Figure S3M,S3N; Kruskal-

Wallis rank sum test). Together, these data indicate loss of bmm delays germline

development. Because we recovered fewer bmm^1 spermatocyte and spermatid clones

14 days after clone induction (Figure 3K,3L; Kruskal-Wallis rank sum test), this effect on

germline development represents a cell-autonomous role for bmm.

$brummer$-dependent regulation of testis triglyceride levels affects

spermatogenesis

ATGL catalyzes the first and rate-limiting step of triglyceride hydrolysis[52,65,66]. Loss of

this enzyme or its homologs leads to excess triglyceride accumulation[21,46,52,54,55] and

shifts in multiple lipid classes[45,67–69]. To determine how loss of *bmm* affects

spermatogenesis, we carried out mass spectrometry (MS)-based untargeted lipidomic

profiling of *bmm[1]* and *bmm[rev]* males. Hierarchical clustering of lipid species suggests

that *bmm[1]* and *bmm[rev]* males show distinct lipidomic profiles (Figure 4A). Overall, we

detected 2464 and 1144 lipid features with high quantitative confidence in positive and

negative ion modes, respectively. By matching experimental *m/z*, isotopic ratio, and

tandem MS spectra to lipid libraries, we confirmed 293 unique lipid species

(Supplemental table 1). We found 107 lipids had a significant change in abundance

between *bmm[1]* and *bmm[rev]* males ($p_{adj} < 0.05$): 85 species were upregulated in *bmm[1]*

males and 22 lipid species were downregulated. Among differentially regulated species

in different lipid classes, triglyceride had the largest residual above expected proportion

($p = 5.00 \times 10^{-4}$ by Pearson's Chi-squared test). This suggests triglyceride is the lipid class

most affected by loss of *bmm* (Figure 4B,4C).

In *bmm[1]* males, most triglyceride species (55/97) were significantly higher.

Because we observed a positive correlation between the fold increase in triglyceride

abundance with both the number of double bonds ($p = 7.52 \times 10^{-8}$ by Kendall's rank

correlation test; Figure S4A) and the number of carbons ($p = 2.77 \times 10^{-10}$ by Kendall's rank

correlation test; Figure S4B), our data align well with *bmm*/*ATGL's* known role in

regulating triglyceride levels[46,47,52] and its substrate preference of long-chain

polyunsaturated fatty acids[65]. While we also detected changes in species such as fatty

acids, acylcarnitine, and membrane lipids (Figure S4C–S4H), in line with recent

Drosophila lipidomic data[70,71], the striking accumulation of triglyceride in *bmm[1]* males

suggested that excess testis triglyceride in *bmm[1]* males contributes to their

spermatogenic defects. To test this, we examined spermatogenesis in *bmm[1]* males

carrying loss-of-function mutations in *midway* (*mdy*). *mdy* is the *Drosophila* homolog of

diacylglycerol O-acyltransferase 1 (*DGAT1*), and whole-body loss of *mdy* reduces

whole-body triglyceride levels[72–74]. Importantly, testes isolated from males lacking both

bmm and *mdy* (genotype *mdy$^{QX25/k03902}$;bmm[1]*) had fewer LD than testes dissected from

bmm[1] males (Figures 4D,S4I; one-way ANOVA with Tukey multiple comparison test).

We found that testes isolated from *mdy$^{QX25/k03902}$;bmm[1]* males were significantly

larger and had more spermatid bundles than testes from *bmm[1]* males (Figure 4E–G;

one-way ANOVA with Tukey multiple comparison test). The elevated number of GSC in

bmm[1] male testes was similarly rescued in *mdy$^{QX25/k03902}$;bmm[1]* males (Figure 4H; one-

way ANOVA with Tukey multiple comparison test). These data suggest that defective

spermatogenesis in *bmm[1]* males can be partly attributed to excess triglyceride

accumulation. Notably, at least some of these defects are cell-autonomous: RNAi-

mediated knockdown of *mdy* in the germline of *bmm[1]* males partially rescued the

defects in testis size (Figure 4I; Kruskal-Wallis rank sum test with Dunn's multiple

comparison test) and GSC variance (Figure S4J; *p*=4.5 x 10^{-5} and 8.2 x 10^{-3} by F-test

from the GAL4- and UAS-only crosses, respectively). Thus, *bmm*-mediated regulation of

testis triglyceride plays a previously unrecognized role in regulating sperm development.

Overall, our study demonstrates a key role for neutral lipids, specifically

triglyceride, in supporting *Drosophila* sperm development. We further show that loss of

bmm, a key regulator of cellular triglyceride homeostasis, delays germline differentiation

leading to an accumulation of early-stage germ cells. While studies in mammals

suggest that testis lipid droplets contain cholesterol and play a role in promoting

steroidogenesis[75,76], we show that *bmm*-dependent regulation of triglyceride-containing

LD also influences sperm development. In nongonadal cell types, triglycerides are a rich

source of fatty acids for cellular ATP production, lipid building blocks to support

membrane homeostasis and growth, and metabolites that can act as signaling

molecules[77,78]. Because all of these factors support normal sperm development, future

studies will need to test how each factor contributes to the spermatogenic defects

caused by excess triglyceride accumulation. Further studies will also be needed to

identify the upstream signaling networks that regulate cellular triglyceride metabolism in

the testis. For example, the insulin/insulin-like growth factor signaling pathway, Target of

rapamycin pathway, and nuclear factor κB/Relish pathway influence *bmm* mRNA levels

in nongonadal cell types[79–85], and contribute to sperm development[9,14,86,87]. Given the

parallel spermatogenic defects we observed in *bmm^1* flies and *ATGL* knockout mice,

these follow-up studies will likely provide vital mechanistic insight into how neutral lipids

and triglycerides support spermatogenesis in other animals.

ACKNOWLEDGEMENTS

We thank Dr. Ronald Kühnlein for *bmm¹* and *bmmʳᵉᵛ* lines[46], Dr. Michael Welte for *UAS-GFP-LD*[36], and Dr. Kaeko Kamei for *bmm-GFP*[57]. We used stocks from the Bloomington *Drosophila* Stock Center (NIH P40OD018537) and Vienna *Drosophila* Resource Center (VDRC). We acknowledge critical resources and information provided by FlyBase[88] (supported by the National Human Genome Research Institute at the U.S. National Institutes of Health (U41 HG000739) and the British Medical Research Council (MR/N030117/1)). This work was supported by the Life Sciences Institutes Imaging Core, supported by the UBC GREx Biological Resilience Initiative. Funding for this study was provided by grants to EJR from the Canadian Institutes for Health Research (PJT-153072), Michael Smith Foundation for Health Research (16876), and the Canadian Foundation for Innovation (JELF-34879). GT was supported by a grant from the Natural Sciences and Engineering Research Council (NSERC; 2018-04648), TH/HY/CW were supported by NSERC (2020-04895), MA was supported by the Jacob's foundation. We would like to acknowledge that our research takes place on the traditional, ancestral, and unceded territory of the Musqueam people; a privilege for which we are grateful.

AUTHOR CONTRIBUTIONS

Conceptualization, C.C. and E.J.R.; Methodology, C.C. and Y.Y.P.; Software, C.C.; Investigation, C.C., H.Y., and Y.Y.P.; Lipidomics, M.A., H.Y., C.W., T.H.; Writing – Original Draft, C.C. and E.J.R.; Writing – Review and Editing, C.C., E.J.R., and Y.Y.P.;

Supervision, E.J.R., G.T., and T.H.; Project administration, E.J.R.; Funding Acquisition,

E.J.R., G.T., and T.H.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1 – Lipid droplets are present in early-stage somatic and germ cells. (A)

Testis lipid droplets (LD) in w^{1118} animals visualized with neutral lipid dye BODIPY.

(A,A') Scale bar=50 µm; (A'',A''') scale bar=15 µm. Asterisk indicates hub in all images.

Arrows point to LD; arrowheads point to spermatocytes in A,B. (B) Testis LD visualized

with BODIPY in newly-eclosed males. Scale bars: main image=50 µm; inset image=10

µm. (C) Testis LD from w^{1118} animals at different times post-eclosion. Scale bars=50

µm. (D) Testis LD visualized with LipidTox Red in animals with somatic cell

overexpression of GFP-LD (*Tj-GAL4>UAS-GFP-LD*). GFP- and LipidTox Red-positive

punctae are somatic LD (D–D''' arrows); LipidTox punctae without GFP indicate

germline LD (D–D''' arrowheads). Scale bars=10 µm. (E) Histogram showing the spatial

distribution of somatic cell LD; error bars represent standard error of the mean (SEM).

(F) Cumulative frequency distributions of somatic LD (blue line, data reproduced from

E), zfh-1-positive somatic cells (zfh-1$^+$ cells, orange line), and Eya-positive somatic cells

(Eya$^+$ cells, grey line). (G) Testis LD indicated by LipidTox Red in males with germline

overexpression of GFP-LD (*nos-GAL4>UAS-GFP-LD*). GFP- and LipidTox Red-positive

punctae indicate germline LD (arrows); LipidTox punctae without GFP indicate non-

germline LD (arrowheads). Scale bars=10 μm. (H) Histogram representing the spatial

distribution of LD within germline; error bars represent SEM. (I) Histogram representing

the spatial distribution of LD and GFP fluorescence (green line) (arbitrary units, a.u.) in

a representative testis of a *bam-GFP* animal (panel J). (J) Testis LD in a *bam-GFP*

animal. Scale bar=50 μm. See also Supplemental Figure 1.

Figure 2 – *bmm* regulates testis lipid droplets in a cell-autonomous manner.　(A)

Testis lipid droplets (LD) indicated by LipidTox Red in *bmm-GFP* animals. Scale

bars=50 μm. Asterisks indicate the hub in all images. (B) Quantification of nuclear GFP

intensity in testes isolated from *bmm-GFP* animals (n=3). Germline stem cell (GSC),

spermatogonia (SG), spermatocyte (SC). (C) Spatial distribution of LD (grey histogram)

and GFP expression (green line) in testes from *bmm-GFP* animals as a function of

distance from the hub (n=3). (D,E) LD near the apical region of the testis in *bmm^rev^* (D)

or *bmm^1^* (E) animals.; (F) LD further away from the apical tip in *bmm^1^* animals. (D–F)

Scale bars=50 μm. (G) Histogram representing testis LD size distribution in *bmm^rev^*

(grey) and *bmm^1^* (orange). (H) Apical tip of the testes is at the left of the graph;

individual dots represent a single LD and its relative position to the hub marked by an

asterisk. Cumulative frequency distribution of the distance between LDs and the apical

tip of the testes are drawn as solid lines. (I) Number of testis LD in *bmm^rev^* (grey) or

*bmm^1^* (orange) in FLP-FRT clones 3 days post-clone induction; dots represent

measurements from a single clone. The number of cells in each cyst (CC) counted is

indicated. See also Supplemental Figure 2.

Figure 3 – A cell-autonomous role for *bmm* in regulating spermatogenesis. Testes

isolated from *bmm^rev* (A) and *bmm^1* (A') animals raised at 29°C stained with phalloidin.

Scale bars=100 µm. (B) Testis size in *bmm^1* and *bmm^rev* animals raised at 29°C. (C)

Spermatid bundles in *bmm^1* and *bmm^rev* testes from animals reared at 29°C. (D,E)

Representative images of *bmm^rev* (D) or *bmm^1* (E) testes stained with DAPI and anti-

Vasa antibody. Arrows indicate germline stem cells (GSC). Scale bar=50 µm. The hub

is marked by an asterisk in all images. (F) GSC number in *bmm^1* and *bmm^rev* testes. (G)

Proportion of GSCs that were either *bmm^1* or *bmm^rev* clones at 3 and 14 days post-

clone induction. (H) Representative images of *bmm^rev* (H) and *bmm^1* (H') testes carrying

bam-GFP; data quantified in Figure S3J. Arrows indicate regions with high Bam-GFP.

Scale bars=50 µm. (I) Representative images of *bmm^rev* (I) or *bmm^1* (I',I'') testes stained

with anti-Vasa antibody. Arrows indicate Vasa-positive cysts in *bmm^1* testis. Panel I'' is

magnified from the boxed region in I'. (I,I') Scale bars=100 µm; (I'') scale bar=50 µm. (J)

Maximum projection of *bmm^rev* (J) or *bmm^1* (J') testes stained with anti-Boule antibody

(green) and DAPI (blue). Scale bars=100 µm. Number of *bmm^1* and *bmm^rev*

spermatocyte clones (K) or post-meiotic clones (L) at 14 days post-clone induction. See

also Supplemental Figure 3.

Figure 4 – Loss of *bmm* disrupts triglyceride homeostasis and leads to

spermatogenic defects. (A) Hierarchical clustering of lipid species detected in *bmm^rev*

and *bmm^1* animals. (B) Histograms showing the proportion of species in each lipid class

with different levels between *bmm^1* and *bmm^rev*. Numbers on histograms indicate the

number of species with differences in abundance. (C) Volcano plot showing fold change

in abundance of triglycerides (green; 97 species) and non-triglyceride lipids (grey; 186

species) in our dataset. (D) Arrows indicate testis LD stained with LipidTox Red in

bmm^rev (D), *bmm^1* (D'), or *mdy^QX25/k03902*; *bmm^1* (D") animals. (E) Whole testes isolated

from *bmm^rev* (E), *bmm^1* (E'), or *mdy^QX25/k03902*;*bmm^1* (E") animals stained with anti-Vasa

antibody (red) and DAPI (blue). Arrowheads indicate spermatid bundles. Scale

bars=100 µm. (F) Testis size in *bmm^rev*, *bmm^1*, and *mdy^QX25/k03902*;*bmm^1* animals.

Spermatid bundles (G) and number of germline stem cells (H) in *bmm^rev*, *bmm^1*, and

mdy^QX25/k03902;*bmm^1* animals. (I) Testis size in animals with germline-specific *mdy*

knockdown (*nos-GAL4>mdy RNAi*; *bmm^1*) compared with controls (*nos-GAL4>+*; *bmm^1*

and *+>mdy RNAi*; *bmm^1*). See also Supplemental Figure 4.

STAR METHODS

RESOURCE AVAILABILITY

Lead contact. Further information and requests for resources and reagents should be directed to, and will be fulfilled by, lead contact Dr. Elizabeth J. Rideout (elizabeth.rideout@ubc.ca).

Materials availability. *Drosophila* strains and their source are listed in the Key Resources table. Further information and requests for resources and reagents should be directed to, and will be fulfilled by, lead contact Dr. Elizabeth J. Rideout (elizabeth.rideout@ubc.ca).

Data and Code availability. All raw data and results of statistical tests reported in this paper are located in Supplementary files 1-4. This paper does not report original code. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Fly husbandry. Fly stocks were maintained at room temperature in 12:12 hour light:dark cycle. Unless otherwise indicated, all flies were raised at 25°C with a density of 50 larvae per 10 mL fly media. Because this project examines sperm development, we used male flies in all experiments. Fly media contained 20.5 g of sucrose (SU10,

Snow Cap), 70.9 g of Dextrose (SUG8, Snow Cap), 48.5 g of cornmeal (AO18006,

Snow Cap), 30.3 g of baker's yeast (NB10, Snow Cap), 4.55 g of agar (DR-820-25F,

SciMart), 0.5 g of calcium chloride dihydrate (CCL302.1, BioShop Canada), 0.5 g of

magnesium sulfate heptahydrate (MAG511.1, BioShop Canada), 4.9 mL of propionic

acids (P1386, Sigma-Aldrich), and 488 µL of phosphoric acid (P5811, Sigma-Aldrich)

per 1L of media. For diets with medium- or long-chain triglyceride, 4 g of coconut oil

(medium chain triglyceride) or olive oil (long chain triglyceride) was added per 100 mL of

media described above prior to cooling. Males were collected and dissected within 24

hours of eclosion unless otherwise indicated. Fixations were performed at room

temperature with 4% paraformaldehyde (CA11021-168, VWR) in PBS for 20 minutes on

a rotating platform followed by washing in PBS twice before staining. Fly strains used in

our study are listed in the Key Resources table.

METHOD DETAILS

Testis cell stage classification and measurements. Cells at an early stage of

development (stem cells and early-stage somatic and germline cells) were located in

the apical region of the testis, and were identified by their small and dense nuclei[89].

GSC were defined as Vasa-positive cells in direct contact with the hub; proliferating

GSC were identified as Vasa-positive cells in direct contact with the hub that were also

phospho-H3 positive. Cells in the testis region occupied by primary spermatocytes were

identified by their large cell size and decondensed chromosome staining occupying

three nuclear domains[89]. Spermatid bundles were identified by their condensed and

needle-shaped nuclei, which roughly corresponds to nuclei with protamine-based

chromatin[3]. Testis size was measured by quantifying the length of a line drawn down

the middle of a testis image; starting from the apical tip of the testis and ending where

the testis meets the seminal vesicle.

FLP-FRT clone induction. Adult males were collected at 3-5 days post-eclosion and

heat-shocked three times at 37°C with a 10 min rest period at room temperature

between heat-shocks. After heat-shock, the flies were incubated at room temperature

until dissection.

Immunohistochemistry. Fixed samples were rinsed three times with blocking solution

containing 0.2% bovine serum albumin (A4503, Sigma-Aldrich), 0.3% Triton-X in PBS,

then blocked for 1 hr in PBS on a rotating platform at room temperature. During the

incubation, the blocking solution was changed every 15 minutes. After blocking, the

sample were resuspended in blocking solution with the appropriate concentration of

primary antibody (see Key Resources table), and incubated overnight at 4°C. Samples

were rinsed three times with blocking solution after removing primary antibody, and

blocked for one hour on a rotating platform in blocking solution. Secondary antibody

was applied in blocking solution and left on the rotating platform at room temperature for

40 min. The sample was then rinsed with blocking solution three more times, and

washed four times for 15 min per wash in blocking solution. Testis samples were

resuspended in Vectashield mounting media with DAPI (H-1200-10, Vector Laboratory)

or SlowFade Diamond mounting media (S36972, Thermo Fisher Scientific) prior to

mounting.

Lipid droplet staining. Fixed testes were briefly permeabilized with 0.1% Triton-X in

PBS for 5 min prior to applying phalloidin. For BODIPY staining, samples were

suspended in PBS containing 10 µg/mL DAPI (2879083-5mg, PeproTech), 1:500

BODIPY 495/503 (Thermo Fisher Scientific D3922), and 1:1000 phalloidin iFluor647

(ab176759, Abcam) or 1:40 phalloidin TexasRed (T7471, Thermo Fisher Scientific). For

staining with LipidTox Red, samples were suspended in PBS containing 10 µg/mL DAPI

(2879083-5mg, PeproTech), 1:200 LipidTox Red (H34476, Thermo Fisher Scientific),

and 1:1000 phalloidin iFluor647 (ab176759, Abcam). For staining free sterols, samples

were prepared as for BODIPY staining with 50 µg/mL filipin in place of BODIPY for 30

min. Samples were incubated on a rotating platform for 40 minutes at room

temperature. After incubation, samples were washed twice with PBS, then resuspended

in SlowFade Diamond mounting media (Thermo Fisher Scientific S36972) prior to

mounting.

Image acquisition and processing. All images were acquired on a Leica SP5 confocal

microscope system with 20X or 40X objectives and subsequently quantified with Fiji

image analysis software[90].

Drosophila lipidomics. *Drosophila* extracts were prepared following the previously

reported protocol[91]. Briefly, 10 *Drosophila* males (~10 mg) were weighed, 300 µL of ice-

cold methanol/water mixture (9:1, v:v) was added to these males, and the samples were

homogenized with glass beads using a bead beater (mini-beadbeater-16, BioSpec,

Bartlesville, Ok, USA). Sample weight was used for sample normalization. Fly lysate

was kept at -20°C for 4 hours for protein precipitation. Then, 900 µL of methyl tert-butyl

ether was added and the solution was shaken for 5 min to extract lipids. To induce

phase separation 285 µL of water was added, followed by centrifugation. The upper

layer was separated, dried, and reconstituted in isopropanol/acetonitrile (1:1, v:v) for

liquid chromatography-mass spectrometry (LC-MS) analysis. The volume of

reconstitution solution was proportional to sample weight for normalization. Quality

control (QC) samples were prepared by pooling 20 µL aliquot from each sample. The

method blank sample was prepared using an identical workflow but without adding

Drosophila.

Drosophila extracts were analyzed on an UHR-QqTOF (Ultra-High Resolution

Qq-Time-Of-Flight) mass spectrometry Impact II (Bruker Daltonics, Bremen, Germany)

interfaced with an Agilent 1290 Infinity II LC Systems (Agilent Technologies, Santa

Clara, CA, USA). LC separation was performed using a Waters reversed-phase (RP)

UPLC Acquity BEH C18 Column (1.7 µm, 1.0 mm ×100 mm, 130 Å) (Milford, MA, USA)

maintained at 30°C. For positive ion mode, the mobile phase A was 60% acetonitrile in

water and the mobile phase B was 90% isopropanol in acetonitrile, both containing 5

mM ammonium formate (pH = 4.8, adjusted by formic acid). For negative ion mode, the

mobile phase A was 60% acetonitrile in water and the mobile phase B was 90%

isopropanol in acetonitrile, both containing 5 mM NH_4FA (pH = 9.8, adjusted by

ammonium hydroxide). The LC gradient for positive and negative ion modes was set as

follow: 0 min, 5% B; 8 min, 40% B; 14 min, 70% B; 20 min, 95% B; 23 min, 95% B; 24

min, 5% B; 33 min, 5% B. The flow rate was 0.1 mL/min. The injection volume was

optimized to 2 μL in positive mode and 5 μL in negative mode using QC sample. The

ESI source conditions were set as follows: dry gas temperature, 220 °C; dry gas flow, 7

L/min; nebulizer gas pressure, 1.6 bar; capillary voltage, 4500 V for positive mode and

3000 V for negative mode. The MS1 analysis was conducted using following

parameters: mass range, 70-1000 m/z; spectrum type: centroid, calculated using

maximum intensity; absolute intensity threshold: 250. Data-dependent MS/MS analysis

parameters: collision energy: 16-30 eV; cycle time, 3 s; spectra rate: 4 Hz when

intensity $< 10^4$ and 12 Hz when intensity $> 10^5$, linearly increased from 10^4 to 10^5.

External calibration was applied using sodium formate to ensure the m/z accuracy

before sample analysis.

The raw LC-MS data were processed using MS-DIAL (ver. 4.38)[92]. The detailed

MS-DIAL parameters are: MS1 tolerance, 0.01 Da; MS/MS tolerance, 0.05; mass slice

width, 0.05 Da; smoothing method, linear weighted moving average; smoothing level, 3

scans; minimum peak width, 5 scans. Lipid features with high quantitative confidence

were selected by the following criteria: retention time was within the gradient elution

time (< 23 min); average intensity in QC samples is larger than 5-fold of the intensity in

method blank sample. Lipid identification was performed by matching experimental

precursor m/z, isotopic ratio and MS/MS spectrum against the LipidBlast libraries

embedded in MS-DIAL. To improve the quantification accuracy, the measured MS

signal intensities were corrected using serial diluted QC samples following the reported

workflow[93].

QUANTIFICATION AND STATISTICAL ANALYSIS

All microscopy images were quantified using Fiji software[90]. For lipid droplet counts, a

single optical slice through the middle of the testis containing the hub was used with the

exception of FLP-FRT experiment where all lipid droplets within a GFP-negative cyst

were counted (Figure 2I).

All statistical analyses were done using R (obtained from https://cran.r-project.org). With

exception of data concerning spatial distribution, and lipidomic data, Shapiro-Wilk test

(via *shapiro.test* in base R) was used to assess normality of distribution prior to testing

for significance. Kruskal-Wallis rank sum test (from the R package *coin*) and Dunn's test

(from the R package *dunn.test*) were used in place of Welch two-sample t-test and

Tukey's multiple comparison test when the assumption of normality was not met. For

testing differences in variance between two populations, F-test (via *var.test* in base R)

was used. For testing differences in spatial distribution, two-sample Kolmogorov-

Smirnov test (via *ks.test* in base R) was used. All *p*-values are indicated in figures;

extremely small *p*-values are listed as $p < 2.2 \times 10^{-16}$.

SUPPLEMENTAL TABLE AND FILES

Supplemental table 1 – Table showing identified lipid species from untargeted

lipidomic analysis.

Supplementary file 1 – Raw data and statistical outputs from Figure 1.

Supplementary file 2 – Raw data and statistical outputs from Figure 2 and

Supplemental figure 2.

Supplementary file 3 – Raw data and statistical outputs from Figure 3 and

Supplemental figure 3.

Supplementary file 4 – Raw data and statistical outputs from Figure 4 and

Supplemental figure 4.

REFERENCES

1. de Cuevas, M., and Matunis, E.L. (2011). The stem cell niche: lessons from the Drosophila testis. Dev. Camb. Engl. *138*, 2861–2869. 10.1242/dev.056242.

2. Demarco, R.S., Eikenes, Å.H., Haglund, K., and Jones, D.L. (2014). Investigating spermatogenesis in Drosophila melanogaster. Methods San Diego Calif *68*, 218–227. 10.1016/j.ymeth.2014.04.020.

3. Fabian, L., and Brill, J.A. (2012). Drosophila spermiogenesis. Spermatogenesis *2*, 197–212. 10.4161/spmg.21798.

4. Brill, J.A., Yildirim, S., and Fabian, L. (2016). Phosphoinositide signaling in sperm development. Semin. Cell Dev. Biol. *59*, 2–9. 10.1016/j.semcdb.2016.06.010.

5. Brill, J.A., Hime, G.R., Scharer-Schuksz, M., and Fuller, M.T. (2000). A phospholipid kinase regulates actin organization and intercellular bridge formation during germline cytokinesis. Development *127*, 3855.

6. Giansanti, M.G., Bonaccorsi, S., Kurek, R., Farkas, R.M., Dimitri, P., Fuller, M.T., and Gatti, M. (2006). The Class I PITP Giotto Is Required for Drosophila Cytokinesis. Curr. Biol. *16*, 195–201. 10.1016/j.cub.2005.12.011.

7. Wong, R., Hadjiyanni, I., Wei, H.-C., Polevoy, G., McBride, R., Sem, K.-P., and Brill, J.A. (2005). PIP2 Hydrolysis and Calcium Release Are Required for Cytokinesis in Drosophila Spermatocytes. Curr. Biol. *15*, 1401–1406. 10.1016/j.cub.2005.06.060.

8. Wong, R., Fabian, L., Forer, A., and Brill, J.A. (2007). Phospholipase C and myosin light chain kinase inhibition define a common step in actin regulation during cytokinesis. BMC Cell Biol. *8*, 15. 10.1186/1471-2121-8-15.

9. Amoyel, M., Hillion, K.-H., Margolis, S.R., and Bach, E.A. (2016). Somatic stem cell differentiation is regulated by PI3K/Tor signaling in response to local cues. Dev. Camb. Engl. *143*, 3914–3925. 10.1242/dev.139782.

10. Fairchild, M.J., Islam, F., and Tanentzapf, G. (2017). Identification of genetic networks that act in the somatic cells of the testis to mediate the developmental program of spermatogenesis. PLOS Genet. *13*, e1007026. 10.1371/journal.pgen.1007026.

11. Inaba, M., Venkei, Z.G., and Yamashita, Y.M. (2015). The polarity protein Baz forms a platform for the centrosome orientation during asymmetric stem cell division in the Drosophila male germline. eLife *4*, e04960. 10.7554/eLife.04960.

12. Krahn, M.P., Klopfenstein, D.R., Fischer, N., and Wodarz, A. (2010). Membrane Targeting of Bazooka/PAR-3 Is Mediated by Direct Binding to Phosphoinositide Lipids. Curr. Biol. *20*, 636–642. 10.1016/j.cub.2010.01.065.

13. Papagiannouli, F., Berry, C.W., and Fuller, M.T. (2019). The Dlg Module and
Clathrin-Mediated Endocytosis Regulate EGFR Signaling and Cyst Cell-Germline
Coordination in the Drosophila Testis. Stem Cell Rep. *12*, 1024–1040.
10.1016/j.stemcr.2019.03.008.

14. Ueishi, S., Shimizu, H., and H. Inoue, Y. (2009). Male Germline Stem Cell Division
and Spermatocyte Growth Require Insulin Signaling in Drosophila. Cell Struct.
Funct. *34*, 61–69. 10.1247/csf.08042.

15. Phan, V.H., Herr, D.R., Panton, D., Fyrst, H., Saba, J.D., and Harris, G.L. (2007).
Disruption of Sphingolipid Metabolism Elicits Apoptosis-Associated Reproductive
Defects in Drosophila. Dev. Biol. *309*, 329–341. 10.1016/j.ydbio.2007.07.021.

16. Steinhauer, J., Gijón, M.A., Riekhof, W.R., Voelker, D.R., Murphy, R.C., and
Treisman, J.E. (2009). Drosophila Lysophospholipid Acyltransferases Are
Specifically Required for Germ Cell Development. Mol. Biol. Cell *20*, 5224–5235.
10.1091/mbc.e09-05-0382.

17. Szafer-Glusman, E., Giansanti, M.G., Nishihama, R., Bolival, B., Pringle, J., Gatti,
M., and Fuller, M.T. (2008). A Role for Very-Long-Chain Fatty Acids in Furrow
Ingression during Cytokinesis in Drosophila Spermatocytes. Curr. Biol. *18*, 1426–
1431. 10.1016/j.cub.2008.08.061.

18. Ben-David, G., Miller, E., and Steinhauer, J. (2015). Drosophila spermatid
individualization is sensitive to temperature and fatty acid metabolism.
Spermatogenesis *5*. 10.1080/21565562.2015.1006089.

19. Jung, A., Hollmann, M., and Schafer, M.A. (2007). The fatty acid elongase NOA is
necessary for viability and has a somatic role in Drosophila sperm development. J.
Cell Sci. *120*, 2924–2934. 10.1242/jcs.006551.

20. Walther, T.C., and Farese, R.V. (2012). Lipid Droplets and Cellular Lipid
Metabolism. Annu. Rev. Biochem. *81*, 687–714. 10.1146/annurev-biochem-061009-
102430.

21. Wat, L.W., Chao, C., Bartlett, R., Buchanan, J.L., Millington, J.W., Chih, H.J.,
Chowdhury, Z.S., Biswas, P., Huang, V., Shin, L.J., et al. (2020). A role for
triglyceride lipase brummer in the regulation of sex differences in Drosophila fat
storage and breakdown. PLOS Biol. *18*, e3000595. 10.1371/journal.pbio.3000595.

22. Bajpai, M., Gupta, G., and Setty, B.S. (1998). Changes in carbohydrate metabolism
of testicular germ cells during meiosis in the rat. Eur. J. Endocrinol. *138*, 322–327.
10.1530/eje.0.1380322.

23. Paniagua, R., Rodriguez, M.C., Nistal, M., Fraile, B., and Amat, P. (1987). Changes
in the lipid inclusion/Sertoli cell cytoplasm area ratio during the cycle of the human
seminiferous epithelium. Reproduction *80*, 335–341. 10.1530/jrf.0.0800335.

24. Mori, H., and Christensen, A.K. (1980). Morphometric analysis of Leydig cells in the
normal rat testis. J. Cell Biol. *84*, 340–354. 10.1083/jcb.84.2.340.

25. Kerr, J.B., and De Kretser, D.M. (1975). CYCLIC VARIATIONS IN SERTOLI CELL
LIPID CONTENT THROUGHOUT THE SPERMATOGENIC CYCLE IN THE RAT.
Reproduction *43*, 1–8. 10.1530/jrf.0.0430001.

26. Demarco, R.S., Uyemura, B.S., D'Alterio, C., and Jones, D.L. (2019). Mitochondrial
fusion regulates lipid homeostasis and stem cell maintenance in the Drosophila
testis. Nat. Cell Biol. *21*, 710–720. 10.1038/s41556-019-0332-3.

27. Casado, M.E., Pastor, O., Mariscal, P., Canfrán-Duque, A., Martínez-Botas, J.,
Kraemer, F.B., Lasunción, M.A., Martín-Hidalgo, A., and Busto, R. (2013). Hormone-
sensitive lipase deficiency disturbs the fatty acid composition of mouse testis.
Prostaglandins Leukot. Essent. Fatty Acids *88*, 227–233.
10.1016/j.plefa.2012.12.005.

28. El-Shehawi, A.M., El-Shazly, S., Ahmed, M., Alkafafy, M., Sayed, S., Farouk, S.,
Alotaibi, S.S., and Elseehy, M.M. (2020). Transcriptome Analysis of Testis from
HFD-Induced Obese Rats (Rattus norvigicus) Indicated Predisposition for Male
Infertility. Int. J. Mol. Sci. *21*, 6493. 10.3390/ijms21186493.

29. Wang, W., Wei, S., Li, L., Su, X., Du, C., Li, F., Geng, B., Liu, P., and Xu, G. (2015).
Proteomic analysis of murine testes lipid droplets. Sci. Rep. *5*, 12070.
10.1038/srep12070.

30. Chen, M., Wang, H., Li, X., Li, N., Xu, G., and Meng, Q. (2014). PLIN1 deficiency
affects testicular gene expression at the meiotic stage in the first wave of
spermatogenesis. Gene *543*, 212–219. 10.1016/j.gene.2014.04.021.

31. Hermo, L., Chung, S., Gregory, M., Smith, C.E., Wang, S.P., El-Alfy, M., Cyr, D.G.,
Mitchell, G.A., and Trasler, J. (2008). Alterations in the testis of hormone sensitive
lipase-deficient mice is associated with decreased sperm counts, sperm motility, and
fertility. Mol. Reprod. Dev. *75*, 565–577. 10.1002/mrd.20800.

32. Masaki, H., Kim, N., Nakamura, H., Kumasawa, K., Kamata, E., Hirano, K., and
Kimura, T. (2017). Long-chain fatty acid triglyceride (TG) metabolism disorder
impairs male fertility: a study using adipose triglyceride lipase deficient mice. MHR
Basic Sci. Reprod. Med. *23*, 452–460. 10.1093/molehr/gax031.

33. de Cuevas, M., and Matunis, E.L. (2011). The stem cell niche: lessons from the
Drosophila testis. Development *138*, 2861–2869. 10.1242/dev.056242.

34. Walther, T.C., Chung, J., and Jr, R.V.F. (2017). Lipid Droplet Biogenesis. Annu.
Rev. Cell Dev. Biol. *33*, 491–510. 10.1146/annurev-cellbio-100616-060608.

35. Boyle, M., and DiNardo, S. (1995). Specification, migration and assembly of the
somatic cells of the Drosophila gonad. Development *121*, 1815–1825.
10.1242/dev.121.6.1815.

36. Yu, Y.V., Li, Z., Rizzo, N.P., Einstein, J., and Welte, M.A. (2011). Targeting the
motor regulator Klar to lipid droplets. BMC Cell Biol. *12*, 9. 10.1186/1471-2121-12-9.

37. Leatherman, J.L., and DiNardo, S. (2008). Zfh-1 Controls Somatic Stem Cell Self-
Renewal in the Drosophila Testis and Nonautonomously Influences Germline Stem
Cell Self-Renewal. Cell Stem Cell *3*, 44–54. 10.1016/j.stem.2008.05.001.

38. Amoyel, M., Hillion, K.-H., Margolis, S.R., and Bach, E.A. (2016). Somatic stem cell
differentiation is regulated by PI3K/Tor signaling in response to local cues. Dev.
Camb. Engl. *143*, 3914. 10.1242/dev.139782.

39. Fabrizio, J.J., Boyle, M., and DiNardo, S. (2003). A somatic role for eyes absent
(eya) and sine oculis (so) in drosophila spermatocyte development. Dev. Biol. *258*,
117–128. 10.1016/S0012-1606(03)00127-1.

40. Chen, D., and McKearin, D.M. (2003). A discrete transcriptional silencer in the bam
gene determines asymmetric division of the Drosophila germline stem cell.
Development *130*, 1159–1170. 10.1242/dev.00325.

41. Insco, M.L., Leon, A., Tam, C.H., McKearin, D.M., and Fuller, M.T. (2009).
Accumulation of a differentiation regulator specifies transit amplifying division
number in an adult stem cell lineage. Proc. Natl. Acad. Sci. U. S. A. *106*, 22311–
22316. 10.1073/pnas.0912454106.

42. Chen, D., Wu, C., Zhao, S., Geng, Q., Gao, Y., Li, X., Zhang, Y., and Wang, Z.
(2014). Three RNA Binding Proteins Form a Complex to Promote Differentiation of
Germline Stem Cell Lineage in Drosophila. PLoS Genet. *10*.
10.1371/journal.pgen.1004797.

43. Gonczy, P., Matunis, E., and DiNardo, S. (1997). bag-of-marbles and benign gonial
cell neoplasm act in the germline to restrict proliferation during Drosophila
spermatogenesis. Development *124*, 4361–4371.

44. Athenstaedt, K., and Daum, G. (2003). YMR313c/TGL3 Encodes a Novel
Triacylglycerol Lipase Located in Lipid Particles of Saccharomyces cerevisiae*. J.
Biol. Chem. *278*, 23317–23323. 10.1074/jbc.M302577200.

45. Chitraju, C., Trötzmüller, M., Hartler, J., Wolinski, H., Thallinger, G.G., Haemmerle,
G., Zechner, R., Zimmermann, R., Köfeler, H.C., and Spener, F. (2013). The impact
of genetic stress by ATGL deficiency on the lipidome of lipid droplets from murine
hepatocytes. J. Lipid Res. *54*, 2185–2194. 10.1194/jlr.M037952.

46. Grönke, S., Mildner, A., Fellert, S., Tennagels, N., Petry, S., Müller, G., Jäckle, H.,
and Kühnlein, R.P. (2005). Brummer lipase is an evolutionary conserved fat storage
regulator in Drosophila. Cell Metab. *1*, 323–330. 10.1016/j.cmet.2005.04.003.

47. Haemmerle, G., Lass, A., Zimmermann, R., Gorkiewicz, G., Meyer, C., Rozman, J.,
Heldmaier, G., Maier, R., Theussl, C., Eder, S., et al. (2006). Defective Lipolysis and
Altered Energy Metabolism in Mice Lacking Adipose Triglyceride Lipase. Science
312, 734–737. 10.1126/science.1123965.

48. Haemmerle, G., Moustafa, T., Woelkart, G., Büttner, S., Schmidt, A., van de Weijer,
T., Hesselink, M., Jaeger, D., Kienesberger, P.C., Zierler, K., et al. (2011). ATGL-
mediated fat catabolism regulates cardiac mitochondrial function via PPAR-α and
PGC-1. Nat. Med. *17*, 1076–1085. 10.1038/nm.2439.

49. Huijsman, E., van de Par, C., Economou, C., van der Poel, C., Lynch, G.S.,
Schoiswohl, G., Haemmerle, G., Zechner, R., and Watt, M.J. (2009). Adipose
triacylglycerol lipase deletion alters whole body energy metabolism and impairs
exercise performance in mice. Am. J. Physiol.-Endocrinol. Metab. *297*, E505–E513.
10.1152/ajpendo.00190.2009.

50. Korbelius, M., Vujic, N., Sachdev, V., Obrowsky, S., Rainer, S., Gottschalk, B.,
Graier, W.F., and Kratky, D. (2019). ATGL/CGI-58-Dependent Hydrolysis of a Lipid
Storage Pool in Murine Enterocytes. Cell Rep. *28*, 1923-1934.e4.
10.1016/j.celrep.2019.07.030.

51. Kurat, C.F., Natter, K., Petschnigg, J., Wolinski, H., Scheuringer, K., Scholz, H.,
Zimmermann, R., Leber, R., Zechner, R., and Kohlwein, S.D. (2006). Obese Yeast:
Triglyceride Lipolysis Is Functionally Conserved from Mammals to Yeast *. J. Biol.
Chem. *281*, 491–500. 10.1074/jbc.M508414200.

52. Zimmermann, R., Strauss, J.G., Haemmerle, G., Schoiswohl, G., Birner-
Gruenberger, R., Riederer, M., Lass, A., Neuberger, G., Eisenhaber, F., Hermetter,
A., et al. (2004). Fat Mobilization in Adipose Tissue Is Promoted by Adipose
Triglyceride Lipase. Science *306*, 1383–1386. 10.1126/science.1100747.

53. Attané, C., Peyot, M.-L., Lussier, R., Poursharifi, P., Zhao, S., Zhang, D., Morin, J.,
Pineda, M., Wang, S., Dumortier, O., et al. (2016). A beta cell ATGL-
lipolysis/adipose tissue axis controls energy homeostasis and body weight via
insulin secretion in mice. Diabetologia *59*, 2654–2663. 10.1007/s00125-016-4105-2.

54. Grönke, S., Müller, G., Hirsch, J., Fellert, S., Andreou, A., Haase, T., Jäckle, H., and
Kühnlein, R.P. (2007). Dual Lipolytic Control of Body Fat Storage and Mobilization in
Drosophila. PLOS Biol. *5*, e137. 10.1371/journal.pbio.0050137.

55. Lee, J.H., Kong, J., Jang, J.Y., Han, J.S., Ji, Y., Lee, J., and Kim, J.B. (2014). Lipid
Droplet Protein LID-1 Mediates ATGL-1-Dependent Lipolysis during Fasting in
Caenorhabditis elegans. Mol. Cell. Biol. *34*, 4165–4176. 10.1128/MCB.00722-14.

56. Tuohetahuntila, M., Molenaar, M.R., Spee, B., Brouwers, J.F., Houweling, M.,
Vaandrager, A.B., and Helms, J.B. (2016). ATGL and DGAT1 are involved in the
turnover of newly synthesized triacylglycerols in hepatic stellate cells. J. Lipid Res.
57, 1162–1174. 10.1194/jlr.M066415.

57. Men, T.T., Thanh, D.N.V., Yamaguchi, M., Suzuki, T., Hattori, G., Arii, M., Huy, N.T.,
and Kamei, K. (2016). A *Drosophila* Model for Screening Antiobesity Agents.
BioMed Res. Int. *2016*, 1–10. 10.1155/2016/6293163.

58. Li, H., Janssens, J., Waegeneer, M.D., Kolluru, S.S., Davie, K., Gardeux, V.,
Saelens, W., David, F., Brbić, M., Leskovec, J., et al. (2021). Fly Cell Atlas: a single-
cell transcriptomic atlas of the adult fruit fly. 2021.07.04.451050.
10.1101/2021.07.04.451050.

59. Xu, T., and Rubin, G.M. (1993). Analysis of genetic mosaics in developing and adult
Drosophila tissues. Development *117*, 1223–1237.

60. Kim, N., Nakamura, H., Masaki, H., Kumasawa, K., Hirano, K., and Kimura, T.
(2017). Effect of lipid metabolism on male fertility. Biochem. Biophys. Res. Commun.
485, 686–692. 10.1016/j.bbrc.2017.02.103.

61. Lasko, P.F., and Ashburner, M. (1988). The product of the Drosophila gene vasa is
very similar to eukaryotic initiation factor-4A. Nature *335*, 611–617.
10.1038/335611a0.

62. Resende, L.P.F., Boyle, M., Tran, D., Fellner, T., and Jones, D.L. (2013). Headcase
Promotes Cell Survival and Niche Maintenance in the Drosophila Testis. PLoS ONE
8, e68026. 10.1371/journal.pone.0068026.

63. Kiger, A.A., White-Cooper, H., and Fuller, M.T. (2000). Somatic support cells restrict
germline stem cell self-renewal and promote differentiation. Nature *407*, 750–754.
10.1038/35037606.

64. Lin, T.Y., Viswanathan, S., Wood, C., Wilson, P.G., Wolf, N., and Fuller, M.T.
(1996). Coordinate developmental control of the meiotic cell cycle and spermatid
differentiation in Drosophila males. Development *122*, 1331–1341.

65. Eichmann, T.O., Kumari, M., Haas, J.T., Farese, R.V., Zimmermann, R., Lass, A.,
and Zechner, R. (2012). Studies on the substrate and stereo/regioselectivity of
adipose triglyceride lipase, hormone-sensitive lipase, and diacylglycerol-O-
acyltransferases. J. Biol. Chem. *287*, 41446–41457. 10.1074/jbc.M112.400416.

66. Schweiger, M., Schreiber, R., Haemmerle, G., Lass, A., Fledelius, C., Jacobsen, P.,
Tornqvist, H., Zechner, R., and Zimmermann, R. (2006). Adipose Triglyceride
Lipase and Hormone-sensitive Lipase Are the Major Enzymes in Adipose Tissue
Triacylglycerol Catabolism *. J. Biol. Chem. *281*, 40236–40241.
10.1074/jbc.M608048200.

67. Missaglia, S., Maggi, L., Mora, M., Gibertini, S., Blasevich, F., Agostoni, P., Moro, L.,
Cassandrini, D., Santorelli, F.M., Gerevini, S., et al. (2017). Late onset of neutral
lipid storage disease due to novel PNPLA2 mutations causing total loss of lipase
activity in a patient with myopathy and slight cardiac involvement. Neuromuscul.
Disord. *27*, 481–486. 10.1016/j.nmd.2017.01.011.

68. Williams, M.L., Coleman, R.A., Placezk, D., and Grunfeld, C. (1991). Neutral lipid
storage disease: a possible functional defect in phospholipid-linked triacylglycerol
metabolism. Biochim. Biophys. Acta BBA - Mol. Basis Dis. *1096*, 162–169.
10.1016/0925-4439(91)90055-E.

69. Yang, L., Liang, J., Lam, S.M., Yavuz, A., Shui, G., Ding, M., and Huang, X. (2020).
Neuronal lipolysis participates in PUFA-mediated neural function and
neurodegeneration. EMBO Rep. *21*. 10.15252/embr.202050214.

70. Nazario-Yepiz, N.O., Fernández Sobaberas, J., Lyman, R., Campbell, M.R.,
Shankar, V., Anholt, R.R.H., and Mackay, T.F.C. (2021). Physiological and
metabolomic consequences of reduced expression of the Drosophila brummer
triglyceride Lipase. PLOS ONE *16*, e0255198. 10.1371/journal.pone.0255198.

71. Giedt, M.S., Thomalla, J.M., Johnson, M.R., Lai, Z.W., Tootle, T.L., and Welte, M.A.
(2021). Adipose triglyceride lipase promotes prostaglandin-dependent actin
remodeling by regulating substrate release from lipid droplets (Cell Biology)
10.1101/2021.08.02.454724.

72. Beller, M., Bulankina, A.V., Hsiao, H.-H., Urlaub, H., Jäckle, H., and Kühnlein, R.P.
(2010). PERILIPIN-Dependent Control of Lipid Droplet Structure and Fat Storage in
Drosophila. Cell Metab. *12*, 521–532. 10.1016/j.cmet.2010.10.001.

73. Buszczak, M., Lu, X., Segraves, W.A., Chang, T.Y., and Cooley, L. (2002).
Mutations in the midway Gene Disrupt a Drosophila Acyl Coenzyme A:
Diacylglycerol Acyltransferase. Genetics *160*, 1511–1518.

74. Martínez, B.A., Hoyle, R.G., Yeudall, S., Granade, M.E., Harris, T.E., Castle, J.D.,
Leitinger, N., and Bland, M.L. (2020). Innate immune signaling in Drosophila shifts
anabolic lipid metabolism from triglyceride storage to phospholipid synthesis to
support immune function. PLOS Genet. *16*, e1009192.
10.1371/journal.pgen.1009192.

75. Freeman, D.A., and Ascoli, M. (1982). Studies on the source of cholesterol used for
steroid biosynthesis in cultured Leydig tumor cells. J. Biol. Chem. *257*, 14231–
14238.

76. Wang, F., Chen, Z., Ren, X., Tian, Y., Wang, F., Liu, C., Jin, P., Li, Z., Zhang, F.,
and Zhu, B. (2017). Hormone-sensitive lipase deficiency alters gene expression and
cholesterol content of mouse testis. Reproduction *153*, 175–185. 10.1530/REP-16-
0484.

77. Walther, T.C., and Farese, R.V. (2012). Lipid droplets and cellular lipid metabolism.
Annu Rev Biochem *81*, 687–714. 10.1146/annurev-biochem-061009-102430.

78. Olzmann, J.A., and Carvalho, P. (2019). Dynamics and functions of lipid droplets.
Nat. Rev. Mol. Cell Biol. *20*, 137–155. 10.1038/s41580-018-0085-z.

79. Birse, R.T., Choi, J., Reardon, K., Rodriguez, J., Graham, S., Diop, S., Ocorr, K.,
Bodmer, R., and Oldham, S. (2010). High-fat-diet-induced obesity and heart
dysfunction are regulated by the TOR pathway in Drosophila. Cell Metab *12*, 533–
544. 10.1016/j.cmet.2010.09.014.

80. Molaei, M., Vandehoef, C., and Karpac, J. (2019). NF-κB Shapes Metabolic
Adaptation by Attenuating Foxo-Mediated Lipolysis in Drosophila. Dev. Cell *49*, 802-
810.e6. 10.1016/j.devcel.2019.04.009.

81. Alic, N., Andrews, T.D., Giannakou, M.E., Papatheodorou, I., Slack, C., Hoddinott,
M.P., Cocheme, H.M., Schuster, E.F., Thornton, J.M., and Partridge, L. (2011).
Genome-wide dFOXO targets and topology of the transcriptomic response to stress
and insulin signalling. Mol Syst Biol *7*, 502. 10.1038/msb.2011.36.

82. Junger, M.A., Rintelen, F., Stocker, H., Wasserman, J.D., Vegh, M., Radimerski, T.,
Greenberg, M.E., and Hafen, E. (2003). The Drosophila forkhead transcription factor
FOXO mediates the reduction in cell number associated with reduced insulin
signaling. J Biol *2*, 20. 10.1186/1475-4924-2-20.

83. Zinke, I., Schutz, C.S., Katzenberger, J.D., Bauer, M., and Pankratz, M.J. (2002).
Nutrient control of gene expression in Drosophila: microarray analysis of starvation
and sugar-dependent response. EMBO J *21*, 6162–6173.

84. Puig, O., and Tjian, R. (2005). Transcriptional feedback control of insulin receptor by
dFOXO/FOXO1. Genes Dev. *19*, 2435–2446. 10.1101/gad.1340505.

85. Kang, P., Chang, K., Liu, Y., Bouska, M., Birnbaum, A., Karashchuk, G., Thakore,
R., Zheng, W., Post, S., Brent, C.S., et al. (2017). Drosophila Kruppel homolog 1
represses lipolysis through interaction with dFOXO. Sci. Rep. *7*, 16369.
10.1038/s41598-017-16638-1.

86. Amoyel, M., Simons, B.D., and Bach, E.A. (2014). Neutral competition of stem cells
is skewed by proliferative changes downstream of Hh and Hpo. EMBO J. *33*, 2295–
2313. 10.15252/embj.201387500.

87. Hof-Michel, S., Cigoja, L., Huhn, S., and Bökel, C. (2020). Innate immune signalling
drives loser cell elimination during stem cell competition in the *Drosophila* testis
(Developmental Biology) 10.1101/2020.03.05.979161.

88. Thurmond, J., Goodman, J.L., Strelets, V.B., Attrill, H., Gramates, L.S., Marygold,
S.J., Matthews, B.B., Millburn, G., Antonazzo, G., Trovisco, V., et al. (2018).

FlyBase 2.0: the next generation. Nucleic Acids Res. *47*, D759–D765.
10.1093/nar/gky1003.

89. White-Cooper, H. (2004). Spermatogenesis. In Drosophila Cytogenetics Protocols
Methods in Molecular Biology., D. S. Henderson, ed. (Humana Press), pp. 45–75.
10.1385/1-59259-665-7:45.

90. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-
source platform for biological-image analysis. Nat. Methods *9*, 676–682.
10.1038/nmeth.2019.

91. Yu, H., Villanueva, N., Bittar, T., Arsenault, E., Labonté, B., and Huan, T. (2020).
Parallel metabolomics and lipidomics enables the comprehensive study of mouse
brain regional metabolite and lipid patterns. Anal. Chim. Acta *1136*, 168–177.
10.1016/j.aca.2020.09.051.

92. Tsugawa, H., Cajka, T., Kind, T., Ma, Y., Higgins, B., Ikeda, K., Kanazawa, M.,
VanderGheynst, J., Fiehn, O., and Arita, M. (2015). MS-DIAL: data-independent
MS/MS deconvolution for comprehensive metabolome analysis. Nat. Methods *12*,
523–526. 10.1038/nmeth.3393.

93. Yu, H., and Huan, T. (2021). Patterned Signal Ratio Biases in Mass Spectrometry-
Based Quantitative Metabolomics. Anal. Chem. *93*, 2254–2262.
10.1021/acs.analchem.0c04113.

SUPPLEMENTAL FIGURE LEGENDS

Supplemental Figure 1 related to Figure 1 – Cholesterol is absent from testis lipid

droplets. (A) Testes stained with BODIPY (A) to detect neutral lipids and Filipin III (A')

to detect free cholesterol. Scale bars=50 µm.

Supplemental Figure 2 related to Figure 2 – *bmm* mRNA levels during

spermatogenesis in germline and somatic lineages. (A) Pseudotime trajectory of

germline (black line) based on single-cell RNA sequencing data[62]. Individual cells are

labeled according to the annotation within the data set. (B) Rolling average of

normalized *bmm* transcript counts in the germline along the trajectory shown in panel A

are plotted as a black line on the upper panel. Composition of cell types mapped on to

the trajectory at each time point is shown at the lower half of panel B. (C) Pseudotime

trajectory of the somatic cells (black line) based on publicly available single-cell RNA

sequencing data[62]. Individual cells are labeled according to the annotation within the

data set. (D) Rolling average of normalized *bmm* transcript counts in somatic cells

plotted as a black line along the trajectory shown in C (upper panel). Composition of cell

types mapped on to the trajectory at each time point (lower panel). (E-H)

Representative images of *bmm^rev^* (E and F) and *bmm^1^* (G and H) testes with somatic

over-expression of *GFP-LD* (*Tj-GAL4>UAS-GFP-LD*). Panel F and H contain magnified

images of the area indicated by the boxes in panel E and G, respectively. In *bmm^rev^*

testes, LD were restricted to a region near the apical tip (E) of the testis in both somatic

(F–F''' arrows) and germline cells (F–F''' arrowheads). In *bmm^1^* testes, LD were present

in both somatic (G–H arrows) and germline cells (G–H arrowheads), near the apical tip

of the testis in a region corresponding to early-stage germ cells, and in the region

corresponding to spermatocytes (G arrows and arrowheads, and H). Scale bars=50 μm;

(F and H) scale bars=20 μm.

Supplemental Figure 3 related to Figure 3 – Additional characterization of testis

development and spermatogenesis in animals lacking *bmm*. (A) Testis size was

smaller in *bmm^1^* mutant animals compared with *bmm^rev^* controls at <24 hr post-eclosion

when raised at 25°C (A; Welch two-sample t-test). (B) The number of spermatid bundles

was significantly lower in *bmm^1^* mutant animals compared with *bmm^rev^* controls at <24

hr post-eclosion when raised at 25°C (Kruskal-Wallis rank sum test). (C) Testis size was

significantly smaller in *bmm¹* mutant males compared with *bmm^rev* control males at 14-

days post-eclosion (Welch two sample t-test). (D) The median number of spermatid

bundles was not significantly different between *bmm¹* mutant males and *bmm^rev* control

males at 14 days post-eclosion (Welch two sample t-test), but 8/27 *bmm¹* testis had no

spermatid bundles, a phenotype we did not observe in age-matched *bmm^rev* males

(0/22) (p=0.0163, Pearson's Chi-squared test). (E) Food supplemented with 4% medium

chain triglyceride (MCT), but not long chain triglyceride (LCT), significantly increased

testis length in *bmm¹* animals but had no effect on this phenotype in *bmm^rev* control

animals (one-way ANOVA with Tukey multiple comparison test). (F) Food

supplemented with 4% medium chain triglyceride significantly increased the number of

spermatid bundles in *bmm¹* testes but had no effect on this phenotype in *bmm^rev* control

animals (one-way ANOVA with Tukey multiple comparison test). (G) Representative

images of *bmm^rev* (G–G') or *bmm¹* (G"–G"') testes stained for Fas3 (G and G") and Vas

(G' and G"'). Scale bars = 25 µm. (H) Quantification of hub area in *bmm^rev* or *bmm¹*

testes showed a significantly larger hub size in *bmm¹* testes (Welch two-sample t-test).

(I) The number of germline stem cells undergoing mitosis (phospho-histone H3$^+$

GSC/total GSC) was not significantly different between *bmm¹* and *bmm^rev* testes

(Kruskal-Wallis rank sum test). (J) The distance between the hub and the first Bam-GFP

positive cyst (Figure 3H) was significantly higher in *bmm¹* testes than in *bmm^rev* testes

(Welch two-sample t-test). (K) All *bmm^rev* testes and most *bmm¹* testes contained

spermatids when raised at 25°C; however, the most advanced stage of

spermatogenesis observed in the majority of *bmm¹* testes isolated from animals reared

at 29°C was the spermatocyte stage. (L) Testes isolated from *bmm¹* animals contained

significantly less Boule-positive area than control testes (Welch two-sample t-test). (M)

Testes isolated from *bmm^1* animals contain fewer individualization complexes than

bmm^{rev} control testes (Kruskal-Wallis rank sum test). (N) Fewer waste bags were

present in testes isolated from *bmm^1* animals compared with *bmm^{rev}* control testes

(Kruskal-Wallis rank sum test).

Supplemental Figure 4 related to Figure 4 – Lipidomic analysis of animals lacking

bmm. (A) Higher fold-changes of triglycerides in *bmm^1* animals were associated with

less saturation in the acyl-groups (Kendall's rank correlation test). (B) Higher fold-

changes of triglycerides in *bmm^1* animals were associated with higher number of

carbons in the acyl-groups (Kendall's rank correlation test). Each dot represents a single

triglyceride species for panel B and C. (C) The volcano plot of identified lipids with

monoglycerides highlighted in blue and diglycerides highlighted in orange. Many

monoglycerides and diglycerides show increase in fold-change in *bmm^1* males. (D) The

number of carbon and the degree of saturation of monoglycerides (MAG) and

diglycerides (DAG) with significant changes in abundance between *bmm^1* and *bmm^{rev}*

males. (E) The volcano plot of identified lipids with fatty acids highlighted in magenta

and acyl-carnitine highlighted in green. Many fatty acids show an increase in fold-

change while many acyl-carnitines show a decrease in fold-change in *bmm^1* males. (F)

The number of carbon and the degree of saturation of fatty acids (FA) and acyl-

carnitines (ACar) with significant changes in abundance between *bmm^1* and *bmm^{rev}*

males. (G) The volcano plot of identified lipids with membrane lipids highlighted in

yellow. (H) The number of carbon and the degree of saturation of membrane lipids with

significant changes in abundance between bmm^1 and bmm^{rev} males. For panel H and I,

PC: phosphatidylcholine; PE: phosphatidylethanolamine; PI: phosphatidylinositol; LPC:

lysophosphatidylcholine; LPE: lysophosphatidylethanolamine; SM: sphingomyelin; PG:

phosphatidylglycerol. (I) Loss of *mdy* function rescued the high number of LD in bmm^1

testes to the level observed in bmm^{rev} testes (one-way ANOVA with Tukey multiple

comparison test). (J) Germline specific loss of *mdy* in bmm^1 animals does not reduce

GSC numbers, but the variance in GSC number is significantly reduced compared to

controls (*nos-GAL4>+*; bmm^1 vs *nos-GAL4>mdy RNAi*; bmm^1: $p=4.5 \times 10^{-5}$; *+>mdy*

RNAi; bmm^1 vs *nos-GAL4>mdy RNAi*; bmm^1: $p=0.0082$ by F-test).

889
890

 KEY RESOURCE TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-Vasa (1:200)	Gift from R. Lehman, MIT	
Anti-Eya (1:50)	Developmental Studies Hybridoma Bank (DSHB)	eya10H6
Anti-zfh1 (1:1000)	Gift from J. Skeath, WUSTL	
Anti-boule (1:1000)	Gift from S. Wasserman, UCSD	
Anti-phospho-histone H3 (1:1000)	Millipore Sigma	05-1354
Experimental models: *Drosophila melanogaster*		
w^1118	Bloomington *Drosophila* stock center	3605
CantonS	Bloomington *Drosophila* stock center	64349
OregonR		
bmm^1	Dr. R. Kühlein	
bmm^rev	Dr. R. Kühlein	
mdy[Qx25], cn[1], bw[1]/CyO, l(2)DTS513[1]	Bloomington *Drosophila* stock center	5095
y[1],w[67c23];P{lacW}Cse1[k03802],mdy[k03902]/CyO	Bloomington *Drosophila* stock center	10536
w[1118];P{GD1749}v6367 (UAS-mdy-RNAi)	Vienna *Drosophila* resource center	6367

REAGENT or RESOURCE	SOURCE	IDENTIFIER
nos-GAL4::VP16	Bloomington *Drosophila* stock center	7303
Tj-GAL4	Gift from D. Godt, University of Toronto	
c587-GAL4	Bloomington *Drosophila* stock center	67747
bam-GFP	[44]	
bmm-GFP	[40]	
GFP-LD	Dr. M. Welte	
P{neoFRT}82B, bmm[1]	This study	
P{neoFRT}82B, bmm[rev]	This study	
bam-GFP, bmm[1]	This study	
bam-GFP, bmm[rev]	This study	
Software and algorithms		
Fiji	https://imagej.net/software/fiji/	
R	https://cran.r-project.org	

892
893

Figure 1 – Lipid droplets are present in early-stage somatic and germline cells.

Figure 2 – *brummer* regulates testis lipid droplets in a cell-autonomous manner.

Figure 3 – A cell-autonomous role of *brummer* in regulating spermatogenesis.

Figure 4 – Loss of *brummer* disrupts triglyceride homeostasis and leads to spermatogenic defects.

Supplemental Figure 1 – Cholesterol is absent from testis lipid droplets.

Supplemental Figure 2 – *brummer* regulates testis lipid droplets in a cell-autonomous manner.

Supplemental Figure 3 – Additional characterization of testis development and spermatogenesis in animals lacking *brummer*.

Supplemental Figure 4 – Lipidomic analysis of animals lacking *brummer*.