TITLE

An important role for triglyceride in regulating spermatogenesis

AUTHORS

Charlotte F. Chao[1&], Yanina-Yasmin Pesch[1&], Huaxu Yu[2], Chenjingyi Wang[2], Maria J.

Aristizabal[3], Tao Huan[2], Guy Tanentzapf[1], Elizabeth J. Rideout[1*]

AFFILIATIONS

[1] Department of Cellular and Physiological Sciences, Life Sciences Institute, The

University of British Columbia, Vancouver, BC, Canada V6T 1Z3

[2] Department of Chemistry, The University of British Columbia, Vancouver, BC, Canada

V6T 1Z1

[3] Department of Biology, Queen's University, Kingston, ON, Canada K7L 3N6

* Corresponding author

[&] Equally contributing authors

[*] Correspondence: elizabeth.rideout@ubc.ca

ABSTRACT

Drosophila is a powerful model to study how lipids affect spermatogenesis. Yet, the contribution of neutral lipids, a major lipid group which resides in organelles called lipid droplets (LD), to sperm development is largely unknown. Emerging evidence suggests LD are present in the testis and that loss of neutral lipid- and LD-associated genes causes subfertility; however, key regulators of testis neutral lipids and LD remain unclear. Here, we show LD are present in early-stage somatic and germline cells within the *Drosophila* testis. We identified a role for triglyceride lipase *brummer* (*bmm*) in regulating testis LD, and found that whole-body loss of *bmm* leads to defects in sperm development. Importantly, these represent cell-autonomous roles for *bmm* in regulating testis LD and spermatogenesis. Because lipidomic analysis of *bmm* mutants revealed excess triglyceride accumulation, and spermatogenic defects in *bmm* mutants were rescued by genetically blocking triglyceride synthesis, our data suggest that *bmm*-mediated regulation of triglyceride influences sperm development. This identifies triglyceride as an important neutral lipid that contributes to *Drosophila* sperm development, and reveals a key role for *bmm* in regulating testis triglyceride levels during spermatogenesis.

35   **KEYWORDS**

36   Triglyceride, lipid droplet, spermatogenesis, testis, *brummer*, adipose triglyceride lipase

37   (ATGL), lipidomics.

INTRODUCTION

Lipids play an essential role in regulating spermatogenesis across animals [1–4].

Studies in *Drosophila* have illuminated key roles for multiple lipid species in regulating

sperm development [5–7]. For example, phosphatidylinositol and its phosphorylated

derivatives participate in diverse aspects of *Drosophila* spermatogenesis including

meiotic cytokinesis [1,8–11], somatic cell differentiation [12], germline and somatic cell

polarity maintenance [13–16], and germline stem cell (GSC) maintenance and

proliferation [17]. Membrane lipids also influence sperm development [18,19], whereas

fatty acids play a role in processes such as meiotic cytokinesis [20] and sperm

individualization [21,22]. While these studies suggest key roles for membrane lipids and

fatty acids during *Drosophila* spermatogenesis, some of which are conserved in

mammals [23–25], much less is known about how neutral lipids contribute to

spermatogenesis.

Neutral lipids are a major lipid group that includes triglyceride and cholesterol

ester, and reside within specialized organelles called lipid droplets (LD) [26]. LD are

found in diverse cell types (*e.g.* adipocytes, muscle, liver, glia, neurons) [27,28,26], and

play key roles in maintaining cellular lipid homeostasis. In nongonadal cell types, correct

regulation of LD contributes to cellular energy production [29–31], sequestration and

redistribution of lipid precursors [32–36], and regulation of lipid toxicity [37–39]. The

importance of LD to normal cellular function in nongonadal cell types is shown by the

fact that dysregulation of LD causes defects in cell differentiation, survival, and energy

production [26,37,40,41]. In the testis, much less is known about the regulation and

function of neutral lipids and LD, and how this regulation affects sperm development.

Multiple lines of evidence suggest a potential role for neutral lipids and LD during

spermatogenesis. First, genes that encode proteins associated with neutral lipid

metabolism and LD are expressed in the testis across multiple species [42–44].

Second, testis LD have been identified in mammals and flies under both normal

physiological conditions [27,44–48] and after mitochondrial stress [49]. Third, loss of

genes associated with neutral lipid metabolism and LD cause subfertility phenotypes in

both flies and mammals [27,50–52]. While studies suggest that mammalian testis LD

contribute to steroidogenesis [53,54], the spatial, temporal, and cell-type specific

requirements for neutral lipids and LD in the testis have not been explored in detail in

any animal. It remains similarly unclear which genes are responsible for regulating

neutral lipids and LD during spermatogenesis.

To address these knowledge gaps, we used *Drosophila* to investigate the

regulation and function of neutral lipids and LD during sperm development. Our detailed

analysis of spermatogenesis under normal physiological conditions revealed the

presence of LD in early-stage somatic and germline cells in the testis. We identified

triglyceride lipase *brummer* (*bmm*) as a regulator of testis LD, and showed that this

represents a cell-autonomous role for *bmm* in the germline. Importantly, we found that

bmm-mediated regulation of testis LD was significant for spermatogenesis, as both

whole-body and cell-autonomous loss of *bmm* caused defects in sperm development.

Given that our lipidomic analysis revealed an excess accumulation of triglyceride in

animals lacking *bmm*, and that genetically blocking triglyceride synthesis rescued many

spermatogenic defects associated with *bmm* loss, our data suggests that *bmm*-

mediated regulation of triglyceride is important for normal *Drosophila* sperm

development. This reveals previously unrecognized roles for neutral lipids such as

triglyceride in regulating spermatogenesis, and for *bmm* in regulating sperm

development under normal physiological conditions. Together, these findings advance

knowledge of the regulation and function of neutral lipids during spermatogenesis.

RESULTS

Lipid droplets are present in early-stage somatic and germline cells

We previously reported the presence of small LD (<1 μm) at the apical tip of the testis

[27], a finding we reproduced in w^{1118} males using neutral lipid stain BODIPY (Figure

1A). These LD were present in the region that contains stem cells, early-stage somatic

cells, and germline cells (Figure 1A-A', arrows). LD were also present in the hub, an

organizing center and stem cell niche in the *Drosophila* testis (Figure 1A''-A''', arrows)

[55], but largely absent within the area occupied by spermatocytes (Figure 1A and A',

arrowheads). This LD distribution was reproduced in two independent genetic

backgrounds and at two additional ages (Figure 1B, 1C). While LD may contain multiple

neutral lipid species [56], cholesterol-binding fluorescent polyene antibiotic filipin III did

not detect cholesterol within testis LD (Figure S1A), suggesting triglyceride is the main

neutral lipid in *Drosophila* testis LD.

Drosophila spermatogenesis requires the co-development and differentiation of

the germline and the somatic lineages [57]. To identify LD in each lineage, we used the

GAL4/UAS system to overexpress a GFP transgene fused to the LD-targeting motif of

motor protein *Klarsicht* (*UAS-GFP-LD*) [58]. Somatic overexpression of *UAS-GFP-LD*

using *Traffic jam (Tj)-GAL4* revealed that the majority of somatic LD in 0-day-old males

were located <30 μm from the hub (Figure 1D, 1E). Because the somatic LD distribution

coincided with a marker for somatic stem cells and their immediate daughter cells (Zinc

finger homeodomain 1, Zfh-1) (Figure 1F; two-sample Kolmogorov-Smirnov test) [59],

but not with a marker for late somatic cells (Eyes absent, Eya) [12,60], our data suggest

LD are present in early somatic cells. Germline overexpression of *UAS-GFP-LD* using

nanos (nos)-GAL4 demonstrated the presence of LD within germline cells near the

apical tip of the testis in 0-day-old males (Figure 1G, 1H). Specifically, the

disappearance of germline LD coincided with peak expression of a GFP reporter that

reflects the expression of Bag-of-marbles (Bam) protein in the testis (Bam-GFP) [61]

(Figure 1I, 1J). Because peak Bam expression signals the last round of transient

amplifying mitotic cell cycle prior to the germline's transition into the meiotic cell cycle

[62–64], our data suggests that germline LD, like somatic LD, are present in cells at

early stages of development.

brummer plays a cell-autonomous role in regulating testis lipid droplets

Adipose triglyceride lipase (*ATGL*) is a critical regulator of neutral lipid metabolism and

LD [65–74]. Loss of *ATGL* in many cell types triggers LD accumulation, and *ATGL*

overexpression decreases LD number [30,67,68,71,73,75,76]. Given that the

Drosophila ATGL homolog *brummer* (*bmm*) regulates testis LD induced by

mitochondrial stress [49], we explored whether *bmm* regulates testis LD under normal

physiological conditions. We first examined *bmm* expression in the testis by isolating

this organ from flies carrying a *bmm* promoter-driven *GFP* transgene (*bmm-GFP*) that

recapitulates many aspects of *bmm* mRNA regulation [77]. GFP expression was present

in the germline of *bmm-GFP* testes, and we found germline GFP levels were higher in

spermatocytes than at earlier stages of sperm development (Figure 2A, 2B; one-way

ANOVA with Tukey multiple comparison test). In further support of this observation, we

analyzed a publicly available single-cell RNA sequencing dataset from the male

reproductive organ [78]. Using pseudotime analysis, we arranged the germline (Figure

S2A) and somatic cells (Figure S2B) based on their annotated developmental trajectory.

The expression pattern of *bmm* in the germline matched our observation with the *bmm-*

GFP reporter (Figure S2C). While levels of the *bmm-GFP* reporter were low in somatic

cells, single-cell RNA sequencing data identified *bmm* expression in the somatic

lineage, with higher expression in cells at later stages of development (Figure S2D).

Additional neutral lipid- and lipid droplet-associated genes such as *lipid storage droplet-*

2, *Seipin*, *Lipin*, and *midway* also showed differential regulation during differentiation in

the testis (Figure S2C, S2D). Combined with our data on the location of testis LD, these

gene expression data suggest that *bmm* upregulation in both somatic and germline cells

during differentiation corresponds to the downregulation of testis LD. Supporting this,

germline GFP levels were negatively correlated with testis LD in *bmm-GFP* flies (Figure

2A, 2C), suggesting regions with higher *bmm* expression had fewer LD.

To test whether *bmm* regulates testis LD, we compared LD in testes from 0-day-

old males carrying a loss-of-function mutation in *bmm* (*bmm¹*) to control male testes

(*bmm^rev)* [67]. *bmm¹* males had significantly more LD across all LD sizes compared with

control males at the apical tip of the testis (Figure 2D–2G; Welch two-sample t-test with

Bonferroni correction) and showed a significantly expanded LD distribution (Figure 2D–

2F, 2H; two-sample Kolmogorov-Smirnov test). This suggests *bmm* normally restricts

LD to the apical tip of the testis, an observation we confirmed in both somatic and

germline lineages using lineage-specific expression of GFP-LD (Figure S3A–S3D).

Importantly, after inducing homozygous bmm^{rev} or bmm^1 clones in the testes using the

FLP-FRT system (Figure 3A, 3B) [79], we found bmm^1 spermatocyte clones had

significantly more LD at 3 days post-clone induction (Figure 3C; Welch two-sample t-

test), a stage at which LD were absent from bmm^{rev} clones. Because we observed no

significant effect of cell-autonomous bmm loss on LD at any other stage of germline

development (Figure 3C), this suggests bmm does not regulate LD at early stages of

germ cell development. Instead, our data suggest bmm plays a role in regulating LD at

the spermatogonia-spermatocyte transition. While we were unable to assess LD in

bmm^1 somatic clones, our data reveals a previously unrecognized cell-autonomous role

for bmm as a key regulator of LD in germline cells.

brummer plays a cell-autonomous role in regulating germline development

To determine the physiological significance of bmm-mediated regulation of testis LD, we

investigated testis and sperm development in males without bmm function. In 0-day-old

bmm^1 males reared at 25°C, testis size was significantly smaller than in age-matched

bmm^{rev} controls (Figure 4A, 4B; Welch two-sample t-test), and the number of spermatid

bundles was significantly lower (Figure 4C; Kruskal-Wallis rank sum test). When the

animals were reared at 29°C, a temperature that exacerbates spermatogenesis defects

associated with changes in lipid metabolism [21], bmm^1 phenotypes were more

pronounced (Figure S3A, S3B; Welch two-sample t-test, Kruskal-Wallis rank sum test).

Defects in testis size were also observed at 14 days post-eclosion; suggesting testis

size defects persist later into the life course (Figure S4C; Welch two-sample t-test). In

contrast, the number of spermatid bundles per testis was not significantly different

between bmm^1 and bmm^{rev} males at this age (Figure S4D; Welch two-sample t-test),

potentially due to a large decrease in the number of spermatid bundles in 14-day-old

bmm^{rev} males (Figure 4C, S4D). Together, these data suggest loss of bmm affects testis

development and spermatogenesis. Similar phenotypes are observed in male mice

without ATGL [52], and supplementing the diet of bmm^1 males with medium-chain

triglycerides (MCT) partially rescues the testis and spermatogenic defects we observed

in flies (Figure S4E, S4F; one-way ANOVA with Tukey multiple comparison test), as it

does in mice [52,80]. This identifies similarities between flies and mice in fertility-related

phenotypes associated with whole-body loss of bmm/ATGL.

188   To explore spermatogenesis in bmm^1 animals, we used an antibody against germ

cell-specific marker Vasa to visualize the germline in the testes of bmm^1 and bmm^{rev}

males (Figure 4D, 4E) [81]. We observed a significant increase in the number of

germline stem cells (GSC) (Figure 4F; Kruskal-Wallis rank sum test) and higher

variability in GSC number in bmm^1 males (p=5.7×10^{-12} by F-test). Given that GSC

number is affected by hub size and GSC proliferation [82,83], we monitored both

parameters in bmm^1 and bmm^{rev} controls. While hub size in bmm^1 testes was

significantly larger than in testes from bmm^{rev} controls (Figure S4G, S4H; Welch two-

sample t-test), the number of phosphohistone H3-positive GSCs, which indicates

proliferating GSCs, was unchanged in bmm^1 animals (Figure S4I; Kruskal-Wallis rank

sum test). While this indicates a larger hub may partly explain bmm's effect on GSC

number, bmm also plays a cell-autonomous role in regulating GSCs, as we recovered a

higher proportion of bmm^1 clones in the GSC pool compared with bmm^{rev} clones at 14

days after clone induction (Figure 4G; Welch two-sample t-test). Given that we detected

no effect of cell-autonomous bmm loss on the number of GSC LD (Figure 3C), more

work will be needed to understand how bmm regulates GSCs at a stage prior to its

effects on LD number. Future studies will also need to confirm whether bmm^1 mutant

GSCs show an increased ability to occupy space at the hub.

Beyond GSCs, we uncovered additional spermatogenesis defects in bmm^1

testes. Peak Bam-GFP expression in germline cells of the testes from 0-day-old bmm^1

and bmm^{rev} males showed GFP-positive cysts were significantly further away from the

hub in bmm^1 testes (Figure 4H, S4J; Welch two-sample t-test). Indeed, 15/18 bmm^1

testes contained Vasa-positive cysts with large nuclei in the distal half of the testis

(Figure 4I, arrows), a phenotype not present in bmm^{rev} testes (0/8) (p=0.0005 by

Pearson's Chi-square test). Because these phenotypes are also seen in testes with

differentiation defects [13,84], we recorded the stage of sperm development reached by

the germline in bmm^1 testes. Most bmm^1 testes contained post-meiotic cells in males

raised at 25°C (Figure S4K); however, germline development did not progress past the

spermatocyte stage in most bmm^1 testes from animals raised at 29°C (Figure S4K).

Testes from bmm^1 males reared at 25°C also had a smaller Boule-positive area (Figure

4J, S4L; Welch two-sample t-test), and fewer individualization complexes and waste

bags (Figure S4M, S4N; Kruskal-Wallis rank sum test). Because Boule-positive area,

individualization complexes, and waste bags indicate germline cells at later stages in

development, these data suggest the loss of bmm causes a reduction in differentiated

cell types. Because we observed significantly fewer bmm^1 spermatocyte and spermatid

clones at 14 days after clone induction (Figure 4K, 4L; p = 0.0496, Kruskal-Wallis rank

sum test), these effects on germline development may represent a cell-autonomous role

for *bmm* in regulating spermatogenesis in this cell type. Given that the statistical

significance of this finding was not as strong as for our other data, future studies should

repeat this experiment with more samples. We also reveal a potential non-cell-

autonomous role for somatic *bmm*. While there was no difference in the ratio of Zfh-1-

positive cells between homozygous clones and heterozygous clones in animals carrying

the *bmm¹* or *bmmʳᵉᵛ* alleles at 14 days post clone induction (Figure S4O; Kruskal-Wallis

rank sum test), the distance from the hub to the Zfh-1 positive clones reside was

significantly decreased in *bmm¹* homozygous clones (Figure S4P; Kruskal-Wallis rank

sum test). Together, these data indicate *bmm* plays a cell-autonomous role in germline

cells, and potentially a non-cell-autonomous role in somatic cells, to regulate

spermatogenesis.

***brummer*-dependent regulation of testis triglyceride levels affects**

spermatogenesis

ATGL catalyzes the first and rate-limiting step of triglyceride hydrolysis [73,85,86]. Loss

of this enzyme or its homologs leads to excess triglyceride accumulation

[27,30,67,73,75] and shifts in multiple lipid classes [66,87–89]. To determine how loss of

bmm affects spermatogenesis, we carried out whole-body mass spectrometry (MS)-

based untargeted lipidomic profiling of *bmm¹* and *bmmʳᵉᵛ* males. Hierarchical clustering

of lipid species suggests that *bmm¹* and *bmmʳᵉᵛ* males show distinct lipidomic profiles

(Figure 5A). Overall, we detected 2464 and 1144 lipid features with high quantitative

confidence in positive and negative ion modes, respectively. By matching experimental

m/z, isotopic ratio, and tandem MS spectra to lipid libraries, we confirmed 293 unique

lipid species (Supplemental table 1). We found 107 lipids had a significant change in

abundance between *bmm¹* and *bmm^rev* males ($p_{adj}<0.05$): 85 species were upregulated

in *bmm¹* males and 22 lipid species were downregulated. Among differentially regulated

species from different lipid classes, triglyceride had the largest residual above expected

proportion (*p*=$5.00×10^{-4}$ by Pearson's Chi-squared test). This suggests triglyceride is the

lipid class most affected by loss of *bmm* (Figure 5B, 5C).

In *bmm¹* males, the majority of triglyceride species (55/97) were significantly

higher in abundance compared with *bmm^rev* control males. Because we observed a

positive correlation between the fold increase in triglyceride abundance with both the

number of double bonds (*p*=$7.52×10^{-8}$ by Kendall's rank correlation test; Figure S5A)

and the number of carbons (*p*=$2.77×10^{-10}$ by Kendall's rank correlation test; Figure

S5B), our data align well with *bmm/ATGL's* known role in regulating triglyceride levels

[67,68,73] and its substrate preference of long-chain polyunsaturated fatty acids [85].

While we also detected changes in species such as fatty acids, acylcarnitine, and

membrane lipids (Figure S5C–S5H), in line with recent *Drosophila* lipidomic data

[90,91], the striking accumulation of triglyceride in *bmm¹* males suggested that excess

testis triglyceride in *bmm¹* males may contribute to their spermatogenic defects. To test

this, we examined spermatogenesis in *bmm¹* males carrying loss-of-function mutations

in *midway* (*mdy*). *mdy* is the *Drosophila* homolog of *diacylglycerol O-acyltransferase 1*

(*DGAT1*), and whole-body loss of *mdy* reduces whole-body triglyceride levels [92–94].

Importantly, testes isolated from males with global loss of both *bmm* and *mdy*

(*mdy*$^{QX25/k03902}$*;bmm*^{1}) had fewer LD than testes dissected from *bmm*^{1} males (Figures

5D, S5I; one-way ANOVA with Tukey multiple comparison test).

We found that testes isolated from *mdy*$^{QX25/k03902}$*;bmm*^{1} males were significantly

larger and had more spermatid bundles than testes from *bmm*^{1} males (Figure 5E–G;

one-way ANOVA with Tukey multiple comparison test). The elevated number of GSCs

in *bmm*^{1} male testes was similarly rescued in *mdy*$^{QX25/k03902}$*;bmm*^{1} males (Figure 5H;

one-way ANOVA with Tukey multiple comparison test). These data suggest that

defective spermatogenesis in *bmm*^{1} males can be partly attributed to excess triglyceride

accumulation. Notably, at least some of the effects of global *mdy* loss on *bmm*^{1} males

can be attributed to the germline: RNAi-mediated knockdown of *mdy* in the germline of

bmm^1 males partially rescued the defects in testis size (Figure 5I; Kruskal-Wallis rank

sum test with Dunn's multiple comparison test) and GSC variance (Figure S5J; *p*=4.5 x

10^{-5} and 8.2 x 10^{-3} by F-test from the GAL4- and UAS-only crosses, respectively). While

future studies will need to test whether germline-specific loss of *mdy* also rescues

spermatid number defects in *bmm*^{1} males, our data suggest *bmm*-mediated regulation

of testis triglyceride plays a previously unrecognized role in regulating sperm

development.

DISCUSSION

In this study, we used *Drosophila* to gain insight into how the neutral lipids, a major lipid

class, contribute to sperm development. We describe the distribution of LD under

normal physiological conditions in the *Drosophila* testis, and show that LD are present

at the early stages of development in both somatic and germline cells. While many

factors are known to regulate LD in nongonadal cell types, we reveal a cell-autonomous

role for triglyceride lipase *bmm* in regulating testis LD during spermatogenesis. In

particular, we identified a requirement for *bmm* in mediating the decrease in LD at the

spermatogonia-spermatocyte transition. This regulation is important for sperm

development, as our data indicates that loss of *bmm* causes a decrease in the number

of differentiated cell types in the testis. This phenotype may be attributed to a delay in

differentiation, a block in differentiation, or to a loss of differentiated cells through cell

death. Future studies will therefore be essential to resolve why *bmm* loss causes a

reduction in differentiated cell types. Nevertheless, these defects in the number of

differentiated cell types can be partially explained by the excess accumulation of

triglyceride in flies lacking *bmm*, as global and cell type-specific inhibition of triglyceride

synthesis rescues multiple spermatogenic defects in *bmm* mutants. Together, our data

reveals previously unrecognized roles for LD and triglycerides during spermatogenesis,

and for *bmm* as an important regulator of testis LD and germline development under

normal physiological conditions.

One key outcome of our study was increased knowledge of LD regulation and

function in the testis. Despite rapidly expanding knowledge of LD in cell types such as

adipocytes or skeletal muscle, less is known about how LD influence spermatogenesis

under normal physiological conditions. In mammals, testis LD contain cholesterol and

play a role in promoting steroidogenesis [95,96]. In flies, we show that LD are present in

the testis, and that excess accumulation of these LD affects sperm development. In

nongonadal cell types, triglycerides provide a rich source of fatty acids for cellular ATP

production, lipid building blocks to support membrane homeostasis and growth, and

metabolites that can act as signaling molecules [26]. Because ATP production, lipid

precursors, and lipid signaling all play roles in supporting normal sperm development

[97,98], future studies will need to determine how each of these processes is affected

when excess triglyceride accumulates in testis LD (Figure 6). It will also be important to

determine whether it is the loss of metabolites produced by *bmm*'s enzymatic action, or

an increase in triglycerides, that leads to the reduction in differentiated cell types during

spermatogenesis. Together, these experiments will provide critical insight into how

triglyceride stored within testis LD contributes to overall cellular lipid metabolism during

spermatogenesis. Because of the parallel spermatogenic defects we observed in *bmm*

mutants and *ATGL*-deficient mice, we expect these mechanisms will also operate in

other species.

A more comprehensive understanding of neutral lipid metabolism during sperm

development will also emerge from studies on the upstream signaling networks that

regulate testis LD and triglyceride. Given that we show an important and cell-

autonomous role for *bmm* in regulating testis LD and triglyceride, future studies will

need to identify factors that regulate *bmm* in the testis. Based on public single-cell

RNAseq data and the *bmm-GFP* reporter strain, our data suggest *bmm* mRNA levels

are differentially regulated between early and later stages of sperm development.

Candidates for mediating this regulation include the insulin/insulin-like growth factor

signaling pathway (IIS), Target of rapamycin (TOR) pathway, and nuclear factor

κB/Relish pathway (NFκB), as all of these pathways influence *bmm* mRNA levels in

nongonadal cell types [99–105]. Beyond mRNA levels, Bmm protein levels and post-

translational modifications may also be differentially regulating during spermatogenesis.

For example, studies show that the proteins encoded by *bmm* homologs in other

animals are regulated by phosphorylation [106], mediated by kinases such as

adenosine monophosphate-activated protein kinase (AMPK) and protein kinase A

(PKA) [107–109]. Importantly, many of these pathways, including IIS, TOR, AMPK,

NFκB and possibly PKA influence *Drosophila* sperm development [110–115]. Identifying

the signaling networks that influence *bmm* regulation during sperm development will

therefore lead to a deeper understanding of how testis LD and triglyceride are

coordinated with physiological factors to promote normal spermatogenesis. Because

pathways such as IIS and AMPK, and others, regulate sperm development in other

species [116–118], these insights may reveal conserved mechanisms that govern the

regulation of cellular neutral lipid metabolism during sperm development.

ACKNOWLEDGEMENTS

We thank Dr. Ronald Kühnlein for *bmm^1* and *bmm^{rev}* lines [67], Dr. Michael Welte for *UAS-GFP-LD* [58], and Dr. Kaeko Kamei for *bmm-GFP* [77]. We used stocks from the Bloomington *Drosophila* Stock Center (NIH P40OD018537) and Vienna *Drosophila* Resource Center (VDRC). We acknowledge critical resources and information provided by FlyBase [119] (supported by the National Human Genome Research Institute at the U.S. National Institutes of Health (U41 HG000739) and the British Medical Research Council (MR/N030117/1)). This work was supported by the Life Sciences Institutes Imaging Core, supported by the UBC GREx Biological Resilience Initiative. Funding for this study was provided by grants to EJR from the Canadian Institutes for Health Research (PJT-153072), Michael Smith Foundation for Health Research (16876), and the Canadian Foundation for Innovation (JELF-34879). GT was supported by a grant from the Natural Sciences and Engineering Research Council (NSERC; 2018-04648), TH/HY/CW were supported by NSERC (2020-04895), MA was supported by the Jacob's foundation. We would like to acknowledge that our research takes place on the traditional, ancestral, and unceded territory of the Musqueam people; a privilege for which we are grateful.

AUTHOR CONTRIBUTIONS

Conceptualization, C.C. and E.J.R.; Methodology, C.C. and Y.Y.P.; Software, C.C.; Investigation, C.C., H.Y., and Y.Y.P.; Lipidomics, M.A., H.Y., C.W., T.H.; Writing – Original Draft, C.C. and E.J.R.; Writing – Review and Editing, C.C., E.J.R., and Y.Y.P.;

Supervision, E.J.R., G.T., and T.H.; Project administration, E.J.R.; Funding Acquisition,

E.J.R., G.T., and T.H.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1 – Lipid droplets are present in early-stage somatic and germ cells. (A)

Testis lipid droplets (LD) in w^{1118} animals visualized with neutral lipid dye BODIPY.

(A,A') Scale bar=50 µm; (A'',A''') scale bar=15 µm. Asterisk indicates hub in all images.

Arrows point to LD; arrowheads point to spermatocytes in A,B. Spermatocytes were

identified as described in methods section. (B) Testis LD visualized with BODIPY in

newly-eclosed males from two wild-type genotypes. Scale bars: main image=50 µm;

inset image=10 µm. (C) Testis LD from w^{1118} animals at different times post-eclosion.

Scale bars=50 µm. (D) Testis LD visualized with LipidTox Red in animals with somatic

cell overexpression of GFP-LD (*Tj-GAL4>UAS-GFP-LD*). GFP- and LipidTox Red-

positive punctae are somatic LD (D–D'' arrows); LipidTox punctae without GFP indicate

germline LD (D–D'' arrowheads). Scale bars=10 µm. (E) Histogram showing the spatial

distribution of somatic cell LD; error bars represent standard error of the mean (SEM).

(F) Cumulative frequency distributions of somatic LD (blue line, data reproduced from

E), zfh-1-positive somatic cells (zfh-1[+] cells, orange line), and Eya-positive somatic cells

(Eya[+] cells, grey line). (G) Testis LD visualized with LipidTox Red in males with germline

overexpression of GFP-LD (*nos-GAL4>UAS-GFP-LD*). GFP- and LipidTox Red-positive

punctae indicate germline LD (arrows); LipidTox punctae without GFP indicate non-

germline LD (arrowheads). Scale bars=10 µm. (H) Histogram representing the spatial

distribution of LD within the germline; error bars represent SEM. (I) Histogram

representing the spatial distribution of LD and GFP fluorescence (green line) (arbitrary

units, a.u.) in a representative testis of a *bam-GFP* animal (panel J). (J) Testis LD in a

bam-GFP animal; arrows point to LD and arrowheads point to spermatocytes. Scale

bar=50 µm. See also Supplemental Figure 1.

Figure 2 – *brummer* regulates lipid droplets in both germline and somatic cells of

the testis. (A-A'''') Testis lipid droplets (LD) indicated by LipidTox Red in *bmm-GFP*

animals. Arrows point to LD in all images. Arrowheads point to spermatocytes. Scale

bars=50 µm. Asterisks indicate the hub in all images. (B) Quantification of nuclear GFP

intensity in testes isolated from *bmm-GFP* animals (n=3). Germline stem cell (GSC),

spermatogonia (SG), spermatocyte (SC). (C) Spatial distribution of LD (grey histogram)

and GFP expression (green line) in testes from *bmm-GFP* animals as a function of

distance from the hub (n=3). (D,E) LD near the apical region of the testis in bmm^{rev} (D)

or bmm^1 (E) animals. (F) LD further away from the apical tip in bmm^1 animals. (D–F)

Scale bars=50 µm. (G) Histogram representing testis LD size distribution in bmm^{rev}

(grey) and bmm^1 (orange). (H) Apical tip of the testes is at the left of the graph;

individual dots represent a single LD and its relative position to the hub marked by an

asterisk. Cumulative frequency distribution of the distance between LDs and the apical

tip of the testes are drawn as solid lines. See also Supplemental Figure 2 and 3.

Figure 3 – *bmm* regulates germline lipid droplets in a cell-autonomous manner.

(A–B) Single confocal slices through a representative testis isolated from an individual

carrying clones induced using the FLP-FRT system at 3 days post-clone induction.

Clones are homozygous for an allele that encodes a functional *bmm* protein product

(*bmm^rev*; A–A''') or a loss-of-function *bmm* allele (*bmm^1*, B–B'''). GFP negative areas

mark homozygous clones in panel A and B; the boxed areas in A, A'' and B, B'' are

shown in A', A''' and B', B''', respectively. In homozygous *bmm^rev* spermatocyte clones

we detected no LD using neutral lipid dye BODIPY (A'' and A'''). In contrast,

spermatocyte clones homozygous for *bmm^1* have detectable LD (B'' and B''', arrows).

Scale bars=50 µm in A,A'' and B,B''; scale bars=10 µm in A',A''' and B',B'''. (C) Number

of testis LD in *bmm^rev* (grey) or *bmm^1* (orange) in FLP-FRT clones 3 days post-clone

induction; dots represent measurements from a single clone. The number of cells in

each cyst (CC) counted is indicated. There were significantly more LD in *bmm^1*

spermatocyte (SC) clones (*p* = 0.026; Welch two-sample t-test) but not at other stages

of development.

Figure 4 – A cell-autonomous role for *bmm* in regulating spermatogenesis. Testes

isolated from *bmm^rev* (A) and *bmm^1* (A') animals raised at 25°C stained with phalloidin.

Scale bars=100 µm. (B) Testis size in *bmm^1* and *bmm^rev* animals raised at 25°C. (C)

Spermatid bundle number in *bmm^1* and *bmm^rev* testes from animals reared at 25°C.

(D,E) Representative images of *bmm^rev* (D) or *bmm^1* (E) testes stained with DAPI and

anti-Vasa antibody. Arrows indicate germline stem cells (GSC). Scale bar=50 µm. The

hub is marked by an asterisk in all images. (F) GSC number in *bmm^1* and *bmm^rev*

testes. (G) Proportion of GSCs that were either bmm^1 or bmm^{rev} clones at 3 and 14

days post-clone induction. (H) Representative images of bmm^{rev} (H) and bmm^1 (H')

testes carrying *bam-GFP*; data quantified in Figure S4J. Arrows indicate regions with

high Bam-GFP. Scale bars=50 μm. (I) Representative images of bmm^{rev} (I) or bmm^1

(I',I") testes stained with anti-Vasa antibody. Arrows indicate Vasa-positive cysts in

bmm^1 testis. Panel I" is magnified from the boxed region in I'. (I,I') Scale bars=100 μm;

(I") scale bar=50 μm. (J) Maximum projection of bmm^{rev} (J) or bmm^1 (J') testes stained

with anti-Boule antibody (green) and DAPI (blue). Scale bars=100 μm. Number of bmm^1

and bmm^{rev} spermatocyte clones (K) or post-meiotic clones (L) at 3- and 14-days post-

clone induction. See also Supplemental Figure 4.

Figure 5 – Loss of *bmm* disrupts triglyceride homeostasis and leads to

spermatogenic defects. (A) Hierarchical clustering of lipid species detected in bmm^{rev}

and bmm^1 animals. (B) Histograms showing the proportion of species in each lipid class

with different levels between bmm^1 and bmm^{rev}. Numbers on histograms indicate the

number of species with differences in abundance. (C) Volcano plot showing fold change

in abundance of triglycerides (green; 97 species) and non-triglyceride lipids (grey; 186

species) in our dataset. (D) Arrows indicate testis LD stained with LipidTox Red in

bmm^{rev} (D), bmm^1 (D'), or $mdy^{QX25/k03902}$; bmm^1 (D") animals. (E) Whole testes isolated

from bmm^{rev} (E), bmm^1 (E'), or $mdy^{QX25/k03902}$;bmm^1 (E") animals stained with anti-Vasa

antibody (red) and DAPI (blue). Arrowheads indicate spermatid bundles. Scale

bars=100 μm. (F) Testis size in bmm^{rev}, bmm^1, and $mdy^{QX25/k03902}$;bmm^1 animals.

Spermatid bundles (G) and number of germline stem cells (H) in bmm^{rev}, bmm^1, and

$mdy^{QX25/k03902}$;bmm^1 animals. (I) Testis size in animals with germline-specific mdy

knockdown (*nos-GAL4>mdy RNAi*; bmm^1) compared with controls (*nos-GAL4>+*; bmm^1

and *+>mdy RNAi*; bmm^1). See also Supplemental Figure 5.

Figure 6 – Model of *bmm*-mediated lipid droplet regulation in the *Drosophila*

testis. Schematic representation summarizing *bmm*-mediated lipid droplet regulation in

the testis during development.

MATERIALS AND METHODS

Materials and Resource availability. *Drosophila* strains and their source are listed in a Key Resources table. Further information and requests for resources and reagents should be directed to, and will be fulfilled by, lead contact Dr. Elizabeth J. Rideout (elizabeth.rideout@ubc.ca).

Data and Code availability. All raw data and results of statistical tests reported in this paper are located in Supplementary files 1-4. This paper does not report original code. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Fly husbandry. Fly stocks were maintained at room temperature in 12:12 hour light:dark cycle. Unless otherwise indicated, all flies were raised at 25°C with a density of 50 larvae per 10 mL fly media. Because this project examines sperm development, we used male flies in all experiments. Fly media contained 20.5 g sucrose (SU10, Snow Cap), 70.9 g Dextrose (SUG8, Snow Cap), 48.5 g cornmeal (AO18006, Snow Cap), 30.3 g baker's yeast (NB10, Snow Cap), 4.55 g agar (DR-820-25F, SciMart), 0.5 g calcium chloride dihydrate (CCL302.1, BioShop Canada), 0.5 g magnesium sulfate heptahydrate (MAG511.1, BioShop Canada), 4.9 mL propionic acids (P1386, Sigma-Aldrich), and 488 µL phosphoric acid (P5811, Sigma-Aldrich) per 1L of media. For diets with medium- or long-chain triglyceride, 4 g of coconut oil (medium chain triglyceride) or olive oil (long chain triglyceride) was added per 100 mL of media described above prior

to cooling. Males were collected and dissected within 24 hours of eclosion unless

otherwise indicated. Fixations were performed at room temperature with 4%

paraformaldehyde (CA11021-168, VWR) in PBS for 20 minutes on a rotating platform

followed by washing in PBS twice before staining. Fly strains used in our study are

listed in a Key Resources table.

Testis cell stage classification and measurements. Cells at an early stage of

development (stem cells and early-stage somatic and germline cells) were located in

the apical region of the testis, and were identified by their small and dense nuclei[120].

GSCs were defined as Vasa-positive cells in direct contact with the hub; proliferating

GSCs were identified as Vasa-positive cells in direct contact with the hub that were also

phospho-H3 positive. Cells in the testis region occupied by primary spermatocytes were

identified by their large cell size and decondensed chromosome staining occupying

three nuclear domains [120]. Spermatid bundles were identified by their condensed and

needle-shaped nuclei, which roughly corresponds to nuclei with protamine-based

chromatin [121]. Hub size was measured by quantifying FasIII-positive area of the

testis. Testis size was measured by quantifying the length of a line drawn down the

middle of a testis image; starting from the apical tip of the testis and ending where the

testis meets the seminal vesicle.

FLP-FRT clone induction. Adult males were collected at 3-5 days post-eclosion and

heat-shocked three times at 37°C for 30 min followed by a 10 min rest period at room

temperature between heat shocks. After heat-shock, the flies were incubated at room

temperature until dissection.

Immunohistochemistry. Fixed samples were rinsed three times with blocking solution

containing 0.2% bovine serum albumin (A4503, Sigma-Aldrich), 0.3% Triton-X in PBS,

then blocked for 1 hr on a rotating platform at room temperature. During the incubation,

the blocking solution was changed every 15 minutes. After blocking, the sample were

resuspended in blocking solution with the appropriate concentration of primary antibody

(see Key Resources table), and incubated overnight at 4°C. Samples were rinsed three

times with blocking solution after removing primary antibody, and blocked for one hour

on a rotating platform in blocking solution. Secondary antibody was applied in blocking

solution and left on the rotating platform at room temperature for 40 min. The sample

was rinsed with blocking solution three more times, and washed four times for 15 min

per wash in blocking solution. Testis samples were resuspended in Vectashield

mounting media with DAPI (H-1200-10, Vector Laboratory) or SlowFade Diamond

mounting media (S36972, Thermo Fisher Scientific) prior to mounting.

Lipid droplet staining. Fixed testes were briefly permeabilized with 0.1% Triton-X in

PBS for 5 min prior to applying phalloidin. For BODIPY (4,4-Difluoro-1,3,5,7,8-

Pentamethyl-4-Bora-3a,4a-Diaza-*s*-Indacene) staining, samples were suspended in

PBS containing 10 µg/mL DAPI (2879083-5mg, PeproTech), 1:500 BODIPY 495/503

(Thermo Fisher Scientific D3922), and 1:1000 phalloidin iFluor647 (ab176759, Abcam)

or 1:40 phalloidin TexasRed (T7471, Thermo Fisher Scientific). For staining with

LipidTox Red, samples were suspended in PBS containing 10 μg/mL DAPI (2879083-

5mg, PeproTech), 1:200 LipidTox Red (H34476, Thermo Fisher Scientific), and 1:1000

phalloidin iFluor647 (ab176759, Abcam). For staining free sterols, samples were

prepared as for BODIPY staining with 50 μg/mL filipin in place of BODIPY for 30 min.

Samples were incubated on a rotating platform for 40 minutes at room temperature.

After incubation, samples were washed twice with PBS, then resuspended in SlowFade

Diamond mounting media (Thermo Fisher Scientific S36972) prior to mounting.

Image acquisition and processing. All images were acquired on a Leica SP5 confocal

microscope system with 20X or 40X objectives and quantified with Fiji image analysis

software[122].

Drosophila lipidomics. *Drosophila* extracts were prepared following the previously

reported protocol[123]. Briefly, 10 *Drosophila* males (~10 mg) were weighed, 300 μL of

ice-cold methanol/water mixture (9:1, v:v) was added to these males, and the samples

were homogenized with glass beads using a bead beater (mini-beadbeater-16,

BioSpec, Bartlesville, Ok, USA). Sample weight was used for sample normalization. Fly

lysate was kept at -20°C for 4 hours for protein precipitation. Then, 900 μL of methyl

tert-butyl ether was added and the solution was shaken for 5 min to extract lipids. To

induce phase separation 285 μL of water was added, followed by centrifugation. The

upper layer was separated, dried, and reconstituted in isopropanol/acetonitrile (1:1, v:v)

for liquid chromatography-mass spectrometry (LC-MS) analysis. The volume of

reconstitution solution was proportional to sample weight for normalization. Quality

control (QC) samples were prepared by pooling 20 μL aliquot from each sample. The

method blank sample was prepared using an identical workflow but without adding

Drosophila.

Drosophila extracts were analyzed on an UHR-QqTOF (Ultra-High Resolution

Qq-Time-Of-Flight) mass spectrometry Impact II (Bruker Daltonics, Bremen, Germany)

interfaced with an Agilent 1290 Infinity II LC Systems (Agilent Technologies, Santa

Clara, CA, USA). LC separation was performed using a Waters reversed-phase (RP)

UPLC Acquity BEH C18 Column (1.7 μm, 1.0 mm ×100 mm, 130 Å) (Milford, MA, USA)

maintained at 30°C. For positive ion mode, the mobile phase A was 60% acetonitrile in

water and the mobile phase B was 90% isopropanol in acetonitrile, both containing 5

mM ammonium formate (pH = 4.8, adjusted by formic acid). For negative ion mode, the

mobile phase A was 60% acetonitrile in water and the mobile phase B was 90%

isopropanol in acetonitrile, both containing 5 mM NH$_4$FA (pH = 9.8, adjusted by

ammonium hydroxide). The LC gradient for positive and negative ion modes was set as

follows: 0 min, 5% B; 8 min, 40% B; 14 min, 70% B; 20 min, 95% B; 23 min, 95% B; 24

min, 5% B; 33 min, 5% B. The flow rate was 0.1 mL/min. The injection volume was

optimized to 2 μL in positive mode and 5 μL in negative mode using QC sample. The

ESI source conditions were set as follows: dry gas temperature, 220 °C; dry gas flow, 7

L/min; nebulizer gas pressure, 1.6 bar; capillary voltage, 4500 V for positive mode and

3000 V for negative mode. The MS1 analysis was conducted using following

parameters: mass range, 70-1000 m/z; spectrum type: centroid, calculated using

maximum intensity; absolute intensity threshold: 250. Data-dependent MS/MS analysis

parameters: collision energy: 16-30 eV; cycle time, 3 s; spectra rate: 4 Hz when

intensity $< 10^4$ and 12 Hz when intensity $> 10^5$, linearly increased from 10^4 to 10^5.

External calibration was applied using sodium formate to ensure the m/z accuracy

before sample analysis.

The raw LC-MS data were processed using MS-DIAL (ver. 4.38)[124]. The

detailed MS-DIAL parameters are: MS1 tolerance, 0.01 Da; MS/MS tolerance, 0.05;

mass slice width, 0.05 Da; smoothing method, linear weighted moving average;

smoothing level, 3 scans; minimum peak width, 5 scans. Lipid features with high

quantitative confidence were selected by the following criteria: retention time was within

the gradient elution time (< 23 min); average intensity in QC samples is larger than 5-

fold of the intensity in method blank sample. Lipid identification was performed by

matching experimental precursor m/z, isotopic ratio and MS/MS spectrum against the

LipidBlast libraries embedded in MS-DIAL. To improve the quantification accuracy, the

measured MS signal intensities were corrected using serial diluted QC samples

following the reported workflow[125].

Quantification and statistical analysis. All microscopy images were quantified using

Fiji software[122]. For lipid droplet counts, a single optical slice through the middle of

the testis containing the hub was used with the exception of FLP-FRT experiment where

all lipid droplets within a GFP-negative cyst were counted (Figure 2I). All statistical

analyses were done using R (obtained from https://cran.r-project.org). With exception of

data concerning spatial distribution, and lipidomic data, Shapiro-Wilk test (via

shapiro.test in base R) was used to assess normality of distribution prior to testing for

significance. Kruskal-Wallis rank sum test (from the R package *coin*) and Dunn's test

(from the R package *dunn.test*) were used in place of Welch two-sample t-test and

Tukey's multiple comparison test when the assumption of normality was not met. For

testing differences in variance between two populations, F-test (via *var.test* in base R)

was used. For testing differences in spatial distribution, two-sample Kolmogorov-

Smirnov test (via *ks.test* in base R) was used. All *p*-values are indicated in figures;

extremely small *p*-values are listed as $p < 2.2 \times 10^{-16}$.

 RESOURCE TABLE

REAGENT or RESOURCE	SOURCE	RESOURCE #
Antibodies		
Anti-Vasa (1:200)	Gift from Dr. R. Lehman, MIT	
Anti-Eya (1:50)	Developmental Studies Hybridoma Bank (DSHB)	eya10H6
Anti-zfh1 (1:1000)	Gift from Dr. J. Skeath, WUSTL	
Anti-boule (1:1000)	Gift from Dr. S. Wasserman, UCSD	
Anti-phospho-histone H3 (1:1000)	Millipore Sigma	05-1354
Experimental models: *Drosophila melanogaster*		
w^1118	Bloomington *Drosophila* stock center	3605
CantonS	Bloomington *Drosophila* stock center	64349
OregonR	Bloomington *Drosophila* stock center	25211
bmm^1	Gift from Dr. R. Kühnlein	
bmm^rev	Gift from Dr. R. Kühnlein	
mdy[Qx25], cn[1], bw[1]/CyO, I(2)DTS513[1]	Bloomington *Drosophila* stock center	5095
y[1],w[67c23];P{lacW}Cse1[k03802],mdy[k03902]/CyO	Bloomington *Drosophila* stock center	10536
w[1118];P{GD1749}v6367 (UAS-mdy-RNAi)	Vienna *Drosophila* resource center	6367

REAGENT or RESOURCE	SOURCE	RESOURCE #
nos-GAL4::VP16	Bloomington *Drosophila* stock center	7303
Tj-GAL4	Gift from Dr. D. Godt, University of Toronto	
c587-GAL4	Bloomington *Drosophila* stock center	67747
Bam-GFP	[40]	
bmm-GFP	Gift from Dr. K. Kamei[57]	
GFP-LD	Gift from Dr. M. Welte[36]	
P{neoFRT}82B, bmm[1]	This study	
P{neoFRT}82B, bmm[rev]	This study	
bam-GFP, bmm[1]	This study	
bam-GFP, bmm[rev]	This study	
Software and algorithms		
Fiji	https://imagej.net/software/fiji/	
R	https://cran.r-project.org	

SUPPLEMENTAL TABLE AND FILES

Supplemental table 1 – Table showing identified lipid species from untargeted

lipidomic analysis.

Supplementary file 1 – Raw data and statistical outputs from Figure 1.

Supplementary file 2 – Raw data and statistical outputs from Figure 2, Supplemental

figure 2, and Supplemental figure 3.

Supplementary file 3 – Raw data and statistical outputs from Figure 3 and

Supplemental figure 4.

Supplementary file 4 – Raw data and statistical outputs from Figure 4 and

Supplemental figure 5.

SUPPLEMENTAL FIGURE LEGENDS

Supplemental Figure 1 related to Figure 1 – Cholesterol is absent from testis lipid

droplets. (A) Testes stained with BODIPY (A) to detect neutral lipids and Filipin III (A')

to detect free cholesterol. Scale bars=50 µm.

Supplemental Figure 2 related to Figure 2 – Expression of *brummer* and selected

lipid metabolic genes during spermatogenesis in germline and somatic lineages.

(A) Pseudotime trajectory of germline (black line) based on single-cell RNA sequencing

data [78]. Individual cells are labeled according to the annotation within the data set. (B)

Pseudotime trajectory of the somatic cells (black line) based on publicly available

single-cell RNA sequencing data [78]. Individual cells are labeled according to the

annotation within the data set. Only one trajectory (branch b) is marked and used for

panel D. (C) Rolling average of normalized transcript counts in the germline along the

trajectory shown in panel A are plotted as a black line on the upper panel. Composition

of cell types mapped on to the trajectory at each time point is shown at the bottom of

panel B. (D) Rolling average of normalized transcript counts in somatic cells plotted as a

black line along the trajectory shown in C (upper panel). Composition of cell types

mapped on to the trajectory at each time point is shown at the bottom of panel D.

Supplemental Figure 3 related to Figure 2 – *brummer* regulates lipid droplets in both germline and somatic cells of the testis. (A-D) Representative images of *bmm^rev^* (A and B) and *bmm^1^* (C and D) testes with somatic over-expression of *GFP-LD* (*Tj-GAL4>UAS-GFP-LD*). Panel F and H contain magnified images of the area indicated by the boxes in panel A and C, respectively. In *bmm^rev^* testes, LD were restricted to a region near the apical tip (A) of the testis in both somatic (B–B''' arrows) and germline cells (B–B''' arrowheads). In *bmm^1^* testes, LD were present in both somatic (C–D arrows) and germline cells (C–D arrowheads), near the apical tip of the testis in a region corresponding to early-stage germ cells and in the region corresponding to spermatocytes. (A,C) Scale bars=50 μm; (B,D) scale bars=20 μm.

Supplemental Figure 4 related to Figure 4 – Additional characterization of testis development and spermatogenesis in animals lacking *bmm*. (A) Testis size was smaller in *bmm^1^* mutant animals compared with *bmm^rev^* controls at <24 hr post-eclosion when raised at 29°C (A; Welch two-sample t-test). (B) The number of spermatid bundles was significantly lower in *bmm^1^* mutant animals compared with *bmm^rev^* controls at <24 hr post-eclosion when raised at 29°C (Kruskal-Wallis rank sum test). (C) Testis size was significantly smaller in *bmm^1^* mutant males compared with *bmm^rev^* control males at 14-days post-eclosion (Welch two-sample t-test). (D) While the median number of spermatid bundles was not significantly different between *bmm^1^* mutant males and *bmm^rev^* control males at 14 days post-eclosion (Welch two-sample t-test), 8/27 *bmm^1^* testis had no spermatid bundles, a phenotype absent in age-matched *bmm^rev^* males

(0/22) (p=0.0163, Pearson's Chi-squared test), suggesting a subtle defect is present.

(E) Food supplemented with 4% medium chain triglyceride (MCT), but not long chain

triglyceride (LCT), significantly increased testis length in *bmm¹* animals but had no effect

on this phenotype in *bmm^rev* control animals (one-way ANOVA with Tukey multiple

comparison test). (F) Food supplemented with 4% medium chain triglyceride

significantly increased the number of spermatid bundles in *bmm¹* testes but had no

effect on this phenotype in *bmm^rev* control animals (one-way ANOVA with Tukey

multiple comparison test). (G) Representative images of *bmm^rev* (G–G') or *bmm¹* (G''–

G''') testes stained for Fas3 (G and G'') and Vas (G' and G'''). Scale bars=25 µm. (H)

Quantification of hub area in *bmm^rev* or *bmm¹* testes showed a significantly larger hub

size in *bmm¹* testes (Welch two-sample t-test). (I) The number of germline stem cells

(GSC) undergoing mitosis (phospho-histone H3⁺ GSC/total GSC) was not significantly

different between *bmm¹* and *bmm^rev* testes (Kruskal-Wallis rank sum test). (J) The

distance between the hub and the first Bam-GFP positive cyst (Figure 3H) was

significantly higher in *bmm¹* testes than in *bmm^rev* testes (Welch two-sample t-test). (K)

All *bmm^rev* testes and most *bmm¹* testes contained spermatids when raised at 25°C;

however, the most advanced stage of spermatogenesis observed in the majority of

bmm¹ testes isolated from animals reared at 29°C was the spermatocyte stage. (L)

Testes isolated from *bmm¹* animals showed a significantly smaller Boule-positive area

than control testes (Welch two-sample t-test). (M) Testes isolated from *bmm¹* animals

contain fewer individualization complexes than *bmm^rev* control testes (Kruskal-Wallis

rank sum test). (N) Fewer waste bags were present in testes isolated from *bmm¹*

animals compared with *bmm^rev* control testes (Kruskal-Wallis rank sum test). (O)

Proportion of Zfh-1 positive clones that were homozygous for either *bmm¹* or *bmm^rev* at

3 and 14 days post-clone induction (Kruskal-Wallis rank sum test). (P) The location of

Zfh-1 positive clones homozygous for either *bmm¹* or *bmm^rev* at 3 and 14 days post-

clone induction measured as the distance from the center of the hub. Clones

homozygous for *bmm¹* locate significantly closer to the hub than the wildtype clones

(Kruskal-Wallis rank sum test).

Supplemental Figure 5 related to Figure 5 – Lipidomic analysis of animals lacking

bmm. (A) Higher fold-changes of triglycerides in *bmm¹* animals were associated with

less saturation in the acyl-groups (Kendall's rank correlation test). (B) Higher fold-

changes of triglycerides in *bmm¹* animals were associated with higher number of

carbons in the acyl-groups (Kendall's rank correlation test). Each dot represents a single

triglyceride species for panel B and C. (C) Volcano plot of identified lipids;

monoglycerides shown in blue and diglycerides shown in orange. Many monoglycerides

and diglycerides show increase in fold-change in *bmm¹* males. (D) The number of

carbon and the degree of saturation of monoglycerides (MAG) and diglycerides (DAG)

with significant changes in abundance between *bmm¹* and *bmm^rev* males. (E) Volcano

plot of identified lipids; fatty acids shown in magenta and acyl-carnitine shown in green.

Many fatty acids show an increase in fold-change while many acyl-carnitines show a

decrease in fold-change in *bmm¹* males. (F) The number of carbon and the degree of

saturation of fatty acids (FA) and acyl-carnitines (ACar) with significant changes in

abundance between *bmm¹* and *bmm^rev* males. (G) Volcano plot of identified lipids;

membrane lipids shown in yellow. (H) The number of carbon and the degree of

saturation of membrane lipids with significant changes in abundance between *bmm¹*

and *bmm^rev* males. For panel G and H, PC: phosphatidylcholine; PE:

phosphatidylethanolamine; PI: phosphatidylinositol; LPC: lysophosphatidylcholine; LPE:

lysophosphatidylethanolamine; SM: sphingomyelin; PG: phosphatidylglycerol. (I) Loss

of *mdy* function rescued the elevated number of LD in *bmm¹* testes to control levels

(one-way ANOVA with Tukey multiple comparison test). (J) Germline-specific loss of

mdy in *bmm¹* animals did not reduce GSC numbers, but the variance in GSC numbers

was significantly rescued (*nos-GAL4>+*; *bmm¹* vs *nos-GAL4>mdy RNAi*; *bmm¹*: *p*=4.5 ×

10^{-5}; *+>mdy RNAi*; *bmm¹* vs *nos-GAL4>mdy RNAi*; *bmm¹*: *p*=0.0082 by F-test).

727
728

REFERENCES

1. Brill JA, Yildirim S, Fabian L. Phosphoinositide signaling in sperm development. Semin Cell Dev Biol. 2016;59: 2–9. doi:10.1016/j.semcdb.2016.06.010

2. Keber R, Rozman D, Horvat S. Sterols in spermatogenesis and sperm maturation. J Lipid Res. 2013;54: 20–33. doi:10.1194/jlr.R032326

3. Rato L, Alves MG, Socorro S, Duarte AI, Cavaco JE, Oliveira PF. Metabolic regulation is important for spermatogenesis. Nat Rev Urol. 2012;9: 330–338. doi:10.1038/nrurol.2012.77

4. Wang C, Huang X. Lipid metabolism and Drosophila sperm development. Sci China Life Sci. 2012;55: 35–40. doi:10.1007/s11427-012-4274-2

5. de Cuevas M, Matunis EL. The stem cell niche: lessons from the Drosophila testis. Dev Camb Engl. 2011;138: 2861–2869. doi:10.1242/dev.056242

6. Demarco RS, Eikenes ÅH, Haglund K, Jones DL. Investigating Spermatogenesis in Drosophila melanogaster. Methods San Diego Calif. 2014;68: 218–227. doi:10.1016/j.ymeth.2014.04.020

7. Fabian L, Brill JA. Drosophila spermiogenesis: Big things come from little packages. Spermatogenesis. 2012;2: 197–212. doi:10.4161/spmg.21798

8. Brill JA, Hime GR, Scharer-Schuksz M, Fuller MT. A phospholipid kinase regulates actin organization and intercellular bridge formation during germline cytokinesis. Development. 2000;127: 3855.

9. Giansanti MG, Bonaccorsi S, Kurek R, Farkas RM, Dimitri P, Fuller MT, et al. The Class I PITP Giotto Is Required for Drosophila Cytokinesis. Curr Biol. 2006;16: 195–201. doi:10.1016/j.cub.2005.12.011

10. Wong R, Hadjiyanni I, Wei H-C, Polevoy G, McBride R, Sem K-P, et al. PIP2 Hydrolysis and Calcium Release Are Required for Cytokinesis in Drosophila Spermatocytes. Curr Biol. 2005;15: 1401–1406. doi:10.1016/j.cub.2005.06.060

11. Wong R, Fabian L, Forer A, Brill JA. Phospholipase C and myosin light chain kinase inhibition define a common step in actin regulation during cytokinesis. BMC Cell Biol. 2007;8: 15. doi:10.1186/1471-2121-8-15

12. Amoyel M, Hillion K-H, Margolis SR, Bach EA. Somatic stem cell differentiation is regulated by PI3K/Tor signaling in response to local cues. Dev Camb Engl. 2016;143: 3914. doi:10.1242/dev.139782

13. Fairchild MJ, Islam F, Tanentzapf G. Identification of genetic networks that act in the
somatic cells of the testis to mediate the developmental program of spermatogenesis.
PLOS Genet. 2017;13: e1007026. doi:10.1371/journal.pgen.1007026

14. Inaba M, Venkei ZG, Yamashita YM. The polarity protein Baz forms a platform for the
centrosome orientation during asymmetric stem cell division in the Drosophila male
germline. eLife. 2015;4: e04960. doi:10.7554/eLife.04960

15. Krahn MP, Klopfenstein DR, Fischer N, Wodarz A. Membrane Targeting of Bazooka/PAR-3 Is
Mediated by Direct Binding to Phosphoinositide Lipids. Curr Biol. 2010;20: 636–642.
doi:10.1016/j.cub.2010.01.065

16. Papagiannouli F, Berry CW, Fuller MT. The Dlg Module and Clathrin-Mediated Endocytosis
Regulate EGFR Signaling and Cyst Cell-Germline Coordination in the Drosophila Testis.
Stem Cell Rep. 2019;12: 1024–1040. doi:10.1016/j.stemcr.2019.03.008

17. Ueishi S, Shimizu H, Inoue YH. Male Germline Stem Cell Division and Spermatocyte Growth
Require Insulin Signaling in Drosophila. Cell Struct Funct. 2009;34: 61–69.
doi:10.1247/csf.08042

18. Phan VH, Herr DR, Panton D, Fyrst H, Saba JD, Harris GL. Disruption of Sphingolipid
Metabolism Elicits Apoptosis-Associated Reproductive Defects in Drosophila. Dev Biol.
2007;309: 329–341. doi:10.1016/j.ydbio.2007.07.021

19. Steinhauer J, Gijón MA, Riekhof WR, Voelker DR, Murphy RC, Treisman JE. Drosophila
Lysophospholipid Acyltransferases Are Specifically Required for Germ Cell Development.
Mol Biol Cell. 2009;20: 5224–5235. doi:10.1091/mbc.e09-05-0382

20. Szafer-Glusman E, Giansanti MG, Nishihama R, Bolival B, Pringle J, Gatti M, et al. A Role for
Very-Long-Chain Fatty Acids in Furrow Ingression during Cytokinesis in Drosophila
Spermatocytes. Curr Biol. 2008;18: 1426–1431. doi:10.1016/j.cub.2008.08.061

21. Ben-David G, Miller E, Steinhauer J. Drosophila spermatid individualization is sensitive to
temperature and fatty acid metabolism. Spermatogenesis. 2015;5.
doi:10.1080/21565562.2015.1006089

22. Jung A, Hollmann M, Schafer MA. The fatty acid elongase NOA is necessary for viability and
has a somatic role in Drosophila sperm development. J Cell Sci. 2007;120: 2924–2934.
doi:10.1242/jcs.006551

23. Rabionet M, Bayerle A, Jennemann R, Heid H, Fuchser J, Marsching C, et al. Male meiotic
cytokinesis requires ceramide synthase 3-dependent sphingolipids with unique
membrane anchors. Hum Mol Genet. 2015;24: 4792–4808. doi:10.1093/hmg/ddv204

24. Santiago Valtierra FX, Peñalva DA, Luquez JM, Furland NE, Vásquez C, Reyes JG, et al. Elovl4
and Fa2h expression during rat spermatogenesis: a link to the very-long-chain PUFAs

typical of germ cell sphingolipids. J Lipid Res. 2018;59: 1175–1189.
doi:10.1194/jlr.M081885

25. Zadravec D, Tvrdik P, Guillou H, Haslam R, Kobayashi T, Napier JA, et al. ELOVL2 controls the
level of n-6 28:5 and 30:5 fatty acids in testis, a prerequisite for male fertility and sperm
maturation in mice. J Lipid Res. 2011;52: 245–255. doi:10.1194/jlr.M011346

26. Walther TC, Farese RV. Lipid Droplets and Cellular Lipid Metabolism. Annu Rev Biochem.
2012;81: 687–714. doi:10.1146/annurev-biochem-061009-102430

27. Wat LW, Chao C, Bartlett R, Buchanan JL, Millington JW, Chih HJ, et al. A role for triglyceride
lipase brummer in the regulation of sex differences in Drosophila fat storage and
breakdown. Hassan BA, editor. PLOS Biol. 2020;18: e3000595.
doi:10.1371/journal.pbio.3000595

28. Thiele C, Spandl J. Cell biology of lipid droplets. Curr Opin Cell Biol. 2008;20: 378–385.
doi:10.1016/j.ceb.2008.05.009

29. Bosma M. Lipid droplet dynamics in skeletal muscle. Exp Cell Res. 2016;340: 180–186.
doi:10.1016/j.yexcr.2015.10.023

30. Grönke S, Müller G, Hirsch J, Fellert S, Andreou A, Haase T, et al. Dual Lipolytic Control of
Body Fat Storage and Mobilization in Drosophila. PLOS Biol. 2007;5: e137.
doi:10.1371/journal.pbio.0050137

31. Rambold AS, Cohen S, Lippincott-Schwartz J. Fatty Acid Trafficking in Starved Cells:
Regulation by Lipid Droplet Lipolysis, Autophagy, and Mitochondrial Fusion Dynamics.
Dev Cell. 2015;32: 678–692. doi:10.1016/j.devcel.2015.01.029

32. Dichlberger A, Schlager S, Maaninka K, Schneider WJ, Kovanen PT. Adipose triglyceride
lipase regulates eicosanoid production in activated human mast cells. J Lipid Res.
2014;55: 2471–2478. doi:10.1194/jlr.M048553

33. Mitsche MA, Hobbs HH, Cohen JC. Patatin-like phospholipase domain–containing protein 3
promotes transfers of essential fatty acids from triglycerides to phospholipids in hepatic
lipid droplets. J Biol Chem. 2018;293: 6958–6968. doi:10.1074/jbc.RA118.002333

34. Rajakumari S, Rajasekharan R, Daum G. Triacylglycerol lipolysis is linked to sphingolipid and
phospholipid metabolism of the yeast Saccharomyces cerevisiae☆. Biochim Biophys Acta
BBA - Mol Cell Biol Lipids. 2010;1801: 1314–1322. doi:10.1016/j.bbalip.2010.08.004

35. Schlager S, Goeritzer M, Jandl K, Frei R, Vujic N, Kolb D, et al. Adipose triglyceride lipase acts
on neutrophil lipid droplets to regulate substrate availability for lipid mediator synthesis.
J Leukoc Biol. 2015;98: 837–850. doi:10.1189/jlb.3A0515-206R

36. Zanghellini J, Natter K, Jungreuthmayer C, Thalhammer A, Kurat CF, Gogg-Fassolter G, et al.
Quantitative modeling of triacylglycerol homeostasis in yeast - metabolic requirement for
lipolysis to promote membrane lipid synthesis and cellular growth: Triacylglycerol
mobilization in yeast. FEBS J. 2008;275: 5552–5563. doi:10.1111/j.1742-
4658.2008.06681.x

37. Bailey AP, Koster G, Guillermier C, Hirst EMA, MacRae JI, Lechene CP, et al. Antioxidant Role
for Lipid Droplets in a Stem Cell Niche of Drosophila. Cell. 2015;163: 340–353.
doi:10.1016/j.cell.2015.09.020

38. Liu L, MacKenzie KR, Putluri N, Maletić-Savatić M, Bellen HJ. The Glia-Neuron Lactate
Shuttle and Elevated ROS Promote Lipid Synthesis in Neurons and Lipid Droplet
Accumulation in Glia via APOE/D. Cell Metab. 2017;26: 719-737.e6.
doi:10.1016/j.cmet.2017.08.024

39. Nguyen TB, Louie SM, Daniele JR, Tran Q, Dillin A, Zoncu R, et al. DGAT1-Dependent Lipid
Droplet Biogenesis Protects Mitochondrial Function during Starvation-Induced
Autophagy. Dev Cell. 2017;42: 9-21.e5. doi:10.1016/j.devcel.2017.06.003

40. Henne M. And three's a party: lysosomes, lipid droplets, and the ER in lipid trafficking and
cell homeostasis. Curr Opin Cell Biol. 2019;59: 40–49. doi:10.1016/j.ceb.2019.02.011

41. Welte MA. Expanding Roles for Lipid Droplets. Curr Biol. 2015;25: R470–R481.
doi:10.1016/j.cub.2015.04.004

42. Casado ME, Pastor O, Mariscal P, Canfrán-Duque A, Martínez-Botas J, Kraemer FB, et al.
Hormone-sensitive lipase deficiency disturbs the fatty acid composition of mouse testis.
Prostaglandins Leukot Essent Fatty Acids. 2013;88: 227–233.
doi:10.1016/j.plefa.2012.12.005

43. El-Shehawi AM, El-Shazly S, Ahmed M, Alkafafy M, Sayed S, Farouk S, et al. Transcriptome
Analysis of Testis from HFD-Induced Obese Rats (Rattus norvigicus) Indicated
Predisposition for Male Infertility. Int J Mol Sci. 2020;21: 6493.
doi:10.3390/ijms21186493

44. Wang W, Wei S, Li L, Su X, Du C, Li F, et al. Proteomic analysis of murine testes lipid
droplets. Sci Rep. 2015;5: 12070. doi:10.1038/srep12070

45. Bajpai M, Gupta G, Setty BS. Changes in carbohydrate metabolism of testicular germ cells
during meiosis in the rat. Eur J Endocrinol. 1998;138: 322–327.

46. Kerr JB, Kretser DMD. CYCLIC VARIATIONS IN SERTOLI CELL LIPID CONTENT THROUGHOUT
THE SPERMATOGENIC CYCLE IN THE RAT. Reproduction. 1975;43: 1–8.
doi:10.1530/jrf.0.0430001

47. Mori H, Christensen AK. Morphometric analysis of Leydig cells in the normal rat testis. J Cell
Biol. 1980;84: 340–354. doi:10.1083/jcb.84.2.340

48. Paniagua R, Rodríguez MC, Nistal M, Fraile B, Amat P. Changes in the lipid inclusion/Sertoli
cell cytoplasm area ratio during the cycle of the human seminiferous epithelium.
Reproduction. 1987;80: 335–341. doi:10.1530/jrf.0.0800335

49. Demarco RS, Uyemura BS, D'Alterio C, Jones DL. Mitochondrial fusion regulates lipid
homeostasis and stem cell maintenance in the Drosophila testis. Nat Cell Biol. 2019;21:
710–720. doi:10.1038/s41556-019-0332-3

50. Chen M, Wang H, Li X, Li N, Xu G, Meng Q. PLIN1 deficiency affects testicular gene
expression at the meiotic stage in the first wave of spermatogenesis. Gene. 2014;543:
212–219. doi:10.1016/j.gene.2014.04.021

51. Hermo L, Chung S, Gregory M, Smith CE, Wang SP, El-Alfy M, et al. Alterations in the testis
of hormone sensitive lipase-deficient mice is associated with decreased sperm counts,
sperm motility, and fertility. Mol Reprod Dev. 2008;75: 565–577. doi:10.1002/mrd.20800

52. Masaki H, Kim N, Nakamura H, Kumasawa K, Kamata E, Hirano K, et al. Long-chain fatty acid
triglyceride (TG) metabolism disorder impairs male fertility: a study using adipose
triglyceride lipase deficient mice. MHR Basic Sci Reprod Med. 2017;23: 452–460.
doi:10.1093/molehr/gax031

53. Freeman DA, Ascoli M. Studies on the source of cholesterol used for steroid biosynthesis in
cultured Leydig tumor cells. J Biol Chem. 1982;257: 14231–14238.

54. Wang F, Chen Z, Ren X, Tian Y, Wang F, Liu C, et al. Hormone-sensitive lipase deficiency
alters gene expression and cholesterol content of mouse testis. Reproduction. 2017;153:
175–185. doi:10.1530/REP-16-0484

55. de Cuevas M, Matunis EL. The stem cell niche: lessons from the Drosophila testis.
Development. 2011;138: 2861–2869. doi:10.1242/dev.056242

56. Walther TC, Chung J, Jr RVF. Lipid Droplet Biogenesis. Annu Rev Cell Dev Biol. 2017;33: 491–
510. doi:10.1146/annurev-cellbio-100616-060608

57. Boyle M, DiNardo S. Specification, migration and assembly of the somatic cells of the
Drosophila gonad. Development. 1995;121: 1815–1825. doi:10.1242/dev.121.6.1815

58. Yu YV, Li Z, Rizzo NP, Einstein J, Welte MA. Targeting the motor regulator Klar to lipid
droplets. BMC Cell Biol. 2011;12: 9. doi:10.1186/1471-2121-12-9

59. Leatherman JL, DiNardo S. Zfh-1 Controls Somatic Stem Cell Self-Renewal in the Drosophila
Testis and Nonautonomously Influences Germline Stem Cell Self-Renewal. Cell Stem Cell.
2008;3: 44–54. doi:10.1016/j.stem.2008.05.001

60. Fabrizio JJ, Boyle M, DiNardo S. A somatic role for eyes absent (eya) and sine oculis (so) in
drosophila spermatocyte development. Dev Biol. 2003;258: 117–128.
doi:10.1016/S0012-1606(03)00127-1

61. Chen D, McKearin DM. A discrete transcriptional silencer in the bam gene determines
asymmetric division of the Drosophila germline stem cell. Development. 2003;130: 1159–
1170. doi:10.1242/dev.00325

62. Insco ML, Leon A, Tam CH, McKearin DM, Fuller MT. Accumulation of a differentiation
regulator specifies transit amplifying division number in an adult stem cell lineage. Proc
Natl Acad Sci U S A. 2009;106: 22311–22316. doi:10.1073/pnas.0912454106

63. Chen D, Wu C, Zhao S, Geng Q, Gao Y, Li X, et al. Three RNA Binding Proteins Form a
Complex to Promote Differentiation of Germline Stem Cell Lineage in Drosophila. PLoS
Genet. 2014;10. doi:10.1371/journal.pgen.1004797

64. Gonczy P, Matunis E, DiNardo S. bag-of-marbles and benign gonial cell neoplasm act in the
germline to restrict proliferation during Drosophila spermatogenesis. Development.
1997;124: 4361–4371.

65. Athenstaedt K, Daum G. YMR313c/TGL3 Encodes a Novel Triacylglycerol Lipase Located in
Lipid Particles of Saccharomyces cerevisiae*. J Biol Chem. 2003;278: 23317–23323.
doi:10.1074/jbc.M302577200

66. Chitraju C, Trötzmüller M, Hartler J, Wolinski H, Thallinger GG, Haemmerle G, et al. The
impact of genetic stress by ATGL deficiency on the lipidome of lipid droplets from murine
hepatocytes. J Lipid Res. 2013;54: 2185–2194. doi:10.1194/jlr.M037952

67. Grönke S, Mildner A, Fellert S, Tennagels N, Petry S, Müller G, et al. Brummer lipase is an
evolutionary conserved fat storage regulator in Drosophila. Cell Metab. 2005;1: 323–330.
doi:10.1016/j.cmet.2005.04.003

68. Haemmerle G, Lass A, Zimmermann R, Gorkiewicz G, Meyer C, Rozman J, et al. Defective
Lipolysis and Altered Energy Metabolism in Mice Lacking Adipose Triglyceride Lipase.
Science. 2006;312: 734–737. doi:10.1126/science.1123965

69. Haemmerle G, Moustafa T, Woelkart G, Büttner S, Schmidt A, van de Weijer T, et al. ATGL-
mediated fat catabolism regulates cardiac mitochondrial function via PPAR-α and PGC-1.
Nat Med. 2011;17: 1076–1085. doi:10.1038/nm.2439

70. Huijsman E, van de Par C, Economou C, van der Poel C, Lynch GS, Schoiswohl G, et al.
Adipose triacylglycerol lipase deletion alters whole body energy metabolism and impairs
exercise performance in mice. Am J Physiol-Endocrinol Metab. 2009;297: E505–E513.
doi:10.1152/ajpendo.00190.2009

71. Korbelius M, Vujic N, Sachdev V, Obrowsky S, Rainer S, Gottschalk B, et al. ATGL/CGI-58-
Dependent Hydrolysis of a Lipid Storage Pool in Murine Enterocytes. Cell Rep. 2019;28:
1923-1934.e4. doi:10.1016/j.celrep.2019.07.030

72. Kurat CF, Natter K, Petschnigg J, Wolinski H, Scheuringer K, Scholz H, et al. Obese Yeast:
Triglyceride Lipolysis Is Functionally Conserved from Mammals to Yeast *. J Biol Chem.
2006;281: 491–500. doi:10.1074/jbc.M508414200

73. Zimmermann R, Strauss JG, Haemmerle G, Schoiswohl G, Birner-Gruenberger R, Riederer M,
et al. Fat Mobilization in Adipose Tissue Is Promoted by Adipose Triglyceride Lipase.
Science. 2004;306: 1383–1386. doi:10.1126/science.1100747

74. Attané C, Peyot M-L, Lussier R, Poursharifi P, Zhao S, Zhang D, et al. A beta cell ATGL-
lipolysis/adipose tissue axis controls energy homeostasis and body weight via insulin
secretion in mice. Diabetologia. 2016;59: 2654–2663. doi:10.1007/s00125-016-4105-2

75. Lee JH, Kong J, Jang JY, Han JS, Ji Y, Lee J, et al. Lipid Droplet Protein LID-1 Mediates ATGL-1-
Dependent Lipolysis during Fasting in Caenorhabditis elegans. Mol Cell Biol. 2014;34:
4165–4176. doi:10.1128/MCB.00722-14

76. Tuohetahuntila M, Molenaar MR, Spee B, Brouwers JF, Houweling M, Vaandrager AB, et al.
ATGL and DGAT1 are involved in the turnover of newly synthesized triacylglycerols in
hepatic stellate cells. J Lipid Res. 2016;57: 1162–1174. doi:10.1194/jlr.M066415

77. Men TT, Thanh DNV, Yamaguchi M, Suzuki T, Hattori G, Arii M, et al. A *Drosophila* Model for
Screening Antiobesity Agents. BioMed Res Int. 2016;2016: 1–10.
doi:10.1155/2016/6293163

78. Li H, Janssens J, Waegeneer MD, Kolluru SS, Davie K, Gardeux V, et al. Fly Cell Atlas: a single-
cell transcriptomic atlas of the adult fruit fly. bioRxiv; 2021. p. 2021.07.04.451050.
doi:10.1101/2021.07.04.451050

79. Xu T, Rubin GM. Analysis of genetic mosaics in developing and adult Drosophila tissues.
Development. 1993;117: 1223–1237.

80. Kim N, Nakamura H, Masaki H, Kumasawa K, Hirano K, Kimura T. Effect of lipid metabolism
on male fertility. Biochem Biophys Res Commun. 2017;485: 686–692.
doi:10.1016/j.bbrc.2017.02.103

81. Lasko PF, Ashburner M. The product of the Drosophila gene vasa is very similar to
eukaryotic initiation factor-4A. Nature. 1988;335: 611–617. doi:10.1038/335611a0

82. Resende LPF, Boyle M, Tran D, Fellner T, Jones DL. Headcase Promotes Cell Survival and
Niche Maintenance in the Drosophila Testis. Bökel C, editor. PLoS ONE. 2013;8: e68026.
doi:10.1371/journal.pone.0068026

83. Kiger AA, White-Cooper H, Fuller MT. Somatic support cells restrict germline stem cell self-
renewal and promote differentiation. Nature. 2000;407: 750–754. doi:10.1038/35037606

84. Lin TY, Viswanathan S, Wood C, Wilson PG, Wolf N, Fuller MT. Coordinate developmental
control of the meiotic cell cycle and spermatid differentiation in Drosophila males.
Development. 1996;122: 1331–1341.

85. Eichmann TO, Kumari M, Haas JT, Farese RV, Zimmermann R, Lass A, et al. Studies on the
substrate and stereo/regioselectivity of adipose triglyceride lipase, hormone-sensitive
lipase, and diacylglycerol-O-acyltransferases. J Biol Chem. 2012;287: 41446–41457.
doi:10.1074/jbc.M112.400416

86. Schweiger M, Schreiber R, Haemmerle G, Lass A, Fledelius C, Jacobsen P, et al. Adipose
Triglyceride Lipase and Hormone-sensitive Lipase Are the Major Enzymes in Adipose
Tissue Triacylglycerol Catabolism *. J Biol Chem. 2006;281: 40236–40241.
doi:10.1074/jbc.M608048200

87. Missaglia S, Maggi L, Mora M, Gibertini S, Blasevich F, Agostoni P, et al. Late onset of
neutral lipid storage disease due to novel PNPLA2 mutations causing total loss of lipase
activity in a patient with myopathy and slight cardiac involvement. Neuromuscul Disord.
2017;27: 481–486. doi:10.1016/j.nmd.2017.01.011

88. Williams ML, Coleman RA, Placezk D, Grunfeld C. Neutral lipid storage disease: a possible
functional defect in phospholipid-linked triacylglycerol metabolism. Biochim Biophys Acta
BBA - Mol Basis Dis. 1991;1096: 162–169. doi:10.1016/0925-4439(91)90055-E

89. Yang L, Liang J, Lam SM, Yavuz A, Shui G, Ding M, et al. Neuronal lipolysis participates in
PUFA-mediated neural function and neurodegeneration. EMBO Rep. 2020;21.
doi:10.15252/embr.202050214

90. Nazario-Yepiz NO, Fernández Sobaberas J, Lyman R, Campbell MR, Shankar V, Anholt RRH,
et al. Physiological and metabolomic consequences of reduced expression of the
Drosophila brummer triglyceride Lipase. Melkani GC, editor. PLOS ONE. 2021;16:
e0255198. doi:10.1371/journal.pone.0255198

91. Giedt MS, Thomalla JM, Johnson MR, Lai ZW, Tootle TL, Welte MA. Adipose triglyceride
lipase promotes prostaglandin-dependent actin remodeling by regulating substrate
release from lipid droplets. Cell Biology; 2021 Aug. doi:10.1101/2021.08.02.454724

92. Beller M, Bulankina AV, Hsiao H-H, Urlaub H, Jäckle H, Kühnlein RP. PERILIPIN-Dependent
Control of Lipid Droplet Structure and Fat Storage in Drosophila. Cell Metab. 2010;12:
521–532. doi:10.1016/j.cmet.2010.10.001

93. Buszczak M, Lu X, Segraves WA, Chang TY, Cooley L. Mutations in the midway Gene Disrupt
a Drosophila Acyl Coenzyme A: Diacylglycerol Acyltransferase. Genetics. 2002;160: 1511–
1518.

1002 94. Martínez BA, Hoyle RG, Yeudall S, Granade ME, Harris TE, Castle JD, et al. Innate immune
1003   signaling in Drosophila shifts anabolic lipid metabolism from triglyceride storage to
1004   phospholipid synthesis to support immune function. PLOS Genet. 2020;16: e1009192.
1005   doi:10.1371/journal.pgen.1009192

1006 95. Freeman DA, Ascoli M. Studies on the source of cholesterol used for steroid biosynthesis in
1007   cultured Leydig tumor cells. J Biol Chem. 1982;257: 14231–14238.

1008 96. Wang F, Chen Z, Ren X, Tian Y, Wang F, Liu C, et al. Hormone-sensitive lipase deficiency
1009   alters gene expression and cholesterol content of mouse testis. Reproduction. 2017;153:
1010   175–185. doi:10.1530/REP-16-0484

1011 97. Walther TC, Farese RV. Lipid droplets and cellular lipid metabolism. Annu Rev Biochem.
1012   2012;81: 687–714. doi:10.1146/annurev-biochem-061009-102430

1013 98. Olzmann JA, Carvalho P. Dynamics and functions of lipid droplets. Nat Rev Mol Cell Biol.
1014   2019;20: 137–155. doi:10.1038/s41580-018-0085-z

1015 99. Birse RT, Choi J, Reardon K, Rodriguez J, Graham S, Diop S, et al. High-fat-diet-induced
1016   obesity and heart dysfunction are regulated by the TOR pathway in Drosophila. Cell
1017   Metab. 2010;12: 533–44. doi:10.1016/j.cmet.2010.09.014

1018 100. Molaei M, Vandehoef C, Karpac J. NF-κB Shapes Metabolic Adaptation by Attenuating
1019   Foxo-Mediated Lipolysis in Drosophila. Dev Cell. 2019;49: 802-810.e6.
1020   doi:10.1016/j.devcel.2019.04.009

1021 101. Alic N, Andrews TD, Giannakou ME, Papatheodorou I, Slack C, Hoddinott MP, et al.
1022   Genome-wide dFOXO targets and topology of the transcriptomic response to stress and
1023   insulin signalling. Mol Syst Biol. 2011;7: 502. doi:10.1038/msb.2011.36

1024 102. Junger MA, Rintelen F, Stocker H, Wasserman JD, Vegh M, Radimerski T, et al. The
1025   Drosophila forkhead transcription factor FOXO mediates the reduction in cell number
1026   associated with reduced insulin signaling. J Biol. 2003;2: 20. doi:10.1186/1475-4924-2-20

1027 103. Zinke I, Schutz CS, Katzenberger JD, Bauer M, Pankratz MJ. Nutrient control of gene
1028   expression in Drosophila: microarray analysis of starvation and sugar-dependent
1029   response. EMBO J. 2002;21: 6162–73.

1030 104. Puig O, Tjian R. Transcriptional feedback control of insulin receptor by dFOXO/FOXO1.
1031   Genes Dev. 2005;19: 2435–2446. doi:10.1101/gad.1340505

1032 105. Kang P, Chang K, Liu Y, Bouska M, Birnbaum A, Karashchuk G, et al. Drosophila Kruppel
1033   homolog 1 represses lipolysis through interaction with dFOXO. Sci Rep. 2017;7: 16369.
1034   doi:10.1038/s41598-017-16638-1

106. Bartz R, Zehmer JK, Zhu M, Chen Y, Serrero G, Zhao Y, et al. Dynamic activity of lipid
droplets: protein phosphorylation and GTP-mediated protein translocation. J Proteome
Res. 2007;6: 3256–3265. doi:10.1021/pr070158j

107. Pagnon J, Matzaris M, Stark R, Meex RCR, Macaulay SL, Brown W, et al. Identification and
functional characterization of protein kinase A phosphorylation sites in the major lipolytic
protein, adipose triglyceride lipase. Endocrinology. 2012;153: 4278–4289.
doi:10.1210/en.2012-1127

108. Narbonne P, Roy R. Caenorhabditis elegans dauers need LKB1/AMPK to ration lipid
reserves and ensure long-term survival. Nature. 2009;457: 210–214.
doi:10.1038/nature07536

109. Ahmadian M, Abbott MJ, Tang T, Hudak CSS, Kim Y, Bruss M, et al. Desnutrin/ATGL is
regulated by AMPK and is required for a brown adipose phenotype. Cell Metab. 2011;13:
739–748. doi:10.1016/j.cmet.2011.05.002

110. Amoyel M, Hillion K-H, Margolis SR, Bach EA. Somatic stem cell differentiation is regulated
by PI3K/Tor signaling in response to local cues. Dev Camb Engl. 2016;143: 3914–3925.
doi:10.1242/dev.139782

111. Amoyel M, Simons BD, Bach EA. Neutral competition of stem cells is skewed by
proliferative changes downstream of Hh and Hpo. EMBO J. 2014;33: 2295–2313.
doi:10.15252/embj.201387500

112. Ueishi S, Shimizu H, H. Inoue Y. Male Germline Stem Cell Division and Spermatocyte
Growth Require Insulin Signaling in Drosophila. Cell Struct Funct. 2009;34: 61–69.
doi:10.1247/csf.08042

113. Hof-Michel S, Cigoja L, Huhn S, Bökel C. Innate immune signalling drives loser cell
elimination during stem cell competition in the *Drosophila* testis. Developmental Biology;
2020 Mar. doi:10.1101/2020.03.05.979161

114. Couderc J-L, Richard G, Vachias C, Mirouse V. Drosophila LKB1 is required for the assembly
of the polarized actin structure that allows spermatid individualization. White-Cooper H,
editor. PLOS ONE. 2017;12: e0182279. doi:10.1371/journal.pone.0182279

115. Steinhauer J, Statman B, Fagan JK, Borck J, Surabhi S, Yarikipati P, et al. Combover interacts
with the axonemal component Rsp3 and is required for Drosophila sperm
individualization. Dev Camb Engl. 2019;146: dev179275. doi:10.1242/dev.179275

116. Tartarin P, Guibert E, Touré A, Ouiste C, Leclerc J, Sanz N, et al. Inactivation of AMPKα1
induces asthenozoospermia and alters spermatozoa morphology. Endocrinology.
2012;153: 3468–3481. doi:10.1210/en.2011-1911

117. Martin-Hidalgo D, Hurtado de Llera A, Calle-Guisado V, Gonzalez-Fernandez L, Garcia-
Marin L, Bragado MJ. AMPK Function in Mammalian Spermatozoa. Int J Mol Sci. 2018;19:
3293. doi:10.3390/ijms19113293

118. Pitetti J-L, Calvel P, Zimmermann C, Conne B, Papaioannou MD, Aubry F, et al. An Essential
Role for Insulin and IGF1 Receptors in Regulating Sertoli Cell Proliferation, Testis Size, and
FSH Action in Mice. Mol Endocrinol. 2013;27: 814–827. doi:10.1210/me.2012-1258

119. Thurmond J, Goodman JL, Strelets VB, Attrill H, Gramates LS, Marygold SJ, et al. FlyBase
2.0: the next generation. Nucleic Acids Res. 2018;47: D759–D765.
doi:10.1093/nar/gky1003

120. White-Cooper H. Spermatogenesis. In: Henderson DS, editor. Drosophila Cytogenetics
Protocols. Totowa, NJ: Humana Press; 2004. pp. 45–75. doi:10.1385/1-59259-665-7:45

121. Fabian L, Brill JA. Drosophila spermiogenesis. Spermatogenesis. 2012;2: 197–212.
doi:10.4161/spmg.21798

122. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.
doi:10.1038/nmeth.2019

123. Yu H, Villanueva N, Bittar T, Arsenault E, Labonté B, Huan T. Parallel metabolomics and
lipidomics enables the comprehensive study of mouse brain regional metabolite and lipid
patterns. Anal Chim Acta. 2020;1136: 168–177. doi:10.1016/j.aca.2020.09.051

124. Tsugawa H, Cajka T, Kind T, Ma Y, Higgins B, Ikeda K, et al. MS-DIAL: data-independent
MS/MS deconvolution for comprehensive metabolome analysis. Nat Methods. 2015;12:
523–526. doi:10.1038/nmeth.3393

125. Yu H, Huan T. Patterned Signal Ratio Biases in Mass Spectrometry-Based Quantitative
Metabolomics. Anal Chem. 2021;93: 2254–2262. doi:10.1021/acs.analchem.0c04113

1093

1094
1095

1096

FIGURE 1

FIGURE 2

C   Number of lipid droplets during germline development

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

Supplemental figure 1 - related to Figure 1

Supplemental Figure 2 - related to Figure 2

Supplemental figure 3 - related to Figure 2

Supplemental figure 4 - related to Figure 4

Supplemental figure 5 - related to Figure 5