Parallels and divergences in landscape genetic and metacommunity patterns in

zooplankton inhabiting soda pans

Zsófia Horváth[1,2,3,*], Tsegazeabe Hadush Haileselasie[2,4], Csaba F. Vad[1,2,3], Robert Ptacnik[3],

Luc De Meester[2,5,6,7]

[1]Institute of Aquatic Ecology, HUN-REN Centre for Ecological Research, Budapest, Hungary

[2]Laboratory of Aquatic Ecology and Evolutionary Biology, KU Leuven, Leuven, Belgium

[3]WasserCluster Lunz, Lunz am See, Austria

[4]Mekelle University, College of Natural and Computational Sciences (CNCS), Department of

Biology Mekelle, Tigray, Ethiopia

[5]Leibniz Institut für Gewässerökologie und Binnenfischerei (IGB), Müggelseedamm 310,

12587 Berlin, Germany

[6]Institute of Biology, Freie Universität Berlin, Königin-Luise-Strasse 1-3, 14195 Berlin,

Germany

[7]Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), Königin-Luise-Str.

2-4, 14195 Berlin, Germany

*Corresponding author: Zsófia Horváth, email: hhzsofia@gmail.com

ORCID: 0000-0003-2857-1094, 0000-0003-4567-620X, 0000-0003-4744-0698, 0000-0001-

7176-7653, 0000-0001-5433-6843

Running head: Landscape genetic and metacommunity patterns

Abstract

Ecological processes maintaining landscape genetic variation and metacommunity structure in

natural landscapes have traditionally been studied in isolation. Their integrated study may

hold important information as to what extent the effect of major ecological processes are

species- or landscape-specific, resulting in a more coherent picture on the spatial organization

of biodiversity. Here, we explicitly compared the relative importance of spatial and

environmental drivers of both cladoceran metacommunity structure as well as landscape

genetic structure of its most widespread member, the water flea *Daphnia magna*, in soda pans

of the Seewinkel region in Austria. This landscape of soda pans is characterized by strong

environmental gradients and unidirectional wind acting as a key dispersal agent among these

temporary habitats. Our study shows both parallels and divergences in the relative importance

of local environmental sorting and spatial connectivity in determining landscape genetic

versus metacommunity structure. The metacommunity is structured primarily by the

environment, while in the *D. magna* metapopulation, the spatial signal is predominant. The

much weaker environmental signal in *Daphnia* can be explained by the fact that the

microsatellite markers are presumably neutral and was confirmed by a per-allele analysis. An

important parallel between metacommunity and landscape genetic structure is the strong

signal of the prevailing wind direction in determining the spatial pattern. This suggests that

for both community assembly in cladocerans and population assembly in *D. magna*, wind

plays an important role in determining connectivity among soda pans, thereby affecting

dispersal and colonization rates, influencing both local species and genetic composition.

Introduction

Dispersal is essential for maintaining connectivity among local populations and communities. It drives landscape-level heterogeneity in genetic and species composition, in interaction with other processes including local environmental sorting (selection) and drift (Vellend and Geber 2005). This implies that at least theoretically, metapopulations and metacommunities are in essence structured by the same processes (Vellend 2016). Drivers of landscape genetic structure are indeed expected to show strong parallels to the drivers of variation in species composition in local communities across landscapes (cf. metacommunity structure; Leibold et al. 2004). In both cases, dispersal is linked to landscape connectivity, and can lead to a pattern of increasing similarities with decreasing geographic distance (i.e., distance decay of similarity). Similarly, in both cases, environmental heterogeneity can structure patterns of convergence and divergence. Vellend (2010, 2016) highlighted the parallels between metacommunity (species sorting, dispersal, neutral dynamics and speciation) and metapopulation processes (natural selection, gene flow, neutral dynamics and mutation). Yet, due to a lack of explicit comparative studies in the same landscapes, little is known on the relative contribution of these processes to shaping landscape genetic variation and species composition.

The explicit comparative test of the drivers of landscape genetic and metacommunity structure requires studies performed in the same landscape with the same statistical tools. In metacommunity ecology, several analyses have been developed to disentangle spatial from environmental effects (Borcard et al. 1992; Peres-Neto et al. 2006), followed by numerous empirical studies that tested the relative role of environmental sorting and dispersal on biodiversity and community composition. Landscape genetics similarly involves testing for the relative importance of local environment and dispersal on genetic metapopulation

structure and gene flow (Manel *et al.* 2003; Guillot *et al.* 2005). While population genetic

studies traditionally focused on geographic distance as a key structuring factor (isolation-by-

distance; Bohonak 1999, 2002), there is increasing evidence for strong environmental

gradients mediating effective gene flow and structuring landscape genetic variation (Nosil et

al. 2005; Orsini et al. 2013; Sexton et al. 2014; Moncada et al. 2021). Given these parallels,

the same multivariate approaches that are applied in metacommunity ecology can deliver

information on the relative importance of spatial and environmental predictors for explaining

not only community composition in a landscape, but also landscape genetic structure (Orsini

et al. 2013). While there is an increasing interest in multi-species approaches (Manel and

Holderegger 2013; Wang and Bradburd 2014; Hein et al. 2021) resulting in a growing number

of multi-species landscape genetic studies (Manier and Arnold 2006; Goldberg and Waits

2010; Frantz et al. 2012; Richardson 2012), explicit comparisons of metacommunity and

metapopulation level patterns are largely lacking.

Here, we compare metacommunity structure of cladoceran zooplankton to landscape

genetic variation in its dominant member, the water flea *Daphnia magna*, in the same set of

habitats, saline temporary waters (soda pans) in the Seewinkel region of Austria (**Fig. 1**).

Specifically, we aim to explicitly test the relative strength of major structuring forces (i.e.,

environment and space) across the two levels of ecological organization. Our study landscape

features strong environmental gradients of salinity and turbidity that are expected to act as

strong selective forces for the local biota (Horváth et al. 2014). At the same time, the flat

lowland area experiences strong unidirectional winds, which is expected to result in a

directional spatial imprint on passively dispersing organisms (Epele et al. 2021; Kling and

Ackerly 2021). The latter was documented for the metacommunity structure of zooplankton in

an earlier study (Horváth et al. 2016). Soda pans therefore represent an attractive model for a

simultaneous analysis of landscape genetic and community variation along strong

environmental and connectivity gradients within a relatively small region.

Materials and Methods

Study area and sample collection

Soda pans are shallow, saline and temporary fishless waterbodies, which generally dry out in

summer. Within Europe, their occurrence is restricted to the Pannonian Plain; the lowlands of

Hungary, eastern Austria, and northern Serbia (**Fig. 1a**; (Boros et al. 2014). Soda pans usually

have a vegetation-free lakebed and no or only scarce emergent vegetation along the shorelines

(**Fig. 1c**). The pans are alkaline and their salinity exhibit a wide gradient from almost

freshwater to hypersaline conditions (Boros et al. 2014). They are generally well-mixed and

not stratified due to their exposure to wind and their high surface to depth ratio. Due to the

absence of fish and the hypertrophic conditions, zooplankton communities are bottom-up

regulated (Horváth et al. 2013, 2014). The predominant environmental gradient, salinity, is a

strong selection factor, and only a limited set of highly tolerant species account for the largest

part of zooplankton biomass (Horváth et al. 2014). This includes *Daphnia magna*, a widely-

used model species in landscape genetic studies, which is at the same time the most

widespread species in these habitats (Tóth et al. 2014).

According to a recent survey (Tóth et al. 2014), there are 23 well-preserved soda pans

in the Seewinkel region of Austria. We collected zooplankton samples from all 23 pans in

spring 2014 (28-29 April), by randomly collecting 20 l of water from the central part of each

habitat. Water was then sieved through a plankton net with a mesh size of 45 μm and

preserved in 70% ethanol on the field. In the lab, we applied subsampling to count the

abundance of Cladocera species in the samples; counting and identifying at least 300

individuals represents a subsample of 0.5-15%, depending on the total density in the sample.

Afterwards, the entire sample was checked for rare species. We also isolated either all or at

least 20 *D. magna* individuals, and preserved them in 96% ethanol until genotyping.

We determined the GPS location of the approximate centre of the pans, and measured

water depth (Z), conductivity, and pH in the field. Water samples were subsequently analysed

in the lab to quantify the concentrations of total suspended solids (TSS), total phosphorus

(TP) and chlorophyll *a* (Chl). TSS was measured gravimetrically by filtering water (1–25 ml,

depending on turbidity) through pre-dried (oven-drying at 105 °C) and pre-weighted glass

fibre filters (GF/F). TP was measured by using persulfate digestion (Clesceri et al. 1999),

followed by the ascorbic acid colorimetric method (Hansen et al. 1999). For Chl, we used

fluorometry with acetone extraction (Arar and Collins 1997), without correcting for

phaeophytin.

During our sampling campaign, 15 out of the 23 pans were found to host *D. magna*

populations (**Fig. 1b**, **Table S1** in Supporting information). We therefore decided to use only

these 15 sites in our comparative landscape genetic and metacommunity analysis. In our

landscape genetic analysis, we thus capture the complete metapopulation of this species

inhabiting soda pans.

Population genetic analysis: DNA extraction and microsatellite amplification

We screened for variation at 13 polymorphic neutral markers (see **Table S1** in Supporting

information; based on a set of 84 primers used by Orsini et al. 2012) in all 15 soda pan *D.*

magna populations. On average, 19 animals per population were genotyped (**Table S1**). We

genotyped 20 individuals from 11 pans; in the remaining four pans 21, 19, 16 and 10

individuals were genotyped. As no more animals were available in our samples, these latter

numbers could not be increased any further.

Genomic DNA was extracted from individual *Daphnia* using the HotShot method

(Truett et al. 2000; Montero-Pau et al. 2008). DNA amplifications were carried out in a final

reaction volume of 11.5 µl containing 1.5 µl of template genomic DNA and 10 µl of a

mixture of 0.2-0.3 µM of each primer (labelled with fluorescent dyes, 8 for multiplex M01

and 5 for M03) and 1x Qiagen Multiplex PCR Master Mix buffer (Qiagen®). PCR

amplification cycling conditions included an initial denaturation step at 95 °C for 15 minutes,

30 cycles of denaturation at 94 °C for 30 seconds, annealing at 54 (for multiplex M03) or 56

°C (for multiplex M01) for 1.5 minutes, extension at 72 °C for 1.5 minutes, and a final

elongation step at 60 °C for 30 minutes.

Amplified PCR products were separated on an ABI3130 capillary sequencer (Applied

Biosystems®). Allele sizes (fragment lengths) were determined by comparison with an

internal Liz500 size standard using the software GeneMapper (v3.7; Applied Biosystems®).

Data analysis

Population genetic structure

Standard measures of genetic variation, including percent of polymorphic loci, total (A) and

mean number of alleles per locus (allelic richness, A_r), expected heterozygosity (H_e),

observed heterozygosity (H_o), inbreeding coefficients (F_{IS}: individuals relative to the

subpopulation; F_{IT}: individuals relative to the total population) and fixation index (F_{ST},

quantifying subpopulation differentiation compared to the total population) were calculated in

R (R Development Core Team 2012) using packages "adegenet" (Jombart 2008), "polysat"

(Clark and Jasieniuk 2011), and "pegas" (Paradis 2010). A locus was considered polymorphic

if the most common allele had a frequency less than 0.99. Deviations from Hardy-Weinberg

equilibrium were tested with Monte Carlo permutation tests (Goudet's G-statistic with 1000

permutations) in "pegas" (Paradis 2010); these tests were carried out separately for all loci

and populations.

167   The extent of genetic differentiation among populations was assessed by comparing

pairwise F_{ST} values (Weir and Cockerham 1984) across all loci, using the package "hierfstat"

in R (Goudet 2005). A global F_{ST} value for the region across all loci and populations was

calculated with the package "polysat" (Clark and Jasieniuk 2011), and its significant

difference from zero was assessed using 1000 permutations.

Genetic and community matrix

For obtaining a data matrix representing genetic differentiation, we first built a PCoA on the

pairwise F_{ST} values (based on Bruvo distances) with the package "ape" for R (Paradis et al.

2004). In cyclically parthenogenetic organisms such as the water flea *D. magna*, clonal

structure might complicate genetic analyses, as multiple individuals might belong to the same

clone. In the soda pan populations, however, all sequenced individuals were characterized by

different multilocus genotypes, and this allowed us to use data on allelic frequencies in the

PCoA analysis. In a second step, we used the scores of all PCoA axes as the dependent matrix

in the subsequent variation partitioning analyses.

181   For the metacommunity data, we also ran a PCoA, this time based on Bray-Curtis

182   dissimilarities of the double square root transformed abundance data of the Cladocera

183   communities (after the exclusion of two singletons), and then extracted the scores of all PCoA

184   axes.

185   *Environmental matrix*

186 Among the environmental predictors, we log-transformed the non-normally distributed ones

187 (Z, TSS, Chl, TP). In soda pans, strong correlations between environmental variables are

188 frequent (Horváth et al. 2014, 2016); hence we first inspected pairwise Pearson's correlations

189 among our predictors (**Table S2** in the Supporting information). Based on that, we performed

190 an *a priori* data reduction by excluding TP from our initial set of explanatory variables

191 (because it was strongly correlated to almost all other predictors), and only retained Z, TSS,

192 Chl, and conductivity. These four predictors were then standardised to zero mean and unit

193 variance.

194 To select the most relevant environmental predictors, we applied an automatic

195 stepwise model selection from an initial RDA (Redundancy Analysis) model including the

196 four standardised environmental explanatory variables and the PCoA axes scores (either

197 resulting from the metapopulation or the metacommunity) as the dependent matrix. We

198 selected for significant predictors by using a permutation test ("ordistep" function in "vegan",

199 with direction="both" and 2000 permutations; Oksanen et al. 2018).

201 *Spatial matrix*

202 For quantifying spatial predictors, we employed spatial eigenvector methods in order to get a

203 more explicit representation of spatial patterns. Because of the strong prevalent wind direction

204 (Horváth et al. 2016), we applied both Moran's Eigenvector Maps (MEM; Dray et al. 2006),

205 as well as Asymmetric Eigenvector Maps (AEM; Blanchet et al. 2008), to identify patterns

206 both with (AEM) and without taking directionality into account (MEM). We used the same

207 connectivity matrix (binary neighbours list) for both sets of eigenvectors, built with a

208 threshold larger than the longest distance between any pair of pans (to enable all possible

209 connections between pans). For the AEM eigenvectors, the dominant wind direction was

added as the vector angle (292.5°) of the directional component (see also **Fig. S1** in

Supporting information). MEM and AEM eigenvectors were computed with the "spdep"

(Bivand et al. 2015) and "adespatial" packages (Dray et al. 2018). To select the most relevant

eigenvectors, we applied automatic stepwise model selection in a similar manner as for

environmental predictors.

Relative role of environmental and spatial predictors

To quantify the relative importance of environmental and spatial predictors, we carried out

variation partitioning analyses with the "vegan" package (Oksanen et al. 2018). Here, the

PCoA axes scores (either resulting from levels of genetic differentiation or community

dissimilarity) were used as the dependent matrix, together with a selected set of

environmental and spatial predictors.

In order to make results the most comparable across the metapopulation and the

metacommunity, we used a merged set of predictors, by including all predictors that proved

significant for at least one of the dependent matrices. For the metacommunity, the only

significant environmental predictor was TSS, while for the metapopulation, no significant or

marginally significant predictor was found. Hence we used TSS as the environmental

predictor for both datasets (in all four analyses). In the case of AEM eigenvectors, one

eigenvector (AEM 10) was significant for the metacommunity, and another (AEM 3) was

marginally significant for the metapopulation. Therefore, we included both in our variation

partitioning models, to use the exact same set of variables for both datasets. By doing so, we

can ensure that lower variation explained by either environment or space in our comparative

analysis does not arise from a difference in the number of predictors considered.

Partitioning analyses on single species and alleles

To explore variation in the spatial and environmental signal in the individual community

members and verify whether the degree of dispersal limitation as observed for individual

species might be linked to their occupancy in the metacommunity, we ran a series of variation

partitioning analyses on each community member (excluding singletons).

For the analyses at the level of individual species, we used the double square root

transformed community matrix (abundance data) after the exclusion of the two singletons. For

individual species, we applied automatic stepwise model selection on linear models. We

selected for significant predictors by using a stepwise model selection by AIC ("stepAIC"

function in "MASS", with direction="both" and 2000 permutations; Ripley et al. 2013).

Given that for the five species, different environmental predictors and eigenvectors proved

important (see **Table S3** and **S4** in Supporting information), we decided not to unify

predictors as carried out above for genetic and community data, and used the respective sets

of significant predictors for each species. The resulting values of pure environmental and

spatial effects (**Table S5** in Supporting information) were related to the regional occupancy of

species using linear models to test the idea that dispersal limitation is more apparent and

species sorting less effective in rare than in common species (Horváth et al. 2019). To ensure

that patterns are not due to any sampling effect (Leibold and Chase 2017; Viana and Chase

2019), we also tested the relationship between total explained variation and regional

occupancies.

To explore whether a similar pattern exists for landscape genetic structure of the

dominant community member, we also ran such analyses for genetic variation within the *D.*

magna metapopulation. We could not run such an analysis at the level of single genotypes

because all sequenced individuals in our study were characterized by different multilocus

genotypes. We thus decided to carry out these analyses at the level of alleles, checking

whether spatial signals are more important in less common compared to more widespread

alleles, quantified as the number of occurrences in the 15-site landscape. This linear

regression involved all alleles after the exclusion of singletons. We did not unify predictors,

because the high number of variation partitioning analyses (one for each of the 131 non-

singleton alleles) resulted in the full set of predictors being selected at least once. We report

the number of times a given environmental or spatial predictor was selected in the 131 models

in the Supporting information (**Fig. S2**).

Isolation-by-distance

To test whether genetic differentiation among populations increases with geographic distance,

we carried out a Mantel test on pairwise measures of genetic differentiation ($F_{ST}/(1 - F_{ST})$),

against pairwise geographic distances with 2000 permutations. Given the expectation that

dispersal rates might be higher in the direction aligned with the wind, we carried out two

Mantel tests, decomposing geographic distances into a component parallel to the predominant

wind direction and a component orthogonal to the predominant wind direction (see also **Fig.**

4b).

Results

The cladoceran metacommunity counted a total of seven species. Apart from two singletons

(*Alona rectangula* and *Megafenestra aurita*), the other species all occurred in at least three

habitats, with *D. magna* being the most frequent species among them (occurring in all 15

habitats). The other species were *Moina brachiata* (13), *Macrothrix hirsuticornis* (8),

Chydorus sphaericus (5), and *Daphnia atkinsoni* (3). Mean species richness (± SD) in the

habitats was 3.07 ± 1.22.

Across the 13 loci, a total of 158 alleles were detected (with a mean ± SD of 12.2 ± 4.2

alleles per locus and 5.7 ± 2.6 alleles per locus in a population) in *D. magna* from the soda

pans (**Table S1** in Supporting information). 27 alleles were singletons while 131 alleles

occurred in at least two habitats. The regional mean allelic richness was 6.08 ± 3.2 (range: 2-

22 alleles per locus in a population, **Table S1** in Supporting information). All loci were

polymorphic. Summaries of genetic diversity parameters are presented in **Table S1** (for each

habitat) and **Table S6** (for each locus) in the Supporting information, together with the

pairwise F_{ST} values of all habitat pairs (**Table S7**). According to Monte Carlo tests, the

observed level of heterozygosity was significantly lower than what we expect under Hardy-

Weinberg equilibrium both for the populations (**Table S1** in Supporting information) and for

the individual alleles (**Table S6** in Supporting information). Global genetic differentiation

among populations was significant (F_{ST}=0.12, t=211.56, df=999, p<0.001).

While symmetric MEMs did not reveal a significant spatial signal in the

metacommunity and the landscape genetic data (**Fig. 2** and **Table S8** in Supporting

information), the asymmetric AEM-based spatial components had a significant pure effect on

explaining variation in both community composition as well as genetic composition (**Fig. 2**

and **Fig. 3** and **Table S8** in Supporting information). The asymmetric spatial components

explained 10% of the variation in the Cladocera metacommunity ($F_{2,11}$=1.907, p=0.04) and

24% of the variation in landscape genetic data ($F_{2,11}$=2.936, p=0.01; **Fig. 2**). The amount of

variation in genetic identity explained by environment (total suspended solids) was much

lower (7%, $F_{1,11}$= 2.050, p=0.07) than both the amount explained by spatial predictors in the

same dataset (24%) and the amount explained in the metacommunity by the same

environmental gradient (25%, $F_{1,11}$= 5.265, p=0.001).

We found a significant relationship between the regional occurrence of species and the

pure explanatory power of local environment for cladoceran metacommunity structure (**Fig.**

3; R^2=0.80, $F_{1,3}$= 11.69, p=0.04), with a higher fraction of variation explained by

environmental gradients in more widespread species. An opposite trend was found for the

relationship between the pure effect of spatial predictors and regional occurrence (**Fig. 3**;

R^2=0.68, $F_{1,3}$= 6.458, p=0.085). No relationship was found between the total amount of

explained variation and regional occurrence (R^2=0.27, $F_{1,3}$= 1.089, p=0.373). In a parallel

analysis on alleles (**Fig. S3** in Supporting information), no effect of regional occurrence was

found on the amount of variation explained by space or environment. The amount of variation

explained by space (variable AEM components depending on the allele; **Fig. S2** in Supporting

information) was relatively high for all alleles, whereas environmental signal was overall very

low.

In the metapopulation, we found no clear isolation-by-distance patterns. Genetic

distances were not related to spatial distances, neither along nor orthogonal to the dominant

wind direction (Mantel's p = 0.23 for along the wind and p = 0.31 for orthogonal distances;

Fig. 4).

Discussion

Our comparative analysis highlights clear divergences and some parallels in the relative roles

of the main ecological processes in structuring the cladoceran zooplankton metacommunity

and the landscape genetic variation of one of its most widespread and abundant species within

the same landscape. It shows that while the processes structuring the assembly of

communities and populations are similar (Vellend 2010, 2016), they do result in different

patterns because of differences in the relative strengths of selection, dispersal or gene flow,

and drift in determining the assembly of communities and populations.

Whereas the structure of the metacommunity inhabiting the habitat network is

significantly influenced by environmental gradients (total suspended solids), this is not the

case for the landscape genetic structure. This may be linked to the fact that microsatellite loci

are presumably neutral, and a more detailed genomic analysis might reveal loci and SNPs that

do show an association with environmental gradients (e.g., predation, as in Chaturvedi et al.

2021). Yet, our data do indicate that natural selection in this system with a strong salinity

gradient is not so strong that it affects the assembly of populations so profoundly that it would

affect the structure of neutral genetic variation (as opposed to e.g., hypersaline lakes; Frisch et

al. 2021).

A key parallel in metacommunity and landscape genetic structure in the study system

is the significant signal of space, and the fact that the explanatory power of space for both

metacommunity and metapopulation is linked to the wind directionality captured in our AEM

based analysis. For both metacommunity and metapopulation structure, we indeed observed

that the amount of variation explained by space is significant in the AEM analysis while not

significant in the MEM-based analyses. This indicates that wind is likely an important

dispersal agent in this system, in line with the strong winds in the region (Horváth et al.

2016).

The large surface to depth ratio of the soda pans (Boros et al. 2014), their temporary

nature exposing sediments and dormant eggs to wind (Vanschoenwinkel et al. 2008; Pinceel

et al. 2015; Boros et al. 2017), the strong winds in the region (Horváth et al. 2016), and the

presence of large numbers of birds that can act as dispersal agents (Szabó et al. 2022) lead to

the expectation that there should be high dispersal rates among habitats in this system. Yet

there are some indications of dispersal limitation. First, a historical analysis of species

diversity decline driven by the disappearance of rare species with increasing loss of habitat

patches suggests non-equilibrium metacommunity structure for these species, reflecting

strong abundance-linked dispersal limitation (Horváth et al. 2019). Our data reveal a similar

pattern of dispersal limitation, given that the environmental signal increases and the spatial

signal declines with increasing abundance of species in the studied metacommunity. Variation

in the abundances of *D. magna*, being the most widespread species, is least explained by

spatial drivers of all species. This is in contrast, however, with the importance of the spatial

signal in explaining the landscape genetic structure of the species. Space is more important in

determining landscape genetic structure than in determining occurrences and abundances of

this species. This increase in spatial signal is not caused by the presence of many rare alleles

that would show a stronger signal of dispersal limitation than more widespread alleles, as

there was no association between the spatial signal and the relative frequency of alleles.

While there is a clear spatial signal for the landscape genetic structure, there is no

pattern of isolation-by-distance. Isolation-by-distance is a common expectation in population

genetics (Slatkin 1987; Bohonak 1999) predicted by theoretical models (Wright 1943; Hartl

and Clark 1997) as dispersal (gene flow) is expected to decline with increasing geographic

distances. The absence of a pattern of isolation-by-distance has nevertheless also been

observed in several other studies (Jenkins et al. 2010; Sexton et al. 2014) including *Daphnia*

(Innes 1991; Vanoverbeke and De Meester 1997) and suggests that the processes determining

landscape genetic structure deviate from standard expectations reflecting ongoing gene flow,

but may for instance also include a signature of patterns of past colonization (Orsini et al.

2013). The spatial signature reflected by the significant contribution of two AEM factors also

does not translate into a difference in the relationship between genetic differentiation and

geographic distance as measured parallel or orthogonal to the dominant wind direction. This

suggests that the impact of wind is differentiated and results in a rather patchy pattern, as it is

also reflected in the patterns that are tested for in the significant AEM factors, being AEM3

and AEM10. Wind directionality thus does influence landscape genetic structure (cf.

difference between MEM and AEM signals), but in a patchy way. This might reflect that the

wind effect is modulated by dispersal mediated by birds. The overall absence of a signal of

isolation-by-distance and the patchy nature of the wind effect may reflect that in the soda pans

in the Seewinkel region, wind does influence directionality of gene flow, but the strong winds

and birds flying from one soda pan to the other might enable equally strong connectivity

among pans that are further away from each other than among neighbouring pans. Birds can

indeed bridge large distances and do not necessarily move only between nearby patches

(Green and Figuerola 2005; Viana et al. 2013; Urgyán et al. 2022). The observed pattern

suggests that short- and long-distance dispersal are indeed equally important at the scale of

this landscape. At the same time, the level of genetic differentiation (global F_{ST} value of 0.12)

is moderate but significant, indicating that gene flow is not so high in this system as to

homogenize genetic variation.

Our comparative analysis of metapopulation genetic and metacommunity structure in

the same landscape of soda pans reveals some clear divergences (relative role of

environmental and spatial drivers) and some parallels (strong indications for a significant

contribution of wind directionality to the pattern) between assembly at these two different

levels of ecological organisation. We believe that such comparative analyses can contribute to

a better understanding of the patterns at each level. In addition, given that population

assembly determines evolutionary potential and that patterns of local adaptation may impact

community assembly and responses to environmental change (Urban et al. 2008; Hendry

2017; Leibold et al. 2022), the joint analysis of identity and trait variation at the level of

populations and communities is important to understand how populations and communities

respond to landscape heterogeneity and environmental change (Govaert et al. 2021, 2022).

Acknowledgements

We thank Adrienn Tóth, Emil Boros, Rudolf Schalli, and Richard Haider for assistance during

sample collection, Christian Preiler for laboratory measurements, and Bart Hellemans for

practical advice on population genetic analyses. Research was supported by the FWO project

G0C3818N, Projekt 06/14216 of the research funding program "International

Communication" of ÖFG (Austrian Research Association), the NKFIH project 2019-2.1.11-

TÉT-2020-00159, and KU Leuven Research Council project C16/2017/002. ZH

acknowledges support from the János Bolyai Research Scholarship of the Hungarian

Academy of Sciences and NKFIH FK project 132095.

References

Arar, E. J., and G. B. Collins. 1997. In-vitro determination of chlorophyll a and pheophytin a

in marine and freshwater algae by fluorescence: Cincinnati. Ohio, National Exposure

Research Laboratory, Office of Research and Development, US Environmental

Protection Agency, Method EPA **445**.

Bivand, R., M. Altman, L. Anselin, R. Assunção, O. Berke, A. Bernat, and G. Blanchet. 2015.

Package 'spdep.' The Comprehensive R Archive Network.

Blanchet, F. G., P. Legendre, and D. Borcard. 2008. Modelling directional spatial processes in

ecological data. Ecological Modelling **215**: 325–336.

Bohonak, A. J. 1999. Dispersal, gene flow, and population structure. The Quarterly Review of

Biology **74**: 21–45.

Bohonak, A. J. 2002. IBD (isolation by distance): a program for analyses of isolation by

distance. Journal of Heredity **93**: 153–154.

Borcard, D., P. Legendre, and P. Drapeau. 1992. Partialling out the Spatial Component of

Ecological Variation. Ecology **73**: 1045–1055. doi:10.2307/1940179

Boros, E., Z. Horváth, G. Wolfram, and L. Vörös. 2014. Salinity and ionic composition of the

shallow astatic soda pans in the Carpathian Basin. Annales de Limnologie -

International Journal of Limnology **50**: 59–69. doi:10.1051/limn/2013068

Boros, E., K. V.-Balogh, L. Vörös, and Z. Horváth. 2017. Multiple extreme abiotic conditions

and trophic state of intermittent soda pans in the Carpathian Basin (Central Europe).

Limnologica **62**: 38–46.

Chaturvedi, A., J. Zhou, J. A. M. Raeymaekers, and others. 2021. Extensive standing genetic

variation from a small number of founders enables rapid adaptation in Daphnia. Nat

Commun **12**: 4306. doi:10.1038/s41467-021-24581-z

Clark, L. V., and M. Jasieniuk. 2011. POLYSAT: an R package for polyploid microsatellite

analysis. Mol Ecol Resour **11**: 562–566. doi:10.1111/j.1755-0998.2011.02985.x

Clesceri, L. S., A. E. Greenberg, and A. D. Eaton. 1999. Standard methods for examination of

water & wastewater.

Dray, S., G. Blanchet, D. Borcard, and others. 2018. Package 'adespatial,.'

Dray, S., P. Legendre, and P. R. Peres-Neto. 2006. Spatial modelling: a comprehensive

framework for principal coordinate analysis of neighbour matrices (PCNM).

Ecological Modelling **196**: 483–493.

Epele, L. B., D. A. Dos Santos, R. Sarremejane, and others. 2021. Blowin' in the wind: Wind

directionality affects wetland invertebrate metacommunities in Patagonia. Global

Ecology and Biogeography **30**: 1191–1203. doi:10.1111/geb.13294

Frantz, A. C., S. Bertouille, M. C. Eloy, A. Licoppe, F. Chaumont, and M. C. Flamand. 2012.

Comparative landscape genetic analyses show a Belgian motorway to be a gene flow

barrier for red deer (Cervus elaphus), but not wild boars (Sus scrofa). Molecular

Ecology **21**: 3445–3457. doi:10.1111/j.1365-294X.2012.05623.x

Frisch, D., C. Lejeusne, M. Hayashi, M. T. Bidwell, J. Sánchez-Fontenla, and A. J. Green.

2021. Brine chemistry matters: Isolation by environment and by distance explain

population genetic structure of Artemia franciscana in saline lakes. Freshwater

Biology **66**: 1546–1559. doi:10.1111/fwb.13737

Goldberg, C. S., and L. P. Waits. 2010. Comparative landscape genetics of two pond-breeding

amphibian species in a highly modified agricultural landscape. Molecular ecology **19**:

3650–3663.

Goudet, J. 2005. Hierfstat, a package for R to compute and test hierarchical F-statistics.

Molecular Ecology Notes **5**: 184–186.

Govaert, L., L. De Meester, S. Rousseaux, S. A. Declerck, and J. H. Pantel. 2021. Measuring

the contribution of evolution to community trait structure in freshwater zooplankton.

Oikos **130**: 1773–1787.

Govaert, L., J. H. Pantel, and L. De Meester. 2022. Quantifying eco-evolutionary

contributions to trait divergence in spatially structured systems. Ecological

Monographs **92**: e1531. doi:10.1002/ecm.1531

Green, A. J., and J. Figuerola. 2005. Recent advances in the study of long-distance dispersal

of aquatic invertebrates via birds. Diversity and Distributions **11**: 149–156.

Guillot, G., A. Estoup, F. Mortier, and J. F. Cosson. 2005. A spatial statistical model for

landscape genetics. Genetics **170**: 1261–1280.

Hansen, H. P., F. Koroleff, K. Grasshoff, K. Kremling, and M. Ehrhardt. 1999. Methods of

seawater analysis. Grasshoff, K., Kremling, K., Ehrhardt, M. Wiley-VCH, Germany.

Hartl, D. L., and A. G. Clark. 1997. Principles of population genetics, Sinauer associates

Sunderland.

Hein, C., H. E. Abdel Moniem, and H. H. Wagner. 2021. Can We Compare Effect Size of

Spatial Genetic Structure Between Studies and Species Using Moran Eigenvector

Maps? Frontiers in Ecology and Evolution **9**.

Hendry, A. P. 2017. Eco-evolutionary Dynamics, Princeton University Press.

Horváth, Z., R. Ptacnik, C. F. Vad, and J. M. Chase. 2019. Habitat loss over six decades

accelerates regional and local biodiversity loss via changing landscape connectance.

Ecology Letters **22**: 1019–1027.

Horváth, Z., C. F. Vad, and R. Ptacnik. 2016. Wind dispersal results in a gradient of dispersal

limitation and environmental match among discrete aquatic habitats. Ecography **39**:

726–732.

Horváth, Z., C. F. Vad, A. Tóth, K. Zsuga, E. Boros, L. Vörös, and R. Ptacnik. 2014.

Opposing patterns of zooplankton diversity and functioning along a natural stress

gradient: When the going gets tough, the tough get going. Oikos **123**: 461–471.

doi:10.1111/j.1600-0706.2013.00575.x

Horváth, Z., C. F. Vad, L. Vörös, and E. Boros. 2013. The keystone role of anostracans and

copepods in European soda pans during the spring migration of waterbirds. Freshwater

Biology **58**: 430–440. doi:10.1111/fwb.12071

Innes, D. J. 1991. Geographic patterns of genetic differentiation among sexual populations of

Daphnia pulex. Canadian Journal of Zoology **69**: 995–1003.

Jenkins, D. G., M. Carey, J. Czerniewska, and others. 2010. A meta-analysis of isolation by

distance: relic or reference standard for landscape genetics? Ecography **33**: 315–320.

doi:10.1111/j.1600-0587.2010.06285.x

Jombart, T. 2008. adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics **24**: 1403–1405.

Kling, M. M., and D. D. Ackerly. 2021. Global wind patterns shape genetic differentiation,

asymmetric gene flow, and genetic diversity in trees. Proceedings of the National

Academy of Sciences **118**: e2017317118.

Leibold, M. A., and J. M. Chase. 2017. Metacommunity ecology, Princeton University Press.

Leibold, M. A., L. Govaert, N. Loeuille, L. De Meester, and M. C. Urban. 2022. Evolution

and community assembly across spatial scales. Annual Review of Ecology, Evolution,

and Systematics **53**: 299–326.

Leibold, M. A., M. Holyoak, N. Mouquet, and others. 2004. The metacommunity concept: a

framework for multi-scale community ecology. Ecology Letters **7**: 601–613.

doi:10.1111/j.1461-0248.2004.00608.x

Manel, S., and R. Holderegger. 2013. Ten years of landscape genetics. Trends in Ecology &

Evolution **28**: 614–621.

Manel, S., M. K. Schwartz, G. Luikart, and P. Taberlet. 2003. Landscape genetics: combining

landscape ecology and population genetics. Trends in Ecology & Evolution **18**: 189–

197.

Manier, M. K., and S. J. Arnold. 2006. Ecological correlates of population genetic structure: a

comparative approach using a vertebrate metacommunity. Proceedings of the Royal

Society of London B: Biological Sciences **273**: 3001–3009.

Moncada, B., J. A. Mercado-Díaz, N. Magain, and others. 2021. Phylogenetic diversity of two

geographically overlapping lichens: isolation by distance, environment, or

fragmentation? Journal of Biogeography **48**: 676–689. doi:10.1111/jbi.14033

Montero-Pau, J., A. Gómez, and J. Muñoz. 2008. Application of an inexpensive and high-

throughput genomic DNA extraction method for the molecular ecology of

zooplanktonic diapausing eggs. Limnology and Oceanography: Methods **6**: 218–222.

Nosil, P., T. H. Vines, and D. J. Funk. 2005. Reproductive isolation caused by natural

selection against immigrants from divergent habitats. Evolution **59**: 705–719.

Oksanen, J., F. G. Blanchet, M. Friendly, and others. 2018. vegan: Community Ecology

Package. R package version 2.5-3.

Orsini, L., K. I. Spanier, and L. De Meester. 2012. Genomic signature of natural and

anthropogenic stress in wild populations of the waterflea Daphnia magna: validation in

space, time and experimental evolution. Molecular Ecology **21**: 2160–2175.

Orsini, L., J. Vanoverbeke, I. Swillen, J. Mergeay, and L. De Meester. 2013. Drivers of

population genetic differentiation in the wild: isolation by dispersal limitation,

isolation by adaptation and isolation by colonization. Molecular ecology **22**: 5983–

5999.

Paradis, E. 2010. pegas: an R package for population genetics with an integrated–modular

approach. Bioinformatics **26**: 419–420.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: analyses of phylogenetics and evolution

in R language. Bioinformatics **20**: 289–290.

Peres-Neto, P. R., P. Legendre, S. Dray, and D. Borcard. 2006. Variation partitioning of

species data matrices: estimation and comparison of fractions. Ecology **87**: 2614–

2625.

Pinceel, T., L. Brendonck, and B. Vanschoenwinkel. 2015. Propagule size and shape may

promote local wind dispersal in freshwater zooplankton—a wind tunnel experiment.

Limnology and Oceanography.

R Development Core Team. 2012. R: A Language and Environment for Statistical

Computing.

Richardson, J. L. 2012. Divergent landscape effects on population connectivity in two co-

occurring amphibian species. Molecular Ecology **21**: 4437–4451.

Ripley, B., B. Venables, D. M. Bates, K. Hornik, A. Gebhardt, D. Firth, and M. B. Ripley.

2013. Package 'MASS.' Cran R **538**.

Sexton, J. P., S. B. Hangartner, and A. A. Hoffmann. 2014. Genetic isolation by environment

or distance: which pattern of gene flow is most common? Evolution **68**: 1–15.

Slatkin, M. 1987. Gene flow and the geographic structure of natural populations. Science **236**:

787–792.

Szabó, B., A. Szabó, C. F. Vad, E. Boros, D. Lukić, R. Ptacnik, Z. Márton, and Z. Horváth.

2022. Microbial stowaways: Waterbirds as dispersal vectors of aquatic pro-and

microeukaryotic communities. Journal of Biogeography **49**: 1286–1298.

doi:10.1111/jbi.14381

Tóth, A., Z. Horváth, C. F. Vad, K. Zsuga, S. A. Nagy, and E. Boros. 2014. Zooplankton of

the European soda pans: fauna, communities and conservation of a unique habitat

type. International Review of Hydrobiology **99**: 255–276.

doi:10.1002/iroh.201301646

Truett, G. E., P. Heeger, R. L. Mynatt, A. A. Truett, J. A. Walker, and M. L. Warman. 2000.

Preparation of PCR-quality mouse genomic DNA with hot sodium hydroxide and tris

(HotSHOT). Biotechniques **29**: 52–54.

Urban, M. C., M. A. Leibold, P. Amarasekare, and others. 2008. The evolutionary ecology of

metacommunities. Trends in Ecology & Evolution **23**: 311–317.

Urgyán, R., B. A. Lukács, R. Fekete, A. Molnár V., A. Nagy, O. Vincze, A. J. Green, and Á.

Lovas-Kiss. 2022. Plants dispersed by a non-frugivorous migrant change throughout

the annual cycle. Global Ecology and Biogeography **Early View**.

doi:10.1111/geb.13608

Vanoverbeke, J., and L. De Meester. 1997. Among-populational genetic differentiation in the

cyclical parthenogen Daphnia magna (Crustacea, Anomopoda) and its relation to

geographic distance and clonal diversity. Hydrobiologia **360**: 135–142.

doi:10.1023/A:1003160903708

Vanschoenwinkel, B., S. Gielen, M. Seaman, and L. Brendonck. 2008. Any way the wind

blows - frequent wind dispersal drives species sorting in ephemeral aquatic

communities. Oikos **117**: 125–134. doi:10.1111/j.2007.0030-1299.16349.x

Vellend, M. 2010. Conceptual synthesis in community ecology. The Quarterly review of

biology **85**: 183–206.

Vellend, M. 2016. The theory of ecological communities, Princeton University Press.

Vellend, M., and M. A. Geber. 2005. Connections between species diversity and genetic

diversity. Ecology Letters **8**: 767–781. doi:10.1111/j.1461-0248.2005.00775.x

Viana, D. S., and J. M. Chase. 2019. Spatial scale modulates the inference of metacommunity

assembly processes. Ecology **100**: e02576. doi:10.1002/ecy.2576

Viana, D. S., L. Santamaría, T. C. Michot, and J. Figuerola. 2013. Migratory strategies of

waterbirds shape the continental-scale dispersal of aquatic organisms. Ecography **36**:

430–438. doi:10.1111/j.1600-0587.2012.07588.x

Wang, I. J., and G. S. Bradburd. 2014. Isolation by environment. Mol Ecol **23**: 5649–5662.

doi:10.1111/mec.12938

Weir, B. S., and C. C. Cockerham. 1984. Estimating F-statistics for the analysis of population

structure. Evolution **38**: 1358–1370.

Wright, S. 1943. Isolation by distance. Genetics **28**: 114.

 Figure captions

 Figure 1 – (a) Map showing the Pannonian Plain as an enclosed lowland between the Alps

and the Carpathian Mountains in East-Central Europe, where dark green shading indicates the

Pannonian biogeographic region (white circles: soda pans). The orange frame on the border

between Hungary and Austria highlights the study region (Seewinkel). (b) The studied 15

soda pans in the Seewinkel region (white circles). (c) An example of the study habitats during

the survey.

Figure 2 – Variation partitioning results obtained for the Cladocera metacommunity and the

Daphnia magna metapopulation. Environmental effect only included the concentration of

total suspended solids (TSS) in all cases. Spatial predictors were selected from eigenvectors

of Moran's Eigenvector Maps (MEM) for symmetric and Asymmetric Eigenvector Maps

(AEM) for wind-related patterns. Bold letters indicate significant pure effects of environment

or space ($p < 0.05$). Results of the F-statistics are given in Table S8.

Figure 3 – Left panel: Pure environmental and spatial component of explained variation in

the abundance of each member of the Cladocera metacommunity ("Dap_atk" – *Daphnia*

atkinsoni, "Chy_sph" – *Chydorus sphaericus*, "Mac_hir" – *Macrothrix hirsuticornis*,

"Moi_bra" – *Moina brachiata*, "Dap_mag" – *Daphnia magna*), plotted against their regional

occurrence. Spatial predictors were selected from eigenvectors of Asymmetric Eigenvector

Maps (AEM). Right panel: Pure environmental and spatial component of explained variation

calculated for the metacommunity of Cladocera and for the genetic variation in the *D. magna*

metapopulation (presented in detail in Figure 2).

Figure 4 – (a) Eigenvectors of Asymmetric Eigenvector Maps (AEM) retained in the

analyses, with their eigenfunction values represented by the size of squares at the spatial

position of each habitat. (b) Calculation of spatial distances along and orthogonal to the

616   dominant wind direction (bold lines illustrate a specific example in each case; above) and

617   their relationship with pairwise F_{ST} values (below).

618   **Figures**

619
620   **Figure 1**

Figure 2

625 **Figure 3**

Figure 4