Evidence for a functional interaction between the respiratory syncytial virus fusion and attachment proteins in the envelope of infectious virus particles.

Tra Nguyen Huong[1], Boon Huan Tan[2,3] and Richard J. Sugrue[1]*

1. School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive, Singapore 637551, Republic of Singapore.

2. Biological Defence Program, DSO National Laboratories, 27 Medical Drive, Singapore 117510, Republic of Singapore.

3. LKC School of Medicine, Nanyang Technological University, 11 Mandalay Road, Singapore 308232, Republic of Singapore.

* Correspondence: rjsugrue@ntu.edu.sg

Abstract

We have examined the interaction between the respiratory syncytial virus (RSV) F and G proteins on the surface of infected cells during multiple cycle infection using a low multiplicity of infection (MOI) model, and on the surface of virus particles that were isolated from infected cells. A combination of the proximity ligation assay (PLA) and confocal microscopy was used to demonstrate the interaction between the F and G proteins within the virus filaments on infected cells. Co-precipitation of the F and G proteins was confirmed using detergent extracts prepared from infected cells and in detergent extracts prepared from purified virus particles. The influence of the G protein in mediating virus spread in the low MOI model was further examined using the recombinant virus isolates rg224RSV (that expresses all virus proteins) and rg224RSV-ΔG (which does not express the G protein). While cells could be initially infected by both viruses, the rg224RSV-ΔG virus exhibited severely impaired localised virus transmission in the multiple cycle infection assay. Collectively these data provide evidence that the F and G proteins interact within the envelope of RSV particles, and suggests that this interaction may promote virus transmission. The interaction between these proteins in a single protein complex represents a potential new target for the development of antivirus strategies and in the development of RSV vaccine candidates.

Introduction.

The mature Respiratory Syncytial Virus (RSV) particle assembles at the apical cell surface as cell-associated filamentous structures [1], and during RSV infection it is the virus filaments that mediate virus transmission [2, 3]. The virus filaments are bounded by a virus envelope in which the G protein and virus fusion (F) protein are inserted. The G protein is the RSV cell-attachment protein [4] and it undergoes extensive glycosylation to form the mature 90 kDa G protein [5]. The F protein initiates fusion of the virus envelope and cell membrane during virus entry into the cell. It is initially expressed as a single polypeptide chain (F0), and it is subsequently cleaved by furin into the F1 (50 kDa) and F2 (20 kDa) subunits [6-8]. Several models have been put forward to explain the mechanism of RSV-induced membrane fusion [9]. It is suggested that the mature F protein initially exists as a labile pre-fusion state, which is subsequently converted into a stable post-fusion molecule during membrane fusion [10, 11]. While the structural basis for the fusion process mediated by the RSV F protein is understood [9, 12], the cue that converts the labile pre-fusion state into the stable post-fusion form during virus infection is not established. In many paramyxoviruses the interaction of the attachment protein with the cell receptor mediates structural changes in the F protein that are required during membrane fusion [13]. In this context we have previously demonstrated that a protein complex involving the F and G protein forms on the surface of infected cells [14]. This earlier study relied on the co-immunoprecipitation of detergent extracts obtained from virus-infected cells. However, this earlier study did not specifically address if this protein complex was incorporated into the envelope of virus filaments (i.e. infectious virus particles). We have therefore extended our earlier findings to determine if this protein complex was formed in the envelope of infectious virus particles. The presence of this complex in virus particles would have important implications towards understanding how the process of F

protein mediated cell entry is regulated. It could also potentially lead to both novel antivirus

strategies that are based on destabilising this interaction and in the development of

recombinant RSV subunit vaccines.

Materials and Methods.

Virus and cell preparation. The rg224RSV-FSG and rg224RSV-FS viruses have been

described previously [15, 16]. The RSV A2, rg224RSV-FSG and rg224RSV-FS viruses were

prepared in HEp-2 cells as described previously [17, 18]. Infections were performed in HEp-

2 cells at the desired multiplicity of infection in DMEM with 2% FCS in a humidified

chamber at 33 ºC with 5% CO_2. The HEp2 cells were maintained in DMEM with 10% FCS

incubated at 37 ºC with 5% CO_2.

RSV multiple cycle virus infection assay. The cell-associated and cell-free (released) virus

infectivity was measured as described previously [19]. Briefly, HEp2 cells that were 95%

confluent were infected with RSV using a multiplicity of infection of between 0.001 and

0.0001 as indicated. At specific time intervals the cell-free virus infectivity was recovered by

harvesting the tissue culture supernatant. The cell-associated virus was recovered by

scrapping the cells into the same volume of pre-warmed DMEM (2% FCS), and the virus

infectivity released from the cells by two cycles of rapid freeze thaw. Contaminating cellular

material was removed from each virus preparation by centrifugation at 5000g for 10 minutes

at 4ºC. The recovered viral infectivity in each fraction was assessed on HEp-2 cells by

microplaque assay.

Microplaque titration of RSV infectivity. The recovered viral infectivity was assessed on

HEp-2 cells using a microplaque assay as described previously [20] but with minor

modifications. Briefly, at 48 hrs post-infection (hpi) the HEp-2 cell monolayers were

incubated with an anti-RSV (Novacastra) and anti-mouse IgG conjugated to Alexa 488. The

stained microplaques were counted using a Nikon Eclipse 80i (Japan) fluorescent microscope.

Antibodies and specific reagents. The anti-mouse and anti-rabbit IgG conjugated to Alexa

488 and Alexa555 (Molecular Probes), anti-NP (Chemicon, San Diago, CA, USA) and Alexa

Fluor™ 488 Phalloidin (ThermoFisher) were purchased. The RSV M2-1 antibody has been

described previously [21-23], and the F protein polyclonal and anti-G were gifts from Jose

Melero (Madrid, Spain) and Geraldine Taylor (IAH, UK) respectively.

Proximity Ligation assay. The cells were fixed using 4% PFA in PBS and then incubated

with anti-G and anti-F antibodies (either in combination or singularly as required). The

proximity Ligation Assay (PLA) was then performed using the Duolink® In Situ Detection

Red as recommended by the manufacturer (Sigma Aldrich). After the PLA reaction was

performed, the cells were then incubated with the anti-G or anti-F antibodies and the anti-

mouse or anti-rabbit IgG conjugated to Alexa 488 as appropriate. Cells were washed with

PBS and mounted on glass slide using mounting media (Dakocytomation Fluorescence

Mounting Media, Dako, USA). The cells were examined using either a Nikon Eclipse 80i

(Japan) fluorescent microscope or by using a Zeiss 710 confocal microscope with Zeiss

software.

Immunofluorescence microscopy. This was performed as described previously [24].

Briefly, the cells were fixed using 4% (w/v) paraformaldehyde in PBS at $4^{o}C$ for 20 mins and

washed using PBS at $4^{o}C$. For Alexa Fluor™ 488 Phalloidin staining the cells were fixed

using 3% (w/v) paraformaldehyde in PBS. The cells were permeabilised using 0.1% (v/v)

triton X100 in PBS at $4^{o}C$ for 15 mins. The cells were stained using the primary and

secondary antibody combinations (Alexa 488 or Alexa 555 as appropriate). Alexa Fluor™

488 Phalloidin staining was performed using the manufacturer's instructions. The cells were

examined using either a Nikon Eclipse 80i (Japan) fluorescent microscope or by using a Zeiss

710 confocal microscope with Zeiss software as appropriate.

Immunoblotting analysis. The cells were washed twice using sterile PBS (at 4^{o}C) and then

extracted directly into Boiling Mix (1%(w/v) SDS, 5% mercaptoethanol (v/v) in 20mM

Tris/HCL, pH 7.5). The extracted cells were immediately heated at 100^{o}C for 2 min and then

clarified by centrifugation (13,000 x g for 2 min). The proteins were separated by SDS-

PAGE and transferred by Western blotting onto nitrocellulose membranes. Protein bands

were then probed with the relevant primary antibody and corresponding secondary antibody

conjugated to HRP, and visualized using the ECL detection system (GE Healthcare).

Surface protein biotinylation. This was performed with 0.5 mg/ml EZ-Link Sulfo-NHS-

LC-LCBiotin (Pierce Biotechnology) solution in PBS pH 8 as described previously [14]. The

cell monolayers were treated with 0.5 mg/ml EZ-Link Sulfo-NHS-LC-LCBiotin (Pierce

Biotechnology) solution in PBS pH 8 for 1 h at 20^{o}C. The cells were washed using PBS with

2mM lysine and then the cells were lysed using ice-cold extraction buffer (PBS pH 8, 1%

Nonidet P-40 (BDH), 1 mM EDTA, 2mM lysine and 2 mM PMSF) at 4°C for 20 min and

clarified by centrifugation (13,000g, 10min) at 4^{o}C. The biotinylated proteins were isolated

by immunoprecipitation and transferred onto PVDF membranes by western blotting and

detected using streptavidin-HRP. Protein bands were quantified by using ImageJ (ver IJ1.46r)

to analyse the protein bands on autoradiographs. Protein profiles were obtained or protein

bands to be quantified were delineated, and the relative intensities determined and compared

with the background intensity in control lanes.

Biotinylation of RSV particles. HEp2 cells were infected with RSV using a multiplicity of

infection of 0.05 and at 60 hpi the infectious virus particles were isolated using

ultracentrifugation as described previously [18]. The virus particles were suspended in PBS

pH 8 with 0.5 mg/ml EZ-Link Sulfo-NHS-LC-LCBiotin for 1 h at 20^{o}C. The reaction was

quenched by adding 20mM lysine and after 10 mins the virus particles recovered by

centrifugation (284,000 × g) for 90 mins at 4°C. The virus particles were suspended in Hanks

buffered saline solution HBSS with 20mM lysine, and extracted using an equal volume of 2%

(v/v) Nonidet P-40 in HBSS with 20mM lysine at 4°C for 20 mins. The detergent extract was

clarified by centrifugation (13,000g, 10min) at 4^{o}C and the proteins isolated by

immunoprecipitation [14].

Results and discussion.

We have previously described a low multiplicity of infection (MOI) model for

examining the multiple cycle infection in RSV-infected cell monolayers [19, 25]. Under these

conditions localised virus transmission in cell monolayers was observed when using a moi of

between 0.001 and 0.0001. This phenomenon is a robust feature of all permissive cell lines

infected with RSV. High mutation rates that are associated with RNA viruses can lead to the

formation of defective interfering (DI) particles that are potentially non-infectious or that

exhibit impaired infectivity [26]. The presence of DI particles in virus preparations have also

been shown to significantly influence the virus transmission processes in permissive cells

[27-29], and unless otherwise specified, in this current study the HEp2 cells were infected

with RSV using a MOI of 0.0001. Under these conditions the DI particles that would be

expected to be present in the initial virus inoculum would be sufficiently diluted out and

individual cells in the monolayer would only be infected with individual and fully infectious

virus particle. This allowed both RSV transmission in the cell monolayer and the interaction

between the F and G proteins to be examined with the minimal involvement of the DI

particles. HEp2 cells were infected with RSV and at between 1 and 3 days-post-infection

(dpi) the cell monolayers were co-stained using anti-G and Alexa Fluor™ 488 Phalloidin and

imaged using immunofluorescence (IF) microscopy (Fig. 1A).While anti-G staining allowed

infected cells to be visualised, the Alexa Fluor™ 488 Phalloidin stained the F-actin and

confirmed the integrity of the cell monolayer. At 1 dpi mainly singly infected cells were

detected within the cell monolayer, while at 2dpi larger infected cell clusters that consisted of

25±15 cells per cluster was detected, and the anti-G staining was restricted to these infected

cell clusters. At 3 dpi enlarged cell clusters were detected, and their appearance correlated

with the start of syncytia formation. At each time of infection the cell-associated (CA) and

cell-free (CF) virus infectivity was separated and quantified by plaque assay (Fig. 1B). At

between 1 and 3 dpi a continuous increase in the CA-virus infectivity was detected, while the

appearance of CF-virus infectivity was only detected by 3 dpi. This pattern of virus spread

confirmed our previous observations demonstrating the localised cell-to-cell transmission of

RSV in cell monolayers [19].

At 2 dpi the HEp2 cells were also co-stained using anti-F and anti-G and the infected

cell clusters examined by IF microscopy (Fig. 2 A-C), which showed strong co-staining

within the virus filaments. Examination of the co-staining pattern in more detail indicated that

while anti-G staining was observed along the whole length of the virus filaments, increased

anti-F staining at the distal end of the virus filaments suggested the increased concentration of

the F protein at the ends of the virus filaments (Fig. 2 B and C). These imaging data suggest a

compartmentalisation of the F protein distribution in the virus filaments, and this is similar to

the observation in RSV VLPs that are formed by the co-expression of the F and G proteins

[30]. At 2 dpi the HEp2 cells were also examined by confocal microscopy (Fig. 3A), which

also indicated strong anti-G and anti-F co-staining within the virus filaments at the distal ends

of the virus filaments. Examination of the pixel distribution at the periphery (Fig. 3B (i) and

(ii))) and top (Fig. 3C (i) and (ii)) of the infected cell clusters confirmed high levels of co-

localisation between the anti-F and anti-G antibody staining within virus filaments (Pearson

coefficient = 0.78±0.04).

The high level of co-localisation was suggestive of an interaction *in situ*, and this

possibility was examined further using the proximity ligation assay (PLA), which is an

immuno-PCR based-assay that allows detection of interacting proteins *in situ* [31]. Although

this assay cannot quantitate the number of interactions involving specific proteins, it is able to

provide a qualitative indication of interacting proteins in intact cells. In the PLA the cells

were incubated with anti-F and anti-G antibodies, followed by secondary antibodies that are

conjugated to complementary DNA sequences that form a PCR primer pair. When the

antibodies are in close proximity, the reaction products appear as distinct bright red spots

when viewed by IF microscopy. HEp2 cells were either mock-infected or infected with RSV,

and at 2 dpi the cells were incubated with anti-G and anti-F antibodies and the PLA reaction

performed. The cells were subsequently labelled with anti-mouse IgG-Alexa 488 to detect the

anti-G staining on the infected cell monolayer, which allowed us to examine the distribution

of the virus filaments on the infected cells in the context of the PLA signal. The PLA signal

was absent on the mock-infected cells (Fig. 4A), and on the infected cell monolayer, the PLA

signal was restricted to the anti-G stained infected cell clusters (Fig. 4B (i) and (ii)). The PLA

signal was absent on the negative control assays in which the PLA performed on infected

cells that were incubated using either only anti-G or anti-F antibody (Fig. 4C (i) and (ii)).

This demonstrated that the positive PLA reaction was not due to antibody cross reactivity.

Collectively, these data indicated a positive PLA reaction in the virus-infected cells, and

provided direct evidence for an interaction between the F and G proteins in the low MOI

multiple cycle infection model.

220   The co-stained infected cells were examined in greater detail using confocal

microscopy to determine the distribution of the PLA signal in the context of the anti-F stained

virus filaments at the cell periphery (Fig. 5A(i)-(iii)). High levels of PLA staining on the anti-

F stained virus filaments was visible and examined the pixel distribution of the PLA signal

and anti-F staining (Fig. 5B(i) and (ii)) at regions of the images where the PLA signal was

present. This revealed high levels of co-localisation between the PLA signal and the anti-G

stained virus filaments (Pearson's correlation coefficient = 0.68±0.05). Similarly, analysis of

the antibody staining intensity profile across the virus filaments (Fig. 5C(i) and (ii)) where

the PLA signal was present showed good correlation between the PLA signal and anti-F

staining intensities. These analyses confirmed the presence of the PLA signal on the anti-F

stained virus filaments, which accounted for greater than 95% of the total PLA signal

detected indicating close association of the F and G proteins in these structures.

232   We examined the interaction of the F and G proteins under low moi conditions using

an established cell surface biotinylation assay that has been described previously [14]. Cells

were either mock-infected or with RSV infected using a moi of 0.001, and at 2 dpi the cells

were surface biotinylated and cell lysates prepared. Analysis of the total biotinylated proteins

in the mock-infected and RSV-infected cell lysates revealed biotinylated proteins up to 200

kDa in size (Fig. 6A(i)). The cell lysates were immunoprecipitated using anti-F antibody

which revealed the appearance of the F1 protein and a 90 kDa protein corresponding in size

to the G protein in the RSV-infected cells lysates. Similarly the appearance of a 90 kDa G

protein and lower levels of the 50 kDa F1 protein were observed by immunoprecipitation of

the RSV-infected cell lysates with anti-G. These biotinylated protein species were not

detected in lysates prepared from mock-infected cells, nor were they detected following

immunoprecipitation with anti-NP (an antibody that recognises the influenza virus

nucleoprotein). The ratio of the G to F1 protein bands was approximately 10:1 and 1:1 in the immunoprecipitation assay using anti-G and anti-F antibodies respectively (Fig. 6A(ii)).

The interaction between the F and G proteins was also examined in virus particles that were isolated from HEp2 cells by ultracentrifugation as described previously [14]. Immunoblotting of the virus preparation with anti-F and anti-G showed proteins of the expected size for the 50 kDa F1 protein and the 90 kDa G protein respectively (Fig. 6B(i)). The virus preparation was surface-biotinylated and the analysis of the total-labelled proteins in the detergent virus extract revealed a prominent protein band at approximately 90 kDa, and lower levels of biotinylated protein species up to 250 kDa were also observed (Fig. 6B(ii)). Immunoprecipitation with anti-F showed the presence of the 50 kDa F1 protein together with a 90 kDa protein whose size was consistent with the G protein, while immunoprecipitation with anti-G showed a prominent 90 kDa G protein species, together with a minor protein band corresponding in size to the F1 subunit. (Fig. 6B(iii)). The absence of biotinylated protein species following immunoprecipitation with anti-NP (non-specific antibody) demonstrated the specificity of the immunoprecipitation assay. The ratio of the G to F1 protein bands was approximately 10:1 and 1:1 in the immunoprecipitation assay using anti-G and anti-F antibodies respectively (6B(iv)) and was similar to that observed in the biotinylation analysis on virus-infected cells described above.

A single protein complex involving the F and G proteins within the envelope of infectious RSV particles provided evidence for a functional linkage between the activity associated with the G protein (cell attachment) and the F protein (membrane fusion). The surface expressed virus glycoproteins were detected by biotinylation of surface accessible lysine residues and the F and G protein may be labelled to varying degrees (e.g. due to differences in the abundance and surface exposure of the lysine residues). It was therefore not

possible to determine the stoichiometry of the F and G proteins in this protein complex.

However, this analysis revealed that the relative amounts of the F and G proteins co-

immunoprecipitated using either anti-F or anti-G differed. While an approximate ratio of 1:1

was noted by immunoprecipitation with anti-F, a lower level of the F1 protein was co-

precipitated by immunoprecipitation with anti-G. The difference in the ratio in the band

intensity observed with the different antibodies suggested that while the F and G protein

interact on the surface of cells, not all the surface expressed G protein interacts with the F

protein. These differences suggested that the F protein may interact with only a subpopulation

of the total G protein that is present in the virus envelope. This conclusion is consistent with

the imaging analysis which indicated increased levels of the F protein at the distal ends of the

virus filaments, and that the co-localisation between anti-F and anti-G staining was mainly

observed at the distal ends of the anti-G stained virus filaments.

In a final analysis we examined if the absence of the G protein impaired RSV

transmission in the low MOI infection model. This was performed by comparing the

transmission of wild-type RSV virus (rg224RSV) that expresses all three virus glycoproteins

with a virus that does not express the G protein (rg224RSV-ΔG). Both viruses were

engineered to express enhanced green fluorescent protein (EGFP), which is an additional

marker to assess infection. The replication characteristics of both viruses have been

previously extensively characterised, and the rg224RSV virus has been shown to exhibit

similar replication characteristics to the non-EGFP expressing parent virus (e.g. [15]). HEp-2

cell monolayers were infected with rg224RSV or rg224RSV-ΔG, and at between 1 and 4 dpi

the cell monolayers were stained using anti-F antibody and visualised using IF microscopy

(Fig. 7A). In rg224RSV-infected cells small clusters of infected cells were detected at 2 dpi

that increased in size by 3 dpi, and by 4 dpi syncytia formation had occurred. The pattern of

spread of the rg224RSV virus in the low MOI infection model was similar with that of RSV

A2 isolate described previously [19]. In contrast, the appearance of small clusters of infected

cells was detected in rg224RSV-ΔG -infected cell monolayers by 2 and 3 dpi, and by 4 dpi

intensely anti-F-stained smaller cell clusters were detected. The smaller clusters of infected

cells indicated that the rg224RSV-ΔG virus exhibited impaired spread in the cell monolayer.

The presence of anti-F stained debris over the cell monolayer was also observed by 3 dpi and

increased by 4 dpi. This was not apparent during the early stages of infection and is

consistent with localised cell damage caused by virus replication in these smaller infected cell

clusters. However, this cell debris was non-infectious since it did not appear to give rise to

infection in the cells that it came into contact with, even up to 6 dpi (Huong et al, unpublished

observations). These data are consistent with reports that RSV lacking the G protein is

impaired in HEp2 cells, and is over attenuated in animal cell models of infection [32]. In a

parallel analysis, the HEp2 cell monolayers infected with rg224RSV or rg224RSV-ΔG were

harvested at between 1 and 4 dpi and examined by immunoblotting using anti-G, anti-M2-1

and anti-F (Fig. 7B), which confirmed the absence of the G protein in rg224RSV-ΔG-infected

cells. Although a small reduction in the F protein was observed in rg224RSV-ΔG-infected

cells (presumably due to the reduced virus spread), the absence of the G protein appeared to

have no general inhibitory effect on the expression of other virus genes and was consistent

with previous observations (e.g. [17]).

Virus filaments have been shown to mediate virus infection and they play a role in

localised cell-to-cell virus transmission [2, 3, 33], and we therefore hypothesised that the

reduced virus spread observed in rg224RSV-ΔG-infected cells could be due to impaired virus

filament formation. The anti-F-stained cells were therefore examined by IF microscopy at 24

hpi (Fig. 8A) and by confocal microscopy at 3dpi (Fig. 8B) to determine if virus filament

formation occurred in the absence of the G protein. On rg224RSV-infected and rg224RSV-

ΔG-infected cells the presence of anti-F-stained virus filaments was detected. However, to

visualise the virus filaments more clearly on rg224RSV-ΔG-infected cells, the sensitivity on

the camera setting had to be reduced due to the apparent increase in anti-F staining intensity

on these cells. This apparent increase in antibody staining intensity was consistent with

accumulation of the F protein within the smaller infected cell clusters. This analysis indicated

that while virus spread was mediated less efficiently in the absence of the G protein, the

absence of the G protein did appear to not inhibit the trafficking of the F protein to the site of

virus assembly. Therefore, we can conclude that a block in virus filament formation could not

account for the observed reduced transmission of rg224RSV-ΔG.

Conclusion.

While our data suggests that the G protein was required for initiating a multiple cycle

RSV infection, it does not appear to be required to infect individual cells in the cell

monolayer. In this context our data is consistent with these previous report that have

examined RSV isolates that do not express the G protein (e.g. [32] [17]). However, these

reports have suggested that the G protein was dispensable for RSV to initiate infection in

some permissive cell types. The apparent contradiction in relation to the established role of

the G protein in mediating cell attachment was subsequently resolved when it was

demonstrated that in these cells the F protein can also bind to susceptible cells independently

of the G protein [34, 35]. In this context, low levels of localised transmission did occur in the

low MOI infection model using rg224RSV-ΔG, however this process was greatly facilitated

by the presence of the G protein. The expression of the recombinant RSV F protein alone is

sufficient to mediate syncytial formation [36, 37], suggesting that the association of the RSV

G protein with the F protein was not required to mediate membrane fusion. However, even in

viruses where a functional interaction between the F and respective attachment protein has

been demonstrated, the expression of the recombinant fusion protein alone can initiate

membrane fusion [38]. It is therefore possible that the association of the G protein with the F

protein may have other consequences. Receptor-mediated fusion involving a functional

association between the respective attachment protein and virus fusion protein is a feature of

many members of the paramyxoviruses [13]. In this paradigm the interaction between the

attachment protein and the cell receptor leads to structural changes in the conformation of the

F protein that initiates membrane fusion. In some cases both functional and structural

interactions between the virus attachment and fusion proteins have been demonstrated [39-

41]. During the process of virus particle assembly the F protein must be prevented from

mediating the premature fusion of the virus envelope and cell membranes, as this would be

expected to lead to an inhibition in virus infectivity i.e. virus inactivation. It is therefore

critical that the F protein is maintained in a pre-fusion form prior to cell membrane insertion

during virus entry. There are likely to be several factors that stabilise the F protein in its pre-

fusion conformation within the virus envelope. Based on our analysis, we can speculate that

the interaction between the F and G proteins may contribute to F protein stabilisation within

the virus envelope, and this may be a factor in preventing premature membrane fusion. This

suggestion is compatible with the results of the low MOI infection model, since although

premature membrane fusion would not be expected to inhibit virus gene expression in

infected cells, it would be expected to impair the localised spread of virus infection within the

cell monolayer. Interestingly, the structure of the F protein that was obtained by co-

crystallisation with an antibody that stabilized the F protein in its pre-fusion form [42] has

demonstrated that interactions with other proteins can potentially stabilise the pre-fusion form

of the RSV F protein. Furthermore the recombinant expressed F protein can be engineered to

form a stable pre-fusion structure with improved and enhanced immunological properties [43,

44]. The interaction between the F and G proteins has also been demonstrated in RSV virus-

like particles (VLPs) [30], suggesting that the formation of a protein complex is an intrinsic

biological property of these proteins and is not dependant on virus infection. Furthermore, the

VLPs containing both the F and G proteins exhibit enhanced neutralising antibody titres

compared with VLPs that only contain the F protein, suggesting that the G protein may

stabilise the F protein in its pre-fusion form [45]. In the context of the infectious virus, we

can hypothesise that in the virus envelope the interaction with the G protein may lead to

stabilization of the F protein in its pre-fusion form. Although a better understanding of the

interaction between the F and G protein should lead to an improved understanding of the

mechanism of virus assembly, this information may also facilitate the development of viable

antiviral strategies that can destabilise this interaction. Furthermore, the interaction of the G

and F proteins may have implications in the development of new RSV vaccine candidates.

The presence of the G protein in such formulations could both generate antibody responses

that block cell attachment [46] as well improve antibody responses against the F protein by

stabilising the F protein in its pre-fusion conformation. In this context, future work will be

needed to examine further the functional significance of the interaction between the F and G

proteins in the virus envelope, and to further define the structural basis for the formation of

this protein complex.

Declarations.

Consent for publication: All authors approved publication

Competing interests: We confirm that the authors have no conflict of interest.

Author contributions. TNH performed all experiments described in the manuscript, TBH

and RJS wrote the manuscript. RJS conceived of the study

Acknowledgements: We thank Mark Peebles (Nationwide Children's Hospital in Columbus,

Ohio) and Peter Collins (National Institute of Allergy and Infectious Diseases) for providing

the g224RSV-FSG and g224RSV-FS viruses. We thank the Ministry of Education, Republic

of Singapore (RG59/12) for providing funding (awarded to RJS).

References.

1.   Roberts SR, Compans RW, Wertz GW: **Respiratory syncytial virus matures at the apical**
surfaces of polarized epithelial cells. *Journal of virology* 1995, **69**(4):2667-2673.
2.   Gower TL, Pastey MK, Peeples ME, Collins PL, McCurdy LH, Hart TK, Guth A, Johnson TR,
Graham BS: **RhoA signaling is required for respiratory syncytial virus-induced syncytium**
formation and filamentous virion morphology. *Journal of virology* 2005, **79**(9):5326-5336.
3.   Ravi LI, Liang L, Wong PS, Brown G, Tan BH, Sugrue RJ: **Increased hydroxymethylglutaryl**
coenzyme A reductase activity during respiratory syncytial virus infection mediates actin
dependent inter-cellular virus transmission. *Antiviral research* 2013, **100**(1):259-268.
4.   Levine S, Klaiber-Franco R, Paradiso PR: **Demonstration that glycoprotein G is the**
attachment protein of respiratory syncytial virus. *The Journal of general virology* 1987, **68 (**
Pt 9):2521-2524.
5.   García-Beato R, Martínez I, Francí C, Real FX, García-Barreno B, Melero JA: **Host cell effect**
upon glycosylation and antigenicity of human respiratory syncytial virus G glycoprotein.
Virology 1996, **221**(2):301-309.
6.   Sugrue RJ, Brown C, Brown G, Aitken J, Mc LRHW: **Furin cleavage of the respiratory syncytial**
virus fusion protein is not a requirement for its transport to the surface of virus-infected
cells. *The Journal of general virology* 2001, **82**(Pt 6):1375-1386.
7.   González-Reyes L, Ruiz-Argüello MB, García-Barreno B, Calder L, López JA, Albar JP, Skehel JJ,
Wiley DC, Melero JA: **Cleavage of the human respiratory syncytial virus fusion protein at**
two distinct sites is required for activation of membrane fusion. *Proceedings of the*
National Academy of Sciences of the United States of America 2001, **98**(17):9859-9864.
8.   Zimmer G, Conzelmann KK, Herrler G: **Cleavage at the furin consensus sequence**
RAR/KR(109) and presence of the intervening peptide of the respiratory syncytial virus
fusion protein are dispensable for virus replication in cell culture. *Journal of virology* 2002,
76(18):9218-9224.
9.   McLellan JS, Ray WC, Peeples ME: **Structure and function of respiratory syncytial virus**
surface glycoproteins. *Current topics in microbiology and immunology* 2013, **372**:83-104.
10.   Matthews JM, Young TF, Tucker SP, Mackay JP: **The core of the respiratory syncytial virus**
fusion protein is a trimeric coiled coil. *Journal of virology* 2000, **74**(13):5911-5920.
11.   Lawless-Delmedico MK, Sista P, Sen R, Moore NC, Antczak JB, White JM, Greene RJ, Leanza
KC, Matthews TJ, Lambert DM: **Heptad-repeat regions of respiratory syncytial virus F1**
protein form a six-membered coiled-coil complex. *Biochemistry* 2000, **39**(38):11684-11695.
12.   McLellan JS, Yang Y, Graham BS, Kwong PD: **Structure of respiratory syncytial virus fusion**
glycoprotein in the postfusion conformation reveals preservation of neutralizing epitopes.
Journal of virology 2011, **85**(15):7788-7796.
13.   Jardetzky TS, Lamb RA: **Activation of paramyxovirus membrane fusion and virus entry.**
Current opinion in virology 2014, **5**:24-33.

14. Low KW, Tan T, Ng K, Tan BH, Sugrue RJ: **The RSV F and G glycoproteins interact to form a**
complex on the surface of infected cells. *Biochemical and biophysical research*
communications 2008, **366**(2):308-313.
15. Techaarpornkul S, Barretto N, Peeples ME: **Functional analysis of recombinant respiratory**
syncytial virus deletion mutants lacking the small hydrophobic and/or attachment
glycoprotein gene. *Journal of virology* 2001, **75**(15):6825-6834.
16. Hallak LK, Collins PL, Knudson W, Peeples ME: **Iduronic acid-containing glycosaminoglycans**
on target cells are required for efficient respiratory syncytial virus infection. *Virology* 2000,
271(2):264-275.
17. Kwilas S, Liesman RM, Zhang L, Walsh E, Pickles RJ, Peeples ME: **Respiratory syncytial virus**
grown in Vero cells contains a truncated attachment protein that alters its infectivity and
dependence on glycosaminoglycans. *Journal of virology* 2009, **83**(20):10710-10718.
18. Radhakrishnan A, Yeo D, Brown G, Myaing MZ, Iyer LR, Fleck R, Tan BH, Aitken J, Sanmun D,
Tang K *et al*: **Protein analysis of purified respiratory syncytial virus particles reveals an**
important role for heat shock protein 90 in virus particle assembly. *Molecular & cellular*
proteomics : MCP 2010, **9**(9):1829-1848.
19. Huong TN, Iyer Ravi L, Tan BH, Sugrue RJ: **Evidence for a biphasic mode of respiratory**
syncytial virus transmission in permissive HEp2 cell monolayers. *Virology journal* 2016,
13:12.
20. Cannon MJ: **Microplaque immunoperoxidase detection of infectious respiratory syncytial**
virus in the lungs of infected mice. *Journal of virological methods* 1987, **16**(4):293-301.
21. Brown G, Jeffree CE, McDonald T, Rixon HW, Aitken JD, Sugrue RJ: **Analysis of the**
interaction between respiratory syncytial virus and lipid-rafts in Hep2 cells during
infection. *Virology* 2004, **327**(2):175-185.
22. McDonald TP, Pitt AR, Brown G, Rixon HW, Sugrue RJ: **Evidence that the respiratory**
syncytial virus polymerase complex associates with lipid rafts in virus-infected cells: a
proteomic analysis. *Virology* 2004, **330**(1):147-157.
23. Rixon HW, Brown G, Aitken J, McDonald T, Graham S, Sugrue RJ: **The small hydrophobic (SH)**
protein accumulates within lipid-raft structures of the Golgi complex during respiratory
syncytial virus infection. *The Journal of general virology* 2004, **85**(Pt 5):1153-1165.
24. Jumat MR, Yan Y, Ravi LI, Wong P, Huong TN, Li C, Tan BH, Wang Y, Sugrue RJ:
Morphogenesis of respiratory syncytial virus in human primary nasal ciliated epithelial
cells occurs at surface membrane microdomains that are distinct from cilia. *Virology* 2015,
484:395-411.
25. Ravi LI, Li L, Wong PS, Sutejo R, Tan BH, Sugrue RJ: **Lovastatin treatment mitigates the pro-**
inflammatory cytokine response in respiratory syncytial virus infected macrophage cells.
Antiviral research 2013, **98**(2):332-343.
26. Klasse PJ: **Molecular determinants of the ratio of inert to infectious virus particles**. *Progress*
in molecular biology and translational science 2015, **129**:285-326.
27. Dropulic B, Hermankova M, Pitha PM: **A conditionally replicating HIV-1 vector interferes**
with wild-type HIV-1 replication and spread. *Proceedings of the National Academy of*
Sciences of the United States of America 1996, **93**(20):11103-11108.
28. Thompson KA, Yin J: **Population dynamics of an RNA virus and its defective interfering**
particles in passage cultures. *Virology journal* 2010, **7**:257.
29. Weidmann M, Sall AA, Manuguerra JC, Koivogui L, Adjami A, Traore FF, Hedlund KO,
Lindegren G, Mirazimi A: **Quantitative analysis of particles, genomes and infectious**
particles in supernatants of haemorrhagic fever virus cell cultures. *Virology journal* 2011,
8:81.

30.   Ravi LI, Tan TJ, Tan BH, Sugrue RJ: **Virus-induced activation of the rac1 protein at the site of**
479   **respiratory syncytial virus assembly is a requirement for virus particle assembly on**
480   **infected cells**. *Virology* 2021, **557**:86-99.

481   31.   Soderberg O, Gullberg M, Jarvius M, Ridderstrale K, Leuchowius KJ, Jarvius J, Wester K,
482   Hydbring P, Bahram F, Larsson LG *et al*: **Direct observation of individual endogenous**
483   **protein complexes in situ by proximity ligation**. *Nature methods* 2006, **3**(12):995-1000.

484   32.   Teng MN, Whitehead SS, Collins PL: **Contribution of the respiratory syncytial virus G**
485   **glycoprotein and its secreted and membrane-bound forms to virus replication in vitro and**
486   **in vivo**. *Virology* 2001, **289**(2):283-296.

487   33.   Gower TL, Peeples ME, Collins PL, Graham BS: **RhoA is activated during respiratory syncytial**
488   **virus infection**. *Virology* 2001, **283**(2):188-196.

489   34.   Karger A, Schmidt U, Buchholz UJ: **Recombinant bovine respiratory syncytial virus with**
490   **deletions of the G or SH genes: G and F proteins bind heparin**. *The Journal of general*
491   *virology* 2001, **82**(Pt 3):631-640.

492   35.   Techaarpornkul S, Collins PL, Peeples ME: **Respiratory syncytial virus with the fusion**
493   **protein as its only viral glycoprotein is less dependent on cellular glycosaminoglycans for**
494   **attachment than complete virus**. *Virology* 2002, **294**(2):296-304.

495   36.   Zimmer G, Budz L, Herrler G: **Proteolytic activation of respiratory syncytial virus fusion**
496   **protein. Cleavage at two furin consensus sequences**. *The Journal of biological chemistry*
497   2001, **276**(34):31642-31650.

498   37.   Zimmer G, Trotz I, Herrler G: **N-glycans of F protein differentially affect fusion activity of**
499   **human respiratory syncytial virus**. *Journal of virology* 2001, **75**(10):4744-4751.

500   38.   Connolly SA, Leser GP, Yin HS, Jardetzky TS, Lamb RA: **Refolding of a paramyxovirus F**
501   **protein from prefusion to postfusion conformations observed by liposome binding and**
502   **electron microscopy**. *Proceedings of the National Academy of Sciences of the United States*
503   *of America* 2006, **103**(47):17903-17908.

504   39.   Hu XL, Ray R, Compans RW: **Functional interactions between the fusion protein and**
505   **hemagglutinin-neuraminidase of human parainfluenza viruses**. *Journal of virology* 1992,
506   **66**(3):1528-1534.

507   40.   Melanson VR, Iorio RM: **Amino acid substitutions in the F-specific domain in the stalk of the**
508   **newcastle disease virus HN protein modulate fusion and interfere with its interaction with**
509   **the F protein**. *Journal of virology* 2004, **78**(23):13053-13061.

510   41.   Mirza AM, Iorio RM: **A mutation in the stalk of the newcastle disease virus hemagglutinin-**
511   **neuraminidase (HN) protein prevents triggering of the F protein despite allowing efficient**
512   **HN-F complex formation**. *Journal of virology* 2013, **87**(15):8813-8815.

513   42.   McLellan JS, Chen M, Leung S, Graepel KW, Du X, Yang Y, Zhou T, Baxa U, Yasuda E,
514   Beaumont T *et al*: **Structure of RSV fusion glycoprotein trimer bound to a prefusion-specific**
515   **neutralizing antibody**. *Science (New York, NY)* 2013, **340**(6136):1113-1117.

516   43.   McLellan JS, Chen M, Joyce MG, Sastry M, Stewart-Jones GB, Yang Y, Zhang B, Chen L,
517   Srivatsan S, Zheng A *et al*: **Structure-based design of a fusion glycoprotein vaccine for**
518   **respiratory syncytial virus**. *Science (New York, NY)* 2013, **342**(6158):592-598.

519   44.   Ngwuta JO, Chen M, Modjarrad K, Joyce MG, Kanekiyo M, Kumar A, Yassine HM, Moin SM,
520   Killikelly AM, Chuang GY *et al*: **Prefusion F-specific antibodies determine the magnitude of**
521   **RSV neutralizing activity in human sera**. *Science translational medicine* 2015,
522   **7**(309):309ra162.

523   45.   Cullen LM, Luo B, Wen Z, Zang L, Durr E, Morrison TG: **The Respiratory Syncytial Virus G**
524   **Protein Enhances the Immune Responses to the RSV F Protein in an Enveloped Virus-like**
525   **Particle Vaccine Candidate**. *bioRxiv : the preprint server for biology*
526   2022:2022.2009.2012.507712.

527   46.   Bergeron HC, Murray J, Nuñez Castrejon AM, DuBois RM, Tripp RA: **Respiratory Syncytial**
528   **Virus (RSV) G Protein Vaccines With Central Conserved Domain Mutations Induce CX3C-**
529   **CX3CR1 Blocking Antibodies**. *Viruses* 2021, **13**(2).

 Figure legends.

 Figure 1. Distribution of the F and G proteins on cells infected using the low multiplicity

 of infection model. HEp2 cell monolayers were RSV-infected using a multiplicity of

 infection (MOI) of 0.0001 and at between 1 and 3 days post-infection (dpi). **(A)** The

 monolayers in the tissue culture dish were fixed using 3% (w/v) paraformaldehyde and

 stained using anti-G followed by anti-mouse IgG conjugated to Alexa 555 and Alexa Fluor™

 488 Phalloidin. The stained cells were then viewed using fluorescence microscopy (objective

 x40 magnification). The infected cell clusters (white arrow) and syncytial formation (*) in the

 monolayer are highlighted, and Alexa Fluor™ 488 Phalloidin staining (F-actin) of the

 monolayer shows the cell monolayer. **(B)** The cell-associated and cell-free virus infectivity

 was recovered at each time of infection as described previously [19] and the infectivity

 assessed by microplaque assay. ND means not detected.

 Figure 2. Distribution of the F and G proteins on cells infected using the low multiplicity

 of infection model. HEp2 cell monolayers were RSV-infected using a MOI of 0.0001 and at

 2 dpi the cells were co-stained with anti-F and anti-G. **(A)** the stained cells were examined

 using phase contrast microscopy to view the cells in the field of view and by

 immunofluorescence (IF) microscopy to visualise the anti-G and anti-F staining on infected

 cells (objective x40 magnification). **(B and C)** As indicated, the stained cells were examined

 using bright-field (BF) microscopy to view the cells in the field of view and using IF

 microscopy to visualise the anti-G and anti-F staining on representative infected cells

 (objective x100 (oil) magnification). In the BF microscopy images the nuclei (N) of cells in

 the field of view are highlighted. In the IF microscopy images the insets are enlarged images

 taken from the cell at the region indicated (*). The anti-G stained (short white arrows) and

anti-G and anti-F co-strained (long white arrows) regions of the virus filaments are also

indicated in the insets.

Figure 3. Distribution of the F and G proteins on cells infected using the low multiplicity

of infection model. HEp2 cell monolayers were RSV-infected using a MOI of 0.0001 and at

2 dpi the cells were co-stained with anti-F and anti-G, and **(A)** the infected cell cluster

examined by confocal microscopy at an optical plane that allows visualisation of the (i) cell

periphery and (ii) cell top. The virus filaments are highlighted (white arrows). **(B and C)** Co-

localisation analysis was performed by examining the pixel distribution in the co-stained cells

at a focal plane that allows the **(B)** the cell periphery and **(C)** the cell top to be examined. **B**(i)

shows a scatter plot of the entire image in **B**(ii), while **C**(i) is a scatter plot of the area

highlighted in **C**(ii) by white circle. The virus filaments are highlighted (white arrows).

Figure 4. The F and G proteins interact on the surface of virus filaments. HEp2 cell

monolayers were **(A)** mock-infected and **(B)** RSV-infected using a multiplicity of infection of

0.0001, and at 2 days-post infection (dpi) the cells were labelled with anti-F and anti-G and

the proximity ligation assay (PLA) performed and the cells stained using anti-G. In each case

the antibody staining and PLA signal is shown. **(B**(i)) highlights the cell top and B(ii) the cell

periphery. **(C)** Cells were also singly labelled using either (i) anti-G or (ii) anti-F and the

PLA performed. The cells were then stained with anti-G and anti-F as indicated.

Figure 5. The PLA signal is localised to virus filaments. HEp2 cell monolayers were

infected with RSV using a multiplicity of infection of 0.0001, and at 2 days-post infection

(dpi) the cells were labelled with anti-F and anti-G and the proximity ligation assay (PLA)

performed. **(A)** The cells were stained using anti-F, and the anti-F staining and PLA signal

was examined by confocal microscopy at a focal plane that allowed the visualisation of the

virus filaments at different magnifications ((i) to (iii)). Inset in (iii) is an enlarged image taken

from each plate. The virus filaments are highlighted (white arrow). **(B)** Co-localisation

analysis was performed by examining the pixel distribution in the anti-F stained cells

following the PLA assay. (i) Is a scatter plot of the region highlighted in (ii). **(C)** Distribution

of staining intensity in the anti-F stained cells following the PLA assay. (i) The intensity

profile of the area demarcated by the white line in (ii).

Fig. 6. Co-precipitation of the F and G proteins in the low multiplicity of infection

model and in purified virus particles. (A) HEp2 cell monolayers were mock-infected (M)

and RSV-infected using a multiplicity of infection of 0.001 (I) and at 48 hr post-infection

(hpi) the cells were surface-biotinylated. (i) the clarified detergent extracts were either

examined directly (total lysates) or immunoprecipitated using anti-G, anti-F or anti-NP as

indicated. Protein species corresponding in size to the F1, F2 and G proteins are also

indicated. (ii) Densitometry analysis of the immunoprecipitation from mock-infected (MI)

and virus-infected (RSV) cells are shown for the immunoprecipitation with anti-G (G), anti-F

(F) and anti-NP (NP). **(B)** The virus particles were harvested as described previously [18] and

(i) The virus preparation was examined by immunoblotting with anti-G (G) and anti-F (F),

and protein bands corresponding in size to the mature G protein and the F1 protein are

indicated. **(ii)** The virus preparation was surface-biotinylated and the proteins in the virus

fraction was analysed by Western blotting. The total surface biotinylated proteins in the

clarified detergent extract are shown. (iii) The surface biotinylated virus was detergent

extracted and immunoprecipitated using anti-F (F), anti-G (G) and anti-NP (NP). Protein

species corresponding in size to the F1 protein and the mature G protein are indicated. (iv)

Densitometry analysis of the immunoprecipitation assays with anti-F (F), anti-G (G) and anti-

NP (NP).

Figure 7. The presence of the G protein is required for efficient localised virus spread in

low multiplicity of infection model. (A) HEp2 cells were infected with either (i) rg224RSV

or (ii) rg224RSV-ΔG using a multiplicity of infection of 0.0001 and at between 1 and 4 days

post-infection (dpi) the cells were stained with anti-F. The anti-F and the green fluorescent

protein (EGFP) staining patterns were imaged using immunofluorescence

microscopy(objective x20 magnification). The infected cell clusters are highlighted (open

white boxes), the single infected cells (arrow heads) and cell debris (longer arrows) are

indicated. **(B)** At between 1 and 4 dpi the rg224RSV- and rg224RSV-ΔG-infected cell

monolayers were extracted in boiling mix and examined by immunoblotting using anti-G,

anti-M2-1 and anti-F. Protein bands corresponding in size to the respective virus proteins are

indicated. M is mock-infected.

Figure 8. The presence of the G protein is not required for trafficking of the F protein to

the sites of virus assembly. (A) Cells were infected using a MOI of 0.0001 and at 24 hrs

post-infection the cells were stained using ant-F and examined by immunofluorescence (IF)

microscopy to visualise the anti-F staining on infected cells (objective x100 (oil)

magnification). **(B)** At 3 dpi the anti-F stained rg224RSV or rg224RSV-ΔG infected cells

were examined by confocal microscopy to detect the presence of virus filaments. The image

of rg224RSV-ΔG-infected cells was recorded using a reduced camera setting that allows the

virus filaments to be more clearly imaged. The virus filaments are highlighted (white arrows).

	Virus titer pfu/ml	
Days post-infection	cell-associated	cell-free
1	2.0×10^2	ND
2	1.4×10^3	ND
3	9.5×10^3	2.1×10^2