Disorganization of basement membrane zone architecture causes impaired melanocyte inhabitation in vitiligo

(15 words)

Fei Yang [1,2§], Lingli Yang [1§*], Yasutaka Kuroda [1,2], Sylvia Lai [1], Yoshito Takahashi [2], Tetsuya Sayo [2*], Takeshi Namiki [3], Kimiko Nakajima [4], Shigetoshi Sano [4], Shintaro Inoue [5], Daisuke Tsuruta [6] and Ichiro Katayama [1]

[1] Department of Pigmentation Research and Therapeutics, Graduate School of Medicine, Osaka Metropolitan University, Osaka 5450051, Japan

[2] Biological Science Research Laboratories, Kao Corporation, Odawara 2500002, Japan

[3] Department of Dermatology, Tokyo Medical and Dental University Graduate School and Faculty of Medicine, Tokyo 1138510, Japan

[4] Department of Dermatology, Kochi Medical School, Kochi University, Kochi 7838505, Japan

[5] Department of Cosmetic Health Science, Gifu Pharmaceutical University, Gifu 5011196, Japan

[6] Department of Dermatology, Graduate School of Medicine, Osaka Metropolitan University, Osaka 5458585, Japan

[§] These two authors contributed equally to this work: Fei Yang, Lingli Yang.

[*] Correspondence: yang.lingli@omu.ac.jp (L.Y.); sayou.tetsuya@kao.com (T.S.); Tel./Fax: +81-6-6556-7618 (L.Y.); Tel./Fax: +81-465-34-6111 (T.S.)

Keywords: fibroblasts; melanocyte; basement membrane; vitiligo; MMP2

Abbreviations: BM, basement membrane; BMZ, basement membrane zone; MMP2-Fb, MMP2-overexpressing fibroblasts; MMP2, matrix metalloproteinase 2.

Abstract (150 words)

Vitiligo is a common but puzzling skin disorder characterized by the selective loss of epidermal melanocytes due to unknown mechanisms. Here, we show patients with vitiligo have disruptions of basement membrane zone (BMZ) architecture including branched, fragmented, and multilayered lamina densa with increased numbers of dermal fibroblasts and overexpression of MMP2 in those cells. Extracts of vitiliginous skin showed increased MMP2 protein expression, with a large portion of the MMP2 proteins in the active form. Intradermal injection of MMP2-overexpressing fibroblasts in K14-SCF transgenic mice induced a vitiligo-like skin appearance and disappearance of melanocytes, which were reversed by coadministration of MMP2 inhibitors. These results suggest that disorganization of the BMZ driven by MMP2 overexpression in dermal fibroblasts may cause the impaired melanocyte inhabitation observed in vitiligo.

Introductions

Vitiligo is a puzzling disorder characterized by selective loss of epidermal melanocytes by an unknown mechanism[1]. Vitiligo was first described 3,500 years ago and is today a common skin disorder with an incidence of 1–2% wordwide[2, 3]. The most striking histological feature of vitiligo is the disappearance of melanocytes in the epidermis. The cellular and molecular mechanisms underlying melanocyte disappearance have attracted much interest from cell biologists and dermatologists. So far, many theories regarding the cause of melanocyte disappearance have been proposed, including autoimmune mechanisms, neural deficiencies, impaired redox status, and reduced melanocyte adhesion[4, 5]. A consensus on the true mechanism remains elusive.

In recent years, research on vitiligo pathogenesis has mainly focused on the importance of the IFNγ-CXCL9/10/CXCR3 axis. Autoreactive CD8[+] T cells and resident memory CD8[+] T cells are suggested to play an important role in the elimination of melanocytes in vitiliginous skin[4, 6]; however, compared with other chronic skin conditions, vitiligo often lacks clinical symptoms and overt signs of inflammation. Histologically, although limited lymphocytic infiltration is observed in the margin of early lesions, skin with established vitiligo lesions frequently displays unremarkable histopathological changes other than a near complete absence of melanocytes. These features make it difficult to understand why the characteristic white patches persist in the absence of overt infiltrates. There is no known cure for vitiligo, and the characteristic white patches have a substantial impact on patients'

quality of life. Clarification of the mechanism behind the loss of melanocytes in the absence of

infiltration or severe inflammation may lead to new therapy options.

Epidermal melanocytes are specialized, pigment-producing cells derived from the neural crest.

Under normal homeostatic conditions, they localize in the basal layer of the epidermis where they are

maintained in a quiescent state and rarely proliferate, keeping a life-long stable ratio of 1:5 with basal

keratinocytes[7]. Physiologically, dysregulation of homeostasis disturbs the epidermal-melanin unit and

triggers pigmentary disorders[7, 8]. Melanocyte homeostasis is intricately regulated by epidermal

keratinocytes, the surrounding stroma, and nearby dermal cells[9]. Cellular physiology and

pathophysiology depend on both cell–cell interactions and cell–matrix interactions, and differentiated

epidermal melanocytes are strictly localized at the basement membrane (BM) and cannot survive if

they leave the basal layers, except in nevi or melanomas[10].

The BM is a thin sheet of highly specialized extracellular matrix[11, 12] located between the epidermis

and the dermis that participates in diverse cellular functions including adhesion, proliferation,

differentiation, and survival[13]. There are three layers within the BM: the lamina lucida, the lamina

densa, and the anchoring fibrils. The lamina densa is a dense, colored layer approximately 30–60 nm

in width that is closest to the underlying connective tissue. The anchoring fibrils are linked to the

papillary dermis and localized in the lower part of the BM. Destruction of extracellular matrix

macromolecules is seen in diseases such as arthritis[14], periodontal disease[15], and dermatological

disorders[16]; however, there has been little research on abnormal BM morphology in vitiligo[17, 18, 19, 20].

Because the BM provides structural support for keratinocytes and melanocytes, it is reasonable to

hypothesize that structural abnormalities of the BM may have an impact on melanocyte disappearance

in vitiligo.

Collagen IV is the most abundant BM component and serves as the backbone of the BM. Laminin,

a matrix protein generated by keratinocytes and fibroblasts, is the second most abundant BM

component[21]. In the BM, laminins are integrated into the type IV collagen scaffold by crosslinking to

assemble the sheet-like BM complex[21]. Basal keratinocytes firmly adhere to the BM matrix via

hemidesmosomes and focal adhesions[21]. Hemidesmosomes link intracellular keratin intermediate

filaments to the underlying BM via integrin α6β4–laminin 332 binding and collagen XVII–collagen

IV binding. Focal adhesions link intracellular actin filaments to the BM via integrin α3β1–laminin 511

binding[21].

Matrix metalloproteinases (MMPs) are a group of proteolytic enzymes that use zinc to control cell

matrix composition. Metalloproteinases and the proteolytic breakdown of BM are essential for normal

biological processes such as embryonic development, neuron growth, hair follicle cycling, bone

remodeling, wound healing, and angiogenesis[22]. Dysregulated metalloproteinase expression leads to

pathological conditions including inflammation, arthritis, cardiovascular diseases, nephritis,

neurological diseases, pulmonary diseases, and cancer[22, 23]. MMP levels are elevated in various tissues

in autoimmune disorders[24], including brain tissue in autoimmune encephalomyelitis[25, 26], cerebrospinal

fluid in multiple sclerosis[27], synovial fluid in rheumatoid arthritis[28], and salivary glands in Sjogren's

Syndrome[29]. MMP2 was first reported by Liotta et al.[30], who found that an enzyme secreted by a

metastatic murine tumor cleaved type IV collagen, producing fragments. The enzyme did not attack

collagen types I, II, III, and V. When it was purified, the enzyme was found to consist of two forms of

approximately 72 kDa and 62 kDa, representing the pro and active forms, respectively[31].

We used electron microscopy, three-dimensional/organotypic culture, and mouse experiments to

reveal a novel mechanism to explain melanocyte disappearance from the BM in vitiligo. Compared

with healthy skin, lesional skin from patients with vitiligo displayed abnormalities in the structure and

components of the BM. The abnormalities were caused by fibroblasts that highly expressed MMP2,

which degraded BM matrix components and induced failure of melanocyte inhabitation. These

findings provide new knowledge about the etiology and pathogenesis of vitiligo and suggest a new

approach to vitiligo treatment.

Results

Basement membrane zone architecture is disorganized in vitiligo

To investigate the BM zone (BMZ) in vitiligo in detail, we performed an ultrastructural analysis

using biopsy specimens from five healthy donors and five patients with generalized vitiligo (two stable,

three progressive). We observed abnormal BM morphology in the samples of both stable and

progressive vitiligo. As shown in **Fig 1a**, the BM of healthy skin contained intact lamina densa with

high electron density, whereas the BM of vitiliginous skin contained branched and fragmented lamina

densa, some of which were multilayered. Furthermore, the anchoring fibrils (**Fig 1a, red asterisks**)

were distributed normally in the BM of healthy skin, but they were scattered and distributed in both

the upper and the lower portion of the branched lamina densa in vitiliginous skin.

Next, we used immunofluorescence staining to investigate the expression and localization of BM

proteins. Skin specimens from five healthy donors and 11 patients with generalized vitiligo (five stable,

six progressive) were stained with antibodies against collagen IV (the main component of BM) and

TRP1 (a melanocyte marker). In healthy skin, the BM had normal morphology (a thin continuous

linear band of collagen IV) and a normal number and distribution of melanocytes (**Fig 1b**). By contrast,

in vitiliginous skin, the BM had various abnormal morphologies, including obvious thickening (**Fig**

1b, upper panel of Vitiligo-L), severe disruption (**Fig 1b, middle panel of Vitiligo-L**), and

multilayering (**Fig 1b, lower panel of Vitiligo-L**), accompanied by the disappearance of melanocytes.

We investigated the ultrastructural localization of collagen IV by immunoelectron microscopy using

an antibody against collagen IV. As expected, in an ultrathin section of healthy skin, we observed intact

BM, with collagen IV-labeled gold-particle deposits located close to the lamina densa (**Fig 1c, white**

arrowhead in left panel) in a continuous BM-like arrangement. By contrast, in an ultrathin section

of vitiliginous skin, we observed conspicuous structural alterations of the BM, with collagen IV-

labeled gold-particle deposits scattered across a wide area not only close to the normal lamina densa

area (**Fig 1c, white arrowhead in right panel**) but also near the branched structures (**Fig 1c, white**

arrow) and multilayered structures (**Fig 1c, white asterisk**) of lamina densa. These results revealed

that the BMZ architecture is disrupted in lesional vitiliginous skin.

Matrix metalloproteinase 2 expression is elevated in the dermis of vitiliginous skin

To ascertain why the BM structure was disrupted, we measured the expression levels of MMP2

(also known as gelatinase A) and MMP9 (also known as gelatinase B), the major proteolytic enzymes

capable of degrading BM collagen, in skin sections from patients with vitiligo. Immunofluorescence

staining was performed on frozen skin sections using anti-MMP2 antibody and anti-MMP9 antibody.

In comparison with healthy skin, lesional vitiliginous skin showed modestly increased MMP9

expression and dramatically increased MMP2 expression in the upper dermis (**Fig 2a and 2b**).

Fibroblasts[32] and vascular endothelial cells[33] are the main MMP2-producing cells in human skin. To

determine which cell types were responsible for the high MMP2 production in vitiliginous skin, we

performed immunolocalization studies on vitiliginous and healthy skin tissues. Frozen sections from

five healthy controls and 11 patients with vitiligo were incubated with anti-MMP2 antibody and then

counterstained with anti-HSP47 (a skin fibroblast marker) and anti-CD31 (a marker of endothelial

cells) antibodies. Remarkably, the number of fibroblasts was significantly increased in the vitiliginous

skin; however, there was no appreciable difference in the number of endothelial cells between the

vitiliginous skin and the healthy skin (**Fig 2c and 2d**). In addition, the percentage of MMP2-positive

fibroblasts was almost four times higher in the vitiliginous skin than in the healthy skin, but there was

no significant difference in the percentage of MMP2-positive endothelial cells (**Fig 2e and 2f**). To

determine whether MMP2-positive fibroblasts were present in the area surrounding the BM, we

performed immunofluorescent labeling with anti-MMP2 antibody and counterstaining with anti-

HSP47 and anti-collagen IV (a marker of BM) antibodies. This revealed that in the vitiliginous skin,

but not the healthy skin, MMP2-positive fibroblasts were located near the BM (**white arrowheads in**

Fig 2g) and also went through the BM to reach the epidermis (**white arrows in Fig 2g**).

Next, we examined the activity of MMP2 by western blot using skin lysates from three patients

with vitiligo (one stable, two progressive) and three healthy controls. In the healthy skin, MMP2 was

slightly expressed and mainly in the pro (inactive) form; however, in the vitiliginous skin, MMP2 was

highly expressed and mainly in the active form (**Fig 2h and 2i**). These results revealed that vitiliginous

skin had increased MMP2 in the dermis, produced mainly by upper-dermis fibroblasts, and the MMP2

was mainly in the active form, suggesting that MMP2-induced BM degradation might be involved in

the production of vitiligo lesions.

Overexpression of MMP2 in dermal fibroblasts results in melanocyte disappearance

in reconstructed human skin equivalents and human *ex vivo* cultured skin explants

To determine whether damage to the BM precedes the disappearance of melanocytes in vitiliginous

skin, we developed an *in vitro* three-dimensional (3D) skin equivalent model. In each 3D skin

equivalent, half of the collagen gel in the dermis layer was prepared with mock-transfected fibroblasts

(Mock-Fbs) and half with MMP2-overexpressing fibroblasts (MMP2-Fbs) (**Fig 3a**). Once the

structured collagen gel was established, melanocytes and keratinocytes were uniformly seeded on top

of the gel. After 1 week of air-lift culture, macroscopically pigmented skin developed on the part of

the gel prepared with Mock-Fbs, whereas no pigmentation developed on the part of gel prepared

with MMP2-Fbs (**Fig 3b**). Immunofluorescent staining of melanocytes and BM revealed that in the

3D skin equivalents prepared with Mock-Fbs, the BM had a typical thin line morphology, and

melanocytes were uniformly distributed among the keratinocytes in the basal layer; however, in the

3D skin equivalents prepared with MMP2-Fbs, the BM appeared damaged, and the melanocytes had

disappeared (**Fig 3c**).

Next, we created an *ex vivo* subcutaneous injection model in which Mock-Fbs or MMP2-Fbs were

subcutaneously inoculated into cultured, full-thickness, fresh skin explants. To track the injected

fibroblasts, we labeled them with PKH fluorescence cell-membrane labeling dye (**Fig 3d**). After *ex*

vivo skin culture, the skin explants were stained with anti-Melan-A and anti-collagen IV antibodies.

In contrast to skin explants injected with PBS or Mock-Fbs, skin explants injected with MMP2-Fbs

displayed obvious BM disruption and melanocyte disappearance (**Fig 3e**). These results suggest that

increased MMP2 expression by dermal fibroblasts is a key factor in melanocyte disappearance in

vitiligo.

Melanocyte disappearance results from decreased adhesion to the BM

We performed immunofluorescent staining to further examine the disappearance of melanocytes

from the damaged BM. The melanocytes were anchored to the BM and located on the basal layer in

the 3D skin equivalents prepared with Mock-Fbs, (**Fig 4a, Mock-Fb**); however, they were detached

from the BM and tended to be located at the suprabasal layer in the marginal zone of the depigmented

3D skin equivalents (**Fig 4a, MMP2-Fb (MZ)**). These detached melanocytes were negative for

cleaved-caspase 3 (**Fig 4a**), suggesting that apoptotic events were not involved in their disappearance.

These data suggested that abnormal adhesion between melanocytes and the BM was involved in the

melanocyte disappearance, so we next focused on melanocyte adhesion.

Cells in the basal layer of the epidermis are bound to the BM mainly through integrins, but the

mechanistic details of melanocyte adhesion to the BM are still unknown. Melanocytes have been

reported to express integrin β1 both *in vitro* and *in vivo*[34]. In addition, discoidin domain receptor 1

(DDR1), a membrane protein, was recently reported to bind collagen IV and mediate the adhesion of

melanocytes to BM[35]. Furthermore, some studies reported that loss of melanocytes in vitiligo resulted

from decreased DDR1 expression in melanocytes[36, 37]. We examined adhesion patterns between

melanocytes and the BM matrix using *in situ* proximity ligation assay (PLA), an antibody-based

technology that enables the visualization of protein–protein interactions. Specifically, we examined

integrin β1–laminin-binding and DDR1–collagen IV binding in 3D skin equivalents. In the PLA, when

a pair of specific primary antibodies (DDR1 and collagen IV; integrin β1 and laminin) is in close

proximity, complementary DNA strands on a corresponding pair of secondary antibodies with PLA

probes engage in rolling circle amplification and generate a single fluorescent signal *in situ*, indicating

the presence of the corresponding protein–protein interaction (**Fig 4b**). We detected PLA signals for

DDR1–collagen IV and integrin β1–laminin along the basal layer of epidermis in 3D skin equivalents

prepared with Mock-Fbs (**Fig 4c**); however, in the marginal zone of depigmented 3D skin equivalents

prepared with MMP2-Fbs, the DDR1–collagen IV and integrin β1–laminin PLA signals were

markedly decreased (**Fig 4c and 4d**).

We confirmed the detachment of melanocytes from the BM by *in situ* PLA of perilesional skin from

patients with vitiligo (**Fig 5a**). Immunofluorescence staining did not show any difference in the

expression levels of DDR1, collagen IV, integrin β1, and laminin between healthy skin and perilesional

vitiliginous skin (**Fig 5b, 5c, and 5f**). We visualized the adhesions between melanocytes and the BM

matrix by *in situ* PLA and co-immunohistochemistry staining of melanocytes with anti-TRP1 antibody

(**Fig 5d and 5e, white rectangle box in right lower panel**). In melanocytes of healthy skin, integrin

β1–laminin interactions were much more predominant than DDR1–collagen IV interactions; however,

both interactions were significantly reduced in melanocytes of perilesional vitiliginous skin (**Fig 5e**

and 5g). These data demonstrated that the two major melanocyte–BM adhesion patterns were

strikingly weakened in vitiliginous skin (**Fig 5h**).

In contrast to the detachment of melanocytes, we did not observe detachment of keratinocytes

(blistering) in perilesional vitiliginous skin despite a weakened focal adhesion signal (integrin β1–

laminin binding) around basal keratinocytes (**Fig 4c; Fig 5e**). Therefore, we examined the expression

levels and binding signals of hemidesmosome proteins. There was no difference in the expression

levels of integrin α6 and laminin between healthy skin and lesional vitiliginous skin (**Fig 6a and 6c**);

however, the expression levels of collagen XVII and collagen IV were significantly increased in the

lesional vitiliginous skin (**Fig 6b and 6c**). We used *in situ* PLA to examine the hemidesmosome

adhesion signal between basal keratinocytes and BM matrix (**Fig 6d**). In basal keratinocytes of healthy

skin, integrin α6–laminin interactions were much more predominant than collagen XVII–collagen IV

interactions (**Fig 6e and 6f**). By contrast, in basal keratinocytes of vitiliginous skin, the integrin α6–

laminin interaction was dramatically reduced, whereas the collagen XVII–collagen IV interaction was

strikingly increased (**Fig 6e and 6f**). These results indicated that the adhesion patterns between basal

keratinocytes and the BM were strikingly altered in lesional vitiliginous skin (**Fig 6g**), and a

compensatory increase in hemidesmosome adhesion via collagen XVII–collagen IV binding may be

the reason there was no skin blistering.

Decreased melanocyte adhesion to the BM is caused by abnormal BM architecture

The expression levels of DDR1, collagen IV, integrin β1, and laminin were similar between

vitiliginous and healthy skin, but the vitiliginous skin displayed decreased DDR1–collagen IV and

integrin β1–laminin interactions (**Fig 5**). To clarify what caused the reduction of these interactions, we

assessed the physical distance between the melanocytes and the BM using electron microscopy (**Fig**

7a). As basal cells are known to attach to the lamina densa of the BM, we measured the width of the

lamina lucida (the distance between the basal cell plasma membrane and the lamina densa) in healthy

skin and vitiliginous skin, but we unexpectedly found no difference between healthy skin and lesional

vitiliginous skin (**Fig 7a and 7c**). We then performed immunoelectron microscopy to determine

whether the decreased binding was due to aberrant localization of adhesion molecules. In healthy skin,

laminin, the predominant adhesion molecule on the BM responsible for melanocyte binding, was

clearly distributed along and confined to the lamina densa; however, in lesional and perilesional

vitiliginous skin, laminin was diffusely spread in the lamina densa and the beneath the disorganized

BM architecture (**Fig 7b**). Furthermore, the amount of laminin confined in the lamina densa was

markedly reduced in vitiliginous skin compared with that in healthy skin (**Fig 7b and 7d**). These data

suggest that laminin was aberrantly distributed in the vitiliginous skin, and the disorganized BMZ

architecture made it difficult for laminin to bind with integrin β1 on epidermal melanocytes.

Inhibition of MMP2 reversed depigmentation in a novel mouse model of vitiligo

based on subcutaneous inoculation of MMP2-Fbs.

We used K14-SCF transgenic mice characterized by the retention of melanocytes in the epidermis

to examine skin color changes in vitiligo. Murine Mock-Fbs and MMP2-Fbs were subcutaneously

transplanted into the back skin of K14-SCF mice (**Fig 8a**). One week after transplantation, no color

change was observed on the backs of mice transplanted with Mock-Fbs; however, depigmentation

spots were observed on the backs of mice transplanted with MMP2-Fbs (**Fig 8b**). Skin tissue from the

mice was subjected to immunofluorescence staining. Mice transplanted with Mock-Fbs or MMP2-Fbs

had many more fibroblasts in their skin tissues than control mice injected with PBS (**Fig 8c**).

Furthermore, skin tissues from mice with Mock-Fb transplantation contained epidermal melanocytes

and intact BM, whereas those from mice with MMP2-Fb transplantation contained no epidermal

melanocytes and had disrupted BM structure (**Fig 8c**). These data indicated that the subcutaneous

MMP2-Fb transplantation in K14-SCF mice provided a vitiligo-like depigmentation model. Next, to

determine whether inhibition of MMP2 could reverse the vitiligo-like depigmentation, we

subcutaneously co-injected MMP2-Fbs and an MMP2 inhibitor, either SB-3CT or tanomastat, into the

back skin of K14-SCF mice (**Fig 8d**). Compared with PBS as a control, both SB-3CT and tanomastat

remarkably inhibited the depigmentation of back skin (**Fig 8e and 8f**). These data suggest that

melanocytes disappeared from the basal layer of the depigmented epidermis as a consequence of

MMP2-induced disruption of the BM structure.

Discussion

The BM of vitiliginous skin has been shown to be morphologically thickened[17, 18], damaged[19], or

branched[20], but it is unknown how the BM structure is disrupted and what role BM disruption plays

in the pathophysiology of vitiligo. We confirmed the abnormal BM structure in vitiliginous skin and,

furthermore, that the disorganized BMZ prevents melanocytes from adhering to the BM via integrin

β1–laminin binding. Moreover, we discovered that the disorganized BM architecture is caused by

increased MMP2 expression in dermal fibroblasts. MMP2-Fbs implanted in 3D skin equivalents and

ex vivo cultured skin explants showed BM structural disruption and melanocyte detachment. K14-SCF

mice developed vitiligo-like skin lesions after subcutaneous inoculation with MMP2-Fbs, and

treatment with MMP2 inhibitors successfully reduced the depigmentation in these mice.

Melanocyte floating and defective melanocyte adhesion in vitiligo were described previously[38, 39],

but the majority of previous studies focused on melanocyte–keratinocyte adhesion. Few investigations

have looked at melanocyte–BM adhesion. We verified melanocyte detachment in vitiligo but found

that neither the expression levels of melanocyte adhesion molecules nor the expression levels of BM

proteins were decreased. Instead, immunoelectron microscopy, histo-immunofluorescence staining,

and PLA analyses revealed that the reduction in melanocyte adhesion was due to disorganization of

the BMZ architecture. These findings imply that the detachment of melanocytes in vitiligo is caused

by an aberrant spatial distribution of adhesion molecules in the BM. In contrast to the decreased

adhesion and detachment of melanocytes, we did not observe any detachment of keratinocytes (i.e.,

blistering), which suggests that a powerful compensating mechanism in keratinocytes enables them to

adhere to the BM via collagen XVII and other molecules in hemidesmosomes, even when the

distribution of laminin is aberrant. Our ultrastructural investigations of vitiliginous skin confirmed the

existence of hemidesmosomes in basal keratinocytes. It is well known that there are no

hemidesmosome complexes in melanocytes, as these structures are exclusive to keratinocytes[40]. The

attachment of melanocytes to the BM is considerably weaker than that of keratinocytes and is mediated

by integrin β1–laminin binding[34] and DDR1–collagen IV binding[35]. Differentiated human

melanocytes are restricted to the BM, and their continued attachment to the BM is essential for their

survival, as they cannot survive in the upper epidermal layers unless they are changed into nevi or

melanomas[35]. Several studies hypothesized that melanocytes disappear in vitiligo because of weak

attachment to the epidermal BM, and they found that expression of the melanocyte–BM adhesion

molecule DDR1 was decreased in vitiligo[36, 37]. However, our *in situ* PLA study demonstrated that the

predominant adhesion between melanocytes and the BM is due to integrin β1–laminin binding, which

was dramatically reduced in vitiliginous skin. The protein expression levels of DDR1, integrin β1,

laminin, and collagen IV were unchanged in vitiliginous skin, which may explain why the reduced

integrin β1–laminin binding was not uncovered previously. Our findings suggest that structural

abnormality of the BM, rather than altered expression levels of adhesion molecules, was mostly

responsible for the decreased adhesion of melanocytes.

The two main MMPs that break down epidermal BM components are MMP2 and MMP9[23, 41].

MMP2 was dramatically elevated in the dermis of lesional vitiliginous skin, regardless of whether the

vitiligo was stable or progressive. Additionally, MMP9 expression was slightly increased in the dermis

of vitiliginous skin. Recent research revealed decreased E-cadherin expression in melanocytes[39] and

increased MMP9 expression in the epidermis of lesional vitiliginous skin[42], suggesting that reduced

E-cadherin levels in melanocytes and MMP9-induced breakdown of E-cadherin on keratinocytes and

melanocytes may cause melanocyte detachment in vitiligo. Our results show, however, that for

melanocytes to inhabit the skin's basal layer, they must adhere not only to neighboring keratinocytes

but also to the BM. In addition, MMP9 should affect not only cell–cell adhesion but also BM assembly.

Reports investigating *in vitro* cultured human melanocytes[43] and immortalized mouse melanoblasts[44]

demonstrated that reduced MMP2 expression in those cells was associated with cell migration and

might therefore be involved in the pathogenesis of vitiligo. However, in light of the aberrant BM

structure that we observed in vitiliginous skin, increased expression of MMP2 in dermal fibroblasts is

probably the most important event in vitiligo pathogenesis *in vivo*.

MMP2 is expressed constitutively by many cell types in culture. In adult tissues, MMP2 has been

identified in keratinocytes, skin fibroblasts, chondrocytes, osteoblasts, endothelial cells, monocytes,

and a number of other cell types[22]. Our immunolocalization analyses demonstrated that the majority

of the increase in MMP2-positive cells in vitiliginous skin was due to MMP2-expressing dermal

fibroblasts. In agreement with this, a recent study using single-cell analysis of lesional vitiliginous

skin revealed that dermal fibroblasts were associated with the loss of melanocytes[45]. Several

researchers have noted abnormal dermal fibroblasts in lesional vitiliginous skin[45, 46]. Kovacs et al.

observed that dermal fibroblasts in vitiliginous skin had a myofibroblast phenotype and a propensity

for premature senescence[47]. There is growing evidence that dermal fibroblasts contribute to the defects

associated with melanocytes in vitiligo, although their exact role remains unclear. It is currently

generally accepted that vitiligo is an autoimmune disease resulting from the destruction of melanocytes

by melanocyte-specific autoreactive $CD8^+$ T lymphocytes, and animal models of vitiligo have been

established accordingly[4, 45]. Despite that, patients with vitiligo frequently lack obvious inflammatory

clinical signs, so we looked for a mechanism other than direct immune cell participation. Our findings

enabled us to establish a new alternative mouse model of vitiligo based on subcutaneous injection of

MMP2-Fbs, which may provide new insights into pathomechanisms of non-immune cell-mediated

vitiligo.

MMP2 expression was elevated in dermal fibroblasts of vitiliginous skin, but the underlying reason

is unknown. There are many reports on paracrine modulation of MMP2 expression. Numerous

cytokines that are increased in lesional vitiliginous skin, including interleukin (IL)-1β[48, 49], IL-6[48],

tumor necrosis factor (TNF)-α[49, 50], and TGF-β[51, 52], have been reported to stimulate MMP2 expression.

Destruction of extracellular matrix macromolecules is seen in other diseases such as arthritis[14],

periodontal disease[15], and dermatological disorders[16], and invading inflammatory cells exacerbate the

pathogenesis of those diseases. These invading immune cells are thought to play an etiologic role

through their ability to produce cytokines that induce MMP expression in resident mesenchymal cells[53].

Damage to the epidermal BM alters interactions between epidermal keratinocytes and the

extracellular matrix[54]. Epidermal keratinocytes interact with extracellular matrix components such as

proteolytic fragments or cleavage sites (neoepitopes) in the surrounding stroma, which affects

epidermal homeostasis and keratinocyte phenotypes[12, 54, 55]. There is emerging evidence that

keratinocytes in lesional vitiliginous skin are genetically and biologically distinct from normal cells[46,

56, 57, 58]. We observed epidermal thickening in lesional vitiliginous skin in accordance with previous

histological findings[56, 59], but the mechanisms responsible for the fundamental abnormalities in

keratinocytes during vitiligo development remain unknown. Previous studies indicated that under

pathological conditions, extensive BM destruction results in enhanced release of cytokines in the

epidermis[54, 55, 60]. We found that disruption of the BMZ architecture altered the expression pattern of

adhesive molecules in basal keratinocytes, and the cytokines IL-1β, TGF-β1, TNF-α, and IL-6 were

highly expressed in the epidermis of vitiliginous skin lesions (**Extended Data Fig 1**). This cytokine

expression from epidermal keratinocytes might upregulate MMP2 expression in dermal fibroblasts

through paracrine regulation.

We also detected elevated MMP2 expression in dermal fibroblasts in stable vitiligo. We believe that

there may be a vicious cycle in which MMP2 destroys BM, and the consequent disruption of the BM

alters the phenotype of keratinocytes, resulting in cytokine production, which in turn upregulates

MMP2 expression in fibroblasts. When the normal BM healing process is interrupted, a pathological

vicious cycle of cytokine and protease release occurs, leading to the development of a permanent and

intractable pathological skin phenotype, such as stable vitiligo. The use of skin needling and laser

ablation in combination with phototherapies or topical medicines such as steroids, tacrolimus, and 5-

fluorouracil to treat stable vitiligo has shown satisfying results[61, 62, 63]. Stimulation of the dermis by

needling or laser ablation may induce biological changes in dermal fibroblasts, BM, and keratinocytes

that halt the pathogenic vicious cycle and repair the BMZ. The vicious cycle described here may

explain why melanocytes cannot return to the BM for a long time after they disappear, but it remains

unclear what triggers the initial event in the cycle. As the body's first line of defense, the NLRP3 and

NLRP1 inflammasomes are activated in response to various external stimuli, such as oxidative stress,

UV radiation, chemical agents, extracellular ATP, and microbial dysbiosis, and the cytokines IL-1β,

IL-6, and TNF-α consequently become elevated in the epidermis[64, 65]. The infiltrating lymphocytes

and other inflammatory cells that have been detected in the early initial stage of vitiligo may also be

capable of producing these cytokines. Ultimately, the cytokines may induce MMP2 expression in

dermal fibroblasts and initiate the vicious cycle of vitiligo development. Nonetheless, further research

is required to test this hypothesis.

We found that MMP2 expression was increased in protein extracts of full skin from vitiligo lesions

compared with that in protein extracts of healthy skin. The MMP2 in the skin extracts was mainly the

active 62-kDa form, indicating that the latent (pro) form of MMP2 in the dermis of lesional skin is

activated locally. Pro-MMP2 is known to be activated by TIMP-2, MT1-MMP, organomercurials, and

several proteases, including MMP7 and MMP3[22]. Further investigation is needed to determine whether

or not these are involved in the MMP2 activation in lesional vitiliginous skin.

Therapeutically, restoring the BM is the most important issue for melanocyte reinhabitation.

Pharmacologic interventions (e.g., anti-inflammatory medications or selective inhibitors of signaling

pathways) or molecular biological interventions (e.g., natural MMP2 inhibitors, monoclonal

antibodies, or siRNAs) that seek to modulate the expression or activation of MMP2 may have an

impact on the prevention and treatment of vitiligo. For example, steroids and selective inhibitors of

the IL-1β, TGF-β, TNF-α, and IL-6 signaling pathways may be partially effective for vitiligo treatment.

JAK inhibitors not only inhibit the production of cytokines by infiltrating $CD8^+$ T lymphocytes but

also have the ability to block some of the cytokine pathways in fibroblasts, which may make them

effective in vitiligo treatment. However, to successfully cure vitiligo, MMP2 expression should be

restored to a normal level. The use of MMP2 inhibitors, which are currently being evaluated in clinical

trials, is a promising therapeutic option for vitiligo; however, the systemic administration of MMP2

inhibitors may serve as a double-edged sword that promotes tissue healing and disturbs melanocyte

homeostasis. Therefore, it is crucial to suppress MMP2 with a selective inhibitor at an appropriate

dosage and preferably via local administration (e.g., topical application) to reduce potential side effects.

In summary, fibroblasts in the dermis of lesional vitiliginous skin express high levels of MMP2,

which disrupts the BM architecture, leading to a breakdown of the integrin β–laminin interactions that

provide the main adhesion between melanocytes and the BM. As a result, melanocytes detach from

the BM and disappear from the epidermis. The impairment of BM structure may lead to functional

changes in epidermal cells and induce keratinocyte abnormality, promoting melanocyte disappearance

by stimulating dermal fibroblasts, as summarized in **Fig 9**. These findings shed new light on the

pathogenesis of vitiligo. It appears that a disorganized BM caused by MMP2 from dermal fibroblasts

is responsible for the loss of melanocytes in vitiligo. Our *in vivo* results suggest that MMP2

suppression may have therapeutic implications for vitiligo; however, the MMP2-Fb–induced

depigmentation in 3D skin equivalents, *ex vivo* skin explants, and a mouse model was inadequate to

examine chronic changes in epidermal keratinocytes. Another limitation of this study is that only

MMP2 and MMP9 were examined. Other proteinases and endoglycosidases from dermal cells, such

as macrophages and mast cells, have the potential to degrade the BMZ, and it is possible that cytokines

and chemokines from abnormal keratinocytes alter these cells, thereby indirectly affecting the

phenotype of fibroblasts. Further research is needed to discover whether additional factors and cells

contribute to the BMZ disruption in vitiligo.

Figure Legends

Figure 1. Disorganization of basement membrane zone architecture in vitiligo

a, Representative ultrastructural images of ultrathin sections of skin from five healthy controls and

five patients with generalized vitiligo. In the right panel, the areas of epidermis and dermis are

represented by Epi and Der, respectively, and the basement membrane zone is indicated by a red line.

In the left panel, the area of the white rectangle is shown enlarged at the lower right. Red asterisk:

anchoring fibril; Epi: epidermis; Der: dermis; BMZ: basement membrane zone. Scale bars: 2 μm. **b,**

Representative immunofluorescence images of skin specimens from five healthy controls and 11

patients with generalized vitiligo stained with anti-collagen IV (green), anti-TRP1 (red), and DAPI

(blue). The areas in the white rectangles are shown in the corresponding right panels. Scale bar: 100

μm. **c,** Representative immunoelectron microscopy images of frozen sections stained with anti-

collagen IV. White arrowhead: collagen IV-labeled gold-particle deposits close to the lamina densa;

White arrow: collagen IV-labeled gold-particle deposits close to the branched structure of lamina

densa; White asterisk: collagen IV-labeled gold-particle deposits close to the layered structure of

lamina densa. Scale bars: 1 μm.

Figure 2. Matrix metalloproteinase 2 expression is elevated in the dermis of

430 **vitiliginous skin**

431 **a,** Representative immunofluorescence images of frozen sections of skin from five healthy controls

432 and 11 patients with generalized vitiligo. MMP2 is stained green, and MMP9 is stained red. Vitiligo-

433 L: lesional vitiliginous skin. **b,** Quantitation of MMP2 and MMP9 expression by fluorescence intensity

434 measurement. **c,** Representative immunofluorescence staining of MMP2, HSP47, and CD31 in skin

435 from healthy controls and patients with vitiligo. **d,** Quantitation of fibroblast and endothelial cell

436 numbers in healthy and vitiliginous skin. **e,** Quantitation of MMP2$^+$ fibroblast and MMP2$^+$ endothelial

437 cell numbers. **f,** The ratios of MMP2$^+$ fibroblasts among all fibroblasts and MMP2$^+$ endothelial cells

438 among all endothelial cells. **g,** Representative immunofluorescence staining of MMP2$^+$ fibroblasts

439 distributed around the BM. MMP2 is stained red; the fibroblast marker HSP47 is stained green; and

440 collagen IV, the main component of BM, is stained blue. White arrowhead: MMP2$^+$ fibroblasts close

441 to the BM; White arrow: MMP2$^+$ fibroblasts passed through the BM. **h,** Western blot analyses showing

442 expression of the pro and active forms of MMP2 in skin lysates from healthy skin and lesional

443 vitiliginous skin. **i,** Quantitation of active MMP2 and total MMP2 (the total of active and pro MMP2)

444 expression according to western blot data. Nuclei are counterstained in blue with DAPI in **a** and **c**.

445 Scale bars in **a** and **c**: 50 μm. Scale bars in **g**: 20 μm.

446 **Figure 3. Overexpression of MMP2 in dermal fibroblasts results in melanocyte**

447 **disappearance in three-dimensional human full-thickness skin equivalent and**

human *ex vivo* cultured skin explants.

a, Schematic flow diagram depicting the reconstructed three-dimensional human skin full-thickness

equivalent model. **b,** Macroscopic images of skin equivalents. **c,** Representative immunofluorescence

staining of the melanocyte marker TRP1 and the BM marker laminin 332 in skin equivalents. TRP1

is stained red, and laminin 332 is stained green. Nuclei are counterstained in blue with DAPI. **d,**

Schematic diagram of the human skin explant culture model. **e,** Representative immunofluorescence

staining of collagen IV and melan-A in cultured human skin explants. Collagen IV is stained blue, and

melan-A is stained green. Intradermally injected fibroblasts are labeled with PKH fluorescent dye in

red. The white rectangle outlines the area of melanocytes and BM, and the detail is shown in the

corresponding lower panel. Scale bars in **c** and **e**: 50 μm. Data in **b**, **d**, **e**, **f,** and **i** are shown as mean ±

standard deviation.

Figure 4. Melanocyte disappearance results from decreased adhesion to the BM in

the three-dimensional skin equivalent model.

a, Representative immunofluorescence staining of TRP1 and cleaved-caspase 3 in three-dimensional

skin equivalents. TRP1 is stained green, and cleaved-caspase 3 is stained red. Mock-Fb indicates skin

equivalent embedded with mock-transfected fibroblasts; MMP2-Fb (MZ) indicates the marginal zone

of skin equivalent embedded with MMP2-overexpressing fibroblasts. Detached melanocytes are

shown by white arrows. **b,** Schematic diagram of the proximity ligation assay and melanocyte antibody

cocktail immunofluorescence co-staining. **c,** The interactions between DDR1 and collagen IV and

between integrin β1 and laminin in skin equivalents are visualized in red fluorescence by proximity

ligation assay. Melanocytes are visualized in green. Nuclei are counterstained in blue with DAPI. The

areas in the white rectangles are shown at the lower right. **d,** Quantitation of the fluorescence intensity

around melanocytes in proximity ligation assays of five skin equivalents. Scale bars in **a** and **c**: 50 μm.

Data in **d** are shown as mean ± standard deviation.

Figure 5. Melanocyte disappearance results from decreased adhesion to the BM in

vitiliginous skin.

a, Representative immunofluorescence staining of TRP1 in healthy skin and perilesional vitiliginous

skin. TRP1 is stained red. Nuclei are counterstained in blue with DAPI. Detached melanocytes are

shown by white arrows. Vitiligo-PL indicates perilesional vitiliginous skin. **b,** Representative

immunofluorescence staining of DDR1, collagen IV, and TRP1 in healthy skin and perilesional

vitiliginous skin. DDR1 is stained green, collagen IV is stained blue, and TRP1 is stained red. The

area in the white rectangle is shown on the lower right. The red lines circling the areas in the enlarged

images indicate melanocytes. **c,** Representative immunofluorescence staining of integrin β1, laminin,

and TRP1 in healthy skin and perilesional vitiliginous skin. Integrin β1 is stained green, laminin is

stained blue, and TRP1 is stained red. The area in the white rectangle is shown on the lower right. **d,**

Schematic diagram of the proximity ligation assay and melanocyte antibody cocktail

immunofluorescence co-staining. **e,** The interactions between DDR1 and collagen IV and between

integrin β1 and laminin are visualized in red fluorescence by proximity ligation assay. Melanocytes

are visualized in green. Nuclei are counterstained in blue with DAPI. The area in the white rectangle

is shown on the lower right. **f,** Quantitation of the immunofluorescence signal around melanocytes in

b and **c**. **g,** Quantitation of the fluorescence intensity around melanocytes in proximity ligation assays

in **e**. **h,** Illustration of melanocyte–BM adhesion in healthy skin and vitiliginous skin. Scale bars in **a,**

b, c, and **e**: 50 μm. Data in **f** and **g** are shown as mean ± standard deviation.

Figure 6. Compensatory enhanced hemidesmosome adhesion in the basal

keratinocytes of vitiliginous skin.

a, Representative immunofluorescence staining of integrin α6, laminin, and TRP1 in healthy skin and

lesional vitiliginous skin. Integrin α6 is stained green, laminin is stained blue, and TRP1 is stained red.

The area in the white rectangle is shown on the lower right. **b,** Representative immunofluorescence

staining of collagen XVII, collagen IV, and TRP1 in healthy skin and lesional vitiliginous skin.

Collagen XVII is stained green, collagen IV is stained blue, and TRP1 is stained red. The area in the

white rectangle is shown on the lower right. **c,** Quantitation of the immunofluorescence signal around

basal keratinocytes in **a** and **b**. **d,** Schematic diagram of the proximity ligation assay. **e,** The interactions

between integrin α6 and laminin and between collagen XVII and collagen IV are visualized in red

fluorescence by proximity ligation assay. Nuclei are counterstained in blue with DAPI. The area in the

white rectangle is shown on the lower right. **f,** Quantitation of the fluorescence intensity around basal

keratinocytes in the proximity ligation assays in **e. g,** Illustration of basal keratinocyte–BM adhesion

in healthy skin and vitiliginous skin. Scale bars in **a, b,** and **e**: 50 μm. Data in **c** and **f** are shown as

mean ± standard deviation.

Figure 7. The decreased adhesion of melanocytes to the BM is due to abnormal BM

architecture.

a, Representative ultrastructural images of ultrathin sections of skin from five healthy controls and

five patients with generalized vitiligo. An enlarged image of the area indicated by the red arrowhead

is shown on the lower right. The red double arrow indicates the lamina lucida. Epi: epidermis; Der:

dermis. Scale bars: 1 μm. **b,** Representative immunoelectron microscopy images of ultrathin skin

sections stained with anti-laminin antibody. The area of the lamina lucida is outlined in yellow. Scale

bars: 1 μm. **c,** Quantitation of the width of the lamina lucida in **a**. Three measurements were performed

per section. **d,** Quantitation of the ratio of the number of laminin gold particles located at the lamina

lucida to the length of the lamina lucida. **e,** Illustration of melanocyte–BM adhesion in healthy skin

and vitiliginous skin. Data in **c** and **d** are shown as mean ± standard deviation.

Figure 8. Subcutaneous inoculation of mice with MMP2-overexpressing fibroblasts

induced a vitiligo phenotype that could be reversed by inhibition of MMP2

a, Schematic diagram of the subcutaneous fibroblast inoculation mouse model. **b,** Macro appearance

of the mice at 3 weeks after fibroblast inoculation. The bottom panels show the corresponding

threshold binarization images of the back skin of the mice. **c,** Representative immunofluorescence

staining of TRP1, HSP47, and collagen IV in mouse skin sections. TRP1 is stained red, HSP47 is

stained green, and collagen IV is stained blue. The area in the white rectangle is shown in the

corresponding lower panel. Scale bars: 50 μm. **d,** Schematic of the experimental procedure to treat

mice with MMP2 inhibitor. **e,** Macroscopic images of the mice at 2 weeks after subcutaneous

inoculation. The bottom panels show the corresponding threshold binarization images of the back skin

of the mice. **f,** Quantitation of the ratio of depigmentation area on the back skin of the mice. Data in **f**

are shown as mean ± standard deviation.

Figure 9. Summary illustration of MMP2-induced melanocyte disappearance in

vitiligo.

Schematic representation of normal skin and vitiliginous skin. Melanocytes are located at the basal

layer of the epidermis interacting with the BM. In normal skin, laminin is normally distributed on

collagen IV. Basal melanocytes attach to the BM mainly by integrin β1–laminin binding. In

vitiliginous skin, collagen IV is degraded into fragments by the increased MMP2$^+$ dermal fibroblasts,

the laminin distribution is disrupted, and it is difficult for basal melanocytes to attach to the BM

through integrin β1–laminin binding. Melanocytes detach from the BM and disappear from the

epidermis. Although integrin β1–laminin binding and integrin α6–laminin binding are diminished in

basal keratinocytes of vitiliginous skin, collagen XVII–collagen IV binding is enhanced as a

compensatory mechanism.

Materials & **Methods**

Human skin specimens

Histo-immunofluorescence staining was performed with frozen biopsy samples obtained from

lesional and perilesional skin of well-defined patients with non-segmental vitiligo (n = 13, 7 in

progressive state and 4 in stable state) (**Table 1**) and from corresponding sites on healthy donors (n =

5; 2 from the abdomen, 2 from an arm, and 1 from the chest). Three-millimeter punch fresh skin biopsy

specimens of lesional skin from confirmed patients with vitiligo (n = 3) and samples from

corresponding sites on healthy donors (n = 3) were used for protein extraction. One-millimeter punch

skin biopsy specimens of lesional skin from confirmed patients with vitiligo (n = 5) and samples from

corresponding sites on healthy donors (n = 5) were used for electron microscopy and immuno-electron

microscopy analyses. The progressive state of vitiligo was defined by the development of new lesions

or the extension of pre-existing lesions in the last 6 months, and the stable state was defined by no

increase in the size of existing lesions and an absence of new lesions in the past 1 year. Written

informed consent was obtained from all participants prior to study inclusion. The study was approved

by the Medical Ethics Committee of Osaka Metropolitan University (No. 4152).

Fluorescent Immunohistochemistry Staining

Human skin tissue and 3D skin samples embedded in OCT compound (Sakura Fineteck, Tokyo,

Japan) were cut into 6-μm sections for immunohistochemistry analyses. After fixation with 4%

paraformaldehyde (163-20145, Wako, Osaka, Japan) for 10 min and blocking with 5% BSA for 30

min at room temperature, these sections were incubated with 1:100 diluted primary antibodies specific

for collagen IV (ab6586, Abcam, Cambridge, MA, USA), TRP1 (SIG-38150, Bio Legend, San Diego,

CA, USA), MMP2 (ab86607, Abcam), MMP9 (ab119906, Abcam), HSP47 (ADI-SPA-470, Enzo Life

Sciences, Farmingdale, NY, USA), CD31 (M0823, Dako, Santa Clara, CA, USA), laminin 332

(ab78286, Abcam), melan-A (SIG-38160, Bio Legend), cleaved-caspase 3 (#9664, Cell Signaling

Technology, Danvers, MA, USA), DDR1 (PA5-29284, Thermo Fischer Scientific, Waltham, MA,

USA), integrin β1 (ab30388, Abcam), Laminin (ab11575, Abcam), integrin α6 (ab20142, Abcam),

and collagen XVII (HPA043673, Sigma-Aldrich, Burlington, MA, USA) and then with the appropriate

secondary antibody (Alex Fluor 488/555/647 anti-rabbit IgG, anti-mouse IgG1, anti-mouse IgG2a,

anti-mouse IgG2b, or anti-goat IgG; 1:1,000, Invitrogen, Grand Island, NY, USA). Nuclei were

counterstained with DAPI (#62248, Thermo Fischer Scientific). Murine skin tissues embedded in OCT

compound were cut into 6-μm sections for immunohistochemistry analyses. After fixation and

blocking, these sections were incubated with 1:100 diluted primary antibodies specific for TRP1 (SIG-

38150, Bio Legend), HSP47 (ADI-SPA-470, Enzo Life Sciences), and collagen IV (ab19808, Abcam),

and then with the appropriate secondary antibody (Alex Fluor 488 anti-mouse IgG2b, Alex Fluor 555

anti-mouse IgG2a, or Alex Fluor 647 anti-rabbit IgG; 1:1,000, Invitrogen). Slides were observed using

an LSM 710 Zeiss confocal laser microscope. Images were captured using Zeiss ZEN 2012 software

(Carl Zeiss Inc., Jena, Germany). Counting of positively stained cells and measurement of mean

fluorescence intensity were carried out using Image J Software (NIH,

http://rsb.info.nih.gov/ij/index.html).

Electron Microscopy

Punch biopsy skin samples (1 mm) were fixed with 2.5% glutaraldehyde in 0.1 M phosphate-

buffered saline (PBS) at pH 7.4 for 2 h and then washed overnight with PBS. Samples were then post-

fixed in 1% O_sO_4 (157-01141, Wako) in PBS for 1h. After washing with DDW, samples were

dehydrated with a graded ethanol series and embedded in EPON (TAAB EPON, 342-2, Nisshin EM,

Tokyo, Japan). Ultrathin sections (70–80 nm) were cut and transferred to grids, dual-stained with

uranyl acetate and lead citrate, and visualized on an FEI Talos F200C G2 Transmission Electron

Microscope (Thermo Fisher Scientific). The width of lamina lucida was measured using Image J

Software (NIH, http://rsb.info.nih.gov/ij/index.html).

Pre-Embedding Immunoelectron Microscopy

Pre-embedding gold enhancement immunogold labeling was performed as detailed previously[66]

with a slight modification. Briefly, skin tissues were fixed in 4% paraformaldehyde (163-20145,

Wako) for 30 min at room temperature and embedded in OCT compound. Frozen sections (7 μm) were

permeabilized in 0.25% saponin (47036, Sigma-Aldrich) for 30 min and then blocked for 30 min in

PBS containing 0.1% saponin, 10% bovine serum albumin (013-25773, Wako), 10% normal goat

serum (16210-072, Thermo Fisher Scientific), and 0.1% cold water fish skin gelatin (G7041, Sigma-

Aldrich). Samples were incubated with anti-collagen IV (1:100; ab19808, Abcam) or anti-laminin

(1:100; ab11575, Abcam) overnight at $4°C$ in a blocking solution and then with colloidal gold-

conjugated anti-rabbit IgG (1.4-nm diameter; 1:200; #2003, Nanoprobes, Yaphank, NY, USA) in a

blocking solution for 2 h at room temperature. The signal was intensified with the Gold Enhancement

EM Kit (#2113, Nanoprobes) for 4 min at room temperature. The specimens were post-fixed in 1%

OsO4, dehydrated in a series of graded ethanol solutions, and embedded in EPON (TAAB EPON,

342-2, Nisshin EM). Ultrathin sections (70 nm) were collected and stained with uranyl acetate and

lead citrate. Images were taken with an FEI Talos F200C G2 Transmission Electron Microscope

(Thermo Fisher Scientific) equipped with an Advanced Microscopy Techniques charge-coupled

device-based camera system. The quantitation analyses of positive gold particles were carried out

using Image J Software (NIH, http://rsb.info.nih.gov/ij/index.html).

Western Blot Analyses

For the preparation of protein samples, punch fresh skin biopsy specimens were extracted as

described previously[67]. Five micrograms of extracted protein was used for western blot analysis with

primary antibody against MMP2 (1:200; ab92536, Abcam). The signal intensity of bands was

quantified using the Image J densitometry software (NIH).

Cell Culture and Establishment of MMP2-Overexpressing Cells

Normal human dermal fibroblasts (Neonatal, Primary, FC-0001, Lifeline Cell Technology, Carlsbad,

CA, USA) were maintained in MEM (M4655, Sigma-Aldrich) with 1% NEAA (139-15651, Wako),

5% FBS (10270-106, Gibco, Thermo Fisher Scientific), 1 mM Sodium Pyruvate (11360-070, Gibco,

Thermo Fisher Scientific), and 1% penicillin/streptomycin (168-23191, Wako). C57BL/6 mouse

dermal fibroblasts (Neonatal, Primary, SCR-M230057, ScienCell Research Laboratories, Carlsbad,

CA, USA) were maintained in Fibroblast Medium-2 (2331, ScienCell Research Laboratories). Cells

were incubated at 37°C in a humidified incubator supplied with 5% CO_2.

Normal human dermal fibroblasts and C57BL/6 mouse dermal fibroblasts (2×10^6 cells/well on 10-

cm plates) were transfected with 20 μg human MMP2 expression or murine MMP2 expression

episomal vector DNA or empty episomal vector DNA (customized pEBMulti-Puro, Fujifilm Wako

Pure Chemical Corporation, Osaka, Japan) mixed with P3000 and Lipofectamine 3000

(Lipofectamine™ 3000, Thermo Fisher Scientific). Successfully transfected cells were selected with

1 μg/ml puromycin according to the manufacturer's instructions. The expression of the MMP2 proteins

in cell lysate and culture medium were confirmed by western blot analyses using antibody against

MMP2 (ab92536, Abcam).

Reconstructed Three-Dimensional Human Full-Thickness Skin Equivalents

A reconstructed 3D human full-thickness skin equivalent model was generated as described

previously[68] with modification. Collagen gels were prepared from an acid collagen solution (IAC-50,

Koken Ltd., Tokyo, Japan) according to the manufacturer's protocol. Then, 2.5×10^5 fibroblasts were

cultured on collagen gel in MEM (M4655, Sigma-Aldrich) with 1% NEAA (139-15651, Wako), 5%

FBS (10270-106, Gibco, Thermo Fisher Scientific), 1 mM sodium pyruvate (11360-070, Gibco,

Thermo Fisher Scientific), and 1% penicillin/streptomycin (168-23191, Wako). This medium was

changed every other day after fibroblast mounting. Two days after the dermal component was prepared,

a cell suspension of 8.4×10^5 keratinocytes and 1.2×10^5 melanocytes in 60 μl medium composed of

a 1:1 mixture of Epilife and Medium 254 was seeded onto the concave surface of the contracted gel.

After 30 min, 2 ml culture medium was added to the inside and outside of the transwell, and the

submerged culture was maintained for 2 days. The culture medium was composed of DMEM/F-12

(3:1 vol/vol; D5921, N4888, Sigma-Aldrich) containing 0.3% FCS, 0.1 mM ethanolamine, 0.1 mM *o*-

phosphoethanolamine, 0.4 μg/ml hydrocortisone, 5 μg/ml insulin, 0.09 mM adenine, 5 μg/ml

transferrin, 20 pM triiodothyronine, 0.64 mM choline chloride, 0.675 mM serine, 20 μg/ml linoleic

acid/BSA, 50 μg/ml ascorbic acid, 7.25 mM glutamine, 10 nM progesterone, 100 I.U./ml penicillin,

and 100 μg/ml streptomycin. The thus-formed early skin equivalents were then raised to the air-liquid

interface and placed in cornification medium composed of a 1:1 mixture of Ham's F-12 and DMEM

supplemented with 0.3% FCS, 0.1 mM ethanolamine, 0.1 mM *o*-phosphoethanolamine, 0.4 μg/ml

hydrocortisone, 5 μg/ml insulin, 0.09 mM adenine, 5 μg/ml transferrin, 20 pM triiodothyronine, 0.64

mM choline chloride, 0.75 mM serine, 20 μg/ml linoleic acid/BSA, 50 μg/ml ascorbic acid, 7.25 mM

glutamine, 0.75 mM calcium chloride, 100 I.U./ml penicillin, and 100 μg/ml streptomycin. The

medium was subsequently changed every other day. One week after the air-liquid interface incubation,

the skin equivalents were harvested and subjected to macroscopic examination and further histology

analyses. All skin equivalent experiments were performed in quintuplicate in five transwells.

To prepare the half-half 3D human skin equivalent, MMP2-overexpressing fibroblasts embedded in

collagen gel were incubated at 37°C for 30 min. The early-formed dermal tissue construct was then

cut into half size and transferred to one side of a culture transwell insert, and freshly prepared collagen

solution containing normal fibroblasts was injected into the other side of the transwell insert. After 2

days of culture, keratinocytes and melanocytes were seeded as described above.

Ex Vivo Human Skin Explant Culture

Skin from the lateral abdomen was obtained with the informed consent of patients undergoing

abdomen reduction surgery, in accordance with the ethical guidelines of the Research Ethics

Committee of Osaka Metropolitan University (No. 4152). The explants were cut (1.0 cm^2) and cultured

at an air-liquid interface on culture inserts (filter pore size 8 μm; #141082, Thermo Fisher Scientific)

with a 1:1 mixture of Ham's F-12 and DMEM supplemented with 0.3% FCS, 0.1 mM ethanolamine,

0.1 mM o-phosphoethanolamine, 0.4 μg/ml hydrocortisone, 5 μg/ml insulin, 0.09 mM adenine, 5

μg/ml transferrin, 20 pM triiodothyronine, 0.64 mM choline chloride, 0.75 mM serine, 20 μg/ml

linoleic acid/BSA, 50 μg/ml ascorbic acid, 7.25 mM glutamine, 0.75 mM calcium chloride, 100

I.U./ml penicillin, and 100 μg/ml streptomycin. The insert was set on a 12-well plate (3401, Corning

Inc., Corning, NY, USA) and cultured at 37°C in a humidified incubator with 5% CO_2 for 7 days.

For fibroblast inoculation, normal fibroblasts and MMP2-overexpressing fibroblasts were labeled

with the red fluorescent dye PKH26 (MINI26-1KT, Sigma-Aldrich) according to the manufacturer's

instructions and then subcutaneously injected into cultured human skin explants at a density of 5×10^6

cells/ml in 100 μl using disposable 35G needles (ReactSystem, Osaka, Japan).

In Situ **Proximity Ligation Assay**

Multicolor *in situ* PLA was performed using the Duolink PLA Multicolor detection kit (Duolink®

PLA® Multicolor Reagent Pack, DUO96000, Sigma-Aldrich) as described previously[69] with slight

modification. Briefly, frozen skin sections were blocked with blocking solution (DUO82007, Duolink,

Sigma-Aldrich) for 30 min at 37 °C and then incubated overnight at 4°C with primary antibodies pre-

conjugated with Red Oligo A (DUO86010A, Sigma-Aldrich) or Red Oligo B (DUO86010B, Sigma-

Aldrich). After washing with Duolink *In Situ* Wash Buffer A (DUO82046, Sigma-Aldrich), ligation

and amplification reactions were performed using a Duolink PLA Multicolor detection kit (DUO96000,

Sigma-Aldrich).

For co-staining with melanocyte marker after amplification and washing, sections were incubated

with the Alexa Fluor 488-conjugated antibody cocktail (MART1 + Tyrosinase + gp100) [Alexa Fluor®

488] (1:100 dilution; NBP2-34681AF488, Novus Biologics, San Jose, CA, USA) for 2 h. After 30 min

of washing with Duolink *In Situ* Wash Buffer B (DUO82048, Sigma-Aldrich), sections were mounted

using Duolink *In Situ* Mounting Medium with DAPI (DUO82040, Sigma-Aldrich). Then, slides were

observed using an LSM 710 Zeiss confocal laser microscope. Images were captured using Zeiss ZEN

2012 software (Carl Zeiss Inc., Jena, Germany). Mean fluorescence intensity measurements were

carried out using Image J Software (NIH, http://rsb.info.nih.gov/ij/index.html).

Animals and *In Vivo* Experimental Procedures

Krt14-Kitl mice, a C57BL/6-background congenic strain that expresses membrane-bound stem

cell factor under the keratin14 promoter to retain epidermal melanocytes and thus have pigmented

skin, were purchased from the RIKEN BioResource Research Center (No. RBRC00694, RIKEN BRC,

Tsukuba, Ibaraki, Japan). PBS, Mock-Fbs, or MMP2-Fbs (5×10^6 cells/ml in 100 µl PBS) was

subcutaneously injected into the back skin of male 7-week-old Krt14-Kitl mice.

For treatment, 5 µM SB-3CT dissolved in DMSO (HY-12354, MedChemExpress, Monmouth

junction, NJ, USA) or 5 µM Tanomastat dissolved in DMSO (T006900, Toronto Research Chemicals,

Toronto, Canada) was subcutaneously co-injected with Mock-Fbs or MMP2-Fbs (5×10^6/ml in 100

µl PBS) into the back skin of Krt14-Kitl mice. Two weeks after these treatments, the back skin of the

mice was depilated and photographed, and the depigmentation area (n = 5 mice per group) was

measured using binarization and different global thresholding values (Image J Software, NIH,

http://rsb.info.nih.gov/ij/index.html). Experiments were conducted in accordance with the Guiding

Principles for the Care and Use of Laboratory Animals, and the experimental protocol used in this

study was approved by the Committee for Animal Experiments at Osaka Metropolitan University

(Permit No. 18042, Osaka, Japan).

Statistical Analysis

Experiments were repeated at least three times. Data are presented as the mean ± standard

deviation. One-way analysis of variance with Dunnet's test or unpaired Student's t-test (two-tailed)

was used for statistical analyses. A value of $p < 0.05$ indicated statistical significance. All statistical

analyses for this study were performed using GraphPad Prism version 8.0.0 for Windows (GraphPad

Software Inc., San Diego, CA, USA).

Table 1. Clinical characteristics of the patients.

Patient No.	Sex	Age (years)	Type	Biopsy site	Activity
1[§]	M	72	Non-segmental vitiligo	Back	Progressive
2[§]	F	49	Non-segmental vitiligo	Lumbar	Stable
3[*]	M	44	Non-segmental vitiligo	Back	Progressive
4[*]	F	33	Non-segmental vitiligo	Axilla	Progressive
5[*]	F	71	Non-segmental vitiligo	Elbow	Stable
6[§]	F	23	Non-segmental vitiligo	Abdomen	Stable
7[§]	M	30	Non-segmental vitiligo	Lateral abdomen	Progressive
8[§]	F	20	Non-segmental vitiligo	Abdomen	Progressive
9	M	68	Non-segmental vitiligo	Back	Progressive
10	F	49	Non-segmental vitiligo	Abdomen	Progressive
11	F	62	Non-segmental vitiligo	Neck	Stable

713 *Samples for full skin protein extract were obtained from these three patients.

714 §Samples for electron microscopy were from these five patients.

References

1.   Le Poole IC, van den Wijngaard RM, Westerhof W, Dutrieux RP, Das PK. Presence or
absence of melanocytes in vitiligo lesions: an immunohistochemical investigation. *J Invest*
Dermatol **100**, 816-822 (1993).

2.   Picardo M*, et al.* Vitiligo. *Nat Rev Dis Primers* **1**, 15011 (2015).

3.   Alikhan A, Felsten LM, Daly M, Petronic-Rosic V. Vitiligo: a comprehensive overview
Part I. Introduction, epidemiology, quality of life, diagnosis, differential diagnosis,
associations, histopathology, etiology, and work-up. *J Am Acad Dermatol* **65**, 473-491
(2011).

4.   Katz EL, Harris JE. Translational Research in Vitiligo. *Front Immunol* **12**, 624517 (2021).

5.   Lerner AB. Vitiligo. *J Invest Dermatol* **32**, 285-310 (1959).

6.   Bergqvist C, Ezzedine K. Vitiligo: A Review. *Dermatology* **236**, 571-592 (2020).

7.   Cichorek M, Wachulska M, Stasiewicz A, Tyminska A. Skin melanocytes: biology and
development. *Postepy Dermatol Alergol* **30**, 30-41 (2013).

8.   Haass NK, Herlyn M. Normal human melanocyte homeostasis as a paradigm for
understanding melanoma. *J Investig Dermatol Symp Proc* **10**, 153-163 (2005).

9.   Santiago-Walker A, Li L, Haass NK, Herlyn M. Melanocytes: from morphology to
application. *Skin Pharmacol Physiol* **22**, 114-121 (2009).

10.   Le Poole IC, van den Wijngaard RM, Westerhof W, Das PK. Tenascin is overexpressed in
vitiligo lesional skin and inhibits melanocyte adhesion. *Br J Dermatol* **137**, 171-178 (1997).

11.   Costell M*, et al.* Perlecan maintains the integrity of cartilage and some basement
membranes. *J Cell Biol* **147**, 1109-1122 (1999).

12.   Erickson AC, Couchman JR. Still more complexity in mammalian basement membranes.
J Histochem Cytochem **48**, 1291-1306 (2000).

13.   Lin CQ, Bissell MJ. Multi-faceted regulation of cell differentiation by extracellular matrix.
FASEB J **7**, 737-743 (1993).

14.   Walakovits LA, Moore VL, Bhardwaj N, Gallick GS, Lark MW. Detection of stromelysin
and collagenase in synovial fluid from patients with rheumatoid arthritis and
posttraumatic knee injury. *Arthritis Rheum* **35**, 35-42 (1992).

15.   Checchi V*, et al.* The Role of Matrix Metalloproteinases in Periodontal Disease. *Int J*
Environ Res Public Health **17**, (2020).

16.   Saarialho-Kere UK, Kovacs SO, Pentland AP, Olerud JE, Welgus HG, Parks WC. Cell-
matrix interactions modulate interstitial collagenase expression by human keratinocytes
actively involved in wound healing. *J Clin Invest* **92**, 2858-2866 (1993).

17.   Fawzy MM, El Maadawi ZM, Hegazy RA, El Fatah NS. Vitiligo - The story from within:
A transmission electron microscopic study before and after narrow-band ultraviolet B.
Ultrastruct Pathol **40**, 265-275 (2016).

18.   Montes LF, Abulafia J, Wilborn WH, Hyde BM, Montes CM. Value of histopathology in
vitiligo. *Int J Dermatol* **42**, 57-61 (2003).

19.   Panuncio AL, Vignale R. Ultrastructural studies in stable vitiligo. *Am J Dermatopathol* **25**,
16-20 (2003).

20.   Breathnach AS, Bor S, Wyllie LM. Electron microscopy of peripheral nerve terminals and
marginal melanocytes in vitiligo. *J Invest Dermatol* **47**, 125-140 (1966).

21.   Has C, Nystrom A. Epidermal Basement Membrane in Health and Disease. *Curr Top*
Membr **76**, 117-170 (2015).

22.   Nagase H, Woessner JF, Jr. Matrix metalloproteinases. *J Biol Chem* **274**, 21491-21494
(1999).

23.   Yong VW, Krekoski CA, Forsyth PA, Bell R, Edwards DR. Matrix metalloproteinases and
diseases of the CNS. *Trends Neurosci* **21**, 75-80 (1998).

24.   Ram M, Sherer Y, Shoenfeld Y. Matrix metalloproteinase-9 and autoimmune diseases. *J*
788   *Clin Immunol* **26**, 299-307 (2006).

790   25.   Song J, *et al.* Focal MMP-2 and MMP-9 activity at the blood-brain barrier promotes
791   chemokine-induced leukocyte migration. *Cell Rep* **10**, 1040-1054 (2015).

793   26.   Agrawal S, *et al.* Dystroglycan is selectively cleaved at the parenchymal basement
794   membrane at sites of leukocyte extravasation in experimental autoimmune
795   encephalomyelitis. *J Exp Med* **203**, 1007-1019 (2006).

797   27.   Leppert D, *et al.* Matrix metalloproteinase-9 (gelatinase B) is selectively elevated in CSF
798   during relapses and stable phases of multiple sclerosis. *Brain* **121 (Pt 12)**, 2327-2334
799   (1998).

801   28.   Tchetverikov I, *et al.* Matrix metalloproteinases-3, -8, -9 as markers of disease activity and
802   joint damage progression in early rheumatoid arthritis. *Ann Rheum Dis* **62**, 1094-1099
803   (2003).

805   29.   Konttinen YT, *et al.* Matrix metalloproteinase (MMP)-9 type IV collagenase/gelatinase
806   implicated in the pathogenesis of Sjogren's syndrome. *Matrix Biol* **17**, 335-347 (1998).

808   30.   Liotta LA, Abe S, Robey PG, Martin GR. Preferential digestion of basement membrane
809   collagen by an enzyme derived from a metastatic murine tumor. *Proc Natl Acad Sci U S A*
810   **76**, 2268-2272 (1979).

812   31.   Collier IE, *et al.* H-ras oncogene-transformed human bronchial epithelial cells (TBE-1)
813   secrete a single metalloprotease capable of degrading basement membrane collagen. *J Biol*
814   *Chem* **263**, 6579-6587 (1988).

816   32.   Kobayashi T, Hattori S, Shinkai H. Matrix metalloproteinases-2 and -9 are secreted from
817   human fibroblasts. *Acta Derm Venereol* **83**, 105-107 (2003).

819   33.   Wang L, *et al.* Matrix metalloproteinase 2 (MMP2) and MMP9 secreted by
820   erythropoietin-activated endothelial cells promote neural progenitor cell migration. *J*
821   *Neurosci* **26**, 5996-6003 (2006).

823   34.   Zambruno G, Marchisio PC, Melchiori A, Bondanza S, Cancedda R, De Luca M.
824   Expression of integrin receptors and their role in adhesion, spreading and migration of
825   normal human melanocytes. *J Cell Sci* **105 (Pt 1)**, 179-190 (1993).

827   35.   Fukunaga-Kalabis M*, et al.* CCN3 controls 3D spatial localization of melanocytes in the
828   human skin through DDR1. *J Cell Biol* **175**, 563-569 (2006).

830   36.   Ricard AS*, et al.* Study of CCN3 (NOV) and DDR1 in normal melanocytes and vitiligo
831   skin. *Exp Dermatol* **21**, 411-416 (2012).

833   37.   Reichert-Faria A, Jung JE, Moreschi Neto V, de Castro CC, Mira MT, Noronha L.
834   Reduced immunohistochemical expression of Discoidin Domain Receptor 1 (DDR1) in
835   vitiligo skin. *J Eur Acad Dermatol Venereol* **27**, 1057-1059 (2013).

837   38.   Gauthier Y, Cario-Andre M, Lepreux S, Pain C, Taieb A. Melanocyte detachment after
838   skin friction in non lesional skin of patients with generalized vitiligo. *British Journal of*
839   *Dermatology* **148**, 95-101 (2003).

841   39.   Wagner RY*, et al.* Altered E-Cadherin Levels and Distribution in Melanocytes Precede
842   Clinical Manifestations of Vitiligo. *J Invest Dermatol* **135**, 1810-1819 (2015).

844   40.   Verrando P, Hsi BL, Yeh CJ, Pisani A, Serieys N, Ortonne JP. Monoclonal antibody GB3,
845   a new probe for the study of human basement membranes and hemidesmosomes. *Exp Cell*
846   *Res* **170**, 116-128 (1987).

848   41.   Kahari VM, Saarialho-Kere U. Matrix metalloproteinases and their inhibitors in tumour
849   growth and invasion. *Ann Med* **31**, 34-45 (1999).

851   42.   Boukhedouni N*, et al.* Type-1 cytokines regulate MMP-9 production and E-cadherin
852   disruption to promote melanocyte loss in vitiligo. *JCI Insight* **5**, (2020).

854   43.   Kumar R, Parsad D, Kanwar AJ, Kaul D. Altered levels of Ets-1 transcription factor and
855   matrix metalloproteinases in melanocytes from patients with vitiligo. *Br J Dermatol* **165**,
856   285-291 (2011).

44.   Lei TC, Vieira WD, Hearing VJ. In vitro migration of melanoblasts requires matrix
metalloproteinase-2: implications to vitiligo therapy by photochemotherapy. *Pigment Cell*
Res **15**, 426-432 (2002).

45.   Xu Z*, et al.* Anatomically distinct fibroblast subsets determine skin autoimmune patterns.
Nature **601**, 118-124 (2022).

46.   Delmas V, Larue L. Molecular and cellular basis of depigmentation in vitiligo patients.
Exp Dermatol **28**, 662-666 (2019).

47.   Kovacs D*, et al.* Vitiligo Skin: Exploring the Dermal Compartment. *J Invest Dermatol* **138**,
394-404 (2018).

48.   Kusano K*, et al.* Regulation of matrix metalloproteinases (MMP-2, -3, -9, and -13) by
interleukin-1 and interleukin-6 in mouse calvaria: association of MMP induction with
bone resorption. *Endocrinology* **139**, 1338-1345 (1998).

49.   Siwik DA, Chang DL, Colucci WS. Interleukin-1beta and tumor necrosis factor-alpha
decrease collagen synthesis and increase matrix metalloproteinase activity in cardiac
fibroblasts in vitro. *Circ Res* **86**, 1259-1265 (2000).

50.   Choi JY, Piao MS, Lee JB, Oh JS, Kim IG, Lee SC. Propionibacterium acnes stimulates
pro-matrix metalloproteinase-2 expression through tumor necrosis factor-alpha in human
dermal fibroblasts. *J Invest Dermatol* **128**, 846-854 (2008).

51.   Overall CM, Wrana JL, Sodek J. Transcriptional and post-transcriptional regulation of 72-
kDa gelatinase/type IV collagenase by transforming growth factor-beta 1 in human
fibroblasts. Comparisons with collagenase and tissue inhibitor of matrix metalloproteinase
gene expression. *J Biol Chem* **266**, 14064-14071 (1991).

52.   Philips N, Keller T, Gonzalez S. TGF beta-like regulation of matrix metalloproteinases by
anti-transforming growth factor-beta, and anti-transforming growth factor-beta 1
antibodies in dermal fibroblasts: Implications for wound healing. *Wound Repair Regen* **12**,
53-59 (2004).

53.   Galis ZS*, et al.* Cytokine-stimulated human vascular smooth muscle cells synthesize a

complement of enzymes required for extracellular matrix digestion. *Circ Res* **75**, 181-189
(1994).
896

897   54.   Yazlovitskaya EM*, et al.* The laminin-binding integrins regulate nuclear factor kappaB-
898   dependent epithelial cell polarity and inflammation. *J Cell Sci* **134**, (2021).
899

900   55.   Breitkreutz D, Koxholt I, Thiemann K, Nischt R. Skin basement membrane: the
901   foundation of epidermal integrity--BM functions and diverse roles of bridging molecules
902   nidogen and perlecan. *Biomed Res Int* **2013**, 179784 (2013).
903

904   56.   Kovacs D*, et al.* Altered epidermal proliferation, differentiation, and lipid composition:
905   Novel key elements in the vitiligo puzzle. *Sci Adv* **8**, eabn9299 (2022).
906

907   57.   Shiu J*, et al.* Multimodal analyses of vitiligo skin identify tissue characteristics of stable
908   disease. *JCI Insight* **7**, (2022).
909

910   58.   Moretti S*, et al.* Keratinocyte dysfunction in vitiligo epidermis: cytokine
911   microenvironment and correlation to keratinocyte apoptosis. *Histol Histopathol* **24**, 849-
912   857 (2009).
913

914   59.   Jung SE, Kang HY, Lee ES, Kim YC. Changes of epidermal thickness in vitiligo. *Am J
915   Dermatopathol* **37**, 289-292 (2015).
916

917   60.   Ryan MC*, et al.* The functions of laminins: lessons from in vivo studies. *Matrix Biol* **15**,
918   369-381 (1996).
919

920   61.   Elshafy Khashaba SA, Elkot RA, Ibrahim AM. Efficacy of NB-UVB, microneedling with
921   triamcinolone acetonide, and a combination of both modalities in the treatment of vitiligo:
922   A comparative study. *J Am Acad Dermatol* **79**, 365-367 (2018).
923

924   62.   Bayoumi W*, et al.* Effect of a preceding laser dermabrasion on the outcome of combined
925   therapy with narrowband ultraviolet B and potent topical steroids for treating
926   nonsegmental vitiligo in resistant localizations. *Br J Dermatol* **166**, 208-211 (2012).
927

928   63.   Ezzedine K, Eleftheriadou V, Whitton M, van Geel N. Vitiligo. *Lancet* **386**, 74-84 (2015).
929

64.   Marie J, *et al.* Inflammasome activation and vitiligo/nonsegmental vitiligo progression. *Br*
J Dermatol **170**, 816-823 (2014).

65.   Moretti S, *et al.* New insights into the pathogenesis of vitiligo: imbalance of epidermal
cytokines at sites of lesions. *Pigment Cell Res* **15**, 87-92 (2002).

66.   Hayashi-Nishino M, Fujita N, Noda T, Yamaguchi A, Yoshimori T, Yamamoto A. A
subdomain of the endoplasmic reticulum forms a cradle for autophagosome formation.
Nat Cell Biol **11**, 1433-1437 (2009).

67.   Yang F, Yang L, Wataya-Kaneda M, Yoshimura T, Tanemura A, Katayama I. Uncoupling
of ER/Mitochondrial Oxidative Stress in mTORC1 Hyperactivation-Associated Skin
Hypopigmentation. *J Invest Dermatol* **138**, 669-678 (2018).

68.   Yang L, *et al.* Microbubble-enhanced ultrasound for gene transfer into living skin
equivalents. *J Dermatol Sci* **40**, 105-114 (2005).

69.   Phoomak C, *et al.* The translocon-associated protein (TRAP) complex regulates quality
control of N-linked glycosylation during ER stress. *Sci Adv* **7**, (2021).

Figure 1

a Transmission electron microscopy (TEM)

Healthy control — Epi, BMZ, Der

Vitiligo-L — Epi, BMZ, Der

b TRP1/Collagen IV/DAPI   Enlarged

Healthy control

Vitiligo-L

c Immunoelectron microscopy (anti-Collagen IV)

Epi, Der — Healthy control   Epi, Der — Vitiligo-L

Figure 2

a
Healthy control | Vitiligo-L   Healthy control | Vitiligo-L
MMP2/DAPI   MMP9/DAPI

b Fluorescence intensity (IntDen/Area)
Healthy control | Vitiligo-L
<0.0001 (MMP2)
0.0003 (MMP9)

c MMP2/DAPI | HSP47/DAPI | CD31/DAPI | MMP2/HSP47/CD31/DAPI
Healthy control
Vitiligo-L

g MMP2/HSP47/Collagen IV
Healthy control — Epi, Der
Vitiligo-L — Epi, Der
Vitiligo-L — Epi, Der

d Number of cells /Area (10⁶ pixel)
Healthy control | Vitiligo-L
<0.0001 (Fibroblasts)
0.1213 (Endothelial cells)

e Number of MMP2⁺ cells /Area (10⁶ pixel)
Healthy control | Vitiligo-L
<0.0001 (Fibroblasts)
<0.0001 (Endothelial cells)

f % of MMP2⁺ cells
Healthy control | Vitiligo-L
<0.0001 (Fibroblasts)
0.2937 (Endothelial cells)

h kD   WB: 5 µg protein (skin extracts)
250, 150, 100, 75, 50, 37
proMMP2
active MMP2
1 2 3   1 2 3
Vitiligo-L   Healthy control

i Integrated Density (x10⁴)
Healthy control | Vitiligo-L
0.0010 (Active MMP2)
0.0009 (Total MMP2)

Figure 3

a

Dermal repopulation

Epidermal proliferation Submerged phase

Epidermal differentiation Air-exposed phase

Seeding MMP2-Fb in collagen gel → Gel contraction → Cutting in half after gel contraction → Transfering the half-gel into a new transwell; adding normal fibroblasts-rich collagen gel → Seeding keratinocytes and melanocytes → Air-liquid interface lifting

Mock-Fb MMP2-Fb Mock-Fb MMP2-Fb

Day 0 — Day 1-Day 2 — Day 2-Day 4 — Day 4-Day 11

Mock-trafected fibroblast (Mock-Fb) | MMP2-OE fibroblast (MMP2-Fb) | Keratinocyte | Melanocyte

b

3D human Pigmented Skin (Macro Image)

Mock-Fb ← → MMP2-Fb

1:3 magnification

1:4.5 magnification

c

Mock-Fb | MMP2-Fb

TRP1/Laminin 332/DAPI

d

PBS

PKH-labeled Mock-Fb

PKH-labeled MMP2-Fb

Healthy donor → Skin biopsy → Intradermal injection → *ex vivo* culture

e

Melan-A/Collagen IV/PKH

PBS | Mock-Fb | MMP2-Fb Field View 1 | Field View 2

Figure 4

a

3D human pigmented skin

Mock-Fb

MMP2-Fb

TRP1/Cleaved-caspase 3/DAPI

Mock-Fb

MMP2-Fb (MZ)

b

Proximity ligation assay: visualizing protein-protein interactions *in situ*

PLA signal
Ligation
Amplication
Detection

Primary antibodies with PLA probes

DDR1 /Integrin β1

Collagen IV /Laminin

10-80 nm

Distance between two target proteins

Co-staining with melanocyte (Alexa Fluor488 conjugated antibody cocktail)

MART-1 Tyrosinase gp100

c

PLA signal/Melanocyte/DAPI

Mock-Fb

Red: DDR1/Collagen IV | Red: Integrin β1/Laminin

MMP2-Fb (MZ)

Red: DDR1/Collagen IV | Red: Integrin β1/Laminin

d

PLA signal around melanocyte

Mock-Fb
MMP2-Fb (MZ)

Integrated Density (units/basal melanocyte)

2×10^5
1.5×10^5
1×10^5
5×10^4

0.0028

0.0021

DDR1 /Collagen IV Complex

Integrin β1 /Laminin Complex

Figure 5

Figure 6

Figure 7

a Transimission electron microscopy (TEM)

Healthy control | Vitiligo-L

b Immunoelectron microscopy (anti-Laminin)

Healthy control | Vitiligo-L | Vitiligo-PL

c Width of lamina lucida (nm)
0.4331
Healthy control | Vitiligo-L

d Number of gold particles located at Lamina lucida /Length of Lamina lucida
<0.0001
0.3280
Healthy control | Vitiligo-L | Vitiligo-PL

e
Healthy control — MC
Vitiligo — MC

ITGβ1 | Laminins
DDR1 | COL IV | BM

Figure 8

a Mock-Fb (fibroblasts with empty vector)
MMP2-Fb (MMP2-OE fibroblasts)

b PBS | Mock-Fb | MMP2-Fb

c TRP1/HSP47/Collagen IV
PBS | Mock-Fb | MMP2-Fb

d MMP2-Fb + MMP2 Inhibitor (SB-3CT; Tanomastat)

e PBS | Mock-Fb | PBS+ | SB-3CT+ | Tanomastat+ | MMP2-Fb

f Depigmentation area (% of total back skin area)
(%)
0.0004
<0.0001
0.0040
0.7845
PBS | Mock-Fb | PBS+MMP2-Fb | SB-3CT+MMP2-Fb | Tanomastat+MMP2-Fb

Figure 9

Normal skin

Perilesional skin   **Lesional skin**

Melanocyte

Basal
Keratinocyte

Melanocyte

Basal
Keratinocyte

🌿 Normal fibroblast   🔵 MMP2⁺ fibroblast   🟣 MMP2   Integrin β1   Integrin α6   Laminin

Melanocyte   Basement Membrane   Collagen IV   Collagen XVII

Supplementary Information

Materials & **Methods**

Fluorescent Immunohistochemistry Staining

Human skin tissue embedded in OCT compound (Sakura Fineteck, Tokyo, Japan) was cut into 6-µm sections for immunohistochemistry analyses. After fixation with 4% paraformaldehyde (163-20145, Wako, Osaka, Japan) for 10 min and blocking with 5% BSA for 30 min at room temperature, these sections were incubated with 1:100 diluted primary antibodies specific for TGF-β1 (ARG66427, Arigobio, Hsinchu, Taiwan), TNF-α (60291-1-Ig, Proteintech, Rosemont, USA), IL-1β (16806-1-AP, Proteintech), and IL-6 (21865-1-AP, Proteintech) and then incubated with the appropriate secondary antibody (Alex Fluor 488/555 anti-rabbit IgG, anti-mouse IgG1, anti-mouse IgG2a; 1:1,000, Invitrogen, Grand Island, NY, USA). Nuclei were counterstained with DAPI (#62248, Thermo Fischer Scientific). Slides were observed using an LSM 710 Zeiss confocal laser microscope. Images were captured using Zeiss ZEN 2012 software (Carl Zeiss Inc., Jena, Germany). Counting of positively stained cells and measurement of mean fluorescence intensity were carried out using Image J Software (NIH, http://rsb.info.nih.gov/ij/index.html).

Figure Legends

Extended Data Fig. 1. Increased cytokine expression in the epidermis of vitiliginous skin

Representative immunofluorescence staining of TGF-β1 (**a**), TNF-α (**b**), IL-1β (**c**), and IL-6 (**d**) in skin sections from healthy controls and lesional skin sections from patients with vitiligo. Nuclei were counterstained in blue with DAPI. The area in the white rectangle is shown on the lower right. Scale bars in **a, b, c**, and **d**: 50 μm.

Extended Data Fig. 1