Control of stereocilia length during development of hair bundles

Jocelyn F. Krey[1], Paroma Chatterjee[1], Julia Halford[1], Christopher L. Cunningham[2],

Benjamin J. Perrin[3], and Peter G. Barr-Gillespie[1,†]

[1]Oregon Hearing Research Center and Vollum Institute, Oregon Health & Science University, Portland,
OR 97239, USA

[2]Pittsburgh Hearing Research Center, University of Pittsburgh School of Medicine, Pittsburgh, PA
15260, USA

[3]Department of Biology, Indiana University-Purdue University Indianapolis, 723 West Michigan Street,
Indianapolis, IN 46202, USA

[†]Corresponding author at gillespp@ohsu.edu

Short title: Control of stereocilia length

Abbreviations: CV, coefficient of variation; F-actin, filamentous actin; Het, heterozygote; IHC, inner hair
cell; KO, knockout; OHC, outer hair cell; P, postnatal day; PSF, point-spread function; SIM, structured
illumination microscopy.

Abstract

Assembly of the hair bundle, the sensory organelle of the inner ear, depends on differential growth of actin-based stereocilia. Separate rows of stereocilia, labeled 1 through 3 from tallest to shortest, lengthen or shorten during discrete time intervals during development. We used lattice structured illumination microscopy and surface rendering of mouse apical inner hair cells to measure stereocilia dimensions during early postnatal development; these measurements revealed a sharp transition at postnatal day 8 between stage III (row 1 and 2 widening; row 2 shortening) and stage IV (final row 1 lengthening and widening). Tip proteins that determine row 1 lengthening did not accumulate simultaneously during stages III and IV; while the actin-bundling protein EPS8 peaked at the end of stage III, GNAI3 peaked several days later—in early stage IV—and GPSM2 peaked near the end of stage IV. To establish the contributions of key macromolecular assemblies to bundle structure, we examined mouse mutants that eliminated tip links ($Cdh23^{v2J}$ or $Pcdh15^{av3J}$), transduction channels ($Tmie^{KO}$), or the row 1 tip complex ($Myo15a^{sh2}$). $Cdh23^{v2J/v2J}$ and $Pcdh15^{av3J/av3J}$ bundles had adjacent stereocilia in the same row that were not matched in length, revealing that a major role of these cadherins is to synchronize lengths of side-by-side stereocilia. Use of the tip-link mutants also allowed us to distinguish the role of transduction from effects of transduction proteins themselves. While levels of GNAI3 and GPSM2, which stimulate stereocilia elongation, were greatly attenuated at the tips of $Tmie^{KO/KO}$ row 1 stereocilia, they accumulated normally in $Cdh23^{v2J/v2J}$ and $Pcdh15^{av3J/av3J}$ stereocilia. These results reinforced the suggestion that the transduction proteins themselves facilitate localization of proteins in the row 1 complex. By contrast, EPS8 concentrates at tips of all $Tmie^{KO/KO}$, $Cdh23^{v2J/v2J}$ and $Pcdh15^{av3J/av3J}$ stereocilia, correlating with the less polarized distribution of stereocilia lengths in these bundles. These latter results indicated that in wild-type hair cells, the transduction complex prevents accumulation of EPS8 at the tips of shorter stereocilia, causing them to shrink (row 2 and 3) or disappear (row 4 and microvilli). Reduced rhodamine-actin labeling at row 2 stereocilia tips of tip-link and transduction mutants suggests that transduction's role is to destabilize actin filaments there. These results suggest that regulation of stereocilia length occurs through EPS8, and that CDH23 and PCDH15 regulate stereocilia lengthening beyond their role in gating mechanotransduction channels.

Character count: 382 characters

Introduction

Sensory hair cells of the inner ear are distinguished by the staircase arrangement of the ~100 stereocilia in their hair bundles, the sensory organelle that converts mechanical stimuli into electrical signals [1, 2]. The staircase architecture enables directional sensitivity of mechanotransduction [3] and increases

transduction sensitivity [4]. In the mature mammalian cochlea, stereocilia of inner hair cells (IHCs) and

outer hair cells (OHCs) are arranged in three rows, albeit with distinct stereocilia lengths and widths [5].

During development, stereocilia in each row differentially widen and lengthen in specific phases [1, 6]. A

complex of GPSM2 and GNAI3, coupled to MYO15A and EPS8 by WHRN, catalyzes lengthening of

stereocilia beyond about a micrometer [7–9]; this complex concentrates exclusively in row 1. These five

proteins undergo phase separation together, and the presence of GPSM2 both enhances phase

separation and promotes F-actin bundling by EPS8 [10, 11], activity that presumably underlies stereocilia

lengthening. Mutations in any of the genes encoding these five proteins leads to hair bundles that have

short stereocilia and a minimal staircase, i.e., only very small changes of stereocilia length in successive

rows [12].

Lengths of the shorter rows of stereocilia are regulated differently. Several other proteins, including

TWF2, EPS8L2, and CAPZB, are found predominantly at row 2 tips [13–15]; these proteins have actin-

filament capping activity, so their presence may slow lengthening. Stereocilia lacking *Capzb* are narrow,

then shorten and disappear [14], suggesting that CAPZB (and its binding partner TWF2) contribute both

to stereocilia widening but also length stability.

Tension in tip links, extracellular filaments that interconnect stereocilia rows, elicits transduction currents

by opening cation-conducting mechanotransduction channels [2, 16]. Transduction regulates the final

dimensions of stereocilia. Blockade of transduction channels elicits stereocilia shortening, suggesting a

dynamic balance between actin polymerization and depolymerization [17]. Establishment of transduction

currents requires the small membrane protein TMIE [18] and either TMC1 or TMC2, two channel-like

proteins [19, 20]. An ordered stereocilia staircase still forms in mouse mutants lacking either *Tmie* or both

Tmc1 and *Tmc2* [18, 19], but row-specific distinctions of stereocilia width and length become more muted

[6, 21]. The presence of transduction-channel proteins also regulates the localization of proteins specific

for row 1 and row 2 [6]. Although row 1 lacks transduction [22], in transduction mutants, GNAI3 and

GPSM2 do not accumulate substantially at row 1 tips in transduction mutants [6]. Significantly,

transduction-channel proteins do localize to row 1 during early postnatal development [23].

Tip links are made up of dimers of CDH23, which project from the side of a taller stereocilium, and dimers

of PCDH15, projecting from tips of short stereocilia [24–26]. The molecular motor MYO7A positions both

molecules and tensions tip links [27–29]. All three of these proteins are members of the Usher I protein

complex [30], and all are necessary for formation of tip links and activation of transduction. Mutant mouse

lines lacking functional genes for *Cdh23* or *Pcdh15* develop ragged staircases, with stereocilia of irregular

lengths but relatively uniform diameters [31–33]. Because tip links gate transduction channels [34],

Cdh23 and *Pcdh15* mutants lose both transduction and membrane tenting [33].

When tip links are under tension, the stereocilia membrane lifts off of the underlying F-actin core, a
phenomenon called membrane tenting [34, 35]; once the membrane is pulled away, actin polymerization
is enhanced, especially on the side of stereocilium where the tip link anchors [17, 36]. This phenomenon
gives row 2 stereocilia a pronounced beveled shape [17, 36]. Tension in tip links thus promotes actin
polymerization in the shorter stereocilia rows, independent of channel gating [37]. Lacking tip links,
shorter stereocilia in homozygous *Cdh23* and *Pcdh15* mutants have rounded tips instead of beveled ones
[33].

F-actin can be depolymerized at stereocilia tips due to the action of the ADF/CFL family members DSTN
and CFL1 [38, 39]; these proteins sever actin filaments, which can lead to dissolution of actin structures
[40]. Proteins in the ADF/CFL family typically associate with WDR1 (AIP1), which is located in stereocilia
[41] and participates in length regulation [38]. DSTN and CFL1 are specifically located at row 2 stereocilia
tips during early postnatal development, and this localization is disrupted in mutants lacking
mechanotransduction [39]. Actin dynamics controlled by DSTN and CFL1 may underlie pruning of the
shortest rows of stereocilia that occurs during hair-bundle maturation [42].

Thus at least five key multimolecular assemblies control stereocilia dimensions: (1) tip links, (2)
transduction channels, (3) the row 1 complex, (4) row 2 cappers, and (5) actin-severing proteins. Each
of these assemblies is activated at different times during development and is responsible for one or more
steps of hair-bundle development. Here we defined with higher precision the changes in stereocilia
dimensions during postnatal development of apical IHCs from postnatal day 0.5 (P0.5) to P21.5.
Moreover, we used mutant mice to define more specifically the role of tip links on bundle development
and the localization of tip-protein complexes. We examined stereocilia dimensions and row protein
localization in *Pcdh15^{av3J}* and *Cdh23^{v2J}* null hair cells, as well as in *Tmie^{KO}* and *Myo15a^{sh2}* null hair cells.
Besides gating transduction channels, we found that the tip-link proteins play an essential role in
regulating stereocilia lengths; they also have a significant impact on accumulation of row-specific proteins
at stereocilia tips, which in turn modulate stereocilia length. These observations complement previous
results and provide us with a more comprehensive molecular understanding of bundle development.

Results

107 **Quantitation of stereocilia actin-core dimensions using lattice structured illumination microscopy**
108
We determined dimensions of stereocilia F-actin cores, labeled with fluorescent phalloidin, which were
imaged in apical IHCs of C57BL/6 mice at precise times during early postnatal development. Our
measurements should reflect native lengths and widths; dimensions of mildly-fixed, phalloidin-labeled
stereocilia are not significantly different from dimensions of live stereocilia labeled with membrane dyes
[43]. We improved resolution over conventional confocal microscopy by using lattice structured

illumination microscopy (lattice SIM) [44–46], which has a point-spread function (PSF) of ~150 nm under

our conditions [46]; typical confocal microscopy PSFs are ~230 nm [47]. We rendered image surfaces

from each phalloidin-stained hair bundle (Fig 1A-F); rendered surfaces are three-dimensional models of

specific structures computed from stacks of images by pre-processing, segmentation, and connected-

component labeling steps. We saved separate surfaces for each row 1, 2, or 3 stereocilium, as well as

for some row 4 stereocilia and apical-surface microvilli (referred together as row 4+ stereocilia). Because

of their relatively large sizes, row 1 and 2 stereocilia could be measured reliably; dimensions of row 3

and 4+ stereocilia and microvilli could also be measured, but required careful choices of rendering

parameters and editing of surfaces after rendering.

Fig 1. **Stereocilia dimensions during development of C57BL/6J apical IHCs.**

A-F, Imaris reconstruction of phalloidin-labeled IHC stereocilia from indicated ages. Stereocilia

surfaces are color-coded according to row; each box is 14 x 35 µm. Imaris reconstructions show

overlapping hair bundles of adjacent cells, presumably an artefact of sample preparation. ***G-N,***

IHC stereocilia dimension measurements using reconstructed stereocilia surfaces. ***G***, Row 1

and 2 stereocilia length during development; center ten stereocilia in each row. Row 1 data were

fit with a line with slope of zero from P0.5-P7.5 followed by an exponential climb from P7.5-

P21.5; row 2 data were fit with an exponential decline. ***H***, Row 1 and 2 stereocilia width (center

ten). In the panel, data for each row were fit with an exponential climb. The data could also be

fit linearly; between P0.5 and P7.5, the slope for row 1 was 0.026 µm · day^{-1} (95% confidence

interval of 0.025-0.026), while the row 2 slope was 0.034 µm · day^{-1} (95% confidence interval of

0.032-0.036). ***I***, Volume per individual row 1 or row 2 stereocilium (center ten). Data were fit with

a linear increase from P0.5-P7.5, following by an exponential climb (row 1) or decline (row 2)

from P7.5-P21.5. ***J***, Row 1 and 2 stereocilia cross-sectional area (center ten). Data were fit with

a linear increase from P0.5-P7.5, followed by an exponential climb (row 1) or decline (row 2)

from P7.5-P21.5. ***K***, Total stereocilia volume per cell using row 1, 2, and 3 stereocilia. Data were

fit with an exponential climb. ***L***, Number of stereocilia per cell. Data were fit with an exponential

decline. ***M***, Length difference between row 1 and row 2 stereocilia in the same column. Data

were fit with a line with slope of zero from P0.5-P7.5, followed by an exponential climb from

P7.5-P21.5. ***N***, Length difference between adjacent (side-by-side) stereocilia in the same row.

Data were fit by exponential declines. For each plot, dimension measurements of stereocilia

from each bundle were averaged to give an individual cell mean; then all means for individual

cells (5-6 cells from each of 3-4 cochleas) were averaged and plotted ± SEM.

For row 1 and row 2, the variability in stereocilia length within each IHC hair bundle (represented by the

coefficient of variation, CV) was high at P0.5 and declined by P7.5 (panel A in S1 Fig). The majority of

length variability at intermediate ages during development was because the peripheral stereocilia in a

row were systematically shorter than the central stereocilia of that row [6], a phenomenon that disappears

by P11 [43] (depicted schematically in panels E-G of S1 Fig). We therefore restricted most of our

measurements to the center ten stereocilia of each row (panel F of S1 Fig), which could be reliably carried

out only with rows 1 and 2. By focusing on these center stereocilia, length variability was reduced

significantly in row 1 and row 2 at all ages except P0.5 and was constant across development (panel B

in S1 Fig). The width CV for each bundle was also significantly lower for row 1 and especially row 2 for
the center ten stereocilia at most ages (panels C-D in S1 Fig).

Stereocilia dimensions during stages III and IV of early postnatal development

Tilney suggested that hair-bundle development in chickens could be divided into four stages [1]. While
these stages are present in apical IHCs of C57BL/6 mice [6], the temporal and spatial resolution of
previous measurements was insufficient to define transition points between the stages. We improved on
those measurements by (a) using the objective measurement approach described above, (b) including a
larger number of samples for each time point, and (c) expanding the number of time points examined
(Fig 1G-N). For length and width measurements, as well as individual stereocilium volumes, we
calculated the mean and CV for the center ten stereocilia of each row of a single bundle; each symbol
(each time point) in Fig 1G-N corresponds to data from 22-24 single hair cells from four different cochleas.
Total stereocilia volume and number measurements were made for all stereocilia in a bundle, also with
a total of 22-24 single-cell measurements from four cochleas.

As previously reported [48], the number of IHC stereocilia decreased between P0.5 and P21.5, eventually
reaching ~17 each of row 1 and row 2 stereocilia (Fig 1L). Row 1 stereocilia length remained constant
between P0.5 and P8.5, then increased exponentially through P21.5, nearly doubling in length over that
time period (Fig 1G). The transition between the flat and exponential regimes was sharp. By contrast,
row 2 lengths were greatest at P0.5; the length then decreased linearly until P8.5, when the length
stabilized (Fig 1G). Row 2 stereocilia widened about 30% faster than row 1 stereocilia (Fig 1H). The
difference in width between the rows at each time point was somewhat less than we previously estimated
using a different quantitation method [6]. The widening of row 1 over the entire developmental period was
fit by an exponential function, while row 2 started with exponential growth and switched to narrowing with
an exponential time course. The more sustained widening in row 1 stereocilia meant that they eventually
became about 20% wider than those in row 2 (Fig 1H).

If new actin filaments were added to a stereocilium at a constant rate, stereocilium cross-sectional area
(but not width) would grow linearly. Indeed, stereocilia cross-sectional area for IHCs increased linearly
between P0.5 and P8.5 (row 1) or P7.5 (row 2); after that time, row 1 area increased exponentially in row
1 and decreased exponentially in row 2 (Fig 1J). Again, the transition between the linear and exponential
regimes occurred sharply around P8 (Fig 1J).

The volume of each center ten row 1 or 2 IHC stereocilium increased linearly until P8.5 (row 1) or P7.5
(row 2); row 1 volume increased exponentially afterwards, while row 2 volume decreased exponentially
(Fig 1I). To measure total stereocilia volume in a hair bundle, we included all stereocilia from rows 1-3;
volume increased exponentially between P0.5 and P21.5 (Fig 1K), with no apparent transitions.

We also measured the difference in length between adjacent IHC stereocilia. The difference in length

between adjacent row 1 and 2 stereocilia pairs—those that are in a single column along the tip-link axis—

was constant at ~1 µm between P0.5 and P7.5, then increased exponentially to over 4 µm by P21.5

(Fig 1M). Unsurprisingly, this growth profile resembled that of row 1 lengthening, and was enhanced by

the minor row 2 shortening over the same period. Immediately adjacent stereocilia (side-by-side) in a row

were very similar in length (Fig 1N). Using the central ten stereocilia of each row, we measured the

absolute value of the difference in length between each pair of adjacent stereocilia, then plotted the mean

and CV for each hair bundle. Row 1 and row 2 stereocilia pairs showed a narrow distribution of length

differences within a bundle, averaging approximately 0.3 µm (Fig 1N).

Using mouse mutants to dissect hair-bundle development

To understand the contribution of CDH23 and PCDH15 to hair-bundle development in IHCs, we

investigated morphological features of hair bundles in postnatal apical IHCs from $Cdh23^{v2J}$ and

$Pcdh15^{av3J}$ mice [32, 49–51]. These mouse lines are referred to here as "tip-link mutants," and they also

lack mechanotransduction because of the loss of tip links [33]. We confirmed that PCDH15

immunoreactivity was lost in $Pcdh15^{av3J/av3J}$ mice and CDH23 immunoreactivity was absent in $Cdh23^{v2J/v2J}$

mice (S2 Fig). To disentangle loss of tip links from loss of transduction, we also evaluated $Tmie^{KO}$ mice,

which represent the "transduction mutant" class; they phenocopy both the $Tmc1;Tmc2$ double knockout

and $Cib2$ knockouts [18, 23]. Phenotypes shared by tip-link and transduction mutants most likely result

from the loss of transduction, while phenotypes specific for tip-link mutants arise from the absence of the

links themselves. We also analyzed $Myo15a^{sh2}$ mice, which represent the "row 1 tip complex mutant"

class; they largely phenocopy $Gpsm2$, $Gnai3$, $Eps8$, and $Whrn$ mutants [9].

We examined hair-bundle structure, stereocilia shape, and presence of interstereocilia links using

scanning electron microscopy of apical IHCs at P8.5, at the end of stage III (Fig 2). In heterozygote

controls from each of the four mouse lines, bundle structure was similar with thick row 1 and 2 stereocilia

and much thinner row 3 stereocilia (Fig 2A-D); moreover, stereocilia length in each row was relatively

uniform. By contrast, in tip-link mutants, while distinct rows 1, 2, and 3 usually could be identified, rows

were less orderly and stereocilia length in each row was irregular (Fig 2E-F). Row 1 and 2 stereocilia

were still relatively thick, and row 3 stereocilia were generally considerably thicker than those of controls

(Fig 2E-F). As previously reported, $Tmie^{KO/KO}$ bundles have stereocilia that are more uniform in length

and width (Fig 2G).

Fig 2. Scanning electron microscopy of mutants showing links and tip profiles.

A-H, Scanning electron micrographs of single IHC hair bundles from P8.5 cochleas of indicated

genotypes. For each genotype, top panel (labeled i) is a profile example and bottom panel (ii) is

a top-down example. Panel widths, 6 µm. Inserts are 0.5 µm wide and are magnified three-fold

on left; arrows indicate tip links. The insets in Ai, Bi, Ci, Di, and Gi all provide examples of

beveled or pointed stereocilia tips; the insets in Ei, Fi, and Hi provide example of rounded tips.

Some links were still present in both $Cdh23^{v2J/v2J}$ and $Pcdh15^{av3j/av3J}$ bundles, but were not

located at the apices of a stereocilia tips. *I-L*, Quantitation of number of stereocilia per hair
bundle from scanning electron micrographs. Rows 1, 2, and 3 separately plotted; additional rows
of stereocilia and microvilli on each cell's apical surface were also counted (row 4+). P values
for unpaired t-tests are indicated.

While tip links were impossible to convincingly distinguish from other links at early developmental ages
using scanning electron microscopy [52], total links in hair bundles were abundant in heterozygote
controls, as well as *Tmie$^{KO/KO}$* and *Myo15a$^{sh2/sh2}$* mutants; by contrast, there were qualitatively fewer total
links in *Cdh23$^{v2J/v2J}$* and *Pcdh15$^{av3J/av3J}$* bundles, as reported previously [33]. Beveled row 2 tips were
common in heterozygote control bundles and in *Tmie$^{KO/KO}$* bundles (Fig 2A-D, G), consistent with tip links
causing membrane tenting and increased actin polymerization [6, 18]. By contrast, row 2 tips of tip-link
mutant stereocilia were rounded (Fig 2E-F), presumably because of the loss of tip links.

We examined scanning electron micrographs to determine the number of stereocilia in each row (Fig 2I-
L). Pooled heterozygotes stereocilia from all four lines had 17.4 ± 0.3 row 1 stereocilia, 18.1 ± 0.3 row 2
stereocilia, 23.1 ± 0.5 row 3 stereocilia, and 51 ± 4 total stereocilia (mean ± SEM; n = 27 cells). The
values of 17-18 row 1 and 2 stereocilia were very close to those estimated by lattice SIM at P7.5 (Fig
1L).

In IHCs homozygous for any of the four mutant alleles, the number of row 1 and row 2 stereocilia did not
change significantly relative to heterozygotes (Fig 2I-L). In tip-link mutants, the number of row 3 stereocilia
decreased significantly (Fig 2I-J); by contrast, the number of stereocilia in row 3 did not change in
Tmie$^{KO/KO}$ hair bundles, although the number of row 4+ stereocilia (stereocilia associated with the bundle
beyond row 3 and microvilli on the apical surface) decreased significantly (Fig 2K).

Increased variability in stereocilia dimensions of tip-link mutants
Although we previously measured stereocilia dimensions in transduction mutants [6], corresponding
dimensions in tip-link mutants have yet to be systematically investigated. We therefore determined length,
width, and volume of mutant IHC stereocilia from rendered lattice SIM images from P7.5, at the end of
stage III, and P21.5, at the end of stage IV (Fig 3A-H). Stereocilia of tip-link mutants were less well
regimented than those in controls, making assignment to rows 1, 2, and 3 more difficult. Proximity to the
bare zone along the lateral edge of a hair bundle was used to assign row 1, with neighboring row
assignments made medially from row 1. For length and width measurements, we calculated the mean
and CV for the central ten stereocilia of each row; these were our single-cell data points (gray symbols).
A cochlea measurement (colored symbols) was derived from mean or CV measurements from 5-6 single
IHCs.

Fig 3. Stereocilia length in IHCs of mutants.
A-H, Imaris reconstruction of phalloidin-labeled stereocilia from *Cdh23^{v2J}* (A-B), *Pcdh15^{av3J}* (C-
D), *Tmie^{KO}* (E-F), and *Myo15a^{sh2}* (G-H) IHCs at P7.5 and P21.5. Stereocilia surfaces are color-

coded according to row; scale bar in A (2 µm) applies to all panels. ***I-P,*** Average stereocilia
length (panels labeled i) and coefficient of variation for length (CV; ii). Length measurements
used reconstructed stereocilia surfaces. Lengths were determined for the center ten stereocilia
of each hair cell, and means for all ten stereocilia were plotted separately for rows 1 and 2 (gray
symbols). The cell means were averaged for each cochlea (5-6 cells per cochlea) and plotted
with colored symbols (4 cochleas per condition) [53]. Nested t-tests were used to compare the
cochlea values for each genotype [54]. P values are indicated and were bolded if <0.010. ***I-J,***
Length and length CV at P7.5 and P21.5 for *Cdh23^{v2J}*. ***M-N,*** Length and length CV at P7.5 and
P21.5 for *Pcdh15^{av3J}*. ***K-L,*** Length and length CV at P7.5 and P21.5 for *Tmie^{KO}*. ***O-P,*** Length and
length CV at P7.5 and P21.5 for *Myo15a^{sh2}*.

At P7.5, average lengths of the central ten stereocilia in rows 1 and 2 changed only modestly between
heterozygote control and *Cdh23$^{v2J/v2J}$* (Fig 3Ii) or *Pcdh15$^{av3J/av3J}$* stereocilia (Fig 3Mi). By contrast, average
CV values for length (within each hair bundle) were significantly higher for mutant stereocilia, especially
those of row 2 (Fig 3Iii and 3Mii). By P21.5, average lengths were still similar between control and mutant
cells (Fig 3Ji and 3Ni); again, CV values were considerably higher for *Cdh23$^{v2J/v2J}$* and *Pcdh15$^{av3J/av3J}$*
mutants, again most notably for row 2 (Fig 3Jii and 3Nii). Thus *Cdh23$^{v2J/v2J}$* and *Pcdh15$^{av3J/av3J}$* mutants
displayed higher variability of lengths of stereocilia in each IHC for rows 1 and 2 than controls, despite
having similar average lengths.

Stereocilia lengths in *Tmie$^{KO/KO}$* mutants at P7.5 were also not significantly different from those of controls
(Fig 3Ki); by P21.5, however, row 1 stereocilia had lengthened considerably less than those of controls
(Fig 3Li). Variability in row 1 and 2 lengths was significantly greater in *Tmie$^{KO/KO}$* stereocilia as compared
to controls (Fig 3Kii, 3Lii), although CV values were much less than those for tip-link mutants.

Stereocilia widths of the central ten stereocilia changed in *Pcdh15$^{av3J/av3J}$*, *Cdh23$^{v2J/v2J}$*, and *Tmie$^{KO/KO}$*
mutants; in each case, rows 1 and 2 widths were similar in controls but row 2 was reduced in width in
mutants (Fig 4Gi-Fi). Width variability increased significantly in tip-link mutants, especially in row 2
(Fig 4Aii-Dii); by contrast, width CV decreased in *Tmie$^{KO/KO}$* mutants (Fig 4Eii-Fii). Row 2 width decreased
in all three mutants, suggesting that transduction promotes widening. However, widths were more
variable in the tip-link mutants, suggesting that CDH23-PCDH15 links between adjacent stereocilia might
also coordinate widening.

Fig 4. Stereocilia width in mutants.
Width measurements used reconstructed stereocilia surfaces. ***A-H,*** Average stereocilia width
(panels labeled i) and coefficient of variation for width (CV; ii) were determined for the center ten
stereocilia of each hair cell (gray symbols), separately for rows 1 and 2. Plotting and statistical
testing were as in Fig 3. ***I-J,*** Width and width CV at P7.5 and P21.5 for *Cdh23^{v2J}*. ***M-N,*** Width
and width CV at P7.5 and P21.5 for *Pcdh15^{av3J}*. ***K-L,*** Width and width CV at P7.5 and P21.5 for
Tmie^{KO}. ***O-P,*** Width and width CV at P7.5 and P21.5 for *Myo15a^{sh2}*.

Adjacent stereocilia length coordination is disrupted in tip-link mutant IHCs

An elevated CV for IHC stereocilia length for tip-link mutants (Fig 3) could arise from systematic variation
in length within a row or from stochastic variation within the row. To probe the source of high CV, we
used Imaris surfaces to measure the differences in length between adjacent (side-by-side) stereocilia of
row 1 or row 2 of $Cdh23^{v2J/v2J}$, $Pcdh15^{av3J/av3J}$, $Tmie^{KO/KO}$, and $Myo15a^{sh2/sh2}$ hair bundles. The average
difference in length was larger in tip-link mutants than in controls at P7.5 (row 1 in Fig 5Ci and row 2 in
Fig 5Ei; compare to Fig 1N), and was particularly large at P21.5 (Fig 5Di, Fi). CV values for the length
differences were high but similar for control and mutant stereocilia (Fig 5Cii-Fii). Because the length
differences were so small, the high CV values may simply reflect these small values relative to a
presumed absolute measurement error of similar magnitude. Altogether, the length-difference results
suggest that length variability is stochastic, and indicated that CDH23 and PCDH15 participate in
coordination of the lengths of adjacent stereocilia, both for row 1 and for row 2.

Fig 5. Adjacent stereocilia length coordination in mutants.

A, Phalloidin-labeled stereocilia from P7.5 $Pcdh15^{av3J}$ IHCs. Adjacent stereocilia in homozygous
mutant hair bundles are irregular in length. Panel widths: 30 μm. **B**, Schematic showing
measurements made. For each pair of adjacent stereocilia in row 1 and, separately, in row 2,
we measured the difference in length between them (side-by-side stereocilia Δ length). For
example, Δ length for the pair indicated in panel B would be |$length_1$ – $length_2$|, or the absolute
value of the difference in lengths. In addition, for each row 1 – row 2 pair in a column, we
measured the difference in length between the two stereocilia (in-column stereocilia Δ length).
The brackets in panel B mark this length difference for the first row 1-row 2 stereocilia pair. **C-**
H, Difference in adjacent stereocilia length for $Cdh23^{v2J}$, $Pcdh15^{av3J}$, $Tmie^{KO}$, and $Myo15a^{sh2}$
mice using reconstructed stereocilia surfaces. Plotting and statistical testing were as in Fig 3. **C-**
D, Difference in length (Δ length) of side-by-side row 1 stereocilia at P7.5 and P21.5. **E-F**,
Δ length of side-by-side row 2 stereocilia at P7.5 and P21.5. **G-H**, Δ length of in-column
(adjacent row 1 and 2) stereocilia at P7.5 and P21.5.

$Tmie^{KO/KO}$ mutant IHCs showed a different pattern for adjacent stereocilia length differences (Fig 5C-F).
At P7.5, adjacent stereocilia length differences were similar in control and mutant hair bundles, both for
adjacent stereocilia in row 1 (Fig 5Ci) or in row 2 (Fig 5Ei) At P21.5, $Tmie^{KO/KO}$ mutants showed much
greater length variability for row 1 stereocilia (Fig 5Di), but not for row 2 (Fig 5Fi). These results suggest
that the lengths of adjacent row 2 stereocilia remain coordinated in $Tmie^{KO/KO}$ mutants, which still have
tip links, although row 1 coordination is reduced.

The difference in length between row 1 and row 2 stereocilia in a column was approximately the same
for tip-link mutants and heterozygotes at P7.5 (Fig 5G) and P21.5 (Fig 5H). By contrast, CV values were
much higher for tip-link mutants at both ages (Fig 5Gii, 5Hii). These results suggest that tip links also
control the relative difference in length between row 1 and row 2, perhaps indirectly through controlling
adjacent stereocilia lengths in the same row.

While *Tmie^(KO/KO)* mutant IHCs had a similar in-column difference in length compared to heterozygotes at
P7.5 (Fig 5G), that difference was much less in mutants than heterozygotes at P21.5 (Fig 5Hi), which
corresponds to the reduced row 1 lengthening seen in these mutants (Fig 3). This reduced difference in
length was accompanied by an increased CV for the difference, perhaps reflecting the reduced
coordination among row 1 stereocilia (Fig 5Hii).

Myo15a interacts genetically with *Pcdh15*
As previously reported, stereocilia of *Myo15a^(sh2/sh2)* IHC hair bundles were very short, and were more
uniform in length and width than *Myo15a^(sh2)*/+ stereocilia (Fig 2G-H). Although they still support
mechanotransduction [12], tips of *Myo15a^(sh2/sh2)* IHCs were rounded (Fig 2H), perhaps because the
minimal staircase prevented tip links from sufficiently tenting membranes off the tips of the shorter rows.
The number of stereocilia in each row of *Myo15a^(sh2/sh2)* bundles did not change appreciably (Fig 2L). As
has been well documented [55], row 1 and 2 stereocilia lengths and widths were significantly reduced for
Myo15a^(sh2/sh2) mutants (Fig 3Oi, Pi; Fig 4Gi, Hi), associated with lower CV values (Fig 3Oii, Pii; Fig 4Gii-
Hii). The adjacent stereocilia length differences for *Myo15a^(sh2/sh2)* at P7.5 (Fig 5C, E) and P21.5 (Fig 5D,
F) were considerably less than those of heterozygote controls. Likewise, the in-column difference in
length for *Myo15a^(sh2/sh2)* mutants was substantially lower than controls at both ages (Fig 5Gi, 5Hi); the CV
for difference in length was increased substantially (Fig 5Hii, 5Hii), which may reflect errors in measuring
these very short stereocilia.

We created double knockout *Pcdh15^(av3J/av3J)*;*Myo15a^(sh2/sh2)* (*Pcdh15;Myo15a* DKO) mice to test for a
genetic interaction between *Pcdh15* and *Myo15a* (Fig 6). In *Pcdh15^(av3J/av3J)*;*Myo15a^(sh2/sh2)* IHC hair
bundles, stereocilia were spread across the apical surface and were packed roughly hexagonally (Fig
6C), albeit not in a precise array as in *Pcdh15^(av3J)*/+;*Myo15a^(sh2/sh2)* (*shaker 2* KO mice). Lengths and widths
of row 1 and 2 *Pcdh15^(av3J/av3J)*;*Myo15a^(sh2/sh2)* stereocilia were similar to those of *Pcdh15^(av3J)*/+;*Myo15a^(sh2/sh2)*
stereocilia (Fig 6D-E), although row 2 stereocilia were significantly shorter in the DKO bundles as
compared to *sh2* KO bundles. *Pcdh15^(av3J/av3J)*;*Myo15a^(sh2/sh2)* stereocilia showed a significant lateral-to-
medial gradient in length, which was less apparent in *Pcdh15^(av3J)*/+;*Myo15a^(sh2/sh2)* bundles. Row 2
stereocilia showed more length and width variation in DKO as compared to *sh2* KO bundles (Fig 6H-I),
but the differences were not large.

Fig 6. *Pcdh15^(av3J)*;*Myo15a^(sh2)* double knockout mice.
A-C, Phalloidin-labeled IHC hair bundles from *Pcdh15^(av3J)*/+;*Myo15a^(sh2)*/+ (Het;Het),
Pcdh15^(av3J)/+;*Myo15a^(sh2/sh2)* (Het;KO), and *Pcdh15^(av3J/av3J)*; *Myo15a^(sh2/sh2)* (KO;KO) cochleas,
Panels labeled i are single x-y planes from image stacks; panels labeled ii are x-z reslices from
those stacks, with the region of the stack indicated with a yellow box in i. ***D-K***, Dimension
measurements using reconstructed stereocilia surfaces. Plotting was as in Fig 3; statistical tests
used one-way ANOVA analysis with the Šidák correction. ***D-E***, Length and CV length
measurements. ***F-G***, Width and width CV measurements. ***H-I***, Difference in length (Δ length)

and CV for Δ length of side-by-side row 1 stereocilia. ***J-K***, Difference in length (Δ length) and

CV for Δ length of in-column (adjacent row 1 and 2) stereocilia.

The difference in length between adjacent stereocilia was similar in DKO hair bundles as compared to

sh2 KO bundles (Fig 6F), with no significant differences in CV (Fig 6J). Likewise, the difference in length

between row 1 and 2 stereocilia in a single column was no different in DKO bundles than in *sh2* KO

bundles (Fig 6G, K). These results suggest that row 1 complex proteins are required to elongate

stereocilia before PCDH15-dependent links can help refine bundle architecture.

Disrupted pruning of stereocilia and microvilli in mutants

Mutations in tip-link genes also impacted the pruning of short rows of IHC stereocilia. In control hair cells,

row 3 stereocilia were obvious at P7.5 (Fig 7A-D) but were much harder to detect by P21.5 (Fig 7E-H).

Stereocilia and microvilli beyond row 3 (row 4+) were present at P7.5 (Fig 7A-D, K) but had nearly

completely disappeared by P21.5 (Fig 7E-H, L). By contrast, not only did the number of stereocilia in

rows 3 and 4+ increase in mutants (Fig 2I-L), but their length and diameter—and thus volume—also

increased in mutants (Fig 7). Row 3 and 4+ stereocilia lengths and widths, measured using Imaris surface

reconstructions, were quite variable since the phalloidin signal in these stereocilia in controls was very

low and therefore challenging to render. Instead, we measured surface volumes, which incorporated both

length and width, as they better reflected the overall F-actin amount in each stereocilium and were more

consistently measured from control IHCs (Fig 7I-L).

Fig 7. Pruning of short stereocilia and microvilli in mutant IHCs.

A-H, Phalloidin labeling of P7.5 (A-D) and P21.5 (E-H) IHC hair bundles (i, heterozygotes; ii,

homozygotes). Arrows indicate extra rows of thicker stereocilia. Panel widths: 20 µm. ***I-***

L, Individual stereocilium volumes (panels labeled i) and total volumes per cell for all stereocilia

in a row (ii). Row 3 (I-J) and row 4+ (K-L) are separately plotted. Volume measurements used

reconstructed stereocilia surfaces. Plotting and statistical testing were as in Fig 3.

In tip-link mutants, the volume of each row 3 stereocilium increased significantly at P7.5 compared to

controls, and that translated into a significant increase in total volume in all stereocilia of row 3 (Fig 7A-

B, I). This increase in individual and total volume for row 3 stereocilia was maintained at P21.5 for

Pcdh15$^{av3J/av3J}$ mutant hair bundles, but not for *Cdh23$^{v2J/v2J}$* mutant bundles (Fig 7E-F, J). This

measurement was one of very few that distinguished the *Cdh23$^{v2J/v2J}$* and *Pcdh15$^{av3J/av3J}$* mutants. Row 4+

individual and total stereocilia volume also increased significantly in tip-link mutant IHCs at P7.5 (Fig 7K),

but was less striking at P21.5 (Fig 7L).

Reduced pruning and increased volume of row 3 and 4+ stereocilia was much more apparent in *Tmie$^{KO/KO}$*

and *Myo15a$^{sh2/sh2}$* hair bundles (Fig 2K-L; Fig 7). As previously reported, *Tmie$^{KO/KO}$* bundles had a

prominent row 3 and often additional rows (Fig 7C, G); *Myo15a$^{sh2/sh2}$* bundles had 4-6 rows of stereocilia

(Fig 7D, H). At P7.5, individual and summed volumes of row 3 (Fig 7Ii-ii) and row 4+ (Fig 7Ki-ii) *Tmie$^{KO/KO}$*

and *Myo15a$^{sh2/sh2}$* stereocilia were substantially and significantly increased over controls; this effect

remained at P21.5, although more dramatically for *Myo15a$^{sh2/sh2}$* than *Tmie$^{KO/KO}$* bundles (Fig 7Ji-ii, Li-ii).

These results showed that pruning of short rows of stereocilia was altered in all mutant bundles.

Total hair-bundle volume does not change in mutants

Total stereocilia volume per hair cell was not significantly different between *Cdh23^{v2J}*/+, *Pcdh15^{av3J}*/+,

Tmie^{KO}/+ heterozygotes and C57BL/6 wild-type mice at P7.5 and P21.5 (Fig 8). Despite the variability in

stereocilia number, length, and width, total stereocilia volumes in *Cdh23$^{v2J/v2J}$*, *Pcdh15$^{av3J/av3J}$*, and

Tmie$^{KO/KO}$ IHC hair bundles were statistically indistinguishable at P7.5 (Fig 8A). At P21.5, differences

between *Cdh23^{v2J}* and *Tmie^{KO}* controls and mutants were modest, albeit statistically significant (Fig 8B).

The similarity in stereocilia volume is broadly consistent with Tilney's proposal that each hair cell has a

constant amount of F-actin in its bundle, but cells deploy it differently depending on other factors [56].

Although *Myo15a* was an outlier at P21.5 (Fig 8B), actin and actin crosslinkers intended for stereocilia

may have been directed instead to cytocauds in the cell body [57].

Fig 8. Total stereocilia volume in mutant IHCs.

A-B, Total stereocilia volume per hair bundle and volume CV for *Cdh23^{v2J}*, *Pcdh15^{av3J}*, *Tmie^{KO}*

and *Myo15a^{sh2}* IHCs. Volume measurements used reconstructed stereocilia surfaces. Plotting

was as in Fig 3; statistical tests used one-way ANOVA analysis with the Šidák correction. ***A***,

P7.5. No significant differences in volume between any of the genotypes. ***B***, P21.5. Modest

differences in total stereocilia volume were seen for *Cdh23^{v2J}*/+ vs. *Cdh23$^{v2J/v2J}$* and *Tmie^{KO}*/+

vs. *Tmie$^{KO/KO}$*; a large difference in volume was seen for *Myo15a^{sh2}* IHCs.

Localization of row 1 tip complex proteins in control and mutant IHCs

Because proteins that localize to row 1 tips control stereocilia length, we examined in greater detail the

appearance of EPS8, GNAI3, and GPSM2 at the tips of C57BL/6 hair bundles (Fig 9) and at mutant

bundle tips (Figs 10-12). EPS8 increased at row 1 and 2 tips during stage III, then showed an initial

decrease and subsequent increase during stage IV (Fig 9I). GNAI3 was relatively low during stage III,

but increased to a peak at P10.5, near the beginning of stage IV (Fig 9J). By contrast, GPSM2 increased

modestly at the end of stage III but shot up substantially during late stage IV, peaking at P15.5 (Fig 9K).

GNAI3 and GPSM2 levels each dropped precipitously by P21.5 (Fig 9J-K). Our results showing

sequential accumulation of EPS8, then GNAI3, and finally GPSM2 at stereocilia tips are not consistent

with the earlier suggestion that the row 1 tip complex assembles simultaneously [9].

Fig 9. Developmental expression of row 1 complex proteins in IHC stereocilia.

A-D, Localization of GPSM2 and EPS8 in P0.5, P7.5, P15.5, and P21.5 IHCs. ***E-F***, Localization

of GNAI3 and EPS8 in P0.5, P7.5, P15.5, and P21.5 IHCs. In A-B and E-F, images are

projections of horizontal slices with partially flattened hair bundles. In C-D and G-H, images are

projections of horizontal slices obtained at the level of row 1 tips only (top) or row 2 tips and row

shafts (bottom). Images in A-D were acquired in the same imaging session and used the same

acquisition, processing, and display parameters; the same holds for images in E-H. Panel

widths: A-H, 30 µm. *I-J*, Quantitation of EPS8, GNAI3, and GPSM2 immunofluorescence at row
1 and 2 tips. Fluorescence was normalized to fluorescence at P7.5 (EPS8 and GNAI3) or P15.5
(GPSM2) to allow comparison between multiple experiments.

As in C57BL/6 stereocilia, in heterozygotes from mutant IHCs, GNAI3, GPSM2, and EPS8 were
concentrated at tips of row 1 stereocilia (Fig 10A-D, I-L). WHRN was also concentrated at row 1 tips in
heterozygotes (Fig 10E-H). In *Pcdh15$^{av3J/av3J}$* IHCs, GNAI3 and GPSM2 remained concentrated at row 1
but EPS8 was seen at all stereocilia tips in similar amounts (Fig 10B, F, J). These results contrasted with
those from *Tmie$^{KO/KO}$* IHCs, where GPSM2 and GNAI3 largely disappeared from hair bundles (Fig 10D,
L); WHRN intensity was reduced, but remained most abundant in row 1 (Fig 10H). EPS8 levels in shorter
rows of *Tmie$^{KO/KO}$* IHCs increased substantially [6], but EPS8 was still most abundant in row 1 (Fig 10D,
H, L).

Fig 10. Row 1 protein localization in mutant IHCs.
A-D, Localization of GNAI3 and EPS8 in *Pcdh15^{av3J}* and *Tmie^{KO}* heterozygotes and
homozygotes at P7.5. *E-H*, Localization of WHRN and EPS8 in *Pcdh15^{av3J}* and *Tmie^{KO}*
heterozygotes and homozygotes at P7.5. *I-L*, Localization of GNAI3 and EPS8 in *Pcdh15^{av3J}*
and *Tmie^{KO}* heterozygotes and homozygotes at P7.5. Arrows in B and D show high levels of row
2 EPS8 but not GNAI3. Images from all pairs of heterozygote and homozygote cochleas (A and
B, C and D, E and F, G and H, I and L) were acquired in the same session and used the same
acquisition, processing, and display parameters. All images are projections of horizontal slices;
projections that allow visualization of the entire hair bundle typically had stereocilia splayed
against the apical surface, especially at P15.5, which resulted in immunofluorescence signal
from stereocilia being overlaid on top of immunofluorescence signal from apical surfaces.
Arrows in B, F, and J show high levels of EPS8 in row 2 stereocilia of *Pcdh15$^{av3J/av3J}$* IHCs.
Asterisks in D and L show reduced levels of GNAI3 and GPSM2 in *Tmie$^{KO/KO}$* IHCs. Panel widths:
30 µm.

Quantitation of row 1 tip complex proteins allowed us to determine the distribution of these proteins
between row 1 and 2 stereocilia more thoroughly (Fig 11-12; S4 Fig). Total EPS8 at stereocilia tips was
not altered in tip-link mutants at P7.5 or P21.5 (Fig 11A-B); EPS8 decreased modestly but not significantly
in *Tmie$^{KO/KO}$* hair bundles. Here we used fluorescence intensity normalized to heterozygote levels for
each age, which reduced signal variability but prevented us from comparing total fluorescence at different
ages.

Fig 11. Distribution of EPS8 at tips of stereocilia in tip-link and transduction mutant IHCs.
A-B, EPS8 fluorescence average intensity per hair bundle for all measured stereocilia (row 1
and 2) for *Cdh23^{v2J}*, *Pcdh15^{av3J}*, and *Tmie^{KO}* hair cells at P7.5 (A) and P21.5 (B). Plotting and
statistical testing were as in Fig 3. *C-F*, Frequency distribution of stereocilia length and EPS8 tip
intensity in *Cdh23^{v2J}* hair cells. *G-J*, Frequency distribution of stereocilia length and EPS8 tip
intensity in *Pcdh15^{av3J}* hair cells. *K-N*, Frequency distribution of stereocilia length and EPS8 tip
intensity in *Tmie^{KO}* hair cells. Stereocilia length measurements (B, D, F, H, J, L) and EPS8 tip
intensities (C, E, G, I, K, M) were from all row 1 and 2 stereocilia; distributions were fit with

double Gaussian functions. ***O-R***, Correlation of normalized EPS8 intensity with stereocilia
length. Data from individual rows were displayed with separate symbol colors but all data were
used for fits; data were fit with $y = a + b \cdot \exp(-k \cdot x)$, where y is stereocilia length, x is EPS
intensity, and a, b, and k are constants.

Frequency distribution plots, using normalized EPS8 intensities (signal at each tip divided by the total
signal for all row 1 and 2 stereocilia measured in the hair bundle), revealed how EPS8 was uniformly
targeted to stereocilia tips in tip-link mutants. Fitting each frequency distribution with a double Gaussian
function, two prominent peaks were detected in $Cdh23^{v2J}/+$ or $Pcdh15^{av3J}/+$ controls (blue data in Fig 11D,
F, H, J). By contrast, $Cdh23^{v2J/v2J}$ and $Pcdh15^{av3J/av3J}$ mutants each displayed a single distribution,
centered at a tip intensity/average = 1 (orange data in Fig 11 D, F, H, J). The convergence to a single
distribution was seen both at P7.5 (Fig 11D, H) and P21.5 (Fig 11F, J). These results confirmed the
relative uniformity of EPS8 at all stereocilia tips in tip-link mutants.

We also plotted the distribution of row 1 and 2 stereocilia lengths for each dataset; in homozygous
mutants, the length distributions were biased towards shorter stereocilia lengths and were broadened,
especially at P21.5 (Fig 11C, E, G, I). These results also show that a consequence of the increased
variability in row 1 and 2 lengths in the tip-link mutants is an increase in stereocilia of intermediate lengths
(Fig 11 C, E, G, I), in between average lengths for row 1 and 2 in the heterozygote controls. Thus, despite
their lack of length coordination within rows, hair bundles of tip-link mutants as a whole contain stereocilia
of more uniform lengths than those of controls.

In $Tmie^{KO/KO}$ hair bundles at P7.5, the EPS8 frequency distribution showed a broad double peak, unlike
the two well separated peaks in the $Tmie^{KO}/+$ control (Fig 11L; see also panel J in S5 Fig); the distribution
was weighted more towards high EPS8 tip/average values than in the tip-link mutant plots. At P7.5,
$Tmie^{KO/KO}$ showed a broad distribution of stereocilia length that was still fit with two Gaussians (Fig 11K).
EPS8 distribution and stereocilia length for $Tmie^{KO/KO}$ at P21.5 were broadly similar to those of
$Cdh23^{v2J/v2J}$ and $Pcdh15^{av3J/av3J}$. These results highlight the effect of loss of CDH23 or PCDH15 on EPS8;
while tip-link mutants have lost transduction, as have $Tmie^{KO/KO}$ bundles, the impact on EPS8 distribution
in the tip-link mutants is somewhat greater.

We also compared the EPS8 intensity at each tip with the length of that stereocilium at P15.5 (Fig 11O-
R). We found that these data were fit well with an exponential, showing that while the length of shorter
stereocilia was correlated with EPS8 intensity, beyond a certain EPS8 intensity, stereocilia length no
longer depended on EPS8 intensity. Loss of PCDH15 caused a broadening of this curve, although the
trend was still the same. Loss of TMIE did not change the shape of the curve but instead lowered the
plateau, reflecting the reduced stereocilia lengthening in these mutants. While more stereocilia had
intermediate EPS8 intensity in both mutants, stereocilia belonging to different rows were scattered along
the curve in the $Pcdh15^{av3J/av3J}$ mutants, reflecting the loss of row organization in these mutants. These

results suggested that EPS8 could influence stereocilia length, but only up to a certain length; this
maximal elongation may be regulated by mechanisms in addition to EPS8 levels.

Total GNAI3, GPSM2, and WHRN fluorescence at stereocilia tips was not affected in tip-link mutants (Fig
10A-B, E-F, I-J; Fig 12A-D; panels A-D in S4 Fig). In $Tmie^{KO/KO}$ mutants, as previously reported [6],
GNAI3, GPSM2, and WHRN fluorescence at stereocilia tips was considerably reduced (Fig 10C-D, G-H,
K-L), with statistically significant reductions for GNAI3 and WHRN at P7.5 (Fig 12A-B) and GPSM2 at
P15.5 (Fig 12C). These results confirm that TMIE (and TMC1/2) affect distribution of GNAI3, GPSM2 and
WHRN directly, not via transduction-mediated ion flux [6]. TMC2 still targeted to stereocilia in
$Pcdh15^{av3J/av3J}$ mutants, but TMC1 disappeared (S3 Fig), suggesting that TMC2 may be sufficient for
localization of GPSM2, GNAI3, and WHRN to row 1 tips.

Fig 12. Distribution of GNAI3 and WHRN at tips of stereocilia in tip-link and transduction
mutant IHCs.

A-B, GNAI3 (A) and WHRN (B) fluorescence average intensity per hair bundle for all measured
stereocilia (row 1 and 2) for $Cdh23^{v2J}$, $Pcdh15^{av3J}$, and $Tmie^{KO}$ hair cells at P7.5. Plotting and
statistical testing were as in Fig 3. ***C-F***, Frequency distribution of GNAI3 and WHRN tip intensity
in $Cdh23^{v2J}$ hair cells. ***G-J*** Frequency distribution of GNAI3 and WHRN tip intensity in $Pcdh15^{av3J}$
hair cells. ***K-N***, Frequency distribution of GNAI3 and WHRN tip intensity in $Tmie^{KO}$ hair cells. Tip
intensities for GNAI3 (E, G, I, K, M, O) and WHRN (F, H, J, L, N, P) were from all row 1 and 2
stereocilia; distributions were fit with double Gaussian functions.

In control hair bundles at P7.5, GNAI3 frequency distributions were very different from those of EPS8;
most stereocilia had little or no GNAI3, but a minority had a considerable amount (blue in Fig 12E and
panel Q in S4 Fig). This distribution matched the immunocytochemistry images, where GNAI3 was found
nearly exclusively at row 1 tips (Fig 10; S4 Fig). In $Cdh23^{v2J/v2J}$ and $Pcdh15^{av3J/av3J}$ mutants, the frequency
distribution was moderately altered; most stereocilia still had little or no GNAI3, but relatively larger
numbers had intermediate levels of signal (orange in Figs 12E and panel Q in S4 Fig). Even though
GNAI3 levels dropped significantly in $Tmie^{KO/KO}$ bundles (Fig 12A), the frequency distribution resembled
that of tip-link mutants (green in Fig 12I). The distribution of WHRN at P7.5 was similar to but broader
than that of GNAI3, and was altered similarly in tip-link and transduction mutants (Figs 12B, F, J and
panels N, R in S4 Fig). At P21.5, distribution of GNAI3 and WHRN among stereocilia tips was not altered
substantially in tip-link mutants (panels O-P and S-T in S4 Fig).

Total GPSM2 intensity was decreased significantly in stereocilia tips of P15.5 $Tmie^{KO/KO}$ mutants,
although EPS8 was not (Fig 12C-D). The distribution of GPSM2 at P15.5 in heterozygote stereocilia (Fig
12G, K) was similar to that of GNAI3 at P7.5 (Fig 12E, I), and the altered distribution in $Pcdh15^{av3J/av3J}$
and $Tmie^{KO/KO}$ mutants also resembled that of GNAI3 (Fig 12G, K).

Plots of stereocilia length against GPSM2 intensity (Fig 12M-P) were fit well with an exponential
relationship for both controls and mutants, similar to that seen with EPS8 (Fig 11O-R). The exponential

constant k was considerably smaller for the GPSM2 fits ($k \approx 3$) than for the EPS8 fits ($k \approx 8$), consistent
with GPSM2 tip accumulation only occurring in the longest stereocilia (i.e., row 1).

Tmie does not interact genetically with *Pcdh15*

We created double knockout *Pcdh15^{av3J/av3J}*;*Tmie^{KO/KO}* (*Pcdh15;Tmie* DKO) mice to clarify interaction
between *Pcdh15* and *Tmie* (S5 Fig). KO;KO hair bundles demonstrated unaltered EPS8 levels and a
single-Gaussian distribution, similar to that in tip-link mutants (panels I-J in S5 Fig). KO;KO bundles also
showed reduced GNAI3 levels and altered distribution as seen in *Tmie^{KO/KO}* bundles (panels K-L in S5
Fig). KO;KO bundles shared features of both Het;KO and KO;Het bundles (panels A-D in S5 Fig).
Mutations in *Pcdh15* thus did not suppress transduction-mutant phenotypes, nor did mutations in *Tmie*
suppress phenotypes seen in tip-link mutants, indicating a lack of genetic interaction for these
phenotypes.

Labeling stereocilia free barbed ends with rhodamine-actin

Row 2 stereocilia shrink during stage III, then stabilize in length during stage IV (Fig 1G). Actin-filament
free ends, measured in permeabilized hair bundles using rhodamine-actin as a probe, were particularly
high at row 2 tips in control bundles during early postnatal development but decreased in cochleas of
transduction mutants or cochleas treated with transduction-channel blockers [39]. Results with tip-link
mutants (Fig 13A-B, G-H, M, P) were very similar to those obtained with *Tmie^{KO/KO}* mutants [39] (Fig 13C-
D, I-J, N, Q). In *Pcdh15^{av3J}*/+ controls, the high level of row 2 labeling by rhodamine-actin seen in control
bundles at P7.5 (Fig 13A, M) had disappeared by P21.5 (Fig 13G, P); row 1 labeling was reduced as well
at P21.5 (Fig 13P). In *Pcdh15^{av3J/av3J}* mutants at P7.5, rhodamine-actin labeling was relatively low, with
row 1 and 2 tips having similar intensities (Fig 13B, M). That pattern was relatively unchanged at P21.5
(Fig 13H, P).

Fig 13. Free barbed end labeling of IHC stereocilia with rhodamine-actin.
A-L, Rhodamine-actin (Rh-actin) and phalloidin labeling of mutant IHC hair bundles at P7.5 and
P21.5. *Pcdh15^{av3J}* (A-B, G-H), *Tmie^{KO}* (C-D, I-J), and *Myo15a^{sh2}* (E-F, K-L). Arrows in A, C, and
E indicate high levels of row 2 rhodamine-actin labeling in P7.5 control IHC bundles; asterisks
in B, D,F, H, and J indicate similar rhodamine-actin levels at all stereocilia tips in P7.5 and P21.5
mutant IHC bundles. Panel widths: 24 µm. ***M-R***, Rh-actin tip intensity and intensity CV in
Pcdh15^{av3}, *Tmie^{KO}*, and *Myo15a^{sh2}* at P7.5 (M-O) and P21.5 (P-R). Plotting and statistical testing
were as in Fig 3.

By contrast, in *Myo15a^{sh2/sh2}* mice, row 1 and row 2 rhodamine-actin labeling was elevated to the intensity
level seen in row 1 from control mice at P7.5 (Fig 13E-F, O); rhodamine-actin labeling then declined
precipitously in both rows by P21.5 (Fig 13K-L, R). Thus, during stage III, actin filament free ends appear
to be suppressed by the row 1 complex but enhanced by transduction; in turn, increased free ends
correlate with row 2 shortening.

CAPZB no longer concentrates at P21.5 stereocilia tips in tip-link mutants

After P10, CAPZB—and to a lesser extent, EPS8L2—shifts from the shafts of all stereocilia to concentrate at row 2 tips in control hair bundles [6]; this shift does not occur in homozygous transduction mutants [6]. We confirmed results in $Tmie^{KO}$ mice (Fig 14C-D, G-H) and showed further that CAPZB no longer concentrated at row 1 or 2 tips in $Pcdh15^{av3J/av3J}$ IHCs (Fig 14A-B, G), instead remaining along stereocilia shafts (asterisks in Fig 14B). By contrast, CAPZB concentrated at tips of both rows in relatively equal amounts in in $Myo15a^{sh2/sh2}$ bundles (arrows in Fig 14F; Fig 14G).

> **Fig 14. Row 2 protein localization in mutant IHCs.**
>
> **A-F**, Localization of CAPZB and EPS8L2 in $Pcdh15^{av3J}$ (A-B), $Tmie^{KO}$ (C-D), and $Myo15a^{sh2}$ (E-F) heterozygous and homozygous IHCs at P21.5. Arrows in A, C, and E show high levels of row 2 CAPZB in heterozygotes. In B, asterisks show CAPZB labeling along stereocilia shafts and arrowheads show strong EPS8L2 labeling at row 1 and 2 tips. Arrows in F show CAPZB at tips of all stereocilia in $Myo15a^{sh2/sh2}$. Panel widths: H-M, 30 μm. **G-H**, Quantitation of CAPZB (G) and EPS8L2 (H) average intensity in row 2 tips divided by the average intensity in row 1 tips in heterozygotes and homozygotes at P21.5. Plotting and statistical testing were as in Fig 3.

EPS8L2 was present at both row 1 and row 2 tips in control hair bundles, with a modest bias towards row 2; EPS8L2 was found at nearly equal levels in both rows of $Pcdh15^{av3J/av3J}$ and $Tmie^{KO/KO}$ IHC bundles (Fig 14A-D, H; arrowheads in Fig 14B). In contrast to CAPZB, EPS8L2 was nearly completely absent from $Myo15a^{sh2/sh2}$ bundles (Fig 14E-F). These results suggested that the presence of transduction and the row 1 complex are required for selective targeting of CAPZB to row 2 tips during late development. The appearance of CAPZB at tips by P21.5 was correlated with the decline in free barbed end labeling at this age for both control and mutant bundles, suggesting that capping proteins, row 1 proteins, and transduction all stabilize F-actin at stereocilia tips during stage IV.

Discussion

The results presented here support a comprehensive model that explains key steps of developmental assembly of the apical IHC hair bundle (Fig 15). By increasing resolution for our measurements of stereocilia dimensions, we defined more accurately Tilney's stages III and IV of stereocilia growth in mouse hair cells and the point of transition between them [1]. These improvements were made possible by boosting sampling frequency and increasing measurement precision using automated analysis. In addition, use of four mouse mutant lines allowed us to determine what role the tip-link proteins, transduction proteins, and the row 1 complex play in bundle development. Together the results provide a picture of several simultaneous processes proceeding in parallel with significant interconnections reinforcing key transitions.

Fig 15. Model for assembly of IHC hair bundle.

A, Left, stereocilia building blocks (combined actin and actin crosslinkers) increase continuously
in IHCs throughout early postnatal development. Right, a spatial gradient of building blocks may
also exist, with building blocks having greater access to row 1 stereocilia than to shorter rows.
B, PCDH15 and CDH23 maintain coordination of adjacent stereocilia lengths, presumably by
triggering actin depolymerization when spontaneous stereocilia elongation interrupts the stable
state. *C-D*, Stage-dependent processes. *C*, EPS8 is present at all stages of stereocilia growth.
In stage II, EPS8 is present at all tips in approximate proportion to stereocilia length. In stage III
and IV, EPS8 levels drop at row 2 tips while they remain constant at row 2 tips. *D*, GPSM2
appears in row 1 stereocilia during stage II, and stimulates the row 1 complex sufficiently to drive
this phase of lengthening. GPSM2 levels drop during stage II, when lengthening is arrested. If
transduction proteins are present in the hair bundle, GPSM2 levels rise during stage IV,
presumably leading to activation of EPS8 and actin-filament elongation. *E*, At row 2, the actin-
depolymerizing proteins DSTN and CFL1 (and their partner WDR1) are found diffusely in
stereocilia. During stage III, transduction leads to localization of DSTN and CFL1 at row 2 tips,
triggering actin-filament turnover and shrinkage of row 2. CAPZB and TWF2 are localized
separately during this stage, but in stage IV both are found together at row 2 tips if transduction
is active.

Steady growth of stereocilia building blocks and local regulation

In chick cochlea, hair bundles with widely varying numbers and dimensions of stereocilia still have the
same total amount of F-actin in their bundles [56]. Similarly, we found that P7.5 bundles from mouse
mutants with differing stereocilia number, length, and width still had the same total stereocilia volume.
Moreover, in wild-type hair cells, the total stereocilia volume per bundle increased monotonically despite
the very different growth behavior of individual stereocilia rows during stages III and IV. While cellular
actin transcript and protein levels do not rise during early postnatal bundle development [58–60] (see
also umgear.org), actin crosslinkers do increase in expression over this period [59, 61] (umgear.org);
moreover, loss of crosslinkers affects stereocilia dimensions [61–63], and FSCN2 overexpression
lengthens stereocilia [64]. A continually increasing supply of actin crosslinkers, including ESPN, FSCN2,
and PLS1, may therefore steadily increase the supply of "building blocks" (i.e., actin plus actin
crosslinkers) for stereocilia growth during early postnatal development.

These results suggest that hair cells continuously synthesize stereocilia building blocks and that local
regulatory mechanisms then control how these building blocks are exploited. These local regulatory
mechanisms must be switched on and off rapidly. Our results showed that during stage III, lengthening
of stereocilia was completely inhibited; cross-sectional area grew linearly during this time, suggesting a
constant rate of addition of actin filaments to the periphery of the stereocilia core. At P8, however, a sharp
transition occurred; lengthening resumed then and proceeded exponentially for the next two weeks.

Stereocilia building blocks appear to be made available to the hair bundle in a lateral to medial gradient;
even when tip links are not present and the row 1 complex is distributed to most stereocilia, a row 1

through row 3 gradient in length was still apparent. This gradient was even apparent in *Myo15a^sh2/sh2*
mutants, which lack the row 1 complex at any tips. A plausible explanation is that building blocks are
more accessible at the fonticulus, the aperture in the cuticular plate that the kinocilium extends through.
This region is not only filled with microtubules and vesicles [65] but also could be a region where soluble
molecules like actin monomers and crosslinkers more freely diffuse. The alternative explanation that
stereocilia growth starts later in shorter stereocilia is not supported by the comparison of row 1 and 2
kinetics in C57BL/6 hair cells (Fig 1).

The row 1 tip complex controls lengthening in stages II and IV
The biochemical properties of the row 1 tip complex, composed of EPS8, MYO15A, WHRN, GNAI3, and
GPSM2, suggest how the complex controls stereocilia length. EPS8 and MYO15A each can control actin
polymerization [66, 67], and early in development, the two proteins target to all stereocilia in amounts
approximately proportionally to stereocilia length [7]. Moreover, together both proteins form a phase-
separated complex that may promote actin polymerization [10]. In the absence of transduction [6] or tip
links (this work), the EPS8-MYO15A complex is no longer exclusively targeted to row 1 and instead
appears at all tips, also in proportion to length. These results suggest that the EPS8-MYO15A complex
is targeted to all tips, but that other events during development lead to their accumulation with the rest of
the complex members only at the tips of row 1.

In stereocilia, as long as functional RGS12 is present [68], GPSM2 and its partner GNAI3 are exclusively
found at row 1 tips [7, 8]; there, they assemble with the EPS8-MYO15A complex [7–9] using WHRN as
an adapter [9]. GPSM2 forms large protein condensates in vitro, both by itself and with the entire row 1
complex [11], and it is reasonable to suggest that GPSM2 drives the formation of similar condensates at
the tips of row 1 stereocilia. When the row 1 complex is formed, GPSM2 activates EPS8's F-actin
bundling activity, which likely promotes stereocilia lengthening [11]. GNAI3 can target to stereocilia tips
independently from other row 1 proteins [7], and may act as an adapter between the membrane at row 1
tips and the rest of the row 1 complex [11].

Previous characterization of the row 1 tip complex has not distinguished its activity during the two
lengthening stages of hair-bundle development [1]. In apical IHCs, stage II—initial lengthening of
stereocilia—occurred prior to our first measurements at P0.5. Tadenev and colleagues demonstrated
that GNAI3 and GPSM2 are indeed required for this initial lengthening stage. First, both proteins appear
at row I tips during stage II [9]. In addition, hair cells that express pertussis toxin, which inactivates all
GNAI isoforms, have very short stereocilia, as do *Gpsm2*-null hair cells [9]. These stereocilia are much
shorter than those of bundles at P0.5, which is at the end of stage II. Bundles lacking EPS8, MYO15A,
or WHRN each show the same short-stereocilia phenotype [69–71].

Although these simple knockouts do not probe later steps in development, our experiments suggest that
GNAI3 and GPSM2 are also required for the final stereocilia lengthening that occurs during stage IV.

GNAI3 and GPSM2 each build up at tips of row 1 stereocilia during stage IV, with GPSM2 peaking later
than GNAI3. $Cdh23^{v2J/v2J}$ and $Pcdh15^{av3J/av3J}$ mutants, which retain GNAI3 and GPSM2 in row 1, have
row 1 lengths nearly equal to those of controls. By contrast, $Tmie^{KO/KO}$ mutants, which do not accumulate
GNAI3 or GPSM2 in row 1 stereocilia during stage IV, have shorter row 1 stereocilia.

Loss of CDH23 or PCDH15 leads to uncoordinated stereocilia lengthening
Our experiments also revealed several specific functions for CDH23 and PCDH15, presumably as
heterotetrametric links, during development of apical IHC hair bundles. While CDH23-PCDH15 tip links
gate transduction channels and transduction impacts bundle development [6], we show here that CDH23
and PCDH15 are also essential for coordinating the lengths of adjacent stereocilia in the same row. While
we cannot rule out effects specific to CDH23 or PCDH15, the similarity in phenotype between $Cdh23^{v2J}$
and $Pcdh15^{av3J}$ mouse lines and demonstrated interaction between them [26, 72, 73] makes it most likely
that roles for these two proteins, like length coordination, require their interaction.

CDH23 and PCDH15 have each been suggested to contribute to the transient lateral links [24, 26, 74,
75], and these links may be essential for coordinating adjacent stereocilia lengths. Transient lateral links
bridge the gap between adjacent stereocilia along their shafts and are prominent in mouse IHCs and
OHCs between E17.5 and P0; they begin to disappear after P0, and are no longer seen after P5 except
at stereocilia tips [76]. Although some links between adjacent stereocilia remain in both $Pcdh15^{av3J/av3J}$
and $Cdh23^{v2J/v2J}$ hair bundles (Fig 2), these are likely ankle links [76, 77].

Transient lateral links are similar to the thread-like links that interconnect the distal microvilli tips in the
intestinal brush border [78]; these links are composed of CDHR2 (PCDH24) and CDHR5, members of
the adhesion complex, and are essential for microvillar clustering and length uniformity [78]. Length
uniformity in microvilli of the brush border has been proposed to arise from stimulation of actin dynamics
through the adhesion complex or by physical constraints, as a link could prevent an adjacent microvillus
from elongating [79]. In IHCs, we propose that dimeric complexes of CDH23 and PCDH15 carry out the
same function throughout development.

Nevertheless, the morphology of $Pcdh15^{av3J};Myo15a^{sh2}$ DKO hair bundles was dissimilar to that of CACO-
2_{BBE} intestinal cells with shRNA knockdown of PCDH24; these latter cells had microvilli of widely differing
lengths that lacked hexagonal arrangement [78]. DKO bundles retained a modest gradient in stereocilia
length, despite the lack of tip links, and adjacent stereocilia were similar in length; while packed less
tightly than either $Pcdh15^{av3J}/+;Myo15a^{sh2}/+$ or $Pcdh15^{av3J}/+;Myo15a^{sh2/sh2}$ stereocilia, packing was still
hexagonal. The retention of length coordination, albeit to very short lengths, suggests that additional
mechanisms contribute to regulation of stereocilia length in addition to the row 1 complex and PCDH15.
Physical constraints of the overlying tectorial membrane probably do not contribute to length regulation;
when the tectorial membrane is eliminated genetically, IHC and OHC bundles retain their normal
morphology and stereocilia lengths are aligned [80, 81].

A system for aligning lengths of adjacent stereocilia may have evolved to produce optimal activation of
transduction channels. Alignment of stereocilia lengths in each row should coordinate transduction
channel gating and increase sensitivity. For small stimuli, the shear between two stereocilia can be
approximated by $x = \gamma \cdot X$ [82], where X is the displacement applied to the hair bundle's tip and γ is a
geometrical gain factor; x is the distance that the gating spring is stretched by the stimulus and hence is
the relevant activation delivered to the transduction channels [83]. In turn, $\gamma = S / L$, where S is the
spacing between stereocilia at their bases and L is the length of the stereocilia at the point of shear. In
wild-type bundles, spacing between adjacent stereocilia in a given row is constant within that row and
from bundle to bundle [82]. By contrast, in tip-link mutants, variability in L of row 2 will produce variability
in x; if a cell has variations in x with a constant X, channel open probabilities will vary, perhaps
considerably. In turn, variability in open probabilities will broaden the displacement-transduction current
relationship of the bundle and reduce sensitivity. Tight control of stereocilia length within a row is thus
required to maintain the hair cell's high sensitivity.

739 Transduction proteins localize the row 1 complex
Our results here confirm our previous suggestion that transduction proteins themselves regulate the
amount of GNAI3 and GPSM2 that accumulate at row 1 tips during stage IV [6]. While little GNAI3 (or
GPSM2) is found in row 1 of $Tmie^{KO/KO}$ mutant hair bundles, GNAI3 and GPSM2 concentrate relatively
normally in row 1 stereocilia in $Cdh23^{v2J/v2J}$ and $Pcdh15^{av3J/av3J}$ mutants. Stage III and IV localization of
these proteins to row 1 tips occurs long after the initial lengthening of stereocilia adjacent to the kinocilium,
so transduction mediated through kinocilia-stereocilia links [84] is likely not responsible. TMIE and the
TMCs do localize to row 1 near tips during early postnatal development [23], and at least TMC2 is present
in row 1 in $Pcdh15^{av3J/av3J}$ mutants (S3 Fig).

Accordingly, GNAI3 or GPSM2 may bind directly or indirectly to TMC channels localized to tips of row 1
stereocilia during development (Fig 15E). This coupling could be mediated by the established CIB2-
WHRN interaction [85]; CIB2 is part of the transduction-channel complex [86–88] and WHRN interacts
with GPSM2 [9, 11]. TMC channels are no longer at row 1 tips in older stereocilia [89], so localization of
GNAI3 and GPSM2 to row 1 by the transduction complex may only be necessary at the beginning of
stage IV to initiate formation of a robust row 1 complex.

754 Transduction controls actin dynamics at row 2 tips
Transduction controls actin-regulatory mechanisms in row 2 differently than those in row 1; during stage
III, row 2 stereocilia shorten when row 1 lengthens, showing they respond differently during development,
and the complement of proteins at their tips differs from that at row 1 tips. Our results confirm the
observations of McGrath and colleagues, who demonstrated that row 2 tips transiently have abundant
free actin filaments ends, but not in $Tmie^{KO/KO}$ mutants nor in the presence of channel blockers [39].
$Pcdh15^{av3J/av3J}$ mutants also lose strong row 2 rhodamine-actin labeling, confirming that transduction
itself—rather than transduction proteins—is responsible for this effect.

Effects of transduction on protein localization likely occur because transduction channels are partially
open at rest during development [90], and because substantial Ca^{2+} entry occurs when channels are
open [16]. The steep local gradient of Ca^{2+} may in turn localize specific proteins to row 2 tips but not
elsewhere in the row 2 stereocilia.

Transduction recruits DSTN (ADF) and CFL1 to stereocilia tips [39], which may depend on local Ca^{2+}
entry (Fig 15F). Moreover, in many contexts, Ca^{2+} influx increases DSTN and CFL1 activity by activating
Ca^{2+}-dependent phosphatases [40]. In tip-link or transduction mutants, not only is the high level of
rhodamine-actin labeling at row 2 tips reduced, but rhodamine-actin labeling at row 1 tips also increases.
These results suggest that actin dynamics are regulated by transduction at both row 1 and row 2 tips, but
in opposite directions. Transduction contributes to the polarization of EPS8 into row 1, as both $Tmie^{KO/KO}$
and $Pcdh15^{av3J/av3J}$ mutants have relatively similar amounts of EPS8 in all tips; in this case, the similarity
in phenotype between the two mutants suggests that transduction currents themselves mediate EPS8
exclusion from the shorter rows. By preventing accumulation of EPS8 at row 2 tips, transduction could
thereby promote its accumulation at row 1, thus contributing to the polarization in actin dynamics between
the two rows.

These results suggest that transduction stimulates actin depolymerization at the tips of the shorter rows
[38, 39]. Moreover, by increasing actin dynamics, transduction thus also controls stereocilia remodeling
(row 2) or resorption (rows 3 and beyond) during development. A key question is why this model, which
suggests that transduction promotes row 2 shortening, is contradicted by experiments showing that
transduction-channel block stimulates row 2 shortening [17]. A plausible explanation for this discrepancy
is that these latter experiments were conducted on stage IV hair bundles [6]. Thus while transduction
promotes row 2 shortening during stage III, changes that occur during stage IV reverse the sign of
transduction's effect. The F-actin capping complex of CAPZB and TWF2 appears during stage IV only if
transduction is present [6], so the effect of blocking transduction during stage IV may be to dissociate the
capping complex and trigger stereocilia shortening.

A model for hair-bundle length regulation during stages III and IV
Our model for development of mouse IHC hair bundles (Fig 15) references Tilney's chick bundle
development model [1]; it also exploits key observations made in developing mouse cochlear hair cells
[7–9, 39] and biochemical studies of the row 1 complex [9, 11]. In this model, mouse bundles undergo
temporally segregated initial lengthening (stage II), widening (stage III) and final lengthening (stage IV)
steps. A key feature of bundle development is the creation of three stereocilia rows, each with unique
dimensions for their stereocilia. Establishment of these rows occurs when symmetry within the bundle is
broken, allowing distinct actin-regulatory mechanisms to control F-actin growth within individual
stereocilia rows.

At least two global mechanisms, not specific to any row, also regulate growth of hair bundles. Our results
suggest that the concentration of building blocks for stereocilia, including G-actin, actin crosslinkers, and
other essential actin cytoskeleton proteins, slowly increases in each hair cell over development. Even
when individual stereocilia rows show sharp transitions in lengthening and widening, there is little
evidence of any alteration in the rate of growth of total stereocilia volume. Thus the cell constantly
supplies stereocilia building blocks, and regulatory steps that are specific to individual rows leads to the
transitions in bundle-assembly modes. In all rows, EPS8 mediates incorporation of these building blocks
into growing stereocilia (Fig 15A). Another global mechanism is coordination of stereocilia length, which
occurs within each row because of the presence of CDH23 and PCDH15, presumably through the
transient lateral links (Fig 15B).

A key event during development of the bundle is the breaking of symmetry between the rows. Asymmetry
is created not only by increasing the length of row 1, but also by limiting the growth of the shorter rows.
The three main symmetry-breaking mechanisms that affect actin regulation are (a) establishment of the
row 1 complex by GNAI3 and GPSM2; (b) the appearance of tip links, which pull on the tips of the shorter
stereocilia but not on row 1 tips; and (c) appearance of transduction in the shorter rows. These
mechanisms in turn control three actin-regulatory processes: first, F-actin bundling and polymerization,
which lengthen or widen stereocilia; second, F-actin depolymerization, which shortens or thins stereocilia;
and finally, F-actin capping, which stabilizes filaments and hence stereocilia length. By locally altering
the balance of these three regulatory processes, stereocilia of varying dimensions can be created.

GPSM2 and GNAI3, which are only recruited to row 1, presumably boost EPS8 F-actin-bundling activity
to lengthen row 1 during stage II (Fig 15E). Given that they do not have GPSM2 and GNAI3, how do the
shorter rows lengthen during this stage? Transduction is not involved, as it only appears in apical IHCs
after P0 [52, 91], after stage II is complete (Fig 15C). These rows likely lengthen during stage II because
tip links are already present [76], and the tension they apply to the tips of the shorter stereocilia rows
causes membrane tenting, actin polymerization, and stereocilia lengthening [36]. This lengthening likely
is mediated by EPS8-MYO15A complexes, which are present at this early stage at row 2 and 3 tips [9].

GPSM2 and GNAI3 also control lengthening during stage IV (Fig 15E). While EPS8 levels are high across
stages III and IV, GPSM2 and GNAI3 levels are low during stage III widening. Towards the end of stage
III, however, GNAI3 begins to accumulate at row 1 tips, reaching a peak during early stage IV. This
accumulation depends on the presence of the transduction complex, and may be due to binding of GNAI3
to the complex at the tip of row 1, where it is transiently during development. GPSM2 appears at row 1
tips later during stage IV (Fig 15E), presumably requiring GNAI3 to establish its presence first, and
recruitment of GPSM2 may be directly responsible for the burst in lengthening that occurs then. During
stage IV, only row 1 of apical IHCs lengthens.

Tip links appear early in development of the hair bundle (Fig 15C), before the appearance of transduction
[48, 52, 92]. Tip-link tension pulls the membrane at the tip of the short stereocilium of the interconnected
pair away from the F-actin core beneath (tip-link tenting), which eliminates the membrane's suppression
of actin polymerization [93, 94]. Actin polymerization thus proceeds as the membrane is tented, allowing
stereocilia lengthening [36]. During stage II, the shorter rows may lengthen simultaneously with row 1,
but stereocilia in the shorter rows will be limited in their maximal length because the molecular motor
pulling on the tip link will run to the end of its track. Thus, bundles formed by the end of stage II have
multiple rows of thin stereocilia with a relatively short stereocilia staircase spacing (Fig 1A). Bundles
around P0 are also often tilted strongly in the negative direction [48, 52], which may occur when tip links
draw all of the stereocilia together. Negatively tilted bundles are especially apparent in OHCs. Because
of the different structures at each of their ends, tip links thus contribute to the distinction between row 1
and the shorter rows.

During stage III, the appearance of transduction (Fig 15C) allows for further divergence of row identity
(Fig 15F). Steady Ca^{2+} influx, due to the transduction channels being partially open at rest, is present at
all rows except row 1 [22]. DSTN and CFL1 are recruited to the tips of shorter stereocilia if transduction
is active, and these proteins may also be activated indirectly by entering Ca^{2+}. Activated DSTN and CFL1,
in turn, increase the turnover of actin filaments at tips of these rows, which may directly lead to both row
2 shortening and pruning (disappearance) of other rows. In addition, EPS8 declines at row 2 tips during
stage III (Fig 15D), further distinguishing these tips from those of row 1 stereocilia. The presence of
dynamic actin at row 2 tips may explain this decline. EPS8 may still target to the tips of shorter rows, but
could be released after filament severing by DSTN and CFL1. The F-actin capping protein EPS8L2
instead concentrates at row 2 tips [6, 15], and its mechanism of localization and function likely differs
from that of EPS8.

Changes in length of one row may affect the availability of building blocks for remaining rows, especially
later in development. For example, without robust lengthening of row 1 (e.g., in *Myo15a*$^{sh2/sh2}$ hair
bundles), more building blocks are available to thicken shorter stereocilia rows, which also stabilizes
them. If depolymerization activity is low due to a lack of transduction currents (e.g., in transduction or tip-
link mutants), supernumerary stereocilia can thicken. In addition, allocation of stereocilia building blocks
appears to extend along a lateral-medial gradient, such that the stereocilia closest to the lateral edge (the
bare zone) receive building blocks at a higher rate than those farthest away. Finally, the taller a
stereocilium, the farther it is from cytoplasmic pools of DSTN and CFL1 and cytoplasmic changes in Ca^{2+}.
Short stereocilia whose filaments are not capped by EPS8, EPS8L2, CAPZB, or TWF2 will be more
susceptible to resorption.

How the mutant phenotypes arise
The mutant phenotypes described here each result from failure in one or more of the three main
symmetry-breaking events that normally occur during development. In *Myo15a*$^{sh2/sh2}$ hair bundles, without

trafficking of the row 1 complex, stereocilia length asymmetry cannot be created; lengthening requires
EPS8, which is transported by MYO15A. The lack of lengthening of the taller rows frees stereocilia
building blocks; these building blocks are subsequently used to thicken rows 3 and 4+, as well as
eventually creating cytocauds. In addition, without mechanisms to uniquely distinguish row 1, all
stereocilia become similar to row 2. For example, all rows show high levels of rhodamine-actin
incorporation at tips and equal levels of accumulation of DSTN and CFL1. Moreover, all rows are equally
thickened and capping proteins, which usually enrich at row 2 (EPS8L2, CAPZB, TWF2), are present at
all rows, presumably slowly pruning of the shorter rows.

While $Tmie^{KO/KO}$ and tip-link mutants both lack transduction, they differ in the establishment of the other
two main symmetry-breaking events. In $Tmie^{KO/KO}$ mutants, members of the row 1 complex are still
present, but neither GPSM2 nor GNAI3 traffic efficiently to row 1 tips in the absence of transduction
proteins, at least during stage IV. Stage IV lengthening of row 1 stereocilia is therefore significantly
reduced as there is no additional stimulation of EPS8 activity in that row by GPSM2. Since row 1 still
grows taller than the other rows, both tip links and the lateral-medial gradient in building blocks contribute
to forming shallow gradients in length and width. In $Cdh23^{v2J/v2J}$ and $Pcdh15^{av3J/av3J}$ tip-link mutants,
members of the row 1 complex still traffic and accumulate at normal levels at tips, so stage IV lengthening
continues to occur and taller stereocilia are created. Because kinocilial links are lost and hair bundles
change in shape, allocation of GNAI3, GPSM2, and WHRN becomes more variable; in some cases, they
are delivered to stereocilia beyond just those bordering the bare zone. More randomized allocation of
row 1 complex members and the varying positions of stereocilia across the lateral-medial resource
gradient may also create length variability.

Due to the loss of tip links, all stereocilia tips in $Cdh23^{v2J/v2J}$ and $Pcdh15^{av3J/av3J}$ hair bundles develop a
rounded shape; the uniquely beveled tip of the shorter rows of both control and $Tmie^{KO/KO}$ stereocilia fails
to develop. EPS8 accumulates at more equal levels across all stereocilia tips in tip-link mutants as
compared to the $Tmie^{KO/KO}$ mutants, which may occur because all stereocilia develop the rounded tip
shape of row 1. In tip-link and transduction mutants, loss of transduction promotes equal but reduced
levels of DSTN/CFL1 activity at all tips. Although $Dstn;Cfl1$ double-knockout mice retain a polarized EPS8
distribution [39], seemingly contradicting aspects of our model, the induction of the double knockout
occurred postnatally, after accumulation of EPS8 already occurred at row 1 tips. Regardless of the
mechanism, EPS8 accumulation and more equalized distribution of DSTN/CFL1 apparently allows some
supernumerary rows to thicken and persist.

Conclusions
Development of the hair bundle is guided by three symmetry-breaking events—the formation of the row 1
complex, the asymmetric distortion of the lower-stereocilium membrane caused by tip-link tension, and
the establishment of mechanotransduction only at stereocilia tips of shorter rows. During stages II, III,
and IV of bundle development, CDH23 and PCDH15 are essential for coordination of stereocilia length

within a row. During stage II, before transduction, lengthening of row 1 elicits lengthening of shorter rows

because tip links under tension stimulate lengthening; in wild-type IHCs, this produces a bundle with

three to four rows and an even staircase. Lengthening of row 1—and hence the short rows too—stops at

the beginning of stage III, when transduction first appears. Stereocilia widen during stage III since building

blocks continue to be supplied. During stage III, row 2 stereocilia (and row 3) shorten modestly due to

the action of DSTN and CFL1 but also due to the exclusion of EPS8. The row 1 complex reassembles

during stage IV in a stepwise manner, dependent on the expression of transduction proteins. Rapid

growth of row 1 stereocilia does not occur until sufficient GPSM2 is present; these stereocilia then

lengthen while those of the shorter rows continue to shrink. Thus the interplay of the row 1 complex and

transduction creates the final apical IHC bundle, with its dramatic length differences between the rows.

Materials and Methods

Ethics statement

All animal studies were performed in accordance with the guidelines established by the Oregon Health & Science University Institutional Animal Care and Use Committee (protocol #IP00000714) and the Johns Hopkins University Institutional Animal Care and Use Committee (protocol #M016M271).

Mice

Mice were housed in individually ventilated cages (IVCs; Thoren) with a maximum of five animals per cage with food and water ad libitum. Breeder pairs were separated 16 to 18 days after crossing and single housed animals were provided with nesting material; pregnant dams were given breeder chow (5058 PicoLab; LabDiet) in place of standard chow (5LOD Irr Rodent; LabDiet). Mice were housed under barrier specific pathogen free conditions in accordance with OHSU IACUC. Mouse pups were assumed to be born at midnight, so animals used on the first day are referred to as P0.5. Data labeled as "P21.5" here refers to mice at weaning age; while most animals were P21.5 (including all animals used in Fig 1), we used a range of ages from P20.5-P25.5. Both male and female pups were used; because newborn mice cannot be sexed without genetic analysis, we used a mixture of sexes in all experiments. We analyzed IHCs from the basal half of the apical ~1/3 of the cochlea (17-33% of the cochlear length, measured from the apex; roughly the 125th IHC from the apex through the 250th), and we refer to them as apical IHCs.

For developmental analysis, we used C57BL/6J mice (RRID:IMSR_JAX:000664, Jackson Laboratories, Bar Harbor, ME). While C57BL/6J mice exhibit progressive hearing loss relative to other strains, at 1.5 months of age (i.e., older than the mice we used), auditory brainstem responses and cochlear hair cell preservation were identical to strains with excellent hearing, such as CBA/Ca [95]. All mutant mice were backcrossed for more than ten generations to C57BL/6J. The $Cdh23^{v2J}$ line has been described previously [33, 50]; the $v2J$ mutation changes a donor splice site and produces stop-codon containing mutant splice forms with that are predicted to produce truncated proteins lacking the transmembrane domain. The $av3J$ mutation in $Pcdh15$ also leads to a premature stop codon prior to the transmembrane domain [33, 49]. Accordingly, both $Cdh23^{v2J}$ and $Pcdh15^{av3J}$ are expected to be full null mutations [33]. The $Tmie^{KO}$ line has been described previously [6, 18], as has $Myo15a^{sh2}$ [55, 69].

Primary antibodies

Anti-GPSM2 Cat# HPA007327 (RRID: AB_1849941) from Sigma-Aldrich (St. Louis, MO). Rabbit polyclonal anti-GNAI3, directed against a C-terminal mouse GNAI3 peptide (KNNLKECGLY), was Cat# G4040 (RRID: AB_259896) from Sigma-Aldrich (St. Louis, MO). Mouse monoclonal anti-EPS8 (clone 15), against mouse EPS8 amino acids 628-821, was Cat# 610143; RRID: AB_397544 from BD Bioscience (San Jose, CA). Rabbit polyclonal anti-WHRN (#8127), against a C-terminal mouse WHRN peptide with an added cysteine (KERDYIDFLVTEFNVML-C), was from the Ulrich Müller lab. Mouse monoclonal anti-CAPZB2, against a C-terminal peptide of human CAPZB2, was Cat# AB6017 (RRID:

AB_10806205) from EMD Millipore (Burlington, MA). Mouse monoclonal anti-EPS8L2, against human
EPS8L2 amino acids 615-715, was Cat# ab57571 (RRID: AB_941458) from Abcam (Cambridge, UK).
Anti-PCDH15 (HL5614) was generated in rabbit and obtained from Zubair Ahmed and anti-CDH23 was
generated in goat by GenScript (Piscataway, NJ) against mouse CDH23 EC15/16 (CATRPAPPDRERQ)
(Indzhykulian et al, 2013).

Immunofluorescence sample preparation
Inner ears from C57BL/6J mice or mutant mice littermates dissected at the indicated ages in cold Hank's
balanced salt solution (Cat# 14025076, Thermo Fisher Scientific) supplemented with 5 mM HEPES, pH
7.4 (dissection buffer). To allow perfusion of the fixative, small openings were made within the periotic
bones.

For measurements of stereocilia dimensions from C57BL/6J or mutant mice using only phalloidin
(Figs 1, 3-8), ears were fixed in 4% formaldehyde (Cat# 1570, Electron Microscopy Sciences) in 1x PBS
for 0.5 – 1 hour at room temperature. Ears were washed in PBS, then cochleas were dissected out from
the periotic bone and the lateral wall was removed. After permeabilization in 0.2% Triton X-100 in 1x PBS
for 10 minutes at room temperature, cochleas were incubated with phalloidin (0.4 U/ml Alexa Fluor 488
phalloidin; Cat# A12379, Thermo Fisher Scientific) in 1x PBS for 12-16 hours at 4°C. Organs were
washed three times in PBS for 5 min per wash and mounted in Vectashield (Cat# H-1000, Vector
Laboratories).

For immunofluorescence using antibodies against row 1 or 2 proteins (Figs 8-11; S4 Fig), ears were fixed
in 4% formaldehyde (Cat# 1570, Electron Microscopy Sciences) in dissection buffer for 20-30 min at
room temperature. As noted previously, row 2 protein antibodies (EPS8L2, CAPZB) required fixation of
at most 20 min, whereas row 1 protein antibodies were not sensitive to fixative duration. Ears were
washed in PBS, then cochleas were dissected out from the periotic bone and the lateral wall was
removed. Cochleas were permeabilized in 0.2% Triton X-100 in 1x PBS for 10 min and blocked in 5%
normal donkey serum (Cat# 017-000-121, Jackson ImmunoResearch) diluted in 1x PBS (blocking buffer)
for 1 hr at room temperature. Organs were incubated overnight at 4°C with primary antibodies in blocking
buffer and then washed three times in 1x PBS. Dilutions were 1:500 for anti-GNAI3; 1:250 for anti-
GPSM2, anti-EPS8, anti-EPS8L2, anti-WHRN, anti-CAPZB. Tissue was then incubated with secondary
antibodies, which were 2 µg/ml donkey anti-rabbit Alexa Fluor 488 (Cat# A21206, Thermo Fisher
Scientific) and 2 µg/ml donkey anti-mouse Alexa Fluor 568 (Cat# A10037, Thermo Fisher Scientific); 1
U/ml CF405 phalloidin (Cat# 00034, Biotium) was also included for the 3-4 hr room temperature
treatment. Tissue was washed three times in PBS and mounted on a glass slide in ~50 µl of Vectashield
and covered with a #1.5 thickness 22 x 22 mm cover glass (Cat# 2850-22, Corning) or a #1.5 thickness
22 x 40 mm cover glass (Cat# CLS-1764-2240, Fisher Scientific).

Immunofluorescence using anti-CDH23 and anti-PCDH15 antibodies was performed as outlined
previously [96]. Briefly, ears were fixed in 4% formaldehyde (Cat# 1570, Electron Microscopy Sciences)
in dissection buffer for 20-30 min at room temperature then washed twice in calcium/magnesium free
HBSS (Cat# 14-175-095m Fisher Scientific). Tissue was blocked for 2 hours in 10% normal donkey
serum (Cat# 017-000-121, Jackson ImmunoResearch) diluted in calcium/magnesium free HBSS with
2 mM EDTA, and this blocking buffer was used for all antibody steps. Tissue was incubated overnight at
4°C with 1:100 anti-CDH23 and 1:200 anti-PCDH15, washed three times with calcium/magnesium-free
HBSS, then incubated for 2-3 hours at room temperature with 2 μg/ml donkey anti-goat Alexa Fluor 488
(Cat# A21206, Thermo Fisher Scientific), 2 μg/ml donkey anti-rabbit Alexa Fluor 568 (Cat# A10037,
Thermo Fisher Scientific), and 1 U/ml CF405 phalloidin (Cat# 00034, Biotium). Tissue was washed three
times in calcium/magnesium-free HBSS and mounted on a glass slide in ~50 μl of Vectashield and
covered with a #1.5 thickness 22 x 22 mm cover glass (Cat# 2850-22, Corning).

Rhodamine-actin labeling
Labeling of free barbed ends of actin filaments was performed as outlined [39]. Inner ears from C57BL/6J
mice or mutant mice littermates dissected at the indicated ages in cold Hank's balanced salt solution
(Cat# 14025076, Thermo Fisher Scientific) supplemented with 5 mM HEPES, pH 7.4 (dissection buffer).
Cochlea were dissected and lateral wall, Reissner's membrane, and tectorial membrane were removed.
Live tissue was then briefly washed in cytoskeletal buffer (20 mM HEPES, pH 7.4, 138 mM KCl, 4 mM
$MgCl_2$, 3 mM EGTA, 1% bovine serum albumin (Cat# A3803, Sigma Aldrich), 2 mM ATP (Cat#
10127531001, Sigma Aldrich), and 0.05% saponin (Cat# S7900, Sigma Aldrich) before incubating for
5 minutes at room temperature in 40 ng/μl rhodamine-labeled actin (Cat# AR05, Cytoskeleton, Inc.)
freshly prepared in cytoskeletal buffer. Samples were then washed in cytoskeletal buffer and fixed for
1 hour at room temperature in 4% formaldehyde (Cat# 1570, Electron Microscopy Sciences) in PBS.
Tissues were rinsed twice in PBS, then incubated for 1 hour at room temperature in 0.4 U/ml Alexa Fluor
488 phalloidin (Cat# A12379, Thermo Fisher Scientific) diluted in PBS with 0.05% saponin. Tissue was
washed three times in PBS and mounted on a glass slide in ~50 μl of Vectashield and covered with a
#1.5 thickness 22 x 40 mm cover glass (Cat# CLS-1764-2240, Fisher Scientific).

Fluorescence microscopy
Structured illumination (SIM) images were acquired at 26-30°C with a 63X 1.4 NA oil immersion lens on
a Zeiss lattice-based Elyra 7 microscope with dual PCO.edge 4.2 sCMOS cameras for detection.
Illumination grid selection and z-spacing was guided by the software and kept consistent across images.
Post-acquisition processing was performed with software-recommended standard filtering for the 488 nm
channel, without baseline subtraction and with "scale to raw" checked. Contrast was manually adjusted
to retain both dim and bright structures due to the high dynamic range of the phalloidin signal. Verification
of channel alignment and measurement of the microscope point-spread function was carried out as
previously described [46].

Airyscan images were acquired at room temperature using a 63x, 1.4 NA Plan-Apochromat objective on
a Zeiss 3-channel LSM980 laser-scanning confocal microscope equipped with an Airyscan.2 detector
and run under ZEISS ZEN (v3.1, 64-bit software; Zeiss) acquisition software. Settings for x-y pixel
resolution, z-spacing, as well as pinhole diameter and grid selection, were set according to software-
suggested settings for optimal Nyquist-based resolution. Raw data processing for Airyscan-acquired
images was performed using manufacturer-implemented automated settings. Display adjustments in
brightness and contrast and reslices and/or maximum intensity Z-projections were made in Fiji software
(https://imagej.net/software/fiji/).

For cochlea imaging, for each stain, 2-4 images were acquired from 1-2 cochlea per genotype per age
for each experiment, and experiments were conducted three to four times. Ears from control and mutant
littermates or from different ages of C57BL/6J mice (of both sexes) were stained and imaged on the same
days for each experiment to limit variability. Genotyping was performed either prior to dissection or
performed on tails collected during dissection for younger animals (<P8). Genotypes were known by the
experimenter during staining and image acquisition. During image acquisition, the gain and laser settings
for the antibody and phalloidin signals were adjusted to reveal the staining pattern in control samples,
and the corresponding KO samples used the same settings. Image acquisition parameters and display
adjustments were kept constant across ages and genotypes for every antibody/fluorophore combination.

Image rendering and measurements of stereocilia dimensions
Surface rendering was carried out with Imaris software (imaris.oxinst.com; versions 9.7.0, 9.8.0, 9.9.0;
Oxford Instruments); the rendered surfaces were stored as sparse voxel octrees [97], which allowed them
to be manipulated and measured using Imaris. A batch processing file was created in Imaris and applied
to each dataset. Within each batch, background subtraction was first performed in the image processing
tab using a filter width of 1 μm for the phalloidin channel for all images. Volumetric surface areas of the
stereocilia in each image were then created from the phalloidin channel using creation parameters guided
by the program and edited as needed so that the phalloidin signal for each stereocilium was accurately
surrounded by a surface. Due to the high dynamic range of the signal, and the low intensity of the signal
in control row 3 or 4 stereocilia, creation parameters were guided to create the most accurate surfaces
for row 1 and 2, which often left some low intensity row 3 or 4 stereocilia undetected. Manual cutting of
any connected stereocilia surfaces was performed so that each stereocilium was modeled by a unique
surface. For any visible stereocilia lacking a surface (most often in rows 3 or 4), a surface was created
using the magic wand tool in the Create tab of Imaris. If the intensity of the stereocilium was too similar
to the phalloidin intensity of the surrounding cuticular plate, a surface could often not be generated. Thus,
row 3 and 4 stereocilia are only sparsely represented in older control hair bundles. Stereocilia surfaces
were labeled as corresponding to each stereocilia row (1, 2, 3, or 4+) and as corresponding to each cell
in the image. Measurements of Volume, BoundingBoxOO (object-oriented) Lengths, Number, and
Position (x,y) for all labeled surfaces were exported from Imaris to Microsoft Excel. BoundingBoxOO

lengths are generated within Imaris by considering the minimal rectangular box which fully encloses each
object with no constraints to the orientation; the dimensions of that box are reported as Lengths A, B, and
C, with A being the shortest principal axis and C the longest principal axis. To measure length and width
of each stereocilium, object-oriented BoundingBoxOO Length C and the average of lengths A and B were
used, respectively. For volume, we used the surface volume measurement provided by Imaris, which
quantifies the amount of space each object occupies. The number of stereocilia and the sum of stereocilia
volume per cell were calculated in Excel from the Imaris output. To measure the center 10 stereocilia, 10
neighboring row 1 and row 2 stereocilia at the center of each bundle were selected within Imaris, and
measurements for this subset were exported to Excel. Position (x,y) data could be used to determine
neighboring stereocilia and calculate differences in dimensions for adjacent stereocilia within each row
or adjacent row 1 and 2 stereocilia (Fig 5).

Three-dimensional rendering of stereocilia and EPS8 and GPSM2 tip staining (Fig 11O-R and Fig 12M-
P) from Airyscan z-stack images was also performed in Imaris using similar methods. Background
subtraction was first performed in the image processing tab using a filter width of 1 µm for all channels
for all images. Volumetric surface areas of the stereocilia in each image were then created from the
phalloidin channel using creation parameters guided by the program and edited as needed so that the
phalloidin signal for each stereocilium was accurately surrounded by a surface. Volumetric surface areas
of the EPS8 or GPSM2 signals in each image were also created in their respective channels using
creation parameters guided by the program and filtered to only include those with at least 0.01 overlap
volume ratio with the phalloidin stereocilia surfaces, so that only EPS8/GPSM2 surfaces at stereocilia
tips were included. All surfaces were manually edited to split any joining surfaces and create any surfaces
that were missed for low intensity phalloidin, EPS8, or GPSM2 signals. Once all surfaces were edited,
the two sets of surfaces were merged and the phalloidin and EPS8 or GPSM2 surfaces corresponding
to each individual stereocilium were joined in the edit tab, creating a surface that encompassed both the
phalloidin signal and tip complex signals for each stereocilium. E Each stereocilium surface was then
labeled according to its row position. Thus, for each stereocilium we measured both the length
(BoundingBoxOO Length C) and the corresponding intensity sum for each channel in the image, with the
channel 1 (568 nm) intensity sum measuring EPS8 amount and the channel 2 (488 nm) measuring
GPSM2 amount. These measurements were exported to Excel, pooled, and then plotted within Prism.

Quantitation of proteins at stereocilia tips
For Fig 9I-K, Fig 11A-N, Fig 12A-L, Fig 13M-R, and Fig 14G-H, quantitation of protein expression at row 1
and 2 tips was carried out after importing Airyscan z-stacks into Fiji. For analysis of C57BL/6J mice
throughout development, CAPZB and EPS8L2 in P21.5 mutants, and rhodamine-actin signal, summed
z-projections of Airyscan stacks were made that included row 1 and row 2 tips in the same projection.
For samples in which row 1 and 2 were in separate Z planes, separate Z-projections were made for row
1 and row 2 tips. Regions of interest (ROIs) were selected at 10 (or more) row 1, and 10 (or more) row 2

tips per hair bundle. ROIs were circles that encompassed most of each tip. Using Fiji's Measure function,
the area and mean gray value were measured for each ROI; measurements were also made outside the
stereocilia and above the epithelium to determine background. The total fluorescence signal in a tip (tip
signal) was thus area • mean gray value • minus total signal for the background. Tip signals below the
background were assigned the value of 0, which were used in calculating averages.

For analysis of mutant mice in Figs 11-12, S4 Fig, and S5 Fig, due to increased length variability, regions
of interest (ROIs) were selected at row 1 or row 2 tips within individual x-y slices from the z-stacks. Images
were kept as multi-channel stacks and phalloidin was used to guide selection at the tip of each
stereocilium. For these images, all row 1 and 2 stereocilia were measured within each cell. Measurements
and calculation of tip signal were otherwise carried out as above. Mean intensities were calculated as the
average tip intensity for all row 1 and row 2 stereocilia within each cell. Mean intensities were normalized
to the average tip intensity for all row 1 and 2 stereocilia from all het cochlea imaged on the same day.
Histograms were created in Prism using frequency distributions of the number of values, with a bin width
of 0.2 µm. The distributions were fit with a least-squares regression for a sum of two Gaussians.

Data presentation and statistical analysis
For quantitation of length, width, and adjacent stereocilia length differences in Fig 1, we derived the data
from the center ten stereocilia from each row. Volumes were instead calculated from Imaris volume
measurements from all stereocilia in each hair bundle. The data were derived from 2 litters per time point,
2 cochleas per litter, 1 image per cochlea, and 3-6 cells per image; thus n (the number of cells) was 12-
24, and N (the number of cochleas examined) was 4. Each point represents the mean ± standard
deviation for all cell averages from all cochleas (i.e., n samples). Blinding of sample age was not possible
as the age was obvious from the bundle and tissue morphology.

For quantitation in Figs 3-8, we compared measurements of dimensions between heterozygote and
knockout hair cells. We carried out volume, length, and width measurements for each stereocilium, then
derived adjacent stereocilia length and volume sum measurements. Blinding of samples was not possible
as the mutant phenotypes are easily discerned. To avoid problems with pseudoreplication [53, 54, 98],
we considered the five hierarchical levels in our data sets—litter, animal, cochlea, cell, and stereocilium.
The two cochleas from each animal were analyzed independently; we did not track individual animals.
Cochlea mounting, intensity of labeling reagents, distance from the coverslip (potentially leading to
spherical aberration and decrease in signal), and exact position along the apical-basal gradient all
substantially affected the measurements and had a greater influence than animal or litter. For length,
width, and volume measurements, measurements from all stereocilia of each row were averaged,
providing a single measurement per cell. For these figures, all data were derived from 2-3 litters per
genotype, 1-2 cochleas per litter, 2 images per cochlea, 4 total cochlea images, 5-6 cells per image, all
row 1 and 2 stereocilia for each hair cell, and some row 3 stereocilia for each hair cell. For each condition,
n was 23-24 and N was 4. For length and width (and their corresponding CVs), we used the nested

(hierarchical) t-test function in Prism to compare the results from heterozygote and knockout cochleas;

while comparing the results from the 4 cochleas per condition, the nested t-test approach takes into

account the structure of the data, i.e., the variance in individual cell measurements for each condition

[54]. For comparisons of dimensions of *Pcdh15^{av3J};Myo15a^{sh2}* genotypes (Fig 6), as well as volume

measurements for mutants and C57BL/6 (Fig 8), we used the nested one-way ANOVA function in Prism,

adjusting for multiple comparisons using the Šidák correction [99].

For comparisons of intensity of row 1 tip proteins (Figs 11-12, S4 Fig), as well as rhodamine-actin labeling

at row 1 and 2 tips (Fig 13) and row 2 protein intensities (Fig 14), we used nested t-tests as above. For

comparisons of intensity of row 1 tip proteins in *Pcdh15^{av3J};Tmie^{KO}* genotypes (S5 Fig), we used ordinary

one-way ANOVA tests with the Tukey correction for multiple comparisons.

Data reporting

Underlying data (e.g., measurements of individual stereocilia in hair bundles) and aggregated data used

to construct figures have been deposited in figshare:

Krey, JF, and Barr-Gillespie, PG. Control of stereocilia length during development of hair
bundles. 2022. figshare. 10.6084/m9.figshare.21632636

Acknowledgements

We thank Jennifer Goldsmith for mouse husbandry and Michael Bateschell for laboratory support. John

Brigande and Ulrich Müller provided helpful comments on the manuscript. We utilized the following core

facilities: lattice SIM, confocal microscopy, and Imaris software access from the OHSU Advanced Light

Microscopy Core @ The Jungers Center, and electron microscopy from the OHSU Multiscale Microscopy

Core. The authors declare no competing financial interests.

References

1. Tilney LG, Tilney MS, DeRosier DJ. Actin filaments, stereocilia, and hair cells: how cells count and measure. Ann. Rev. Cell Biol. 1992;8: 257–274.

2. Qiu X, Müller U. Sensing sound: Cellular specializations and molecular force sensors. Neuron. 2022S0896–6273(22)00857.

3. Shotwell SL, Jacobs R, Hudspeth AJ. Directional sensitivity of individual vertebrate hair cells to controlled deflection of their hair bundles. Ann N Y Acad Sci. 1981;374: 1–10.

4. Hudspeth AJ. Hair-bundle mechanics and a model for mechanoelectrical transduction by hair cells. Soc. Gen. Physiol. Ser. 1992;47: 357–370.

5. Fettiplace R. Hair cell transduction, tuning, and synaptic transmission in the mammalian cochlea. Compr Physiol. 2017;7: 1197–1227.

6. Krey JF, Chatterjee P, Dumont RA, O'Sullivan M, Choi D, Bird JE, et al. Mechanotransduction-dependent control of stereocilia dimensions and row identity in inner hair cells. Curr Biol. 2020;30: 442–454.e7.

7. Tarchini B, Tadenev AL, Devanney N, Cayouette M. A link between planar polarity and staircase-like bundle architecture in hair cells. Development. 2016;143: 3926–3932.

8. Mauriac SA, Hien YE, Bird JE, Carvalho SD, Peyroutou R, Lee SC, et al. Defective Gpsm2/Gαi3 signalling disrupts stereocilia development and growth cone actin dynamics in Chudley-McCullough syndrome. Nat Commun. 2017;8: 14907.

9. Tadenev ALD, Akturk A, Devanney N, Mathur PD, Clark AM, Yang J, et al. GPSM2-GNAI specifies the tallest stereocilia and defines hair bundle row identity. Curr Biol. 2019;29: 921–934.e4.

10. Lin L, Shi Y, Wang M, Wang C, Lu Q, Zhu J, et al. Phase separation-mediated condensation of Whirlin-Myo15-Eps8 stereocilia tip complex. Cell Reports. 2021;34: 108770.

11. Shi Y, Lin L, Wang C, Zhu J. Promotion of row 1–specific tip complex condensates by Gpsm2-Gαi provides insights into row identity of the tallest stereocilia. Science Advances. 2022;8: eabn4556.

12. Stepanyan R, Frolenkov GI. Fast adaptation and Ca^{2+} sensitivity of the mechanotransducer require myosin-XVa in inner but not outer cochlear hair cells. J Neurosci. 2009;29: 4023–4034.

13. Peng AW, Belyantseva IA, Hsu PD, Friedman TB, Heller S. Twinfilin 2 regulates actin filament lengths in cochlear stereocilia. J Neurosci. 2009;29: 15083–15088.

14. Avenarius MR, Krey JF, Dumont RA, Morgan CP, Benson CB, Vijayakumar S, et al. Heterodimeric capping protein is required for stereocilia length and width regulation. J Cell Biol. 2017;216: 3861–3881.

15. Furness DN, Johnson SL, Manor U, Ruttiger L, Tocchetti A, Offenhauser N, et al. Progressive hearing loss and gradual deterioration of sensory hair bundles in the ears of mice lacking the actin-binding protein Eps8L2. Proc Natl Acad Sci U S A. 2013;110: 13898–13903.

16. Fettiplace R, Kim KX. The physiology of mechanoelectrical transduction channels in hearing. Physiol Rev. 2014;94: 951–986.

17. Vélez-Ortega AC, Freeman MJ, Indzhykulian AA, Grossheim JM, Frolenkov GI. Mechanotransduction current is essential for stability of the transducing stereocilia in mammalian auditory hair cells. eLife. 2017;6:

18. Zhao B, Wu Z, Grillet N, Yan L, Xiong W, Harkins-Perry S, et al. TMIE is an essential component of
the mechanotransduction machinery of cochlear hair cells. Neuron. 2014;84: 954–967.

19. Kawashima Y, Geleoc GS, Kurima K, Labay V, Lelli A, Asai Y, et al. Mechanotransduction in mouse
inner ear hair cells requires transmembrane channel-like genes. J Clin Invest. 2011;121: 4796–4809.

20. Pan B, Akyuz N, Liu XP, Asai Y, Nist-Lund C, Kurima K, et al. TMC1 forms the pore of
mechanosensory transduction channels in vertebrate inner ear hair cells. Neuron. 2018;99: 736–
753.e6.

21. Beurg M, Cui R, Goldring AC, Ebrahim S, Fettiplace R, Kachar B. Variable number of TMC1-
dependent mechanotransducer channels underlie tonotopic conductance gradients in the cochlea.
Nat Commun. 2018;9: 2185.

22. Beurg M, Fettiplace R, Nam JH, Ricci AJ. Localization of inner hair cell mechanotransducer channels
using high-speed calcium imaging. Nat Neurosci. 2009;12: 553–558.

23. Cunningham CL, Qiu X, Wu Z, Zhao B, Peng G, Kim YH, et al. TMIE Defines Pore and Gating
Properties of the Mechanotransduction Channel of Mammalian Cochlear Hair Cells. Neuron.
2020;107: 126–143.e8.

24. Ahmed ZM, Goodyear R, Riazuddin S, Lagziel A, Legan PK, Behra M, et al. The tip-link antigen, a
protein associated with the transduction complex of sensory hair cells, is protocadherin-15. J
Neurosci. 2006;26: 7022–7034.

25. Siemens J, Lillo C, Dumont RA, Reynolds A, Williams DS, Gillespie PG, et al. Cadherin 23 is a
component of the tip link in hair-cell stereocilia. Nature. 2004;*428*: 950–955.

26. Kazmierczak P, Sakaguchi H, Tokita J, Wilson-Kubalek EM, Milligan RA, Müller U, et al. Cadherin 23
and protocadherin 15 interact to form tip-link filaments in sensory hair cells. Nature. 2007;449: 87–
91.

27. Boeda B, El-Amraoui A, Bahloul A, Goodyear R, Daviet L, Blanchard S, et al. Myosin VIIa, harmonin
and cadherin 23, three Usher I gene products that cooperate to shape the sensory hair cell bundle.
EMBO J. 2002;21: 6689–6699.

28. Senften M, Schwander M, Kazmierczak P, Lillo C, Shin JB, Hasson T, et al. Physical and functional
interaction between protocadherin 15 and myosin VIIa in mechanosensory hair cells. J Neurosci.
2006;26: 2060–2071.

29. Grati M, Kachar B. Myosin VIIa and sans localization at stereocilia upper tip-link density implicates
these Usher syndrome proteins in mechanotransduction. Proc Natl Acad Sci U S A. 2011;108:
11476–11481.

30. Richardson GP, de Monvel JB, Petit C. How the genetics of deafness illuminates auditory physiology.
Annu Rev Physiol. 2011;73: 311–334.

31. Holme RH, Steel KP. Stereocilia defects in waltzer (Cdh23), shaker1 (Myo7a) and double
waltzer/shaker1 mutant mice. Hear Res. 2002;169: 13–23.

32. Lefevre G, Michel V, Weil D, Lepelletier L, Bizard E, Wolfrum U, et al. A core cochlear phenotype in
USH1 mouse mutants implicates fibrous links of the hair bundle in its cohesion, orientation and
differential growth. Development. 2008;135: 1427–1437.

33. Alagramam KN, Goodyear RJ, Geng R, Furness DN, van Aken AF, Marcotti W, et al. Mutations in
protocadherin 15 and cadherin 23 affect tip links and mechanotransduction in mammalian sensory
hair cells. PLoS One. 2011;6: e19183.

34. Assad JA, Shepherd GMG, Corey DP. Tip-link integrity and mechanical transduction in vertebrate
hair cells. Neuron. 1991;7: 985–994.

35. Kachar B, Parakkal M, Kurc M, Zhao Y, Gillespie PG. High-resolution structure of hair-cell tip links.
Proc. Natl. Acad Sci. USA. 2000;97: 13336–13341.

36. Rzadzinska AK, Schneider ME, Davies C, Riordan GP, Kachar B. An actin molecular treadmill and
myosins maintain stereocilia functional architecture and self-renewal. J. Cell Biol. 2004;164: 887–
897.

37. Caberlotto E, Michel V, de Monvel JB, Petit C. Coupling of the mechanotransduction machinery and
F-actin polymerization in the cochlear hair bundles. Bioarchitecture. 2011;1: 169–174.

38. Narayanan P, Chatterton P, Ikeda A, Ikeda S, Corey DP, Ervasti JM, et al. Length regulation of
mechanosensitive stereocilia depends on very slow actin dynamics and filament-severing proteins.
Nat Commun. 2015;6: 6855.

39. McGrath J, Tung CY, Liao X, Belyantseva IA, Roy P, Chakraborty O, et al. Actin at stereocilia tips is
regulated by mechanotransduction and ADF/cofilin. Curr Biol. 2021;31: 1141–1153.e7.

40. Bamburg JR. Proteins of the ADF/cofilin family: essential regulators of actin dynamics. Annual review
of cell and developmental biology. 1999;15: 185.

41. Krey JF, Sherman NE, Jeffery ED, Choi D, Barr-Gillespie PG. The proteome of mouse vestibular hair
bundles over development. Sci Data. 2015;2: 150047.

42. Barr-Gillespie PG. Assembly of hair bundles, an amazing problem for cell biology. Mol Biol Cell.
2015;26: 2727–2732.

43. Miller KK, Atkinson P, Mendoza KR, Ó Maoiléidigh D, Grillet N. Dimensions of a living cochlear hair
bundle. Front Cell Dev Biol. 2021;9: 742529.

44. Betzig E. Excitation strategies for optical lattice microscopy. Opt Express. 2005;13: 3021–3036.

45. Richter V, Piper M, Wagner M, Schneckenburger H. Increasing resolution in live cell microscopy by
structured illumination (SIM). Applied Sciences. 2019;9: 1188.

46. Krey JF, Liu C, Belyantseva IA, Bateschell M, Dumont RA, Goldsmith J, et al. ANKRD24 organizes
TRIOBP to reinforce stereocilia insertion points. Journal of Cell Biology. 2022;221: e202109134.

47. Cole RW, Jinadasa T, Brown CM. Measuring and interpreting point spread functions to determine
confocal microscope resolution and ensure quality control. Nat Protoc. 2011;6: 1929–1941.

48. Waguespack J, Salles FT, Kachar B, Ricci AJ. Stepwise morphological and functional maturation of
mechanotransduction in rat outer hair cells. J Neurosci. 2007;27: 13890–13902.

49. Alagramam KN, Murcia CL, Kwon HY, Pawlowski KS, Wright CG, Woychik RP. The mouse Ames
waltzer hearing-loss mutant is caused by mutation of Pcdh15, a novel protocadherin gene. Nat Genet.
2001;27: 99–102.

50. Di Palma F, Holme RH, Bryda EC, Belyantseva IA, Pellegrino R, Kachar B, et al. Mutations in Cdh23,
encoding a new type of cadherin, cause stereocilia disorganization in waltzer, the mouse model for
Usher syndrome type 1D. Nat Genet. 2001;27: 103–107.

51. Zheng QY, Scarborough JD, Zheng Y, Yu H, Choi D, Gillespie PG. Digenic inheritance of deafness
caused by 8J allele of myosin-VIIA and mutations in other Usher I genes. Hum Mol Genet. 2012;21:
2588–2598.

52. Lelli A, Asai Y, Forge A, Holt JR, Geleoc GS. Tonotopic gradient in the developmental acquisition of
sensory transduction in outer hair cells of the mouse cochlea. J Neurophysiol. 2009;101: 2961–2973.

53. Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK. SuperPlots: Communicating reproducibility and
variability in cell biology. J Cell Biol. 2020;219: e202001064.

54. Eisner DA. Pseudoreplication in physiology: More means less. J Gen Physiol. 2021;153:
e202012826.

55. Hadi S, Alexander AJ, Vélez-Ortega AC, Frolenkov GI. Myosin-XVa controls both staircase
architecture and diameter gradation of stereocilia rows in the auditory hair cell bundles. J Assoc Res
Otolaryngol. 2020;21: 121–135.

56. Tilney LG, Tilney MS. The actin filament content of hair cells of the bird cochlea is nearly constant
even though the length, width//number of stereocilia vary depending on the hair cell location. J. Cell
Biol. 1988;107: 2563–2574.

57. Beyer LA, Odeh H, Probst FJ, Lambert EH, Dolan DF, Camper SA, et al. Hair cells in the inner ear of
the pirouette and shaker 2 mutant mice. J Neurocytol. 2000;29: 227–240.

58. Shin JB, Longo-Guess CM, Gagnon LH, Saylor KW, Dumont RA, Spinelli KJ, et al. The R109H variant
of fascin-2, a developmentally regulated actin crosslinker in hair-cell stereocilia, underlies early-onset
hearing loss of DBA/2J mice. J Neurosci. 2010;30: 9683–9694.

59. Avenarius MR, Saylor KW, Lundeberg MR, Wilmarth PA, Shin JB, Spinelli KJ, et al. Correlation of
actin crosslinker and capper expression levels with stereocilia growth phases. Mol Cell Proteomics.
2014;13: 606–620.

60. Zhu Y, Scheibinger M, Ellwanger DC, Krey JF, Choi D, Kelly RT, et al. Single-cell proteomics reveals
changes in expression during hair-cell development. Elife. 2019;8:

61. Krey JF, Krystofiak ES, Dumont RA, Vijayakumar S, Choi D, Rivero F, et al. Plastin 1 widens
stereocilia by transforming actin filament packing from hexagonal to liquid. J Cell Biol. 2016;215: 467–
482.

62. Zheng L, Sekerkova G, Vranich K, Tilney LG, Mugnaini E, Bartles JR. The deaf jerker mouse has a
mutation in the gene encoding the espin actin-bundling proteins of hair cell stereocilia and lacks
espins. Cell. 2000;102: 377–385.

63. Sekerkova G, Richter CP, Bartles JR. Roles of the espin actin-bundling proteins in the morphogenesis
and stabilization of hair cell stereocilia revealed in CBA/CaJ congenic jerker mice. PLoS Genet.
2011;7: e1002032.

64. Roy P, Perrin BJ. The stable actin core of mechanosensory stereocilia features continuous turnover
of actin cross-linkers. Mol Biol Cell. 2018;29: 1856–1865.

65. Kachar B, Battaglia A, Fex J. Compartmentalized vesicular traffic around the hair cell cuticular plate.
Hear Res. 1997;107: 102–112.

66. Hertzog M, Milanesi F, Hazelwood L, Disanza A, Liu H, Perlade E, et al. Molecular basis for the dual
function of Eps8 on actin dynamics: bundling and capping. PLoS Biol. 2010;8: e1000387.

67. Gong R, Jiang F, Moreland ZG, Reynolds MJ, de Los Reyes SE, Gurel P, et al. Structural basis for
tunable control of actin dynamics by myosin-15 in mechanosensory stereocilia. Sci Adv. 2022;8:
eabl4733.

68. Akturk A, Day M, Tarchini B. RGS12 polarizes the GPSM2-GNAI complex to organize and elongate
stereocilia in sensory hair cells. Science Advances. 2022;8: eabq2826.

69. Probst FJ, Fridell RA, Raphael Y, Saunders TL, Wang A, Liang Y, et al. Correction of deafness in shaker-2 mice by an unconventional myosin in a BAC transgene. Science. 1998;280: 1444–1447.

70. Manor U, Disanza A, Grati M, Andrade L, Lin H, Di Fiore PP, et al. Regulation of stereocilia length by myosin XVa and whirlin depends on the actin-regulatory protein Eps8. Curr Biol. 2011;21: 167–172.

71. Mburu P, Mustapha M, Varela A, Weil D, El-Amraoui A, Holme RH, et al. Defects in whirlin, a PDZ domain molecule involved in stereocilia elongation, cause deafness in the whirler mouse and families with DFNB31. Nat Genet. 2003;34: 421–428.

72. Sotomayor M, Weihofen WA, Gaudet R, Corey DP. Structure of a force-conveying cadherin bond essential for inner-ear mechanotransduction. Nature. 2012;492: 128–132.

73. Choudhary D, Narui Y, Neel BL, Wimalasena LN, Klanseck CF, De-la-Torre P, et al. Structural determinants of protocadherin-15 mechanics and function in hearing and balance perception. Proc Natl Acad Sci U S A. 2020;117: 24837–24848.

74. Michel V, Goodyear RJ, Weil D, Marcotti W, Perfettini I, Wolfrum U, et al. Cadherin 23 is a component of the transient lateral links in the developing hair bundles of cochlear sensory cells. Dev Biol. 2005;280: 281–294.

75. Gillespie PG, Dumont RA, Kachar B. Have we found the tip link, transduction channel, and gating spring of the hair cell? Curr Opin Neurobiol. 2005;15: 389–396.

76. Goodyear RJ, Marcotti W, Kros CJ, Richardson GP. Development and properties of stereociliary link types in hair cells of the mouse cochlea. J Comp Neurol. 2005;485: 75–85.

77. Michalski N, Michel V, Bahloul A, Lefevre G, Barral J, Yagi H, et al. Molecular characterization of the ankle-link complex in cochlear hair cells and its role in the hair bundle functioning. J Neurosci. 2007;27: 6478–6488.

78. Crawley SW, Shifrin DAJ, Grega-Larson NE, McConnell RE, Benesh AE, Mao S, et al. Intestinal brush border assembly driven by protocadherin-based intermicrovillar adhesion. Cell. 2014;157: 433–446.

79. Crawley SW, Mooseker MS, Tyska MJ. Shaping the intestinal brush border. J Cell Biol. 2014;207: 441–451.

80. Legan PK, Lukashkina VA, Goodyear RJ, Kössi M, Russell IJ, Richardson GP. A targeted deletion in alpha-tectorin reveals that the tectorial membrane is required for the gain and timing of cochlear feedback. Neuron. 2000;28: 273–285.

81. Avan P, Le Gal S, Michel V, Dupont T, Hardelin JP, Petit C, et al. Otogelin, otogelin-like, and stereocilin form links connecting outer hair cell stereocilia to each other and the tectorial membrane. Proc Natl Acad Sci U S A. 2019;116: 25948–25957.

82. Jacobs RA, Hudspeth AJ. Ultrastructural correlates of mechanoelectrical transduction in hair cells of the bullfrog's internal ear. Cold Spring Harb. Symp. Quant. Biol. 1990;55: 547–561.

83. Corey DP, Hudspeth AJ. Kinetics of the receptor current in bullfrog saccular hair cells. J. Neurosci. 1983;3: 962–976.

84. Kindt KS, Finch G, Nicolson T. Kinocilia mediate mechanosensitivity in developing zebrafish hair cells. Dev Cell. 2012;23: 329–341.

85. Riazuddin S, Belyantseva IA, Giese APJ, Lee K, Indzhykulian AA, Nandamuri SP, et al. Alterations of the CIB2 calcium-and integrin-binding protein cause Usher syndrome type 1J and nonsyndromic deafness DFNB48. Nature genetics. 2012;44: 1265–1271.

86. Giese APJ, Tang YQ, Sinha GP, Bowl MR, Goldring AC, Parker A, et al. CIB2 interacts with TMC1
and TMC2 and is essential for mechanotransduction in auditory hair cells. Nat Commun. 2017;8: 43.

87. Liang X, Qiu X, Dionne G, Cunningham CL, Pucak ML, Peng G, et al. CIB2 and CIB3 are auxiliary
subunits of the mechanotransduction channel of hair cells. Neuron. 2021;109: 2131–2149.e15.

88. Jeong H, Clark S, Goehring A, Dehghani-Ghahnaviyeh S, Rasouli A, Tajkhorshid E, et al. Structures
of the TMC-1 complex illuminate mechanosensory transduction. Nature. 2022;610: 796–803.

89. Kurima K, Ebrahim S, Pan B, Sedlacek M, Sengupta P, Millis BA, et al. TMC1 and TMC2 localize at
the site of mechanotransduction in mammalian inner ear hair cell stereocilia. Cell Rep. 2015;12:
1606–1617.

90. Johnson SL, Kennedy HJ, Holley MC, Fettiplace R, Marcotti W. The resting transducer current drives
spontaneous activity in prehearing mammalian cochlear inner hair cells. J Neurosci. 2012;32: 10479–
10483.

91. Beurg M, Goldring AC, Ricci AJ, Fettiplace R. Development and localization of reverse-polarity
mechanotransducer channels in cochlear hair cells. Proc Natl Acad Sci U S A. 2016;113: 6767–6772.

92. Tilney LG, Cotanche DA, Tilney MS. Actin filaments, stereocilia and hair cells of the bird cochlea. VI.
How the number and arrangement of stereocilia are determined. Development. 1992;116: 213–226.

93. Peskin CS, Odell GM, Oster GF. Cellular motions and thermal fluctuations: the Brownian ratchet.
Biophys J. 1993;65: 316–324.

94. Prost J, Barbetta C, Joanny JF. Dynamical control of the shape and size of stereocilia and microvilli.
Biophys J. 2007;93: 1124–1133.

95. Hequembourg S, Liberman MC. Spiral ligament pathology: a major aspect of age-related cochlear
degeneration in C57BL/6 mice. J Assoc Res Otolaryngol. 2001;2: 118–129.

96. Indzhykulian AA, Stepanyan R, Nelina A, Spinelli KJ, Ahmed ZM, Belyantseva IA, et al. Molecular
remodeling of tip links underlies mechanosensory regeneration in auditory hair cells. PLoS Biol.
2013;11: e1001583.

97. Laine S, Karras T. Efficient sparse voxel octrees. IEEE Trans Vis Comput Graph. 2011;17: 1048–
1059.

98. Colegrave N, Ruxton GD. Using Biological Insight and Pragmatism When Thinking about
Pseudoreplication. Trends Ecol Evol. 2018;33: 28–35.

99. Šidák Z. Rectangular confidence regions for the means of multivariate normal distributions. Journal
of the American Statistical Association. 1967;62: 626–633.

1385

Supporting Information Legends

S1 Fig. Variability in stereocilia length and width measurements.

A, CVs for stereocilia length from all stereocilia at indicated ages. Blue lines in panels A-D show
developmental trends for row 1 length or width CV. *B*, CVs for stereocilia length from center ten
stereocilia at indicated ages. P values from unpaired Student's t-tests comparing all row 1
stereocilia versus center ten row 1 stereocilia: P0.5, 0.1861; P1.5-P21.5, <0.0001. P values from
unpaired Student's t-tests comparing length CVs for all row 2 stereocilia versus center ten row 2
stereocilia: P0.5, 0.4933; P1.5-P21.5, <0.0001. *C*, CVs for stereocilia width from all stereocilia
at indicated ages. *D*, CVs for stereocilia width from center ten stereocilia at indicated ages.
P values from unpaired Student's t-tests comparing length CVs for all row 1 stereocilia versus
center ten row 1 stereocilia: P0.5, 0.0473; P1.5, <0.0001; P2.5, 0.0893; P3.5-P4.5, <0.0001;
P5.5, 0.0001; P7.5, <0.0001; P10.5, 0.0038; P13.5, 0.0823; P16.5, 0.0126; P21.5, 0.118.
P values from unpaired Student's t-tests comparing all row 2 stereocilia versus center ten row 2
stereocilia: P0.5-P21.5, <0.0001. *E-G*, Models of stereocilia arrangement. A peripheral-to-
central gradient of stereocilia length for row 1 and row 2 was seen at P7.5 (F; red gradient
symbol), rather than uniform lengths (E; red line). Stereocilia lengths largely equalized by P21.5
(G).

S2 Fig. CDH23 and PCDH15 immunoreactivity in mouse tip-link mutants.

A-B, CDH23 and PCDH15 localization in *Cdh23^{v2J}/+* (A) and *Cdh23$^{v2J/v3J}$* (B) IHCs. *C-D*, CDH23
and PCDH15 localization in *Pcdh15^{av3J}/+* (C) and *Pcdh15$^{av3J/av3J}$* (D) IHCs. Panel widths: 30 μm.

S3 Fig. Localization of TMC1-HA and TMC2-Myc in *Pcdh15$^{av3J/av3J}$* mutant IHCs.

A, TMC1-HA in P7 *Tmc1$^{HA/HA}$;Pcdh15^{av3J}/+* cochlea at the hair-bundle level. *B*, TMC1-HA in
same P7 *Tmc1$^{HA/HA}$;Pcdh15^{av3J}/+* cochlea at the soma level. *C*, TMC1-HA in P7
Tmc1$^{HA/HA}$;Pcdh15$^{av3J/av3J}$ cochlea (bundles). *D*, TMC1-HA in same P7 *Tmc1$^{HA/HA}$;Pcdh15$^{av3J/av3J}$*
cochlea (somas). *E*, TMC2-Myc in P4 *Tmc1$^{Myc/Myc}$;Pcdh15^{av3J}/+* cochlea (bundles). *F*, TMC2-
Myc in P4 *Tmc1$^{Myc/Myc}$;Pcdh15$^{av3J/av3J}$* cochlea (bundles). Panel widths: 35 μm.

S4 Fig. Localization of GNAI3 and WHRN in *Cdh23$^{v2J/v2J}$* and *Pcdh15$^{av3J/av3J}$* mutant IHCs.

A-L, Localization of row 1 complex proteins in *Cdh23^{v2J}* and *Pcdh15^{av3J}* heterozygotes and
homozygotes IHCs. Panel widths: 30 μm. *A-B*, GNAI3 and EPS8 in *Cdh23^{v2J}/+* and *Cdh23$^{v2J/v2J}$*
at P7.5. *C-D*, WHRN and EPS8 in *Cdh23^{v2J}/+* and *Cdh23$^{v2J/v2J}$* at P7.5. *E-F*, GNAI3 and EPS8
in *Cdh23^{v2J}/+* and *Cdh23$^{v2J/v2J}$* at P21.5. *G-H*, WHRN and EPS8 in *Cdh23^{v2J}/+* and *Cdh23$^{v2J/v2J}$*
at P21.5. *I-J*, GNAI3 and EPS8 in *Pcdh15^{av3J}/+* and *Pcdh15$^{av3J/av3J}$* at P21.5. *K-L*, WHRN and
EPS8 in *Pcdh15^{av3J}/+* and *Pcdh15$^{av3J/av3J}$* at P21.5. *M-P*, GNAI3 and WHRN normalized
fluorescence average intensity per hair bundle for all measured stereocilia (row 1 and 2) for
Cdh23^{v2J} and *Pcdh15^{av3J}* IHCs at P7.5 and P21.5. Intensities were normalized to the
heterozygote average for each genotype pair. Plotting and statistical testing were as in Fig 3. *Q-*
T, Frequency distribution of GNAI3 and WHRN tip intensity in *Cdh23^{v2J}* hair cells at P7.5 and
P21.5.

S5 Fig. *Pcdh15^(av3J)*;*Tmie^(KO)* double knockout mice.

A-D, Scanning electron micrographs showing P8.5 IHC hair bundles from *Pcdh15^(av3J)*/+;*Tmie^(KO)*/+

(Het;Het)   *Pcdh15^(av3J)*/+;*Tmie^(KO/KO)*   (Het;KO),   *Pcdh15^(av3J/av3J)*;*Tmie^(KO)*/+   (KO;Het),   and

Pcdh15^(av3J/av3J);*Tmie^(KO/KO)* (KO;KO) cochleas. ***E-H***, Localization of GNAI3 and EPS8 in IHCs of

Pcdh15^(av3J);*Tmie^(KO)* genotypes at P7.5. ***I***, EPS8 fluorescence average intensity per bundle for all

measured stereocilia in bundles of each genotype. ***J,*** Frequency distribution of EPS8 tip intensity

in bundles of each genotype. Het;Het and Het;KO distributions were fit with double Gaussians;

KO;Het and KO;KO distributions were fit with single Gaussians. ***K***, GNAI3 fluorescence average

intensity per bundle for all measured stereocilia in bundles of each genotype. ***L,*** Frequency

distribution of GNAI3 tip intensity in bundles of each genotype. Distributions from each genotype

were fit with double Gaussians. For statistical comparisons in I and K, we used ordinary one-

way ANOVA tests with the Tukey correction.

1436

Figure 1
Krey et al.

Figure 1

Figure 2
Krey et al.

Figure 2

Figure 3
Krey et al.

Figure 3

Figure 4
Krey et al.

Figure 4

Figure 5
Krey et al.

Figure 5

Figure 6
Krey et al.

Figure 6

Figure 7
Krey et al.

Figure 7

Figure 8
Krey et al.

Figure 8

Figure 9
Krey et al.

Figure 9

Figure 10
Krey et al.

Figure 10

Figure 11
Krey et al.

Figure 11

Figure 12
Krey et al.

Figure 12

Figure 13
Krey et al.

Figure 13

Figure 14
Krey et al.

Figure 14

Figure 15
Krey et al.

Figure 15