Utilizing co-abundances of antimicrobial resistance genes to identify potential co-selection in the resistome

Authors:

Hannah-Marie Martiny[1]

Patrick Munk[1]

Christian Brinch[1]

Frank M. Aarestrup[1]

M. Luz Calle[2]

Thomas N. Petersen[1]

Affiliations:

[1]Research Group for Genomic Epidemiology, Technical University of Denmark, Kongens Lyngby, Denmark

[2]Biosciences Department, Faculty of Sciences and Technology, University of Vic - Central University of Catalonia, Vic, Spain

Abstract

The rapid spread of antimicrobial resistance (AMR) is a threat to global health, and the nature of co-occurring antimicrobial resistance genes (ARGs) may cause collateral AMR effects once antimicrobial agents are used. Therefore, it is essential to identify which pairs of ARGs co-occur. Given the wealth of NGS data available in public repositories, we have investigated the correlation between ARG abundances in a collection of 214,095 metagenomic datasets. Using more than $6.76 \cdot 10^8$ read fragments aligned to ARGs to infer pairwise correlation coefficients, we found that more ARGs correlated with each other in human and animal sampling origins than in soil and water environments. Furthermore, we showed that the correlations serve as risk profiles of resistance co-occurring to critically important antimicrobials. Using these profiles, we found several key ARGs indirectly but

28 strongly selecting for ARGs of critical importance, such as tetracycline ARGs correlating with

29 most forms of resistances. In conclusion, this study highlights the important ARG players

30 indirectly involved in shaping the resistomes of various environments that can serve as

31 monitoring targets in AMR surveillance programs.

Introduction

Antimicrobial resistance (AMR) is one of the biggest threats to human and animal health[1,2],

and it is widely acknowledged that the misuse of antimicrobials has accelerated the

dissemination and prevalence of antimicrobial resistance genes (ARGs)[3]. Most attempts to

reduce the burden of AMR have focused on reducing the use of single classes of

antimicrobial agents considered of critical importance[4]. Despite considerable efforts in

various settings, such as livestock, these regulations have not significantly reduced the

spread of ARGs[5,6]. It is known that even after banning the use of specific antimicrobials, the

ARGs conferring resistance will still be prevalent in the environment[7]. Furthermore, studies

have shown that ARGs can be indirectly selected if they co-occur with an ARG conferring

resistance to the antimicrobial, causing the selective pressure[6,8–10]. We have also recently

observed that changes in the selective pressure of even a single antimicrobial agent

influence several ARGs in pig metagenomes[10]. It is, however, not known whether this is due

to co-selection due to the genetic linkage of ARGs, the presence of ARGs in the same

bacterial clones or because different bacterial species with different ARGs are co-selected in

a microbial network.

Most studies associating AMU and AMR have, however, focused on one antimicrobial agent

at that time and how the use impacts the development of resistance to that agent and not

to other unrelated antimicrobials. This needs to be understood better to stop these

collateral damage effects[11,12]. Co-occurrence of microbes has been evaluated in soil[13,14] and

marine environments[15,16], whereas ARG co-occurrences have been studied in sewage

sludge[17], freshwater[18], marine[19], swine[7], and cattle[20]. However, most of these studies have

only been evaluated on a smaller scale and not always using the same methods. There are

currently a large collection of next-generation sequencing datasets from metagenomic

samples available in public repositories, providing an excellent resource to quantify the

prevalence of ARGs by analyzing the read abundances [21–24].

In this study, we have studied the co-abundance of sequencing reads aligned to ARGs to

assess how resistance to one specific antimicrobial agent is linked to the abundance of

another class of antimicrobial. With a collection of 214,095 metagenomic datasets[25], we

examined the correlation between pairwise ARG read abundances with a compositional

approach[26,27] using SparCC (Sparse Correlations for Compositional data)[28]. Our results

demonstrated that many ARG pairs interact but are highly specific to the environment. We

believe that these interactions provide a new foundation to understand how ARGs are being

co-selected independently of the microbial context, and the findings can be used to design

targeted interventions to limit the spread of AMR.

Methods

Data collection and pre-processing

We have previously described in detail the process of downloading and analyzing 214,095

metagenomic samples[23,25], but in brief: We downloaded raw sequencing reads corresponding

to 442 Tbp from metagenomic samples deposited in the European Nucleotide Archive[29] that

were uploaded between 2010-01-01 and 2020-01-01 and had at least 100,000 reads and were

shotgun sequenced. The raw sequencing reads were quality-checked with FASTQC v.0.11.15

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and trimmed with BBduk2

36.49[30]. The trimmed reads were then globally aligned using KMA[31] 1.2.21 against two

reference sequence databases: ResFinder[32] (downloaded 25-01-2020) and Silva[33] (version 38,

downloaded 16-01-2020). ResFinder is a database of 3,085 acquired ARGs, whereas Silva is a

16S/18S rRNA database of 2,225,272 sequences.

Homology-reduced ResFinder

Since there is a possibility that KMA might assign reads to closely related homologs differently

across samples, we decided to homology reduce the 3,085 ARGs of ResFinder using

USEARCH[34] 11.0.667 with 90% nucleotide identity, producing a total of 716 ARG groups. For

each of these groups, the read counts for the genes in that group were first adjusted by the

length of the gene and then aggregated together. These new group counts are used in the

correlation analysis described in the next section. For the rest of the manuscript, we refer to

these 716 ARG clusters as simply ARGs.

Calculating relative abundances

92   For a category label, we calculated the relative abundance of fragment counts assigned to

93   different genes or classes as:

$$\text{Relative abundance}(x) = \frac{\kappa}{\sum n_i} n_i$$

95   where x is the label, n_i is the count of read fragments assigned to gene i, and $\kappa = 100$ is a

96   scaling constant.

Inferring pairwise correlations with SparCC

The SparCC algorithm[28] was used to obtain correlations using pairs of log-ratio transformed

ARG read counts to infer linear Pearson correlations. SparCC obtains linear Pearson

correlations and p-values through an iterative approach that adjusts for spurious

correlations and lowers the false discovery rate. The ARG-ARG correlations were inferred as

the average over 50 iterations, and one-sided pseudo p-values were obtained through a

bootstrapping procedure of 100 rounds. In each bootstrapping round, the input count

matrix was shuffled and correlations were averaged over 10 iterations to infer one-sided p-

values to test whether the correlation for the observed data was statistically significant.

Correlations values ≥ 0.6 with p-values ≤ 0.01 were selected for further analysis. We

implemented SparCC to run on GPUs on the Danish National Supercomputer for Life

Sciences (https://www.computerome.dk).

We ran SparCC on the entire dataset of the 214K metagenomic samples and subsets of

samples grouped by sampling host and environment, where at least 800 samples existed.

Due to inconsistent labeling of the sampling sources, we made new source groups, as shown

in Table 1. We only consider genes for SparCC analysis that are present in at least 10

samples with a minimum read fragment count of 50. In total, 11 different correlation

matrices were made, i.e., one for each of the source groups listed in Table 1. The correlation

networks were visualized in R 4.1.0[35] with packages igraph[36], qgraph[37], and ggraph[38] using

the Fruchterman-Reingold layout algorithm[39].

Source group	Sampling source label(s)	Number of samples	Number of samples with ARGs
Air	Air metagenome	914	870
Chicken	*Gallus gallus* (1,219)*,* chicken gut metagenome (4)	1,223	1,215
Cow	*Bos taurus* (872)*,* cow dung metagenome (14)	886	824
Dog	*Canis lupus familiaris*	3,439	3,182
Freshwater	Freshwater metagenome	4,494	585
Human	*Homo sapiens*	95,003	57,239
Marine	Marine metagenome	30,002	5,444
Mouse	*Mus musculus* (1,462), mouse metagenome (50), mouse gut metagenome (2,435)	3,947	3,909
Pig	*Sus scrofa* (673)*, Sus scrofa domesticus* (355)*,* pig metagenome (2,129), pig gut metagenome (72)	3,229	3,461
Soil	Soil metagenome	6,533	2,822
All		214,095	119,206

Table 1: Grouped labels for hosts and environments. The parenthesis after each sampling source label denotes the number of samples assigned to that label if the group consisted of multiple labels.

Network comparisons

The topology of the different networks is described using different metrics: the number of nodes (N) and edges (E), the global clustering coefficient, network density, edge density, and the number of components. The global clustering coefficient, or the graph transitivity, measures the density of node triplets in the network[40]. The network density is calculated as $2E/N(N-1)$ as given in Parente et al. (2018)[41]. Edge density is the number of edges over the number of possible edges[42]. The average correlation between ARGs of two antimicrobial classes was calculated using Fisher's z-transformation on the correlation values, averaging the z-scores and converting it back to a correlation score with the inverse Fisher transformation[43].

Data and code availability

The matrix of raw ARG read counts is available at https://zenodo.org/record/6919377. The

code used to run the analysis and create figures and the output files of SparCC are available

at https://github.com/hmmartiny/global_resistome_correlations. Classifications of

antimicrobial importance were retrieved from the 6th revision of critically important

antimicrobials for human medicine from

https://www.who.int/publications/i/item/9789241515528, accessed 2022-10-10.

Results

This study investigated the correlation of pairwise ARG abundances across a highly diverse set of 214,095 metagenomic samples. The samples included in the 214K metagenomic collection come from a variety of backgrounds, such as the sampling period spanning between 2000 and 2020 (Figure S1a), primarily sequenced on Illumina platforms (Figure S1b), and hailing from all over the world (Figure S1c). We observed that not all samples contained sequence read fragments aligned to ARGs, so we only included those that did in the correlation analyses (n = 119,206; Table 1). Looking at the various sampling sources, read fragments aligned to ARGs conferring resistance to tetracycline and beta-lactams were very common in host-associated sources, whereas phenicol resistance was more frequently observed in environmental sources (freshwater, marine, and soil) (Table 2). *catA1* was the most dominant gene in the environmental samples, especially in marine samples, whereas various *tet* genes had high relative abundances in livestock or human samples: *tet(W)* in chicken and pig samples; *tet(Q)* in cow, human, and pig metagenomes. *blaTEM-52B* accounted for more than 30% of the read fragments assigned to ARGs in air metagenomes (Figure S2).

Balancing the sparsity and network complexity

We selected ARGs to infer abundance-based correlations by considering how sparse the input count matrix was and how that sparsity affected the results of SparCC. Out of the 716 ARGs that could be included, we observed that without filtering the raw counts, SparCC found that most of the ARGs correlated with each other, even if the count of read fragments was low. Based on this observation, we decided to require that for an ARG to be included, it had to have a minimum count across a specific number of samples for a group. Applying these two filters, the number of correlations inferred for each sampling group decreased as compared to the no-filter results (Table S1). We decided to pick filter settings to balance the amount of sparsity allowed in the sample groups and the number of correlation coefficients, which ended up being a minimum fragment count of 50 across at least 10 samples.

Analysis of large-scale metagenomic correlation networks

Based on our filter settings, we constructed a global network using the correlation coefficients for the entire collection of metagenomes, with each node representing an ARG and each edge representing a pairwise ARG connection (correlation ≥ 0.6, p-value < 0.01, Figure S3). The global network, nicknamed 'all', contained 225 ARGs connected through 2,344 correlation edges (Figure 1a). As this all-network was hard to interpret due to the many highly interconnected ARGs, we also inferred pairwise ARG correlations in specific sampling groups (Table 1). The genes that were part of these networks were found to correlate with varying degrees of strength (Figure 1a-b). For example, the human network contained many correlation coefficients, but most were less than 0.8 (Figure 1c). Another example is the marine network, where only a few ARGs were found to correlate, but with values above 0.9 (Figure 1b-c). Despite the networks reflecting the composition of the various environments, we still observed overlaps between which ARGs were found to correlate. One example is that all the correlations inferred from the pig metagenomes also existed in the human metagenomic network (Figure 1d).

There was a limited number of correlations for ARGs encoding resistance to fluoroquinolones, steroid antibiotics (fusidic acid), colistin, and rifampicin. On the other hand, beta-lactam, tetracycline, and aminoglycoside ARGs had many correlations with each other and with other classes (Figure S4, Figure S6). Despite resistance to some antimicrobial classes being the most abundant, the ARGs did not always correlate to many others. For example, tetracycline ARGs were the most abundant in dog metagenomes, but no correlations were inferred for these ARGs. Similarly, in the human samples where aminoglycoside and beta-lactam ARGs were less abundant than tetracycline ARGs, aminoglycoside and beta-lactam resistance genes had a higher number of correlations coefficients were reported (Table 2, Figure S4).

On the level of ARG abundance, we observed that just because an ARG was highly abundant in a sampling group, it did not automatically mean that it correlated to many other ARGs. The highly abundant *catA1* gene in marine (97.4% of all ARG reads), freshwater (55.1%), and soil samples (84.7%) (Figure S2) did only correlate with one or two other genes in the water environmental networks, and none in the soil network. On the other hand, *catA1* did seem

to be correlated with 15 other genes in the pig correlation network despite not being highly

abundant in that group of samples (Figure S5a). *mef(A)_1* accounted for 15.9% of the reads

aligned to ARGs in cow samples and 6.63% in pigs (Figure S2) and was also strongly

correlating with other genes, which mainly conferred resistance to aminoglycosides,

(fluoro)quinolones and tetracyclines (Figure S5b). *tet(L)_4* only accounted for 4.01% of the

read fragments aligned to ARGs in metagenomes of chicken origins (Figure S2) but was

shown to correlate in its abundance with the abundance of 8 other ARGs, e.g., with a

correlation of 0.77 with *lnu(A)_1 (*Figure S5c)*.

The hidden signals between ARGs profile the potential risk of co-selection in different

environmental contexts

Using the correlations between ARGs in the various environments (Figure 1), we calculated

the average correlation between ARGs of different antimicrobial classes (Figure S6). These

average correlations can then serve as profiles to assess the risk of indirectly selecting ARGs

that gives resistance to different ARGs through co- and cross-resistance. These risk profiles

can then be used to judge the strength of interactions upon using one antimicrobial in each

setting. Upon constructing these profiles, we observed that the strength and the number of

correlations highly depend on the antimicrobial classes and the environmental

contextFigure S6. We hypothesize that if an important (IA) or highly important antimicrobial

(HIA) is used, first, the resistance to the antimicrobial class will likely flourish, and, secondly,

through co- and cross-resistance, so will ARGs conferring resistance to other classes,

including those that are critically important antimicrobials (CIA).

Figure 2 shows two risk profiles for ARG correlations for two highly critical important

antimicrobials glycopeptide and macrolide. Correlations between ARGs of glycopeptide

resistance to other resistance classes were much rarer than those connected with macrolide

ARGs (Figure S6) and those correlations that were observed were relatively low

(correlation<0.8, Figure 2). Different vancomycin resistance cassettes were responsible in

different environments, namely *VanHAX*, *VanC2*, and *VanX_bc* in human samples and

VanHDX and *VanC1XY* in pig samples (Figure S7a). *VanHAX* only has one correlation,

whereas the remaining five correlate with many different ARGs.

On the contrary, ARGs conferring resistance to macrolide were much more interactive with

other classes of resistance genes in all networks and had strong correlations with specific

classes (correlation > 0.9, Figure 2, Figure S6). The macrolide ARGs co-abundant with other

ARGs were many but separated into distinct network clusters (Figure S7b). For example,

mef(A) and *msr(D)* were usually found together in different environments, both in small and

large clusters.

While Figure 2 highlights how CIA ARGs interact, it is just as important to investigate what

ARGs of less critically important antimicrobials correlate with. As seen in Figure 3, ARGs for

pleuromutilin resistance (IA) and for tetracycline resistance (HIA) were found to interact

with many other classes, including those that are critically important. For example,

pleuromutilin ARGs are few but well connected (Figure 3, Figure S6), as seen with the

connections with *lsa(E)* and *cfr(C)* (Figure S8a)*.* Tetracycline ARGs correlated with the

abundance of multiple ARGs, such as those conferring resistance to lincosamides,

macrolides, and phenicols (Figure 3). While there were many ARGs for tetracycline

resistance, they often correlated to the same ARGs (Figure S8b).

Discussion

Considering the complexity of microbiomes, studying how microbial composition shapes the

distribution of ARGs is a challenging task but one that could shed light on how ARGs

indirectly select one another. However, with the high-throughput sequencing technologies

and many metagenomic datasets available in public repositories, it is now much more

feasible to extract the patterns of how ARGs co-occur without knowing their microbial

origin. Using our recently published collection of 214K metagenomic datasets[25], we have

inferred correlations of pairwise ARG abundances to profile which types of resistance

influence the shape of resistome and which genes are the key players. To the best of our

knowledge, this is the first study to relate ARG abundances on such a large and broad scale.

Our correlation networks revealed that not all ARGs are connected, as we observed that the

interactions were largely shaped by the composition of the environmental resistome.

Nonetheless, we did not always observe strong correlations if an ARG was highly abundant.

The all-network with correlation for all metagenomic samples was the most complex, which

we speculate is due to both the wide variety of sampling sources and the

overrepresentation of human metagenomes (Table 1). We found the differences in the

animal and environmental networks much more interesting, as they reflect the dynamics of

ARG abundances under exposure to different antimicrobials (Figure 1). On the level of which

antimicrobial class an ARG confers resistance to, we could also observe that some resistance

classes had more and stronger correlations (Figure S6). There are several cases of our

observed co-abundant ARGs that have been reported in the literature, and some of the

strongest correlation pairs have been detected due to the ARGs sitting together in the same

genome. For example, the gene cassette *vanHAX* has been found together with *msr(C)* and

aac(6') in genomes of human isolates[52] and *mef(A)* linked with *tet(O)*[53] and *mdf(A)* with

blaTEM, aph(6), sul2, and, tet(A)[54] (Figure S7).

As highlighted in Figure 2 and Figure 3, we argue that the correlations can serve as a way to

profile the risk of co-selection of ARGs occurring in a setting if exposed to an antimicrobial.

Some of these relationships between ARGs of different antimicrobial classes have been

observed in other studies.

Antimicrobials have been classified differently to reflect their importance to human

medicine, of which glycopeptides and macrolides are CIA with the highest priorities.

Glycopeptide and macrolide resistance has previously been linked genetically[6,8] in pigs,

where we also can report the presence of correlations between ARGs of glycopeptide and

macrolide resistances not only in pigs but also in human and mouse environments (Figure

2). Pleuromutilin and tetracycline antimicrobials have less critical importance of

antimicrobials [44], but many correlations for ARGs of pleuromutilin and tetracycline exist in

various networks (Figure 3) suggest that there are still risks associated with the use of these

two. Tetracycline resistance has been found to occur together with resistance to

macrolides[45–47], aminoglycosides[45,48], folate pathway antagonists[49,50], lincosamide[10], and

beta-lactams[48], to name a few studies. This high connectivity of tetracycline ARGs seems in

line with our results, as this specific group of ARGs was connected to almost all classes of

antimicrobials in most of our networks (Figure 3, Figure S8b). The variety of connections to

antimicrobials of less importance to human health should be more in focus, as our results

show that there are risks of critical antimicrobial resistances emerging from the enrichment
of less essential resistance genes.

Following this line of thought of focusing on ARGs that gives resistance to less critically
important antimicrobials, a recent study by Tarek and Garner (2022)[51] proposed to create a
monitoring framework based on isolated components, or clusters, in correlation networks.
They constructed a correlation network encompassing ARG abundances in samples from
wastewater treatment plants and argued that representative ARG members from each
cluster in their network should be monitoring targets. Our networks suggest that limiting to
only a few ARGs would fail to capture the complete picture in some environments, such as
human microbiomes (Figure 1). However, as our risk profiles show, limiting the set of
monitoring targets by only focusing on what happens to ARG abundances during exposure
to one type of antimicrobials would be a way to implement a monitoring system (Figures
Figure 2Figure 3, Figure S7-Figure S8). If a monitoring system is implemented, it would need
to be updated regularly to show the changes in antimicrobial usage and ARGs co-
occurrences since the correlations we have inferred in this study only reflects the current
and past usage of antimicrobials.

In order to use correlation networks for surveillance of AMR, more work is needed to
confirm that the observed interactions do indeed exist in nature[55]. We have defined
interactions as being indirect since we need to determine the degree that co-abundance is
explained by physical linkage on a genome or plasmid or sharing of a cell. To investigate the
direction of correlation, for example, whether *vanHAX* influences *msr(C)* or the other way
around (Figure S7a), the SPIEC-EASI[56] method could be used to infer such directional
dependences. A directional correlation could be included in a risk profile.

We have only included ARGs of different resistance classes that are co-abundant in our
analysis, but the abundance of bacterial reads could also have been analyzed. By including
the bacterial counts, it would likely be possible to understand how the dynamics of
microbiomes and resistomes are linked. Similarly, redoing the alignment procedure to
include the count of reads aligned to other drivers of resistance, e.g., mobile genetic
elements, could also lead to the discovery of new patterns. There has been much work on

linking the prevalence of ARGs with antimicrobial usage (AMU) in various settings[6,8,10], and

incorporating AMU could confirm our speculation that many of the co-abundances are

driven by selective pressures of various antimicrobials. Pooling the various data types

together in this kind of analysis would be a complicated task that nevertheless would shed

light on how the abundance and selective pressures contribute to the shape of resistomes.

By utilizing the wealth of information on ARG abundances available in a collection of 214K

metagenomic datasets, we have studied the co-abundance of ARGs to discover how these

interactions shape the prevalence of resistances in different environments. The inferred

correlation networks provide insights into how two resistance types indirectly and species-

independent select for each other in different habitats. Our results further highlight that

there are instances of genes of one type of resistance often co-occurring with many other

types of resistance and that the environmental context plays an important role, revealing

them as important targets in surveillance programs to limit their impact on global health.

References

1.   O'Neill, J. & The Review on Antimicrobial Resistance (Chaired by Jim O'Neill). Tackling

Drug-Resistant Infections Globally: Final Report and Recommendations. *Rev.*

Antimicrob. Resist. 1–80 (2016).

2.   Murray, C. J. *et al.* Global burden of bacterial antimicrobial resistance in 2019: a

systematic analysis. *Lancet* **399**, 629–655 (2022).

3.   Holmes, A. H. *et al.* Understanding the mechanisms and drivers of antimicrobial

resistance. *Lancet* **387**, 176–187 (2016).

4.   Collignon, P., Powers, J. H., Chiller, T. M., Aidara-Kane, A. & Aarestrup, F. M. World

health organization ranking of antimicrobials according to their importance in human

medicine: a critical step for developing risk management strategies for the use of

antimicrobials in food production animals. *Clin. Infect. Dis.* **49**, 132–141 (2009).

5.   Jensen, H. H. & Hayes, D. J. Impact of Denmark's ban on antimicrobials for growth

promotion. *Curr. Opin. Microbiol.* **19**, 30–36 (2014).

6.   Aarestrup, F. M. *et al.* Effect of abolishment of the use of antimicrobial agents for

growth promotion on occurrence of antimicrobial resistance in fecal enterococci from

food animals in Denmark. *Antimicrob. Agents Chemother.* **45**, 2054–2059 (2001).

7.   Sun, J. *et al.* Development of aminoglycoside and β-lactamase resistance among

intestinal microbiota of swine treated with lincomycin, chlortetracycline, and

amoxicillin. *Front. Microbiol.* **5**, 580 (2014).

8.   Aarestrup, F. M. Characterization of glycopeptide-resistant Enterococcus faecium

(GRE) from broilers and pigs in Denmark: Genetic evidence that persistence of GRE in

pig herds is associated with coselection by resistance to macrolides. *J. Clin. Microbiol.*

38, 2774–2777 (2000).

9.   Looft, T. *et al.* In-feed antibiotic effects on the swine intestinal microbiome. *Proc.*

Natl. Acad. Sci. U. S. A. **109**, 1691–1696 (2012).

10.   Andersen, V. D. *et al.* Predicting effects of changed antimicrobial usage on the

abundance of antimicrobial resistance genes in finisher' gut microbiomes. *Prev. Vet.*

Med. **174**, 104853 (2020).

11.   Peterson, L. R. Squeezing the antibiotic balloon: The impact of antimicrobial classes

on emerging resistance. *Clin. Microbiol. Infect. Suppl.* **11**, 4–16 (2005).

12.   Paterson, D. L. 'Collateral damage' from cephalosporin or quinolone antibiotic

therapy. *Clin. Infect. Dis.* **38**, 341–345 (2004).

13.   Barberán, A., Bates, S. T., Casamayor, E. O. & Fierer, N. Using network analysis to

explore co-occurrence patterns in soil microbial communities. *ISME J.* **6**, 343–351

(2012).

14.   Yuan, M. M. *et al.* Climate warming enhances microbial network complexity and

stability. *Nat. Clim. Chang.* **11**, 343–348 (2021).

15.   Zhang, Z. *et al.* The large-scale spatial patterns of ecological networks between

phytoplankton and zooplankton in coastal marine ecosystems. *Sci. Total Environ.* **827**,

154285 (2022).

16.   Li, L. G., Xia, Y. & Zhang, T. Co-occurrence of antibiotic and metal resistance genes

revealed in complete genome collection. *ISME J.* **11**, 651–662 (2017).

17.   Ju, F. *et al.* Antibiotic resistance genes and human bacterial pathogens: Co-

occurrence, removal, and enrichment in municipal sewage sludge digesters. *Water*

Res. **91**, 1–10 (2016).

18.   Pan, X., Lin, L., Zhang, W., Dong, L. & Yang, Y. Metagenome sequencing to unveil the

resistome in a deep subtropical lake on the Yunnan-Guizhou Plateau, China. *Environ.*

Pollut. **263**, 114470 (2020).

19.   Wu, J. *et al.* Prevalence and distribution of antibiotic resistance in marine fish farming

areas in Hainan, China. *Sci. Total Environ.* **653**, 605–611 (2019).

20.   Liu, J. *et al.* The fecal resistome of dairy cattle is associated with diet during nursing.

Nat. Commun. **10**, (2019).

21.   Munk, P. *et al.* Abundance and diversity of the faecal resistome in slaughter pigs and

broilers in nine European countries. *Nat. Microbiol.* **3**, 898–908 (2018).

22.   Hendriksen, R. S. *et al.* Global monitoring of antimicrobial resistance based on

metagenomics analyses of urban sewage. *Nat. Commun.* **10**, (2019).

23.   Martiny, H.-M. *et al.* Global distribution of mcr gene variants in 214,095 metagenomic

samples. *mSystems* (2022). doi:10.1128/msystems.00105-22

24.   Karkman, A., Do, T. T., Walsh, F. & Virta, M. P. J. Antibiotic-Resistance Genes in Waste

Water. *Trends Microbiol.* **26**, 220–228 (2018).

25.   Martiny, H.-M., Id, P. M., Id, C. B., Aarestrup, F. M. & Petersenid, T. N. A curated data

resource of 214K metagenomes for characterization of the global antimicrobial

resistome. *PLOS Biol.* **20**, e3001792 (2022).

26.   Aitchison, J. The Statistical Analysis of Compositional Data. *J. R. Stat. Soc. Ser. B* **44**,

139–160 (1982).

27.   Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V. & Egozcue, J. J. Microbiome

datasets are compositional: And this is not optional. *Front. Microbiol.* **8**, 1–6 (2017).

28.   Friedman, J. & Alm, E. J. Inferring Correlation Networks from Genomic Survey Data.

PLoS Comput. Biol. **8**, 1–11 (2012).

29.   Leinonen, R. *et al.* The European nucleotide archive. *Nucleic Acids Res.* **39**, 44–47

(2011).

30.   Bushnell, B. BBMap. (2014).

31.   Clausen, P. T. L. C., Aarestrup, F. M. & Lund, O. Rapid and precise alignment of raw

reads against redundant databases with KMA. *BMC Bioinformatics* **19**, 1–8 (2018).

32.   Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J.*

Antimicrob. Chemother. **67**, 2640–2644 (2012).

33.   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: Improved data

processing and web-based tools. *Nucleic Acids Res.* **41**, 590–596 (2013).

34.   Edgar, R. C. & Bateman, A. Search and clustering orders of magnitude faster than

BLAST. *Bioinformatics* **26**, 2460–2461 (2010).

35.   Team, R. C. R: A Language and Environment for Statistical Computing. (2021).

36.   Csardi, G. & Nepusz, T. The igraph software package for complex network research.
424   *InterJournal* **Complex Sy**, 1695 (2006).

425   37.   Epskamp, S., Cramer, A. O. J., Waldorp, L. J., Schmittmann, V. D. & Borsboom, D.
426   qgraph: Network Visualizations of Relationships in Psychometric Data. *J. Stat. Softw.*
427   **48**, 1–18 (2012).

428   38.   Pedersen, T. L. ggraph: An Implementation of Grammar of Graphics for Graphs and
429   Networks. (2021).

430   39.   Fructherman, T. M. . & Reingold, E. M. Graph Drawing by Force-directed Placement.
431   *Softw. - Pract. Exp.* **21**, 1129–1164 (1991).

432   40.   Barrat, A., Barthélemy, M., Pastor-Satorras, R. & Vespignani, A. The architecture of
433   complex weighted networks. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 3747 (2004).

434   41.   Parente, E., Zotta, T., Faust, K., De Filippis, F. & Ercolini, D. Structure of association
435   networks in food bacterial communities. *Food Microbiol.* **73**, 49–60 (2018).

436   42.   Wasserman, S. & Faust, K. *Social network analysis: Methods and applications.*
437   (Cambridge University Press, 1994).

438   43.   Silver, N. C. & Dunlap, W. P. Averaging Correlation Coefficients: Should Fisher's z
439   Transformation Be Used? *J. Appl. Psychol.* **72**, 146–148 (1987).

440   44.   Group, W. H. O. A., Surveillance, I. & Resistance, A. *WHO | WHO list of Critically*
441   *Important Antimicrobials (CIA)*.

442   45.   Arredondo, A., Blanc, V., Mor, C., Nart, J. & León, R. Tetracycline and multidrug
443   resistance in the oral microbiota: differences between healthy subjects and patients
444   with periodontitis in Spain. *J. Oral Microbiol.* **13**, (2021).

445   46.   Nielsen, H. U. K. *et al.* Tetracycline and macrolide co-resistance in Streptococcus
446   pyogenes: Co-selection as a reason for increase in macrolide-resistant S. pyogenes?
447   *Microb. Drug Resist.* **10**, 231–238 (2004).

448   47.   Ardic, N., Ozyurt, M., Sareyyupoglu, B. & Haznedaroglu, T. Investigation of
449   erythromycin and tetracycline resistance genes in methicillin-resistant staphylococci.
450   *Int. J. Antimicrob. Agents* **26**, 213–218 (2005).

451   48.   Stepanauskas, R. *et al.* Coselection for microbial resistance to metals and antibiotics
452   in freshwater microcosms. *Environ. Microbiol.* **8**, 1510–1514 (2006).

453   49.   Gao, P., Munir, M. & Xagoraraki, I. Correlation of tetracycline and sulfonamide

454   antibiotics with corresponding resistance genes and resistant bacteria in a

455   conventional municipal wastewater treatment plant. *Sci. Total Environ.* **421–422**,

456   173–183 (2012).

457   50.   Zhang, M. Q., Yuan, L., Li, Z. H., Zhang, H. C. & Sheng, G. P. Tetracycline exposure

458   shifted microbial communities and enriched antibiotic resistance genes in the aerobic

459   granular sludge. *Environ. Int.* **130**, 104902 (2019).

460   51.   Tarek, M. H. & Garner, E. A proposed framework for the identification of indicator

461   genes for monitoring antibiotic resistance in wastewater: Insights from metagenomic

462   sequencing. *Sci. Total Environ.* 158698 (2022). doi:10.1016/J.SCITOTENV.2022.158698

463   52.   Freitas, A. R. *et al.* High-Resolution Genotyping Unveils Identical Ampicillin-Resistant

464   Enterococcus faecium Strains in Different Sources and Countries: A One Health

465   Approach. *Microorganisms* **10**, (2022).

466   53.   Giovanetti, E., Brenciani, A., Lupidi, R., Roberts, M. C. & Varaldo, P. E. Presence of the

467   tet(O) gene in erythromycin- and tetracycline-resistant strains of Streptococcus

468   pyogenes and linkage with either the mef(A) or the erm(A) gene. *Antimicrob. Agents*

469   *Chemother.* **47**, 2844–2849 (2003).

470   54.   Trongjit, S. & Chuanchuen, R. Whole genome sequencing and characteristics of

471   Escherichia coli with coexistence of ESBL and mcr genes from pigs. *PLoS One* **16**, 1–17

472   (2021).

473   55.   Blanchet, F. G., Cazelles, K. & Gravel, D. Co-occurrence is not evidence of ecological

474   interactions. *Ecol. Lett.* **23**, 1050–1063 (2020).

475   56.   Kurtz, Z. D. *et al.* Sparse and Compositionally Robust Inference of Microbial Ecological

476   Networks. *PLOS Comput. Biol.* **11**, e1004226 (2015).

Resistance class	All	Air	Dog	Chicken	Cow	Freshwater	Human	Marine	Mouse	Pig	Soil
Aminoglycoside (144)	9.12	17.64	8.42	11.51	5.24	16.70	8.51	0.08	10.12	14.08	1.58
Aminoglycoside/Fluoroquinolone /Macrolide/Phenicol/Rifampicin/ Tetracycline (1)	1.08	0.63	0.52	0.12	0.07	0.34	1.08	0.20	1.20	0.17	0.02
Aminoglycoside/Fluoroquinolone /Quinolone (1)	0.19	<0.01	<0.01	0.09	-	0.46	0.15	<0.01	0.01	<0.01	<0.01
Beta_lactam (227)	20.55	41.60	20.07	0.79	1.64	8.44	18.50	1.85	3.02	5.59	1.89
Folate_pathway_antagonist (37)	3.20	2.92	0.97	2.19	0.41	5.45	3.13	0.36	4.33	1.03	0.82
Fosfomycin (28)	1.07	4.36	0.01	0.20	<0.01	1.29	1.49	<0.01	0.22	0.03	0.19
Glycopeptide (27)	0.29	<0.01	-	0.01	<0.01	0.01	0.58	<0.01	0.72	0.02	0.01
Lincosamide (9)	3.33	1.88	29.00	3.42	8.29	0.36	2.02	<0.01	4.01	11.61	0.71
Lincosamide/Macrolide (1)	<0.01	<0.01	-	<0.01	<0.01	<0.01	<0.01	<0.01	-	-	-
Lincosamide/Macrolide/ Streptogramin (44)	12.33	6.12	1.35	14.51	2.26	0.32	17.71	0.03	8.19	9.52	5.95
Lincosamide/Macrolide/ Streptogramin/Tetracycline (1)	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.07	<0.01	<0.01
Lincosamide/Oxazolidinone/ Phenicol/Pleuromutilin/ Streptogramin (3)	0.49	0.41	0.01	1.36	0.50	<0.01	0.45	<0.01	1.64	1.25	0.04
Lincosamide/Pleuromutilin/ Streptogramin (7)	0.27	0.48	0.18	0.09	0.02	0.02	0.34	<0.01	<0.01	0.21	0.05
Lincosamide/Streptogramin (2)	0.28	0.12	<0.01	0.02	<0.01	0.01	0.38	<0.01	0.06	<0.01	<0.01
Macrolide (24)	3.04	1.86	5.78	0.36	16.71	1.90	2.53	<0.01	4.22	6.83	0.25

Macrolide/Streptogramin (4)	1.10	1.69	1.01	0.12	0.39	1.18	1.43	<0.01	1.83	0.48	0.11
Macrolide/Tetracycline (1)	<0.01	<0.01	-	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01
Nitroimidazole (9)	0.32	<0.01	0.04	0.01	0.05	<0.01	0.56	<0.01	0.02	0.14	<0.01
Oxazolidinone/Phenicol (1)	0.02	0.03	-	0.10	<0.01	<0.01	<0.01	<0.01	-	0.09	<0.01
Oxazolidinone/Phenicol/ Tetracycline (1)	0.07	0.02	<0.01	1.54	<0.01	<0.01	<0.01	<0.01	0.03	0.04	<0.01
Phenicol (33)	9.21	6.19	0.36	6.10	4.33	55.89	3.33	97.37	0.14	0.81	85.04
Pleuromutilin (1)	0.06	0.02	-	<0.01	<0.01	0.39	0.06	<0.01	0.01	0.07	<0.01
Polymyxin (10)	1.19	0.30	<0.01	0.14	<0.01	1.85	1.26	<0.01	0.18	0.14	0.12
Quinolone (17)	0.11	<0.01	-	0.01	-	0.18	0.06	<0.01	-	<0.01	<0.01
Rifampicin (6)	0.06	0.09	-	<0.01	<0.01	<0.01	0.03	<0.01	<0.01	<0.01	<0.01
Steroid_antibacterial (2)	0.15	<0.01	-	3.30	<0.01	<0.01	<0.01	<0.01	<0.01	0.11	0.02
Streptogramin (11)	32.48	13.61	32.27	53.99	60.05	5.20	36.39	0.06	59.97	47.76	3.18
Tetracycline (64)	<0.01	-	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01

478   *Table 2: Relative abundance of read fragments aligned to each resistance class (row) for each sampling source (column). The relative abundance is the number of read fragments aligned to the group out of all*

479   *fragments aligned to ARGs. For each resistance class, the parenthesis shows the number of ARGs belonging to that category.*

Figure 1: *The resistome networks consisting of correlations (edges) between pairs of ARGs (nodes).* **a**. *Each correlation network is visualized, where each ARG node is colored by resistance class, and edges are colored by the correlation coefficient value.* **b**. *Metrics of the correlation networks reveal how interconnected and complex the networks are. No. stands for number of.* **c**. *Distribution of included correlation coefficients in each network. Figure S3 shows the distribution of all inferred correlations and their p-values.* **d**. *A heatmap showing how similar the content of the two networks is. In the upper half of the heatmap, the number of overlapping ARG nodes is shown. An overlap means that a specific ARG has a correlation in both networks, ignoring what it is co-abundant with. In the lower half of the heatmap, the number of overlapping correlation edges is shown. An overlapping edge is defined as whether the correlation coefficient is reported in both networks, regardless of the value of the coefficient.*

Figure 2: **Correlation profiles for ARGs conferring resistance to the critically important antimicrobial classes glycopeptides (left) and macrolides (right).** *Each column shows the average correlation from, e.g., macrolide ARGs to ARGs for other antimicrobial classes. The circle is colored by the average correlation, where a white circle indicates no statistically significant correlations of ARGs observed between the two antimicrobial classes.*

Figure 3: **Correlation profiles for ARGs conferring resistance to important pleuromutilins (left) and highly important tetracyclines (right).** *Each column shows the average correlation from, e.g., tetracycline ARGs to ARGs for other antimicrobial classes. The circle is colored by the average correlation, where a white circle indicates no statistically significant correlations of ARGs observed between the two antimicrobial classes.*

Supplementary materials

*Figure S1: **Metagenomic origins colored by the sampling group**. a. Overview of sampling year for metagenomic samples colored by their sampling origin. 100 samples were taken before 2000, and 84,238 did not have a valid sampling date. b. Overview of the amount of sequencing reads available for each sampling origin. c. Sampling locations of metagenomic samples used in the correlation analysis were split by their sampling source. Number of samples with no coordinates available; All: 83,361; Air: 16; Dog: 3,159; Chicken: 570; Cow: 262; Freshwater: 61; Human: 40,003; Marine: 363; Mouse: 2,842; Pig: 320; Soil: 477. The 'Other' label refers to those that are not in one of the source-specific networks but are included in the 'All' network.*

505 *Figure S2:* **Top 10 most abundant ARGs per sampling group.** *A barplot showing the ARGs having the highest 10 relative*
506 *abundances per sampling source colored by resistance class.*

| | No filters | | | minF | minS | | minF | minS | | minF | minS | | minF | minS | |
| | | | | 25 | 10 | | 50 | 10 | | 50 | 20 | | 50 | 50 | |
Source	ARG cols	#nodes	#edges	ARG cols	#nodes	#edges	ARG cols	#nodes	#edges	ARG cols	#nodes	#edges	ARG cols	#nodes	#edges
All	716	443	34,815	459	275	4,786	425	257	2,970	313	165	320	313	165	320
Air	436	565	66,612	34	14	10	19	9	6	0	-	-	0	-	-
Chicken	391	634	92,563	145	123	289	123	191	226	69	56	87	69	56	87
Cow	445	614	92,881	94	75	242	85	68	198	37	25	57	37	25	57
Dog	158	72	85	45	25	35	45	25	35	45	25	35	45	25	35
Freshwater	579	589	43,054	56	40	74	36	24	33	1	-	-	1	-	-
Human	692	474	53,408	307	179	1,445	278	158	902	201	91	120	201	91	120
Marine	668	13	8	24	7	5	10	7	5	4	-	-	4	-	-
Mouse	326	549	80,254	55	45	122	41	35	107	25	21	70	25	21	70
Pig	408	634	87,754	136	114	864	118	99	606	61	52	175	61	52	175
Soil	614	510	22,415	72	53	85	47	28	37	3	-	-	3	-	-

507 *Table S1: **Filtering the counts impacts the number of input ARG columns, as well as the resulting number of correlations.** minF is the minimum number of fragments available for an ARG in*

508 *one sample, and minS is the minimum number of samples to fulfill the minF requirement for the column to be included in the inference procedure*

510 *Figure S3: **Distribution of correlations and p-values produced by SparCC for each data grouping**. Each point is a correlation*

511 *between two ARGs and is colored by whether the point was selected if the p-value < 0.01 and correlation ≥ 0.6.*

*Figure S4: **The relative abundance, number of ARGs, and the number of correlations for each resistance class in each sampling group.** The relative abundance in green shows the percentage of read fragments for each class. In grey is the number of ARGs for each resistance class. Orange coloring indicates the number of correlation coefficients inferred for ARGs in a resistance class.*

Figure S5: **Networks illustrating how one ARG interacts differently depending on the sampling environments**. *The three genes catA1_1, mef(A)_1, and tet(L)_4 are selected to illustrate that the genes correlate more or less with other genes in their abundance in some habitats.*

Figure S6: The average correlation between ARGs of different resistance classes in the upper triangle of the heatmaps with the lower half shows
the correlation coefficients between the two classes. Note that a correlation coefficient between two ARGs might be present in more than one
tile, as some ARGs confer resistance to multiple classes of antimicrobials.

523
524
525

529   *Figure S7: **Highlights of interactions between ARGs of a. glycopeptide resistance and b. macrolide resistance**. The ARG node size shows*

530   *whether the corresponding ARG gives resistance to the antimicrobial class in focus. Only correlation edges and ARGs nodes are colored if they*

531   *correlate with the highlighted ARGs; otherwise, they are colored grey. The coloring schemes for nodes and edges are given in Figure 1a.*

Figure S8: **Highlights of interactions between ARGs of a. pleuromutilin resistance and b. tetracycline resistance.** *The ARG node size shows whether the corresponding ARG gives resistance to the antimicrobial class in focus. Only correlation edges and ARGs nodes are colored if they correlate with the highlighted ARGs; otherwise, they are colored grey. The coloring schemes for nodes and edges are given in* Figure 1a.