Subtle introgression footprints at the end of the speciation continuum in a clade of *Heliconius* butterflies

Quentin Rougemont* (1) Bárbara Huber* (2,3,8), Simon Martin (4), Annabel Whibley (2), Catalina Estrada (3,5), Darha Solano (3), Robert Orpet (3,6), W. Owen McMillan (3), Brigitte Frérot (7), Mathieu Joron (1,2)

Centre d'Ecologie Fonctionnelle et Evolutive, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

Institut de Systématique, Evolution et Biodiversité, Muséum National d'Histoire Naturelle, Paris, France
Smithsonian Tropical Research Institute, Balboa, Panama
Department of Zoology, University of Cambridge, United Kingdom
Department of Life Sciences, Imperial College London, Silwood Park, London, UK
Department of Entomology, Washington State University, Wenatchee, Washington, USA
INRAE, CNRS, IRD, UPEC, Sorbonne Université, Institute of Ecology and Environmental Sciences of Paris, Université de Paris, 78000 Versailles, France
Instituto de Ciencias Ambientales y Ecológicas (ICAE), Universidad de Los Andes, Mérida, Venezuela

* Co-first authors
Corresponding authors: quentinrougemont@orange.fr, babahuber@gmail.com, mathieu.joron@cefe.cnrs.fr

Abstract

Quantifying gene flow between lineages at different stages of the speciation continuum is central to understanding speciation. *Heliconius* butterflies have undergone an adaptive radiation in wing colour patterns driven partly by natural selection for local mimicry. They are also known for assortative mating based on wing pattern. Therefore, wing patterns may facilitate speciation. Here, we take a multifaceted approach to explore speciation and species boundaries between closely-related species *H. hecale* and *H. ismenius.* We focus our research in geographic regions where the two are mimetic and contrast this with geographic regions where they do not mimic each other. To examine population history and patterns of gene flow, we developed a four population model accounting for linked selection. This model suggests that the two species have remained isolated for the majority of their history, yet with a small amount of gene exchange. Accordingly, signatures of genomic introgression were small except at a major wing pattern allele. Additionally, behavioural assays suggest that other cues determine strong sexual isolation. Tests for chemical differentiation between species identified major differences in putative pheromones which likely mediate mate choice and the maintenance of species differences. Our results show that strong divergence and adaptive introgression at wing pattern loci may not reveal the whole *Heliconius* speciation history. In a clade where assortative mating is readily triggered by wing pattern differences, we show that this alone may not lead to speciation and that the accumulation of other barriers to gene flow are important in the completion of speciation.

Introduction

Understanding the modalities of species formation and the maintenance of species barriers is central to evolutionary biology. Toward this goal, an understanding of the demographic history and quantifying gene flow and isolation among populations, is fundamental. Recent empirical population genomics studies have documented the near ubiquity of the heterogeneous landscape of differentiation across a continuum of increasing divergence (Ravinet et al. 2017). These landscapes can arise under divergence hitchhiking, where continuous gene flow will impede genetic differentiation outside of areas involved in local adaptation (Via and West 2008; Feder et al. 2012; Flaxman et al. 2013). Unfortunately, similar patterns can arise under models of secondary contact, where gene flow erodes past genetic differentiation in regions without barrier loci (Barton and Bengtsson 1986). Another complicating factor is the effect of selection at linked sites, where hitchhiking of neutral alleles linked to a selective sweep can also contribute to a heterogeneous landscape (Noor and Bennett 2009; Cruickshank and Hahn 2014). Local variation in recombination and in the density of selected sites within the region determine the intensity of linked selection (Kaplan et al. 1989; Nordborg et al. 1996; Payseur and Nachman 2002; Burri 2017). Linked selection reduces polymorphism at sites harbouring positive as well as deleterious variants and at surrounding sites, especially in areas of low recombination (Hill and Robertson 1966; Charlesworth et al. 1993). Linked selection can be modelled as a local reduction in effective population size (Ne), although this is a simplified approach. A modelling approach that jointly allows for local genomic variation in effective population size and migration rate can improve our understanding of the demographic processes at play during population divergence.

Heliconius butterflies constitute a clade composed of numerous hybridising species, providing an excellent system to understand the contribution of gene flow, linked selection and demography to the genomic landscape of divergence (Martin et al. 2019; Van Belleghem et al. 2020). Many colour pattern morphs within a species are common across the radiation with, for instance, up to 7 sympatric forms within a single population of *H. numata* and over 25 distinct wing pattern populations in the *H. melpomene* or *H. erato* lineages (Mallet and Gilbert 1995). *Heliconius* butterflies are unpalatable to predators and advertise their toxicity through their colour pattern and are classic examples of Müllerian mimicry, where distasteful species converge a common warning signal (Sheppard et al. 1985). The loci responsible for colour pattern variation have been identified using a combination of genetic mapping, GWAS, gene expression analyses and more recently functional knockouts. These studies have revealed that a few highly conserved genes interacting to modulate much of the wing pattern variation both within and between *Heliconius* species. This includes *i)* the gene *optix,* which controls the distribution of red-orange pattern elements (Reed et al. 2011; Zhang et al. 2017; Huber et al. 2015; Lewis et al. 2019), *ii)* the gene *cortex* which controls the presence and position of white and yellow pattern elements (Livraghi et al. 2021), *iii)* the gene *wntA* which controls melanic patterning across the wing (Martin et al. 2012; Moest et al. 2019; Fenner et al. 2020; Van

Belleghem et al. 2020) and *iv)* and *aristaless 1*, which is a white/yellow switch (reviewed in McMillan et al. 2019).

Here, we focus on a pair of closely-related species *H. hecale* and *H. ismenius* which belong to the so-called silvaniform lineage and have a complex history of wing pattern mimicry and diversification (Huber et al. 2015). In Eastern Panama, *H. hecale melicerta* and *H. ismenius boulleti* are perfect co-mimics of each other and both display a pattern made of orange (proximal) and yellow elements (distal) bordered by a thick black margin and a black wing tip. In Western Panama, *H. hecale zuleika* and *H. ismenius clarescens* resemble each other but participate in different local mimicry rings. *H. h. zuleika* shows a pattern composed of a largely black forewing with yellow dots and an orange hindwing and mimics *H. hecalesia formosus, Eueides procula* and various highly distasteful butterflies in the tribe Ithomiini, notably *Tithorea tarricina*. In contrast, *H. i. clarescens* has largely orange wings striped with black and yellow and does not seem to belong to any obvious mimicry ring (Rosser et al. 2015). Previous studies have demonstrated that *Heliconius* co-mimics share the same habitats and that details in the colour patterns matters in butterflies species recognition (Mallet and Gilbert 1995). However, chemical signalling has been suggested to also be involved in maintaining species integrity between pairs of sympatric co-mimetic species where colour-based recognition may be compromised (Mérot et al. 2015, González-Rojas et al. 2020) including in the species we studied here (Mann et al. 2017,). The contrasting situation observed here, *i.e.* perfect mimicry vs. non-mimicry (hereafter considered as co-mimics vs non co-mimics) offers a unique opportunity to study the role of gene flow in the evolution of mimicry. We tested the hypothesis of adaptive introgression of wing patterning genes among co-mimics, and the expectation of local gene flow to be stronger between co-mimics than between non co-mimics. We combined whole genome analyses, as well as behavioural and chemistry assays to understand the modalities of gene flow among species. We aimed at i) reconstructing the divergence history of *H. hecale* and *H. ismenius,* taking into account barriers to gene flow and linked selection in a newly developed four-population model that includes powerful statistics for gene flow detection, ii) quantifying the extent of fine scale variation in gene flow along the genome and its role in the maintenance of co-mimicry and convergence in wing pattern among *H. hecale melicerta* and *H. ismenius boulleti,* and iii) evaluating the role of sex, behavior and chemical factors in the maintenance of the strong isolation observed along the genome of this species complex.

Results

Strong divergence between species

A total of 73 individuals were sampled and genotyped using whole genome sequencing **(Fig 1A and Fig 1B, Table S01)**. These include the two perfect co-mimics *H. hecale melicerta* and *H ismenius boulleti* from Eastern Panama (n = 10) and their two sympatric but non-mimetic counterparts *H. hecale zuleika* and *H. ismenius clarescens* from Western Panama (n = 9). In addition, *H. hecale felix* (n = 4)*, H. hecale clearei* (n = 4)*, and H. ismenius telchinia* (n = 2) were included to test for gene flow and assess

global population structure. To further evaluate introgression and population structure, *H. pardalinus* (n = 15) and *H. numata silvana* (n = 17) specimens were included in our analyses as well as *H. melpomene* (n = 12). These species are all closely related outgroups to *H. hecale* and *H. ismenius*.

To estimate the structure of genetic variation among species and populations, we performed a principal component analysis (PCA) on the whole genome data. This revealed an unambiguous separation of the two species, *H. hecale* and *H. ismenius*, along the first axis. The second axis separated individuals of *H. hecale* into three geographic clusters corresponding to 1) *H. hecale felix*, 2) *H. hecale clearei* and 3) a group comprising *H. hecale zuleika* and *H. hecale melicerta* individuals (**Fig 1C**). High levels of net sequence divergence between species (D_A range = 0.16 to 0.22) and differentiation (F_{ST} range = 0.87 to 0.92) were observed at coding sites and mirror these observations. Above a level of net divergence (D_A) greater than 0.1, the probability of gene flow was found to be reduced in a recent study across the speciation continuum (Roux et al. 2016). Accordingly, measures of relative genetic differentiation (F_{ST}, **Fig 2A**) and of absolute genetic divergence (D_{XY}, **Fig 2B**) across the whole genome were high both between co-mimic and non co-mimic populations.

These landscapes were conserved across species comparisons, even when including more distant groups of species from the present study (**Fig S01**). Recombination landscapes were conserved among species (**Fig S02**) which likely explained the correlated levels of diversity and divergence at a broad spatial scale as detailed in **Fig S03-S04**, although a full investigation of the role of recombination in shaping divergence landscape was outside the scope of this paper. Genetic differentiation and divergence between subpopulations of *H. hecale* and *H. ismenius* (**Fig 2A** and **Fig 2B**, green and chocolate lines) were lower than between species comparison, as expected under ongoing gene flow, high effective population size, and/or very recent divergence. Interestingly, a few peaks of high differentiation (especially F_{ST}) and divergence between *H. h. zuleika and H. h. melicerta* were observed in an ocean of low divergence (**Fig 2B**). Accordingly, shared peaks of F_{ST} and D_{XY} (above the 80% quantile) involved a total of 55 genes spread on 11 chromosomes.

To better assess evolutionary divergence, and test whether the phylogeny followed a simple bifurcating tree or a network with gene flow, we reconstructed the phylogeny using Treemix (Pickrell and Pritchard 2012) and included all outgroups (*H. numata* and *H. pardalinus*, **Fig 1D**). Including migration events only modestly improved the model fit (**Fig S05**). Accordingly, these migration events likely occurred in the ancient past, on ancestral branches between *H. ismenius* and *H. hecale*, as well as from *H. pardalinus* to *H. hecale* or from *H. pardalinus* into *H. numata*. This analysis did not show evidence of gene flow between co-mimics *H. h. melicerta* and *H. i. boulleti*. This network revealed higher drift in *H. ismenius* a result that translates into slightly increased Tajima's D values (**Fig 1E**) as compared to *H. hecale* or to other species with a higher effective size such as *H. pardalinus* or *H. numata* (De Cara et al. 2022).

Figure 1. Sampling, population structure and evolutionary relationship among populations. (**A**) Fine scale sampling design and (**B**) broad scale sampling design as well as phenotypic difference between mimics and non co-mimic species. (**C**) Population structure among sampled species and (**D**) evolutionary tree highlighting genetic exchange among populations. (**E**) Tajima's D in 50 kb windows along the genome. The black dot represents the mean value +/- 1 standard deviation.

Figure 2. Conserved and high divergence between species. (**A**) Genetic differentiation (F$_{ST}$) landscape along each chromosome between sympatric co-mimics (*H. h. melicerta* vs *H. i. boulleti*, red line) and between non co-mimics species (*H. h. zuleika* vs *H. i. clarescens*, blue line). Genetic differentiation between subspecies of each species is also shown (*H. h. melicerta* vs *H. h. zuleika*, chocolate; *H. i. boulleti* vs *H. i. clarescens*, green). (**B**) Absolute genetic divergence (D$_{XY}$) along the genome. The same colour coding is used. In general, divergence and differentiation landscapes are so conserved between co-mimics and non co-mimics that they overlap. (**C**) Landscape of genetic diversity (π) observed along the genome for each species. Again, landscapes within species are highly conserved. (**D**) Population scaled recombination landscape along the genome for *H. hecale* and *H. ismenius.* Computation averaged in 250 kb windows.

Nearly complete isolation is maintained through evolutionary time

Strong differences in levels of genetic diversity can complicate estimating gene flow. Therefore, we assessed levels of genetic diversity π and Tajima's D in 50kb windows. As expected from previous research on the silvaniform clade (De Cara et al. 2022), genetic diversity was approximately 2.85 times as high in *H. hecale* (range = 0.0024 to 0.00248) as in *H. ismenius* (range = 0.00080 to 0.00087). Negative Tajimas' D were observed in both *H. hecale* and *H. ismenius* with values approximately 3-4 times lower in *H. ismenius* than *H. hecale,* indicating a stronger recovery or expansion following a bottleneck. These strong differences in patterns of diversity suggest that effective population sizes are markedly different in the two species and that *H. ismenius* may not be at equilibrium. These conditions may further complicate the interpretation of traditional tests to detect gene flow. Thus, we tested whether gene flow occurred during the species demographic history by 1) inferring the demographic history of the species and comparing scenarios with or without gene flow, 2) developing a four-population ABC model that integrates ABBA-BABBA like

statistics (see **Methods**) and 3) integrating the confounding effect of barriers to gene flow as well as linked selection (Ewing and Jensen 2016; Roux et al. 2016; Schrider et al. 2016).

We first used a Random-Forest procedure (ABC-RF hereafter) (Pudlo et al. 2016) and an ABC neural-network (Csillery et al. 2012) to compare statistically a model of divergence followed by gene flow (SC or Secondary Contact) to a model of divergence without gene flow (SI or Strict Isolation, **Fig S06**). Statistical comparison was performed hierarchically to improve interpretability. First, we compared SI models integrating the effect of linked selection (modelled as a local reduction in Ne, SI-*Nhetero*) versus a null model without local reduction in *Ne* along the genome (SI-*Nhomo*). Similarly we compared SC models integrating both the confounding effect of linked selection as well as barriers to gene flow, locally reducing the effective migration rate (*m*), against models integrating only one of the two effects and against a null model ignoring these confounders (resulting in four alternative models, namely SC-*NhomoMhomo* for homogeneous effective population size and homogeneous gene flow, SC-*Mhetero* for heterogeneous effective population size only, SC-*Nhetero* for heterogeneous migration only and SC- *MheteroNhetero* for both effects).

Both the ABC and ABC-RF strongly support a model of linked selection [SI-*Nhetero* (posterior probability ~ 1, **Table S02**)] as well as the different models of secondary contact [SC (-*Nhetero, -Mhetero, -MheteroNhetero,* posterior probability ~ 1, **Table S03**)] as compared to the alternative null model. Further analyses based on the relationship between diversity, divergence, differentiation and recombination metrics further support the impact of linked selection (see **Fig S03-S04**). In addition, ABC-RF confusion matrix indicated that our classifier was able to accurately discriminate among these alternative models (OOB-prior rate = 6% for SI, **Table S02**). As expected, due to the similarity of the heterogeneous versions of the SC models, the OOB-prior rate increased to approximately 33% when considering these models (**Table S03**). Yet, the strong support for these heterogeneous models in our empirical data, echoes the general observation in other species (Charlesworth and Jensen, 2021) and we therefore chose to make our next comparison by considering the model including both linked selection and barriers to gene flow (SC-*MheterNhetero*).

We compared seven distinct Secondary Contact models differing in the directionality of gene flow (**Fig S06**). Namely we tested for unidirectional migration from *H hecale melicerta* to *H. ismenius boulleti* (model SC$_A$), the reverse direction (model SC$_C$), from *H hecale zuleika* to *H. ismenius clarescens* (model SC$_B$), the reverse (model SC$_D$), for bidirectional gene flow between co-mimics (model SC$_{AC}$) and between non co-mimics (model SC$_{BD}$). The seventh and last model includes all possible directions of migration (model SC$_{AC-BD}$). Here, our model selection procedure clearly rejected models with multiple directions of migration p(SCac-bd) = 0.01, p(SC$_{AC}$)= 0.06, p(SC$_{BD}$) = 0.04 in favour of the model SC$_C$ and SC$_B$. The same results were obtained with ABC-RF (**Table S03**). A final comparison between SCc and SC$_B$ yielded stronger support for SC$_C$ both in the ABC and ABC-RF approach (**Table S03**). Our final model choice test involved a comparison of the SI-*Nhetero* versus SC$_C$*MheteteroNHetero*. Here, our two classifiers disagreed: The ABC-RF inferred the SC$_C$*MheterNhetero* as the best model (*p = 0.91*) whereas the ABC classified SI-

Nhetero as the best (*p = 0.85*). This highlights some fundamental limits to ABC inferences (Robert et al. 2011; **Discussion**). Further investigations revealed that the ABC approach was sensitive to the choice of the applied tolerance but that both the ABC and ABC-RF display a high accuracy (Methods, **Table S04, Fig S07**). Therefore, we focus our investigation on the SC model.

We next examined the data to determine if it supported a model where the two species diverged in allopatry (expected under SC) or in the face of ongoing gene flow as expected under models of isolation with migration (IM, **Fig S06**) or divergence with initial migration (AM, **Fig S06**). To address this question, we constructed two IM models, one with gene flow between co-mimics and one with gene flow between non co-mimics (**Fig S06**) and a model of ancient migration (AM) and compared the best IM models against AM. The best supported model was that of gene flow between co-mimics (p = 0.54). Yet, this IM model was clearly rejected in favour of the AM model (p(AM) = 0.83). Therefore, we compared this AM model against SCc and found higher support for AM against SC (p(SC) = 0.87, supporting the hypothesis of allopatric divergence. All these comparisons were highly robust **(Table S04)**.

We next estimated parameters under the two best models, namely SI and SCc. Larger credible intervals were obtained under SC than SI **(Table S05, Fig S07-08).** Under the SI models effective population size estimated with the ABC for both *H. hecale* and *H. ismenius* were large, being ~20M in *H. hecale* and less than half this figure in *H. ismenius* (**Fig 3**), concordant with observed levels of genetic diversity (π) in these two species. Differences in mean effective population sizes within subspecies were minor (**Fig 3**). Estimates of gene flow were largely symmetric between *H. hecale* subpopulations and approximately ten times as strong as in *H. ismenius* (**Fig 3**). Interestingly, estimates of effective population size increased largely under models of SC so that the difference between species was not detected any more (**Table S05**). Split times estimates displayed narrow credible intervals (**Fig S08**), and accordingly, *H. hecale zuleika* and *H. hecale melicerta* would have diverged the most recently (~1.3 million years ago). *H. ismenius boulleti* likely diverged approximately 1.6 My ago from *H. ismenius clarescens.* Yet, considering uncertainty in parameters estimates, it is possible that all subpopulations diverged approximately at the same time (**Table S05**). Finally, *H. hecale* most likely diverged from *H. ismenius* approximately 3.6 My ago. According to our secondary contact model, migration from *H. hecale melicerta* into *H. ismenius boulleti* started approximately 130 KyA [CI = 1.5KyA– 2MyA] at a rate m = 2.15e-05 [95%CI: 5.8e-7 – 7.3e-5]). Other parameters, namely the shape of the beta distribution used to quantify effective population size reduction (Ne) and barriers to gene flow (m) were also difficult to estimate (see also **Fig. S04** for further analyses of linked selection). In all cases, the beta distribution associated with Ne suggested that effective population size would have been strongly reduced in regions affected by linked selection. The ABC-RF procedure generally failed to generate any informative posterior estimation, in spite of a very high accuracy on simulated data (R^2 = 0.96; **Table S06, Fig S07**) suggesting overfitting. We also attempted to estimate parameters through extreme gradient boosting (Xgboost, Chen et al. 2022). This method displayed modest accuracy on trained data (R^2 = 0.635) but also generally failed to estimate parameters (**Table S06**).

Figure 3. Most likely scenario of divergence inferred from ABC and ABC-RF. The most likely model was obtained through both ABC and ABC-RF. Parameters were estimated with ABC. Arrows display migration rate. The dotted green arrow displays possible migration (inferred with ABC-RF only) between the co-mimics. Split time estimated in years assuming four generations per year. All models included the effect of selection at linked sites and of barriers to gene-flow.

Localised, asymmetric and scattered adaptive introgression between co-mimics

We used the ABBA-BABA family of tests, namely D (Green et al. 2010), f_d (Martin et al. 2015), and D_{FOIL} (Pease and Hahn 2015), to test for introgression globally and in windows among co-mimics and with other closely related species. At the genome scale, we used all possible donor taxa, including other subspecies such as *H. h. felix, H. i. telchinia* and more distant species (*H. pardalinus* and *H. numata silvana).* As expected from our ABC modelling approach, we found limited evidence for gene flow among the co-mimics at a genome wide scale based on both D, f_d or f_dM (**Table S07).** Using the more distantly related *H. pardalinus* as a P1 suggested introgression between *H. h. melicerta* and *H. i. boulleti* as well as between *H. h. zuleika* and *H. i. clarescens.* However, in all these cases, admixture proportions (f_d and f_dM) across the whole genome were not significant. For instance, with introgression from a single individual in a set of four individuals in each population, the maximal f_d value is 0.125 (Van Belleghem et al. 2021).

Next, we looked at f_d values along the genome assessing the topology P1 = *H. i. clarescens*, P2 = *H. i. boulleti,* P3 = *H. h. melicerta* (**Fig 4A**) to test for allele sharing between the co-mimics (P2 and P3) and contrasting this to non co-mimic pair (*H. i. boulleti, H. i. clarescens, H. h. zuleika)* (**Fig 4B**). In our co-mimics pair comparisons, there were a total of 26 windows with f_d>0.125 (mean length = 15kb), with a striking ~55 kb peak at the gene *optix* on chromosome 18 whose role on wing patterning is well established.

Moreover, strong peaks on chromosome 14 and chromosome 19 were also found (**Fig 4A)**. For the non co-mimics comparisons, we observed a total of 35 shared windows (mean length = 14kb). We observed a very high peak on chromosome 7 which corresponds to a ~34 kb block containing four genes of unknown function in *H. melpomene.* As in the analysis of co-mimics, a peak is found on chromosome 19 but the signal on chromosome 18 involving *optix* was not observed (**Fig 4B**), suggesting that the wing pattern alleles were not shared between these subspecies. This result was robust and using more divergent species to increase our power to detect gene flow between always showed the existence of a shared introgression signal (**Fig 4A, Fig 4B**) at chromosome 19 and an introgression/shared allele signal at *optix* on chromosome 18 among co-mimics but no signal on chromosome 7 suggesting a recent introgression event between *H. h. zuleika* and *H. i. clarescens* (**Fig S09-S11**).

Figure 4. Localised, asymmetric and stronger introgression between co-mimics (A) than non co-mimics (B). 10kb sliding windows f_d values are shown for each comparison, with each chromosome being colored in dark and light blue. The f_d topology is displayed on the right. Significant D_{FOIL} values are displayed as blue and red stars for each significant introgression direction and window along the genome. The corresponding D_{FOIL} topologies and directions are displayed on the right.

More introgression between co-mimics than non co-mimics

To test our hypothesis of gene flow in relation to mimicry more directly, we used the D_{FOIL} statistics. Unlike the above ABBA-BABA tests, which necessitate performing separate comparisons between different P1, P2, P3 species and then comparing the number of introgressed windows inferred for each tested, Dfoil relies on a four population tree (and an outgroup), making it possible to test the excess of gene flow directly. To do so, we tested the topology P1, P2, P3, P4, O in which P1 = *H. h. melicerta,* P2 = *H. h. zuleika,* P3 = *H. i. boulleti* and P4 = *H. i. clarescens.* Under stronger gene flow between co-mimics we should expect an excess of P1 → P3, and P3 → P1 over P2 → P4 and P4 → P2.

Figure 5. More introgression between co-mimics than non co-mimics revealed by D$_{FOIL}$ analyses. (**A**) Number of significant D$_{FOIL}$ tests between each possible direction. (**B**) Topology tested in the D$_{FOIL}$ analysis.

Since this analysis can only handle a single sequence per population, we performed all possible comparisons between individuals from each species/population and we performed tests in 200 kb windows as it follows a χ^2 distribution over such a window size. We observed a total of 481 significant windows in either the P1 → P3 direction or P3 → P1 direction versus 218 significant windows when considering P2/P4 supporting the idea that gene flow between co-mimics is stronger (**Figure 4, Figure 5, Table S08**). We observed twice as many significant tests in the P12 → P3 direction as in the P12 → P4 direction (P12 being the ancestral *H. ismenius* branch), suggesting that there has been significant gene flow in ancestral populations, whose footprint may have been removed by selection and recombination.

In order to confirm our previous results, we attempted to refine the distribution of introgression tracts using topology weighting by weighting the support for each fifteen different topology among the four species with *twisst* (Martin and Van Belleghem 2017). This analysis largely confirmed that the majority of the genome (84%) support the species tree (topology 3, **Fig S12**) and this increases to 98.8% when considering additional topologies compatible with the species tree (topology 15 & 10). Overall, one window on chromosome 18 around *optix* was indicative of introgression between co-mimics (topology 9, 12 & 13) and received a weight close to 40% (all topologies are displayed in **Fig S12**). Another topology on chromosome 11 received a weight close to 30%. Considering gene flow between *H. i. clarescens* and *H. h. zuleika, two* windows on chromosome 15 and 08, received a weight of ~32 and 34%, respectively. All remaining topologies were equally compatible with gene flow between all species, incomplete lineage sorting or selection against introgression, as expected due to the strength of the species barriers (**Fig S13**).

Courtship behaviour in *H. hecale*

Considering the high levels of genomic divergence and putative reproductive isolation, we investigated mating behaviour in con- and heterospecific encounters. *Heliconius hecale* displays a simple sequence of behavioural units **(Fig 6A; Table S09)**; see also Crane (1957). In captivity, the aerial phase where both sexes are flying is reduced and based mostly on visual signals (Tinbergen et al. 1942; Pliske 1975; Brower 1996). The aerial-ground phase, where the male is flying stationarily while the female is sitting, is prolonged (as often seen in patrolling species; Scott 1972) and may represent the stage where both chemical, visual, and tactile cues are evaluated. The ground-ground phase represents the last part of the sequence, with physical contact between sexes and mating attempts. Five main steps characterise the *H. hecale* courtship **(Fig 6A** and **Table S09)**. At any time, the female may reject the male by adopting a typical rejection posture (Obara 1964; Rutowski and Schaefer 1984). Minor and atypical steps were also recorded and were consistently supported by previous studies (e. g. direct contact between sensorial organs, such as antennae and legs with wings and head; see ethogram in **Table S09)**, but their importance is difficult to evaluate due to their low frequency..

Interspecific recognition relies on short-range cues

To investigate the signals involved in interspecific recognition, we contrasted inter to intraspecific courtship trials. We observed very strong premating isolation between species and no interspecific mating. This pattern contrasted with intraspecific trials where mating occurred within the first 15 minutes in 75% of our trials (**Fig 6B**). Intraspecific trials showed a gradual transition between main steps, suggesting a gradually increasing selectivity across the courtship sequence. 72.4% of *localization* events were followed by *hovering* events, and 57.6% of hovering events elicited m*ating attempts* in males. In total, 22.5% of such attempts led to mating. By contrast, interspecific trials showed high numbers of approaches but very few were followed by active courtship. Heterospecific confusion occurs at long distances but species discrimination is efficient at short range. *Female rejection* is much higher in interspecific experiments, suggesting that both sexes assess the identity and quality of their mates at short range. All interspecific mating *attempting* events (n = 7 by 4 distinct males) were strongly rejected by the tested female (data not shown). Males approached conspecific and heterospecific female wing models equally (*localization*; G=0.002 P=0.96; **Fig 7A, Table S10**). However, males of both species performed the *hovering* behaviour more often on conspecific than on heterospecific models (G=21.323 P<0.001; **Fig 7B, Table S11**) yet *H. h. melicerta* had a higher propensity to court conspecific models than H. *i. boulleti* males (**Fig 7B**). Finally, we explored the role that color played in female *H. h. melicerta* behavior by altering the wing pattens of male *H. h. melicerta.* In these experiments, males with unaltered (*sham*) vs. modified (*treated*) wing patterns (n=10 each) showed equal progression in courtship (**Fig 7C**), and female rejection rates were not significantly different (n_{sham}=59, $n_{treated}$=78, χ^2=2.635 P=0.10; **Fig 7C**).

Figure 6. Interspecific recognition relies on short-range cues. (**A**) Main steps of courtship and mating behaviour in *H. hecale.* Courting steps named after the nomenclature used for *Bicyclus anynana* (Nieberding et al. 2008). *Female rejection* is more frequent at later stages. (**B**) Comparison of inter- and intraspecific male-female encounters. The number of events in each main courtship step is shown for interspecific trials (dashed line; *H. h. melicerta* and *H. i. boulleti* tested together, n=16) and intraspecific trials (solid line; only *H. hecale* tested, n=24). Table: total number of events at each behavioural unit registered across the totality of the trials. Y-axis: number of events at each step averaged by the number of trials.

Figure 7. Wing cues contribute to mate discrimination by males but not by females. (**A** and **B**) Male choice of *H. h. melicerta* and *H. i. boulleti* between female wings models of both species. Probability of males approaching (**A**) shortly (*localization*) and (**B**) sustainedly (*hovering*) the *H. i. boulleti* female wing models, where 1 means a complete choice of *boulleti* models and 0 a preference for *H. h. melicerta* models. A total of 42 *H. h. melicerta* males and 35 *H. i. boulleti* males were tested. Error bars show support limits equivalent to 95% confidence intervals. (**C**) Female choice experiments based on visual cues in *H. hecale.* The total events recorded in each of the main courtship steps during 3 hours long trials are shown on the

bar chart. Each *H. hecale* female could choose between two males of its own species and race. One of them was *treated* by modifying its colour pattern (yellow and white patterns on both sides of the forewings blacked with *Sharpie*). The other male (called *sham*) was black-painted on the black regions of the forewing. Only *H. hecale melicerta* male wings are shown here.

H. hecale melicerta and *H. ismenius boulleti* differ in their chemical blends

To understand premating isolation in this pair of species, we investigated chemical differences between them. Chemical extracts male and female abdominal glands, wings, and cuticle revealed a high variance in the composition and abundance of chemical compounds both between and within sexes and species. Consistently, certain peaks differed qualitatively or quantitatively between categories, most importantly from claspers (*C*) and male wing overlap (*OW*) (**Fig 8AI, 8BI**). Non-supervised multivariate analyses showed a good discrimination between males of the two species, and between males and females, but could not distinguish between females of the two species (perMANOVA for abdominal glands, both sexes together: $F=14.4$, $df=24$, $p=0.000$; perMANOVA for wings, both sexes together: $F=7.6$, $df=17$, $p=0.000$; perMANOVA for claspers: $F=19.6$, $df=11$, $p=0.002$, and for wings: $F=18.7$, $df=10$, $p=0.003$, males only; perMANOVA for glands: $F=1.1$, $df=12$, $p=0.322$, and for wings: $F=0.7$, $df=6$, $p=0.845$, females only; NMDS plots, **Fig 8AII, 8BII)**.

All compounds were associated with a unique numerical label (**Tables S12, S13 and S14**). A tentative identification of most of them (including major peaks of interest) was achieved either at the compound or at the class-level (**Table S15**) and, for male wing compounds, it closely matched the identification by Mann et al. (2017), even if they used a different solvent for the chemical extractions. Hydrocarbons are the best represented class of compounds in our extracts, and the abundance of several hydrocarbons consistently differed between species. We found that the claspers of males were chemically more diverse than the wing androconia (see also González-Rojas et al. 2020 and Darragh et al. 2020). Also, most compounds discriminating males of the two species were present or more concentrated in *H. h. melicerta* and absent or less concentrated in *H. i. boulleti* (**Fig 8AI** and **8BI**, **Table S12** and **Table S13**). In *H. h. melicerta* claspers, (*E*)-β-ocimene (compound number 3; **Fig. 8AI, Table S12**) dominated (~46.9 % of total), but was nearly absent in *H. i. boulleti*. We did not find α-ionone, previously found to be specific to *H. ismenius* claspers (Estrada et al. 2011); this may indicate differences in the subspecies of *H. ismenius* analysed. One compound of low concentration was exclusive to *H. i. boulleti* males (compound number 35), and three compounds were more concentrated in *H. i. boulleti*'s claspers (heneicosene [compound number 36], tricosene [number 48], and compound number 42) (**Fig 8AI** and **Table S12**). In total, forty-two compounds differed significantly in their concentration between the male genitalia of the two species (**Fig 8AI; Table S12**). Most of these variables were strongly correlated with the *H. hecale-H. ismenius* separation (**Fig S14**). Finally, the *indval* index outlined forty-four compounds significantly associated with either species, but especially with *melicerta* males (**Table S12**). A total of twenty-nine compounds are male-specific (**Table S12**).

Figure 8. Comparison of the chemical cocktails extracted from three different tissues between *H. hecale melicerta* and *H. ismenius boulleti*. Chemical composition was analysed for abdominal glands (*C* and *G*) (part **A**), wing overlap of both forewings and hindwings together (*OW*) (part **B**) and the cuticle in the base of the wings (*B*) (part **C**). In each part **A**, **B** and **C**, section **I** shows a heatmap indicating the log-transformed [$\log_{10}(x+1)$] relative concentration of each compound (horizontal bands) for males and females of both species (columns). Single individuals are labelled under each column. A unique numeric label is indicated for some compounds on the left margin. Section **II** shows non-metric multidimensional scaling (NMDS) ordination of the chemical cocktails of *H. h. melicerta* males (MMC, MMOW, MMB) and females (MFG, MFOW, MFB), and *H. i. boulleti* males (BMC, BMOW, BMB) and females (BFG, BFOW, BFB). Each dot represents a sample of any of these categories. Stress values are shown on the upper part of the graphs. See Table S12-S14 for details on the concentration of the compounds separately. Also see Table S15 for a tentative identification of some of those compounds.

Female abdominal glands (*AG*) showed a few small peaks **(Fig 8AI)**. Two low-concentration compounds differed between species. One is present only in *H. i. boulleti* glands (compound number 77; **Fig 8AI, Table S12**), and is also exclusive to the cuticle of *H. i. boulleti* females (**Table S14**). This

compound is also present on female wings of both species (**Table S13**). The second female gland compound is exclusive to *H. h. melicerta* (compound number 24; **Fig 8AI, Table S12**), but also found in male claspers of *H. h. melicerta*. Nevertheless, possibly due to their low general abundance, neither compound is significantly different between the females of both species (Wilcoxon rank-sum test; **Table S12**) or indicative of species status (*indval* index; **Table S12**).

In male wing extracts, two compounds dominate in *H. h. melicerta* and are absent or found in trace amounts in *H. i. boulleti* (**Fig 8BI** and **Table S13**). One is the alkane heneicosane (~57.1 % of total; number 38) and the other is hexahydrofarnesyl acetone (~7.87 % of total; number 22). This notable interspecific difference registered for these two compounds was also found by Mann et al. (2017). Here, seven compounds are found in higher abundance in *H. h. melicerta* (**Fig 8BI** and **Table S13**) and contribute to the *H. h. melicerta-H. i. boulleti* separation (**Fig 8BII, Fig S15**). The *indval* index identified six compounds associated with *H. h. melicerta* (**Table S13**). Female wing extracts showed fewer compounds of low abundance, highly conserved between species. Only one low concentration compound was exclusive to *H. h. melicerta* wings (2-eicosanyl-5-heptyl-tetrahydrofuran [compound number 79]) although this difference is not significant (**Table S13**). Overall, two wing compounds are male-specific and one low-concentration compound is female-specific (**Table S13**).

Cuticular extracts (*B*) have fewer compounds than other tissues, and are highly conserved between species and sexes (**Fig 8CI, Table S14**). The clustering by sex or species is not significant (NMDS plot, **Fig 8CII**; perMANOVA for all categories together: F=1.6, df=18, p=0.132; perMANOVA for males only: F=1.9, df=8, p=0.140; perMANOVA for females only: F=2.3, df=9, p=0.08).

Discussion

Introgression among species is known to play a role in mimicry adaptation in *Heliconius* butterflies because of relatively permeable species boundaries following species divergence (Martin et al. 2013; Jay et al. 2018; Martin et al. 2019; Van Belleghem et al. 2021, *Heliconius* Genome Consortium 2012). *H. hecale* and *H. ismenius* fall at the end of the species continuum proposed in Roux et al. (2016) with levels of net sequence divergence as high 0.16 to 0.22. Models based on extant patterns of genome wide variation suggest that the two species likely diverged in allopatry over 3.5 million years ago and then came into secondary contact (i.e. currently exchanged some gene). Despite very strong premating isolation and divergence in male chemical signals, these genomics observations reinforce the growing realization that species barriers are surprisingly permeable to introgression even among taxa that diverged in the relatively distant past. Our result contrasts with a previous study (Roux et al. 2016) that concluded that none of the six pairs of species with similar or higher levels of divergence exchange genes.

Nonetheless, gene flow between *H. hecale* and *H. ismenius* is much reduced relative to the pervasive gene flow observed between the similarly ecologically, behaviorally, and chemically distinct *H.*

melpomene and *H. cydno* (Bull *et al.*, 2006; Kronforst *et al.*, 2006; Martin *et al.*, 2013). The reported acceleration of genome-wide divergence accumulation towards later stages of speciation (Kronforst *et al.*, 2013) seems to be illustrated here. *Heliconius hecale* and *H. ismenius* have diverged for ~3,6 My, which is ~1,5 My longer than *H. melpomene* and *H. cydno* (Kozak *et al.*, 2015). This longer history of divergence may explain the few localised windows of introgression along the genome (*e.g.* chromosome 18, 19, 07), introgression having likely been eroded by recombination and removed by selection in most of the genome (Brandvain et al. 2014; Juric et al. 2016). This is supported by a negative correlation between levels of absolute sequence divergence (D_{XY}) and recombination. In addition, the modelling approach and the observed relationship between differentiation-based statistics and recombination indicate a contribution of linked selection to the heterogeneous landscape of divergence (Cutter and Payseur 2013), which is consistent with growing evidence that linked selection may strongly influence divergence patterns during speciation (Burri et al. 2015; Burri 2017; Rougemont and Bernatchez 2018; Van Belleghem et al. 2021; reviewed in Jensen and Charlesworth 2021).

Although generally low overall, localized introgression events were nearly twice as high in subpopulations of each species that share the same warning color pattern. This was also supported by our D_{FOIL} analysis revealing that the most significant tests of introgression involved co-mimics. One of the strongest signals we observed was around the transcription factor *optix*. The gene *optix* is known to be involved in determining red-orange colour pattern elements. Intraspecific differences in red pattern on hindwings in *H. hecale* and in *H. ismenius* is largely modulated by a QTL linked to optix (Huber et al. 2015), which explained a large proportion of the extent of black vs. red/orange color, while forewing diferences and pattern boundaries in black and yellow elements are largely modulated by a QTL near the gene WntA (Huber et al. 2015; Concha et al. 2019). Despite a still incomplete understanding of how the major *Heliconius* color pattern loci function and interact, our genealogical analysis here suggests that one allele may have been, at least partially, transferred among species, bringing support for its implication in pattern evolution under mimicry selection. A limit of this study is that while both the f_d and D_{FOIL} consistently suggest a shared origin of *optix*, the topology inferred from *twisst* was not entirely consistent with an origin as a result of recent introgression.

Speciation in *Heliconius* is multifactorial (Mérot et al. 2017), so the strong genomic divergence we observe likely results from the built-up and strengthening of multiple barriers to gene flow. Color patterns often show conspicuous differences between closely related *Heliconius* species. This trait appears to be important in some aspects of mating behaviour, including male approach (Haussman & al. 2021). In the case of speciation between the *Heliconius hecale* and *H. ismenius* lineages, the shift in wing patterns is however less striking than in many studied cases. The general *melicerta/boulleti* pattern is similar to other races from Western Ecuador (*H. h. australis, H. i. metaphorus*), Northern Andes (*H. h. rosalesi, H. i. tilletti*), but also in the disjunct region of Lower Amazon (*H. h. metellus, H. h. novatus*), suggesting that this wing pattern could be fairly ancient in *H. hecale*. *H. hecale zuleika* and *H. ismenius clarescens*, have very

distinguishable patterns but share similar colours (Figure 1). Not surprisingly, given the similarity in patterns, our behavioral data suggest visual confusion of males in the approach behavior. Nonetheless, short-range cues seem to be very important : males quickly adjust their behavior when in close contact with females, males that court females are strongly rejected by females, and no heterospecific mating was observed in our experiments.

Consistent with a role for short range signals in reproductive isolation, the chemical profiles of male *H. ismenius* and *H. hecale* are both extremely diverse and highly divergent. In contrast, female chemical profiles are nearly identical. This has been observed by in other *Heliconius* species (Darragh et al. 2017; González-Rojas et al. 2020; Bergström and Lundgren 1973; Schulz et al. 1993; Andersson et al. 2007; Costanzo and Monteiro 2007; Nieberding et al. 2008) and suggests a role for selection acting on short range chemical signals produced by male wings and genitalia and assessed by females during courtship. Nevertheless, males approach both con- and heterospecific female wings at the same rate, yet they hover over conspecifics at a higher rate, highlighting the use of short-range cues to determine whether or not to court. It is possible that these cues are visual and may reflect subtle differences between species in pattern or how pattern is perceived (Dell'Aglio et al; 2018).

The chemical profiles reveal differences in compounds with known roles in chemical communication in other species : (*E*)-β-ocimene, present in male *H. hecale* genitalia but nearly absent in *H. ismenius* is also known from *H. melpomene, H. erato,* and *H. numata* where it may act as an antiaphrodisiac transferred to females during copulation (Mérot et al. 2015, Darragh et al. 2017, González-Rojas et al. 2020; Schulz et al. 2008; Estrada et al. 2011; but see Klein and Araújo 2010). The wings and claspers of *H. h. melicerta* males, but not those of *H. i. boulleti*, expresses highly volatile compounds ((*E*)-β-ocimene, hexahydrofarnesyl acetone), consistent with a putative role as an air-borne pheromone. The latter is very abundant in the androconial bouquet of *H. timareta florencia* androconia and *H. melpomene malleti* claspers (González-Rojas et al. 2020). Other compounds tend to be non-volatile, but could play a role during courtship (Millar 2000; Jurenka et al. 2003; Ferveur 2005; Hay-Roe et al. 2007; Dapporto 2007; Heuskin et al. 2014) either at close range (Rutowski 1991; Vane-Wright and Boppré 1993) or via direct contact (Crane 1955; Mega and Araújo 2010; Klein and Araújo 2010, Loudon and Koehl 2000).

H. hecale larvae feed on *Passiflora vitifolia,* whereas *H. ismenius* larvae feed on *P. quadrangularis* and *P. ambigua*. Differences in host plant utilisation, might play an important role in microhabitat segregation as has been suggested in other insects (Mérot et al. 2015; Wood and Keese, 1990; Via et al. 2000; Emelianov et al. 2001; Thomas et al. 2003), either by reducing the probability of encounter or by influencing chemical profiles. Neither of these alternatives have been explored much in *Heliconius* and it is possible that differences in larval host plants help to reduce competition and allow the two species to co-occur. Incipient species of *Heliconius* that have not evolved differences in host plant preference tend to have parapatric distribution, even when assortative mating is strong (see McMillan et al., 1997).

Taken together, our results indicated that the pair of species studied here falls in the range of the speciation continuum where reproductive isolation is nearly complete, and may involve the recruitment of multiple barriers to hybridization, including chemical signatures. Contrary to other pairs of taxa showing a high level of incomplete lineage sorting throughout the genome, here the genomes are indeed well sorted. Yet we were able to detect localised segment with increased gene flow among co-mimics as compared to non co-mimics, including around the wing patterning gene *optix* (Reed et al. 2011, Martin et al. 2014; Zhang et al. 2017; Huber et al. 2015). The small length of introgressed ancestry suggests a distant event of adaptive introgression at *optix*, although we could not entirely rule out incomplete lineage sorting. If this introgression is indeed relatively old, mimicry might therefore have evolved at a time where the species were less isolated genetically, followed by further chemical differentiation and the near completion of speciation. A better understanding of the process of accumulation of barriers to gene flow requires an accurate understanding of the trajectory of key genes determining hybridization during the history of lineage divergence.

Materials and methods

Specimens.

In Eastern Panama, *H. hecale* and *H. ismenius* are excellent co-mimics of each other (subspecies *H. h. melicerta* and *H. i. boulleti*) and both display a pattern made of orange (proximal) and yellow elements (distal) bordered by a thick black margin and a black wing tip. In Western Panama, those species display distinct patterns and join different mimicry rings. *H. h. zuleika* shows a pattern with a largely black forewing and prominent yellow dots, and a merely orange hindwing. In contrast, *H i. clarescens* shows a largely orange wing pattern with black and yellow stripes alternating in the distal part of the forewing.

For genomic sequencing, three to six specimens of each subspecies were collected from Darién (East of Panama) and near David, Chiriquí (West of Panama), totalling eighteen specimens (**Fig 1** and **Table S1**). Bodies were preserved in a NaCl-saturated DMSO solution and wings in paper envelopes. In addition, four samples of *H. hecale felix* collected in Peru, four *H. hecale clearei* from Venezuela and two *H. ismenius telchinia* from Panama were used four broader context in PCA and structure analyses and as control allopatric populations of the study species in our test of gene flow in ABBA-BABBA related methods (**Fig 1** and **Table S1**). Closely related silvaniforms namely 15 *H. pardalinus* and 17 *H. numata silvana* (**Fig 1A**) were also used when assessing gene flow in ABBA-BABBA related methods (detailed in table S1). Finally, a set of 12 *H. melpomene* individuals were used as an outgroup. For behavioural and chemical data *H. h. melicerta* and *H. i. boulleti* specimens were collected in Darién (Eastern Panama) and around Gamboa, Colón (Central Panama), while *H. h. zuleika* specimens were collected from Chiriquí (Western Panama) (**Fig 1**). Stocks of these races were reared at the Smithsonian Tropical Research Institute in Gamboa (Panama) (Huber et al. 2015).

Population Genomics

DNA extraction. DNA was isolated from preserved bodies using the DNeasy Blood and Tissue Kit (Qiagen). Separate Illumina paired-end libraries were generated according to the manufacturer's protocol (Illumina Inc.). Each library was shotgun sequenced to an average coverage of ~25X on an Illumina Hi-Seq 2000 with 2 x 100-base read length.

Genotyping and variant calling. Reads were trimmed using fastp (Chen et al. 2018), aligned to the *H. melpomene v2.5* genome with bwa-mem v.0.7.13 (Li 2013) and filtered with samtools (Li et al. 2009) requiring a minimum quality of 20. Duplicates were removed using picard (http://broadinstitute.github.io/picard/). SNP calling was then performed using GATK V4.1.9 (DePristo et al. 2011). We included variables and invariable sites across the whole genome. Following GATK Best Practices, we marked all sites that did not match the following criterion: MQ < 30, QD < 2, FS > 60, MQRankSum < -20, ReadPosRankSum < 10, ReadPosRankSum > 10. The dataset with variants and invariants sites was used for ABC analyses below. We then extracted SNPs from the dataset and used it for all the remaining analyses. This dataset was filtered to keep SNPs with a minor allele count of 2. Genotypes without a mean depth between 5 and 60 (*i.e.* mean+2 standard deviations) and a genotype quality above 30 were set as missing resulting in ~5.5 millions SNPs.

Phasing and recombination rate estimates. Phasing was performed using beagle v.5.1 (Browning and Browning 2007) with default parameters. We then used LDHat software (McVean et al. 2002) to estimate effective recombination rates ($\rho=4.Ne.r$ where r represents the recombination rate per generation and *Ne* is the effective population size) along the genome. The genome was splitted in chunks of 2,000 SNPs with overlapping windows of 500 SNPs to compute recombination rate and data were then merged together. Recombination rates were averaged into 250-kb windows using a custom Python script. Recombination rate was inferred for each species separately (*H. hecale, H. ismenius)* as well as for outgroups (*H. numata, H. pardalinus)* to analyse how conserved the landscape are along the divergence continuum (see Supplementary Results).

Genetic diversity (π) and Tajima's D were measured for each species along 50 kb windows after converting whole genome vcf file into fasta using seq_stat (https://github.com/QuentinRougemont/PiNPiS).

Population relationship. We assessed genetic relationship among individuals with a PCA using the R package ade4 (Dray and Dufour 2007) based on a LD-pruned dataset, further thinned to keep SNPs spaced at least 500 bp from each other (365K SNPs). We tested population treeness in Treemix (Pickrell and Pritchard 2012) with the same LD-pruned dataset and allowing up to 20 migration events. We chose the model that explained the highest level of variance when adding a migration edge. We performed 500 bootstrap replicates of the best model to obtain robustness of the nodes.

Global demographic history. We tested if the species have diverged in the presence or absence of gene flow through historical times. To do so coalescent simulations in an ABC framework were used. Our

models account for the confounding effect of linked selection, locally reducing the effective population size, and of barriers to gene flow, locally reducing the effective migration rate (Roux et al. 2016; Rougemont and Bernatchez 2018). We extend the two populations models previously developed to a four population model, representative of the four focal population studied. It includes ABBA-BABBA related statistics explicitly developed for detecting gene flow at different time scales. We performed a statistical comparison of a model of divergence without gene flow (SI) against allopatric divergence followed by a secondary contact (SC) (**Fig S06 A-B**). The best of the two models was then compared against models of sympatric divergence with gene flow, namely Isolation w. Migration (IM) and Ancient Migration (AM) (**Fig S06 C-D**). The SI model assumes that an ancestral population of size N_{ANC} splits instantaneously at time Tsplit into two daughter populations of constant effective population size N_{12} and N_{34} and no gene flow. Afterwards population will continue to diverge to give rise to two subpopulations (*H. h. zuleika* and *H. h melicerta*, of size N_1 and N_2) at time T_{12} and two other subpopulations (*H. i. boulleti* and *H. i. clarescens*, of size N_3 and N_4) at time T_{34} (**Fig S06**). These subpopulations were connected by continuous gene flow, $M = 4N_0.m$ with $M_{x \leftarrow y}$ being the number of migrants from population Y into population X at each generation. Under the SC model the two species diverged under a period of strict isolation before undergoing a secondary contact from Tsc until present. We tested all directionality of secondary contacts from a single directionality of migration to a set of fully connected populations (**Fig S06 B**) which enable to test the hypothesis of gene flow either between co-mimics (direction A <-> C in **Fig S06 B)** or between non co-mimics (direction B <-> D in **Fig S06 B**). Under AM the instantaneous split of the ancestral population is followed by a period of gene flow from Tsplit until the split of the two daughter population. The gene flow is stopped when the first pairs of populations splitted in two subpopulations (**Fig S06 C**). As for the SI, we allowed gene flow between each subpopulation within *H. hecale* and *H. ismenius* respectively. In the IM there is continuous gene flow following the split of the ancestral population into two. Following the split of the subpopulations within species we then allowed for gene flow either between co-mimics (direction A <-> C in **Fig S06 D**) or between the non- co-mimics (direction B <-> D in **Fig S06 D**). In all models, migration priors were sampled independently for each direction, enabling asymmetric introgression between species or populations.

Coalescent simulation. For each model, 1×10^6 simulations of datasets matching the sample size of each locus was performed using msnsam (Ross-Ibarra et al. 2008) a modified version of the coalescent simulator ms (Hudson 2002). Selected markers can provide additional information regarding a species history (Bierne et al. 2013; Roux et al. 2013). Therefore, we restricted our analyses to the CDS from the *Heliconius melpomene* genome, using a total of 6,600 informative loci falling in those CDS and passing our filtering criterion. Large and uninformative priors were used. Effective population size for all current and ancestral populations were sampled uniformly on independent intervals [0–40,00,000]. Priors for the divergence time (T_{split}) were uniformly sampled on the interval [0–5,000,000], so that T_{12}, T_{34} and the various T_{sc} were constrained to be chosen within this interval. For the homogeneous version of the migration rate, M was sampled on the interval [0–40] independently for each direction of migrations. We then used beta

distribution hyperprior to model the heterogeneity in effective population sizes and migration rate (Roux et al. 2016; Rougemont and Bernatchez 2018). Heterogeneity in migration was modelled using a beta distribution as a hyperprior shaped by two parameters; α randomly sampled on the interval [0–20] and βrandomly sampled on the interval [0–200]. A value was then independently assigned to each locus. For the heterogeneity of Ne, α percent of loci evolved neutrally and share a value uniformly sampled on the interval [0–1]. Therefore 1–α percent of loci were assumed to evolve non neutrally. Their Ne values were thus drawn from a beta distribution defined on the interval [0–20], as defined on the homogeneous version. Ne values were independently drawn for N_{anc}, N_1, and N_2 ,N_3 ,N_4, but shared the same shape parameters (Roux et al. 2016). Priors were generated using a Python version of priorgen software (Ross-Ibarra et al. 2008). We used the mutation rate of 2.9×10^{-9} mutations per bp and generation estimated in *H. melpomene* (Keightley et al. 2015).

Summary statistics. A modified version of mscalc was used to compute summary statistics (Ross-Ibarra et al. 2008; Ross-Ibarra et al. 2009; Roux et al. 2011). This version, coded in pypy for faster computation, provided the average and standard deviation of statistics computed over loci. We included, the number of biallelic sites, the number of fixed differences between species X and Y (sf), the number of exclusively polymorphic positions in species X (sx), the number of shared biallelic positions between species X and Y (ss), the levels population genetic diversity (Tajima's π and Watterson's   ) and Tajima's D as well pairwise statistics between X and Y populations: levels of population differentiation (F_{ST}), population absolute and net nucleotide divergence (D_{XY} and D_A) respectively, the smallest divergence measured between one individual from X and one from Y (minDiv), the highest divergence measured between one individual from X and one from Y (maxDiv) as well as a range of statistics known to be informative with regards to introgression, namely ABBA-BABBA' D statistics (Durand et al. 2011), Fd statistics and Dfoil (Pease and Hahn 2015). Finally, between population Pearson correlation coefficients in genetic diversity (π and   ) were also computed.

Random Forest and Neural Networks model choice. Two approaches were used to perform model choice. First, posterior probabilities were computed using a classic ABC procedure. We retained the 1000 "best" simulations (out of 1 million) that were weighted by an Epanechnikov kernel peaking when Sobs = Ssim and a neural network that performs a non-linear multivariate regression where the model is a response variable to be inferred. Fifty feedforward neural networks were used for training with 15 hidden layers in the regression. Ten replicates ABC model choices were performed to obtain the average posterior probability of each model. The R package "abc" (Csilléry et al. 2012) was used for the model choice procedure. Because classical ABC may suffer from the curse of dimensionality, especially for a complex model with many summary statistics as here, we also used a Random Forest approach for model choice (*e.g.* (Rougemont et al. 2016). This approach has the advantage of directly assessing the proportion of correct model choice by the algorithm. We constructed forest made of 500 trees and included a linear discrimination analysis as additional summary statistics as implemented in the packages ABCRF (Pudlo et

al. 2016). All available summary statistics were used to construct a confusion matrix, estimate classification error and perform the model choice. The IM and AM models were compared between each other and to other models using only the ABCRF procedure. The whole pipeline is available at https://github.com/QuentinRougemont/ABC_4pop

Parameter Inference. The posterior probabilities of each parameter value were estimated using the neural network procedure with nonlinear regressions of the parameters on the summary statistics using 50 feed-forwards neural networks and 15 hidden layers after a logit transformation of the parameters. A tolerance of 0.001 was applied. The ABC-RF was also used to estimate demographic parameters of the best models using 1,000 trees to build a forest.

Genomic differentiation and divergence analyses to test for linked selection. Divergence among species and populations was measured by estimating nucleotide diversity (π), levels of divergence (D_{XY}) and of genetic differentiation (F_{ST}) in windows of 250kb along the genome using Python scripts from Martin et al. (2013). Patterns of heterogeneous genome-wide divergence can be due to genetic hitchhiking of neutral alleles linked to selective sweeps (Smith and Haigh, 1974) or to background selection (BGS, Charlesworth et al. 1993). These combined effects, referred to as linked selection, reduces polymorphism at sites closely linked to advantageous or deleterious variants (Payseur and Nachman 2002). The intensity of selection on linked loci is mostly modulated by variation in local recombination rate and by gene density (Kaplan et al. 1989). Under linked selection, diversity (π, D_{XY}) and differentiation (F_{ST}) metrics are expected to be positively and negatively correlated with genome-wide variation in recombination rate respectively (Payseur and Nachman 2002). We therefore tested for correlation among π, D_{XY} and F_{ST} separately among species to test for the conservation of these landscapes individually (**Fig S03 A-C**). We then used a PCA to obtain a synthetic view across all populations and species of our estimates of π, D_{XY} and F_{ST} separately. Next, Spearman correlations based on each (z-transformed) variable projected on the PC1 axis separately were carried out to look at correlation among variables (**Fig S03 D, Fig S04**).

Testing for interspecific gene flow. Gene flow was evaluated between species (*H. hecale* vs. *H. ismenius*), between co-mimics (*H. h. melicerta* vs. *H. i. boulleti*) and between non co-mimics (*H. h. zuleika* vs. *H. i. clarescens*) using the same ABBA-BABA test as implemented in our ABC pipeline, namely we calculated D over the whole genome (Green et al. 2010; Durand et al. 2011) and f_d (Martin et al. 2013) over the whole genome and in windows. Given three taxa *P1, P2, P3* where *P1* and *P2* are sister taxa, and an outgroup *O* with the following topology [(*P1, P2*), *P3*], *O* we tested for gene flow between *P2* (the target population) and *P3* the donor population. We used *H. melpomene* as outgroup and tested two configurations to quantify gene flow between the co-mimics *H. h. melicerta* and *H. i. boulleti* on the one hand, and between the non-mimetic pair *H. h. zuleika* and *H. i. clarescens* on the other hand. In addition, we tested other more distant P1 species namely *H. pardalinus* and *H. numata* as well as *H. ismenius telchinia.* A 1Mb block jack-knifing approach was implemented to calculate the mean and variance of *D* and f_d, and to test whether *D* differed

significantly from zero. We calculated f_d statistics in 10kb sliding windows along the genome using the custom scripts developed by Martin et al. (2013).

We complemented our analyses with the D_{FOIL} statistics (Pease and Hahn, 2015), that enable testing the topology [(P1,P2)(P3,P4)],O), with a potential higher power to test the directionality of introgression than the above test. In addition, this test enables the direct consideration of the four co-mimics and non co-mimics species in a single test. As for our ABC pipeline, we modified the original D_{FOIL} statistics to work on allele frequency of multiple individuals instead of the sequence of a single sample.

Finally, we quantify the different topology relationship between mimics and non co-mimics using the tree weighting approach implemented in Twistt (Martin and Van Belleghem 2017). We tested the proportion of each of the 15 different possible topologies (detailed in Fig S12), with a particular focus on topology reflecting gene flow between the co-mimics species (*H. h. melicerta and H. h. boulleti)*, or gene flow between the non co-mimics (*H. h. zuleika vs H. i. clarescens*). The complete method, which considers all subtrees, was used to estimate the exact weighting to the full tree. We then classified topologies that reflect: the true species trees (1 topology), partial tree with incongruence in *H. hecale* only (2 topologies), partial tree with incongruence in *H. ismenius* only (2 topologies) or perfect geography (Geographic perfect (*H. i. boulleti + H. h. melicerta* AND *H. i. clarescens + H. h. zuleika* together, 1 topology):), Geographic partial East (*H. i. boulleti + H. h. melicerta* together NOT *H. i. clarescens + H. h. zuleika*): 2 topologies corresponding to higher gene flow between co-mimics, Geographic partial West (*H. i. clarescens + H. h. zuleika* together NOT bou+mel) : 2 topologies, Incongruent (boul+zul AND/OR cla+mel) : 5 topologies (but see Fig S12).

Behavioural and Chemistry analyses

Courtship description. To define the sequence and prevalence of behavioural steps during courtship and mating in *H. hecale* butterflies, behaviour was video-recorded until mating happened (successful courtships, n=4) or until at least one whole sequence including all steps but mating had been registered (unsuccessful courtships, n=13). Our observations were supported by other behavioural experiments in this (detailed below) and previous studies concerning other species in the subfamily Heliconiinae (Klein and Araújo 2010; Crane 1955, 1957; Rutowski and Schaefer, 1984; Mega and Araújo, 2010).

Inter- and intraspecific sexual interactions. To unravel the signals involved in species recognition between *H. hecale melicerta* and *H. ismenius boulleti,* we performed no-choice interspecific encounter experiments. Three mature (>5 days old) conspecific males of one species were put in a 2x2x2m cage with one newly-emerged (a few hours old) heterospecific female. All male courtship events and female responses occurring during 15-min trials were recorded on video. Main courtship steps (explained below) were registered for 9 trials with *H. h. melicerta* males and 7 with *H. i. boulleti* males. Intraspecific no-choice trials were run as controls using *H. hecale* (n=24), and involved two mature males and two virgin females of each of the two races *H. h. melicerta* and *H. h. zuleika.*

Male choice using female wing models. To determine the importance of female wing cues in courtship decisions by males, choice experiments were performed where males of one species were presented with female wing models of both species. We registered *localization* events (close approach to models) and *hovering* events (sustained stationary flight close to the models for >=3 seconds). Sexually mature, reared (n=44) or field-caught (n=33) males were tested multiple times, on different days, to obtain independent behavioural registrations for each individual. A total of 38 *H. h. melicerta* and 32 *H. i. boulleti* were characterised by three registrations, plus 4 *H. h. melicerta* and 3 *H. i. boulleti* males had only two registrations each. Models were made with real unwashed female wings, thus including visual and chemical signals. Wings were connected to a plastic wire and moved in a fluttering motion inside a 2x2x2 m cage. Trials were run for 5 minutes, presenting two models simultaneously, and switching model positions in the middle. *Localization* and *hovering* were analysed separately. The relative probability of males courting *boulleti* female models rather than *melicerta* models was estimated using a likelihood function as follows (Edwards 1972; Jiggins et al. 2001):

$\ln(L) = \Sigma\ n_x bou \ln(P_y mel) + n_x mel \ln(1-P_y mel)$

where $P_y mel$ denotes the probability of courtship by male *y* towards *melicerta* wings, $n_x bou$ the total number of events of male *x* directed towards *boulleti* models, and $n_x mel$ towards *melicerta* models. The solver option in Excel (Microsoft) was used to estimate the probabilities of male courtship by numerically searching for values of $P_y mel$ that maximised $\ln(L)$. Support limits (asymptotically equivalent to 95% confidence intervals) were assessed by looking for values that decreased $\ln(L)$ by two units (Edwards, 1972). To compare courtship across species, we compared a model with equal relative probabilities

(*Pmel=Pbou*) to a model with probabilities calculated separately for each species (*Pmel≠Pbou*), using a likelihood ratio test assuming a χ^2-distribution with one degree of freedom for the test statistic $G=2\Delta\ln(L)$.

Female choice based on colour. To assess whether visual cues are important in mating acceptance by *Heliconius* females, we performed experiments involving a virgin *H. hecale* female (subspecies *melicerta* or *zuleika*) and two mature males of their own subspecies with two different colouration treatments (**Fig 7C**). *Treated* males were painted with black ink, using a *Sharpie* pen, following Kemp (2007), to cover the white and yellow patches of dorsal and ventral forewings. *Sham* males, or controls, were painted on the black region of the forewing, covering a similar surface as for *treated* males but avoiding pattern modification. Experiments were run in 1x2x2m cages. In this case females were over one day old, thought to be better able to reject courting males than just-emerged females. A total of 26 trials were performed (13 with *H. h. zuleika* females and 13 with *melicerta* females) resulting in 20 successful courtships. Direct observation of sexual behaviour was carried out until mating occurred or for a maximum period of 3 hours, and all main courting events were registered.

Chemical analyses. To evaluate differences in chemical signalling between *H. h. melicerta* and *H. i. boulleti*, chemical compounds were extracted from three different tissues. In males, wing regions where forewing and hindwing overlap (*OW*), and abdominal glands (claspers, *C*) have been proposed to contain pheromone-producing glands (androconia) releasing volatile signals by *Heliconius* males (Emsley 1963; Brown 1981; Schulz et al. 2008; Estrada et al. 2011; Darragh et al. 2017). Similarly, male's genital claspers are often open during courtship, putatively diffusing volatile signals perceived by the female (Schulz et al. 2008; Estrada et al. 2011). Finally, female rejection might involve the evagination of abdominal glands, which in mated females contain volatile anti-aphrodisiacs compounds (e.g. Rutowski and Schaefer 1984, Schulz et al. 2008). Our description of courtship behaviour suggests they might be used both in *H. hecale* and *H. ismenius*. In virgin females, abdominal glands (*G*) were analysed given their outstanding role in producing sex pheromones in several species of Lepidoptera. Additionally, the cuticle of the body close to the base of the wings (*B*), was analysed because of the presence of small bristles which might be involved in the production of chemical signals (*i. e*. hairpencils, Boppré, 1978; Bacquet et al. 2015) diffused by the wings. Tissues from 3-7 individuals were extracted separately from virgin females and mature males of both species, right after sacrificing the individuals. Pieces of tissue were immersed into 130-200 µl of hexane containing dodecane 100 ng/µl as internal standard. Extractions were not stopped. Elution samples were analysed with gas chromatography coupled to mass spectrometry (GC-MS) with a Bruker Autosampler SCION SQ 436-GC, using a non-polar Rtx-5 MS fused silica capillary column (0.32 mm i.d., 30 m long, 0.25 µm thick film) and helium as the gas carrier. Volumes of 0.5 µL were injected automatically in a split-less mode with the injector temperature at 250°C (see Mérot et al. 2015 for more details). Peak area was used as a measure of compound abundance relative to the concentration of dodecane in the sample. Since chemical receptor sensitivity is unknown for these butterflies, we refrained from excluding low-concentration peaks from our descriptions. However, peaks that were too low for

quantification were treated as missing data for a given extract. Quantification values were log-transformed [$\log_{10}(x+1)$], allowing comparison of peaks with large differences in abundance (Lecocq et al. 2013).

Statistical analysis of chemical data. Chemical similarities and differences between species and sexes were visualised with heatmap graphs on the relative abundance of the compounds present in each tissue separately, using *Matlab R2012a Student Version*. Since chemical bouquets are likely to be perceived as composite signals, multivariate analyses were performed on the concentration of all compounds present in each tissue separately, where each compound is treated as a variable. For each tissue, analysis of compositional similarity of chemical cocktails between species was performed with sexes pooled as well as separating by sex, using non-supervised non-metric multidimensional scaling (NMDS) ordination implemented in the *vegan* (Oksanen et al. 2020) packages in *R*. Prior to NMDS, the Bray-Curtis distance matrix of the chemical composition was computed. The appropriate number of dimensions was chosen for each dataset by plotting a "scree diagram" (stress versus number of dimensions). The significance of chemical differences between species or sexes was assessed by a discriminant analysis on the Bray-Curtis similarity matrix, using a permutational multivariate analysis of variance (perMANOVA; Anderson, 2001) implemented in the *vegan* package (Oksanen et al. 2013), with 1000 permutations. Whenever categories were significantly different, partial least squares discriminant analysis (PLS-DA) was applied to the dataset using *R* packages *mixOmics* and *DiscriMiner* to identify variables (*i.e.* compounds) most diagnostic of a given category. The *indval* index, which gives the probability of association between a compound and a given category, was also estimated by computing indicator values that suggest which compounds are indicative of a given category. For that purpose, the *indval* function in the *R* package *labdsv* (Roberts 2013) was used, following Heuskin et al. (2014). Finally, a comparison of the per-compound concentration was performed with a Kruskal-Wallis test coupled with a pairwise Wilcoxon rank-sum test, and Bonferroni correction for multiple testing. To facilitate the identification of the compounds, linear retention indices (LRI) were calculated for each compound relative to the retention times of a mix of alkane standards (C10-C26). So far, a partial number of compounds were tentatively identified by comparison of retention indices and mass spectra with those of authentic reference standards (Sigma-Aldrich Corporation), with published databases (Wiley Registry of Mass Spectral data-7[th] edition and NIST Chemistry WebBook; NIST Chemistry WebBook 2015) and with compounds previously identified (Estrada et al. 2011; Schulz et al. 1998; Schulz et al. 2008). Other peaks of interest remain to be identified and the current identification of some of the compounds remains to be confirmed before manipulative assays of their role in species recognition between *H. hecale* and *H. ismenius* can be envisioned.

Data Availability

Raw data are available on NCBI (see details in Table S1). Filtered vcf will be deposited on Zenodo. Scripts to reproduce all genomics analyses are available on the first author github page.

References

Aeschbacher S, Selby JP, Willis JH, Coop G. 2017. Population-genomic inference of the strength and timing of selection against gene flow. *Proceedings of the National Academy of Sciences* 114:7061–7066.

Anderson MJ. 2001. Permutation tests for univariate or multivariate analysis of variance and regression. *Can J Fish Aquat Sci* 58: 626–639.

Andersson J, Borg-Karlson A-K, Vongvanich N, Wiklund C. 2007. Male sex pheromone release and female mate choice in a butterfly. *J Exp Biol* 210: 964–970.

Bacquet PM, Brattström O, Wang HL, Allen CE, Löfstedt C, Brakefield PM, Nieberding CM. 2015. Selection on male sex pheromone composition contributes to butterfly reproductive isolation. Proc Biol Sci. Apr 7;282(1804):20142734. doi: 10.1098/rspb.2014.2734. PMID: 25740889; PMCID: PMC4375863.

Barton N, Bengtsson BO. 1986. The barrier to genetic exchange between hybridising populations. *Heredity* 57:357–376.

Bergström G, Lundgren L. 1973. Androconial secretion of three species of butterflies of the genus *Pieris* (Lep., Pieridae). *Zoon Suppl* 1: 67–75.

Bierne N, Gagnaire P-A, David P. 2013. The geography of introgression in a patchy environment and the thorn in the side of ecological speciation. *Current Zoology* 59:72–86.

Boppré M. 1978. Chemical communication, plant relationships, and mimicry in the evolution of danaid butterflies. *Entomologia Experimentalis et Applicata*. 24(3):64-77. DOI: 10.1007/bf02385074.

Boppré M. 1984. Chemically mediated interactions between butterflies.

Brandvain Y, Kenney AM, Flagel L, Coop G, Sweigart AL. 2014. Speciation and Introgression between *Mimulus nasutus* and *Mimulus guttatus*. *PLOS Genetics* 10:e1004410.

Brower AVZ. 1996. A new mimetic species of *Heliconius* (Lepidoptera: Nymphalidae), from southeastern Colombia, revealed by cladistic analysis of mitochondrial DNA sequences. *Zool J Linn Soc* 116: 317–332.

Brown KS. 1981. The Biology of *Heliconius* and Related Genera. *Annu Rev Entomol* 26: 427–457.

Browning SR, Browning BL. 2007. Rapid and Accurate Haplotype Phasing and Missing-Data Inference for Whole-Genome Association Studies By Use of Localized Haplotype Clustering. *The American Journal of Human Genetics* 81:1084–1097.

Bull V, Beltrán M, Jiggins CD, McMillan WO, Bermingham E, Mallet J. 2006. Polyphyly and gene flow between non-sibling *Heliconius* species. *BMC Biol* 4: 11.

Burri R, Nater A, Kawakami T, Mugal CF, Olason PI, Smeds L, Suh A, Dutoit L, Bureš S, Garamszegi LZ, et al. 2015. Linked selection and recombination rate variation drive the evolution of the genomic landscape of differentiation across the speciation continuum of *Ficedula* flycatchers. *Genome Res* 25:1656–1665.

Burri R. 2017. Linked selection, demography and the evolution of correlated genomic landscapes in birds and beyond. *Mol Ecol* 26:3853–3856.

Charlesworth B, Morgan MT, Charlesworth D. 1993. The Effect of Deleterious Mutations on Neutral Molecular Variation. *Genetics* 134:1289–1303.

Charlesworth B, Jensen JD. 2021. Effects of Selection at Linked Sites on Patterns of Genetic Variability. Annual Review of Ecology, Evolution, and Systematics 52:177–197.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 34:i884–i890.

Chen T., He T, Benesty M, V. Khotilovich, Y. Tang, H. Cho, K. Chen, R. Mitchell, I Cano, T. Zhou, M. Li, J. Xie, M. Lin, Y. Geng, Y. Li and J. Yuan, 2022,(2022) xgboost: Extreme Gradient Boosting, R package version 1.6.0.1},

Cornuet J-M, Pudlo P, Veyssier J, Dehne-Garcia A, Gautier M, Leblois R, Marin J-M, Estoup A. 2014. DIYABC v2.0: a software to make approximate Bayesian computation inferences about population history using single nucleotide polymorphism, DNA sequence and microsatellite data. *Bioinformatics* 30:1187–1189.

Costanzo K, Monteiro A. 2007. The use of chemical and visual cues in female choice in the butterfly *Bicyclus anynana*. *Proc R Soc B Biol Sci* 274: 845–851.

Crane J. 1955. Imaginal behaviour of a Trinidad butterfly, *Heliconius erato hydara* Hewitson, with special reference to the social use of color. *Zool N Y* 40: 167–196.

Crane J. 1957. Imaginal behaviour in butterflies of the family Heliconiidae: changing social patterns and irrelevant actions. *Zool N Y* 42: 135–145.

Cruickshank TE, Hahn MW. 2014. Reanalysis suggests that genomic islands of speciation are due to reduced diversity, not reduced gene flow. *Molecular Ecology* 23:3133–3157.

Csilléry K, François O, Blum MGB. 2012. abc: an R package for approximate Bayesian computation (ABC). *Methods in Ecology and Evolution* 3:475–479.

Cutter AD, Payseur BA. 2013. Genomic signatures of selection at linked sites: unifying the disparity among species. *Nat Rev Genet* 14:262–274.

Dapporto L. 2007. Cuticular lipid diversification in *Lasiommata megera* and *Lasiommata paramegaera*: the influence of species, sex, and population (Lepidoptera: Nymphalidae). *Biol J Linn Soc* 91: 703–710.

Darragh K, Vanjari S, Mann F, Gonzalez-Rojas MF, Morrison CR, Salazar C, Pardo-Diaz C, Merrill RM, McMillan WO, Schulz S, Jiggins CD. 2017. Male sex pheromone components in *Heliconius* butterflies released by the androconia affect female choice. PeerJ. Nov 7;5:e3953. doi: 10.7717/peerj.3953. PMID: 29134139; PMCID: PMC5680698.

Darragh K, Montejo-Kovacevich G, Kozak KM, Morrison CR, Figueiredo CME, Ready JS, Salazar C, Linares M, Byers KJRP, Merrill RM, McMillan WO, Schulz S, Jiggins CD. 2020. Species specificity and intraspecific variation in the chemical profiles of *Heliconius* butterflies across a large geographic range. Ecol Evol. Apr 3;10(9):3895-3918. doi: 10.1002/ece3.6079. PMID: 32489619; PMCID: PMC7244815.

de Cara MÁR , Jay P , Rougemont Q, Chouteau M, Whibley A, Huber B, Piron-Prunier F, Rogner-Ramos R, Freitas AVL, Salazar C, Silva-Brandão KL, Texeira-Torres T, Joron T, (20222). Balancing selection at a wing pattern locus is associated with major shifts in genome-wide patterns of diversity and gene flow in a butterfly. bioRxiv 2021.09.29.462348; doi: https://doi.org/10.1101/2021.09.29.462348

Dell'Aglio, D.D., Troscianko, J., McMillan, W.O., Stevens, M. and Jiggins, C.D. (2018), The appearance of mimetic *Heliconius* butterflies to predators and conspecifics. Evolution, 72: 2156-2166.

DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del Angel G, Rivas MA, Hanna M, et al. 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet* 43:491–498.

Dray S, Dufour A-B. 2007. The ade4 Package: Implementing the Duality Diagram for Ecologists. *Journal of Statistical Software* 22:1–20.

Durand EY, Patterson N, Reich D, Slatkin M. 2011. Testing for Ancient Admixture between Closely Related Populations. *Molecular Biology and Evolution* 28:2239–2252.

Edwards AWF. 1972. *Likelihood*. Cambridge, UK: Cambridge University Press.

Emelianov I, Drès M, Baltensweiler W, Mallet J. 2001. Host-induced assortative mating in host races of the larch budmoth. *Evol Int J Org Evol* 55: 2002–2010.

Emsley MG. 1963. A morphological study of imagine Heliconiinae (Lep., Nymphalidae) with a consideration of the evolutionary relationships within the group. *Zool N Y* 48: 85–130.

Estrada C, Jiggins CD. 2008. Interspecific sexual attraction because of convergence in warning colouration: is there a conflict between natural and sexual selection in mimetic species? *J Evol Biol* 21: 749–760.

Estrada C, Schulz S, Yildizhan S, Gilbert LE. 2011. Sexual Selection Drives the Evolution of Antiaphrodisiac Pheromones in Butterflies.

Ewing GB, Jensen JD. 2016. The consequences of not accounting for background selection in demographic inference. *Molecular Ecology* 25:135–141.

Feder JL, Egan SP, Nosil P. 2012. The genomics of speciation-with-gene-flow. *Trends Genet* 28:342–350.

Fenner J, Benson C, Rodriguez-Caro L, Ren A, Papa R, Martin A, Hoffmann F, Range R, Counterman BA. 2020. Wnt Genes in Wing Pattern Development of Coliadinae Butterflies. Frontiers in Ecology and Evolution 8. Available from: https://www.frontiersin.org/articles/10.3389/fevo.2020.00197

Ferveur JF. 2005. Cuticular hydrocarbons: their evolution and roles in *Drosophila* pheromonal communication. *Behav Genet* 35: 279–295.

Flaxman SM, Feder JL, Nosil P. 2013. Genetic Hitchhiking and the Dynamic Buildup of Genomic Divergence During Speciation with Gene Flow. *Evolution* 67:2577–2591.

Gagnaire P-A, Pavey SA, Normandeau E, Bernatchez L. 2013. The genetic architecture of reproductive isolation during speciation-with-gene-flow in lake whitefish species pairs assessed by RAD sequencing. *Evolution* 67:2483–2497.

González-Rojas MF, Darragh K, Robles J, Linares M, Schulz S, McMillan WO, Jiggins CD, Pardo-Diaz C, Salazar C. 2020. Chemical signals act as the main reproductive barrier between sister and mimetic *Heliconius* butterflies. Proc. R. Soc. B 287: 20200587. http://dx.doi.org/10.1098/rspb.2020.0587

Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W, Fritz MH-Y, et al. 2010. A Draft Sequence of the Neandertal Genome. *Science* 328:710–722.

Hausmann A.E., Kuo CY., Freire M, Rueda-M N, Linares M, Pardo-Diaz C, Salazar C. and Merrill RM. 2021, Light environment influences mating behaviours during the early stages of divergence in tropical butterflies. Proc. R. Soc. B.2882021015720210157

Hay-Roe MM, Lamas G, Nation JL. 2007. Pre- and postzygotic isolation and Haldane rule effects in reciprocal crosses of *Danaus erippus* and *Danaus plexippus* (Lepidoptera: Danainae), supported by differentiation of cuticular hydrocarbons, establish their status as separate species. *Biol J Linn Soc* 91: 445–453.

Heuskin S, Vanderplanck M, Bacquet P, Holveck M-J, Kaltenpoth M, Engl T, *et al.* 2014. The composition of cuticular compounds indicates body parts, sex and age in the model butterfly *Bicyclus anynana* (Lepidoptera). *Chem Ecol* 2: 37.

Hewitt G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* 405:907–913.

Hill WG, Robertson A. 1966. The effect of linkage on limits to artificial selection. *Genetics Research* 8:269–294.

Huber B, Whibley A, Poul YL, Navarro N, Martin A, Baxter S, Shah A, Gilles B, Wirth T, McMillan WO, et al. 2015. Conservatism and novelty in the genetic architecture of adaptation in *Heliconius* butterflies. *Heredity* 114:515–524.

Hudson RR. 2002. Generating samples under a Wright–Fisher neutral model of genetic variation. *Bioinformatics* 18:337–338.

Jay P, Whibley A, Frézal L, Rodríguez de Cara MÁ, Nowell RW, Mallet J, Dasmahapatra KK, Joron M. 2018. Supergene Evolution Triggered by the Introgression of a Chromosomal Inversion. *Current Biology* 28:1839-1845.e3.

Jiggins CD, Naisbit RE, Coe RL, Mallet J. 2001. Reproductive isolation caused by colour pattern mimicry. *Nature* 411: 302–305.

Jiggins CD, Estrada C, Rodrigues A. 2004. Mimicry and the evolution of premating isolation in *Heliconius melpomene* Linnaeus. *J Evol Biol* 17: 680–691.

Jurenka RA, Subchev M, Abad J-L, Choi M-Y, Fabrias G. 2003. Sex pheromone biosynthetic pathway for disparlure in the gypsy moth, *Lymantria dispar. Proc Natl Acad Sci U S A* 100: 809–814.

Juric I, Aeschbacher S, Coop G. 2016. The Strength of Selection against Neanderthal Introgression. *PLOS Genetics* 12:e1006340.

Kaplan NL, Hudson RR, Langley CH. 1989. The "hitchhiking effect" revisited. *Genetics* 123:887–899.

Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, Davey JW, Jiggins CD. 2015. Estimation of the spontaneous mutation rate in *Heliconius melpomene*. *Mol Biol Evol* 32:239–243.

Kemp DJ. 2007. Female butterflies prefer males bearing bright iridescent ornamentation. *Proc R Soc B Biol Sci* 274: 1043–1047.

Klein AL, Araújo AM de. 2010. Courtship behavior of *Heliconius erato phyllis* (Lepidoptera, Nymphalidae) towards virgin and mated females: conflict between attraction and repulsion signals? *J Ethol* 28: 409–420.

Kozak KM, Wahlberg N, Neild AFE, Dasmahapatra KK, Mallet J, Jiggins CD. 2015. Multilocus Species Trees Show the Recent Adaptive Radiation of the Mimetic *Heliconius* Butterflies. *Syst Biol* 64: 505–524.

Kronforst MR, Young LG, Blume LM, Gilbert LE. 2006. Multilocus Analyses of Admixture and Introgression Among Hybridizing *Heliconius* Butterflies. *Evolution* 60: 1254–1268.

Kronforst MR, Hansen MEB, Crawford NG, Gallant JR, Zhang W, Kulathinal RJ, et al. 2013. Hybridization Reveals the Evolving Genomic Architecture of Speciation. *Cell Rep* 5: 666–677.

Lecocq T, Vereecken NJ, Michez D, Dellicour S, Lhomme P, Valterová I, et al. 2013. Patterns of Genetic and Reproductive Traits Differentiation in Mainland vs. Corsican Populations of Bumblebees. *PLoS ONE* 8: e65642.

Leroy T, Rousselle M, Tilak M-K, Caizergues AE, Scornavacca C, Recuerda M, Fuchs J, Illera JC, De Swardt DH, Blanco G, et al. 2021. Island songbirds as windows into evolution in small populations. *Current Biology* 31:1303-1310.e4.

Lewis JJ, Geltman RC, Pollak PC, Rondem KE, Van Belleghem SM, Hubisz MJ, Munn PR, Zhang L, Benson C, Mazo-Vargas A, et al. 2019. Parallel evolution of ancient, pleiotropic enhancers underlies butterfly wing pattern mimicry. *Proceedings of the National Academy of Sciences* 116:24174–24183.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv:1303.3997 [q-bio]* [Internet]. Available from: http://arxiv.org/abs/1303.3997

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25:2078–2079.

Livraghi L, Hanly JJ, Van Bellghem SM, Montejo-Kovacevich G, van der Heijden ES, Loh LS, Ren A, Warren IA, Lewis JJ, Concha C, et al. 2021. *cortex* cis-regulatory switches establish scale colour identity and pattern diversity in *Heliconius. eLife 10:e68549*.

Lloyd VJ, Nadeau NJ. 2021. The evolution of structural colour in butterflies. *Current Opinion in Genetics & Development* 69:28–34.

Loudon C, Koehl MA. 2000. Sniffing by a silkworm moth: wing fanning enhances air penetration through and pheromone interception by antennae. *J Exp Biol* 203: 2977–2990.

Mallet J, Gilbert Le Jr. 1995. Why are there so many mimicry rings? Correlations between habitat, behaviour and mimicry in *Heliconius* butterflies. *Biological Journal of the Linnean Society* 55:159–180.

Mallet, J., W. O. McMillan and C. D. Jiggins. (1998) Mimicry and warning color at the boundary between microevolution and macroevolution. In Endless Forms: Species and Speciation, D. Howard and S. Berlocher, eds. Oxford University Press.

Mallet J, Beltrán M, Neukirchen W, Linares M. 2007. Natural hybridization in heliconiinae butterflies: the species boundary as a continuum. *BMC Evol Biol* 7: 28.

Mann F, Vanjari S, Rosser N et al. 2017. The Scent Chemistry of *Heliconius* Wing Androconia. *J Chem Ecol* 43, 843–857. https://doi.org/10.1007/s10886-017-0867-3

Martin A, Papa R, Nadeau NJ, et al. 2012. Diversification of complex butterfly wing patterns by repeated regulatory evolution of a *Wnt* ligand. *PNAS*. Jul;109(31):12632-12637. DOI: 10.1073/pnas.1204800109. PMID: 22802635; PMCID: PMC3411988.

Martin SH, Dasmahapatra KK, Nadeau NJ, Salazar C, Walters JR, Simpson F, Blaxter M, Manica A, Mallet J, Jiggins CD. 2013. Genome-wide evidence for speciation with gene flow in *Heliconius* butterflies. *Genome Res.* 23:1817–1828.

Martin SH, Davey JW, Jiggins CD. 2015. Evaluating the Use of ABBA–BABA Statistics to Locate Introgressed Loci. *Molecular Biology and Evolution* 32:244–257.

Martin SH, Van Belleghem SM. 2017. Exploring Evolutionary Relationships Across the Genome Using Topology Weighting. *Genetics* 206:429–438.

Martin SH, Davey JW, Salazar C, Jiggins CD. 2019. Recombination rate variation shapes barriers to introgression across butterfly genomes. *PLOS Biology* 17:e2006288.

Mavárez J, Salazar CA, Bermingham E, Salcedo C, Jiggins CD, Linares M. 2006. Speciation by hybridization in *Heliconius* butterflies. *Nature* 441: 868–871.

McVean G, Awadalla P, Fearnhead P. 2002. A coalescent-based method for detecting and estimating recombination from gene sequences. *Genetics* 160:1231–1241.

Mega NO, Araújo AM de. 2010. Analysis of the mating behavior and some possible causes of male copulatory success in *Dryas iulia alcionea* (Lepidoptera, Nymphalidae, Heliconiinae). *J Ethol* 28: 123–132.

Melo MC, Salazar C, Jiggins CD, Linares M. 2009. Assortative mating preferences among hybrids offers a route to hybrid speciation. *Evolution* 63: 1660–1665.

Merrill, R.M., Dasmahapatra, K.K., Davey, J.W., Dell'Aglio, D.D., Hanly, J.J., Huber, B., Jiggins, C.D., Joron, M., Kozak, K.M., Llaurens, V., Martin, S.H., Montgomery, S.H., Morris, J., Nadeau, N.J., Pinharanda, A.L., Rosser, N., Thompson, M.J., Vanjari, S., Wallbank, R.W.R. and Yu, Q. (2015), The diversification of Heliconius butterflies: what have we learned in 150 years?. J. Evol. Biol., 28: 1417-1438.

Mérot C, Mavárez J, Evin A, Dasmahapatra KK, Mallet J, Lamas G, *et al.* 2013. Genetic differentiation without mimicry shift in a pair of hybridizing *Heliconius* species (Lepidoptera: Nymphalidae). *Biol J Linn Soc* 109: 830–847.

Mérot C, Frérot B, Leppik E, Joron M. Beyond magic traits: Multimodal mating cues in *Heliconius* butterflies. 2015. *Evolution* Nov; 69(11):2891-904. doi: 10.1111/evo.12789. Epub 2015 Oct 29. PMID: 26513426.

Merrill RM, Gompert Z, Dembeck LM, Kronforst MR, McMillan WO, Jiggins CD. 2011. Mate preference across the speciation continuum in a clade of mimetic butterflies. *Evol Int J Org Evol* 65: 1489–1500.

Mérot C, Salazar C, Merrill RM, Jiggins CD, Joron M. 2017. What shapes the continuum of reproductive isolation? Lessons from *Heliconius* butterflies. Proc. R. Soc. B 284: 20170335.

Millar JG. 2000. Polyene hydrocarbons and epoxides: A Second Major Class of Lepidopteran Sex Attractant Pheromones. *Annu Rev Entomol* 45: 575–604.

Moest M, Belleghem SMV, James JE, Salazar C, Martin SH, Barker SL, Moreira GRP, Mérot C, Joron M, Nadeau NJ, et al. 2020. Selective sweeps on novel and introgressed variation shape mimicry loci in a butterfly adaptive radiation. *PLOS Biology* 18:e3000597.

Morris J, Hanly JJ, Martin SH, Van Belleghem SM, Salazar C, Jiggins CD, Dasmahapatra KK. 2020. Deep Convergence, Shared Ancestry, and Evolutionary Novelty in the Genetic Architecture of Heliconius Mimicry. *Genetics* 216:765–780.

Nadeau NJ, Pardo-Diaz C, Whibley A, et al. 2016. The gene *cortex* controls mimicry and crypsis in butterflies and moths. *Nature*. Jun;534(7605):106-110. DOI: 10.1038/nature17961. PMID: 27251285; PMCID: PMC5094491.

Nieberding CM, de Vos H, Schneider MV, Lassance J-M, Estramil N, Andersson J, et al. 2008. The Male Sex Pheromone of the Butterfly *Bicyclus anynana*: Towards an Evolutionary Analysis. *PLoS ONE* 3: e2751.

NIST Chemistry WebBook. 2015. NIST Standard Reference Database Number 69 at <http://webbook.nist.gov>

Noor M a. F, Bennett SM. 2009. Islands of speciation or mirages in the desert? Examining the role of restricted recombination in maintaining species. *Heredity (Edinb)* 103:439–444.

Nordborg M, Charlesworth B, Charlesworth D. 1996. The effect of recombination on background selection. *Genetics Research* 67:159–174.

Obara Y. 1964. Mating behavior of the cabbage butterfly, *Pieris rapae crucivora*. II. The 'mate-refusal posture' of the female. *Zool Mag Dobutsugaku-Zasshi* 73: 175–178.

Oksanen FJ, et al. 2017. Vegan: Community Ecology Package. R package Version 2.4-3. https://CRAN.R-project.org/package=vegan

Pardo-Diaz C, Salazar C, Baxter SW, Merot C, Figueiredo-Ready W, Joron M, *et al.* 2012. Adaptive Introgression across Species Boundaries in *Heliconius* Butterflies. *PLoS Genet* 8: e1002752.

Payseur BA, Nachman MW. 2002. Natural selection at linked sites in humans. *Gene* 300:31–42.

Pease JB, Hahn MW. 2015. Detection and Polarization of Introgression in a Five-Taxon Phylogeny. *Systematic Biology* 64:651–662.

Pickrell JK, Pritchard JK. 2012. Inference of Population Splits and Mixtures from Genome-Wide Allele Frequency Data. *PLOS Genetics* 8:e1002967.

Pliske TE. 1975. Courtship Behavior of the Monarch Butterfly, *Danaus plexippus* L. *Ann Entomol Soc Am* 68: 143–151.

Pudlo P, Marin J-M, Estoup A, Cornuet J-M, Gautier M, Robert CP. 2016. Reliable ABC model choice via random forests. *Bioinformatics* 32:859–866.

Ravinet M, Faria R, Butlin RK, Galindo J, Bierne N, Rafajlović M, Noor M a. F, Mehlig B, Westram AM. 2017. Interpreting the genomic landscape of speciation: a road map for finding barriers to gene flow. *Journal of Evolutionary Biology* 30:1450–1477.

Reed RD, Papa R, Martin A, Hines HM, Counterman BA, Pardo-Diaz C, Jiggins CD, Chamberlain NL, Kronforst MR, Chen R, Halder G, Nijhout HF, McMillan WO. 2011. *optix* drives the repeated convergent evolution of butterfly wing pattern mimicry. *Science.* Aug 26;333(6046):1137-41. doi: 10.1126/science.1208227. Epub 2011 Jul 21. PMID: 21778360.

Robert CP, Cornuet J-M, Marin J-M, Pillai NS. 2011. Lack of confidence in approximate Bayesian computation model choice. *Proceedings of the National Academy of Sciences* 108:15112–15117.

Roberts, D.W. 2012. Ordination and Multivariate Analysis for Ecology. R Package Version 1.0-5. http://CRAN.R-project.org/package=labdsv

Ross-Ibarra J, Wright SI, Foxe JP, Kawabe A, DeRose-Wilson L, Gos G, Charlesworth D, Gaut BS. 2008. Patterns of Polymorphism and Demographic History in Natural Populations of *Arabidopsis lyrata*. *PLOS ONE* 3:e2411.

Ross-Ibarra J, Tenaillon M, Gaut BS. 2009. Historical Divergence and Gene Flow in the Genus *Zea*. *Genetics* 181:1399–1413.

Rosser, N., Kozak, K.M., Phillimore, A.B. *et al.* Extensive range overlap between heliconiine sister species: evidence for sympatric speciation in butterflies?. *BMC Evol Biol* **15**, 125 (2015). https://doi.org/10.1186/s12862-015-0420-3.

Rougemont Q, Bernatchez L. 2018. The demographic history of Atlantic salmon (*Salmo salar*) across its distribution range reconstructed from approximate Bayesian computations. *Evolution* 72:1261–1277.

Rougemont Q, Roux C, Neuenschwander S, Goudet J, Launey S, Evanno G. 2016. Reconstructing the demographic history of divergence between European river and brook lampreys using approximate Bayesian computations. *PeerJ* 4:e1910.

Rougeux C, Bernatchez L, Gagnaire P-A. 2017. Modeling the Multiple Facets of Speciation-with-Gene-Flow toward Inferring the Divergence History of Lake Whitefish Species Pairs (*Coregonus clupeaformis*). *Genome Biol Evol* 9:2057–2074.

Roux C, Castric V, Pauwels M, Wright SI, Saumitou-Laprade P, Vekemans X. 2011. Does Speciation between Arabidopsis halleri and *Arabidopsis lyrata* Coincide with Major Changes in a Molecular Target of Adaptation? *PLOS ONE* 6:e26872.

Roux C, Fraïsse C, Romiguier J, Anciaux Y, Galtier N, Bierne N. 2016. Shedding Light on the Grey Zone of Speciation along a Continuum of Genomic Divergence. *PLOS Biology* 14:e2000234.

Roux C, Tsagkogeorga G, Bierne N, Galtier N. 2013. Crossing the Species Barrier: Genomic Hotspots of Introgression between Two Highly Divergent *Ciona intestinalis* Species. *Molecular Biology and Evolution* 30:1574–1587.

Rutowski RL, Schaefer J. 1984. Courtship behaviour of the gulf fritillary *Agraulis vanillae* Nymphalidae. *J Lepidopterists Soc* 38: 23–31.

Rutowski RL. 1991. The evolution of male mate-locating behavior in butterflies. *Am Nat*: 1121–1139.

Schrider DR, Shanku AG, Kern AD. 2016. Effects of Linked Selective Sweeps on Demographic Inference and Model Selection. *Genetics* 204:1207–1223.

Schulz S, Boppré M, Vane-Wright Ri. 1993. Specific mixtures of secretions from male scent organs of African milkweed butterflies (Danainae). *Philos Trans R Soc B Biol Sci* 342: 161–181.

Schulz S, Beccaloni G, Nishida R, Roisin Y & Mcneil, JN. 1998. 2,5 dialkyltetrahydrofurans, common component of the cuticular lipids of Lepidoptera. Zeitschrift fur Naturforsch. 53c, 107–116.

Schulz S, Estrada C, Yildizhan S, Boppré M, Gilbert LE. 2008. An antiaphrodisiac in *Heliconius melpomene* butterflies. *J Chem Ecol* 34: 82–93.

Scott J. 1972. Mating of butterflies. *J Res Lepidoptera* 11: 99–127.

Sheppard PM, Turner JRG, Brown KS, Benson WW, Singer MC. 1985. Genetics and the evolution of muellerian mimicry in *Heliconius* butterflies. *Phil. Trans. R. Soc. Lond. B*308433–610.

Smith JM, Haigh J. 1974. The hitch-hiking effect of a favourable gene. Genetics Research 23:23–35.

Stankowski S, Chase MA, Fuiten AM, Rodrigues MF, Ralph PL, Streisfeld MA. 2019. Widespread selection and gene flow shape the genomic landscape during a radiation of monkeyflowers. *PLOS Biology* 17:e3000391.

Tinbergen N, Meeuse BJD, Varossieau LKB, Boerema uWW. 1942. Die Balz des Samtfalters, *Eumenis* (=*Satyrus*) semele (L.). *Z Für Tierpsychol* 5: 182–226.

Thomas Y, Bethenod MT, Pelozuelo L, Frérot B, Bourguet D. 2003. Genetic Isolation Between Two Sympatric Host-Plant Races of the European Corn Borer, *Ostrinia nubilalis* Hubner I. Sex Pheromone, Moth Emergence Timing, and Parasitism. *Evolution* 57: 261–273.

Van Belleghem SM, Alicea Roman PA, Carbia Gutierrez H, Counterman BA, Papa R. 2020. Perfect mimicry between *Heliconius* butterflies is constrained by genetics and development. *Proceedings of the Royal Society B*: Biological Sciences 287:20201267.

Van Belleghem SM, Cole JM, Montejo-Kovacevich G, Bacquet CN, McMillan WO, Papa R, Counterman BA. 2021. Selection and isolation define a heterogeneous divergence landscape between hybridizing *Heliconius* butterflies. Evolution 75:2251–2268.

Vane-Wright RI, Boppré M. 1993. Visual and Chemical Signalling in Butterflies: Functional and Phylogenetic Perspectives. *Phil Trans R Soc Lond B* 340: 197–205.

Via S, Bouck AC, Skillman S. 2000. Reproductive isolation between divergent races of pea aphids on two hosts. II. Selection against migrants and hybrids in the parental environments. *Evol Int J Org Evol* 54: 1626–1637.

Via S, West J. 2008. The genetic mosaic suggests a new role for hitchhiking in ecological speciation. *Molecular Ecology* 17:4334–4345.

Westerman EL, VanKuren NW, Massardo D, Tenger-Trolander A, Zhang W, Hill RI, Perry M, Bayala E, Barr K, Chamberlain N, et al. 2018. *Aristaless* Controls Butterfly Wing Color Variation Used in Mimicry and Mate Choice. *Current Biology* 28:3469-3474.e4.

Wood TK, Keese MC. 1990. Host-Plant-Induced Assortative Mating in *Enchenopa* Treehoppers. *Evolution* 44: 619–628.

Wu C-I. 2001. The genic view of the process of speciation. *Journal of Evolutionary Biology* 14:851–865.

Zhang L, Mazo-Vargas A, Reed RD. 2017. Single master regulatory gene coordinates the evolution and development of butterfly color and iridescence. *Proceedings of the National Academy of Sciences* 114:10707–10712.