1   **Title:** Greengenes2 enables a shared data universe for microbiome studies
2

3   **Authors:** Daniel McDonald[1], Yueyu Jiang[2], Metin Balaban[3], Kalen Cantrell[4], Qiyun Zhu[5,6],
4   Antonio Gonzalez[1], James T. Morton[7], Giorgia Nicolaou[8], Donovan H. Parks[9], Søren Karst[10],
5   Mads Albertsen[11], Phil Hugenholtz[9], Todd DeSantis[12], Siavash Mirarab[2], Rob Knight[1,4,13,14,*]
6
Affiliations:
[1] Department of Pediatrics, UC San Diego School of Medicine, La Jolla, CA, USA
[2] Department of Electrical and Computer Engineering, UC San Diego, La Jolla, CA, USA
[3] Bioinformatics and System Biology Program, UC San Diego, La Jolla, CA, USA
[4] Department of Computer Science and Engineering, UC San Diego, La Jolla, CA, USA
[5] School of Life Sciences, Arizona State University, Tempe, AZ, USA
[6] Biodesign Center for Fundamental and Applied Microbiomics, Arizona State University,
Tempe, AZ, USA
[7] Biostatistics & Bioinformatics Branch, Eunice Kennedy Shriver National Institute of Child
Health and Human Development, National Institutes of Health, Bethesda, MD, USA
[8] Halicioglu Data Science Institute, UC San Diego, La Jolla, CA, USA
[9] Australian Centre for Ecogenomics, School of Chemistry and Molecular Biosciences, The
University of Queensland, St Lucia, QLD 4072, Australia
[10] Center for Microbial Communities, Aalborg University, Aalborg, Denmark
[11] Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark
[12] Department of Informatics, Second Genome, Brisbane, CA, USA
[13] Center for Microbiome Innovation, Jacobs School of Engineering, UC San Diego, La Jolla,
CA, USA
[14] Department of Bioengineering, UC San Diego, La Jolla, CA, USA
[*] Corresponding author
27
28   Address correspondence to Rob Knight (robknight@eng.ucsd.edu)
29

Abstract:

16S rRNA and shotgun metagenomics studies typically yield different results, traditionally

thought to be due to biases in amplification. We show that differences in reference phylogeny

are more important. By inserting sequences into a whole-genome phylogeny, we show that 16S

rRNA and shotgun metagenomic data generated from the same samples agree in principal

coordinates space, taxonomy, and in phenotype effect size when analyzed with the same tree.

Body:

Shotgun metagenomics and 16S rRNA gene amplicon (16S) studies are widely used in

microbiome research, but investigators using different methods typically find their results hard to

reconcile. This lack of standardization across methods limits the utility of the microbiome for

reproducible biomarker discovery.

A key problem is that whole-genome resources and rRNA resources depend on different

taxonomies and phylogenies. For example, Web of Life (WoL) [1] and the Genome Taxonomy

Database (GTDB) [2] provide whole-genome trees that cover only a small fraction of known

bacteria and archaea, while SILVA [3] and Greengenes [4] are more comprehensive but not fully

linked to genome records.

We reasoned that an iterative approach could yield a massive reference tree that unifies these

different data layers. We began with a whole-genome catalog of 15,953 bacterial and archeal

genomes evenly sampled from NCBI, and reconstructed an accurate phylogenomic tree by

summarizing evolutionary trajectories of 380 global marker genes using the new workflow

uDance. This work, namely Web of Life version 2 (WoL2), represents a significant upgrade from

the previously released WoL1 (10,575 genomes) [1]. Then, we added 18,356 full-length rRNA

amplicons from the Living Tree Project January 2022 release [5] and 1,725,274 near-complete

16S rRNA genes from Karst et al. [6] and the EMP500 [7] with uDance v1.1.0, then added all full-

length 16S sequences from GTDB r207, and finally inserted 23,113,447 short V4 16S rRNA

Deblur v1.1.0 [8] amplicon sequence variants from Qiita (retrieved Dec. 14, 2021) [9] as well as

mitochondria and chloroplast 16S from SILVA v138 using DEPP v0.3 [10], including everything

from the Earth Microbiome Project [11] and American Gut Project/Microsetta [12] (Fig. 1A). Our use

of uDance ensured the genome-based relationships are kept fixed and relationships between

full-length 16S sequences are inferred. For short fragments, we kept genome and full length

relationships fixed and inserted fragments independently from each other. Following

deduplication and quality control on fragment placement, this yielded a tree covering 21,074,442

sequences from 31 different Earth Microbiome Project Ontology (EMPO) EMPO_3

environments, of which 46.5% of species-level leaves were covered by a complete genome.

Taxonomic labels were decorated onto the phylogeny using tax2tree v1.1 [4]. The input taxonomy

for decoration used GTDB r207, combined with the Living Tree Project January 2022 release.

Taxonomy was harmonized prioritizing GTDB including preserving the polyphyletic labelings of

GTDB (see also Online Methods). Taxonomy will be updated every six months using the latest

versions of GTDB and LTP.

Our resource is much larger than past resources in its phylogenetic coverage, as compared to

the last release of Greengenes (Fig. 1B), SILVA (Fig. S1A) or GTDB (Fig. S1B). However,

because our amplicon library is linked to environments labeled with Earth Microbiome Project

Ontology (EMPO) categories, we can easily identify the environments that contain samples that

can fill out the tree. Because MAG assembly efforts can only cover abundant taxa, we plotted

for each EMPO category the amount of new branch length added to the tree by taxa whose

minimum abundance is 1% in each sample (Fig. 1C). The results show which environment

types on average will best yield new metagenome assembled genomes (MAGs), and also show

which environments harbor individual samples that will have a large impact when sequenced.

Past efforts to reconcile 16S and shotgun datasets have led to non-overlapping distributions and

only techniques such as Procrustes analysis can even show relationships between the results

[13]. On two large human stool cohorts [12,14] where both 16S and shotgun data were generated on

the same samples, we find that Bray-Curtis [15] (non-phylogenetic) ordination fails to reconcile at

the feature level (Fig. 1D) and is poor at the genus level (Fig. 1E, S1C). However, UniFrac [16], a

phylogenetic method, used with our Greengenes2 tree provides far better concordance (Fig. 1F,

S1D).

Figure 1. (A) The Greengenes2 phylogeny rendered using Empress [17] with amplicon sequence

variant multifurcations collapsed, tip color indicating representation in the American Gut Project

(AGP), the Earth Microbiome Project (EMP), both or neither, and with the top 20 represented

phyla depicted in the outer bar. (B) The same collapsed phylogeny, colored by the presence or

absence of a best BLAST [18] hit from Greengenes 13_8 99% OTUs. The bar depicts the same

coloring as the tips. (C) Earth Microbiome Project samples and the amount of novel branch

length, normalized by the total backbone branch length, added to the tree through amplicon

sequence variant fragment placement. (D) Bray Curtis applied to paired 16S V4 rRNA amplicon

sequence variants and whole genome shotgun samples from The Healthy Microbiome Diet

 Initiative subset of The Microsetta Initiative. (E) Same data as (D) but computing Bray Curtis on

 genus collapsed data. (F) Same data as (D-E) but using weighted UniFrac.

We also find that the per-sample shotgun and 16S taxonomy concordances are excellent even

to the species level. We first computed taxonomy profiles for shotgun data using the Woltka

pipeline [19]. Using a Naive Bayes classifier from q2-feature-classifier v2022.2 [20] to compare

GTDB r207 taxonomy results at each level against SILVA v138 (Fig. 2A) or Greengenes v13_8

(Fig. 2B), no species-level reconciliation was possible. In contrast, Greengenes2 provided

excellent concordance at the genus level (Pearson r=0.85) and good concordance at the

species level (Pearson r=0.65) (Fig. 2C). Interestingly, the tree is now sufficiently complete that

exact matching of 16S ASVs followed by reading the taxonomy off the tree performs even better

than the Naive Bayes Classifier (Naive Bayes; Pearson r=0.54 at species, r=0.84 at genus).

Finally, a critical reason to assign taxonomy is downstream use of biomarkers and indicator

taxa. Microbiome science has been described as having a reproducibility crisis [21], but much of

this problem stems from incompatible methods [22]. We initially used the The Human Diet

Microbiome Initiative (THDMI) dataset, which is a multipopulation expansion of The Microsetta

Initiative [12] that contains samples with paired 16S and shotgun preparations, to test whether a

harmonized resource would provide concordant rankings for the variables that affect the human

microbiome similarly. Using Greengenes2, the concordance was good with Bray-Curtis (Fig. 2D;

Pearson r^2=0.56), better using UniFrac with different phylogenies (SILVA 138 and

Greengenes2; Fig S1E; Pearson r^2=0.77), and excellent with UniFrac on the same phylogeny

(Fig. 2E; Pearson r^2=0.87). We confirmed these results with an additional cohort [14] (Fig. S1FG).

Intriguingly, the ranked effect sizes across different cohorts were concordant.

Figure 2. (A-C) Per sample taxonomy comparisons between 16S and whole genome shotgun

profiles from THDMI. The solid bar depicts the 50th percentile, the dashed lines are 25th and

75th percentiles. (A) 16S taxonomy assessed with SILVA 138 using the default q2-feature-

classifier Naive Bayes model. (B) 16S taxonomy assessment with Greengenes 13_8 using the

default q2-feature-classifier Naive Bayes model. (C) 16S taxonomy assessment performed by

reading the lineages directly from the phylogeny or through Naive Bayes trained on the V4

132   *regions of the Greengenes2 backbone. (D-E) Effect size calculations performed with Evident on*

133   *paired 16S and whole genome shotgun samples from THDMI. Calculations performed at*

134   *maximal resolution, using ASVs for 16S and genome identifiers for shotgun. (D) Bray Curtis*

135   *distances. (E) Weighted normalized UniFrac.*

Taken together, these results show that use of a consistent, integrated taxonomic resource

dramatically improves the reproducibility of microbiome studies using different data types, and

allows variables of large versus small effect to be reliably recovered in different populations.

ONLINE METHODS

Phylogeny construction

Web of Life version 2 [1] (a tree inferred using genome-wide data) was used as the starting

backbone. Full length 16S sequences from the Living Tree Project [5], full length mitochondria

and chloroplast from SILVA 138 [3], full length 16S from GTDB r207 [2], full length 16S from Karst

et al [6], and full length 16S from the EMP 500 [7] (samples selected and sequenced specifically for

Greengenes2) were collected and deduplicated. Sequences were then aligned using UPP [23]

and gappy sequences with less than 1000bp were removed. The resulting set of 321,210

unique sequences were used with uDance v1.1.0 to update the Web of Life 2 (WoL2) backbone.

Briefly, uDance updates an existing tree with new sequences and (unlike placement methods)

also infers the relationship of existing sequences. uDance has two modes: one that allows

updates to the backbone and one that keeps the backbone fixed. In our analyses, we kept the

backbone tree (inferred using genomic data) fixed. To extend the genomic tree with 16S data,

we identified 13,249 genomes in the WoL2 backbone tree with at least one 16S copy and used

them to train a DEPP model with the weighted average method detailed below to handle

multiple copies. We then used DEPP to insert all 16S copies of all genomes into the backbone

and measured the distance between the genome position and the 16S position. We removed

copies that were placed far further than others, as identified using a 2-means approach with

centroids equals to at least 13 branches. We repeated this process a second round. Then, for

every remaining genome, we selected as its representative the copy with the minimum

placement error and computing the consensus when there were ties. At the end, we are left with

12,344 unique 16S sequences across all the WoL2 genomes. For tree inference, uDance used

IQ-TREE2 [24] in fast tree search with model GTR+Γ after removing duplicate sequences.

Next, we collected 16S V4 ASVs from Qiita [9] using redbiom [25] (query performed December 14,

2021) from contexts "Deblur_2021.09-Illumina-16S-V4-90nt-dd6875", "Deblur_2021.09-Illumina-

16S-V4-100nt-50b3a2", "Deblur_2021.09-Illumina-16S-V4-125nt-92f954", "Deblur_2021.09-

Illumina-16S-V4-150nt-ac8c0b", "Deblur_2021.09-Illumina-16S-V4-200nt-0b8b48",

"Deblur_2021.09-Illumina-16S-V4-250nt-8b2bff" and aligned them to the existing 16S alignment

of sequences in WoL2 using UPP, setting the maximum alignment subset size to 200 (to help

with scalability). The collected 16S V4 ASVs are aligned to the V4 region of the existing

"backbone" alignments. A DEPP model was then trained on the full length 16S sequences from

the backbone. DEPP constructs a Neural network model that embeds sequences in high

dimensional spaces such that embedded points resemble the phylogeny in their distances.

Such a model then allows insertion of new sequences into a tree using distance-based

phylogenetic insertion method APPLES-2 [26]. The ASVs from redbiom were then inserted into

the backbone using the trained DEPP model. To enable analyses of large datasets, we used a

clustering approach with DEPP: we trained an ensemble of DEPP models corresponding to

different parts of the tree and used a classifier to detect the correct subtree. During training, for

species with multiple 16S, all the copies are mapped to the same leaf in the backbone tree. To

train the DEPP models with multiple sequences mapped to a leaf, each site in the sequences is

encoded as a probability vector of four nucleotides across all the copies.

Integrating the GTDB and Living Tree Project taxonomies

GTDB and Living Tree Project are not directly compatible due to differences in their curation. As

a result, it is not always possible to map a species from one resource to the other, either

because parts of a species lineage are not present, are described using different names, or

have an ambiguous association due to polyphyletic taxa in GTDB. GTDB is actively curated,

while LTP generally uses the NCBI taxonomy. To account for these differences, we first mapped

any species that had a perfect species name association and revised its ancestral lineage to

match GTDB. Next, we generated lineage rewrite rules using the GTDB record metadata.

Specifically, we limited the metadata to records which are GTDB representatives and NCBI type

material, and then defined a lineage renaming from the recorded NCBI taxonomy to the GTDB

taxonomy. These rewrite rules were applied from most to least specific taxa, and through this

mechanism we could revise much of the higher ranks of LTP. We then identified incertae sedis

records in LTP which we could not map, removed their lineage strings and did not attempt to

provide taxonomy for them, instead opting to rely on downstream taxonomy decoration to

resolve their lineages. Next, any record which was ambiguous to map was split into a secondary

taxonomy for use in backfilling in the downstream taxonomy decoration. Finally, we

instrumented numerous consistency checks in the taxonomy through the process to capture

inconsistent parents in the taxonomy hierarchy, consistent numbers of ranks in a lineage and

ensuring the resulting taxonomy was a strict hierarchy.

Taxonomy decoration

The original tax2tree algorithm was not well suited for a large volume of species level records in

the backbone, as the algorithm requires an internal node to place a name. If two species are

siblings, the tree would lack a node to contain the species label for both taxa. To account for

this, we updated the algorithm to insert "placeholder" nodes with zero branch length as the

parents of backbone records, which could accept these species labels. We further updated

tax2tree to operate directly on .jplace data [27], preserving edge numbering of the original edges

prior to adding "placeholder" nodes. To support LTP records which could not be integrated into

GTDB, we instrumented a secondary taxonomy mode for tax2tree. Specifically, following the

standard decoration, backfilling and name promotion procedures, we determine on a per record

basis for the secondary taxonomy what portion of the lineage is missing, and place the missing

labels on the placeholder node. We then issue a second round of name promotion using the

existing tax2tree methods.

The actual taxonomy decoration occurs on the backbone tree, which contains only full length

16S records, and does not contain the amplicon sequence variants (ASV). This is done as ASV

placements are independent, do not modify the backbone, and would substantially increase the

computational resources required. After the backbone is decorated, fragment placements from

DEPP are resolved using a multifurcation strategy using the balanced-parentheses library

(https://github.com/biocore/improved-octo-waddle/).

Phylogenetic collapse for visualization

We are unaware of phylogenetic visualization software that can display a tree with over

20,000,000 tips. To produce the visualizations in figure 1, we reduced the dimension of the tree

by collapsing fragment multifurcations to single nodes, dropping the tree to 522,849 tips.

MAG target environments

A feature table for the 27,015 16S rRNA V4 90nt Earth Microbiome Project samples was

obtained from redbiom. The amplicon sequence variants (ASV) were filtered to the overlap of

ASVs present in Greengenes2. Any feature with < 1% relative abundance within a sample was

removed. The feature table was then rarefied to 1,000 sequences per sample. The amount of

novel branch length was then computed, per sample, by summing the branch length of each

ASV's placement edge. The per sample branch length was then normalized by the total tree

branch length (excluding length contributed by ASVs).

Per sample taxonomy correlations

All comparisons used the THDMI 16S and Woltka processed shotgun data. These data were

accessed from Qiita study 10317, and filtered the set of features which overlap with

Greengenes2 using the QIIME 2[28] q2-greengenes2 plugin. 16S taxonomy was assessed using

either a traditional Naive Bayes classifier with q2-feature-classifier and default references from

QIIME 2 2022.2, or by reading the lineage directly from the phylogeny. To help improve

correlation between SILVA and Greengenes2, and Greengenes and Greengenes2, we stripped

polyphyletic labelings from those data; we did not strip polyphyletic labels from the phylogenetic

taxonomy comparison or the Greengenes2 16S vs. Greengenes2 WGS Naive Bayes

comparison. Shotgun taxonomy was determined by the specific observed genome records.

Once the 16S taxonomy was assigned, those tables as well as the WGS Woltka WoL version 2

table were collapsed at the species, genus, family, order, and class levels. We then computed a

minimum relative abundance per sample in the THDMI dataset. In each sample, we removed

any feature, either 16S or WGS, below the per sample minimum (i.e., max(min(16S),

min(WGS))), forming a common minimal basis for taxonomy comparison. Following filtering,

Pearson correlation was computed per sample using SciPy[29]. These correlations were

aggregated per 16S taxonomy assignment method, and by each taxonomic rank. The 25th, 50th

and 75th percentiles were then plotted with Matplotlib[30].

Principal coordinates

THDMI Deblur 16S and Woltka processed shotgun sequencing data, against WoL version 2,

were obtained from Qiita study 10317. Both feature tables were filtered against Greengenes2

2022.10, removing any feature not present in the tree. For the genus collapsed plot (figure 1e),

both the 16S and WGS data features were collapsed using the same taxonomy. For all three

figures, the 16S data were subsampled, with replacement, to 10,000 sequences per sample.

The WGS data were subsampled, with replacement, to 1,000,000 sequences per sample. Bray

Curtis and Weighted UniFrac, and PCoA were computed using q2-diversity 2022.2. The

resulting coordinates were visualized with q2-emperor [31].

Effect size calculations

Similar to principal coordinates, the THDMI data were rarefied to 9,000 and 2,000,000

sequences per sample for 16S and WGS respectively. Bray Curtis and weighted normalized

UniFrac were computed on both sets of data. The variables for THDMI were subset to those

with at least two category values having more than 50 samples. For UniFrac with SILVA, figure

S1E, we performed fragment insertion using q2-fragment-insertion [32] into the standard QIIME 2

SILVA reference, followed by rarefaction to 9,000 sequences per sample, and then computed

weighted normalized UniFrac.

For FinRISK, the data were rarefied to 1,000 and 500,000 sequences per sample for 16S and

WGS. A different depth was used to account for the overall lower amount of sequencing data for

FinRISK. As with THDMI, the variables selected were reduced to those with at least two

category values having more than 50 samples.

Support for computing paired effect sizes is part of the QIIME2 Greengenes2 plugin, q2-

greengenes2, which performs effect size calculations using Evident

(https://github.com/biocore/evident/).

Data access

The official location of the Greengenes2 releases is http://ftp.microbio.me/greengenes_release/.

The data are released under a BSD-3 clause license. A QIIME 2 plugin is available to facilitate

use with the resource, which can be obtained from https://github.com/biocore/q2-greengenes2/.

Taxonomy construction, decoration, and release processing is part of

https://github.com/biocore/greengenes2. uDance release v1.1.0 is available at GitHub:

https://github.com/balabanmetin/uDance. Phylogeny insertion using DEPP is available at

https://github.com/yueyujiang/DEPP; the trained model accessioned with Zenodo at

10.5281/zenodo.7416684. The THDMI data are part of Qiita study 10317, and EBI accession

PRJEB11419. The FinRISK data are available under EGAD00001007035. Finally, an interactive

website to explore the Greengenes2 data is available at https://greengenes2.ucsd.edu.

Acknowledgements

This work was supported in part by NSF XSEDE BIO210103, NSF RAPID 20385.09, NIH

1R35GM14272, NIH U19AG063744, NIH U24DK131617, NIH DP1-AT010885 and Emerald

Foundation 3022. JTM was funded by the intramural research program of the Eunice Kennedy

Shriver National Institute of Child Health and Human Development (NICHD).

Figure S1. (A) Best BLAST hit for SILVA 138 against Greengenes2. (B) Best BLAST hit for

GTDB r207 SSU sequences against Greengenes2. (C) The FinRISK 16S and WGS data

combined, collapsed to genus, with Bray Curtis computed followed by Principal Coordinates

Analysis, colored by technical preparation. (D) The same data as (C) but using weighted

UniFrac. (E) Effect sizes of the THDMI data using the SILVA 138 phylogeny for 16S data, and

the Greengenes2 phylogeny for WGS data. (F) Effect sizes of the FinRISK data using Bray

Curtis. (G) The same data as (E) but using Weighted UniFrac.

REFERENCES

1. Zhu, Q. *et al.* Phylogenomics of 10,575 genomes reveals evolutionary proximity between domains Bacteria and Archaea. *Nat. Commun.* **10**, 5477 (2019).

2. Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Res.* **50**, D785–D794 (2022).

3. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* **41**, D590–6 (2013).

4. McDonald, D. *et al.* An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J.* **6**, 610–618 (2012).

5. Ludwig, W. *et al.* Release LTP_12_2020, featuring a new ARB alignment and improved 16S rRNA tree for prokaryotic type strains. *Syst. Appl. Microbiol.* **44**, 126218 (2021).

6. Karst, S. M. *et al.* High-accuracy long-read amplicon sequences using unique molecular identifiers with Nanopore or PacBio sequencing. *Nat. Methods* **18**, 165–169 (2021).

7. Shaffer, J. P. *et al.* Standardized multi-omics of Earth's microbiomes reveals microbial and metabolite diversity. *Nat Microbiol* **7**, 2128–2150 (2022).

8. Amir, A. *et al.* Deblur Rapidly Resolves Single-Nucleotide Community Sequence Patterns. *mSystems* **2**, (2017).

9. Gonzalez, A. *et al.* Qiita: rapid, web-enabled microbiome meta-analysis. *Nat. Methods* **15**, 796–798 (2018).

10. Jiang, Y., Balaban, M., Zhu, Q. & Mirarab, S. DEPP: Deep learning enables extending species trees using single genes. *bioRxiv* (2021) doi:10.1101/2021.01.22.427808.

11. Thompson, L. R. *et al.* A communal catalogue reveals Earth's multiscale microbial diversity. *Nature* **551**, 457–463 (2017).

12. McDonald, D. *et al.* American Gut: an Open Platform for Citizen Science Microbiome

Research. *mSystems* **3**, (2018).

13. Human Microbiome Project Consortium. Structure, function and diversity of the healthy

human microbiome. *Nature* **486**, 207–214 (2012).

14. Salosensaari, A. *et al.* Taxonomic signatures of cause-specific mortality risk in human gut

microbiome. *Nat. Commun.* **12**, 2671 (2021).

15. Bray, J. R. & Curtis, J. T. An ordination of the upland forest communities of southern

Wisconsin. *Ecol. Monogr.* **27**, 325–349 (1957).

16. Sfiligoi, I., Armstrong, G., Gonzalez, A., McDonald, D. & Knight, R. Optimizing UniFrac with

OpenACC Yields Greater Than One Thousand Times Speed Increase. *mSystems* **7**,

e0002822 (2022).

17. Cantrell, K. *et al.* EMPress Enables Tree-Guided, Interactive, and Exploratory Analyses of

Multi-omic Data Sets. *mSystems* **6**, (2021).

18. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment

search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

19. Zhu, Q. *et al.* Phylogeny-Aware Analysis of Metagenome Community Ecology Based on

Matched Reference Genomes while Bypassing Taxonomy. *mSystems* **7**, e0016722 (2022).

20. Bokulich, N. A. *et al.* Optimizing taxonomic classification of marker-gene amplicon

sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* **6**, 90 (2018).

21. Schloss, P. D. Identifying and Overcoming Threats to Reproducibility, Replicability,

Robustness, and Generalizability in Microbiome Research. *MBio* **9**, (2018).

22. Sinha, R. *et al.* Assessment of variation in microbial community amplicon sequencing by

the Microbiome Quality Control (MBQC) project consortium. *Nat. Biotechnol.* **35**, 1077–

1086 (2017).

23. Nguyen, N.-P. D., Mirarab, S., Kumar, K. & Warnow, T. Ultra-large alignments using

phylogeny-aware profiles. *Genome Biol.* **16**, 124 (2015).

24. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference
in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

25. McDonald, D. *et al.* redbiom: a Rapid Sample Discovery and Feature Characterization
System. *mSystems* **4**, (2019).

26. Balaban, M., Jiang, Y., Roush, D., Zhu, Q. & Mirarab, S. Fast and accurate distance-based
phylogenetic placement using divide and conquer. *Mol. Ecol. Resour.* **22**, 1213–1227
(2022).

27. Matsen, F. A., Hoffman, N. G., Gallagher, A. & Stamatakis, A. A format for phylogenetic
placements. *PLoS One* **7**, e31009 (2012).

28. Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data
science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).

29. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat.*
Methods **17**, 261–272 (2020).

30. Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Comput. Sci. Eng.* **9**, 90–95 (2007).

31. Vázquez-Baeza, Y., Pirrung, M., Gonzalez, A. & Knight, R. EMPeror: a tool for visualizing
high-throughput microbial community data. *Gigascience* **2**, 16 (2013).

32. Janssen, S. *et al.* Phylogenetic Placement of Exact Amplicon Sequences Improves
Associations with Clinical Information. *mSystems* **3**, (2018).

382