A Systematic Review and Meta-analysis on the Transcriptomic Signatures in Alcohol Use Disorder – a Translational Approach

Marion M Friske*[1], Eva C Torrico[1], Anna M Borruto[1], Maximilian JW Haas[1], Francesco Giannone[1], Andreas-Christian Hade[2,3], Mari-Anne Philips[4], Wolfgang H Sommer[1], Rainer Spanagel[1]

[1]Institute of Psychopharmacology, Central Institute of Mental Health, Mannheim, University of Heidelberg, Germany.

[2]Department of Pathological Anatomy and Forensic Medicine, University of Tartu, Estonia.

[3]Forensic Medical Examination Department, Estonian Forensic Science Institute, Tallinn, Estonia.

[4]Department of Physiology, Institute of Biomedicine and Translational Medicine, University of Tartu, Tartu, Estonia.

*Corresponding author:

Marion M. Friske

marion.friske@zi-mannheim.de

Abstract

Alcohol Use Disorder (AUD) is a complex mental disease with a large treatment need. However, the clinically available therapeutic approaches are limited and thus new treatment targets are warranted. One approach to discover new treatment targets is transcriptomic profiling of key brain regions of the addiction neurocircuitry in animal models of AUD and/or in post-mortem brain tissue from deceased AUD patients. Unfortunately, those studies suffer from large heterogeneity and low sample sizes. Here we used a cross-species meta-analysis that integrates the entire literature on this topic to make more robust conclusions on transcriptomic changes in AUD. Here, we report on differentially expressed genes (DEGs) in 108 rat brain and 380 human brain samples, respectively in one of the key regions of AUD – the prefrontal cortex. We found 50 differentially expressed genes (DEGs) in rat brain, which were mainly associated with pro-oncogenic molecules, signal transduction as well as oxidative stress and inflammation. When we compared these findings with the meta-analytic results in human post-mortem PFC, we found that across species Hsd11b1 is significantly down-regulated. This finding is also confirmed by our cross-species data integrational approach. In summary, our data suggests that down-regulation of Hsd11b1 enzyme and stress-related pathways is a conserved mechanism for the alcohol dependent phenotype across different species, which might explain increased cortisol levels in AUD and therefore serves as a new treatment target.

Keywords: transcriptomics, Meta-Analysis, Alcohol Use Disorder, alcohol, post-mortem, brain tissue, rodents, human, prefrontal cortex.

Introduction

Alcohol Use Disorder (AUD) is a complex mental disease that causes annually more than three million deaths worldwide, which represents 5.3% of deaths per year in total (WHO, 2019). Alcohol has been classified as the most harmful drug in a comparison of the most commonly abused licit and illicit drugs, considering both harm to the user and others (Nutt et al., 2010). Importantly, alcohol is acting systemically and consequently its detrimental effects affect all organ systems. In addition, there are numerous closely linked physiological and psychological comorbidities of AUD such as major depressive disorders (MDD), antisocial disorders, as well as liver and kidney injuries (McHugh et al., 2019; Rocco et al., 2014; Helle et al., 2019; Rumgay et al., 2021).

A key goal of AUD research is to identify the underlying neuroadaptations and molecular targets that lead to alcohol addiction in order to repurpose existing drugs or develop new medications (Burnette et al., 2022; Kranzler & Soyka, 2018; Witkiewitz et al., 2019). Previous studies suggested molecular pathways in diverse brain regions that might be impaired by chronic alcohol consumption and might be involved in the establishment of AUD (Abrahao et al., 2017, Gilpin& Koob, 2008). A growing body of research supports the neuroimmune hypothesis of AUD, which describes the effects of chronic alcohol intake on inflammation and neuromodulation (Mayfield & Harris, 2017, Erickson et al., 2019). Also, AUD has been associated with increased oxidative stress and impaired cell proliferation leading to increased risk for certain cancer types (Seitz & Stickel, 2007; Tsermpini et al., 2022). In addition, chronic alcohol consumption and AUD have been shown to cause disruptions in the circadian rhythm (Davis et al., 2018) and clock genes have been linked to addictive behavior (Perreau-Lenz & Spanagel, 2015). Long-term alcohol consumption, AUD and withdrawal are inducing stress-related pathways and also stressful experiences are elevating

alcohol consumption (Perreau-Lenz & Spanagel, 2015; Spanagel et al., 2014). Consequently, AUD is a common comorbidity of patients with high stress-associated psychological disorders, such as PTSD and anxiety disorders (Debell et al., 2014; Gilpin et al., 2015). Nevertheless, even though numerous studies have been conducted to explore the effects of high alcohol consumption in general as well as compulsive alcohol drinking correlated to AUD, medical approaches for AUD therapy are still limited and only partially effective. So far, naltrexone, acamprosate and disulfiram are the only FDA-approved treatments for AUD (Witzkiewtz et al., 2019), whereby 60% of all AUD patients experience at least one relapse period during or after therapy, irrespective of the treatment itself (Mayernhoff & Durazzo, 2010; Kirshenbaum et al., 2009; Witkiewtz 2011). Furthermore, it is interesting to note that some of the most promising therapeutic mechanisms identified by preclinical research have failed in clinical trials (Schwandt et al., 2016; Heilig et al., 2019; Kwako et al., 2015; Heilig et al., 2016). As a result of these substantial translational failures, pharmaceutical industries ceased to invest efforts in the development of new possible medications (Hyman et al., 2014), leading to the still open debate on the utility of animal models for preclinical testing of novel therapies (Hyman, 2012; Heilig et al., 2021).

Animal research is characterized by several limitations, which can make the interpretation of the results obtained from the studies challenging. A major limitation is the lack of statistical power often caused by insufficient sample size, which can produce unreliable results (Button et al., 2013). Additionally, inter-study variability (due to e.g., breeding facility, experimenter, laboratory conditions, species/strain, experimental protocols) leads to diverse outcomes among published studies. These limitations impact the translatability of preclinical research. Meta-analyses are a powerful tool for at least partially overcoming such limitations (Sena et al., 2014). Hence, in this study we aim to perform systematic literature

screening and meta-analysis on transcriptome-wide studies focusing on brain samples derived from rodents exposed to ethanol. Even though no known animal model of addiction fully mimics the human condition, certain animal models are better suited to investigate specific components of the addiction process in a clinically relevant way (Lynch et al., 2010). We therefore focused our attention on the well-established post-dependent animal model, obtained via chronic intermittent alcohol exposure (CIE), that mimics the natural process in which humans develop alcohol addiction: over a prolonged period of time and after several intermittent drinking episodes interspersed with withdrawal periods (Meinhardt & Sommer, 2015). This model results in robust and stable behavioral symptoms of alcohol dependence that persist during prolonged abstinence and is characterized by long-lasting neuroadaptations that mediate such symptoms (Heilig et al., 2017). Importantly, we also applied the same procedure to transcriptome-wide studies performed on human AUD patients and healthy controls to identify commonly altered genes and gene networks among species. We chose the prefrontal cortex (PFC) as the brain region of interest for this meta-analysis. Converging lines of evidence from human and animal studies have revealed that the PFC is a key structure for performing highest-order cognitive functions (Abernathy et al., 2010; Avchalumov et al., 2021; Ball et al., 2011). Results gained over the past decade suggest that chronic alcohol exposure profoundly impairs this brain region (Abernathy et al., 2010; Pahng et al., 2017; Schacht et al., 2013; Charlton et al., 2019), leading to a dysfunctional control over compulsive behavior (Bechara, 2005; Pahng et al., 2017; Timme et al., 2022; de Paiva Lima et al., 2017). Even after abstinence, alcohol-induced cognitive dysfunctions can affect top-down inhibitory control, decision-making, and behavioral flexibility (Trantham-Davidson & Chandler, 2015; Catafau et al., 1999; Grüsser et al., 2004). In addition, as part of the extended reward system, it has a central role in the development

and maintenance of AUD (George & Koob, 2010, Goldstein & Volkow, 2011, Wise & Koob, 2014). Notably, as a stress-sensitive brain region, the PFC also plays a crucial role in the withdrawal state and relapse-like behavior (Blaine et al., 2016; Blaine & Sinha, 2017; Seo & Sinha, 2015; George et al., 2012), suggesting the functional dysregulation of this brain region as a hallmark of early relapse risk in alcoholic patients.

We believe that our findings might deliver a profound basis for further studies to understand the mechanisms of AUD and to develop new therapeutic approaches that could potentially be more successful than the currently available medications on the market.

Materials and Methods

Systematic Literature Screening

The systematic literature screening was performed as described in the PRISMA Guidelines (Page et al., 2021). Briefly, two researchers independently screened the literature with predefined keywords for PubMed and EMBASE that were determined to specify four main categories of interest: alcohol, animal model of alcohol dependence, gene expression, and brain (Fig. 1). For post-dependent rodent studies, the following keywords were used: (alcohol OR ethanol) AND (rats OR mice OR rat OR mouse) AND (ribonucleic acid OR RNA OR RNA-seq OR mRNA OR ncRNA OR miRNA OR siRNA OR piRNA OR snRNA OR microRNA OR expression OR gene OR cDNA OR transcription OR transcript OR transcripts OR transcriptional OR transcriptomic OR translation OR transcriptome OR transcriptomics OR translation) AND (post-dependent OR "post dependent" OR postdependent OR cie OR "chronic intermittent"). For human post-mortem studies, we used the following keywords: ("post-mortem" OR "postmortem" OR "human" OR "patients") AND ("brain") AND ("alcohol" OR "alcoholic" OR "alcohol dependence" OR "alcoholism" OR "AUD" OR "alcohol use disorder" OR "alcohol addiction") AND ("transcriptome" OR "transcriptomic" OR "transcriptomics" OR "gene-expression" OR "gene expression" OR "mRNA" OR "messenger RNA"). The literature screening was performed using the SysRev screening tool developed by the CAMARADES research group at the University of Edinburgh (www.syrf.org.uk). After each screening step (title/abstract and full text), the researchers compared the respective resulting studies. In case of disagreement, a third independent party chose whether the study was included or excluded for the next screening step. The general structure of the predefined keywords, as well as the complete PRISMA flow diagrams with resulting number of studies for each screening step for rodent and human study screening is depicted in Fig.1.

Since the majority of the identified studies was working on tissue derived from the PFC, we decided to focus on this brain region as scope of our analysis.

A Alcohol/ Ethanol AND Species-specific & model-defining terms AND RNA-specific terms AND brain

B Identification of **rodent** studies via databases and registers

Screening

Records identified from:
Databases (n = 656)
PubMed (n = 192)
EMBASE (n = 464)

Records removed before screening:
Duplicate records (n = 122)

Identification

Records screened
(n = 534)

Records excluded
(n = 307)

Abstracts screened
(n = 227)

Records excluded
(n = 48)

Full text screening
(n = 179)

Included

Studies included in review
(n = 22)
Studies included in quantitative analysis
(n = 5) → 7 datasets, 106 individuals

Inclusion criteria:
rodent (mice/rats), alcohol exposure (≥2 weeks), post-dependence (≥3 days abstinence prior to death), transcriptome-wide RNA data from brain tissue, high quality RNA samples, EtOH-naïve controls

Exclusion criteria:
adolescence, prenatal alcohol exposure, too short exposure/ abstinence time, additional treatments, ethanol-preferring lines/ selectively bred for high ethanol intake

C Identification of **human** studies via databases and registers

Screening

Records identified from:
Databases (n = 715)
PubMed (n = 290)
EMBASE (n = 425)

Records removed before screening:
Duplicate records (n = 157)

Identification

Records screened
(n = 593)

Records excluded
(n = 418)

Abstracts screened
(n = 175)

Records excluded
(n = 114)

Full text screening
(n = 61)

Included

Studies included in review
(n = 23)
Studies included in quantitative analysis
(n = 8) → 8 datasets, 380 samples

Inclusion criteria:
human post-mortem brain tissue, diagnosed AUD (e.g. DSM-IV), transcriptomewide RNA data, high quality RNA samples, healthy

Exclusion criteria:
other psychiatric disorders as comorbiditeis, co-abuse of other drugs, brain injury as cause of death, age < 18 years

Fig. 1: General structure for keyword design for systematic literature research in PubMed and EMBASE representing the four categories based on which we specified the keywords (**A**)

and the resulting number of studies for rodent studies (n=5 studies, 7 datasets) (**B**) as well as human post-mortem brain studies (n=8 studies) (**C**). Two rodent studies each included two datasets fulfilling our criteria, which explains the higher number of datasets compared to studies. Since in both species, the PFC was predominantly used in the original studies, we focused the analysis on this brain region. Before the screening, duplicates were removed using EndNote. The template for overview of the search procedure and the resulting studies is taken from www.prisma-statement.org.

Only studies providing transcriptome-wide RNA-expression data, such as microarray or RNA-Seq data, were considered for the meta-analysis. The Summary Statistics – including the full profile of all detected genes from the sequencing experiment – of the eventually resulting studies, were either downloaded from the original publication or the corresponding authors were contacted to request the respective datasets. Details of the included studies are listed in Suppl. Tab. 1. and Suppl. Tab. 2.

As usually included in systematic reviews and meta-analyses, we aimed to perform an assessment of risk of bias (RoB) for the original studies. Unfortunately, inadequate reporting was impeding us to access full information necessary for proper RoB assessment.

Meta-Analysis

All statistical analyses were performed using the statistical program R Version 4.1.2. Rodent datasets and human post-mortem datasets were meta-analyzed separately and then compared in later steps. As reported previously, analyzing microarray datasets from scratch can be challenging due to high diversity in processing platforms, methods and analysis softwares (Kontou et al., 2018). So far, there has been no standardized analysis

methodology available to combine all these diverse datasets into one generalized analysis pipeline. Therefore, we decided to combine the summary statistics representing log2 fold-changes and corresponding p-values of all detected genes per study instead. Additionally, due to the diverse analysis platforms used, the estimation of the effect sizes was inconsistent across the original studies, which was suggesting to not perform an effect size combinatorial approach in this study. According to a previously published guideline on gene expression meta-analysis (Toro-Domínguez et al, 2021), we eventually decided to perform weighted Stouffer's p-value combination using the R package DExMA (Villatoro-Garcia et al., 2022). P values of log2-fold change calculations from original studies were used as input and studies were weighted by the square root of sample size. We set the threshold for inclusion of a gene to 60 %, meaning that each gene included in the meta-analysis was represented by at least ⅔ or 4/7 of the original studies for rodent and human datasets respectively. Before conducting the meta-analysis, all datasets were annotated with the most current databases mouse4302.db, rae230a.db, illuminaHuman4.db, org.Hs.db, EnsDb.Hsapiens.v86 to ensure comparability of the gene names. Multiple entries for one gene within a study were grouped using the collapseRows function from the R package WGCNA (Langfelder and Horvath, 2008; Miller et al., 2011). Since in the mouse and rat genome not all genes have common names, we decided to use biomaRt for annotating the rodent genes with the mouse orthologue to ensure we are not missing out on data points due to the potential divergent naming of same genes (Durinck et al., 2009). The individual analysis steps as well as the packages for human and rodent meta-analysis in use can be found in Fig. 2. Significant DEGs were defined by a threshold of $p_{corr}<0.05$.

Fig. 2: Analysis pipeline for meta-analysis in animal and human post-mortem PFC tissue, including data pre-processing, meta-analysis and downstream analysis investigating potential enrichment in pathways and cell types.

In addition, a consistency index for the differentially expressed genes was estimated considering fold-change direction and the value of the fold-change per gene that was significant in the meta-analysis. Therefore, the index was calculated for all respective genes considering its values within each original study as described in Eq. 1. The corresponding values can be found in Suppl. Tab. 3 and Suppl. Tab. 5 in the respective column.

$$\text{Consistency index} = (\underline{\hspace{6cm}})$$

Eq. 1: Calculation of consistency index per gene that was detected as significant in the meta-analyses of rodent and human datasets to observe the level of consistency in the direction of dysregulation across the original publications in comparison with the meta-analysis results.

For the rodent analysis, we additionally performed subgroup analysis considering mouse and rat datasets separately to observe potential species-specific effects contributing to the main effects of the meta-analysis results.

Subgroup analysis rodent datasets

To determine the species-specific impact on the outcome of the meta-analysis of post-dependent animals, we performed a subgroup analysis by dividing the original studies according to the included species. Afterwards, we performed a meta-analysis within the subgroups and compared the results with the DEGs derived from the meta-analysis of the two rodent species together. The threshold for considering a gene for the analysis was appearance in at least 60 % of the datasets in the respective subgroup. Significant DEGs were defined by a threshold of $p_{corr}<0.05$.

Down-stream analysis

To identify potential pathways associated with the DEGs identified in the meta-analysis, we performed gene set enrichment analysis (GSEA) with GO-terms and KEGG-pathways using the R package ClusterProfiler (Wu et al., 2021) and g:Profiler (Raudvere et al., 2019). Additionally, we analyzed potential enrichment of the DEGs for specific cell-types. Therefore, we performed cell-type enrichment analysis using the R package LRcell (Ma, 2021).

Overlap between human and animal DEGs

Since animal and human genes differ in their nomenclature, we first identified the human orthologues of the animal DEGs using the Babelgene R package and then compared these

converted DEGs with the human dataset to find potential overlaps in the results. In addition, we also compared pathway analysis results for both species to identify possible commonly dysregulated pathways in the animal model as well as in the human situation. For comparison, significant DEGs were defined by a threshold of $p_{corr} < 0.1$.

Cross-species data integration

The VennDiagram approach is a first step to descriptively compare and show potential overlaps of the results coming from the species-specific meta-analyses. To further investigate which genes are conserved for the alcoholic phenotype across species, we performed a cross-species integrational meta-analytic approach. Therefore, we used the "homology method", where we first transferred the human genes coming from the initial Summary Statistics per study to their respective rodent orthologues as reported previously (Schall & Latham, 2021). Afterwards, separate appearance indices for human and rodent species were calculated as the sum of the number of studies a gene occurred within one species (Eq.2 (1)). Then, the cross-species appearance indices per gene were combined to an overall appearance index as described in Eq. 2 (2). The cross-species appearance index cut-off was set to 0.8, which is representing 60% of rodent and human studies each to ensure that the genes included in the cross-species meta-analysis are included in a representative number of studies. To avoid over-representation of rodent studies, the upper limit for inclusion was set to a cross-species appearance index of 1.25.

(1) Species-specific appearance index = $\sum_i^n \quad index(i) = study1(i) + study2(i) + \ldots + studyn(i)$

(2) Cross-species appearance index = $\left(\frac{Appearance\ index\ rodents}{Appearance\ index\ humans}\right)$

Eq. 2: (1) Calculation of the species-specific appearance index considering the species separately. Let i be a gene measured in the RNA quantification experiment from the original

studies and n be the number of studies per species. If a gene was targeted in the original publication, the value 1 was assigned to the study, whereas 0 was assigned to the study, whenever the gene was not measured. The species-specific appearance index was calculated for each gene and each species separately. (2) Equation of the cross-species appearance index, which represents an overall index considering all species in combination. These indices were used as a cut-off value for integrating genes into the cross-species meta-analytical approach, considering also the proportion of rodent and human studies to avoid over-representation of one species.

Including genes with a cross-species appearance index between 0.8 and 1.25, combined meta-analysis of rodent and human datasets was performed using rodent orthologues for the human datasets and the R package DExMA for meta-analysis. The resulting significant DEGs ($p < 0.05$), were included in further downstream analyses as described in the section "down-stream analysis".

Results

Meta-Analysis in PFC of post-dependent rodents discovered six genes that were not discovered by the included original studies.

Systematic literature screening resulted in seven studies matching the research criteria for this study including the PFC as the brain region of interest (Fig. 1). Since two of these studies included more than one dataset that is matching our criteria, this was leading to nine datasets derived from seven studies. Out of those, we were able to retrieve seven datasets derived from five studies, which resulted in 106 individuals in total. Three datasets were conducted from rats (n= 42 individuals) and four datasets contain mouse tissue (n= 64 individuals). Detailed information about the included studies can be extracted from Suppl. Tab. 1 and the inclusion and exclusion criteria in Fig. 1. First, we observed the distribution of the significant DEGs per original study to get an impression of the heterogeneity across these studies (Fig. 3), where we found 11-38% of the DEGs overlapping with at least one additional original study, while the remaining 62-89% of DEGs were represented by only one original study. This finding describes a high heterogeneity across the original studies based on their reported DEGs. When we compared these genes with the outcome of the meta-analysis that resulted in 50 significant DEGs (p_{corr}<0.05), we observed 44 genes (corresponding to 88% of all significant DEGs of the meta-analysis) overlapping with at least one original study, while six genes (corresponding to 12% of the gene set) were found as solely significant in the meta-analysis (Fig. 3). These six genes are Rnaseh1, Dr1, Lancl2, Mc5r, Scn11a, and Tmem161b. To our knowledge, most of these genes have not been related to chronic alcohol consumption or AUD in previous studies. However, Lancl2 has been suggested as a candidate gene of interest underlying the chromosome 4 QTL for alcohol consumption in alcohol-preferring iP rats and non-alcohol-preferring iNP rats

(Spence et al., 2013). Scn11a has previously been correlated to the altered function of encoded sodium channels, leading to hyperexcitability in nociception as a response to excessive alcohol consumption (Maingret et al., 2008).

Fig. 3: Venn Diagram comparing the significant DEGs ($p_{corr}<0.05$) identified by the original studies and comparison of these genes with the significant DEGs derived from meta-analysis. No gene was found to be significantly dysregulated among all the datasets. This figure solely represents original studies that reported significant DEGs for PFC; for meta-analysis all datasets were included, regardless of the presence of reported significant results for this brain region. Please note that for Meinhardt et al., 2013 and Smith et al., 2016 we included two datasets per study with timepoints that were focusing on either diverse exposure length or abstinence duration. Further details about the original studies can be extracted from Suppl. Tab. 1. Colors of the ellipses were chosen randomly to provide proper visualization and comparability of the overlaps across the datasets.

Although observation of the heterogeneity of DEGs across the original studies did not suggest species-specific differences in the distribution of the transcriptomic patterns, we performed subgroup analysis, considering mouse and rat studies as respective subgroups (Fig.4, Suppl. Tab. 4). Interestingly, we did not find any DEG overlapping between the species that were not found in the meta-analysis results as well. One gene, namely Nidogen-2 (Nid2) (up-reg., cell adhesion), appeared significant across all three datasets: mouse subgroup, rat subgroup and the meta-analysis. In the main analysis, we identified six DEGs that were not significant in any of the original datasets. In the subgroup analysis, we found eleven DEGs that did appear significant only in the meta-analysis dataset, but not in the subgroup analyses (Fig. 4). Despite Tmem116, which was significant in the rodent subgroup analysis and in the meta-analysis, the previously identified unique DEGs of the meta-analysis remained unique by this type of analysis as well (Fig.4). Additionally, six DEGs appeared as unique for the meta-analysis results, which are Birc7 (up-reg., apoptosis), Dio1 (down-reg., thyroid hormone activity), Itga7 (up-reg., autophagy), Pnpla3 (down-reg., balance of energy usage/storage in adipocytes), Tat (down-reg., tyrosine metabolism) and Ttc7b (up-reg., plasma membrane). For rat datasets, 30 genes (corresponding to 4 % of all DEGs of this dataset) overlapped with the results of the main analysis, whereas for mice 13 genes (corresponding to 6% of all DEGs of this dataset) overlapped with the main analysis. When comparing the number of DEGs from the mice subgroup analysis (n=4 datasets, all J57BL/6 strain) to the rat subgroup analysis (n=3 datasets, all Wistar strain), the mice subgroup shows less than ⅓ of the number of DEGs derived from the rat subgroup.

Fig. 4: Subgroup analysis considering mouse and rat studies as individual subgroups and comparison of the resulting genes with the main findings of the meta-analysis. Venn diagram is depicting the significant DEGs ($p_{corr}<0.05$) of the separate analyses per species and of the meta-analysis. Rat subgroup analysis revealed 715 DEGs, mouse subgroup analysis resulted in 202 DEG, and meta-analysis across both species led to 50 DEGs in total. One gene, Nid-2, is commonly significant in all three datasets. The percentage of genes per overlay is calculated with respect to the total number of DEGs detected across all three datasets (n= 972 genes). All included mouse studies worked with the J57BL/6 strain, whereas all rat studies work with the Wistar strain. Additional information can be found in Suppl. Tab. 4. Colors of the circles were chosen randomly to provide proper visualization and comparability of the overlaps.

Thus, in total our analysis of rodent PFC identified 50 DEGs, we next were interested whether these genes were enriched for specific pathways. Therefore, we performed pathway enrichment analyses using mouse orthologs and mouse terms derived from the

most commonly used databases (Supplementary). Signal transduction-related pathways as well as nitrogen metabolic pathways were among the most significant pathways.

Additionally, if we consider individual genes and their involvement in pathways based on the literature, it seems that apoptosis and general cell maintenance controlling genes are downregulated whereas genes that have previously been correlated to cancer, such as Fgf16, Cabin1, Usp35 and Atg101 (He et al, 2021; Zhang et al., 2018; Wang et al., 2022; Zhang et al., 2022) were up-regulated. This finding suggests a higher prevalence of tumor development in ethanol-exposed rodents. In addition, the gene Tat – encoding for Tyrosine aminotransferase –, which is associated with tyrosine breakdown, is up-regulated. Tyrosine is directly correlated to dopamine biosynthesis. Therefore, enhanced tyrosine breakdown might lead to significant dopamine decrease. Previous studies have shown that alcohol-dependent animals as well as AUD patients have decreased dopamine levels, which were associated with withdrawal and the likelihood of relapse (Dahchour & Ward, 2022; Hansson et al., 2019).

In addition, we observed potential enrichment of the DEGs derived from rodent meta-analysis in PFC for cell-type specific patterns, which resulted in enrichment for layer 6 neurons (Suppl. Fig. 1).

Meta-Analysis in PFC of human post-mortem samples results in 278 DEGs enriched in immune- regulatory and pro-oncogenic pathways.

Systematic literature screening resulted in ten studies for quantitative analysis for human post-mortem PFC, where we were able to conduct eight datasets in total, resulting in 380 samples (Fig. 1). Since all included studies were using samples derived from the NSWBTRC at

the University of Sydney, there might be an overlap among the patient samples, which might lead to over-representation of certain individuals. To date, unfortunately, we were not able to correct for this potential bias in our analysis and therefore we were considering each sample as unique.

When we observed the total distribution of significant DEGs (p < 0.1) across the original studies, we found that 50-84 % of DEGs represented solely by the original studies and therefore were only significant in one publication (Fig. 5). When additionally considering the DEGs derived from the meta-analysis, the identified significant DEGs (p < 0.1) had an overlap of 37 % with at least one of the original studies and one gene – Mt1x, encoding for Metallothionein 1X – was overlapping with two original datasets. Fig. 5 shows the Venn Diagram representing the overlap of all significant DEGs reported in the original studies as well as the significant DEGs derived from the meta-analysis.

Fig. 5: Venn Diagram representing the overlap of significant DEGs reported in the original studies (p<0.1) and DEGs derived by meta-analysis (p<0.1). No gene was found to be significantly dysregulated among all the datasets. This figure solely represents original

studies that reported significant DEGs for PFC (n=4), while for meta-analysis all PFC datasets (n=8) were included, regardless of the number of reported significant results in the original publication. Detailed information about the included publications can be found in Suppl. Tab. 2. Colors of the ellipses were chosen to provide proper visualization and comparability of the overlaps coming from different studies.

Next, we tested the significant DEGs (p<0.05) of the meta-analysis for enrichment in specific pathways (Suppl. Tab. 6). Generally, DEGs were enriched in GO-terms associated with inflammation, cell adhesion, and signal transduction, whereas enriched KEGG-pathways were associated with apoptosis, cell differentiation, and inflammatory pathways. When we then ordered the DEGs according to their fold-change direction into activated and repressed genes, we found activated NF-κB expression, and toll-like receptor signaling, as well as cytokine storm and pro-oncogenic pathways (Suppl. Tab. 6). Down-regulated DEGs were enriched for suppressed metabolic processes, neuroactive ligand-receptor interaction and glycerolipid metabolism, as well as signal transduction-related pathways and cellular membrane stability (Suppl. Tab. 6).

Cell-type specific enrichment of the DEGs derived from meta-analysis in human post-mortem brain tissue resulted in a tendency for enrichment in oligodendrocytes (p= .12; Suppl. Fig. 2).

Overlap DEGs derived from a meta-analysis in rodents exposed to ethanol and human post-mortem brain tissue suggests a conserved mechanism for the alcoholic phenotype.

To identify common alterations in gene expression across species, we first determined the human orthologues of the DEGs derived from the rodent datasets. Fig. 6 depicts the results from the comparison of these DEGs across human and rodent species. The overlap between human and rodent meta-analysis results in six commonly dysregulated genes: Adcy3, Hsd11b1, Itm2c, Oga, Rab26 and Rnf122. Interestingly, these genes were at least partially also dysregulated in the same direction in both human and rodent meta-analysis results. However, when we consider the subgroup analysis from the rodents, Hsd11b1 was only significant in the rats and pointing towards an up-regulation, whereby a combination of the rodent datasets is still remaining significant, but leading to a total down-regulation of Hsd11b1 (Suppl. Tab. 4). Interestingly, to our knowledge these six overlapping genes have so far not been studied with an intense interest in the context of alcohol consumption and AUD in the brain.

Fig. 6: Venn Diagram representing the comparative overlap of significant DEGs ($p_{FDR}<0.1$) derived from the meta-analysis in PFC of post-dependent rodents and AUD patients respectively results in six genes being significantly dysregulated. Across these species, Hsd11b1 and Rnd122 were consistently down-regulated, and Itm2 was consistently up-

regulated. Further information can be extracted from Suppl. Tab. 4. Colors of the circles were chosen randomly to provide proper visualization and comparability of the overlaps.

Cross-species data integration confirms the importance of dysregulation in the cortisol pathway for the alcoholic phenotype.

Eventually, we aimed to find genes that are conserved for the alcoholic phenotype across species. Therefore, we performed a cross-species integrational meta-analytic approach using cross-species appearance indices to ensure equal contribution of the species (Eq. 2). After applying the cross-species appearance indices for every homologous gene detected in the original experiments, 8513 genes from eight human and seven rodent original datasets were included. Meta-analysis resulted in 31 significant DEGs (p < 0.05), 22 up-regulated and 9 down-regulated. Hsd11b1 – the gene that showed highest significance in the comparison between the species-specific meta-analyses (Fig. 6) – also appeared in this cross-species meta-analysis as the most significantly down-regulated gene (Suppl. Tab. 7).

Down-stream pathway enrichment analysis using the 31 DEGs resulting from cross-species data integration resulted in enrichment in signal transduction-related pathways, such as glutamatergic signal transduction, signal receptor binding, and focal adhesion (Supplementary). Additionally, enrichment in general biological processes involving protein biosynthesis and metabolism was detected.

Cell-type enrichment analysis resulted in significant enrichment for endothelial cells, astrocytes, oligodendrocytes and polydendrocytes (p_{FDR}=0.05; Suppl. Fig. 3).

Discussion

We performed a meta-analysis on transcriptome-wide datasets derived from post-dependent rodent and post-mortem human brain tissue in the context of alcohol dependence and AUD in a comprehensive manner. (i) Meta-analysis in post-dependent rodent PFC led to 50 DEGs, whereby six of these genes were not identified before in the original studies included in the analysis. (ii) The DEGs in rodents were mainly related to immune-regulatory and cancer-related pathways and showed enrichment in layer 6 neurons. (iii) Meta-analysis in human post-mortem PFC resulted in more than 278 DEGs, which showed association with inflammatory, cell adhesion and apoptosis pathways and enrichment in oligodendrocytes. (iv) Cross-species comparison and integration suggest a novel mechanism to explain increased cortisol levels in alcohol addiction via down-regulation of Hsd11b1 as a conserved mechanism for the alcohol dependent phenotype.

The observation of the DEGs identified in the respective original rodent studies pointed out a high heterogeneity, since 62-92% of the identified significant genes were found dysregulated in only one original study. One of the general aspects of performing meta-analyses is to observe an effect across several original studies, considering heterogeneity introduced by e.g., diverse study design, and including a higher sample size for the analysis to increase statistical power. The varying study design, such as the differences in length of alcohol exposure and duration of abstinence before sacrifice of the animals, might be a potential explanation for the heterogeneous DEG distribution among the original studies. Also, it is known that mice and rats show differing transcriptomic patterns in response to alcohol exposure (Contet et al., 2012). Additionally, individuals derived from different breeders can differ genetically and physiologically. It has been shown in the past, that the

microbiome has an impact on gene expression in the brain via the gut-brain-axis (Margolis et al., 2021) and that the microbiome of laboratory animals is influenced by a variety of factors (Ericsson & Franklin, 2021). Nevertheless, this heterogeneity across the original studies points out the importance of our meta-analysis: genes being identified as significantly dysregulated despite these inter-study variances might represent targets of particular interest, as their dysregulation was detected regardless of the chronic alcohol treatment paradigm. Also, in the clinical situation of human AUD patients and the post-mortem brain samples included in the pre-clinical analyses, the background of the patients varies in terms of drinking duration and relapse phases, environment, genotype, period of abstinence before death and psychological aspects that influence the transcriptomic profile. Therefore, observing transcriptomic changes on a heterogeneous background might deliver promising targets that could be representative for the general alcohol dependent phenotype that might be useful for treatment of a broad spectrum of AUD patients.

In the meta-analysis of alcohol-dependent rodent PFC, we found 50 DEGs (p<0.05), including six genes that do not overlap with the previous findings of any of the original studies. Interestingly, to our knowledge, most of these uniquely identified genes have not yet been addressed in the alcohol field. Tmem161b has previously been associated with reward processing in alcohol dependence in a GWAS on patients with major depressive disorder (Muench et al., 2018). Lancl2 was suggested as a candidate gene of interest underlying the chromosome 4 QTL for alcohol consumption in iP and iNP rats (Spence et al., 2013). In addition, Scn11a has been found to cause altered functions of encoded sodium channels that are correlated with hyperexcitability and nociception in response to alcohol consumption (Maingret et al., 2008; Yang et al., 2020). However, when we performed subgroup analysis considering the rodent species, mouse and rat as separate subgroups, we

identified eleven unique DEGs that were represented by the meta-analysis results solely and not in any of the subgroup-specific results. Within these eleven genes, five out of the six unique genes from the previous comparison of the meta-analysis with the results of the original studies were replicated. Also, the number of non-overlapping DEGs in the meta-analysis decreased when comparing these resulting genes with the DEGs from subgroup analyses. One gene was found to be significant in both of the subgroups as well as in the meta-analysis, which is Nid2, a protein-coding gene that is involved in cell adhesion. Nid2 is essential for the plasticity and excitability of neurons, neuronal network establishment and maintenance (Vasudevan et al., 2010). In our analysis, Nid2 was significantly up-regulated in post-dependent rodents, which therefore might lead to increased neuronal plasticity and excitability. Also, an increase in Nid2 due to aging has been previously found in humans (Keable et al., 2016). Nid2 has been suggested as a potential tumor marker for ovarian, esophagus and colon cancer (Kuk et al., 2010; Chai et al., 2018; Yu et al., 2019) and therefore, its over-expression might also give evidence for an elevated risk for tumor development in these animals.

In general, we observed differences between the results coming from subgroup analysis and meta-analysis. Subgroup-specific differences leading to stronger effects coming from one subgroup only and potentially evening-out effects coming from opposite effects in mice and rats are enlightened by this type of additional analysis. The different outcomes of the subgroup analyses compared to the results from the meta-analysis in terms of the number of DEGs and the general overlap might be due to the settings of the analysis. To avoid the bias caused by genes that might have strong effects, but were only detected in a small number of studies, we set the threshold for including a gene in the analyses to 60%.

Pathway analysis of the DEGs related to rodent PFC meta-analysis resulted in enrichment for dysregulated basic cellular regulatory pathways, such as general cellular catabolic processes, voltage-gated channel regulatory activity, cell membrane regulatory aspects and signal transduction. In addition, the DEGs of the rodent PFC were enriched for pathways related to nitrogen biosynthesis. Nitrogen is an essential element for biosynthesis of amino acids and therefore general protein synthesis. Also, nucleotides –the smallest compartment of DNA and RNA molecules– central coenzymes such as NAD, and hormones have nitrogen implemented as an essential element. Therefore, impairment of nitrogen biosynthesis is supposedly having a broad influence on cellular mechanisms and metabolisms. As central molecule of the urea cycle, nitrogen biosynthesis dysregulation is not only having impact on digestion-related pathways across the whole organism, but also urea cycle disorders have previously been found to impair brain functions, mainly on astrocytes, and therefore lead to disabilities in cognitive functions (Gropman et al., 2013). When we then split the DEGs of the meta-analysis according to the direction of the fold-change, we found a potentially elevated risk for cancer development, since some genes that are known to be pro-oncogenic molecules were up-regulated. This finding is adding to the previous studies that found an increased risk for cancer in alcohol-exposed rodents and AUD patients (Chan et al., 2014; Rumgay et al., 2021).

When we looked into a potential cell type-specific enrichment of the DEGs identified by meta-analysis, we found significant enrichment for layer 6 neuronal cells in post-dependent rodents. These days, more and more single-cell sequencing data are published, which are pointing out that different cell types play an important role in health and disease. Neurons in general, and especially cortical neurons, have been previously shown to be associated

with immune system deficiency caused by alcohol use disorders (Erickson et al., 2019). Layer 6 is the deepest layer of the mammalian cerebral cortex and contains the highest variety of neuronal cells, including pyramidal, inhibitory and excitatory neurons. Previous studies reported the effects of ethanol on layer 6 neurons mainly in the context of alcohol-exposure during brain development and fetal ethanol exposure. Therefore, a generally reduced size and altered morphology of prefrontal cortex layer 6 neurons were found in mice exposed to ethanol during fetal development (Louth et al., 2018). In addition, the synaptic activity of layer 6 neurons of the somatosensory cortex is altered by prenatal ethanol exposure in mice (Delatour et al., 2020). However, even though this specific type of neuronal cells seems to have been only in the scope of studies focusing on the influence of ethanol exposure during the developmental stage, our data suggest that alcohol consumption might have an impact on this cell type also at the adult stage.

In the meta-analysis of human post-mortem brain samples derived from the PFC, we identified a high heterogeneity among the reported DEGs from the original studies and the meta-analysis results, as well. This heterogeneity might at least partially be explained by the inclusion of different subjects per original study, which are potentially representing diverse genotypes, drinking patterns, and environmental aspects. In addition, the time span in which these data were obtained is more than 20 years and therefore RNA detection methods as well as analysis methods have changed throughout the years, which might introduce a technical bias into our analyses. However, since we analyzed Summary Statistics from the original studies and all human samples included in our analysis were originally derived from the NSWBTRC at the University of Sydney, Australia, we were not able to balance out the bias due to the potential over-representation of specific individuals that

have been analyzed in more than one of the original studies. Therefore, we have to consider that potentially patients are over-represented in our analysis.

Our analysis of human PFC resulted in a total number of 268 DEGs ($p < 0.05$), where more than half of these genes were uniquely identified by the meta-analysis and did not appear as significant in the original studies. Downstream analysis showed enrichment in several pathways. GO-term and KEGG pathway analyses were associated with inflammation, cell adhesion and differentiation as well as apoptosis. When we ordered the DEGs from the meta-analysis by their fold-change direction, pathways for neuro-inflammation such as NF-κB and cytokine storm as well as pro-oncogenic pathways were associated with the up-regulated DEGs. Suppressed pathways associated with the DEGs are e.g., neuroactive ligand-receptor interaction, signal transduction and cellular membrane stability processes. GSEA led to enhanced glutamatergic signal transmission, which in the past has been shown to be associated with enhanced drug seeking (Koob & Volkow, 2010). The NMDA receptor is one of the ionotropic glutamate receptors and is deactivated by acute alcohol consumption (Hoffman et al., 1989), but activated by chronic alcohol exposure and withdrawal (Smothers et al., 1997). Enhanced NMDA-induced calcium invasion is hypothesized to contribute to alcohol-related neuronal loss in the PFC due to increased excitotoxicity (Iorio et al., 1993; Bliss et al., 2014). Up-regulation of the GABA-ergic synapses in our meta-analysis also contributes to previous findings, which state that the GABA(A)-receptors might enhance tolerance to anxiolytic, sedative and positive reinforcing effects during chronic alcohol consumption (Kumar et al., 2004; Kumar et al., 2009). Dopamine (DA) and related signal transmission and pathways have been the scope of alcohol-related research for decades, where it was identified as directly affected by alcohol consumption and these impairments of the DA system were associated with impaired cognitive function due to alcohol addiction

(Trantham-Davidson & Chandler, 2015). In this scenario, our results point towards increased dopaminergic synaptic transmission in AUD patients (Suppl. Tab. 6). Furthermore, we were also able to obtain findings that were not studied before: we observed down-regulation of steroid-related metabolic pathways and receptor binding. Neurosteroids have potentially inhibitory functions leading to anxiolytic, neuroprotective, memory-related and cognitive effects (Baulieu, 1991; Rupprecht & Holsboer, 1999; Reddy, 2010). Impairment of neurosteroids in the PFC have been associated with mood disorders (Qi et al., 2018), particularly in depression (Romeo et al., 1998). Even though neurosteroids have not been studied extensively in the field of AUD, we believe that they represent potential targets for future research, especially when considering the high proportion of AUD patients with additional psychiatric comorbidities (Swendsen and Merikangas, 2000).

As already found in the rodent data, an elevated risk for cancer development has been confirmed also in the human analysis. In addition, several studies have shown that the immune system is strongly impaired by long-term alcohol consumption and neuro-inflammation has been stated as a risk factor for additional psychiatric disorders in AUD patients (Erickson et al., 2019). Also, NF-κB and cytokine storm-related genes have previously been described to be enhanced during chronic alcohol consumption (Pascual et al., 2021; Mayfield et al., 2013).

In human post-mortem AUD tissue, we found cell type-specific enrichment for oligodendrocytes, which is supporting previous findings from other labs (Lewohl et al., 2000; Mayfield et al., 2002; Brenner et al., 2020). Impairment of oligodendrocytes in human AUD patients has been associated with reduced white matter volume and cognitive malfunction (Xin et al., 2020). This cell type is mainly regulating myelinating processes. In terms of AUD,

myelin damage and impairment of oligodendrocytes have been identified in the past and the potential correlating risk for AUD patients to develop comorbid neurodegenerative diseases such as Parkinson's Disease or Alzheimer's Disease has been discussed (Rice & Gu, 2019; Guo et al., 2021).

In this comparative study, we aimed to observe potential commonalities between the post-dependent animal model of alcohol dependence (chronic intermittent ethanol exposure; CIE) and the patient situation, where we found six significant DEGs overlapping in the direct comparison: Adcy3, Hsd11b1, Itm2c, Oga, Rab26 and Rnf122. Among these six genes, Hsd11b1 showed the highest significance in the comparative approach and is also appearing as the most significantly down-regulated gene in the cross-species data integrative approach. Hsd11b1 is encoding a protein that is involved in the catabolism of cortisol. Down-regulation of this gene might lead to an increase in cortisol levels and therefore potentially induce chronic stress reaction. Furthermore, the transcript encoding CRH (Corticotropin-releasing hormone) was significantly downregulated in human PFC and also in the integrative cross-species meta-analysis. CRH is the central activator of the HPA (hypothalamic-pituitary-adrenal) axis, which is the main regulatory system of the body's response to stress. Accumulating evidence confirms an increase of CRF in the brain after acute alcohol exposure. The effect of chronic alcohol exposure has not been so clear, but several studies support decrease of CRF mRNA in the brain after chronic alcohol exposure (Schreiber and Gilpin, 2018). The current study supports the cross-species effect of decreased CRF in the PFC after chronic alcohol exposure that might be resulted from negative feedback of chronically increased cortisol in long-term excess alcohol use. Since the PFC is generally known to be a stress-sensitive brain region, these dysregulations might even

have a broader impact on the whole individual. Stress and increased cortisol levels are implemented in the process of establishing AUD and are also one of the key aspects of acute and long-term withdrawal (Anderson et al., 2016; Becker, 2017; Heilig & Koob, 2007). Previous studies have suggested that increased cortisol concentrations are induced by increased cortisol biosynthesis (Lu & Richardson, 2014), whereas our results suggest a new pathway of escalated alcohol consumption leading to reduced cortisol catabolism via reduced Hsd11b1 presence and an increased cortisol level as a consequence.

On a genome-wide level, human GWAS studies support significant association between excessive alcohol use and the corticotropin receptor gene *Crhr1* (Sanchez-Roige et al., 2018; Gelernter et al, 2019; Brazel et al, 2019) but no GWAS evidence for alcohol use has been currently found for the Hsd11b1 gene in these recent high throughput studies. However, an SNP in the promoter region of the Hsd11b1 gene, rs10779511, is significantly associated with Aspartate aminotransferase (AST) levels (Sakaue et al, 2021). Elevated AST levels in the serum, have been shown to be one of the strongest diagnostic markers for history of excess alcohol consumption as elevated AST to alanine transferase (ALT) ratio is a useful index for distinguishing nonalcoholic steatohepatitis from alcoholic liver disease (Sorbi et al, 1999). Also, in the liver of AUD patients, down-regulation of Hsd11b and its analogs has been shown to induce a hypercortisolism state, which was suggested to be associated with alcoholic liver disease (Ahmed et al., 2008).

Adcy3 encodes for adenylate cyclase 3 and has previously been found to be strongly associated with obesity (Wu et al., 2016). In addition, it was suggested to have a regulatory function in the immune system, as it has been found to be dysregulated in systemic inflammatory diseases, such as inflammatory bowel disease and ulcerative colitis (Hulur et al., 2015). Even though, so far there has been found no association of Adcy3 with AUD or

related phenotypes, its importance to obesity and immune regulatory function links to well-known comorbidities of AUD. The finding of Adcy3 dysregulation is also supported by the cross-species data integrative approach, where Adcy3 is significantly up-regulated.

Itm2c encodes for integral membrane protein 2 c, which is inhibiting the proliferation and projection development of neuronal cells. Previous studies have shown interaction of this gene in Alzheimer's Disease (Dolfe et al., 2018) and it seems to be highly expressed in the rodent brain (Choi et al., 2001). However, to our knowledge, this gene and the corresponding protein have not been studied extensively in general and have also not been addressed in the field of alcohol research so far.

The Oga gene – encoding for O-GlcNAcase – has previously been suggested to have an emerging role in several cancer types via an atypical gene expression alteration mechanism (Parker et al., 2021). This finding is contributing to the previously reported increased risk of AUD patients to develop several cancer types (Cao & Giovannucci, 2016), even though this gene has not been found to be associated with alcohol abuse or dependence so far. In brain-specific disorders, Oga has been found to be associated with intelligence (Savage et al., 2018), cognitive performance (Lee et al., 2018), Alzheimer's disease (Kulminski et al., 2022), Educational attainment (Okbay et al., 2016), ADHD and autism spectrum disorder (Rao et al, 2022). Additionally, Oga was also significantly dysregulated in the cross-species integrative meta-analysis, underlining its potentially important role in AUD and the development of common AUD- related comorbidities.

Also the recently discovered gene Rab26, encoding for the Ras-related protein Rab-26, is contributing to the increased risk for cancer development in AUD patients, based on our analysis. Rab26 is a member of the Ras protein family, which is generally categorized as an oncogene family. Ras proteins are generally involved in regulation of endothelial cell

permeability and induction of autophagy and therefore, when dysregulated, contribute to cancer development and progression (Dong et al., 2018).

Rnf122 was characterized as a ubiquitin ligase and therefore it is involved in protein degradation, including its own degradation process. In addition, it is associated with the calcium-modulating cyclophilin ligand (Peng et al., 2010). Previous Genome-wide association study (GWAS) identified SNPs in Rnf122 as susceptibility markers for attention deficit hyperactivity disorder (ADHD) and also found up-regulation of the Rnf122 RNA levels in the respective patient group (Garcia-Martinez et al.,2017). However, based on our literature research it seems like Rnf122 just recently got attention from researchers and therefore this gene also requires further investigation into its involvement in physiological and mental diseases, including AUD.

Even though these genes identified by cross-species overlap were the only ones identically labeled across the species, there are also genes coming from similar gene families that are commonly altered in the rodent as well as the human meta-analysis e.g., up-regulation of genes encoding for integrins and Lancl gene family members.

In the cross-species integrative approach, we aimed to observe the commonly dysregulated transcriptomic changes in a combined analysis. Here, we observed 31 DEGs, which also partially confirmed previously identified DEGs coming from species-specific meta-analyses, considering rodent and human data separately. The increased number of DEGs compared to the comparative approach of the separate species-specific meta-analyses (Fig. 6) could be explained by increased statistical power due to the strongly increased input. While for species-specific meta-analyses seven rodent and eight human datasets were used respectively, in the cross-species approach the input increased to 15 studies in total.

Pathway enrichment analysis pointed towards signal transduction-related pathways, including glutamatergic signaling. Alcohol is known to impact the glutamatergic system and glutamatergic neurons have been suggested as a promising target for psychopharmacotherapy of AUD patients (Holmes et al., 2013; Alasmari et al., 2018).

This translational approach comparing the effects of chronic alcohol consumption on transcriptomic signatures in animal models of alcohol dependence and AUD patient confirms previous findings and also enlightens potentially new targets for further study in the alcohol addiction field. Based on our results that point towards an overlay of transcriptomic signatures between the animal model of alcohol dependence and human AUD brain tissue, we suggest further investigation into stress-related pathways involving Hsd11b1 in the brain. Our results suggest a new mechanism to explain increased cortisol levels in the alcohol addicted phenotype via down-regulation of Hsd11b1. Therefore, manipulation of Hsd11b1 expression and activity in the AUD patient might lead to long-lasting decrease in relapse risk, which so far is the biggest challenge in AUD psychopharmacotherapy.

Funding: Financial support for this work was provided by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) with the Graduiertenkolleg GRK2350/1 to RS, TRR 265 (A05 and B02) to RS (Heinz et al., 2020), SFB1158 (B04) to RS. Also supported by the German Federal Ministry of Education and Research (BMBF), "A systems-medicine approach towards distinct and shared resilience and pathological

mechanisms of substance use disorders" (01ZX01909) (to RS), and the Ministry for Science, Research and Art of Baden-Wuerttemberg (MWK) for the 3R-Center Rhein-Neckar (to RS).

Acknowledgment

The authors thank the CAMARADES research group in Edinburgh for support on the study design and the systematic literature screening process. In addition, we want to thank Iris Reinhardt, Dr. Emanuel Schwarz and Dr. Han Chao for their support on statistical analysis. We also would like to acknowledge Prof. R. Dayne Mayfield for providing very helpful input and contacts. Thanks to Dr. Rick Bernardi for professional English editing. Finally, we are very grateful to all the authors of the original studies that made it possible to include these datasets in our analysis. Without this involvement, it would not have been possible to conduct the study to this extent. The authors report no biomedical financial interests or potential conflict of interest.

Authors contribution

MMF, AMB, FG and RS designed the study. MMF, AMB and FG designed keywords for the systematic literature screening. MMF, ECT, AMB and FG performed systematic literature screening and data extraction. MMF and WHS contacted the authors in case datasets were not published including all necessary information. MMF and ECT designed the analysis pipeline. MMF, ECT and MJWH performed data analysis. ACH and MAP provided previously unpublished data. MAP provided scientific support. MMF, AMB, WHS, and RS wrote the manuscript. All authors have read and approved the manuscript.

Data Availability Statement

The meta-analysis in rodents (Open Science Framework: https://doi.org/10.17605/OSF.IO/ TF8R4) as well as in human post-mortem brain tissue (Prospero: CRD42020192453) has been pre-registered in advance. As mentioned in the methods section, the analysis strategy was deviating from the pre-registered methods, since the obtained data were not enabling us to perform the initially planned random effect size combination.

Datasets as well as analysis pipelines will be publicly available from the day of publication onwards.

References

Abernathy, K., J. Chandler, and J. J. Woodward. "Alcohol and the Prefrontal Cortex." *Int Rev Neurobiol* 91 (2010): 289-320.

Ahmed, A. , S. Saksena, M. Sherlock, S. P. Olliff, E. Elias, and P. M. Steward. "Induction of Hepatic 11beta-Hydroxysteroid Dehydrogenase Type 1 in Patients with Alcoholic Liver Disease." *Clin Endocrinol.* 68, no. 6 (2008): 898-903.

Alasmari, F., S. Goodwani, R. E. McCullumsmith, Y. Sari. "Role of glutamatergic system and mesocorticolimbic circuits in alcohol dependence." *Prog Neurobiol.* 171, no.1 (2018):32-49.

Anderson, R. I. , M. F. Lopez, and H. C. Becker. "Stress-Induced Enhancement of Ethanol Intake in C57bl/6j Mice with a History of Chronic Ethanol Exposure: Involvement of Kappa Opioid Receptors." *Front Cell Neurosci.* 10 (2016): 45.

Avchalumov, Y. , R. J. Oliver, W. Trenet, R. E. H. Osorno, B. D. Sibley, D. C. Purohit, C. Contet, *et al.* "Chronic Ethanol Exposure Differentially Alters Neuronal Function in the Medial Prefrontal Cortex and Dentate Gyrus." *Neuropharmacology* 185, no. 1 (2021): 108438.

Barbier, E., J. D. Tapocik, N. Juergens, C. Pitcairn, A. Borich, J. R. Schank, H. Sun, *et al.* "DNA Methylation in the Medial Prefrontal Cortex Regulates Alcohol-Induced Behavior and Plasticity." *J Neurosci* 35, no. 15 (2015): 6153-6164.

Baulieu, E. E. "Neurosteroids: A New Function in the Brain." *Biol Cell* 71, no. 1-2 (1991): 3-10.

Bechara, A. "Decision Making, Impulse Control and Loss of Willpower to Resist Drugs: A Neurocognitive Perspective." *Nat Neurosci* 8, no. 11 (2005): 1458-63.

Becker, H. C. "Influence of Stress Associated with Chronic Alcohol Exposure on Drinking." *Neuropharmacology* 122 (2017): 115-26.

Blaine, S. K. , V. Milivojevic, H. Fox, and R. Sinha. "Alcohol Effects on Stress Pathways: Impact on Craving and Relapse Risk." *The Canadian Journal of Psychiatry* 61, no. 3 (2016): 145-53.

Brazel, D. M., Y. Jiang, J. M. Hughey, V. Turcot, X. Zhan, et al. "Exome Chip Meta-analysis Fine Maps Causal Variants and Elucidates the Genetic Architecture of Rare Coding Variants in Smoking and Alcohol Use." *Biol Psychiatry.* 85, no. 11 (2019):946-955

Brenner, E. , G. R. Tiwari, M. Kapoor, Y. Liu, Y. Brock, and R. D. Mayfield. "Single Cell Transcriptome Profiling of the Human Alcohol-Dependent Brain." *Hum Mol Genet* 29, no. 7 (2020): 1144-53.

Briggs, F. "Organizing Principles of Cortical Layer 6." *Front Neural Circuits* 12, no. 4 (2010): 3.

Button, K. S. , J. Ioannidis, C. Mokrysz, B. A. Nosek, J. Flint, E. S. Robinson, and M. R. Munafò. "Power Failure: Why Small Sample Size Undermines the Reliability of Neuroscience.". *Nature Reviews Neuroscience* 14, no. 5 (2013): 365-76.

Carlson, M. "Mouse4302.Db: Affymetrix Affymetrix Mouse Genome 430 2.0 Array Annotation Data (Chip Mouse4302). R Package Version 3.13.0.". (2021).

———. "Org.Hs.Eg.Db: Genome Wide Annotation for Human. R Package Version 3.15.0." (2022).

———. "Rae230a.Db: Affymetrix Affymetrix Rae230a Array Annotation Data (Chip Rae230a). R Package Version 3.13.0." *2021*.

Cao, Y., E. L. Giovannucci. "Alcohol as a risk factor for cancer." *Semin Oncol Nurs* 32, no. 3 (2016):325-331.

Catafau, A. M. , A. Etcheberrigaray, J. Perez de los Cobos, M. Estorch, J. Guardia, A. Flotats, L. Berná*, et al.* "Regional Cerebral Blood Flow Changes in Chronic Alcoholic Patients Induced by Naltrexone Challenge During Detoxification." *J Nucl Med* 40, no. 1 (1999): 19-24.

Chai, A. W. Y. , A. K. L. Cheung, W. Dai, J. M. Y. Ko, N. P. Y. Lee, K. T. Chan, S. Y.-K. Law, and M. L. Lung. "Elevated Levels of Serum Nidogen-2 in Esophageal Squamous Cell Carcinoma." *Cancer Biomark. Sect. A Dis. Mark.* 21, no. 3 (2018): 583-90.

Chan, I. S. , C. D. Guy, M. V. Machado, A. Wank, V. Kadiyala, G. Michelotti, S. Choi*, et al.* "Alcohol Activates the Hedgehog Pathway and Induces Related Procarcinogenic Processes in the Alcohol-Preferring Rat Model of Hepatocarcinogenesis." *Alcohol Clin Exp Res* 38, no. 3 (2014): 787-800.

Charlton, A. J. , C. May, S. J. Luikings, E. L. Burrows, J. H. Kim, A. J. Lawrence, and C. J. Perry. "Chronic Voluntary Alcohol Consumption Causes Persistent Cognitive Deficits and Cortical Cell Loss in a Rodent Model." *Sci Rep* 9, no. 1 (2019): 18651.

Choi, S. C. , J. Kim, T. H. Kim, S. Y. Cho, S. S. Park, K. D. Kim, and S. H. Lee. "Cloning and Characterization of a Type Ii Integral Transmembrane Protein Gene, Itm2c, That Is Highly Expressed in the Mouse Brain." *Mol Cells.* 12, no. 3 (2001): 391-97.

Contet, C. "Gene Expression under the Influence: Transcriptional Profiling of Ethanol in the Brain." *Curr Psychopharmacol* 1, no. 4 (2012): 301-14.

Dahchour, A. , and R. J. Ward. "Changes in Brain Dopamine Extracellular Concentration after Ethanol Administration; Rat Microdialysis Studies." *Alcohol and Alcoholism* 57, no. 2 (2022): 165-75.

Davis, B. T. , R. M. Voigt, M. Shaikh, C. B. Forsyth, and A. Keshavarzian. "Circadian Mechanisms in Alcohol Use Disorder and Tissue Injury." *Alcohol Clin Exp Res* 42, no. 4 (2018): 668-77.

Debell, F., N. T. Fear, M. Head, S. Batt-Rawden, N. Greenberg, S. Wessely, L. Goodwin. "A systematic review of the comorbidity between PTSD and alcohol misuse." *Soc Psychiatry Psychiatr Epidemiol.* 49, no. 9 (2014): 1401-1425

de Paiva Lima, C. , D. A. da Silva E Silva, S. Damasceno, A. F. Ribeiro, C. S. Rocha, A. H. B. de Matos, D. Correia, R. Boerngen-Lacerda, and A. L. B. Godard. "Loss of Control over the Ethanol Consumption: Differential Transcriptional Regulation in Prefrontal Cortex." *J Neurogenet* 31, no. 3 (2017): 170-77.

Delatour, L. C. , P. W. L. Yeh, and H. H. Yeh. "Prenatal Exposure to Ethanol Alters Synaptic Activity in Layer V/Vi Pyramidal Neurons of the Somatosensory Cortex." *Cerebral Cortex* 30, no. 3 (2020): 1735-51.

Dong, W., B. He, H. Qian, Q. Liu, D. Wang, J. Li, Z. Wei, Z. Wang, Z. Xu, G. Wu, G. Qian, G. Wang. "RAB26-dependent autophagy protects adherens junctional integrity in acute lung injury." *Autophagy* 14, no. 10 (2018):1677-1692.

Dolfe, L. , S. Tambaro, H. Tigro, M. Del Campo, J. J. M. Hoozemans, B. Wiehager, C. Graff*, et al.* "The Bri2 and Bri3 Brichos Domains Interact Differently with Aβ42 and Alzheimer Amyloid Plaques." *J Alzheimers Dis Rep* 2, no. 1 (2018): 27-39.

Dolgalev, I. "Babelgene: Gene Orthologs for Model Organisms in a Tidy Data Format. R Package Version 22.9." (2022).

Durinck, S., P. Spellman, E. Birney, and W. Huber. "Mapping Identifiers for the Integration of Genomic Datasets with the R/Bioconductor Package Biomart.". *Nature Protocols* 4, no. 1184-1191 (2009).

Erickson, E. K. , E. K. Grantham, A. S. Warden, and R. A. Harris. "Neuroimmune Signaling in Alcohol Use Disorder." *Pharmacol Biochem Behav* 177 (2019): 34-60.

Ericsson, A. C., C. L. Franklin. "The gut microbiome of laboratory mice: considerations and best practices for translational research." Mamm Genome 32, no. 4 (2021): 239-250.

Farris, S. P., D. Arsappan, S. Hunicke-Smith, R. A. Harris, and R. D. Mayfield. "Transcriptome organization for chronic alcohol abuse in human brain." *Mol Psychiatry* 20, no. 11 (2015): 1438-47.

Garcia-Martinez, I. , C. Sánchez-Mora, M. Soler Artigas, P. Rovira, M. Pagerols, M. Corrales, E. Calvo-Sánchez*, et al.* "Gene-Wide Association Study Reveals Rnf122 Ubiquitin Ligase as a Novel Susceptibility Gene for Attention Deficit Hyperactivity Disorder." *Sci Rep* 7, no. 1 (2017): 5407.

Gelernter J., N. Sun, R. Plolimanti, R. H. Rietrzak, D. F. Levey, Q Lu, Y. Hu et al. "Genome-wide Association Study of Maximum Habitual Alcohol Intake in >140,000 U.S. European and African American Veterans Yields Novel Risk Loci." Biol Psychiatry. 86, no. 5 (2019):365-376.

George, O. , and G. F. Koob. "Individual Differences in Prefrontal Cortex Function and the Transition from Drug Use to Drug Dependence." *Neurosci Biobehav Rev.* 35, no. 2 (2010): 232-47.

George, O. , C. Sanders, J. Freiling, E. Grigoryan, S. Vu, C. D. Allen, E. Crawford, C. D. Mandyam, and G. F. Koob. "Recruitment of Medial Prefrontal Cortex Neurons During Alcohol Withdrawal Predicts Cognitive Impairment and Excessive Alcohol Drinking." *PNAS* 109, no. 44 (2012): 18156-61.

Gilpin, N. W., M. A. Herman, M. Roberto. "The central amygdala as an integrative hub for anxiety and alcohol use disorders." *Biol Psychiatry* 77, no. 10 (2015): 859-869.

Gropman, A. L., M. Summar, J. V. Leonard. "Neurological implications of urea cycle disorders." *J Inherit Metab Dis.* 30, no. 6 (2007):865-879.

Guo, F. , Y.-F. Zhang, K. Liu, X. Huang, R.-X. Li, S.-Y. Wang, F. Wang*, et al.* "Chronic Exposure to Alcohol Inhibits New Myelin Generation in Adult Mouse Brain." *Front Cell Neurosci.* 15 (2021): 732602.

Hade, A. C., M. A. Philips, E. Reimann, T. Jagomae, K. L. Eskla, T. Traks, E. Prans, *et al.* "Chronic Alcohol Use Induces Molecular Genetic Changes in the Dorsomedial Thalamus of People with Alcohol-Related Disorders." *Brain Sci* 11, no. 4 (2021): 435.

Hansson, A. C. , G. Gründer, N. Hirth, H. R. Noori, R. Spanagel, and W. H. Sommer. "Dopamine and Opioid Systems Adaptation in Alcoholism Revisited: Convergent Evidence from Positron Emission Tomography and Postmortem Studies." *Neurosci Biobehav Rev.* 106 (2019): 141-64.

He, W. , X. Liu, Z. Luo, L. Li, and X. Fang. "Fgf16 Regulated by Mir-520b Enhances the Cell Proliferation of Lung Cancer." *Open Med* 16, no. 1 (2021): 419-27.

Heilig, M. , E. Augier, S. Pfarr, and W. H. Sommer. "Developing Neuroscience-Based Treatments for Alcohol Addiction: A Matter of Choice?". *Transl Psychiatry* 9, no. 1 (2019): 255.

Heilig, M. , E. Barbier, A. L. Johnstone, J. Tapocik, M. W. Meinhardt, S. Pfarr, ..., and W. H. Sommer. "Reprogramming of Mpfc Transcriptome and Function in Alcohol Dependence." *Genes, brain and behavior* 16, no. 1 (2017): 86-100.

Heilig, M. , and G. F. Koob. "A Key Role for Corticotropin-Releasing Factor in Alcohol Dependence." *Trends Neurosci* 8 (2007): 399-406.

Heilig, M., J. MacKillop, D. Martinez, J. Rehm, L. Leggio, and L. J. M. J. Vanderschuren. "Addiction as a Brain Disease Revised: Why It Still Matters, And. The Need for Consilience." *Neuropharmacology* 46, no. 10 (2021): 1715-23.

Heilig, M., W. H. Sommer, and R. Spanagel. "The Need for Treatment Responsive Translational Biomarkers in Alcoholism Research." *Curr Top Behav Neurosci* 28 (2016): 151-71.

Heinz, A., F. Kiefer, M. N. Smolka, T. Endrass, C. Beste, A. Beck, S. Liu*, et al.* "Addiction Research Consortium: Losing and Regaining Control over Drug Intake (Recode)-from Trajectories to Mechanisms and Interventions." *Addict Biol* 25, no. 2 (2020): e12866.

Helle, A. C., A. L. Watts, T. J. Trull, and K. J. Sher. "Alcohol Use Disorder and Antisocial and Borderline Personality Disorder." *Alcohol Research: Current Reviews* 40, no. 1 (2019).

Hoffman, P. L. , C. S. Rabe, F. Moses, and B. Tabakoff. "N-Methyl-D-Aspartate Receptors and Ethanol: Inhibition of Calcium Flux and Cyclic Gmp Production." *J Neurochem* 52, no. 6 (1989): 1937-40.

Holmes, A, R. Spanagel, J. H. Krystal. "Glutamatergic targets for new alcohol medications." *Psychopharmacology* 229, no. 3 (2013):539-554.

Hulur, I., E. R. Gamazon, A. D. Skol, R. M. Xicola, X. Llor, K. Onel, N. A. Ellis, S. S. Kupfer. "Enrichment of inflammatory bowel disease and colorectal cancer risk variants in colon expression quantitative trait loci." *BMC Genomics* 16, no. 1 (2015): 138

Hyman, S. E. "Revitalizing Psychiatric Therapeutics." *Neuropharmacology* 39, no. 1 (2014): 220-29.

Hyman, S. E. "Revolution Stalled." *Sci Transl Med.* 4, no. 155 (2012): 155cm11.

Kapoor, M., J. Wang, S. P. Farris, Y. Liu, J. McClintick, I. Gupta, J. L. Meyers, *et al.* "Analysis of whole genome-transcriptomic organization in brain to identify genes associated with alcoholism." *Transl Psychiatry* 9, 89 (2019).

Keable, A. , K. Fenna, H. M. Yuen, D. A. Johnston, N. R. Smyth, C. Smith, R. Al-Shahi Salman*, et al.* "Deposition of Amyloid Beta in the Walls of Human Leptomeningeal Arteries in Relation to Perivascular Drainage Pathways in Cerebral Amyloid Angiopathy." *Biochim Biophys Acta.* 1862, no. 5 (2016): 1037-46.

Kirshenbaum, A. P. , D. M. Olsen, and W. K. Bickel. "A Quantitative Review of the Ubiquitous Relapse Curve." *J Subst Abuse Treat* 36 (2009): 8-17.

Kontou, P., A. Pavlopoulou, G. Braliou, S. Bogiatzi, N. Dimou, S. Bangalore, and P. Bagos. "Identification of Gene Expression Profiles in Myocardial Infarction: A Systematic Review and Meta-Analysis." *BMC Medical Genomics* 11, no. 1 (2018): 109.

Koob, G. F., and N. D. Volkow. "Neurocircuitry of Addiction." *Neuropsychopharmacology* 35, no. 1 (2010): 217-38.

Kuk, C., C. G. Gunawardana, A. Soosaipillai, H. Kobayashi, L. Li, Y. Zheng, and E. P. Diamandis. "Nidogen-2: A New Serum Biomarker for Ovarian Cancer." *Clin Biochem.* 438, no. 4-5 (2010): 355-61.

Kuluminski, A. M., E. Loiko, Y. Loika, I. Culminskaya. "Pleiotropic predisposition to Alzheimer's disease and educational attainment: insights from the summary statistics analysis." *Geroscience* 44, no. 1 (2022):265-280

Kumar, S. , R. L. Fleming, and A. L. Morrow. "Ethanol Regulation of Gamma-Aminobutyric Acid a Receptors: Genomic and Nongenomic Mechanisms." *Pharmacol Ther* 101, no. 3 (2004): 211-26.

Kumar, S., R. Porcu, D. F. Werner, D. B. Matthews, J. L. Diaz- Granados, R. S. Helfand, and A. L. Morrow. "The Role of Gaba(a) Receptors in the Acute and Chronic Effects of Ethanol: A Decade of Progress." *Psychopharmacology* 205, no. 4 (2009): 529-64.

Kwako, L. W. , P. A. Spagnolo, M. L. Schwandt, A. Thorsell, D. T. George, R. Momenan, D. E. Rio*, et al.* "The Corticotropin Releasing Hormone-1 (Crh1) Receptor Antagonist Pexacerfont in Alcohol Dependence: A Randomized Controlled Experimental Medicine Study." *Neuropsychopharmacology* 40, no. 5 (2015): 1053-63.

Lee, J. J., R. Wedow, A. Okbay et al. "Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals." *Nat Genet.* 50, no. 8 (2018): 1112-1121.

Lewohl, J. M. , L. Wang, M. F. Miles, L. Zhang, P. R. Dodd, and Harris R. A. "Gene Expression in Human Alcoholism: Microarray Analysis of Frontal Cortex." *Alcohol Clin Exp Res* 12 (2000): 1873-82.

Liu, J., J . M. Lewohl, R. A. Harris, V. R. Iyer, P. R. Dodd, P. K. Randall, and R. D. Mayfield. "Patterns of gene expression in the frontal cortex discriminate alcoholic from nonalcoholic individuals." *Neuropsychopharmacology* 31, no. 7 (2006): 1574-82.

Louth, E. L. , H. D. Luctkar, K. A. Heney, and C. D. C Bailey. "Developmental Ethanol Exposure Alters the Morphology of Mouse Prefrontal Neurons in a Layer-Specific Manner." *Brain Research* 1678 (2018): 94-105.

Lu, Y.-L., H. N. Richardson. "Alcohol, stress hormones, and the prefrontal cortex: A proposed pathway to the dark side of addiction." *Neuroscience* 277, no.1 (2014):139-151

Lynch, W. J., K. L. Nicholson, M. E. Dance, R. W. Morgan, and P. L. Foley. "Animal Models of Substance Abuse and Addiction: Implications for Science, Animal Welfare, and Society." *Comp Med* 60, no. 3 (2010): 177-88.

Ma, W. "Lrcell: Differential Cell Type Change Analysis Using Logistic/Linear Regression. R Package Version 1.2.0." (2021).

Manzardo, A. M., S. Gunewardena, and M. G. Butler. "Over-expression of the miRNA cluster at chromosome 14q32 in the alcoholic brain correlates with suppression of predicted target mRNA required for oligodendrocyte proliferation." *Gene* 526, no. 2 (2013): 356-63.

Margolis, K. G., J. F. Cryan, E. A. Mayer. "The microbiota-gut-brain axis: from motility to mood." Gastroenterology 160, no. 5 (2021): 1486-1501.

Maingret, F., B. Coste, F. Padilla, N. Clerc, M. Crest, S. M. Korogod, and P. Delmas. "Inflammatory Mediators Increase Nav1.9 Current and Excitability in Nociceptors

through a Coincident Detection Mechanism." *J Gen Physiol.* 131, no. 3 (2008): 211-25.

Mayernhoff, D. J. , and T. C. Durazzo. "Predicting Relapse and Treatment Outcome in Alcohol Use Disorders: Time for an Integrative Biopsychosocial Approach." *Alcohol Clin Exp Res* 34 (2010): 289A-90A.

Mayfield, J. , and R. A. Harris. "The Neuroimmune Basis of Excessive Alcohol Consumption." *Neuropsychopharmacology* 42, no. 2 (2017): 376.

Mayfield, J., L. Ferguson, and R. A. Harris. "Neuroimmune Signaling: A Key Component of Alcohol Abuse." *Curr Opin Neurobiol.* 23, no. 4 (2013): 513-20.

Mayfield, R. D. , J. M. Lewohl, P. R. Dodd, A. Herlihy, J. Liu, and Harris R. A. "Patterns of Gene Expression Are Altered in the Frontal and Motor Cortices of Human Alcoholics." *J Neurochem* 81, no. 4 (2002): 802-13.

McHugh, R. K., and R. D. Weiss. "Alcohol Use Disorder and Depressive Disorders." *Alcohol Research: Current Reviews* 40, no. 1 (2019).

Meinhardt, M. W., and W. H. Sommer. "Postdependent State in Rats as a Model for Medication Development in Alcoholism." *Addiction Biology* 20, no. 1 (2015): 1-21.

Meinhardt, M. W., A. C. Hansson, S. Perreau-Lenz, Bauder-Wenz, O. Staehlin, M. Heilig, C. Harper, *et al.,* "Rescue of Infralimbic mGlur2 Deficit Restores Control Over Drug-Seeking Behavior in Alcohol Dependence." *J Neurosci* 33, no. 7 (2013): 2794-2806.

Miller, J. A., C. Cai, P. Langfelder, D. H. Geschwind, S. M. Kurian, D. R. Salomon, S. Horvath. „Strategies for aggregating gene expression data: The collapseRows R function." BMC Bioinformatics, no. 12 (2011): 322.

Muench, C. , M. Schwandt, J. Jung, C. R. Cortes, R. Momenan, and F. W. Lohoff. "The Major Depressive Disorder Gwas-Supported Variant Rs10514299 in Tmem161b-Mef2c Predicts Putamen Activation During Reward Processing in Alcohol Dependence." *Transl. Psychiatry* 8, no. 131 (2018).

Nutt, D. J. , L. A. King, and L. D. Phillips. "Drug Harms in the Uk: A Multicriteria Decision Analysis." *The Lancet* 376, no. 9752 (2010): 1558-65.

Okbay A., J. P. Beauchamp, M. A. Fotana, J. J. Lee, T. H. Pers et al. "Genome-wide association study identifies 74 loci associated with educational attainment." *Nature* 533, no. 7604 (2016):539-542

Osterndorff-Kahanek, E. A., H. C. Becker, M. F. Lopez, S. P. Farris, G. R. Tiwari, Y. O. Nunez, A. R. Harris, and R. D. Mayfield. "Chronic Ethanol Exposure Produces Time- and Brain Region-Dependent Changes in Gene Coexpression Networks." *PLoS ONE* 10, no. 3: e0121522.

Page, M. J. , J. E. McKenzie, P. M. Bossuyt, I. Boutron, T. C. Hoffmann, C. D. Mulrow, L. Shamseer, *et al.* "The Prisma 2020 Statement: An Updated Guideline for Reporting Systematic Reviews." *BMJ* 372 (2021): n71.

Pahng, A. R. , M. A. McGinn, R. I. Paulsen, and S. Edwards. "The Prefrontal Cortex as a Critical Gate of Negative Affect and Motivation in Alcohol Use Disorder." *Curr Opin Behav Sci.* (2017).

Pahng, A. R. , M. A. McGinn, R. I. Paulsen, and S. Edwards. "The Prefrontal Cortex as a Critical Gate of Negative Affect and Motivation in Alcohol Use Disorder." *Curr Opin Behav Sci.* (2017).

Parker, M. P., K. R. Peterson, C. Slawson. "O-GlcNAcylation and O-GlcNAc Cycling Regulate Gene Transcription: Emerging Roles in Cancer." *Cancers* 13, no. 7 (2021):1666.

Pascual, M., M. Calvo-Rodriguez, L. Núñez, C. Villalobos, J. Urena, and C. Guerri. "Toll-Like Receptors in Neuroinflammation, Neurodegeneration, and Alcohol-Induced Brain Damage." *IUBMB Life* 73, no. 7 (2021): 900-15.

Peng, Z. , T. Shi, and D. Ma. "Rnf122: A Novel Ubiquitin Ligase Associated with Calcium-Modulating Cyclophilin Ligand." *BMC Cell Biol.* 11 (2010): 41.

Perreau-Lenz, S. , and R. Spanagel. "Clock Genes × Stress × Reward Interactions in Alcohol and Substance Use Disorders." *Alcohol* 49, no. 4 (2015): 351-57.

Ponomarev, I., S. Wang, L. Zhang, R. A. Harris, and R. D. Mayfield. "Gene Coexpression Networks in Human Brain Identify Epigenetic Modifications in Alcohol Dependence." *J Neurosci* 32, no. 5 (2012): 1884-1897.

Qi, X., S. Luchetti, R. W. H. Verwer, A. A. Sluiter, M. R. J. Mason, J. Zhou, and D. F. Swaab. "Alterations in the Steroid Biosynthetic Pathways in the Human Prefrontal Cortex in Mood Disorders: A Post-Mortem Study." *Bran Pathol* 28, no. 4 (2018): 536-47.

Rainer, J. . "Ensdb.Hsapiens.V86: Ensembl Based Annotation Package. R Package Version 2.99.0." (2017).

Rao, S., A. Baranova, Y. Yao, J. Wang, F. Zhang. "Genetic Relationships between Attention-Deficit/Hyperactivity Disorder, Autism Spectrum Disorder, and Intelligence." *Neuropsychobiology* 28, no. 1 (2022):1-13.

Rao, X., K. S. Thapa, A. B. Chen, H. Lin, H. Gao, J. L. Reiter, K. A. Hargreaves, *et al.* "Allele-specific expression and high-throughput reporter assays reveal functional genetic variants associated with alcohol use disorders." *Mol Psychiatry* 26, no. 4 (2021): 1142-1151.

Raudvere U., L. Kolberg, I. Kuzmin, T. Arak, P. Adler, H. Peterson, J. Vilo. "g:Profler: a web server for functional enrichment analysis and conversions of gene lists (2019 update)". *Nucleic Acids Research* 47, no. W1 (2019):W191-W198.

Reddy, D. S. "Neurosteroids: Endogenous Role in the Human Brain and Therapeutic Potentials." *Prog Brain Res* 186 (2010): 113-37.

Rice, J., and C. Gu. "Function and Mechanism of Myelin Regulation in Alcohol Abuse and Alcoholism." *BioEssays* 41, no. 7 (2019): e1800255.

Rocco, A. , D. Compare, D. Angrisani, M. Sanduzzi Zamparelli, and G. Nardone. "Alcoholic Disease: Liver and Beyond." *World J Gastroenterol.* 20, no. 40 (2014): 14652-59.

Romeo, E. , A. Stroehle, G. Spalletta, F. di Michele, B. Hermann, F. Holsboer, A. Pasini, and R. Rupprecht. "Effects of Antidepressant Treatment on Neuroactive Steroids in Major Depression." *Am J Psychiatry* 155, no. 7 (1998): 910-13.

Rumgay, H. , N. Murphy, P. Ferrari, and I. Soerjomataram. "Alcohol and Cancer: Epidemiology and Biological Mechanisms." *Nutrients* 13, no. 9 (2021): 3173.

Rupprecht, R. , and F. Holsboer. "Neuroactive Steroids: Mechanisms of Action and Neuropsychopharmacological Perspectives." *Trends Neurosci* 22, no. 9 (1999): 410-16.

Sakaue S., M. Kanai, Y. Tanigawa et al., "A cross-population atlas of genetic associations for 220 human phenotypes." *Nat Genet.* 53, no. 10 (2021):1415-1424.

Sanchez-Roige, S., A. A. Palmer, P. Fontanillas, S. L. Elson et al. "Genome-Wide Association Study Meta-Analysis of the Alcohol Use Disorders Identification Test (AUDIT) in Two Population-Based Cohorts." *Am J Psychiatry.* 176, no. 2 (2019):107-118.

Savage J. E., P. R. Jansen, S. Stringer, et al., "Genome-wide association meta-analysis in 269,867 individuals identifies new genetic and functional links to intelligence." Nat Genet. 50, no. 7 (2018):912-919

Schacht, J. P. , R. F. Anton, and H. Myrick. "Functional Neuroimaging Studies of Alcohol Cue Reactivity: A Quantitative Meta-Analysis and Systematic Review." *Addiction Biology* 8, no. 1 (2017): 121-33.

Schall, P. Z., K. E. Latham. "Cross-species meta-analysis of transcriptome changes during the morula-to-blastocyst transition: metabolic and physiological changes take center stage." Am J Physiol Cell Physiol. 321, no. 6 (2021):C913-C931.

Schreiber A. L., N. W. Gilpin. Coricotropin-Releasing Factor (CRF) Neurocircuitry and Neuropharmacology in Alcohol Drinking. Handb Exp Pharmacol. 248, no. 1 (2018):435-471.

Schwandt, M. L. , C. R. Cortes, L. E. Kwako, D. T. George, R. Momenan, R. Sinha, D. E. Grigoriadis*, et al.* "The Crf1 Antagonist Verucerfont in Anxious Alcohol-Dependent Women: Translation of Neuroendocrine, but Not of Anti-Craving Effects ". *Neuropsychopharmacology* 41, no. 12 (2016): 2818-29.

Seitz, H. K. , and F. Stickel. "Molecular Mechanisms of Alcohol-Mediated Carcinogenesis." *Nature Reviews Cancer* 7, no. 8 (2007): 599-612.

Sena, E. S., G. L. Currie, S. K. McCann, M. R. Macleod, and D. W. Howells. "Systematic Reviews and Meta-Analysis of Preclinical Studies: Why Perform Them and How to Appraise Them Critically.". *Journal of Cerebral Blood Flow & Metabolism* 34, no. 5 (2014): 737-42.

Seo, D. , and R. Sinha. "Neuroplasticity and Predictors of Alcohol Recovery." *Alcohol Res* 37, no. 1 (2015): 143-52.

Smith, M. L., M. F. Lopez, A. R. Wolen, H. C. Becker, and M. F. Miles. "Brain regional gene expression network analysis identifies unique interactions between chronic ethanol exposure and consumption." *PLoS ONE* 15, no. 5 (2020): e0233319.

Smith, M. L., M. F. Lopez, K. J. Archer, A. R. Wolen, H. C. Becker, and M. F. Miles. "Time-Course Analysis of Brain Regional Expression Network Responses to Chronic Intermittent Ethanol and Withdrawal: Implications for Mechanisms Underlying Excessive Ethanol Consumption." *PLos ONE* 11, no. 1 (2016): e0146257.

Smothers, C. T. , J. J. Mrotek, and D. M. Lovinger. "Chronic Ethanol Exposure Leads to a Selective Enhancement of N-Methyl-D-Aspartate Receptor Function in Cultured Hippocampal Neurons." *J Pharmacol Exp Ther* 283, 3, no. 1214-1222 (1997).

Sorbi, D., J. Boynton, K. D. Lindor. "The Ratio of Aspartate Aminotransferase to Alanine Aminotransferase: Potential Value in Differentiating Nonalcoholic Steatohepatitis from Alcoholic Liver Disease." *Am J of Gastroenterol.* 94, no. 4 (1999):1018-1022.

Spanagel, R., H. R. Noori, M. Heilig. "Stress and alcohol interactions: animal studies and clinical significance." *Trends Neurosci.* 37, no. 4 (2014):219-227.

Spence, J. P. , D. Lai, A. Shekar, L. G. Carr, T. Foroud, and T. Liang. "Quantitative Trait Locus for Body Weight Identified on Rat Chromosome 4 in Inbred Alcohol-

Preferring and -Nonpreferring Rats: Potential Implications for Neuropeptide Y and Corticotrophin Releasing Hormone 2." *Alcohol* 47, no. 1 (2013): 63-67.

Svendsen, J. D. , and K. R. Merikangas. "The Comorbidity of Depression and Substance Use Disorders." *Clin Psychol Rev* 20, no. 2 (2000): 173-89.

Timme, N. M. , B. Ma, D. Linsenbardt, E. Cornwell, T. Galbari, and C. C. Lapish. "Compulsive Alcohol Drinking in Rodents Is Associated with Altered Representations of Behavioral Control and Seeking in Dorsal Medial Prefrontal Cortex." *Nat Commun* 13, no. 1 (2022): 3990.

Toro-Domínguez, D. , J. A. Villatoro-García, J. Martorell-Marugán, Y. Román-Montoya, M. E. Alarcón-Riquelme, R. Carmona-Sáez. "A survey on gene expression meta-analysis: methods and applications." *Briefings in Bioinformatics* 22, no. 2 (2021):1694-1705.

Trantham-Davidson, H. , and L. J. Chandler. "Alcohol-Induced Alterations in Dopamine Modulation of Prefrontal Activity." *Alcohol* 49, no. 8 (2015): 773-79.

Tsermpini, E. E., A. P. Ilješ, and V. Dolžan. "Alcohol-Induced Oxidative Stress and the Role of Antioxidants in Alcohol Use Disorder: A Systematic Review." *Antioxidants* 11, no. 7 (2022): 1374.

Vasudevan, A. , M. S. P. Ho, M. Weiergräber, R. Nischt, T. Schneider, A. Lie, N. Smyth, and R. Köhling. "Basement Membrane Protein Nidogen-1 Shapes Hippocampal Synaptic Plasticity and Excitability." *Hippocampus* 20, no. 5 (2010): 608-20.

Villatoro-García, J. , and P. Carmona-Sáez. "Dexma: Differential Expression Meta-Analysis. R Package Version 1.4.4.". (2022).

Wang, F., J. Gelernter, and H. Zhang. "Differential Expression of miR-130a in Postmortem Prefrontal Cortex of Subjects with Alcohol Use Disorders." *J Addict Res Ther* 4, no. 155 (2013): 18179.

Wang, W. , M. Wang, Y. Xiao, Y. Wang, L. Ma, L. Guo, X. Wu, X. Lin, and P. Zhang. "Usp35 Mitigates Endoplasmic Reticulum Stress-Induced Apoptosis by Stabilizing Rrbp1 in Non-Small Cell Lung Cancer." *Mol Oncol.* 16, no. 7 (2022): 1572-90.

Witkiewitz, K. "Predictors of Heavy Drinking During and Following Treatment." *Psychol Addict Behav* 25 (2011): 426-38.

Witkiewitz, K. , Z. Littenand, and L. Leggio. "Advances in the Science and Treatment of Alcohol Use Disorder." *Science Advances* 5 (2019): eaax4043.

Wu, L., C. Shen, M. S. Ahmed, C.-G. Östenson, H. F. Gu. "Adenylate cyclase 3: a new target for anti-obesity drug development." *Obes Rev.* 17, no. 9 (2016): 907-914.

Wu, T., E. Hu, S. Xu, M. Chen, P. Guo, Z. Dai, T. Feng, *et al.* "Clusterprofiler 4.0: A Universal Enrichment Tool for Interpreting Omics Data.". *The Innovation* 2, no. 3 (2021): 100141.

Yang, L. , L. Li, H. Tang, T. Ma, Y. Li, X. Zhang, X. Shi, and J. Y. Liu. "Alcohol-Aggravated Episodic Pain in Humans with Scn11a Mutation and Aldh2 Polymorphism." *Pain* 161, no. 7 (2020): 1470-82.

Xin, W., and J. R. Chan. "Myelin Plasticity: Sculpting Circuits in Learning and Memory." *Nat Rev Neurosci.* 21, no. 12 (2020): 682-94.

Yu, Z.-H. , Y.-M. Wang, Y.-Z. Jiang, S.-J. Ma, Q. Zhoung, Y.-Y. Wan, and X.-W. Wang. "Nid2 Can Serve as a Potential Prognosis Prediction Biomarker and Promotes the Invasion and Migration of Gastric Cancer." *Pathol Res Pract.* 15, no. 10 (2019): 152553.

Zhang, L. , L. Zhou, M. Shi, Y. Kuang, and L. Fang. "Downregulation of Mirna-15a and Mirna-16 Promote Tumor Proliferation in Multiple Myeloma by Increasing Cabin1 Expression." *Oncol Lett.* 15, no. 1 (2018): 1287-96.

Zhang, Z.-J. , K.-P. Wang, J.-P. Huang, C. Jin, H. Jiang, L. Xiong, Z.-Y. Chen, *et al.* "Comprehensive Analysis of the Potential Immune-Related Biomarker Atg101 That Regulates Apoptosis of Cholangiocarcinoma Cells after Photodynamic Therapy." *Front Pharmacol.* 3, no. 13 (2022): 857774.

GO:MF

Term name	Term ID	p_{adj}	$-\log_{10}(p_{adj})$
myosin VI light chain binding	GO:0070858	2.858×10^{-2}	
corticotropin-releasing hormone activity	GO:0017045	2.858×10^{-2}	
proteoglycan sulfotransferase activity	GO:0050698	2.858×10^{-2}	
7-beta-hydroxysteroid dehydrogenase (NADP+) activity	GO:0047022	2.858×10^{-2}	
oxytocin receptor activity	GO:0004990	2.858×10^{-2}	
11-beta-hydroxysteroid dehydrogenase (NADP+) activity	GO:0070524	2.858×10^{-2}	
[protein]-3-O-(N-acetyl-D-glucosaminyl)-L-serine O-N...	GO:0102167	2.858×10^{-2}	
[protein]-3-O-(N-acetyl-D-glucosaminyl)-L-threonine ...	GO:0102166	2.858×10^{-2}	
cortisol dehydrogenase activity	GO:0102196	2.858×10^{-2}	
[protein]-3-O-(N-acetyl-D-glucosaminyl)-L-serinal]-th...	GO:0102571	2.858×10^{-2}	
11-beta-hydroxysteroid dehydrogenase [NAD(P)] activity	GO:0003845	3.174×10^{-2}	
beta-N-acetylglucosaminidase activity	GO:0016231	3.174×10^{-2}	
formyltetrahydrofolate dehydrogenase activity	GO:0016155	3.174×10^{-2}	
chondroitin 6-sulfotransferase activity	GO:0008459	3.174×10^{-2}	
chylomicron binding	GO:0071748	3.174×10^{-2}	
beta-3 adrenergic receptor binding	GO:0031699	3.174×10^{-2}	
myosin VI binding	GO:0070853	3.174×10^{-2}	
ornithine decarboxylase activity	GO:0004586	3.174×10^{-2}	
signaling receptor binding	GO:0005102	3.772×10^{-2}	
aldehyde dehydrogenase (NADP+) activity	GO:0033721	4.078×10^{-2}	
RNA polymerase I core binding	GO:0001042	4.078×10^{-2}	
corticotropin-releasing hormone receptor 2 binding	GO:0043431	4.389×10^{-2}	
corticotropin-releasing hormone receptor 1 binding	GO:0051430	4.389×10^{-2}	
thrombospondin receptor activity	GO:0070053	4.389×10^{-2}	
calcium- and calmodulin-responsive adenylate cyclase ...	GO:0008294	4.389×10^{-2}	
chondroitin sulfotransferase activity	GO:0070326	4.389×10^{-2}	
lipoprotein lipase activity	GO:0004465	4.752×10^{-2}	
chondroitin sulfotransferase activity	GO:0034481	4.752×10^{-2}	
G protein-coupled receptor binding	GO:0001664	4.752×10^{-2}	
heparan sulfate binding	GO:1904399	4.752×10^{-2}	
ErbB-3 class receptor binding	GO:0043125	4.885×10^{-2}	
N-acetylglucosamine 6-O-sulfotransferase activity	GO:0001517	4.885×10^{-2}	
beta-N-acetylhexosaminidase activity	GO:0004563	4.885×10^{-2}	
corticotropin-releasing hormone receptor binding	GO:0051429	4.885×10^{-2}	
neuregulin binding	GO:0038132	4.885×10^{-2}	

1 to 35 of 35

GO:BP

Term name	Term ID	p_{adj}	$-\log_{10}(p_{adj})$
estrous cycle	GO:0044849	1.537×10^{-3}	
learning or memory	GO:0007611	1.537×10^{-3}	
cognition	GO:0050890	1.546×10^{-3}	
phosphorus metabolic process	GO:0006793	1.626×10^{-3}	
phosphate-containing compound metabolic process	GO:0006796	1.626×10^{-3}	
regulation of behavior	GO:0050795	3.572×10^{-3}	
regulation of protein metabolic process	GO:0051246	3.572×10^{-3}	
behavior	GO:0007610	3.572×10^{-3}	
negative regulation of multicellular organismal process	GO:0051241	4.909×10^{-3}	
memory	GO:0035296	5.178×10^{-3}	
tube development	GO:0035295	5.178×10^{-3}	
biological regulation	GO:0065007	5.178×10^{-3}	
regulation of protein modification process	GO:0031399	5.935×10^{-3}	
organonitrogen compound metabolic process	GO:1901564	6.314×10^{-3}	
multicellular organismal process	GO:0032501	7.133×10^{-3}	
cellular cycle	GO:0042698	1.382×10^{-2}	
regulation of catalytic activity	GO:0050790	1.382×10^{-2}	
glucocorticoid biosynthetic process	GO:0006704	1.382×10^{-2}	
positive regulation of protein modification process	GO:0031401	1.382×10^{-2}	
regulation of biological quality	GO:0065008	1.385×10^{-2}	
transport	GO:0006810	1.385×10^{-2}	
positive regulation of protein metabolic process	GO:0051247	1.385×10^{-2}	
NADPH regeneration	GO:0006740	1.385×10^{-2}	
glial cell differentiation	GO:0010001	1.385×10^{-2}	
regulation of multicellular organismal process	GO:0051239	1.385×10^{-2}	
regulation of norepinephrine secretion	GO:0014061	1.385×10^{-2}	
N-acetylglucosamine metabolic process	GO:0006044	1.385×10^{-2}	
multicellular organismal response to stress	GO:0033555	1.385×10^{-2}	
norepinephrine secretion	GO:0048243	1.385×10^{-2}	
dentate gyrus development	GO:0021542	1.385×10^{-2}	
digestive system process	GO:0022600	1.385×10^{-2}	
positive regulation of neuron migration	GO:2001234	1.527×10^{-2}	
regulation of biological process	GO:0050789	1.527×10^{-2}	
associative learning	GO:0008306	1.595×10^{-2}	
regulation of molecular function	GO:0065009	1.595×10^{-2}	
establishment of localization	GO:0051234	1.601×10^{-2}	
digestion	GO:0007586	1.743×10^{-2}	
glucocorticoid metabolic process	GO:0008211	1.743×10^{-2}	
limbic system development	GO:0021761	1.775×10^{-2}	
negative regulation of T cell activation	GO:0050868	1.820×10^{-2}	
norepinephrine transport	GO:0015874	1.820×10^{-2}	
protein metabolic process	GO:0019538	1.820×10^{-2}	
response to ketone	GO:1901654	1.820×10^{-2}	
glucosamine-containing compound metabolic process	GO:1901071	1.864×10^{-2}	
positive regulation of ion transport	GO:0043270	1.910×10^{-2}	
regulation of NMDA receptor activity	GO:2000310	1.913×10^{-2}	
negative regulation of leukocyte cell-cell adhesion	GO:1903038	2.097×10^{-2}	
adrenal gland development	GO:0030325	2.097×10^{-2}	
animal organ development	GO:0048513	2.103×10^{-2}	
homeostatic process	GO:0042592	2.274×10^{-2}	
regulation of protein modification process	GO:1901137	2.294×10^{-2}	
gliogenesis	GO:0042063	2.294×10^{-2}	
positive regulation of secretion by cell	GO:1903532	2.294×10^{-2}	
response to pain	GO:0048265	2.458×10^{-2}	
acylglycerol homeostasis	GO:0055090	2.458×10^{-2}	
negative regulation of lymphocyte activation	GO:0051250	2.458×10^{-2}	
positive regulation of metabolic process	GO:0009893	2.458×10^{-2}	
locomotion	GO:0040011	2.458×10^{-2}	
organophosphate metabolic process	GO:0019637	2.458×10^{-2}	
triglyceride homeostasis	GO:0070328	2.458×10^{-2}	
positive regulation of synaptic transmission, glutamat...	GO:0051968	2.488×10^{-2}	
response to stimulus	GO:0050896	2.488×10^{-2}	
regulation of phosphate metabolic process	GO:0019220	2.539×10^{-2}	
regulation of phosphorus metabolic process	GO:0051174	2.539×10^{-2}	
positive regulation of motor neuron migration	GO:1905485	2.582×10^{-2}	
regulation of motor neuron migration	GO:1905463	2.582×10^{-2}	
regulation of lateral motor column neuron migr...	GO:2000278	2.582×10^{-2}	
positive regulation of heat generation	GO:0031651	2.582×10^{-2}	
regulation of lateral motor column neuron migration	GO:1902076	2.582×10^{-2}	
positive regulation of secretion	GO:0051047	2.582×10^{-2}	
regulation of transport	GO:0051050	2.582×10^{-2}	
obsolete regulation of histone phosphorylation	GO:0033127	2.582×10^{-2}	
carbohydrate derivative metabolic process	GO:1901135	2.582×10^{-2}	
positive regulation of biological process	GO:0048518	2.582×10^{-2}	
response to stress	GO:0006950	2.582×10^{-2}	
regulation of locomotion	GO:0040012	2.582×10^{-2}	
regulation of primary metabolic process	GO:0080090	2.582×10^{-2}	
protein unfolding	GO:0043335	2.582×10^{-2}	
learning	GO:0007612	2.637×10^{-2}	
developmental process	GO:0032502	2.651×10^{-2}	
NADP metabolic process	GO:0006739	2.692×10^{-2}	
amino sugar metabolic process	GO:0006040	2.692×10^{-2}	
localization	GO:0051179	2.692×10^{-2}	
negative regulation of leukocyte activation	GO:0002695	2.719×10^{-2}	
forebrain development	GO:0030900	2.745×10^{-2}	
organic substance transport	GO:0071702	2.764×10^{-2}	
anatomical structure development	GO:0048856	2.772×10^{-2}	
regulation of ion transport	GO:0043269	2.789×10^{-2}	
cell junction assembly	GO:0034329	2.814×10^{-2}	
cell junction organization	GO:0034330	2.844×10^{-2}	
negative regulation of T cell differentiation	GO:0045581	2.844×10^{-2}	
negative regulation of cell-cell adhesion	GO:0071636	2.855×10^{-2}	
regulation of protein phosphorylation	GO:0001932	2.855×10^{-2}	
behavioral fear response	GO:0001662	3.083×10^{-2}	
regulation of neuron migration	GO:2001222	3.059×10^{-2}	
behavioral defense response	GO:0002209	3.083×10^{-2}	
regulation of cell activation	GO:0050866	3.083×10^{-2}	
regulation of digestive system process	GO:0044058	3.083×10^{-2}	
positive regulation of protein phosphorylation	GO:0001934	3.083×10^{-2}	

1 to 100 of 246

GO:CC

Term name	Term ID	p_{adj}	$-\log_{10}(p_{adj})$
microvillus	GO:0005902	1.152×10^{-3}	
actin-based cell projection	GO:0098858	1.396×10^{-2}	
endomembrane system	GO:0012505	2.135×10^{-2}	
early endosome	GO:0005769	3.058×10^{-2}	
ricin complex	GO:0101767	3.058×10^{-2}	
cytoplasm	GO:0005737	3.058×10^{-2}	
integral component of plasma membrane	GO:0005887	3.101×10^{-2}	
intrinsic component of plasma membrane	GO:0031226	3.285×10^{-2}	
cell projection	GO:0042995	3.285×10^{-2}	
apical part of cell	GO:0045177	3.496×10^{-2}	
ERBB3:ERBB2 complex	GO:0038143	3.496×10^{-2}	
extracellular region	GO:0005576	3.593×10^{-2}	
ciliary membrane	GO:0060170	3.593×10^{-2}	
plasma membrane region	GO:0098590	4.265×10^{-2}	
basal plasma membrane	GO:0009925	4.717×10^{-2}	
cell periphery	GO:0071944	4.717×10^{-2}	
plasma membrane bounded cell projection	GO:0120025	4.717×10^{-2}	
basal part of cell	GO:0045178	4.717×10^{-2}	

1 to 18 of 18

REAC

Term name	Term ID	p_{adj}	$-\log_{10}(p_{adj})$
Syndecan interactions	REAC:R-HSA-...	2.832×10^{-2}	
Assembly of active LPL and LIPC lipase complexes	REAC:R-HSA-...	2.832×10^{-2}	

1 to 2 of 2

CORUM

Term name	Term ID	p_{adj}	$-\log_{10}(p_{adj})$
S4-synectin-RhoGDI1 complex	CORUM:6227	5.566×10^{-2}	

1 to 1 of 1