Movement of large African savanna herbivores is shaped by intricate interactions

between behavioural modes and the spatial-temporal environment

Hans Linssen[1,2,3], Henrik J. de Knegt[1] & Jasper A.J. Eikelboom[1]

[1]Wildlife Ecology and Conservation Group, Wageningen University and Research, Droevendaalsesteeg

3a, 6708 PB, Wageningen, The Netherlands

[2]Theoretical and Computational Ecology Group, Institute for Biodiversity and Ecosystem Dynamics,

University of Amsterdam, P.O. Box 94240, 1090 GE, Amsterdam, The Netherlands

[3]Department of Animal Ecology, Netherlands Institute of Ecology, Droevendaalsesteeg 10, 6708 PB,

Wageningen, The Netherlands

Correspondence: Hans Linssen h.j.linssen@uva.nl

Abstract

1. Animal movement arises from a complex interaction between an animal and its heterogeneous
    environment. The large amount of variables involved in that interaction complicates the ecological
    inference thereof. In order to better understand movement, the myriad of involved variables can be
    grouped into behavioural, spatial, and temporal components of the movement process (related to
    the questions of *why*, *where* and *when* to move), which should be addressed simultaneously.

2. We studied the influence of spatial environmental variation on fine-scale movement trajectories of
    eland antelope, blue wildebeest and plains zebra in a South-African game reserve, in simultaneous
    consideration with temporal rhythms and behavioural movement modes.

3. We described the animals' movement process as multi-mode random walks using hidden Markov
    models (HMMs), where both the probabilities of switching between modes as well as the mode-
    dependent speed and turning angle distributions were a function of time of day, tree cover and
    terrain slope.

4. The fitted HMMs clustered the trajectories of all three species into a fast and directed "transit" mode and a slow and tortuous "encamped" mode. All movement properties displayed a strong diel pattern, with the effects of tree cover and slope on speed being largest in the evening when animals generally moved faster. Wildebeest and zebra in their transit mode moved faster with increasing tree cover whereas eland moved slower, but only so on steep terrain. In the encamped mode, tree cover had almost no effect on speed, but a strong effect on tortuosity. The animals also generally moved more tortuously with higher tree cover, even more so on sloping terrain.

5. Due to our integrated approach, we showed that the influence of spatial environmental heterogeneity on animal movement varies greatly in diel cycles and between movement modes, reflecting variation in the animals' habitat selection and decision making throughout the day. Thus, focusing on the interactions between *why*, *when* and *where* to move is needed to more truthfully describe and understand animal movement.

Keywords: behavioural thermoregulation; diurnal cycle patterns; foraging; GPS tracking; predator-prey interactions; state space modelling; ungulate movement; wildlife activity

1 Introduction

Movement is ubiquitous to animal life and central to ecological phenomena such as territoriality and home range dynamics (Kie et al., 2002; Moorcroft et al., 1999), predator-prey dynamics (Abrams, 2007; Flaxman & Lou, 2009), distribution patterns (Bailey et al., 1996) and population dynamics (Armsworth & Roughgarden, 2005; Morales et al., 2010; Revilla & Wiegand, 2008). Studying what drives the movement of animals through their environment is thus key to understanding emergent movement properties and their ecological consequences (Morales et al., 2010; Schick et al., 2008). However, this understanding is hampered by the complexity of the interactions between animals and their environment. Namely, similar movement patterns might emerge from different underlying processes that require different ecological interpretations. For example, animals might move slowly through thick vegetation because it physically impedes their movement, because they forage, hide from predators, or seek shelter

from solar radiation. Moreover, the response of animals to their environment may change over time, e.g.

via diel, lunar and seasonal patterns (e.g. de Knegt et al., 2021). Therefore, the animal movement process

has generally been divided into several components in an effort to alleviate the interpretation issues: an

internal motive regarding *why* to move and its temporally and spatially varying environment influencing

when and *where* to move (Nathan et al., 2008).

First, "*why to move*" relates to the internal motives for movement (e.g. reproduction, resource acquisition

and threat avoidance), which are often manifested in different behaviours (e.g. mating, foraging and

fleeing) that produce distinct movement modes (e.g. distributions of speed and turning angle)

(McClintock et al., 2012; Morales et al., 2004; Nathan et al., 2008). Distinguishing different movement

modes in animal trajectories has yielded insights in different foraging strategies within individuals

(McClintock et al., 2017), predator-prey interactions (Forester et al., 2007) and effects of anthropogenic

disturbance on behaviour (van Beest et al., 2019). Acknowledging the internal motives and their

emergent movement modes is thus key to the interpretation of animal movement, specifically regarding

how this movement interacts with the environment (Jonsen et al., 2006; McClintock et al., 2012, 2017;

Morales et al., 2004).

Second, regarding *"when to move"*, animal movement often shows pronounced temporal variation and

even cyclic patterns. For example, seasonal migratory movements (e.g. Dingle, 2014; Dingle & Drake,

2007) have long been recognized, as well as the influence of lunar rhythms (e.g. Polansky et al., 2010;

Riotte-Lambert & Matthiopoulos, 2020). Moreover, many animals show strong diel activity patterns

(e.g. Ensing et al., 2014; Forester et al., 2007; Green & Bear, 1990; Massé & Côté, 2013; Owen-Smith

& Goodall, 2014; Vazquez et al., 2019), which can arise directly due to variation in light and thermal

conditions (Dussault et al., 2004) or indirectly due to predation risk and human activity (Ensing et al.,

2014; Fischhoff et al., 2007; Owen-Smith & Traill, 2017; Skarin et al., 2010). Temporal periodicity is

thus a central characteristic in the movements of animals (Riotte-Lambert et al., 2013).

Third, the focus on *"where to move"* relates the movement of animals to the characteristics of their

spatial surroundings. Especially the effects of vegetation and topography on the movement of large

terrestrial herbivores have received much attention. Vegetation-rich environments often coincide with

lower movement speeds (Avgar et al., 2013; de Knegt et al., 2007, 2021; Frair et al., 2005; Fryxell et

al., 2008). Topography can influence movement through energetic costs associated with moving through

sloping or rugged terrain (Dailey & Hobbs, 1989; Parker et al., 1984; White & Yousef, 1978). Perhaps

surprisingly, ungulates have been found to increase movement speed on slopes, possibly to reduce their

time spent in energetically costly surroundings (Frair et al., 2005). Furthermore, ungulates can select

slopes for foraging since elevation differences increase the diversity of encountered vegetation and

thereby the chance of finding high quality forage (Leblond et al., 2010). Daily cycles can also interact

strongly with the landscape to affect animal movement (Ager et al., 2003; Fischhoff et al., 2007; Leblond

et al., 2010). For example, ungulates can select areas with high vegetation cover during daytime and

more open areas during the night to decrease predation risk (Ager et al., 2003; Leblond et al., 2010).

These examples illustrate that relationships between spatial environmental characteristics and

movement patterns are conditional on the context of internal motive and time, as this context determines

the costs and benefits that these spatial environmental characteristics represent.

Internal motives, temporal rhythms and spatial environmental heterogeneity are thus three intricately

linked components of the animal movement process. However, animal movement research has often

been hampered by the focus on univariate analyses (Eikelboom et al., 2020), while ideally the

relationships between animal movement and each of these three components should be modelled and

interpreted within each other's context (Nathan et al., 2008). Only when interpreting animal movement

relative to these three components acting in concert can we advance our understanding of the processes

shaping animal movement.

Here we study how movement modes, diel patterns and spatial environmental heterogeneity interact to

shape the movement of large ungulates by modelling the movement process explicitly as a composite

of these components. We use a large amount of fine-scale (10 minute resolution) GPS tracking data of

eland antelope, blue wildebeest and plains zebra in a South-African game reserve to fit Hidden Markov Models (HMMs) with input from the time of day, tree cover and terrain slope. HMMs can describe the movement process as a multi-mode random walk comprised of step lengths and turning angles, with input from both temporal variation and environmental heterogeneity (McClintock et al., 2020), thereby making them an ideal model for our purpose. We aim to quantify the interactive effects of movement mode (as a proxy for internal motive), diel patterns and spatial environmental heterogeneity on movement properties, and how these differ between our study species. In doing so, we address the questions of *why*, *when* and *where* to move, thereby trying to advance our understanding of the movements of our study species and demonstrating ways to analyse these components in concert.

2 Materials and methods

2.1 Animals and study area

From June to August 2017, 34 eland antelope (*Taurotragus oryx*), 34 blue wildebeest (*Connochaetes taurinus*) and 35 plains zebra (*Equus quagga*) were captured in Welgevonden Game Reserve, Limpopo, South Africa (24°13'S, 27°54'E), and equipped with neck collars containing GPS and accelerometer sensors. For more details about the collaring process and sensors, see de Knegt et al. (2021). The animals were released in a fenced study area of 1200 hectares at the northern edge of the reserve (Fig. 1). As a part of the Waterberg mountain massif and Bushveld ecoregion (Mucina et al., 2006), the study area is characterised by nutrient poor sandy soils in the lower northern part and rocky, flat hilltop plateaus dissected by ravines extending towards the south. Woody vegetation varies from mixed to open dry deciduous woodland, with the rocky plateaus being the most open.

Figure 1. Overview of the study area and its environmental heterogeneity with A) terrain slope (radians) and B) fractional tree cover. Numbers at the isoclines denote elevation from sea level (m).

2.2 Animal movement data

The GPS sensors collected spatial positions with a frequency ranging between 15 and 2 minutes, depending on the amount of body activity as determined by the accelerometer (see de Knegt et al., 2021). The position data were corrected, smoothed and modelled to regular ten-minute resolution trajectories as described in de Knegt et al. (2021). Due to sensor failures, we selected from each species only the ten individuals with the most recorded data in the period from September to December 2017 (the first four months of the study period, during which the sensors collected most data). Our subset contained a total of 357,815 positions across 30 individuals of the three species, comprising 481 bouts (i.e. an uninterrupted and regular sequence of positions of a single individual) with a median bout length of 496 positions (3.4 days).

2.3 Environmental data

A digital elevation model of the study area with resolution 2x2 m was available, from which terrain slope (hereafter referred to as slope) was calculated (Fig. 1A). We used aerial images of the study area from August 2013 to create an orthomosaic with a 0.25x0.25 m resolution, which we then used to classify the pixels into trees, grass, bare ground and other/built up area using the Semi-Automatic Classification Plugin for QGIS (Congedo, 2021). After having applied an isotropic Gaussian kernel smoother with a 10 m standard deviation to the classified pixels, we derived fractional tree cover

(hereafter referred to as tree cover) and fractional grass cover maps of the study area at a 2x2 m

resolution (Fig. 1B). Tree cover was strongly negatively correlated with grass cover ($r = 0.95$), so we

decided to use only tree cover in our analyses and consider it a measure for the inverse of grass cover as

well. We extracted tree cover and slope values for each animal position.

2.4 Movement model structure

We constructed a Hidden Markov Model (HMM) for each species separately to examine how their

movement properties were shaped by movement modes, diel patterns and environmental heterogeneity.

We described the movement process as a random walk consisting of steps (distances between subsequent

positions) with lengths s and turning angles φ (change between subsequent movement headings). Since

intervals between steps were constant at ten minutes, we refer to s as speed throughout. s and φ were

fitted to *gamma* (Γ) and *von Mises* (*vM*) distributions, respectively, which were unique to each discrete

movement mode z. Thus, z was defined as a unique combination of three movement parameters: the

mean μ and standard deviation σ of the speed (converted from the shape and scale parameter of the Γ

distribution for easier interpretation) and the *vM* concentration parameter κ of the turning angle (the *vM*

mean was fixed at 0 since there was no reason to assume a movement bias to either the right or left

direction). μ, σ and κ were in turn modelled as functions of several covariates using a log-link (see

section *Movement model covariates*). Mode switching between the steps was governed by switching

matrix M, where the off-diagonal elements are logit-linear functions of covariates. We considered a

three-mode, two-mode and a one-mode HMM, i.e. 3x3, 2x2 and 1x1 dimensions for M respectively.

Thus, the HMM had the general form:

175
$$M = \begin{bmatrix} \gamma_{1\to1} = 1 - \sum_{z=2}^{n} \gamma_{1\to z} & \cdots & \gamma_{1\to n} = f_a(\boldsymbol{\theta}_a, \boldsymbol{\xi}_a) \\ \vdots & \ddots & \vdots \\ \gamma_{n\to1} = f_b(\boldsymbol{\theta}_b, \boldsymbol{\xi}_b) & \cdots & \gamma_{n\to n} = 1 - \sum_{z=1}^{n-1} \gamma_{n\to z} \end{bmatrix}$$

176
$$s_{z=i} \sim \Gamma\left(\mu = f_c(\boldsymbol{\theta}_{c,i}\,\boldsymbol{\xi}_c), \sigma = f_d(\boldsymbol{\theta}_{d,i}, \boldsymbol{\xi}_d)\right)$$

177
$$\phi_{z=i} \sim vM\left(\mu = 0, \kappa = f_e(\boldsymbol{\theta}_{e,i}, \boldsymbol{\xi}_e)\right)$$

179   Where $\gamma_{i \to j}$ is the probability of switching from mode $z = i$ to $z = j$ between consecutive steps, f_n are

180   functions for step switching probabilities and Γ and vM parameters, $\boldsymbol{\theta}_n$ are sets of to-be-fitted coefficients

181   of function f_n (and for mode $z = i$ in case of $\boldsymbol{\theta}_{n,i}$), and $\boldsymbol{\xi}_n$ are sets of covariates (see next section).

2.5 Movement model covariates

Within the HMMs, we modelled the parameters μ, σ and κ and the mode switching probabilities $\gamma_{i \to j}$

each as a function of three covariates. Two of those were the spatial environmental covariates tree cover

and slope. The third covariate was a constructed daily cyclic pattern covariate (DCC) that describes the

diel patterns in μ, σ and κ (driven by e.g. ambient temperature, solar position, perceived predation risk

and internal diel rhythm). To determine those diel patterns, we binned the GPS position data per time of

day (144 bins of 10 minutes) and fitted Γ distributions (μ and σ) to speeds and vM distributions (κ) to

turning angles within a centred moving window of three bins wide. This yielded empirical diel patterns

in parameters μ, σ and κ (Fig. S1). We constructed DCC by first rescaling and averaging the empirical

diel patterns in μ, σ and κ to one aggregate diel pattern per species (Fig. 2), given that these separate

empirical diel patterns were strongly correlated within species (Fig. S2). We then fitted to that aggregate

diel pattern a mixture of a cosine and four wrapped normal (*WN*) distributions with 24 h periods to

compute the final DCC values per species, with the wrapped normal distributions representing two peaks

and two lows per day that were fixed at the peaks and lows of the aggregate diel patterns (Fig. 2). Thus,

DCC had the following form:

$$DCC = \alpha + \beta \cos(t_{day} + \gamma) + \sum_{n=1}^{4} \delta_n f_{WN}(\mu_n, \sigma_n)$$

201   Where t_{day} is the time of day (range $0 - 2\pi$), f_{WN} is the probability density function of the *WN* distribution

202   with a 24 h period (describing 1 out of 4 peaks or lows per day in the aggregate diel pattern), and δ_n are

203   scalars for those distributions. As a result, each animal position had a corresponding DCC value, which

204   depended on the time of day and the species (Fig. 2). We then scaled tree cover, slope and DCC to a

 mean of 0 and standard deviation of 1, and included the three variables and their two-way interactions

as covariates of μ, σ and κ in the HMMs. We included tree cover, slope and DCC without interactions

as covariates of the mode switching probabilities $\gamma_{i \to j}$.

Figure 2. Aggregate diel patterns averaged from μ, σ and κ for eland, wildebeest and zebra (solid line). We fitted

a daily cycle covariate (DCC, dashed line) to those aggregate diel patterns, which summarizes the diel activity

patterns per species as expressed in movement properties μ, σ and κ. DCC is a mixture distribution of a cosine and

four wrapped normal distributions with 24 h periods.

2.6 Model fitting workflow

We fitted the HMM per species and per unique number of movement modes separately in eleven

iterations (Table 1), by numerical maximisation of the log-likelihood function, using the *fitHMM* routine

from the R package *momentuHMM* (McClintock & Michelot, 2018). For the initial iteration, we fitted

a baseline HMM without any covariates to speed μ and σ, turning angle κ and mode switching

probabilities $\gamma_{i \to j}$. For the second to tenth iterations, we added covariates and/or interactions between

covariates to the distribution parameters and switching probabilities (Table 1). The full model in the

eleventh iteration included tree cover, slope and DCC as covariates to $\gamma_{i \to j}$, and these three variables and

their two-way interactions as covariates to μ, σ and κ. The fitted parameter values from each iteration

were used as starting values for the next; starting values for other parameters were set to 0 (note that all

covariates had been scaled to mean 0 and standard deviation 1). For each iteration, four sets of

perturbations of those starting values β were sampled from $\sim N(\beta, \sigma = 0.25 \cdot \beta)$. The model fit with the

highest likelihood out of all starting sets was taken to be the optimal fit and subsequently used to initiate the model fit of the next iteration. However, in practice all sets of fitted models converged to almost identical coefficient values and model likelihoods. We used the Akaike information criterion (AIC) to assess model parsimony throughout the workflow. The model structure and fitting workflow were identical for all three species and number of modes.

All analyses were performed in R 3.6.1 (R Core Team, 2019), using the packages *raster* 3.4-5 (Hijmans, 2021) for map plotting and covariate extraction and *momentuHMM* 1.5.1 (McClintock & Michelot, 2018) for model fitting and calculation of model AIC and stationary mode probabilities.

3 Results

3.1 Hidden Markov Models

We explored Hidden Markov Models (HMMs) with one, two and three movement modes. The single-mode HMMs were far inferior to the two-mode HMMs in terms of model fit, with the best single-mode models being outperformed by the worst two-mode models (ΔAIC = 74,117; 105,847 and 146,886 for eland, wildebeest and zebra respectively) and even more by the best two-mode models (ΔAIC = 85,049; 114,821 and 154,458 for eland, wildebeest and zebra respectively). The best three-mode HMMs were better still than the best two-mode HMMs in terms of model fit, albeit with a smaller difference (ΔAIC = 32,093; 29,170 and 27,381 for eland, wildebeest and zebra respectively). Both the two- and three-mode HMMs detected two modes that were qualitatively similar in terms of speed and turning angle distributions, with the three-mode HMMs detecting an additional slow and directional mode (Fig. S3). This third mode responded very little to both the temporal and spatial environmental variables, while the response displayed by the other two modes was qualitatively similar for both the two- and three-mode HMMs (Figs. S4-S6). Therefore, we visualized only the results from the two-mode HMMs in the main text. However, to strengthen the reliability of our analysis, we only formulated results and conclusions that were supported by both the two- and three-mode HMMs (Wagenmakers et al., 2022). The likelihood of all two-mode HMMs, which were fitted in eleven iterative steps for each species, are

presented in Table 1. The models including only one DCC covariate (step 6) were superior to all

previous models including tree cover and slope, indicating that DCC was the dominant explanatory

variable, accounting for a lot of temporal variation which is not accounted for by tree cover and slope

alone. For each species the full model (step 11), which includes all covariates and interactions, was the

most parsimonious model according to AIC value. We therefore analysed the data further using the full

models.

Table 1. Overview of iteratively fitted two-mode HMMs and their ΔAIC per species. ΔAIC was calculated for

each species separately relative to the best model of that species. tc = tree cover, sl = terrain slope, DCC = daily

cycle covariate, μ, σ and κ are movement parameters and $\gamma_{i \to j}$ = mode switching probability. Powers of 2 indicate

all two-way interactions among the covariates between brackets. The AIC of the full models (step 11) was

1,305,636; 1,304,360 and 1,286,972 for eland, wildebeest and zebra respectively.

Predictor set	Model step	μ, σ and κ covariates	$\gamma_{i \to j}$ covariates	ΔAIC Eland	ΔAIC Wildebeest	ΔAIC Zebra
none	1	–	–	10,932	8,974	7,572
spatial	2	tc, sl	–	10,354	8,451	5,782
	3	tc, sl	tc, sl	10,278	8,405	5,727
	4	(tc, sl)^2	–	10,233	8,267	5,362
	5	(tc, sl)^2	tc, sl	10,157	8,219	5,327
temporal	6	DCC	–	882	978	2,299
	7	DCC	DCC	822	918	2,240
spatial + temporal	8	tc, sl, DCC	–	256	678	787
	9	tc, sl, DCC	tc, sl, DCC	145	432	677
	10	(tc, sl, DCC)^2	–	114	241	67
	11	(tc, sl, DCC)^2	tc, sl, DCC	0	0	0

3.2 Movement modes

All species displayed one faster, more directed mode and one slower, more tortuous mode (Fig. S3). We

labelled these as "transit" and "encamped" movement modes, thereby avoiding to project too specific

behavioural annotations on the modes. For example, while slower and more tortuous movement could be associated with foraging, actual foraging behaviour cannot be concluded from speed and turning angle alone due to overlap of those movement properties among different behaviours (McClintock et al., 2020). We considered "transit" and "encamped" to be reasonable abstractions while retaining spatially descriptive meaning. Overall, zebra moved markedly faster in the transit mode compared to the other two species (zebra: 0.62 ± 0.50SD km/h, eland: 0.25 ± 0.28SD km/h, wildebeest: 0.32 ± 0.30SD km/h) and wildebeest moved markedly slower in the encamped mode compared to the other two species (zebra: 0.13 ± 0.13SD km/h, eland: 0.10 ± 0.17SD km/h, wildebeest: 0.04 ± 0.05SD km/h). Contrarily, the movement tortuosity was more similar between the two modes for wildebeest (transit mode $\kappa = 1.41$, encamped mode $\kappa = 0.89$) and zebra (transit mode $\kappa = 4.80$, encamped mode $\kappa = 1.19$) than for eland (transit mode $\kappa = 18.00$, encamped mode $\kappa = 0.68$).

3.3 Movement related to covariates

The main effects of time of day, tree cover and slope only partly described speed (parameters μ and σ) and turning angle (parameter κ). Speed differed greatly throughout the day, being low just before sunrise and peaking around sunrise and in the evening (Fig. 3; the sunrise peak is not pictured). Higher tree cover resulted in slower eland movement, but faster wildebeest and zebra movement (Figs. S7-S9). Furthermore, all species moved slower at steeper slopes, with their movement tortuosity increasing slightly for eland but decreasing for wildebeest and zebra. Time of day, tree cover and slope interacted to varying degrees to more accurately predict the movement parameters (Fig. 3). On flat terrain, effects of tree cover on speed were generally small to negligible for all three species. On steeper terrain, an increase in tree cover caused eland to move slower in both modes, but wildebeest and zebra in the transit mode to move faster. This interaction was generally more pronounced in the evening (18:00) than in the morning (04:00) and at noon. Likewise, tree cover had an effect on movement tortuosity predominantly on steeper terrain, where it generally caused animals to move more tortuous (Fig. 4). In their transit mode, eland and zebra moved more tortuous than wildebeest, but eland tortuosity was not affected by tree cover or slope. Movement tortuosity was quite stable throughout the day for eland and zebra, but for wildebeest it differentiated more between the internal modes in the evening than in the morning.

Figure 3. Model predictions of movement speed μ as a result of tree cover and slope in the morning (04:00, top row), at noon (12:00, middle row) and in the evening (18:00, bottom row) for eland, wildebeest and zebra. The times were chosen to capture a dip, baseline and peak in DCC (see Fig. 2). Blue and red indicate the transit and encamped mode respectively. Colour mapping within blue and red indicates a progression of slope from 0 (light) till 0.15 (the 90th percentile of slope values in the complete dataset; dark). Small ticks at the inside of tree cover axes mark the 10th, 20th … 90th percentile values of tree cover in the respective data sets. Note that the y-axis scales differ between species. Confidence intervals have been omitted for visual clarity, but they were generally small. For speed σ, see Fig. S5.

Figure 4. Model predictions of turning angle concentration as a result of tree cover and slope in the morning (04:00, top row), at noon (12:00, middle row) and in the evening (18:00, bottom row) for eland, wildebeest and zebra. Concentration parameter κ (which ranges from 0-infinity) has been converted to ρ (range 0-1) in this figure for visualization purposes. The times were chosen to capture a dip, baseline and peak in DCC (see Fig. 2). Blue and red indicate the transit and encamped mode respectively. Colour mapping within blue and red indicates a progression of slope from 0 (light) till 0.15 (the 90th percentile value of slope in the complete dataset; dark). Small ticks at the inside of tree cover axes mark the 10th, 20th … 90th percentile values of tree cover in the respective data sets. Confidence intervals have been omitted for visual clarity, but they were generally small.

3.4 Movement mode occupancy

Time of day had little influence on the stationary mode probabilities of all species (Fig. S10). Wildebeest showed some diel variation, being more likely to move in its encamped state but the difference being smaller in the evening than before sunrise. Tree cover also had little influence on the mode probabilities of eland and wildebeest, while zebra became slightly more likely to move in their transit mode when in areas with a higher tree cover. On steeper slopes, wildebeest became considerably less likely to display their transit mode, while the other species did not.

4 Discussion

We investigated how the movements of three large African savanna ungulate species arise from the interplay between movement modes and temporal and spatial environmental heterogeneity, thereby adressing the questions of *why*, *when* and *where* to move. We show that these three components of the movement process explained a large amount of variation in the emergent movement properties, but especially that there were strong interactions between those components. All movement properties displayed a strong diel pattern, with the effects of tree cover and slope on speed being largest in the evening when animals generally moved faster. Wildebeest and zebra in their "transit" mode moved faster with increasing tree cover whereas eland moved slower, but only so on steep terrain. In the "encamped" mode, tree cover had almost no effect on speed, but a strong effect on tortuosity. The animals also generally moved more tortuous with higher tree cover, even more so on sloping terrain. Tree cover did not influence the probability of being in a certain movement mode, and slope made only wildebeest more likely to be in the encamped mode. All our results and conclusions were supported by both the two- and three-mode HMMs, thereby disregarding those results for which the models did not agree and thus strengthening the reliability of our conclusions (Wagenmakers et al., 2022).

The daily cycle covariate (DCC), our proxy for temporal variation on a diel scale, was the most important factor in explaining variation in the movement properties (Table 1, Fig. 3). There was some level of correlation between variation in DCC on the one hand and spatial environmental variation (tree cover

and slope) on the other hand (Fig. S1, Table S1). This implies that DCC in our analysis captured both

the *direct*, space-invariant, effect of time of day on the movement properties of the animals (i.e.

endogenous rhythms and/or rhythms that are related to temporal variation without a clear spatial

pattern), as well as an *indirect* effect of time of day through temporal variation in habitat selection, given

that habitat influenced the movement properties as well (Figs. 3-4). The purely spatial influence of

environmental heterogeneity on movement (i.e. the fitted time-invariant influence of the tree cover and

slope covariates on movement) was small compared to both the temporal DCC proxy and the

interactions between DCC and the spatial covariates. These results imply that the movements of our

study species are driven mostly by processes that operate along temporal and spatial-temporal

dimensions, rather than along a time-invariant spatial dimension (i.e. behavioural temporal patterns that

arise from either internal diel rhythms or temporal variation in habitat selection throughout the day, or

both).

The purely temporal, space-invariant patterns likely arise from various diel determinants, such as

variation in ambient temperature and perceived predation risk (Palmer et al., 2022). High ambient

temperatures generally suppress movement activity (Ditmer et al., 2018; Owen-Smith, 1998) and many

ungulates show crepuscular activity cycles (Ensing et al., 2014; Forester et al., 2007; Green & Bear,

1990; Vazquez et al., 2019), resting through midday when temperatures are highest (Ben-Shahar, 1987;

Owen-Smith et al., 2010; Owen-Smith & Goodall, 2014; Shrestha et al., 2014). Eland, wildebeest and,

to a lesser extent, zebra in our study conformed to this diel pattern (Fig. 2). Furthermore, movement as

an aspect of antipredator behaviour can show diel patterns in response to daily variability in predation

risk (Bonnot et al., 2013; Fischhoff et al., 2007; Owen-Smith & Traill, 2017; Ross et al., 2013). Activity

during the night can be suppressed by higher predation risk, yielding activity peaks around dawn and

dusk (Crawford et al., 2019). Although our study area was largely predator-free, predator-avoidance

behaviour due to perceived predation risk in low-visibility areas (Riginos, 2015; Riginos & Grace, 2008)

might still have influenced movement in a pattern similar to that of temperature.

Wildebeest and zebra moved faster with increasing tree cover in their transit mode, but not in their

encamped mode. The movement of eland in relation to the environment differed less between its modes,

with eland generally moving slower with increasing tree cover. These patterns can have arisen through

differences in predation risk between the species. Eland is much larger than wildebeest and zebra and

adult elands are less vulnerable to predation from all but the largest predators. Leopards were present in

the study area and they select somewhat bushy surroundings for ambushing their prey (Balme et al.,

2007). Areas with higher tree or shrub cover may thus present a larger perceived risk of predation than

open grassland for wildebeest and zebra, whereas for eland the difference in perceived risk between the

habitats would be smaller. Differences in diet could be another possible explanation for these movement

differences between the species: grazing wildebeest and zebra generally forage in areas with high grass

cover (Groom & Harris, 2010) (i.e. low tree cover in our study area), and may thus increase their speed

in areas with higher tree cover due to the lower likelihood of encountering available forage there. For

eland, a mixed feeder (McNaughton & Georgiadis, 1986), bushy areas provide forage and may hence

cause less of a speed increase.

The animals generally moved slower in their transit mode with increasing slope. Previous research has

found deer more likely to move faster on steeper terrain, assumedly due to energetic costs rendering

slopes less preferable to reside on (Frair et al., 2005). Our results show that responses to spatial

environmental heterogeneity depend on the movement mode and hence partly on the motive for moving.

Especially when in transit, slopes pose increased energetic costs or injury risk associated with moving

through such terrain (Dailey & Hobbs, 1989; Parker et al., 1984; White & Yousef, 1978). Moreover,

tree cover interacted with slope to make eland and wildebeest move slower and more tortuously.

Interestingly, on flat terrains tree cover had very little influence, indicating that movement resistance

caused by the environment can accumulate from different components of that environment, which

underpins the importance of considering multiple sources of spatial environmental heterogeneity

simultaneously when investigating their effect on movement.

The discrete movement modes combined various behaviours that resemble each other in terms of movement properties, rather than perfectly representing single behaviours. The encamped mode might for example include resting, foraging and slow explorative movements, where the occurence of these behaviours may vary during the day. The behaviour(s) that are represented by the fitted movement modes can thus differ between models and, in our case, species. This could explain why wildebeest movement in the encamped mode was much slower than that of the other species, as it perhaps did not include e.g. foraging behaviour in contrast to the other two species. This is supported further by the fact that wildebeest, unlike eland and zebra, became much more likely to be in the transit mode on sloping terrain, as highly sloping terrain might be less suitable foraging habitat. The effect of slope on stationary mode probability was smaller for zebra and virtually absent for eland, and tree cover had no significant effect on any of the species. Moreover, mode occupancy in eland and zebra showed very little variation throughout the day, but wildebeest showed some (Fig. S10). This suggests that the diel patterns in movement properties were for a large part caused by within-mode behavioural variability instead of diel patterns in mode switching. These results indicate that, in line with our finding that diel patterns are a much stronger driver than spatial (time-invariant) environmental heterogeneity, movement of our study species is predominantly driven by the animals' pre-existing internal motives (e.g. to translocate, feed or rest), which vary during the day and thus lead to temporal variation in spatial behaviour.

5 Conclusion

We demonstrated how behavioural movement modes and temporal and spatial environmental heterogeneity (the different components of the movement process relating to the quesions of *why*, *when* and *where* to move) interact in an intricate way to drive ungulate movement. Within a movement mode, the movements of our study species were driven largely by temporal patterns in their behaviour that resulted from internal diel rhythms and temporal variation in habitat selection throughout the day, as opposed to time-invariant environmental heterogeneity. Future research should specifically investigate to what extent factors such as temperature and predation risk interact to drive diel variation in habitat selection, to further assess the role of habitat selection in determining movement properties throughout

the day. Moreover, studying animals in larger or multiple areas and combining data from multiple ungulate species would allow for a comparison with the specific effects and interactions that we found, which will provide insight in the ungulate movement process across a wider range of applications. Understanding the drivers of animal movement, and ultimately of ecological phenomena that emerge from it, critically depends on considering the various components of the movement process in concert and especially the interactions between them.

Acknowledgments

This research was funded by the Dutch Research Council (NWO program "Advanced Instrumentation for Wildlife Protection"). We thank Welgevonden Game Reserve for the animal monitoring, MTN South Africa for the telecommunication and IBM for the data warehousing.

Conflict of interest

All authors declare to have no conflict of interest.

Author contributions

H.L., H.J.d.K. and J.A.J.E. conceived the ideas for the study and designed its methods; H.J.d.K. and J.A.J.E. provided the data; H.L. analysed the data and led the writing of the manuscript. All authors (H.L., H.J.d.K. and J.A.J.E.) contributed critically to the drafts and gave final approval for publication.

Data availability statement

Our data and code will be made publicly available via the 4TU.ResearchData repository after acceptance of the manuscript.

References

Abrams, P. A. (2007). Habitat Choice in Predator-Prey Systems: Spatial Instability due to Interacting Adaptive Movements. *The American Naturalist*, *169*(5), 581–594. https://doi.org/10.1086/512688

Ager, A. A., Johnson, B. K., Kern, J. W., & Kie, J. G. (2003). Daily and Seasonal Movements and Habitat Use by Female Rocky Mountain Elk and Mule Deer. *Journal of Mammalogy*, *84*(3), 1076–1088. https://doi.org/10.1644/BBa-020

Armsworth, P. R., & Roughgarden, J. E. (2005). The Impact of Directed versus Random Movement on Population Dynamics and Biodiversity Patterns. *The American Naturalist*, *165*(4), 449–465. https://doi.org/10.1086/428595

Avgar, T., Mosser, A., Brown, G. S., & Fryxell, J. M. (2013). Environmental and individual drivers of animal movement patterns across a wide geographical gradient. *Journal of Animal Ecology*, *82*(1), 96–106. https://doi.org/10.1111/j.1365-2656.2012.02035.x

Bailey, D. W., Gross, J. E., Laca, E. A., Rittenhouse, L. R., Coughenour, M. B., Swift, D. M., & Sims, P. L. (1996). Mechanisms that result in large herbivore grazing distribution patterns. *Journal of Range Management*, *49*(5), 386–400.

Balme, G., Hunter, L., & Slotow, R. (2007). Feeding habitat selection by hunting leopards Panthera pardus in a woodland savanna: Prey catchability versus abundance. *Animal Behaviour*, *74*(3), 589–598. https://doi.org/10.1016/j.anbehav.2006.12.014

Ben-Shahar, R. & F. N. (1987). Comparison of the diurnal activity patterns of blue wildebeest and red hartebeest. *South African Journal of Wildlife Research*, *17*(2), 49–53. https://doi.org/10.10520/AJA03794369_3498

Bonnot, N., Morellet, N., Verheyden, H., Cargnelutti, B., Lourtet, B., Klein, F., & Hewison, A. J. M. (2013). Habitat use under predation risk: Hunting, roads and human dwellings influence the spatial behaviour of roe deer. *European Journal of Wildlife Research*, *59*(2), 185–193. https://doi.org/10.1007/s10344-012-0665-8

Congedo, L. (2021). Semi-Automatic Classification Plugin: A Python tool for the download and

processing of remote sensing images in QGIS. *Journal of Open Source Software*, *6*(64), 3172.

https://doi.org/10.21105/joss.03172

Crawford, D. A., Cherry, M. J., Kelly, B. D., Garrison, E. P., Shindle, D. B., Conner, L. M., Chandler,

R. B., & Miller, K. V. (2019). Chronology of reproductive investment determines predation

risk aversion in a felid-ungulate system. *Ecology and Evolution*, *9*(6), 3264–3275.

https://doi.org/10.1002/ece3.4947

Dailey, T. V., & Hobbs, N. T. (1989). Travel in alpine terrain: Energy expenditures for locomotion by

mountain goats and bighorn sheep. *Canadian Journal of Zoology*, *67*(10), 2368–2375.

https://doi.org/10.1139/z89-335

de Knegt, H. J., Eikelboom, J. A. J., van Langevelde, F., Spruyt, W. F., & Prins, H. H. T. (2021).

Timely poacher detection and localization using sentinel animal movement. *Scientific Reports*,

11(1), 4596. https://doi.org/10.1038/s41598-021-83800-1

de Knegt, H. J., Hengeveld, G. M., van Langevelde, F., de Boer, W. F., & Kirkman, K. P. (2007).

Patch density determines movement patterns and foraging efficiency of large herbivores.

Behavioral Ecology, *18*(6), 1065–1072. https://doi.org/10.1093/beheco/arm080

Dingle, H. (2014). Seasonal migration patterns. In H. Dingle (Ed.), *Migration: The Biology of Life on*

the Move (pp. 163–182). Oxford University Press.

https://doi.org/10.1093/acprof:oso/9780199640386.003.0009

Dingle, H., & Drake, V. A. (2007). What Is Migration? *BioScience*, *57*(2), 113–121.

https://doi.org/10.1641/B570206

Ditmer, M. A., Moen, R. A., Windels, S. K., Forester, J. D., Ness, T. E., & Harris, T. R. (2018).

Moose at their bioclimatic edge alter their behavior based on weather, landscape, and

predators. *Current Zoology*, *64*(4), 419–432. https://doi.org/10.1093/cz/zox047

Dussault, C., Ouellet, J.-P., Courtois, R., Huot, J., Breton, L., & Larochelle, J. (2004). Behavioural

responses of moose to thermal conditions in the boreal forest. *Écoscience*, *11*(3), 321–328.

https://doi.org/10.1080/11956860.2004.11682839

Eikelboom, J. A. J., de Knegt, H. J., Klaver, M., van Langevelde, F., van der Wal, T., & Prins, H. H.

T. (2020). Inferring an animal's environment through biologging: Quantifying the

environmental influence on animal movement. *Movement Ecology*, *8*(1), 40.

https://doi.org/10.1186/s40462-020-00228-4

Ensing, E. P., Ciuti, S., Wijs, F. A. L. M. de, Lentferink, D. H., Hoedt, A. ten, Boyce, M. S., & Hut, R.

A. (2014). GPS Based Daily Activity Patterns in European Red Deer and North American Elk

(Cervus elaphus): Indication for a Weak Circadian Clock in Ungulates. *PLOS ONE*, *9*(9),

e106997. https://doi.org/10.1371/journal.pone.0106997

Fischhoff, I. R., Sundaresan, S. R., Cordingley, J., & Rubenstein, D. (2007). Habitat use and

movements of plains zebra (Equus burchelli) in response to predation danger from lions.

Behavioral Ecology, *18*(4), 725–729. https://doi.org/10.1093/beheco/arm036

Flaxman, S. M., & Lou, Y. (2009). Tracking prey or tracking the prey's resource? Mechanisms of

movement and optimal habitat selection by predators. *Journal of Theoretical Biology*, *256*(2),

187–200. https://doi.org/10.1016/j.jtbi.2008.09.024

Forester, J. D., Ives, A. R., Turner, M. G., Anderson, D. P., Fortin, D., Beyer, H. L., Smith, D. W., &

Boyce, M. S. (2007). State–Space Models Link Elk Movement Patterns to Landscape

Characteristics in Yellowstone National Park. *Ecological Monographs*, *77*(2), 285–299.

https://doi.org/10.1890/06-0534

Frair, J. L., Merrill, E. H., Visscher, D. R., Fortin, D., Beyer, H. L., & Morales, J. M. (2005). Scales of

movement by elk (Cervus elaphus) in response to heterogeneity in forage resources and

predation risk. *Landscape Ecology*, *20*(3), 273–287. https://doi.org/10.1007/s10980-005-2075-

8

Fryxell, J. M., Hazell, M., Börger, L., Dalziel, B. D., Haydon, D. T., Morales, J. M., McIntosh, T., &

Rosatte, R. C. (2008). Multiple movement modes by large herbivores at multiple

spatiotemporal scales. *Proceedings of the National Academy of Sciences*, *105*(49), 19114–

19119. https://doi.org/10.1073/pnas.0801737105

Green, R. A., & Bear, G. D. (1990). Seasonal Cycles and Daily Activity Patterns of Rocky Mountain

Elk. *The Journal of Wildlife Management*, *54*(2), 272–279. https://doi.org/10.2307/3809041

Groom, R., & Harris, S. (2010). Factors affecting the distribution patterns of zebra and wildebeest in a

resource-stressed environment. *African Journal of Ecology*, *48*(1), 159–168.

https://doi.org/10.1111/j.1365-2028.2009.01097.x

Hijmans, R. J. (2021). *raster: Geographic Data Analysis and Modeling* (3.5-11). https://CRAN.R-

project.org/package=raster

Hilborn, R., & Mangel, M. (1997). *The Ecological Detective*. Princeton University Press.

Jonsen, I. D., Myers, R. A., & James, M. C. (2006). Robust hierarchical state–space models reveal diel

variation in travel rates of migrating leatherback turtles. *Journal of Animal Ecology*, *75*(5),

1046–1057. https://doi.org/10.1111/j.1365-2656.2006.01129.x

Kie, J. G., Bowyer, R. T., Nicholson, M. C., Boroski, B. B., & Loft, E. R. (2002). Landscape

Heterogeneity at Differing Scales: Effects on Spatial Distribution of Mule Deer. *Ecology*,

83(2), 530–544. https://doi.org/10.1890/0012-9658(2002)083[0530:LHADSE]2.0.CO;2

Leblond, M., Dussault, C., & Ouellet, J.-P. (2010). What drives fine-scale movements of large

herbivores? A case study using moose. *Ecography*, *33*(6), 1102–1112.

https://doi.org/10.1111/j.1600-0587.2009.06104.x

Massé, A., & Côté, S. D. (2013). Spatiotemporal variations in resources affect activity and movement

patterns of white-tailed deer (Odocoileus virginianus) at high density. *Canadian Journal of

Zoology*, *91*(4), 252–263. https://doi.org/10.1139/cjz-2012-0297

McClintock, B. T., King, R., Thomas, L., Matthiopoulos, J., McConnell, B. J., & Morales, J. M.

(2012). A general discrete-time modeling framework for animal movement using multistate

random walks. *Ecological Monographs*, *82*(3), 335–349. https://doi.org/10.1890/11-0326.1

McClintock, B. T., Langrock, R., Gimenez, O., Cam, E., Borchers, D. L., Glennie, R., & Patterson, T.

A. (2020). Uncovering ecological state dynamics with hidden Markov models. *Ecology

Letters*, *23*(12), 1878–1903. https://doi.org/10.1111/ele.13610

McClintock, B. T., London, J. M., Cameron, M. F., & Boveng, P. L. (2017). Bridging the gaps in

animal movement: Hidden behaviors and ecological relationships revealed by integrated data

streams. *Ecosphere*, *8*(3), e01751. https://doi.org/10.1002/ecs2.1751

McClintock, B. T., & Michelot, T. (2018). momentuHMM: R package for generalized hidden Markov models of animal movement. *Methods in Ecology and Evolution*, *9*(6), 1518–1530. https://doi.org/10.1111/2041-210X.12995

McNaughton, S. J., & Georgiadis, N. J. (1986). Ecology of African Grazing and Browsing Mammals. *Annual Review of Ecology and Systematics*, *17*, 39–65.

Moorcroft, P. R., Lewis, M. A., & Crabtree, R. L. (1999). Home Range Analysis Using a Mechanistic Home Range Model. *Ecology*, *80*(5), 1656–1665. https://doi.org/10.2307/176554

Morales, J. M., Haydon, D. T., Frair, J., Holsinger, K. E., & Fryxell, J. M. (2004). Extracting More Out of Relocation Data: Building Movement Models as Mixtures of Random Walks. *Ecology*, *85*(9), 2436–2445. https://doi.org/10.1890/03-0269

Morales, J. M., Moorcroft, P. R., Matthiopoulos, J., Frair, J. L., Kie, J. G., Powell, R. A., Merrill, E. H., & Haydon, D. T. (2010). Building the bridge between animal movement and population dynamics. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1550), 2289–2301.

Mucina, L., Rutherford, M. C., Powrie, L., AG, R., KGT, C., Lötter, M. C., DB, H., C, B., Bredenkamp, G. J., Vlok, J. H. J., Euston-Brown, D., Jürgens, N., Preez, P., A, le, Schmiedel, U., CR, S., & N, van. (2006). Vegetation Atlas of South Africa, Lesotho and Swaziland. In *The Vegetation of South Africa, Lesotho and Swaziland* (pp. 748–790). South African National Biodiversity Institute.

Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D., & Smouse, P. E. (2008). A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences*, *105*(49), 19052–19059. https://doi.org/10.1073/pnas.0800375105

Owen-Smith, N. (1998). How high ambient temperature affects the daily activity and foraging time of a subtropical ungulate, the greater kudu (Tragelaphus strepsiceros). *Journal of Zoology*, *246*(2), 183–192. https://doi.org/10.1111/j.1469-7998.1998.tb00147.x

Owen-Smith, N., Fryxell, J. M., & Merrill, E. H. (2010). Foraging theory upscaled: The behavioural ecology of herbivore movement. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1550), 2267–2278. https://doi.org/10.1098/rstb.2010.0095

Owen-Smith, N., & Goodall, V. (2014). Coping with savanna seasonality: Comparative daily activity patterns of African ungulates as revealed by GPS telemetry. *Journal of Zoology*, *293*(3), 181–191. https://doi.org/10.1111/jzo.12132

Owen-Smith, N., & Traill, L. W. (2017). Space use patterns of a large mammalian herbivore distinguished by activity state: Fear versus food? *Journal of Zoology*, *303*(4), 281–290. https://doi.org/10.1111/jzo.12490

Palmer, M. S., Gaynor, K. M., Becker, J. A., Abraham, J. O., Mumma, M. A., & Pringle, R. M. (2022). Dynamic landscapes of fear: Understanding spatiotemporal risk. *Trends in Ecology & Evolution*, *37*(10), 911–925. https://doi.org/10.1016/j.tree.2022.06.007

Parker, K. L., Robbins, C. T., & Hanley, T. A. (1984). Energy Expenditures for Locomotion by Mule Deer and Elk. *The Journal of Wildlife Management*, *48*(2), 474–488. https://doi.org/10.2307/3801180

Polansky, L., Wittemyer, G., Cross, P. C., Tambling, C. J., & Getz, W. M. (2010). From moonlight to movement and synchronized randomness: Fourier and wavelet analyses of animal location time series data. *Ecology*, *91*(5), 1506–1518. https://doi.org/10.1890/08-2159.1

R Core Team. (2019). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. https://cran.r-project.org/bin/windows/base/old/3.6.1/

Revilla, E., & Wiegand, T. (2008). Individual movement behavior, matrix heterogeneity, and the dynamics of spatially structured populations. *Proceedings of the National Academy of Sciences*, *105*(49), 19120–19125. https://doi.org/10.1073/pnas.0801725105

Riginos, C. (2015). Climate and the landscape of fear in an African savanna. *Journal of Animal Ecology*, *84*(1), 124–133. https://doi.org/10.1111/1365-2656.12262

Riginos, C., & Grace, J. B. (2008). Savanna Tree Density, Herbivores, and the Herbaceous Community: Bottom-up Vs. Top-down Effects. *Ecology*, *89*(8), 2228–2238. https://doi.org/10.1890/07-1250.1

Riotte-Lambert, L., Benhamou, S., & Chamaillé-Jammes, S. (2013). Periodicity analysis of movement recursions. *Journal of Theoretical Biology*, *317*, 238–243. https://doi.org/10.1016/j.jtbi.2012.10.026

Riotte-Lambert, L., & Matthiopoulos, J. (2020). Environmental Predictability as a Cause and Consequence of Animal Movement. *Trends in Ecology & Evolution*, *35*(2), 163–174. https://doi.org/10.1016/j.tree.2019.09.009

Ross, J., Hearn, A. J., Johnson, P. J., & Macdonald, D. W. (2013). Activity patterns and temporal avoidance by prey in response to Sunda clouded leopard predation risk. *Journal of Zoology*, *290*(2), 96–106. https://doi.org/10.1111/jzo.12018

Schick, R. S., Loarie, S. R., Colchero, F., Best, B. D., Boustany, A., Conde, D. A., Halpin, P. N., Joppa, L. N., McClellan, C. M., & Clark, J. S. (2008). Understanding movement data and movement processes: Current and emerging directions. *Ecology Letters*, *11*(12), 1338–1350. https://doi.org/10.1111/j.1461-0248.2008.01249.x

Shrestha, A. K., van Wieren, S. E., van Langevelde, F., Fuller, A., Hetem, R. S., Meyer, L., de Bie, S., & Prins, H. H. T. (2014). Larger antelopes are sensitive to heat stress throughout all seasons but smaller antelopes only during summer in an African semi-arid environment. *International Journal of Biometeorology*, *58*(1), 41–49. https://doi.org/10.1007/s00484-012-0622-y

Skarin, A., Danell, Ö., Bergström, R., & Moen, J. (2010). Reindeer movement patterns in alpine summer ranges. *Polar Biology*, *33*(9), 1263–1275. https://doi.org/10.1007/s00300-010-0815-y

van Beest, F. M., Mews, S., Elkenkamp, S., Schuhmann, P., Tsolak, D., Wobbe, T., Bartolino, V., Bastardie, F., Dietz, R., von Dorrien, C., Galatius, A., Karlsson, O., McConnell, B., Nabe-Nielsen, J., Olsen, M. T., Teilmann, J., & Langrock, R. (2019). Classifying grey seal behaviour in relation to environmental variability and commercial fishing activity—A multivariate hidden Markov model. *Scientific Reports*, *9*(1), 5642. https://doi.org/10.1038/s41598-019-42109-w

Vazquez, C., Rowcliffe, J. M., Spoelstra, K., & Jansen, P. A. (2019). Comparing diel activity patterns of wildlife across latitudes and seasons: Time transformations using day length. *Methods in Ecology and Evolution*, *10*(12), 2057–2066. https://doi.org/10.1111/2041-210X.13290

642   Wagenmakers, E.-J., Sarafoglou, A., & Aczel, B. (2022). One statistical analysis must not rule them

643   all. *Nature*, *605*(7910), 423–425. https://doi.org/10.1038/d41586-022-01332-8

644   White, R. G., & Yousef, M. K. (1978). Energy expenditure in reindeer walking on roads and on

645   tundra. *Canadian Journal of Zoology*, *56*(2), 215–223. https://doi.org/10.1139/z78-031

646