1   # Neural, physiological and behavioral correlates of empathy for
2   # pain in Tourette syndrome
3

Ronja Weiblen[1,2], Carina Robert[1], Pauline Petereit[1], Marcus Heldmann[1,3,4], Thomas F. Münte[1,3,4],

Alexander Münchau[4,5], Kirsten Müller-Vahl[6], Ulrike M. Krämer[1,3,4]

[1] Department of Neurology, University of Lübeck, Germany

[2] Department of Psychiatry and Psychotherapy, University of Lübeck, Germany

[3] Department of Psychology, University of Lübeck, Germany

[4] Center of Brain, Behavior and Metabolism, University of Lübeck, Germany

[4] Institute of Systems Motor Science, University of Lübeck, Germany

[5] Department of Psychiatry, Social Psychiatry and Psychotherapy, Hannover Medical School,

Hannover, Germany

* Corresponding authors:

Ronja Weiblen and Ulrike M. Krämer

Department of Neurology, University of Lübeck

Ratzeburger Allee 160

23538 Lübeck, Germany

ronja.weiblen@neuro.uni-luebeck.de and ulrike.kraemer@uni-luebeck.de

Acknowledgements:

The study was supported by the Deutsche Forschungsgemeinschaft (FOR 2698; KR3691/8-1). RW

was funded with a PhD scholarship by the foundation of the German economy (SDW).

26   **Abstract**

Persons with Tourette syndrome (TS) show altered social behaviors, such as echophenomena and

increased personal distress in emotional situations. These symptoms may reflect an overactive mirror

neuron system (MNS), causing both increased automatic imitation and a stronger tendency to share

others' emotions. To test this, we measured the individual level of echophenomena with a video protocol

and experimentally induced empathy for pain in 25 participants with TS and 25 matched controls.

In the empathy for pain paradigm, pictures of hands and feet in painful or neutral situations were

presented, while we measured participants' EEG and skin conductance response (SCR). Changes in

somatosensory mu suppression during the observation of the pictures, as well as pain ratings and SCR

were compared between groups and correlated with the occurrence of echophenomena, self-reported

empathy and clinical measures.

Our TS sample showed significantly more echophenomena than controls, but the groups showed no

behavioral differences in empathic abilities. However, controls, but not patients with TS, showed the

predicted increased mu suppression when watching painful compared to neutral actions. While

echophenomena were present in all persons with TS, the hypothesis of an overactive MNS in TS could

not be substantiated. On the contrary, the TS group showed a noticeable lack of mu attenuation in

response to pain stimuli, questioning the normal functioning of the MNS in empathy for pain situations.

In conclusion, the MNS remains an important area of study for TS, but overactivity of the system could

not explain echophenomena and social symptoms of TS.

Keywords: Tourette Syndrome, Empathy, Mirror Neuron System, Mu Suppression, Echophenomena

Introduction

Gilles de la Tourette Syndrome (TS) is a neuropsychiatric disorder defined by the presence of motor

and vocal tics (DSM 5, Falkai et al., 2018). Of note however, many TS symptoms are inherently social,

socially relevant and/or dependent on the social context. One example are so-called echophenomena,

which are defined as the automatic imitation of actions, sounds or language without explicit awareness

(Ganos et al., 2012). Together with other symptoms such as coprophenomena (Freeman et al., 2009)

or non-obscene socially inappropriate behaviors (NOSI, Kurlan et al., 1996), echophenomena indicate

that social aspects are central to TS, and that individuals with TS are very sensitive to their social

environment. This has led Albin (2018) to suggest his theory of TS as a disorder of the Social Decision

Making network. It has also instigated research on the specific changes in social cognition in TS, but

results so far are inconclusive (Baron-Cohen et al., 1997; Channon et al., 2012; Drury et al., 2012; Eddy

et al., 2010b, 2010a, 2011, 2017; Sprengelmeyer et al., 1997). For example, individuals with TS

reported increased personal distress but decreased perspective taking in the Interpersonal Reactivity

Index, a measure of trait empathy (Eddy et al., 2015). Based on this and other findings, Eddy (2018)

suggested that TS is characterized by increased automatic sharing of others' actions and emotions and

at the same time reduced mentalizing, possibly reflecting an overactive mirror neuron system (MNS).

Similarly, it has been speculated that an overactive MNS might be the cause of echophenomena in TS,

as it is critically involved in movement imitation (Ganos et al., 2012; Heyes & Catmur, 2022; Jonas et

al., 2010). The MNS has been implied in many higher order social processes such as action

understanding, imitation, perspective taking and empathy (Bonini et al., 2022), thus providing a

framework to explain echophenomena as well as irregularities in social behavior in TS. First neural

evidence supports this theory. The temporal parietal junction, which is supposedly part of a top-down

control network modulating the MNS (Brass et al., 2009), shows irregular activity in TS during social

cognition tasks (Eddy et al., 2016, 2017). This supports the idea that control of the MNS might be

impaired in TS, leading to its increased activation. Even though the neural basis of changes in social

cognition as well as echophenomena might be the same, to our knowledge there are currently no

studies investigating social cognition and echophenomena in the same sample. Here, we set out to test

the hypothesis that the shared neural basis of those social symptoms of TS is an overactive MNS. We

measured behavioral and neural correlates of empathy for pain in TS and related these to the patients'

level of experimentally induced echophenomena.

A well-established neural signature of the human MNS, which is measurable with EEG, is

somatosensory mu suppression. Decrease of mu frequency amplitude has been shown in both action

observation as well as action execution (for a meta-analysis, see Fox et al., 2016), confirming the

mirroring property. Furthermore, increased mu suppression has been linked with empathic abilities in

several studies looking at empathy for pain (Cheng et al., 2008; Gallo et al., 2018; Peled-Avron et al.,

2018; Perry et al., 2010). To test, whether an overactive MNS is the basis of echophenomena as well

as altered social behaviors in TS, we thus administered an established empathy for pain paradigm

(Hoenen et al., 2015) in a group of TS patients and age-, sex- and education matched controls. Our

obtained measures were somatosensory mu suppression as a putative marker for the MNS, the skin

conductance response (SCR) as an autonomous response to other's pain, and behavioral evaluations

of the experienced pain as a marker of empathic abilities. To pull apart empathy and perspective taking,

we manipulated not only the observed pain-level (painful vs. neutral situation), but also the pain

sensitivity of the observed actor (normal vs. enhanced pain sensitivity). In TS, we expected an increase

in behavioral empathy, associated with stronger mu suppression and increased SCR to painful stimuli

as neurophysiological markers of increased personal distress. We predicted that deficits in

downregulating the automatic sharing of the actors´ pain should result in impaired differentiation

between one´s own and the other's pain sensitivity in TS. Finally, we investigated the relation between

experimentally induced echophenomena and behavioral and neurophysiological measures of empathy.

We expected that the hypothesized overactive MNS in individuals with TS would lead to increased

mirroring of others' actions, evident in echophenomena, as well as automatic sharing of emotional

states and less distinction between own and other's perspective, observable as an increase in empathy

and decreased perspective taking.

Methods

Participants

Data of N = 50 participants were collected from February 2020 to August 2021. Of those, n = 25 had

been diagnosed with TS. They were recruited from the specialized TS outpatient clinic in the Center for

Integrative Psychiatry at the University Medical Center Schleswig- Holstein, Campus Lübeck and from

the Dept. of Psychiatry, Hannover Medical School, where they received a clinical diagnosis by

experienced neurologists (A.M., K. M.-V.). Patients were diagnosed according to DSM-5 criteria

(American Psychiatric Association, 2013). Based on the collected clinical measures, four TS

participants were excluded from all data analyses due to severe depression (n = 2), ongoing substance

use disorder (n = 1), and subpar IQ (n = 1). Of the remaining n = 21 participants with TS (17 male, mean

age (SD) = 33.0 years (14.0), mean years of education (SD) = 15.7 years (2.7)), four were taking

antipsychotics (one each was taking pimozide, olanzapine, aripiprazole and amisulpride) and three

were medicated with cannabinoids. Dosages were stable in all patients. Three were left-handed.

Additionally, n = 25 healthy controls (HC) (20 male, mean age (SD) = 33.6 years (13.6), mean

years of education (SD) = 15.8 years (2.2)) were individually matched to the TS participants regarding

sex, age and education. They were recruited by means of poster and online advertisements as well as

the university email lists. All controls were free of clinically relevant psychiatric and neurologic

symptomatology at the time of testing. One control participant was left-handed. All participants gave

their written informed consent to participate in the study in accordance with the Declaration of Helsinki

(1964). The local ethics committee gave their approval of the overall project (17-167).

Clinical, personality and intelligence measures

For matching purposes, we measured general intelligence in all participants with four subtests of the

Wechsler Adult Intelligence Scale (WAIS-IV; Hartman, 2009; subtests: similarities, picture completion,

arithmetic, coding). We further assessed psychiatric symptomatology with the Mini International

Neuropsychiatric Interview (M.I.N.I.; Sheehan et al., 1998). Symptoms of obsessive compulsive

disorder (OCD) were scored using the Yale Brown Obsessive Compulsive Scale (YBOCS; Goodman

et al., 1989) and the Obsessive-Compulsive Inventory (OCI; Foa et al., 2002). Severity of attention

deficit hyperactivity disorder (ADHD) symptoms was scored using the ADHD-Index scale of the Conners

Adult ADHD Rating Scale (CAARS; Conners et al., 1999). Depressive symptoms were measured with

the revised version of the Beck-Depression-Inventory (BDI-II; Beck et al., 2009).

To measure TS symptom severity, several questionnaires were administered on the day of the

measurement. Tic severity was assessed using the Yale Global Tic Severity Score (YGTSS; Leckman

et al., 1989) and the Adult Tic Questionnaire (ATQ; Abramovitch et al., 2015), and premonitory urges

using the Premonitory Urge for Tics Scale (PUTS; Woods et al., 2005). Lifetime occurrence of TS

symptoms was assessed with the Diagnostic Confidence Index (DCI) (Robertson et al., 1999). For all

participants, standardized video recordings were taken and scored using the Modified Rush Video-

Based Tic Rating Scale (MRVS, Goetz et al., 1999), which provides a total tic score ranging from 0 to

20.

To assess self-reported empathy in our participants, the Interpersonal Reactivity Index (IRI;

Davis, 1980) was used. This measure includes affective as well as cognitive aspects of empathy and

is divided into four subscales, three measuring aspects of affective (fantasy, empathic concern and

personal distress) and one measuring cognitive empathy (perspective taking).

All questionnaires were administered in their German versions. Structured interviews (YGTSS,

YBOCS, M.I.N.I, DCI) were done by trained interviewers, the rest of the questionnaires were filled out

by the participants either online in the week prior to the measurement or in paper and pencil format on

the day of the measurement. TS symptom questionnaires were always administered on the day of the

measurement.

Procedure

Each participant was asked not to take any drugs or alcohol 24 hours prior to the appointment, but

exemptions were made for necessary medications. Testing started with the IQ test, followed by two

short video recordings for the MRVS (10 min) and the "Echometer" (5 min, explained below).

Participants were then prepared for the EEG and SCR measurement and underwent two paradigms, a

paradigm investigating provoked speech errors (50 min, unrelated to the present topic of investigation

and not included in this paper) and the empathy for pain paradigm (25 min, explained below). The order

of the two paradigms was counterbalanced across participants. After the EEG measurement,

participants took a short break. Finally, we administered the structured interviews (M.I.N.I, Y-BOCS),

all participants filled out the empathy questionnaire (IRI) and TS participants filled out the

aforementioned TS symptom questionnaires. Before leaving, participants were debriefed by the

experimenters. The whole appointment lasted around five to six hours. Participants were compensated

with 15 Euros per hour.

Echometer

To assess participants' tendency for echophenomena, we used a similar paradigm as Finis and

colleagues (2012). In the paradigm, the participants observe short video clips while they are

simultaneously videotaped, to capture any imitation of movements in the videos. Six different video

clips with a duration of three seconds (1 s fixed-image, 1 s tic of a person with TS or spontaneous

movement of an actor, 1 s fixed-image) were presented. The video clips showed head and shoulder

view segments of one of three individuals with TS (2 male) or one of three actors (2 male). Video clips

of the individuals with TS stemmed from previous ten-minute recordings of another study and showed

naturally occurring tics. Video clips of the actors were created to match those of the patients and showed

spontaneous movements. Each clip included one movement; for spontaneous movements: eyes

widening, one eyebrow up and tongue protrusion; for tics: mouth opening, eyebrows up and eyes rolling.

Each video was presented five times in a fixed order with a fixation cross (5 s) in between each clip.

Thus, each of the 30 trials lasted eight seconds.

173   Participants were knowingly videotaped using a camera placed on top of the screen. They were

instructed to watch the video clips and to fixate on the fixation cross shown in between the video clips.

To assure attention, participants were asked to remember which person performed which movement.

After the last video clip, the question "Which movement was performed by this person?" was posed,

and participants were requested to perform the correct movement. Participants were not instructed to

imitate any movements during the task and thus were blinded to the aim of the paradigm.

Empathy for pain paradigm

Our empathy for pain paradigm was similar to that of Hoenen and colleagues (2015). Participants were

seated in front of a computer screen and asked to watch 160 pictures. Each picture was presented for

2.5 s followed by a 2.5 s fixation cross (Fig.1). The stimulus set consisted of 80 pictures depicting painful

Figure 1. Empathy for pain paradigm **a:** Condition matrix with examples of stimuli. **b:** Trial sequence.
Only 25% of the trials were rated.

or neutral situations, taken from the Visually induced pain empathy repository (VIPER; Paulus et al.,

2017). The pictures were presented in four blocks, each consisting of 40 pictures (20/20 painful/neutral).

Before each block, information about the pain sensitivity of the actors in the pictures was provided. In

two blocks, participants were told that the actors have a normal pain sensitivity. In the other two blocks,

participants were told that the actors have enhanced pain sensitivity, resulting in pain experience even

after mild tactile stimulation. The blocks were presented in counterbalanced order and pictures in the

blocks were randomized per participant. In 25% of the trials, participants were asked to rate the

painfulness of the situation for the actor and for themselves if they were in that situation. All participants

rated the same stimuli. Overall, participants watched 80 pictures twice, once in each pain sensitivity

condition, resulting in 160 trials. Overall duration of the paradigm was 25 minutes.

Electrophysiological recordings

Electroencephalographic data (EEG) were recorded with 59 Ag/AgCl electrodes placed on an elastic

cap according to the international 10–20 system using BrainVision Recorder (BrainProducts GmbH)

with the Brainamp MR plus amplifier. An online reference was placed on the left earlobe, and an off-

line reference electrode was placed on the right earlobe. Horizontal and vertical EOG was recorded

with four electrodes placed next to the outer corners of the eyes and above and below the left eye.

Sampling rate was 500 Hz, and data were recorded with an online high-pass filter of 0.016 Hz and a

notch filter at 50 Hz. Impedances were kept below 10 kΩ.

Skin conductance response (SCR) was recorded using two electrodes attached to the medial

phalanges of the index and middle fingers of the left hand. The online reference was placed on the left

forearm and recorded with the same amplifier (BrainAmp ExG, BrainProdcts GmbH). As SCR data were

measured simultaneously with EEG data, the sampling rate was 500 Hz, and an online high-pass filter

of 0.016 Hz as well as a notch filter of 50 Hz were applied.

Behavioral data analysis

Videotapes of the participants during the Echometer were rated by two independent raters, who were

blinded to the observed videos. We calculated echophenomena frequency for exact repetitions of the

observed action (i.e., eye rolling when observing eye rolling) as well as movements in the same body

part (i.e., eye movements when observing eye rolling) in the 8 seconds after onset of the video. For

group comparisons, we calculated independent t-tests for both measures.

For the pain ratings in the empathy for pain paradigm, we calculated mixed ANOVAs for both

the rating of others' pain as well as the rating of one´s own supposed pain. Group was used as a

between group factor (TS vs. HC) and action (painful vs. neutral) and sensitivity (normal vs. enhanced)

served as within subject factors. For correlating behavioral results with other measures, we calculated

the four difference values for each combination of conditions (e.g., difference of action (painful – neutral)

in normal sensitivity condition or difference of sensitivity (enhanced – normal) in painful condition) to

get a measure of the magnitude of the action or sensitivity effect in either condition. All described

correlations are Pearson's correlations

EEG data analysis

All data preprocessing was done in MATLAB®, using the *eeglab* toolbox (Delorme & Makeig, 2004).

During offline processing, data were down-sampled to 250 Hz, re-referenced to left earlobe and filtered

with a high-pass filter at 0.5 Hz and a low-pass filter at 44 Hz. The signal was segmented into epochs

from -500 ms to 2500 ms around the beginning of stimulus presentation. Epochs with obvious

movement artifacts due to voluntary movements or tics were manually removed as well as channels

with very noisy data. Eye movements and further muscle artifacts and channel noise were corrected

using independent component analysis (ICA). Visual screening and the automated classifier ICLabel

(Pion-Tonachini et al., 2019) were used to classify components. On average, 5.9 components were

removed per participant. After the ICA, the epoch baseline (-500 0) was removed from each epoch.

Then, epochs with remaining artifacts were manually rejected and previously rejected channels were

interpolated. On average, 27 epochs were excluded (range: 1 – 86) and 0.7 channels interpolated

(range: 0 – 6) per participant. Finally, we calculated the surface Laplacian as suggested by Perrin and

colleagues (1989) and implemented in MATLAB® by Cohen (2014).

Participants with more than 50 % of rejected trials or less than 15 trials per combination of

conditions were excluded from further analyses. We chose this relatively high threshold, as we had to

exclude many trials in the TS group due to movement artifacts because of tics. We had already

anticipated this and thus chose a high number of trials to begin with. One participant from the TS group

was excluded and none of the healthy controls. Thus, analyses of EEG data are conducted on n = 20

TS patients and n = 25 healthy controls.

On the preprocessed data, we did a time-frequency analysis. We calculated trial-averaged

power for each subject, channel and experimental condition for linearly spaced frequencies from 1 Hz

to 30 Hz using Morlet wavelet convolution with a variable number of wavelet cycles. Wavelet cycles

increased logarithmically from 4 cycles to 15 cycles as a function of frequency. We report decibel (dB)

normalized data, using -400 to -100 ms pre stimulus as a baseline.

247   For the statistical analysis, we took the results of the time-frequency analysis and calculated

mean power in the mu frequency band (8–12 Hz) for the time window of interest, 500 to 1500ms after

stimulus onset. We chose this time window based on the suggestions of Whitmarsh and colleagues

(2011), who underlined the importance of not including the time directly after stimulus onset as to not

confound the modulation of ongoing alpha activity and stimulus-onset evoked responses. As we were

interested in sensorimotor mu suppression, we focused our analysis on a cluster of central electrodes,

including C1, C2, C3, C4, Cz, CPz, CP1, CP2, CP3 and CP4. The occipital cluster, including O1, O2

and Oz, served as a control. Power changes were then compared using a mixed ANOVA with group

(TS vs. HC) as a between subject factor and action (painful vs. neutral), sensitivity (normal vs.

enhanced), region (central vs. occipital) and laterality (right vs. middle vs. left) as within subject factors.

Additionally, we calculated Bayesian statistics for the time frequency data, which enabled us to make

statements for the null hypothesis. Calculations were done using JASP (Version 0.16.4). We report the

results of the Bayesian model averaging (BFincl) implemented as analysis of effects across matched

models. We interpret the Bayes factors according to common guidelines, suggesting values > 3 to

provide moderate and values > 10 to provide strong support for the alternative hypothesis. Values <

0.33 provide moderate and values < 0.1 strong support for the null hypothesis (van Doorn et al., 2021).

263   For correlating EEG results with other measures, we used the same approach as with the

264   behavioral data and calculated the four difference values for each combination of conditions.

265   To check, whether there are group differences in mu power independent of our task

manipulations, we looked at absolute and relative mu power in all central clusters, separately for the

time of stimulus presentation and the baseline. Absolute power spectral density of the mu band (8-

12Hz) was obtained by applying a fast Fourier transform and averaging mu power per cluster and

participant. Relative mu power was calculated by dividing mu power spectral density by the absolute

power of all frequencies within 1 and 30 Hz. Group differences were tested by a mixed ANOVA with

group (TS vs. HC) as a between subject factor and region (central right vs. central middle vs. central

left) as a within subject factor.

SCR data analysis

Analysis of skin conductance was performed using the MATLAB® toolbox *Ledalab* (Benedek &

Kaernbach, 2010). The signal was downsampled to 25 Hz and then smoothed with a Gaussian low-

pass filter. Obvious artifacts were removed manually. Then, the signal was deconvolved using a

continuous decomposition analysis (CDA) with four optimizations. This separates the signal into

continuous traces of tonic and phasic activity. We proceeded to analyze the phasic driver component,

by extracting averaged event-related responses via area under the curve, between 1 to 4 seconds after

stimulus onset. Event-related phasic SCR were range normalized and log transformed across all trials

within each subject. This ensured that individual variability in the signal magnitude would not bias further

analyses. SCR per trial were then averaged within conditions. One participant from each group was

excluded, as the data quality did not allow measurements of any SCR. Thus, analyses of SCR data

were conducted on n = 20 TS patients and n = 24 healthy controls. SCR were analyzed similarly to the

EEG data with a mixed ANOVA with the factors group, action and sensitivity. For correlation analyses,

difference scores were calculated analogous to the behavioral and EEG data.

Results

Demographics, clinical and questionnaire results

Sex, age, and IQ did not differ between TS patients and healthy controls (Table 1), demonstrating

successful matching procedures. TS patients showed significantly more obsessive-compulsive

symptoms, depressive symptoms and symptoms of ADHD. Severity of TS symptoms in our TS group

was comparable to similar studies (i.e.: Brandt et al., 2016; Eddy et al., 2011, 2013; Jonas et al., 2010).

Regarding empathic abilities, TS participants reported significantly lower fantasy abilities in the IRI. The

two groups did not differ regarding their trait empathic concern, personal distress and perspective taking

(Fig. 2a). Additionally, we had hypothesized a correlation between tic severity and trait empathy.

Indeed, the YGTSS total tic score correlated with self-reported personal distress in the empathy

questionnaire (IRI; r = 0.57, p =.009), with higher tic severity relating to more reported personal distress

(Fig. 2b). Other subscales of the IRI did not significantly correlate with the YGTSS total tic score.

304   Table 1:

305   *Overview of personality and clinical measures*

Questionnaire	TS	HC	*t*	*df*	*p*	Cohen's *d*
Sex	17 m, 4 f	20 m, 5 f				
Age	33.00(14.04)	33.56(13.61)	0.14	42.16	.892	0.04
IQ	106.20(9.06)	104.38(10.97)	0.62	43.99	.541	0.18
Obsessive-compulsive behavior						
YBOCS thoughts	3.48(4.13)	0.20(1.00)	-3.55	21.97	.002	-1.14
YBOCS actions	4.24(4.00)	0.20(1.00)	-4.51	22.11	< .001	-1.44
YBOCS total	7.71(7.18)	0.40(2.00)	-4.52	22.61	< .001	-1.45
OCI	19.71(14.69)	8.88(7.24)	-3.08	28.02	.005	-0.96
Depression	11.14(10.69)	5.00(4.84)	-2.44	26.89	.021	-0.77
ADHD	11.29(6.82)	6.64(4.70)	-2.64	34.55	.012	-0.81
Empathy (IRI)						
Fantasy	10.05(3.72)	12.29(3.00)	2.17	36.35	.037	0.67
Empathic Concern	13.5(2.65)	13.8(2.45)	0.39	39.35	.698	0.12
Personal Distress	9.80(3.11)	10.12(3.10)	0.34	40.83	.733	0.10
Perspective Taking	14.05(3.36)	14.56(2.43)	0.57	33.59	.573	0.18
TS Symptomatology						
MRVS total score	11.90(4.20)	2.44(2.29)	-9.25	29.74	< .001	-2.87
YGTSS Total motor tic score	13.81(3.39)					
YGTSS Total vocal tic score	8.38(6.06)					
YGTSS Total tic score	22.19(7.44)					
YGTSS Global severity score	43.00(15.65)					
ATQ	37.71(25.50)					
PUTS	21.14(4.98)					
DCI	60.86(18.02)					

Note. Questionnaire data separated by group. Average scores plus standard deviations in brackets. m
= male, f = female. IQ, Intelligence Quotient as measured by subtests of the Wechsler Adult Intelligence
Scale (WAIS-IV). YBOCS, Yale Obsessive Compulsive Inventory. OCI, Obsessive Compulsive
Inventory. ADHD, Attention Deficit Hyperactivity Disorder as measured by the Conners Adult ADHD
Rating Scale. IRI, Interpersonal Reactivity Index. MRVS, Modified Rush Video-Based Tic Rating Scale.
YGTSS, Yale Global Tic Severity Scale. ATQ, Adult Tic Questionnaire. PUTS, Premonitory Urges for
Tics Scale. DCI, Diagnostic Confidence Index. TS, Tourette Syndrome. HC, Healthy Controls.
313

Behavioral Task Results

The analysis of echophenomena yielded clear group differences for both exact imitations (t(1,32.58) =

-3.49, p = .001, d = -1.11) as well as movements in the same body part (t(1,35.58) = -3.98, p < .001, d

= -1.24). As expected, participants of the TS group showed a higher tendency to echo what they observe

than the healthy controls (Fig. 2c).

Figure 2. Clinical and behavioral results. **a:** Group comparisons in the Interpersonal Reactivity Index
(IRI). **b:** Correlation in the TS group of the subscale personal distress in the IRI and total tic score as
measured by the Yale Global Tic Severity Scale (YGTSS). **c:** Number of echoes in Echometer. Both
exact replications of the observed movement and movements in the same body part were considered
for the analyses
325
326   In the empathy for pain paradigm, we compared the pain ratings per condition and found an

interaction of action and sensitivity on ratings of others' pain (F(1,44) = 21.69, p < .001, η^2 = .02), with

all four pairwise comparisons being significant. When looking at the ratings of the pain that participants

themselves would feel in the observed situation, we again found an interaction of action and sensitivity

(F(1,44) = 4.16, p = .047, η^2 < .01). However, pairwise comparisons revealed no differences between

sensitivity conditions in both painful and neutral stimuli. This shows that both HC and TS performed as

expected: They rated the other's pain higher in the painful condition compared to the neutral condition;

and they rated the other's pain higher in the enhanced pain sensitivity condition compared to the normal

pain sensitivity condition. When rating their own imagined pain, participants made no difference

between sensitivity conditions but rated painful stimuli as more painful than neutral ones. However, we

did not find any group effects in the ratings. The three-way interaction of group, action and sensitivity

on the rating of others' pain was not significant ($F(1,44) = 0.13$, $p = .72$, $\eta^2 < .01$, Fig. 3), although the

differences between sensitivity conditions in the TS group were nominally smaller. On the behavioral

level, the experimental manipulations were thus successful, but none of the hypothesized group

differences were confirmed.

Figure 3. Pain ratings in empathy for pain paradigm. **a:** Rating of painfulness for the actor in the picture
from „not at all" to „very" under the normal (yellow) and enhanced (blue) sensitivity condition for both
healthy controls (HC, left side) and Tourette patients (TS, right side). **b:** Rating of own imagined pain in
the depicted situation on same scale.
345

Electrophysiological results

The mu suppression in the empathy for pain paradigm showed an interaction of all factors (group,

action, sensitivity, region & laterality): $F(2,82) = 3.37$, $p = .039$, $\eta^2 < .01$ (Fig. 4). To understand this

interaction, we calculated ANOVAs separately for central and occipital sites. At occipital electrodes, we

found a main effect of action ($F(1,43) = 13.80$, $p = .001$, $\eta^2 = .01$), indicating more pronounced alpha

suppression after observing painful compared to neutral actions. At the central electrodes, we found a

significant interaction of group, action and sensitivity ($F(1,41) = 6.52$, $p = .014$, $\eta^2 < .01$, BF = 4.58).

Following ANOVAs for each of the groups revealed no significant interaction of action and sensitivity in

the TS group ($F(1,17) = 2.24$, $p = .153$, $\eta^2 < .01$, BF = 2.62), nor any other effects. In HC we found an

interaction of action and sensitivity ($F(1,24) = 5.11$, $p = .033$, $\eta^2 < .01$, BF = 2.26). HC showed a simple

main effect of action in the normal sensitivity condition ($F(1) = 5.34$, $p = .03$) but not in the enhanced

sensitivity condition ($F(1) = 0.03$, $p = .86$). Finally, there were no group differences in absolute or relative

mu power, neither during the baseline period (absolute: $F(1,43) = 0.07$, $p = .790$, $\eta^2 < .01$; relative:

$F(1,43) = 0.14$, $p = .707$, $\eta^2 < .01$) nor during the stimulus presentation (absolute: $F(1,43) = 1.98$, $p =$

.166, $\eta^2 = .03$; relative: $F(1,43) = 0.76$, $p = .387$, $\eta^2 = .02$).

361   Summarizing, we found the expected increase of mu suppression after observing painful

compared to neutral pictures in our control group but only in the normal pain sensitivity condition. TS

participants did not show any significant modulation of mu suppression by the experimental

manipulations. Note however that the results of the Bayesian analyses show that the evidence for the

interaction effect (action and sensitivity) within the HC and TS groups were similarly equivocal though.

This speaks for an overall weak effect of the pain manipulation, and no clear evidence for a lacking

pain-related mu modulation in TS. However, the moderate evidence for the interaction of experimental

conditions with group speaks for a reduced mu-effect in TS compared to the controls. Finally, the groups

had the same level of absolute and relative mu power during the baseline, making it unlikely that

differences in baseline mu activity account for the different results in the groups.

371
372   **Figure 4.** Baseline corrected mu power (8 – 12 Hz) in empathy for pain paradigm. *Top row:* Spatial
373   distribution of mu power differences (painful-neutral) in time-window of interest (500 – 1500 ms). *Bottom
374   row:* Time course of mu power in right central cluster for the painful (red) and neutral (black) condition,
375   separately for the controls (HC, left columns) and Tourette patients (TS, right columns) and for normal
376   and enhanced sensitivity. Time-point 0 refers to the picture onset.
377
Analysis of the skin conductance response revealed a significant interaction of group, action

and sensitivity ($F(1,42) = 5.40$, $p = .025$, $\eta^2 = .01$; Fig. 5). ANOVAs per group revealed a main effect of

action in the TS group ($F(1,19) = 4.39$, $p = .050$, $\eta^2 = .01$), with participants showing an enhanced skin

conductance response after watching painful actions. For the healthy controls we found an interaction

of action and sensitivity ($F(1,23) = 6.24$, $p = .020$, $\eta^2 = .01$), with nominally stronger responses after

painful pictures in the enhanced sensitivity condition. Pairwise comparisons were not significant.

Correlations of echophenomena with empathy measures

Based on our hypothesis on a shared mechanism behind echophenomena and empathy, we expected

the number of echophenomena to correlate with our behavioral, physiological and neural indices of

empathic abilities. To test this, we focused on the difference of action (painful – neutral) in the normal

sensitivity condition. However, echophenomena did not correlate with trait personal distress ($r = 0.22$,

$p = .15$), behavioral ($r = 0.27$, $p = .08$), physiological (SCR: $r = 0.11$, $p = .50$) or neural empathy measures

(mu: $r = 0.15$, $p = .35$). Additionally, we hypothesized a correlation between tic severity and empathy.

However, no correlations could be found between tic severity measures of the YGTSS or the MRVS

video protocol with behavioral, neural or physiological empathy measures.

Figure 5. Skin conductance response in empathy for pain paradigm. Plotted are the time courses of
the adjusted phasic driver for the painful (red) and neutral (black) pictures, separately for the controls
(HC, left columns) and Tourette patients (TS, right columns) and for normal and enhanced sensitivity.
Time-point 0 refers to the picture onset. Statistical analysis is based on our window of interest (1500 -
3000 ms).
398
Discussion

In this study, we tested the hypothesis that echophenomena in TS are related to increased mirror

neuron system activity and increased automatic sharing of others' emotional states. As expected, we

found that participants with TS showed more echophenomena than controls. However, we did not find

any empathy related behavioral differences between TS and HC and no differences in trait empathy or

skin conductance between our groups. Interestingly, the TS group showed reduced pain-related mu

suppression, indicating, contrary to our hypothesis, decreased mirror neuron system activity in

response to others' emotional states.

Echophenomena and trait empathy in TS

In line with a previous study (Finis et al., 2012), we found a strong difference between TS and controls

in the frequency of echophenomena induced by our Echometer paradigm. This was true for both exact

replications of the observed movement as well as movements with the same body part. Furthermore,

all participants with TS showed echoes, which stresses the high prevalence of echophenomena in this

patient group. This is in contrast to previous, lower estimates of its prevalence, usually ranging below

fifty percent (Cavanna et al., 2011; Lees et al., 1984). The difference is most likely due to the mode of

measurement, as prevalence estimates in these studies were based on self-report or clinicians'

observations. It is thus important for future explorations of the neural basis of echophenomena to

experimentally measure their occurrence. Additionally, echoes were not only seen in the TS patient

group but also, with much lower frequency, in the healthy controls, underlining that imitating others is a

normal process and not per se pathological (Ganos et al., 2012). In conclusion, our TS sample exhibited

the expected increased tendency for echophenomena compared to controls and should thus, according

to our hypothesis, show further symptoms of altered social behaviors and an overactive MNS.

421   One of the previous findings which had prompted our hypothesis was the report of increased

self-reported personal distress, combined with reduced perspective-taking in the Interpersonal

Reactivity Index (IRI) in a relatively large TS sample (Eddy et al., 2015). The IRI personal distress

subscale measures unease and anxiety in negative emotional settings and is related to poor social

functioning, whereas the perspective taking subscale measures the tendency to think about the other´s

point of view and is correlated with higher social functioning (Davis, 1983). We utilized the same

questionnaire but could not replicate the results in our sample. TS participants in our study did not report

increased personal distress or reduced perspective taking compared to the controls. We did find an

increase of reported personal distress with higher tic severity, but this was also contrary to what Eddy

and colleagues reported (2015), as they found a *negative* correlation between tics and personal

distress. Although our sample was smaller compared to theirs, our effect sizes were also considerably

smaller, which makes it unlikely that we missed the effect only because of a smaller sample.

Furthermore, our sample included far less medicated participants, but comorbidities were comparable

across both samples. TS participants in our study did report lower fantasy scores compared to controls,

indicating a decreased tendency to imaginatively transpose themselves into fictional situations. Overall,

we found group differences in self-reported empathy but those were not consistent with the previous

report of Eddy and colleagues (2015). Additionally, the results of the empathy self-report did not

correlate with the tendency for echophenomena.

Empathy for pain in TS

In the empathy for pain paradigm, both groups performed similarly and in line with the task manipulation,

showing that task instructions were understood by both groups. We had hypothesized increased

painfulness ratings after painful stimuli for our TS group, which we could not confirm. However, one

could argue that the pain ratings in our paradigm did not reflect only automatic sharing of the experience

of the actor but also cognitive understanding of the situation. Furthermore, the pain stimuli were not

overly graphic and did not show obvious signs of extreme pain, like blood, broken limbs, painful facial

expressions or body language, and might thus not elicit strong discomfort in the viewer. Hence, our

paradigm might not have been sensitive enough to detect subtle behavioral differences between the

groups. Moreover, TS might still show altered neural processing even in the absence of obvious

behavioral differences.

Physiologically, increased distress due to increased sharing should have been visible in a

stronger skin conductance response in the TS group. However, we found no such group differences. If

anything, results suggested a slightly reduced modulation of skin conductance response by the

sensitivity manipulation in the TS group. Previously, the magnitude of SCR to observing other's in pain

has been linked to prosocial behavior (Hein et al., 2011) and implicit race bias (Forgiarini et al., 2011),

with higher SCRs corresponding to increased empathy and prosocial behavior. In conclusion, our

results could indicate slightly reduced empathy in TS compared to controls.

In terms of the neural data, several studies have probed mu suppression under different

conditions or in different groups with an almost identical setup as ours: participants observing pictures

of body parts in painful or neutral situations (Arnett et al., 2019; Hoenen et al., 2015; Li et al., 2017;

Yang et al., 2009). Those studies consistently find increased mu suppression during the observation of

painful compared to neutral pictures. The magnitude of mu suppression is interpreted as a measure of

empathy and is modulated by several factors, with less mu suppression being observed in men than

women (Yang et al., 2009), during negative mood compared to positive or neutral mood (Li et al., 2017)

and after reported parental emotional invalidation (Arnett et al., 2019). In line with those findings, our

healthy controls showed the expected pain-related mu suppression over the right sensorimotor cortex.

This was modulated by our additional manipulation of the actors' pain sensitivity. Suppression only

differed between painful versus neutral stimuli when the pain sensitivity of the actor was normal. This

effect vanished under the enhanced pain sensitivity condition, indicating that controls processed both

kinds of stimuli as painful in this condition. Our results thus mirror those of Hoenen and colleagues

(2015), who originally developed this paradigm and found increased mu suppression in trials where

participants were informed that the actors have an increased pain sensitivity. In a similar setup,

participants observed hands either being pricked by a needle or being touched by a Q-tip (Perry et al.,

2010). Participants were asked to assume either normal pain reactions or switched pain reactions,

where needle pricks were not painful, but Q-tip touches were. When participants assumed normal pain

reactions, the classic increase of mu suppression in painful over neutral situations was observed.

However, when participants assumed switched pain reactions, mu suppression was equally large in

painful and neutral situations. Those results combined with what we observed in the healthy controls

indicate that mu suppression is not only elicited by automatic empathic reactions to the observed stimuli

but can be modulated by top-down processes, making it possible to feel empathy for pain even in

situations that would not be painful for the participants themselves. Interestingly, TS participants did not

show any significant pain-related attenuation of mu power, although Bayesian analyses indicated

equivocal evidence for and against this effect. However, this group difference in pain-related mu

suppression speaks against our hypothesis of increased automatic sharing in TS, but rather points in

the opposite direction. One might argue that a generally overactive MNS in TS could lead to less

pronounced mu attenuation during our paradigm. However, the TS group showed similar absolute and

relative mu power during the baseline as well as averaged across all stimulus presentations. This

questions that the group differences can be attributed to baseline differences and a generally overactive

MNS.

Is an overactive MNS in TS causing echophenomena and changes in empathy?

What are the implications for our hypothesis that an overactive MNS is the basis of both altered social

behaviors and echophenomena? First of all, we did not find any alterations in empathic behavior in the

TS group and can thus not draw any conclusion on possible causes of altered social behavior in TS.

Second, we did not find evidence that an overactive MNS is at the core of the clearly increased tendency

for echophenomena in our TS sample. On the contrary, we found less MNS activity in the TS group

during the empathy for pain task.

Our results might thus support an alternative hypothesis about the basis of echophenomena.

Rather than echophenomena being indicative of a generally increased tendency to imitate, they might

reflect the failure to inhibit imitation in contexts where it is not useful (Ganos et al., 2012). Thus, in

situations where imitation has a social function and can be found in healthy adults, we find no increase

of imitation in TS compared to controls. It is in situations where imitation serves no purpose, such as

our Echometer paradigm, that we find increased imitation in TS. This also fits with previous results in

an automatic imitation paradigm (Jonas et al., 2010). Both TS and HC showed a comparable modulation

of reaction times by incongruent movements compared to congruent movements, indicating similar

automatic imitation tendencies. In conclusion, in social contexts where it is beneficial, such as an

empathy paradigm, imitation or automatic sharing of the other´s state does not seem to be increased

in TS compared to controls. This interpretation is further supported by the fact that we found no

relationship between the extent of echophenomena or tic severity and the behavioral and neural results

in the empathy for pain paradigm.

However, not only did we not find an increase in automatic sharing in our empathy paradigm,

our results even pointed in the opposite direction, with TS participants showing less pain-related mu

suppression than controls. This fits to an additional hypothesis brought forward by Ganos and

colleagues (2012). They propose that people with TS compensate the deficient inhibition in motor

networks with increased activity in inhibitory circuits outside the motor network, presumably prefrontal

areas. They thus assume that compensatory activity in the TS brain leads to less responsiveness to

experimental manipulations. It is feasible that due to the failure to inhibit tics or echophenomena in

inappropriate contexts, the TS brain overcompensates by generally decreasing the responsiveness to

stimuli. This fits to recent Go/No-Go studies in TS, which found reduced task-related activations

compared to healthy controls (Niccolai et al., 2016; Thomalla et al., 2014). In one study, task-related

BOLD signal change was decreased in cortical motor areas but not in other areas in people with TS.

This was accompanied by poorer task performance compared to healthy controls (Thomalla et al.,

2014). In another Go/No-Go task, participants with TS showed less beta suppression in the

sensorimotor area but enhanced beta power in parieto-occipital brain regions compared to healthy

controls, but task performance was the same in both groups (Niccolai et al., 2016). Similarly, in the

aforementioned automatic imitation task (Jonas et al., 2010), where TS and HC showed no differences

in the interference effect, people with TS exhibited less movement facilitation by observing congruent

biological movements than healthy controls. However, they showed more interference than healthy

controls when observing incongruent biological movements compared to the observation of non-

biological movement, hinting to an overactive imitation inhibition system that is active during the task.

To conclude, our findings of decreased mu modulation in TS compared to the controls might be

explained by increased inhibition. However, it is important to note, that none of these studies or our

study provided direct neural evidence for increased inhibitory activity.

Limitations

Our groups were of moderate size, which might have led us to miss small group differences. The results

of Bayes statistics underlined this problem, as we found moderate evidence for group differences in

pain-related mu suppression in the interaction, but overall weak effects of the pain modulation in both

groups. The small magnitude of pain-related mu suppression might be partly due to our sample being

predominantly male, as male participants were found to exhibit less mu suppression than females in

previous research (Yang et al., 2009). Another caveat regarding our results is the heterogeneity of our

TS sample. TS is often accompanied by several comorbidities, and this was the same in our group.

Compared to the controls, the TS group showed more symptoms of depression, anxiety, ADHD and

OCD, all of which could have influenced our results. However, as most individuals with TS suffer from

symptoms of one or more of these comorbidities, we considered it best to include them to create a

representative sample for this clinical group. Finally, our empathy paradigm was relatively easy and

might thus not be suitable to measure subtle group differences in behavioral empathy.

Conclusions

Together, our results speak against an overactive MNS and against increased levels of empathy in TS.

In contrast, our findings indicate reduced pain-related neural processing in TS, although this had no

behavioral impact in the current task. As our TS sample evinced high levels of echophenomena, the

underlying neurocognitive processes of this symptom remain to be established. Overall, as our study,

as well as previous studies (Eddy et al., 2016, 2017; Neuner et al., 2010), imply altered processing of

stimuli in social contexts, future research should continue to explore the neural basis of social symptoms

and echophenomena in TS.

References

Abramovitch, A., Reese, H., Woods, D. W., Peterson, A., Deckersbach, T., Piacentini, J., Scahill, L., & Wilhelm, S. (2015). Psychometric Properties of a Self-Report Instrument for the Assessment of Tic Severity in Adults With Tic Disorders. *Behavior Therapy*, *46*(6), 786–796. https://doi.org/10.1016/j.beth.2015.06.002

Albin, R. L. (2018). Tourette syndrome: A disorder of the social decision-making network. *Brain*, *141*(2), 332–347. https://doi.org/10.1093/brain/awx204

American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders* (Fifth Edition). American Psychiatric Association. https://doi.org/10.1176/appi.books.9780890425596

Arnett, K., Roach, A., Elzy, M., & Jelsone-Swain, L. (2019). Childhood emotional invalidation and right hemispheric mu suppression during a pain empathy task: An EEG study. *Social Neuroscience*, *14*(2), 236–250. https://doi.org/10.1080/17470919.2018.1441905

Baron-Cohen, S., Jolliffe, T., Mortimore, C., & Robertson, M. (1997). Another advanced test of theory of mind: Evidence from very high functioning adults with autism or Asperger syndrome. *Journal of Child Psychology and Psychiatry*, *38*(7), 813–822.

Beck, A., Steer, R., & Brown, G. (2009). *Beck-Depressions-Inventar Revision: Deutsche Bearbeitung von M. Hautzinger/F. Keller/C. Kühner*. Pearson.

Benedek, M., & Kaernbach, C. (2010). A continuous measure of phasic electrodermal activity. *Journal of Neuroscience Methods*, *190*(1), 80–91. https://doi.org/10.1016/j.jneumeth.2010.04.028

Bonini, L., Rotunno, C., Arcuri, E., & Gallese, V. (2022). Mirror neurons 30 years later: Implications and applications. *Trends in Cognitive Sciences*. https://doi.org/10.1016/j.tics.2022.06.003

Brandt, V. C., Patalay, P., Bäumer, T., Brass, M., & Münchau, A. (2016). Tics as a model of over-learned behavior—Imitation and inhibition of facial tics. *Movement Disorders*, *31*(8), 1155–1162.

Brass, M., Ruby, P., & Spengler, S. (2009). Inhibition of imitative behaviour and social cognition. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1528), 2359–2367. https://doi.org/10.1098/rstb.2009.0066

Cavanna, A. E., Critchley, H. D., Orth, M., Stern, J. S., Young, M.-B., & Robertson, M. M. (2011). Dissecting the Gilles de la Tourette spectrum: A factor analytic study on 639 patients. *Journal*

585   *of   Neurology,   Neurosurgery   &   Psychiatry*,   *82*(12),   1320–1323.

586   https://doi.org/10.1136/jnnp.2010.225029

587   Channon, S., Drury, H., Gafson, L., Stern, J., & Robertson, M. M. (2012). Judgements of social

588   inappropriateness in adults with Tourette's syndrome. *Cognitive Neuropsychiatry*, *17*(3), 246–

589   261. https://doi.org/10.1080/13546805.2011.590689

590   Cheng, Y., Yang, C.-Y., Lin, C.-P., Lee, P.-L., & Decety, J. (2008). The perception of pain in others

591   suppresses somatosensory oscillations: A magnetoencephalography study. *NeuroImage*,

592   *40*(4), 1833–1840. https://doi.org/10.1016/j.neuroimage.2008.01.064

593   Cohen, M. X. (2014). *Analyzing Neural Time Series Data: Theory and Practice* (1st ed.). The MIT Press.

594   Conners, C. K., Erhardt, D., & Sparrow, E. (1999). Adult ADHD rating scales: Technical manual.

595   *Toronto, Ontario, Canada: Multi-Health Systems*.

596   Davis, M. H. (1980). *A multidimensional approach to individual differences in empathy*.

597   Davis, M. H. (1983). Measuring individual differences in empathy: Evidence for a multidimensional

598   approach.   *Journal   of   Personality   and   Social   Psychology*,   *44*(1),   113–126.

599   https://doi.org/10.1037/0022-3514.44.1.113

600   Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG

601   dynamics including independent component analysis. *Journal of Neuroscience Methods*,

602   *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

603   Drury, H., Channon, S., Barrett, R., Young, M.-B., Stern, J. S., Simmons, H., & Crawford, S. (2012).

604   Emotional processing and executive functioning in children and adults with Tourette's

605   syndrome. *Child Neuropsychology*, *18*(3), 281–298.

606   Eddy, C. M. (2018). Social cognition and self-other distinctions in neuropsychiatry: Insights from

607   schizophrenia and Tourette syndrome. *Progress in Neuro-Psychopharmacology and Biological*

608   *Psychiatry*, *82*, 69–85. https://doi.org/10.1016/j.pnpbp.2017.11.026

609   Eddy, C. M., Cavanna, A. E., & Hansen, P. C. (2017). Empathy and aversion: The neural signature of

610   mentalizing in Tourette syndrome. *Psychological Medicine*, *47*(3), 507–517.

611   Eddy, C. M., Cavanna, A. E., Rickards, H. E., & Hansen, P. C. (2016). Temporo-parietal dysfunction in

612   Tourette syndrome: Insights from an fMRI study of Theory of Mind. *Journal of Psychiatric*

613   *Research*, *81*, 102–111.

Eddy, C. M., Macerollo, A., Martino, D., & Cavanna, A. E. (2015). Interpersonal reactivity differences in

Tourette   syndrome.   *Psychiatry   Research*,   *228*(3),   932–935.

https://doi.org/10.1016/j.psychres.2015.05.070

Eddy, C. M., Mitchell, I. J., Beck, S. R., Cavanna, A. E., & Rickards, H. (2010a). 024 Social and

economic reasoning in Tourette syndrome. *Journal of Neurology, Neurosurgery & Psychiatry*,

81(10), e9–e10. https://doi.org/10.1136/jnnp.2010.217554.24

Eddy, C. M., Mitchell, I. J., Beck, S. R., Cavanna, A. E., & Rickards, H. (2011). Social reasoning in

Tourette   syndrome.   *Cognitive   Neuropsychiatry*,   *16*(4),   326–347.

https://doi.org/10.1080/13546805.2010.538213

Eddy, C. M., Mitchell, I. J., Beck, S. R., Cavanna, A. E., & Rickards, H. E. (2010b). Altered Attribution

of Intention in Tourette's Syndrome. *The Journal of Neuropsychiatry and Clinical

Neurosciences*, *22*(3), 348–351. https://doi.org/10.1176/jnp.2010.22.3.348

Eddy, C. M., Rickards, H. E., Critchley, H. D., & Cavanna, A. E. (2013). A controlled study of personality

and   affect   in   Tourette   syndrome.   *Comprehensive   Psychiatry*,   *54*(2),   105–110.

https://doi.org/10.1016/j.comppsych.2012.07.004

Falkai, P., Wittchen, H.-U., Döpfner, M., Gaebel, W., Maier, W., Rief, W., Saß, H., & Zaudig, M. (Eds.).

(2018). *Diagnostisches und statistisches Manual psychischer Störungen DSM-5®* (2.

korrigierte Auflage). Hogrefe. https://doi.org/10.1026/02803-000

Finis, J., Moczydlowski, A., Pollok, B., Biermann-Ruben, K., Thomalla, G., Heil, M., Krause, H., Jonas,

M., Schnitzler, A., & Münchau, A. (2012). Echoes from childhood-imitation in Gilles de la

Tourette Syndrome: Echoes from Childhood-Imitation in GTS. *Movement Disorders*, *27*(4),

562–565. https://doi.org/10.1002/mds.24913

Foa, E. B., Huppert, J. D., Leiberg, S., Langner, R., Kichic, R., Hajcak, G., & Salkovskis, P. M. (2002).

The Obsessive-Compulsive Inventory: Development and validation of a short version.

Psychological Assessment, *14*(4), 485.

Forgiarini, M., Gallucci, M., & Maravita, A. (2011). Racism and the Empathy for Pain on Our Skin.

Frontiers in Psychology, *2*. https://www.frontiersin.org/articles/10.3389/fpsyg.2011.00108

Fox, N. A., Bakermans-Kranenburg, M. J., Yoo, K. H., Bowman, L. C., Cannon, E. N., Vanderwert, R.

E., Ferrari, P. F., & van IJzendoorn, M. H. (2016). Assessing human mirror activity with EEG

mu   rhythm: A   meta-analysis.   *Psychological   Bulletin*,   *142*(3),   291–313.

https://doi.org/10.1037/bul0000031

Freeman, R. D., Zinner, S. H., Müller-Vahl, K. R., Fast, D. K., Burd, L. J., Kano, Y., Rothenberger, A.,

Roessner, V., Kerbeshian, J., & Stern, J. S. (2009). Coprophenomena in Tourette syndrome.

Developmental Medicine & Child Neurology, *51*(3), 218–227. https://doi.org/10.1111/j.1469-

8749.2008.03135.x

Gallo, S., Paracampo, R., Müller-Pinzler, L., Severo, M. C., Blömer, L., Fernandes-Henriques, C.,

Henschel, A., Lammes, B. K., Maskaljunas, T., Suttrup, J., Avenanti, A., Keysers, C., &

Gazzola, V. (2018). The causal role of the somatosensory cortex in prosocial behaviour. *ELife*,

7, e32740. https://doi.org/10.7554/eLife.32740

Ganos, C., Ogrzal, T., Schnitzler, A., & Münchau, A. (2012). The pathophysiology of

echopraxia/echolalia: Relevance to Gilles De La Tourette syndrome. *Movement Disorders*,

27(10), 1222–1229. https://doi.org/10.1002/mds.25103

Goetz, C. G., Pappert, E. J., Louis, E. D., Raman, R., & Leurgans, S. (1999). Advantages of a modified

scoring method for the Rush Video-Based Tic Rating Scale. *Movement Disorders: Official

Journal of the Movement Disorder Society*, *14*(3), 502–506.

Goodman, W. K., Price, L. H., Rasmussen, S. A., Mazure, C., Fleischmann, R. L., Hill, C. L., Heninger,

G. R., & Charney, D. S. (1989). The Yale-Brown Obsessive Compulsive Scale. I. Development,

use,   and   reliability.   *Archives   of   General   Psychiatry*,   *46*(11),   1006–1011.

https://doi.org/10.1001/archpsyc.1989.01810110048007

Hartman, D. E. (2009). Wechsler Adult Intelligence Scale IV (WAIS IV): Return of the Gold Standard.

Applied Neuropsychology, *16*(1), 85–87. https://doi.org/10.1080/09084280802644466

Hein, G., Lamm, C., Brodbeck, C., & Singer, T. (2011). Skin Conductance Response to the Pain of

Others   Predicts   Later   Costly   Helping.   *PLOS   ONE*,   *6*(8),   e22759.

https://doi.org/10.1371/journal.pone.0022759

Heyes, C., & Catmur, C. (2022). What Happened to Mirror Neurons? *Perspectives on Psychological

Science*, *17*(1), 153–168. https://doi.org/10.1177/1745691621990638

Hoenen, M., Lübke, K. T., & Pause, B. M. (2015). Somatosensory mu activity reflects imagined pain

intensity of others. *Psychophysiology*, *52*(12), 1551–1558.

Jonas, M., Thomalla, G., Biermann-Ruben, K., Siebner, H. R., Müller-Vahl, K., Bäumer, T., Gerloff, C.,
Schnitzler, A., Orth, M., & Münchau, A. (2010). Imitation in patients with Gilles de la Tourette
syndrome—A behavioral study. *Movement Disorders*, *25*(8), 991–999.
https://doi.org/10.1002/mds.22994

Kurlan, R., Daragjati, C., Como, P. G., McDermott, M. P., Trinidad, K. S., Roddy, S., Brower, C. A., &
Robertson, M. M. (1996). Non-obscene complex socially inappropriate behavior in Tourette's
syndrome. *Journal of Neuropsychiatry and Clinical Neurosciences*, *8*(3), 311–317.

Leckman, J. F., Riddle, M. A., Hardin, M. T., Ort, S. I., Swartz, K. L., Stevenson, J., & Cohen, D. J.
(1989). The Yale Global Tic Severity Scale: Initial testing of a clinician-rated scale of tic severity.
Journal of the American Academy of Child & Adolescent Psychiatry, *28*(4), 566–573.

Lees, A. J., Robertson, M., Trimble, M. R., & Murray, N. M. (1984). A clinical study of Gilles de la
Tourette syndrome in the United Kingdom. *Journal of Neurology, Neurosurgery & Psychiatry*,
47(1), 1–8. https://doi.org/10.1136/jnnp.47.1.1

Li, X., Meng, X., Li, H., Yang, J., & Yuan, J. (2017). The impact of mood on empathy for pain: Evidence
from an EEG study. *Psychophysiology*, *54*(9), 1311–1322.

Neuner, I., Kupriyanova, Y., Stöcker, T., Huang, R., Posnansky, O., Schneider, F., Tittgemeyer, M., &
Shah, N. J. (2010). White-matter abnormalities in Tourette syndrome extend beyond motor
pathways. *NeuroImage*, *51*(3), 1184–1193. https://doi.org/10.1016/j.neuroimage.2010.02.049

Niccolai, V., van Dijk, H., Franzkowiak, S., Finis, J., Südmeyer, M., Jonas, M., Thomalla, G., Siebner,
H. R., Müller-Vahl, K., Münchau, A., Schnitzler, A., & Biermann-Ruben, K. (2016). Increased
beta rhythm as an indicator of inhibitory mechanisms in tourette syndrome. *Movement
Disorders*, *31*(3), 384–392. https://doi.org/10.1002/mds.26454

Paulus, F. M., Müller-Pinzler, L., Walper, D., Marx, S., Hamschmidt, L., Rademacher, L., & Einhäuser,
W. (2017). *Image database to supplement "Paulus, F.M. et al. Pain empathy but not surprise
in response to unexpected action explains arousal related pupil dilation." (VIPER database)*
[Data set]. Zenodo. https://doi.org/10.5281/zenodo.322426

Peled-Avron, L., Goldstein, P., Yellinek, S., Weissman-Fogel, I., & Shamay-Tsoory, S. G. (2018).
Empathy during consoling touch is modulated by mu-rhythm: An EEG study. *Neuropsychologia*,
116, 68–74. https://doi.org/10.1016/j.neuropsychologia.2017.04.026

Perrin, F., Pernier, J., Bertrand, O., & Echallier, J. F. (1989). Spherical splines for scalp potential and

current density mapping. *Electroencephalography and Clinical Neurophysiology*, *72*(2), 184–

187. https://doi.org/10.1016/0013-4694(89)90180-6

Perry, A., Bentin, S., Bartal, I. B.-A., Lamm, C., & Decety, J. (2010). "Feeling" the pain of those who are

different from us: Modulation of EEG in the mu/alpha range. *Cognitive, Affective, & Behavioral*

Neuroscience, *10*(4), 493–504. https://doi.org/10.3758/CABN.10.4.493

Pion-Tonachini, L., Kreutz-Delgado, K., & Makeig, S. (2019). ICLabel: An automated

electroencephalographic independent component classifier, dataset, and website.

NeuroImage, *198*, 181–197. https://doi.org/10.1016/j.neuroimage.2019.05.026

Robertson, M. M., Banerjee, S., Kurlan, R., Cohen, D. J., Leckman, J. F., McMahon, W., Pauls, D. L.,

Sandor, P., & Wetering, B. J. M. van de. (1999). The Tourette Syndrome Diagnostic Confidence

Index: Development and clinical associations. *Neurology*, *53*(9), 2108–2108.

https://doi.org/10.1212/WNL.53.9.2108

Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker,

R., & Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): The

development and validation of a structured diagnostic psychiatric interview for DSM-IV and

ICD-10. *The Journal of Clinical Psychiatry*, *59 Suppl 20*, 22-33;quiz 34-57.

Sprengelmeyer, R., Young, A. W., Pundt, I., Sprengelmeyer, A., Calder, A. J., Berrios, G., Winkel, R.,

Vollmöeller, W., Kuhn, W., Sartory, G., & Przuntek, H. (1997). Disgust implicated in obsessive-

compulsive disorder. *Proceedings of the Royal Society B: Biological Sciences*, *264*(1389),

1767–1773.

Thomalla, G., Jonas, M., Bäumer, T., Siebner, H. R., Biermann-Ruben, K., Ganos, C., Orth, M.,

Hummel, F. C., Gerloff, C., Müller-Vahl, K., Schnitzler, A., & Münchau, A. (2014). Costs of

control: Decreased motor cortex engagement during a Go/NoGo task in Tourette's syndrome.

Brain, *137*(1), 122–136. https://doi.org/10.1093/brain/awt288

van Doorn, J., van den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., Etz, A., Evans, N. J.,

Gronau, Q. F., Haaf, J. M., Hinne, M., Kucharský, Š., Ly, A., Marsman, M., Matzke, D., Gupta,

A. R. K. N., Sarafoglou, A., Stefan, A., Voelkel, J. G., & Wagenmakers, E.-J. (2021). The JASP

guidelines for conducting and reporting a Bayesian analysis. *Psychonomic Bulletin & Review*,

28(3), 813–826. https://doi.org/10.3758/s13423-020-01798-5

Whitmarsh, S., Nieuwenhuis, I. L. C., Barendregt, H. P., & Jensen, O. (2011). Sensorimotor Alpha

Activity is Modulated in Response to the Observation of Pain in Others. *Frontiers in Human*

Neuroscience, *5*. https://doi.org/10.3389/fnhum.2011.00091

Woods, D. W., Piacentini, J., Himle, M. B., & Chang, S. (2005). Premonitory Urge for Tics Scale (PUTS):

Initial psychometric results and examination of the premonitory urge phenomenon in youths

with Tic disorders. *Journal of Developmental & Behavioral Pediatrics*, *26*(6), 397–403.

Yang, C.-Y., Decety, J., Lee, S., Chen, C., & Cheng, Y. (2009). Gender differences in the mu rhythm

during empathy for pain: An electroencephalographic study. *Brain Research*, *1251*, 176–184.

https://doi.org/10.1016/j.brainres.2008.11.062