Purging due to self-fertilization does not prevent accumulation of expansion load

Leo Zeitler*, Christian Parisod, Kimberly J. Gilbert*

December 19, 2022

Department of Biology, University of Fribourg, Chemin du Musée 10, 1700 Fribourg, Switzerland

* Corresponding authors; e-mail: kimberly.gilbert@unifr.ch, leo.zeitler@unifr.ch

Preprint:

Classification: Biological Sciences, Evolution

Keywords: range expansion, *Arabis alpina*, demographic history, mating system, distribution of fitness effects

Abstract

Populations at expanding edges of species distributions face multiple challenges during colonization. Two important difficulties are lack of mates (Allee effects) and elevated mutational burden (expansion load). Self-fertilization impacts both of these factors: providing mate reassurance and potentially purging recessive deleterious variants. Populations at range margins are often observed as enriched for higher self-fertilization rates suggesting that selfing, may be favored under expansion conditions. Using forward-time, individual-based simulations, we disentangle the effects of range expansion versus selfing to understand the potential benefits from mate reassurance and purging of deleterious alleles. We find that selfers do expand faster, but expansion load still accumulates due to demography, regardless of the mating system. The severity of variants contributing to this load, however, differs across mating system: higher selfing rates allow purging of large-effect recessive variants leaving a burden consisting of small- to intermediate-effect alleles. We test these predictions in the mixed-mating, flowering plant *Arabis alpina*, using whole-genome sequences sampled across an area of both a transition from outcrossing to selfing and from glacial refugia to expanded populations. These empirical results indicate an accumulation of expansion load along with evidence of purging in selfing populations. Our simulations, which disentangle the effects of inbreeding due to range expansion versus due to self-fertilization along with the concordance between both simulated and empirical results suggest that an important beneficial factor for the evolution of selfing during range expansions is the effect of purging, but that purging is not sufficient to prevent load accumulation due to demographic effects.

Significance

Range expansions are a pervasive demographic effect across the tree of life. When populations at species range margins expand, they suffer from an accumulation of deleterious alleles due to repeated founder effects. Could self-fertilization, which is considered an evolutionary dead-end, be favored under these conditions, not only to allow for faster expansion due to assured mating but also due to the purging effect that selfing provides against deleterious alleles? Combining simulations and whole-genome data from the mixed-mating alpine rock-cress, which has expanded since the last glaciation, we find that selfing does not prevent the accumulation of deleterious load, however purging does still act to remove the most severe variants, indicating that selfing provides this additional benefit during range expansions.

Introduction

The demographic history of species range expansions is widespread among species across the globe (Davis and Shaw 2001; Parmesan 2006). Difficulties associated with the process of colonization during a species range expansion lead to interesting evolutionary questions of how these populations manage to adapt and continue spreading. Individuals colonizing previously unoccupied environments face limited mate or pollinator availability, resulting in slower expansion as a result of these Allee effects (Moeller *et al.* 2012; Dennis 1989; Courchamp *et al.* 1999; Stephens and Sutherland 1999; Hallatschek and Nelson 2008). Furthermore, expanded populations are known to accumulate deleterious load due to reduced efficacy of selection and high genetic drift in small founding populations, as a result of gene surfing (Klopfstein *et al.* 2006; Burton and Travis 2008; Hallatschek and Nelson 2010). Fixation of deleterious alleles at the range front leads to expansion load which has the potential to temporarily halt population growth or cause local extinction at the boundaries of the species range (Peischl *et al.* 2013, 2015; Gilbert *et al.* 2017, 2018). Evidence of elevated load due to expansion is well-documented empirically, including in humans (Henn *et al.* 2015, 2016; Peischl *et al.* 2018), plants (González-Martínez *et al.* 2017; Willi *et al.* 2018), and experimental bacterial populations (Hallatschek and Nelson 2010; Bosshard *et al.* 2017). Whether some organisms are able to overcome the burden of expansion load through adaptive measures during expansion is, however, unclear.

Numerous plant species are known to have expanded their ranges after the last glaciation, and it is widely observed that many of these species exhibit a transition to self-fertilization ('selfing') at range edges (Barrett 2002). Selfing can reduce fitness through the expression of inbreeding depression (Charlesworth

and Charlesworth 1987; Reusch 2001; Barrett 2013), and is often considered to be an evolutionary dead end, as it can lead to mutational meltdown (Lynch *et al.* 1995; Goodwillie *et al.* 2005; Igic and Busch 2013). Yet the observation of enrichment for selfing at species range edges suggests that there is some advantage gained with this mating system under the demographic and evolutionary conditions that range expansions impose. Simulations have likewise predicted that selfing should evolve at species range edges during expansion (Encinas-Viso *et al.* 2020).

Selfing has three major evolutionary impacts on populations: reducing effective population size, reducing the effective recombination rate, and removing or reducing Allee effects. Selfing can remove Allee effects by assuring mating success in low density populations (Baker 1967; Lloyd and Schoen 1992), and therefore lead to faster expansion speeds, already evidenced by some studies of organisms with uniparental reproduction (Pannell and Barrett 1998; Eriksson and Rafajlović 2021). This is a clear and expected advantage during species range expansion, as colonization can be faster. However, the reduction in N_e along with reduced effective recombination rate due to selfing should be disadvantageous and exacerbated when compounded with the already reduced diversity due to founder effects during range expansion. Some evidence suggests that range expansions may facilitate a transition to selfing by depleting the genetic load at the edge and reducing inbreeding depression (Pujol *et al.* 2009). Because selfing greatly increases homozygosity, it creates the potential for exposing recessive deleterious alleles to selection and thus purging them from the population (Ohta and Cockerham 1974; Charlesworth 1992; Glémin 2007; Pujol *et al.* 2009). Purging therefore has the potential to counteract the accumulation of expansion load. Yet, Glémin (2007) concluded that purging might only be a short term effect of selfing, and that over long evolutionary time scales fixation would be the dominant effect in selfers. Evidence suggests that in small populations purging by self-fertilization is less feasible, and that small-effect deleterious variants can still contribute to an increase in genetic load (Wang *et al.* 1999). The distribution of selection coefficients compared to population size is thus a major factor for the evolution of mating systems (Bataillon and Kirkpatrick 2000; Glémin 2003). As a consequence, one important change in the population genetic signature of a transition from outcrossing to selfing is a shift in the observed distribution of fitness effects (DFE) for mutations segregating in the population to reflect less efficient selection against weakly deleterious additive variants and purging of strongly deleterious recessive variants (Barrett *et al.* 2014; Laenen *et al.* 2018; Arunkumar *et al.* 2015). Whether this prediction holds when a species range expansion occurs concurrently with a mating system shift has, to our knowledge, not been explored.

In combination with simulations of a range expansion and mating system shift, we additionally investigate an empirical system matched to this demographic and evolutionary history. The perennial arctic-alpine plant *Arabis alpina* L., is a species known to have been subject to range expansions and contractions in response to the repeated quaternary climate oscillations (Ansell *et al.* 2008). The Italian peninsula is a known refugium for populations during the last glacial maximum (Ansell *et al.* 2008), after which the species recolonized alpine habitats across Europe and concurrently with this expansion evolved a mating system of predominant self-fertilization (Tedder *et al.* 2015). In Italy, populations of *A. alpina* are predominantly outcrossing with high genetic diversity (Ansell *et al.* 2008; Laenen *et al.* 2018), while in the French and Swiss Alps, populations are more homozygous and selfing (Ansell *et al.* 2008; Buehler *et al.* 2012). Given its demographic history of range expansion and the variation in outcrossing rates, we study the combined effect of a range expansion and an evolutionary transition in mating system from natural populations across Italy and the western Alps.

Here, we test the hypothesis that purging due to selfing may provide an evolutionary benefit during species range expansions, in the face of accumulating deleterious expansion load. We investigate how selfing affects genetic diversity, mutation load, the observed DFE, and the speed of colonization during a range expansion. Using individual-based simulations, we model a range expansion where different degrees of self-fertilization are introduced mid-expansion and compare to a obligately outcrossing control. We focus in particular on the dynamics of load accumulation with varying selfing rates and include lethal, mildly deleterious and beneficial mutations with different dominance coefficients. By characterizing the distribution of selection coefficients of expanded populations, we highlight differences in the efficacy of selection across mutation severity. We also compare the buildup of genetic load across selfing rate scenarios and estimate load under different assumptions of dominance. We then test if our predictions from simulations are equivalently detectable in empirical data from natural populations of *A. alpina* in the Italian-Alpine expansion zone. We find that rapid purging due to self-fertilization is common for

highly deleterious recessive variants, but does not prevent the accumulation of genetic load in expanded populations. Our results mark a vital step towards deciphering the interplay of complex population demography and mating system on the fate of genetic diversity and mutation load, helping us to better understand how populations react to changing environmental conditions.

Results

Selfing leads to faster expansion

Using individual-based forward-time simulations in SLiM v3.7.1 (Haller and Messer 2019), we modelled a range expansion across a one-dimensional linear landscape. We simulated obligate outcrossing followed by a shift to self-fertilization in the 25th deme of the landscape (out of 50 total demes), with selfing rates σ of $0.5, 0.95$, or 1, or continued obligate outcrossing until the landscape is fully colonized (Figure 1A).

One expected benefit of selfing during range expansion is increased expansion speed, which we observed in our simulation results. We compared the number of generations required to cross the landscape among selfing rates and found that mixed mating and obligate selfing populations had a faster expansion speed compared to obligate outcrossing populations (see Figure 1B). The differences in generation time among different selfing rates was minor (mean generation times for $\sigma = 0.5, 0.95$, and 1, respectively: $792 \, (SD = 78.5)$, $800 \, (SD = 51.4)$, $777 \, (SD = 70.7)$) compared to the notable difference in expansion time of obligate outcrossers (990 generations $(SD = 69.8)$).

Range expansion increases genetic load and decreases diversity in simulations

To test if and how selfing modifies the outcomes of a range expansion in our simulations, we examined genetic diversity in expanded populations and across selfing rates. Both outcrossing and selfing edge populations showed large reductions in diversity due to the expansion. Outcrossers retained the highest nucleotide diversity for neutral sites ($\pi_{edge,\sigma=0} = 3.646 \times 10^{-5}$), while with increasing self-fertilization rates, populations showed further reductions in nucleotide diversity ($\pi_{edge,\sigma=1} = 3.410 \times 10^{-6}$, Figure 1C). Core populations which never experienced expansion and always outcrossed had the highest nucleotide diversity ($\pi_{core} = 9.235 \times 10^{-5}$).

Genetic load is predicted to be higher in expanded populations, so we next examined how selfing modulates this outcome of a range expansion. We compared mean fitness ($\overline{\omega}$) in core versus edge populations after the simulated expansion was complete. In all cases, the range expansion reduced fitness at the edge due to expansion load (Figure 1D). In the obligate outcrossing case ($\sigma = 0$), we observed a reduction of fitness from core to edge of 29.1%. Interestingly, selfers showed negligible differences in load accumulation relative to outcrossers, with at most a mean reduction in fitness of 31.7% for obligate selfers. We observed an increase in the proportion of loci fixed for deleterious alleles in all expanded populations, with these proportions increasing for higher selfing rates (Figure S1A). Similarly, we found that mean counts of deleterious loci increased from core to edge as well as from lower to higher rates of selfing (Figure S1B), whereas counts of deleterious alleles showed less to no clear pattern from core to edge and among selfing rates (Figure S1C).

To understand why and how self-fertilization seemingly had no impact on purging genetic load during a range expansion, we examined demes over time and space to disentangle the effects of inbreeding due to demography versus inbreeding due to increased self-fertilization (Figure 2). Outcrossing deme 24 exhibited a mean observed heterozygosity level of $\overline{H}_{24} = 1.47 \times 10^{-4}$ when it was first colonized during the expansion, i.e., when it was the edge of the species range. Beyond deme 24 the mating system shifts to selfing and we observed a continual loss of heterozygosity at the expanding front. When the expansion front reached deme 35, $\overline{H}_{35}$ for 95% selfers rapidly decreased to 7.05×10^{-7}. For outcrossers, however, heterozygosity exhibited a more gradual rate of reduction across the course of expansion, reaching $\overline{H}_{35} = 1.19 \times 10^{-4}$ by the time deme 35 was colonized (Figure 2A, S2). We examined how diversity recovered in deme 35 over time since its colonization until the end of the simulation and found that outcrossers recovered to higher levels ($\overline{H}_{35}$ at the end of the simulation 2.27×10^{-4}) than selfers (for 95% selfing $\overline{H}_{35} = 7.72 \times 10^{-6}$).

Figure 1: Simulation schematic for 1-D landscapes with stepping-stone migration (A). A shift in the rate of self-fertilization occurs in the center of the landscape (blue triangle). The number of generations needed to cross the landscape from the core to deme 50 (B). Expansion time was lower for all selfing rates but higher for obligate outcrossing. Mean nucleotide diversity (C) was reduced outside of the core, with a greater reduction for higher selfing: $\overline{\pi}_{core} = 9.235 \times 10^{-5}, \overline{\pi}_{\sigma=0} = 3.646 \times 10^{-5}, \overline{\pi}_{\sigma=0.5} = 1.274 \times 10^{-5}, \overline{\pi}_{\sigma=0.95} = 4.608 \times 10^{-6}, \overline{\pi}_{\sigma=1} = 3.410 \times 10^{-6}$. Relative fitness (D) decreased from core to edge with similar values across outcrossers and selfers: $\overline{\omega}_{\sigma=0} = 0.709, \overline{\omega}_{\sigma=0.5} = 0.702, \overline{\omega}_{\sigma=0.95} = 0.676, \overline{\omega}_{\sigma=1} = 0.683$, reflecting a relative loss of fitness as compared to the core of $29.1\%, 29.8\%, 32.4\%$, and 31.7% respectively for $\sigma = 0, 0.5, 0.95$, and 1.

Figure 2: Observed heterozygosity (A), count of lethal alleles (B), and the rate of fitness change (C-D) are shown through time for the expanding range edge (green) as compared to change over time within one interior deme, stationary on the landscape (blue, deme 35). For selfing rates 50% and 100%, see Figures S2 and S3. Panels (C) and (D) show the rate of fitness change as measured over 100-generation intervals, separately for outcrossers and selfers. A value of 1 indicates no change in fitness over 100 generations while values above 1 indicate increasing fitness and values below 1 indicate fitness loss. The vertical dashed line indicates the point in time where the mating system shifts to selfing. This shift occurs at deme 25 on the landscape, and since there is variation across simulation replicates in the generation time taken to expand to deme 25, we plotted all values relative to this time point for each replicate shown ($n = 20$ replicates per selfing rate scenario). Each point is the value from a single simulation replicate and lines are loess (span=0.2) fitted curves across all replicates.

Fitness loss despite genetic purging

We counted the number of lethal alleles per individual and observed a reduction in the count of lethals that corresponds with the reduction in heterozygosity (Figure 2A-B). Lethal alleles were only reduced when the shift to selfing occurred, and we observed the same pattern at every simulated selfing rate. Obligate outcrossers did not exhibit a reduction in lethal alleles, showing no evidence of purging. The largest reduction in lethal alleles occurred for the shift to the highest selfing rate ($\sigma = 1$) with a 91.69% drop in lethal alleles, while our lowest simulated selfing rate ($\sigma = 0.5$) still showed a strong effect of purging lethal alleles with a 83.74% reduction (Figure S3).

The rate of change of fitness in a given deme, measured at a focal generation (t) compared to 100 generations prior ($t - 100$), showed a consistent loss of fitness over time due to range expansion as well as some fitness recovery in populations behind the expanding front (Figure 2C, D). Edge demes which recently underwent the shift to selfing exhibited a drastic reduction in fitness relative to equivalent outcrossers. This high rate of fitness loss exhibited by selfers is temporary and only lasts for between 45-235 generations, after which the rate of fitness loss recovers to the same rate as that observed in expanding obligate outcrossers: still below one on average and accumulating expansion load.

We also investigated the impact the range expansion and mating shift had on the realized distribution of selection coefficients. Overall, we found the greatest proportion of deleterious mutations in the weak to intermediate bin of selection coefficients ($-0.0001 \leq s < -0.001$), with just below 60% of all sites falling into this class (Figure 3). The next most deleterious bin ($-0.001 \leq s < -0.01$) contained about 30% of sites, while about 10% of sites are in the weakest selection coefficient bin. Lethal alleles made up a small proportion of segregating sites, as expected given the small proportion defined in the simulation parameters. Within these small numbers of severely deleterious variants, there was a consistent trend for a reduction of lethals from core to edge of nearly 50% for outcrossers and significantly further reduction for all rates of selfing, increasing from a nearly 75% reduction from core to edge for 50% selfers to more than 75% for 100% selfers (Figure 3 inset). The pattern of reduced proportions of deleterious sites as selfing rate increases holds in both the lethal category as well as the second-most deleterious allele class. We consistently observed the reverse pattern in the remaining weaker effect bins, with proportions of weakly deleterious sites slightly increasing at range edges and more so with higher selfing rates. This observation is consistent with more efficient removal of highly deleterious alleles and mutation accumulation at sites with smaller absolute selection coefficients.

Reduced genetic diversity and elevated load in expanded selfing *A. alpina* populations

To test if our observations for genetic diversity and load accumulation from simulations are matched in natural populations, we used the mixed mating plant *Arabis alpina*, which underwent a range expansion concurrently with a shift to higher self-fertilization rates. Using 191 newly sampled and sequenced short-read genomes from Italy and France combined with publicly available data from Switzerland and across Europe (Laenen *et al.* 2018; Rogivue *et al.* 2019), we examined differences across the species range in 527 individuals at high resolution.

Population structure results showed expected clustering by regions (Figure S4), matching the geography of sampled populations from Abruzzo in southern Italy, the Apuan Alps in northern Italy, the French Alps, and the Swiss Alps (Figure 4A). Previously sampled individuals from Italy, France, and Switzerland that we combined with our newly sampled individuals also consistently clustered within the same geographic regions. Samples of fewer individuals from more widely across Europe showed reasonable structuring among Greek, Spanish, and Scandinavian populations. Inference of the demographic history of these populations showed a history of population bottlenecks and recovery consistent with the northward expansion following the last glacial maximum and the shift to selfing (see Figure S5).

To reveal how the expansion and self-fertilization impacts key diversity parameters, we calculated individual inbreeding coefficients (F) and nucleotide diversity (π). We found the highest inbreeding coefficients in mixed mating and highly selfing populations outside of Italy, and reduced inbreeding in the Apuan Alps and Abruzzo (means for Swiss, French, Apuan Alps and Abruzzo, respectively: $\bar{F} = 0.51, 0.75, 0.43, 0.37$). The highest overall inbreeding coefficients were estimated in populations from Spain ($\bar{F} = 0.96$) and France (Br, $\bar{F} = 0.89$, Figure 4B). Swiss populations had the greatest standard deviation (Pa, $SD(F) = 0.26$), and Italian populations had the lowest mean value (Am, $\bar{F} = 0.34$). For nucleotide

Figure 3: The observed distribution of selection coefficients from simulations at the end of expansion. The core deme (blue) is compared to edge demes (green) for obligate outcrossing (darkest color) versus higher selfing rates (lighter colors). Error bars indicate 0.05 and 0.95-quantiles across the 20 simulation replicates. The inset panel emphasizes the degree to which the proportion of sites in the lethal category changes over mating system scenario from core to edge.

diversity, we found high values in the Abruzzo region of Italy ($\overline{\pi}_s = 0.00767$). Genetic diversity reduced when moving north to the French Alps ($\overline{\pi}_s = 0.00333$) and Switzerland ($\overline{\pi}_s = 0.00357$, Figure 4C).

We calculated R'_{xy} to assess the accumulation of derived deleterious alleles, using 270,889 SNPs annotated as deleterious and 2380 as loss of function (LoF) variants, classified by SNPeff (Cingolani *et al.* 2012). R'_{xy} is a pairwise statistic that compares the count of derived alleles found in one population relative to another, and avoids reference bias introduced by branch shortening (Do *et al.* 2015). $R'_{xy} > 1$ indicates that population X has more derived alleles of a given class than population Y relative to the neutral expectation, while $R'_{xy} < 1$ would indicate fewer derived alleles in population X. For deleterious sites we found that alpine populations had more derived alleles compared to Italian populations (Figure 4D), indicating an increase in genetic load from south to north. Within the Alps, all Swiss populations had reduced derived allele frequencies compared to France, while relative to the Apuan Alps in northern Italy, only few Swiss populations exhibited reduced derived allele counts, potentially suggesting that the shift to selfing has begun to alleviate the accumulation of expansion load. For LoF loci, signals of both purging and accumulation were detectable. Some pairwise population comparisons showed an increase in number of LoF alleles from south to north (e.g., nearly all Apuan × Abruzzo comparisons), while others showed mixed results, depending on the focal populations (e.g., French × Apuan, Swiss × French). All population comparisons of Swiss Alps × Apuan Alps showed reduced LoF allele counts.

We next used the SNPs annotated as putatively deleterious to examine the accumulation of genetic load in our expanded populations. We assessed the predictability of two different load proxies using known fitness and load values from our simulation results: an additive model which counted individual deleterious alleles, or a recessive model which counted individual homozygous loci. The correlation between mean fitness and load proxy was better for the recessive model ($R^2 = 0.82, P < 0.001$) than the additive model ($R^2 = 0.10, P < 0.001$, Figure S6), supporting the use of this load proxy even in the presence of selfing. The additive model also predicted load more poorly in a supplemental set of simulations using only fully additive mutations (see Figure S7). Estimating load from these models in the empirical dataset indicated higher load in expanded, selfing populations from France and Switzerland compared to core Italian populations, using the recessive model (Figure 5A). The additive model found slightly reduced load for alpine populations compared to Italian ones (Figure 5B).

To further understand the mutational burden within our *A. alpina* populations, we estimated the distribution of fitness effects of new mutations (DFE) using `fitdadi` (Kim *et al.* 2017), which corrects for demographic history by first fitting a best demographic model to the data (Figure S5). `fitdadi` surprisingly estimated a similar DFE across all of our sampled populations (Figure S8), with a large proportion of strongly deleterious sites at or above 60%, around 20% of sites in the weakest selection class, and approximately 5% in each of the two intermediate selection classes. The proportions varied only marginally across core Italian populations as well as across expanded French and Swiss populations. We additionally examined the fixation of deleterious alleles across our populations, within classes of neutral, deleterious, or LoF sites (Figure S9). Fixation of all sites increased from Italy in the south to France in the north, but then decreased from France to Switzerland, reminiscent of our R'_{xy} results suggesting more purging in Swiss populations.

Figure 4: Sampling sites of *A. alpina* in the Italian-Alpine expansion zone (A). Inbreeding coefficients for individuals across sampled populations, including Spain, Scandinavia, and Greece (B). The distribution of nucleotide diversity estimated for Italian and alpine populations (C), with diamonds indicating group means. R'_{xy} values for deleterious (purple) and LoF (orange) loci (D). $R'_{xy} > 1$ indicates an accumulation of derived alleles at deleterious or LoF sites relative to neutral ones, while $R'_{xy} < 1$ indicates the opposite.

Discussion

In this study, we investigated the impact of selfing on the accumulation of genetic load during a species range expansion. We used simulations to disentangle the reduction in effective population size at expanding fronts due to demography versus the reduction due to self-fertilization. We then compared our expectations for the impact of selfing in empirical data from natural populations having undergone both a range expansion and a mating system shift. Because selfing reduces the effective recombination rate within populations as well as genetic diversity, it is expected to be generally maladaptive for evolution and adaptation. However, conditions at the expanding edge of a species range may particularly favor the evolution of this mating system. And the compounded effects of reduced diversity due to selfing at range edges may even provide an additional benefit of purging homozygous recessive deleterious mutations.

Figure 5: Genetic load in *A. alpina* populations as inferred from counts of deleterious loci using a recessive model (A) versus counts of deleterious alleles using an additive model (B). Loci are classified as putatively deleterious by SNPeff (see Methods).

One clear advantage that we corroborate is that selfing provides mate reassurance (Igic and Busch 2013) and leads to faster spread over geographic space. Despite similar losses in fitness from core to range edge for both outcrossers and all selfing rate scenarios, selfers still colonized the landscape faster than outcrossers. This result adds to the general prediction of Baker's Law, that selfing may be advantageous in mate-limited environments (Baker 1967). Since we do not have equivalently expanded outcrossing populations in our *A. alpina* dataset nor a generation time measurable in contemporary time, we cannot empirically investigate this question. To fully understand the benefits of mate reassurance from selfing, it may be fruitful for future empirical studies to focus on organisms with well-documented expansion times and mating system shifts, or potentially take advantage of laboratory experiments with expansions of outcrossers versus selfers under controlled conditions, for example with mixed mating species of *Caenorhabditis*.

Our main interest in comparing simulation and empirical results is to understand the dynamics of load accumulation during range expansion when selfing evolves. Another potential major benefit of selfing is the opportunity for purging due to increased homozygosity. Our simulation results show that expansion load always accumulates at range edges, regardless of selfing rate and with equivalent severity to outcrossers. This seems to suggest that purging due to selfing offers no additional benefit during species range expansion. However, when looking at the distribution of effect sizes for variants segregating within populations, we do also detect significant effects of purging unique to selfers whereby lethal-effect alleles are successfully and rapidly removed from the population. Purging was most pronounced in obligate selfers, where within only 30-150 generations lethal alleles are removed from the population and remain at low levels for the remainder of the simulations (Figure 2B). This effect is also notable when examining the distribution of mutational effect sizes at the end of the simulations, where selfers exhibited major reductions in lethal alleles (Figure 3). This matches previous theoretical predictions that with increased homozygosity lethal mutations should be efficiently be removed and the number of segregating lethals lowered (Kirkpatrick and Jarne 2000; Hedrick 2002).

Purging does not, however, allow these populations to escape the burden inflicted by expansion load and their demographic past. Load still accumulates in all population expansions regardless of mating system, but how this load is expressed is what differs among mating systems. The overall burden experienced by selfing populations consists of more small-effect deleterious variants, which accumulate to a greater extent as compared to obligate outcrossing populations. Previous simulation studies have also highlighted how small effect variants are much more difficult to purge (Wang *et al.* 1999; Willis 1999), and therefore can still result in an accumulation of expansion load.

In our empirical *A. alpina* results we found similar signatures of both load accumulation and genetic purging in expanded populations. The recessive load model indicates that French and Swiss expanded populations have accumulated genetic load, through higher counts of putatively deleterious sites. Moreover, the additive model shows a small decrease in deleterious counts for our expanded populations. This suggests that negative selection has purged some diversity from these populations, since otherwise allele counts should remain at constant levels across all populations if only genetic drift is acting and not selection (Peischl and Excoffier 2015). However, our empirical DFE inferences only detected minor trends of reduced proportions of sites in the most deleterious class for expanded Swiss and French populations and equivalently support a bimodal-shaped DFE reported in Laenen *et al.* (2018). Whether this reflects true minor differences in the DFEs among these populations, or a lack of proper inferential ability is difficult to know. Comparing genetic load in populations with complex population demographic histories is still challenging (Simons and Sella 2016; Brandvain and Wright 2016), largely because dominance coefficients are unknown and different DFEs may respond differently to estimation accuracy (Gilbert *et al.* 2022) or simply differently to demographic change (Glémin 2003; Balick *et al.* 2015). Other approaches which attempt to directly infer mutational effects require intensive and in-depth analyses, emphasizing the need for improved inference methods or further in-depth investigation to accurately and efficiently identify effect sizes of variants in natural populations and therefore gain a complete picture of the genetic architecture of mutation load. Still, our R'_{xy} analyses provide additional support for our load estimates across populations, showing an accumulation of derived deleterious alleles relative to neutral alleles in all expanded populations relative to Italian populations. Only further along our expansion axis, in Swiss populations, is there some evidence that further accumulation of load is purged due to selfing.

The concurrence between our simulated and empirical results give striking insights into the inter-

actions of demographic change due to range expansion with recombination and diversity changes due to self-fertilization. We observe that during the process of range expansion, newly colonized demes are highly subject to drift and founder events, as extensively discussed in previous studies on range expansions (Excoffier *et al.* 2009; Slatkin and Excoffier 2012; Peischl *et al.* 2013, 2015). On top of this, the added impact of self-fertilization exacerbates losses of heterozygosity at expanding range edges. Our simulations indicate that this loss of diversity can be recovered after the expansion front has passed, when migration and population growth allow for increased efficiency of selection in larger and more diverse populations, as previously described in Gilbert *et al.* (2018). We find new insights here, however, that this recovery is much slower for selfing populations, supporting the widely held idea that selfing should only be favored at range edges and that outcrossing may replace selfing after a range expansion has occurred. Given the result that our empirical populations still exhibit signatures of genetic load is then equally interesting, since these populations are expected to have had many generations for recovery since they were directly on an expanding edge. The dynamics of when selfing evolves and can be favored across a species range is examined *in silico* by Encinas-Viso *et al.* (2020), showing that unless recombination rates are high enough, outcrossing individuals will outcompete selfing populations once the expansion edge has passed. Whether our sampled alpine populations populations have also began shifting back towards increased outcrossing is currently unknown and an avenue of investigation which will be interesting to pursue in the future.

Studying range expansions in plant species offers unique insights into the combination of mating system evolution combined with the evolutionary processes occurring during species range expansions. Previous work in *A. alpina* has also evidenced increased load with high selfing and bottlenecks in Scandinavian populations (Laenen *et al.* 2018), however we highlight previously unidentified evidence for purging of strongly deleterious alleles in intermediate to highly selfing continental populations within the French and Swiss Alps, in addition to expansion load still incurred. Our results are also similar to those found in other plant range expansions where selfing is observed at the range edge. Notably, this is the case in *A. alpina*'s close relative *Arabidopsis lyrata* (Willi *et al.* 2018) as well as in *Mercurialis annua* (González-Martínez *et al.* 2017) where expansion load has been indicated. Our study has uniquely also identified signature of purging due to selfing, which is a known expectation from theoretical predictions (Glémin 2003, 2007), but to our knowledge not thoroughly investigated in empirical systems. Future studies could likewise benefit from direct estimates of fitness across the species range, through crosses and common garden studies.

Population genetic simulations help us to better understand interactions of effects that are difficult to assess or disentangle in empirical populations. Here, we have only explored a finite parameter space and constrained our simulations to simplified demographic models. Since we were only interested in the eventual signatures resulting from selfing evolution during a range expansion, we modeled the loss of self-incompatibility as a sudden shift in the probability of selfing at one location on the landscape. However, in nature, the shift to self-fertilization is expected to occur gradually over time, e.g, due to a reduction in S-allele diversity (Charlesworth and Charlesworth 1979; Vallejo-Marín and Uyenoyama 2004; Porcher and Lande 2005; Encinas-Viso *et al.* 2020). Even with our sudden evolution of selfing imposed in the middle of the landscape, we expect that the same observed qualitative results of purging strong-effect recessive deleterious alleles and loss of heterozygosity would still occur, just more gradually through time. The intermediate rate of selfing we tested could also be considered an earlier transitional state of a species range expansion on its way to evolving higher selfing rates. In a gradual shift to selfing, initial S-allele diversity would be reduced but outcrossing still frequent, and intermediate selfing rates would be a transient state as populations shift to higher selfing and faster expansion. While we focused on the speed and purging benefits of selfing during a range expansion, we did not address a potential third factor impacting expansions: the necessity to locally adapt to unfamiliar environments. Populations must often adapt to novel or fluctuating environments during expansion, e.g., during glacial cycles (Hewitt 2004) or as soil conditions change over altitude or photoperiod conditions change over latitude. Adaptation requires sufficient genetic variation to match the local environment sufficiently for population growth to be sustainable (Kirkpatrick and Barton 1997; Polechová and Barton 2015). For populations that expand to follow an environment they are already adapted to, this difficulty is less relevant. For example, species expanding post-glaciation are believed to have followed the receding ice sheets as suitable habitat that they were pre-adapted to was slowly revealed. However, it is still likely that some aspect of environmental

conditions are always novel as organisms move over space, necessitating some level of adaptation. Our results importantly highlight how the DFE is expected to differ among outcrossed versus selfed expanding populations, creating contrasting genetic architectures within the genome. Such differences in genetic architectures, i.e., few large-effect or many small-effect loci, for adaptive and maladaptive sites along with differing effective recombination rates across selfing rates are likely to interact with adaptation over changing landscapes and result in different adaptive potentials among populations. And in the future, as anthropogenically-induced climate change causes more rapid changes across the landscape, the likelihood of being able to track moving environmental optima is expected to become more difficult, necessitating more rapid adaptation and emphasizing the importance of studying range expansions and shifts and the evolutionary processes involved.

Conclusions

Range expansions are known to increase genetic drift and fixation of deleterious alleles, reducing fitness as a consequence. Self-fertilization further reduces N_e, which allows for a higher rate of fixation of weaker deleterious mutations compared to outcrossers. However, as predicted by Glémin (2007) this process can also allow for short-term purging. We investigated whether this purging is realized during species range expansions and if selfing can thus be beneficial in this evolutionary context. We described two significant factors in our simulations: first, the purging of lethal alleles is indeed observed in selfing populations, and second, this purging is not sufficient to prevent the fitness loss incurred by expansion load. Weak effect mutations accumulate to a larger extent due to the range expansion, leaving a visible signature in the DFE. Furthermore, in natural populations of *A. alpina*, we see consistent effects of purging as well as load accumulation despite the evolution of selfing. Together, this demonstrates that self-fertilization can alter the signature of genetic load in expanded populations, and identifies purging as an additional benefit of selfing along with mate reassurance. Future studies in empirical systems will hopefully be able to distinguish expanded outcrossing versus expanded selfing populations to further validate our results, as much remains to be learned of the interaction between mating system evolution and demographic history of populations. Improved understanding of these important processes will be vital for further insight into how natural populations will (or will not) be able to disperse and adapt in the face of global climate change and anthropogenic forces experienced in natural habitats.

Material and Methods

We conduct simulations of a species range expansion and compare to an empirical dataset from the plant *Arabis alpina* to understand the dynamics of purging and mutation load accumulation in a system where self-fertilization has evolved. To understand whether selfing acts as an evolutionary advantage during expansion by purging deleterious alleles that otherwise accumulate, we focus on tracking genetic load in both simulated and empirical data. Though simplified from reality, our simulations have the important advantage of knowing true fitness and mutational effects within every individual to best understand the dynamics of load accumulation and purging during range expansion.

Simulations

To simulate a range expansion with a shift in mating system we conducted individual-based, forward time simulations, using a non-Wright-Fisher model in SLiM v3.7.1 (Haller and Messer 2019). We modeled the range expansion across a one-dimensional, linear landscape of 50 demes with a stepping-stone migration model (Figure 1A). Each simulation started with a single initial core deme populated with individuals that then underwent repeated bottlenecks and founder events as they colonized the remaining empty 49 demes. The core population was initiated at carrying capacity $K = 5000$, and prior to expanding we ran a burn-in for $4N$ generations. Generations were discrete and non-overlapping, and after the burn-in was complete we opened the landscape for expansion, introducing migration that allowed individuals to move into either adjacent deme. We defined a forward migration rate of $m = 0.05$ per generation and reflecting boundaries at the ends of the landscape in the core and deme 50. All subsequent demes outside the core

had a carrying capacity of $K = 200$. Once the last deme reached carrying capacity and 100 additional generations passed, we stopped the simulation.

To test the effect of increased self-fertilization during the expansion we conducted a set of control simulations where all individuals in all demes are obligately outcrossing and compared these to scenarios where selfing begins at deme 25. In demes $25 - 50$ of these selfing simulations, we set the self-fertilization rate σ to either 0.5, 0.95, or 1, creating three different scenarios of selfing rates to compare to the full outcrossers. In a given deme, individuals to be selfed were chosen with probability σ each generation. We modeled logistic population growth with a Beverton-Holt model, where the expected number of total offspring per deme for the next generation is given by $N_{t+1} = \frac{RN_t}{1+N_t/M}$, where $M = \frac{K}{R-1}$, growth rate $R = 1.2$, N_t is the deme's census size in the current generation t, and K is the carrying capacity of the focal deme. For each parent, we expected less fit individuals to produce fewer offspring and thus implemented a fecundity selection model, where the expected number of offspring for individual i is approximately Poisson distributed (Peischl *et al.* 2015).

Each individual was modeled as a diploid genome consisting of 1×10^7 base pairs (bp) with a recombination rate of 1×10^{-8} per bp per generation. We simulated neutral, beneficial, deleterious, and lethal mutations at a per base pair mutation rate of 7×10^{-8} per generation occurring at relative proportions of 0.25, 0.001, 0.649 and 0.1, respectively. For deleterious and beneficial mutations, selection coefficients were drawn from an exponential function with mean -0.001 or 0.01, respectively, and lethal alleles had a selection coefficient of -1. Dominance coefficients were set to $h = 0.3$ for beneficial and deleterious alleles, and 0.02 for lethal mutations. In a supplementary set of simulations we tested for the effect of full additivity using $h = 0.5$ for non-lethal mutations. These parameters for our simulated distribution of selection and dominance coefficients are drawn from our best knowledge as described widely in the literature, with partial dominance and more recessive lethal alleles (Keightley 1994; Eyre-Walker and Keightley 2007; Halligan and Keightley 2009; Agrawal and Whitlock 2011). We replicated every parameter combination 20 times.

We recorded fitness and calculated summary statistics during the expansion to track the impact of demographic change in combination with selfing rates. In every deme we measured nucleotide diversity for neutral variants, π, mean observed heterozygosity along the genome, H, counts of lethal and deleterious alleles and recorded mean fitness, $\overline{\omega}$, every five generations. This allowed us to compare changes in fitness and allele counts over time, contrasting them with the same statistic 100 generation in the past. We also examined changes in these summary statistics in specific locations across the landscape during and after the expansion had completed: the core (deme 1), the deme prior to the mating shift (deme 24), the deme ten demes past the facultative mating shift (deme 35, to avoid effects of migration from outcrossers), and the end of the landscape (deme 50). We characterized the composition of load in core and edge populations after the expansion by examining the realized distribution of selection coefficients. To do this, we categorized selection coefficients in four discrete bins ($s \in \{(0.0001, 0), (0.001, 0.0001], (1, 0.001], -1\}$). To further characterize load, we calculated the proportion of fixed deleterious alleles, and applied models often used to compare approximated genetic load in empirical populations (Simons and Sella 2016) to our simulated data: we estimated additive load by counting the total number of deleterious alleles per individual, assuming $h = 0.5$, and recessive load by counting the total number of homozygous deleterious loci per individual, assuming $h = 0$. We then compared these values with realized fitness, all of which are known for the simulations.

Arabis alpina dataset

We compared our theoretical results to an empirical dataset of *Arabis alpina* by combining publicly available data (Laenen *et al.* 2018; Rogivue *et al.* 2019) with newly sampled and sequenced genomes. Our dataset focused on sampling four regions with four populations each, consisting of 15-18 individuals. One exception is northern Italy where two nearby populations (Ca & Gf) of 8 individuals each contributed to five total populations from the region. Sampling spans the range expansion from southern Italy north into the French and Swiss Alps and capturing the transition in mating system from outcrossing to selfing. We collected leaf tissue on silica gel from 198 wild *A. alpina* plants in the Apennine Mountains in central Italy, the Apuan Alps in northern Italy, and the western Alps in France during the summer of June 2021. We extracted DNA with the Qiagen DNeasy Plant Mini Kit (Qiagen, Inc., Valencia, CA, USA)

and constructed libraries using Illumina TruSeq DNA PCR-Free (Illumina, San Diego, CA, USA) or Illumina DNA Prep, and sequenced on a Illumina NovaSeq 6000 (paired-end). All sampled individuals are described in more detail in Supplementary Table 1 and are available publicly at NCBI SRA accession PRJNA773763. We combined this dataset with previously published *A. alpina* short-read genomes of 306 individuals sampled from Switzerland (Rogivue *et al.* 2019) and 36 sampled widely across Europe (Laenen *et al.* 2018). For quality control of the reads, we used FastQC (`http://www.bioinformatics.babraham.ac.uk/projects/fastqc`) and MultiQC (Ewels *et al.* 2016). We trimmed reads using trimmomatic 0.39 (Bolger *et al.* 2014) and aligned them to the *A. alpina* reference genome (Jiao *et al.* (2017), version 5.1, `http://www.arabis-alpina.org/refseq.html`) using `bwa mem` 0.7.17 (Li and Durbin 2009). We calculated coverage for the whole dataset with mosdepth (Pedersen and Quinlan 2018), averaging at 13.98 (18.61 for new samples, Supplementary Table 1). We called variant and invariant sites using `freebayes` 1.3.2 (Garrison and Marth 2012). Additional filters were applied in bcftools (Danecek *et al.* 2021), retaining only sites with a maximum missing fraction of 0.2, and removing any variant sites with estimated probability of not being polymorphic less than phred 20 (QUAL>=20). Finally, we removed 13 individuals with greater than 30% missing calls or low coverage (Br22, Br06, Cc05, St15, Am01, Br18, Br24, Ma28, Pa9, Pi9, Pi95, Pi40, Ma97). The final dataset combined had 3,179,432 SNPs, with 43,268,666 invariant sites for 527 individuals from 31 populations, which includes 191 individuals of the 17 newly sampled populations in the Italy-Alps expansion zone.

Population genetic analyses

We inferred the ancestral state of alleles using the close relative of *A. alpina*, *Arabis montbretiana*, by aligning the reference sequences of *A. alpina* with *A. montbretiana* (Madrid *et al.* 2021) using `last` (Kiełbasa *et al.* 2011). To confirm that our samples from across Europe matched the expected population structuring based on known demographic history, we ran `admixture` v1.3.0 (Alexander *et al.* 2009) from $K = 2$ to $K = 15$ on the full sample set but with SNPs pruned for LD using `bcftools +prune` (Danecek *et al.* 2021, R^2 cutoff 0.3 in a window of 1000 sites). We calculated nucleotide diversity per population in 1Mbp windows using `pixy` (Korunes and Samuk 2021, version 1.2.6.beta1, 10.5281/zenodo.6032358) and inbreeding coefficients for each individual with `ngsF` (6 iterations, Vieira *et al.* 2013). To format the input file for `ngsF`, we randomly sampled 100,000 biallelic SNPs and extracted genotype likelihoods using bcftools (Danecek *et al.* 2021).

To calculate putative genetic load, we first predicted derived, deleterious alleles with SNPeff (Cingolani *et al.* 2012). SNPeff estimates how deleterious a variant may be based on whether its mutation causes an amino acid change, at varying levels of importance (Cingolani *et al.* 2012). We used the categories "nonsense" and "missense" as the definition for deleterious sites from SNPeff, "none" and "silent" annotations were used as neutral predictions and "LOF" annotations as loss-of-function mutations ("LoF") after running the program with the `-formatEff` option. Using these annotations, we calculated R'_{xy} as described in Do *et al.* (2015) with R_{xy} for derived allele counts of LoF or deleterious sites over $R_{xy}^{normalization}$ for neutral sites to avoid reference bias. Further, we estimated jackknife confidence intervals using pseudo values from 100 contiguous blocks and assuming normal distributed values. We also estimated genetic load from SNPeff predictions, using the same recessive and additive models as in the simulations, by counting either the total number of deleterious loci or derived alleles at deleterious sites. Finally, we estimated the empirical DFE for every population using `fitdadi` (Kim *et al.* 2017) in 100 replicated runs (see Figures S5, S8 for supplementary methods and results).

Data and code availability

Statistical analyses were conducted in `R` v.4.1.3 (R Core Team 2018), unless otherwise specified. Genetic data is archived at NCBI SRA (accession PRJNA773763). Code and simulation output is available on GitHub (`https://github.com/LZeitler/selfing_expansion`).

Acknowledgments

We thank Drs. Marco Andrello, Michele Di Musciano, Marta Binaghi, Paola Morini, Marco Caccianiga, Alessandro Alessandrini, Rodolfo Gentili, and Enzo Bona for essential help in finding wild populations of *A. alpina* to sample in Italy and France. We thank Dr. Pamela Nicholson and the team at the Bern NGS Platform for sequencing and troubleshooting help as well as Ryan Gutenkunst for help troubleshooting the dadi analyses. We thank Stephan Peischl for useful feedback on the manuscript and Xuejing Wang for statistical advice. This research was funded by Swiss National Science Foundation Ambizione grant #PZ00P3_185952 to KJG.

Author Contributions

LZ contributed to further developing the study's idea, designed, wrote, and ran the simulations, led the empirical analyses, participated in sample collection and DNA extractions, and wrote the manuscript. CP contributed to further developing the study's idea and to editing the manuscript. KJG conceived the original idea for the study, coordinated the study, contributed to designing the simulations, participated in sample collection and DNA extractions, and contributed to writing and editing the manuscript.

References

Agrawal, A. F. and M. C. Whitlock, 2011 Inferences About the Distribution of Dominance Drawn From Yeast Gene Knockout Data. Genetics **187**: 553–566.

Alexander, D. H., J. Novembre, and K. Lange, 2009 Fast model-based estimation of ancestry in unrelated individuals. Genome Research **19**: 1655–1664.

Ansell, S. W., M. Grundmann, S. J. Russell, H. Schneider, and J. C. Vogel, 2008 Genetic discontinuity, breeding-system change and population history of *Arabis alpina* in the Italian Peninsula and adjacent Alps. Molecular Ecology **17**: 2245–2257.

Arunkumar, R., R. W. Ness, S. I. Wright, and S. C. H. Barrett, 2015 The Evolution of Selfing Is Accompanied by Reduced Efficacy of Selection and Purging of Deleterious Mutations. Genetics **199**: 817–829.

Baker, H. G., 1967 Support for Baker's Law-As a Rule. Evolution **21**: 853–856.

Balick, D. J., R. Do, C. A. Cassa, D. Reich, and S. R. Sunyaev, 2015 Dominance of Deleterious Alleles Controls the Response to a Population Bottleneck. PLOS Genetics **11**: e1005436.

Barrett, S. C. H., 2002 The evolution of plant sexual diversity. Nature Reviews Genetics **3**: 274–284.

Barrett, S. C. H., 2013 IV.8. Evolution of Mating Systems: Outcrossing versus Selfing. In *IV.8. Evolution of Mating Systems: Outcrossing versus Selfing*, pp. 356–362, Princeton University Press.

Barrett, S. C. H., R. Arunkumar, and S. I. Wright, 2014 The demography and population genomics of evolutionary transitions to self-fertilization in plants. Philosophical Transactions of the Royal Society B: Biological Sciences **369**: 20130344.

Bataillon, T. and M. Kirkpatrick, 2000 Inbreeding depression due to mildly deleterious mutations in finite populations: Size does matter. Genetics Research **75**: 75–81.

Bolger, A. M., M. Lohse, and B. Usadel, 2014 Trimmomatic: A flexible trimmer for Illumina sequence data. Bioinformatics **30**: 2114–2120.

Bosshard, L., I. Dupanloup, O. Tenaillon, R. Bruggmann, M. Ackermann, *et al.*, 2017 Accumulation of Deleterious Mutations During Bacterial Range Expansions. Genetics **207**: 669–684.

Brandvain, Y. and S. I. Wright, 2016 The Limits of Natural Selection in a Nonequilibrium World. Trends in Genetics **32**: 201–210.

Buehler, D., R. Graf, R. Holderegger, and F. Gugerli, 2012 Contemporary gene flow and mating system of *Arabis alpina* in a Central European alpine landscape. Annals of Botany **109**: 1359–1367.

Burton, O. J. and J. M. J. Travis, 2008 The frequency of fitness peak shifts is increased at expanding range margins due to mutation surfing. Genetics **179**: 941–950.

Charlesworth, B., 1992 Evolutionary rates in partially self-fertilizing species. The American Naturalist **140**: 126–148.

Charlesworth, D. and B. Charlesworth, 1979 The evolution and breakdown of S-allele systems. Heredity **43**: 41–55.

Charlesworth, D. and B. Charlesworth, 1987 Inbreeding Depression and its Evolutionary Consequences. Annual Review of Ecology and Systematics **18**: 237–268.

Cingolani, P., A. Platts, L. L. Wang, M. Coon, T. Nguyen, *et al.*, 2012 A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. Fly **6**: 80–92.

Courchamp, F., T. Clutton-Brock, and B. Grenfell, 1999 Inverse density dependence and the Allee effect. Trends in Ecology & Evolution **14**: 405–410.

Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, *et al.*, 2021 Twelve years of SAMtools and BCFtools. GigaScience **10**: giab008.

Davis, M. B. and R. G. Shaw, 2001 Range Shifts and Adaptive Responses to Quaternary Climate Change. Science **292**: 673–679.

Dennis, B., 1989 Allee effects: Population growth, critical density, and the chance of extinction. Natural Resource Modeling **3**: 481–538.

Do, R., D. Balick, H. Li, I. Adzhubei, S. Sunyaev, *et al.*, 2015 No evidence that selection has been less effective at removing deleterious mutations in Europeans than in Africans. Nature Genetics **47**: 126–131.

Encinas-Viso, F., A. G. Young, and J. R. Pannell, 2020 The loss of self-incompatibility in a range expansion. Journal of Evolutionary Biology **33**: 1235–1244.

Eriksson, M. and M. Rafajlović, 2021 The Effect of the Recombination Rate between Adaptive Loci on the Capacity of a Population to Expand Its Range. The American Naturalist **197**: 526–542.

Ewels, P., M. Magnusson, S. Lundin, and M. Käller, 2016 MultiQC: Summarize analysis results for multiple tools and samples in a single report. Bioinformatics **32**: 3047–3048.

Excoffier, L., M. Foll, and R. J. Petit, 2009 Genetic Consequences of Range Expansions. Annual Review of Ecology, Evolution, and Systematics **40**: 481–501.

Eyre-Walker, A. and P. D. Keightley, 2007 The distribution of fitness effects of new mutations. Nature Reviews Genetics **8**: 610–618.

Garrison, E. and G. Marth, 2012 Haplotype-based variant detection from short-read sequencing. arXiv:1207.3907 [q-bio] .

Gilbert, K. J., S. Peischl, and L. Excoffier, 2018 Mutation load dynamics during environmentally-driven range shifts. PLOS Genetics **14**: e1007450.

Gilbert, K. J., N. P. Sharp, A. L. Angert, G. L. Conte, J. A. Draghi, *et al.*, 2017 Local Adaptation Interacts with Expansion Load during Range Expansion: Maladaptation Reduces Expansion Load. The American Naturalist **189**: 368–380.

Gilbert, K. J., S. Zdraljevic, D. E. Cook, A. D. Cutter, E. C. Andersen, *et al.*, 2022 The distribution of mutational effects on fitness in *Caenorhabditis elegans* inferred from standing genetic variation. Genetics **220**: iyab166.

Glémin, S., 2003 How Are Deleterious Mutations Purged? Drift Versus Nonrandom Mating. Evolution **57**: 2678–2687.

Glémin, S., 2007 Mating Systems and the Efficacy of Selection at the Molecular Level. Genetics **177**: 905–916.

González-Martínez, S. C., K. Ridout, and J. R. Pannell, 2017 Range Expansion Compromises Adaptive Evolution in an Outcrossing Plant. Current Biology **27**: 2544–2551.e4.

Goodwillie, C., S. Kalisz, and C. G. Eckert, 2005 The Evolutionary Enigma of Mixed Mating Systems in Plants: Occurrence, Theoretical Explanations, and Empirical Evidence. Annual Review of Ecology, Evolution, and Systematics **36**: 47–79.

Hallatschek, O. and D. R. Nelson, 2008 Gene surfing in expanding populations. Theoretical Population Biology **73**: 158–170.

Hallatschek, O. and D. R. Nelson, 2010 Life at the Front of an Expanding Population. Evolution **64**: 193–206.

Haller, B. C. and P. W. Messer, 2019 SLiM 3: Forward Genetic Simulations Beyond the Wright–Fisher Model. Molecular Biology and Evolution **36**: 632–637.

Halligan, D. L. and P. D. Keightley, 2009 Spontaneous Mutation Accumulation Studies in Evolutionary Genetics. Annual Review of Ecology, Evolution, and Systematics **40**: 151–172.

Hedrick, P. W., 2002 Lethals in Finite Populations. Evolution **56**: 654–657.

Henn, B. M., L. R. Botigué, C. D. Bustamante, A. G. Clark, and S. Gravel, 2015 Estimating the mutation load in human genomes. Nature Reviews Genetics **16**: 333–343.

Henn, B. M., L. R. Botigué, S. Peischl, I. Dupanloup, M. Lipatov, *et al.*, 2016 Distance from sub-Saharan Africa predicts mutational load in diverse human genomes. Proceedings of the National Academy of Sciences **113**: E440–E449.

Hewitt, G. M., 2004 Genetic consequences of climatic oscillations in the Quaternary. Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences **359**: 183–195.

Igic, B. and J. W. Busch, 2013 Is self-fertilization an evolutionary dead end? New Phytologist **198**: 386–397.

Jiao, W.-B., G. G. Accinelli, B. Hartwig, C. Kiefer, D. Baker, *et al.*, 2017 Improving and correcting the contiguity of long-read genome assemblies of three plant species using optical mapping and chromosome conformation capture data. Genome Research **27**: 778–786.

Keightley, P. D., 1994 The distribution of mutation effects on viability in *Drosophila melanogaster*. Genetics **138**: 1315–1322.

Kiełbasa, S. M., R. Wan, K. Sato, P. Horton, and M. C. Frith, 2011 Adaptive seeds tame genomic sequence comparison. Genome Research **21**: 487–493.

Kim, B. Y., C. D. Huber, and K. E. Lohmueller, 2017 Inference of the Distribution of Selection Coefficients for New Nonsynonymous Mutations Using Large Samples. Genetics **206**: 345–361.

Kirkpatrick, M. and N. H. Barton, 1997 Evolution of a Species' Range. The American Naturalist **150**: 1–23.

Kirkpatrick, M. and P. Jarne, 2000 The Effects of a Bottleneck on Inbreeding Depression and the Genetic Load. The American Naturalist **155**: 154–167.

Klopfstein, S., M. Currat, and L. Excoffier, 2006 The Fate of Mutations Surfing on the Wave of a Range Expansion. Molecular Biology and Evolution **23**: 482–490.

Korunes, K. L. and K. Samuk, 2021 Pixy: Unbiased estimation of nucleotide diversity and divergence in the presence of missing data. Molecular Ecology Resources **21**: 1359–1368.

Laenen, B., A. Tedder, M. D. Nowak, P. Toräng, J. Wunder, *et al.*, 2018 Demography and mating system shape the genome-wide impact of purifying selection in *Arabis alpina*. Proceedings of the National Academy of Sciences **115**: 816–821.

Li, H. and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics (Oxford, England) **25**: 1754–1760.

Lloyd, D. G. and D. J. Schoen, 1992 Self- and Cross-Fertilization in Plants. I. Functional Dimensions. International Journal of Plant Sciences **153**: 358–369.

Lynch, M., J. Conery, and R. Bürger, 1995 Mutational Meltdowns in Sexual Populations. Evolution **49**: 1067–1080.

Madrid, E., E. Severing, E. de Ansorena, C. Kiefer, L. Brand, *et al.*, 2021 Transposition and duplication of MADS-domain transcription factor genes in annual and perennial *Arabis* species modulates flowering. Proceedings of the National Academy of Sciences **118**: e2109204118.

Moeller, D. A., M. A. Geber, V. M. Eckhart, and P. Tiffin, 2012 Reduced pollinator service and elevated pollen limitation at the geographic range limit of an annual plant. Ecology **93**: 1036–1048.

Ohta, T. and C. C. Cockerham, 1974 Detrimental genes with partial selfing and effects on a neutral locus*. Genetics Research **23**: 191–200.

Pannell, J. R. and S. C. H. Barrett, 1998 Baker's Law Revisited: Reproductive Assurance in a Metapopulation. Evolution **52**: 657–668.

Parmesan, C., 2006 Ecological and Evolutionary Responses to Recent Climate Change. Annual Review of Ecology, Evolution, and Systematics **37**: 637–669.

Pedersen, B. S. and A. R. Quinlan, 2018 Mosdepth: Quick coverage calculation for genomes and exomes. Bioinformatics **34**: 867–868.

Peischl, S., I. Dupanloup, A. Foucal, M. Jomphe, V. Bruat, *et al.*, 2018 Relaxed Selection During a Recent Human Expansion. Genetics **208**: 763–777.

Peischl, S., I. Dupanloup, M. Kirkpatrick, and L. Excoffier, 2013 On the accumulation of deleterious mutations during range expansions. Molecular Ecology **22**: 5972–5982.

Peischl, S. and L. Excoffier, 2015 Expansion load: Recessive mutations and the role of standing genetic variation. Molecular Ecology **24**: 2084–2094.

Peischl, S., M. Kirkpatrick, and L. Excoffier, 2015 Expansion Load and the Evolutionary Dynamics of a

Species Range. The American Naturalist **185**: E81–E93.

Polechová, J. and N. H. Barton, 2015 Limits to adaptation along environmental gradients. Proceedings of the National Academy of Sciences **112**: 6401–6406.

Porcher, E. and R. Lande, 2005 Loss of Gametophytic Self-Incompatibility with Evolution of Inbreeding Depression. Evolution **59**: 46–60.

Pujol, B., S.-R. Zhou, J. Sanchez Vilas, and J. R. Pannell, 2009 Reduced inbreeding depression after species range expansion. Proceedings of the National Academy of Sciences **106**: 15379–15383.

R Core Team, 2018 R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing.

Reusch, T. B. H., 2001 Fitness-consequences of geitonogamous selfing in a clonal marine angiosperm (*Zostera marina*). Journal of Evolutionary Biology **14**: 129–138.

Rogivue, A., R. R. Choudhury, S. Zoller, S. Joost, F. Felber, *et al.*, 2019 Genome-wide variation in nucleotides and retrotransposons in alpine populations of *Arabis alpina* (Brassicaceae). Molecular Ecology Resources **19**: 773–787.

Simons, Y. B. and G. Sella, 2016 The impact of recent population history on the deleterious mutation load in humans and close evolutionary relatives. Current Opinion in Genetics & Development **41**: 150–158.

Slatkin, M. and L. Excoffier, 2012 Serial Founder Effects During Range Expansion: A Spatial Analog of Genetic Drift. Genetics **191**: 171–181.

Stephens, P. A. and W. J. Sutherland, 1999 Consequences of the Allee effect for behaviour, ecology and conservation. Trends in Ecology & Evolution **14**: 401–405.

Tedder, A., S. Carleial, M. Gołębiewska, C. Kappel, K. K. Shimizu, *et al.*, 2015 Evolution of the selfing syndrome in *Arabis alpina* (Brassicaceae). PLOS ONE **10**: e0126618.

Vallejo-Marín, M. and M. K. Uyenoyama, 2004 On the Evolutionary Costs of Self-Incompatibility: Incomplete Reproductive Compensation Due to Pollen Limitation. Evolution **58**: 1924–1935.

Vieira, F. G., M. Fumagalli, A. Albrechtsen, and R. Nielsen, 2013 Estimating inbreeding coefficients from NGS data: Impact on genotype calling and allele frequency estimation. Genome Research **23**: 1852–1861.

Wang, J., W. G. Hill, D. Charlesworth, and B. Charlesworth, 1999 Dynamics of inbreeding depression due to deleterious mutations in small populations: Mutation parameters and inbreeding rate. Genetical Research **74**: 165–178.

Willi, Y., M. Fracassetti, S. Zoller, and J. Van Buskirk, 2018 Accumulation of Mutational Load at the Edges of a Species Range. Molecular Biology and Evolution **35**: 781–791.

Willis, J. H., 1999 The Role of Genes of Large Effect on Inbreeding Depression in *Mimulus guttatus*. Evolution **53**: 1678–1691.

Supplementary

Figure S1: The proportion of sites fixed for deleterious alleles (A), the mean counts of deleterious loci (B), and the mean counts of deleterious alleles (C), all assessed at the end of the simulations for core and edge populations across selfing rates. Whiskers indicate the 1.5 interquartile range.

Figure S2: Trajectories for the mean observed heterozygosity over relative time, as described in Figure 2A, but now including all simulated selfing rates.

Figure S3: Trajectories for the mean count of lethal alleles over relative time, as described in Figure 2B, but now including all simulated selfing rates.

Figure S4: Results from $K = 2$ to $K = 15$ from admixture analyses run on the combined empirical dataset across Europe. The lowest CV error is for $K = 14$, however it is most useful to compare the populations structure across values of K to see how well this matches known geography and demographic history of the populations. We observe clean distinctions among our geographic regions sampled (indicated above the bar plots), with evidence for some gene flow across geographic space as one observes higher K values.

Figure S5: We inferred the demographic history of each of our newly sampled populations of *A. alpina* along with the densely sampled Swiss populations using `dadi`. This is a necessary step to account for the demography when inferring the DFE with `fitdadi`. This also allowed us to confirm if this newly inferred demographic history is consistent with past studies in *A. alpina*. The best-fitting models for our populations, based on AIC, were "bottlegrowth" models, indicating a past bottleneck followed by exponential growth (Es, Ca, Gf, Gz, Po), three epoch models, indicating a bottleneck followed by a sudden size change (Pa, Pi, Ma, Am, Br, Cc, Ga, Gs, La, Mv, Se), and the standard neutral model (St). Populations St and Gz were the only instances where competing models fitted approximately equally well (see Supplementary Table 2), therefore results for these population should be interpreted with caution. With the exception of Es, all Alpine populations best fit to three epoch models. Central Italian populations (light blue) show the most historic bottlenecks and the largest ancestral populations sizes. This is consistent with this region of highly outcrossing plants being subject to the last glacial maximum. Northern Italian populations (dark blue) show more recent bottlenecks and reduced ancestral sizes relative to central Italy, potentially reflecting their expansion northward. French and Swiss Alpine populations both showed the most recent bottlenecks and the smallest historic population sizes, consistent with both their shift to selfing and their more recent range expansion. Depleted genetic diversity along the axis of an expanding species range is expected (Pujol *et al.* 2009; Peischl *et al.* 2013, 2015), as is decreased N_e due to inbreeding and thus loss of diversity (Keller and Waller 2002; Charlesworth and Willis 2009). These demographic inferences thus match our understanding of both the mating system shift and the range expansion that these populations experienced.

Figure S6: Observed (known) mean fitness from simulations for core (green), interior (orange, purple) and edge (pink) demes compared to the inverse of the count of deleterious loci (A), both after the range expansion is complete. The count of deleterious loci serves as a proxy for recessive load, which we find best correlates to fitness, compared to the additive model (B), where load is predicted by counting alleles. Results are for simulations with $h = 0.3$ for non-lethal deleterious mutations.

Figure S7: Recessive (A) and additive (B) genetic load compared with known simulated fitness to infer load when all non-lethal deleterious mutations are perfectly additive ($h = 0.5$). Data is from a supplementary set of simulations with these dominance parameters. This repeats the same analyses as Figure S6, except now for simulations with additive mutations. This result again finds that the recessive model predicts load better ($R^2 = 0.70, P < 0.001$) than the additive model ($R^2 = 0.20, P < 0.001$).

Figure S8: We inferred the DFE of each *A. alpina* population in the Italy-Alps expansion zone using `fitdadi` (Kim *et al.* 2017) from `dadi` (Gutenkunst *et al.* (2009) in python 3.8.12. We used the SNPeff annotation to construct polarized site-frequency spectra for neutral and deleterious sites after subsampling to a maximum population size of 20 individuals. To estimate demographic parameters, we tested the default single population demographic models (standard neutral model, two-epoch, growth, bottlegrowth, three-epoch) and two models accounting for inbreeding (standard neutral with inbreeding, two-epoch with inbreeding). We assumed a per base pair mutation rate of $\mu = 7 \times 10^{-9}$ per generation, ran the default optimization for 100 replicates, and selected the best fit parameters within each demographic model based on likelihood and the best fit demographic model based on AIC. For `fitdadi`, we additionally assumed $L_{ns}/L_s = 2.85$, dominance coeffient $h = 0.3$ and estimated the DFE for each model in 100 optimizations. We then chose the best-fit DFE optimization based on likelihood for each population for the previously chosen demographic model.

DFE results from *A. alpina* populations across the Italian-Alpine range expansion for outcrossing populations from Abruzzo (light blue) and the Apuan Alps (dark blue) are compared to the selfing populations that have undergone range expansions into the French Alps (light green) and the Swiss Alps (dark green). We found mean proportions across all populations of 65.4% and 24.8% in the weakest and strongest selection classes, respectively. Less than 5% of sites segregated in the two intermediate selection classes. These proportions varied only marginally between core Italian populations (mean proportions 22.6% and 67.9% for weakest and strongest classes, respectively) and between expanded French and Swiss populations (means proportions 27.4% and 62.6% for weakest and strongest class).

Figure S9: Fixation of predicted neutral (dark purple), deleterious (light purple) and loss of function (LoF, orange) sites per population. Y-axis shows the proportion of fixes sites in focal population and allele category. We found that neutral sites fixed at the highest proportions (mean 0.505%), while LoF sites were at the smallest proportions fixed (mean 0.314%), indicative of their highly deleterious effect. French populations Br and La had the highest overall fixation proportions of any class (0.948%), while samples from the Abruzzo region had the lowest (0.228%). Swiss population showed intermediate neutral fixation but LoF proportions similar to Italian populations.

References

Charlesworth, D. and J. H. Willis, 2009 The genetics of inbreeding depression. Nature Reviews Genetics **10**: 783–796.

Gutenkunst, R. N., R. D. Hernandez, S. H. Williamson, and C. D. Bustamante, 2009 Inferring the Joint Demographic History of Multiple Populations from Multidimensional SNP Frequency Data. PLOS Genetics **5**: e1000695.

Keller, L. F. and D. M. Waller, 2002 Inbreeding effects in wild populations. Trends in Ecology & Evolution **17**: 230–241.

Kim, B. Y., C. D. Huber, and K. E. Lohmueller, 2017 Inference of the Distribution of Selection Coefficients for New Nonsynonymous Mutations Using Large Samples. Genetics **206**: 345–361.

Peischl, S., I. Dupanloup, M. Kirkpatrick, and L. Excoffier, 2013 On the accumulation of deleterious mutations during range expansions. Molecular Ecology **22**: 5972–5982.

Peischl, S., M. Kirkpatrick, and L. Excoffier, 2015 Expansion Load and the Evolutionary Dynamics of a Species Range. The American Naturalist **185**: E81–E93.

Pujol, B., S.-R. Zhou, J. Sanchez Vilas, and J. R. Pannell, 2009 Reduced inbreeding depression after species range expansion. Proceedings of the National Academy of Sciences **106**: 15379–15383.