Calcium flux through ER-TGN contact sites facilitates cargo export

Running title: Secretory cargo sorting requires Ca[2+] flux in ER/TGN contact sites

Bulat R. Ramazanov[1]†, Rosaria Di Martino[2]†, Abhishek Kumar[3], Anup Parchure[1], Yeongho Kim[1],

Oliver Griesbeck[4], Martin A. Schwartz[1,3,5], Alberto Luini[2]†, Julia von Blume[1]†*

[1]Department of Cell Biology, Yale University School of Medicine, New Haven, CT

[2] Institute of Biochemistry and Cell Biology, National Research Council, Via Pietro Castellino

111, 80131 Naples, Italy

[3]Yale Cardiovascular Research Center, Cardiovascular Yale School of Medicine, Yale University,

[4] Max Planck Institute of Neurobiology, Martinsried, Germany

[5] Department of Biomedical Engineering, Yale University, New Haven, CT

† Authors have contributed equally to the work

*Corresponding author

Julia von Blume

333 Cedar Street

New Haven, CT 06510

USA

Yale email: julia.vonblume@yale.edu

Phone: +1 475 325 8124

Summary

The current study demonstrates the trafficking of Cab45 clients relies on Ca^{2+} flux between the endoplasmic reticulum (ER) and trans-Golgi Network (TGN). This process requires the activity of IP3 receptors, present in ER membranes, and depends on the integrity of the membrane contact site between these two organelles.

Abstract

Ca^{2+} influx into the trans-Golgi Network (TGN) promotes secretory cargo sorting by the Ca^{2+}-ATPase SPCA1 and the luminal Ca^{2+} binding protein Cab45. Cab45 oligomerizes upon a local Ca^{2+} influx, and Cab45 oligomers sequester and separate soluble secretory cargo from the bulk flow of proteins in the TGN. However, how this Ca^{2+} flux into the lumen of the TGN is achieved remains elusive, as the cytosol has a very low steady-state Ca^{2+} concentration. The TGN forms membrane contact sites (MCS) with the Endoplasmic Reticulum (ER), whereby the close apposition of the two organelles allows protein-mediated exchange of molecular species such as lipids. Here we show that TGN export of Cab45 clients requires the integrity of ER-TGN MCS and IP3R-dependent Ca^{2+} fluxes in the MCS, suggesting Ca^{2+} transfer between these organelles. Using a MCS targeted Ca^{2+} FRET sensor module, we measure the Ca^{2+} flow in these sites in real-time. These data show for the first time that ER-TGN MCS facilitate Ca^{2+} transfer required for SPCA1-dependent cargo sorting and export from the TGN and thus solves a fundamental question in cell biology.

Introduction

Protein secretion is a fundamental process and facilitates the integrity and cell-cell communication of multiple-tissue organisms (Uhlen et al., 2015). Secreted proteins are synthesized in the Endoplasmic Reticulum (ER) and transported to the Golgi apparatus in COPII coated vesicle (Barlowe and Miller, 2013; Gillon et al., 2012; Zanetti et al., 2013). Upon reaching the Golgi apparatus, these cargo molecules further transit from the *cis* to *trans* Golgi cisterna and finally reach trans Golgi Network (TGN) (De Matteis and Luini, 2008; Di Martino et al., 2019; Ford et al., 2021; Guo et al., 2014; Kienzle and von Blume, 2014). Protein transport between Golgi cisternae and towards the TGN is still highly debated and several models have been proposed (Dunlop et al., 2017; Dunphy and Rothman, 1985; Glick et al., 1997; Glick and Luini, 2011; Glick and Nakano, 2009; Lujan and Campelo, 2021; Nakano and Luini, 2010). At the TGN, these proteins are sorted and packed into different transport carriers to reach their destination such as the cell surface, the endosomal system, or secretory granules in specialized cells (Mostov and Cardone, 1995; Stalder and Gershlick, 2020; Tang, 2001).

Protein sorting at the TGN is a complex process involving physical features of the cargoes themselves and elements of the local environment and so remains a poorly understood process for many proteins. Understanding how soluble secretory proteins can be accurately and specifically sorted to the plasma membrane is a particular challenge. There are no cargo receptors that connect these soluble molecules to the TGN membrane (Kienzle and von Blume, 2014; Pakdel and von Blume, 2018; Ramazanov et al., 2021).

Our work and others have shown that Calcium (Ca^{2+}) is a major regulator of cargo sorting at the TGN. Secretory Pathway ATPase 1 (SPCA1) pumps Ca^{2+} from the cytoplasm into the TGN lumen in an ATP-dependent manner (Kienzle et al., 2014; Lebreton et al., 2021; Lissandron

et al., 2010; Missiaen et al., 2007; Pizzo et al., 2011; Pizzo et al., 2010; Sepulveda et al., 2008;

von Blume et al., 2011; Wong et al., 2013). In response to luminal Ca^{2+} influxes, the Golgi

resident protein Cab45 oligomerizes and captures cargo molecules before they are packed into

sphingomyelin-rich vesicles that bud from the TGN (Crevenna et al., 2016; Deng et al., 2018;

Scherer et al., 1996; von Blume et al., 2012). Despite the importance of Ca^{2+} in these processes,

the source of the Ca^{2+} that is pumped into the TGN lumen by SPCA1 is unknown as the cytosolic

Ca^{2+} concentrations are limited to the low nanomolar scale (Berridge et al., 2003).

The TGN forms MCSs with the ER where lipids are transferred to mediate non-vesicular

inter organelle communication (Masone et al., 2019; Venditti et al., 2020; Venditti et al., 2019).

Endoplasmic reticulum to TGN membrane contact sites (ER-TGN MCSs) contain tethering

proteins such as vesicle-associated membrane proteins A and B (VAPA and VAPB) and lipid

transfer proteins such as Oxysterol-binding protein 1 (OSBP1) (Lehto and Olkkonen, 2003).

OSBP1 possesses dual organelle targeting motifs as it is able to bind the ER proteins VAPA and

VAPB through an acidic track (FFAT) domain, and it is also able to bind to the TGN through

Pleckstrin homology (PH) domain, which recognizes Phosphatidylinisitol-4-phosphate (PI4P) or

Arf1-GTP on TGN membrane (Kawano et al., 2006; Kumagai and Hanada, 2019; Mesmin et al.,

2017). At the ER/TGN interface, OSBP1 counter transports PI4P from the TGN and cholesterol

from the ER in a the process mediated by the oxysterol-binding domain (OBD) of OSBP1 that

binds PI4P or cholesterol in a mutually exclusive manner (Mesmin et al., 2013).

Recent studies report the importance of lipid transfer in ER-TGN MCSs in regulating

protein export from the TGN (Wakana et al., 2021). In addition to their implication in lipid transfer,

MCSs allow Ca^{2+} transfer between the ER and other organelles such as the mitochondria (Kelly,

1985; Pfeffer and Rothman, 1987; Rizzuto et al., 1993; Rizzuto et al., 1998). We therefore

hypothesized that ER-TGN MCSs could provide Ca^{2+} to facilitate SPCA1 and Cab45-dependent

cargo sorting at the TGN. Here we show that the trafficking of Cab45 clients relies on the IP3

receptor (IP3R), present in ER membranes, and on the integrity of the ER-TGN MCSs. We

generated a MCS specific sensor to measure Ca^{2+} flux within these sites, revealing that this Ca^{2+}

flux plays an essential role for cargo export from the TGN and is dependent on tethering between

ER and TGN. With these data we solve a major unresolved question in cell biology.

Results and discussion

Inhibition of IP3R delays TGN export of Cab45 clients

Our previous work demonstrates that sorting soluble secretory proteins requires a transient SPCA1-mediated Ca^{2+} influx into the TGN to facilitate Cab45 oligomerization (Crevenna et al., 2016; Deng et al., 2018; von Blume et al., 2012). The cytosolic Ca^{2+} concentrations at steady state are in the low nanomolar range thus may not provide the amount of Ca^{2+} ions required to promote sorting of Cab45 clients (Crevenna et al., 2016; Pizzo et al., 2011). Therefore, we hypothesized that with the established MCS between the ER and the TGN, and the ER being the largest Ca^{2+} store within the cell, it is a likely candidate to provide Ca^{2+} for sorting at TGN, and this process could be mediated through an IP3 receptor (IP3R)-dependent mechanism.

To investigate if the release of Ca^{2+} from the ER has an impact on the sorting and export of soluble secretory cargo molecules from at TGN, we analyzed trafficking and secretion of the well-established Cab45-clients: cartilage oligomeric protein (COMP) and Lysozyme C (LyzC) (von Blume 2011; von Blume 2012) in the presence and absence of the IP3R antagonist (2-APB) (Maruyama et al., 1997). We used the retention using selective hooks (RUSH) system to quantify the trafficking and packaging of COMP or LyzC, respectively, into secretory vesicles in HeLa cells in the presence of 2-APB (70 μM) or DMSO (control) (Boncompain et al., 2012). To this end, HeLa cell lines were transfected with RUSH-constructs containing COMP-EGFP or LyzC-EGFP to analyze intracellular trafficking of EGFP-fused proteins at different time points after synchronous release from the ER by biotin addition (**Figure 1A**). We observed simultaneous export of COMP-EGFP from the ER in control and 2-APB treated cells (**Figure 1B [0, 20 min]**). However, the appearance of cytosolic vesicles (TGN carriers) was significantly delayed in cells treated with 2-APB compared to control cells at later time points (**Figure 1B, C [30, 40, 60 min]**).

To confirm the observed phenotype on a cell population, we performed RUSH experiments

using COMP-EGFP as cargo in HeLa line treated with DMSO (control) or 2-APB with subsequent

FACS analysis. We used this assay to quantify the intracellular accumulation of EGFP-COMP in

DMSO or 2-APB treated cells. Cells were fixed at 0, 30, 60 and 120 min after biotin addition and

10^4 cells for each time-point were analyzed by FACS. We calculated the average arithmetical value

for fluorescent intensity of COMP-EGFP obtained from the FL-1 channel for each sample. The

arithmetical average values for fluorescence intensity of COMP-EGFP from 2-APB treated cells

were 1.8 and 2.1-fold higher than in control cells after 60 and 120 minutes after biotin addition,

respectively, indicating a longer residence of the EGFP-COMP inside cells in 2-APB treated cells

confirming our microscopy observations (**Figure 1D, Supplementary S1A, S1C**).

To correlate these results with the actual TGN exit of the cargo molecules, we applied live-

cell imaging of the exiting EGFP-tagged cargo molecules in the presence of the GALNT1-BFP

TGN marker. We measured time-dependent changes in fluorescence intensity of EGFP-tagged

protein within ROI of TGN defined by the GALNT1-BFP signal. These results showed that the

reduction in the number of vesicles in 2-APB treated cells in the RUSH experiments was consistent

with prolonged residence of cargo in the TGN compared to DMSO treated cells (**Figure 1E**). To

quantify this phenotype, we applied a non-linear regression function of the intensity values on the

plot shown on Figure 1D (**Supplementary S1E**). In addition, we calculated span values for each

curve representing changes of LyzC-EGFP intensity within Golgi ROI. The span was defined as

the difference between the fluorescence intensity of EGFP at the starting point and predicted

plateau for each curve, representing a change in fluorescence intensity during the experiment. The

span value calculated for LyzC-EGFP expressing cells in the presence of 2-APB exhibited a two-

fold decrease compared to control cells (span value for 2-APB and DMSO samples were 0.8 and

1.5 respectively), indicating a significant defect in TGN export of LyzC-EGFP in these cells.

To further validate that inhibition of IP3R reduces secretion of Cab45 client LyzC from the

cells, we performed a secretion assay. We generated stable cell lines expressing EGFP tagged

LyzC under a constitutive promoter by lentiviral transduction of HeLa cells. Secretion assays were

performed in a complete growth medium, and secreted LyzC-EGFP in the supernatant was

immunoprecipitated using GFP trap agarose beads prior to analysis by western blotting. The

western blot analysis revealed a 48% reduction in secretion LyzC-EGFP into the cell culture media

in 2-APB treated condition compared to DMSO treated control cells (**Figure 1F**).

These data suggested that the activity of IP3 receptors impacts the TGN export and

secretion of Cab45-clients. Previous work has shown that Golgi localized IP3 receptors have no

impact on SPCA1 dependent Ca^{2+} uptake (Wong et al., 2013), therefore we speculated that Ca^{2+}

flow between these organelles might be facilitated by ER-TGN MCS (Ramazanov et al., 2021).

MCS integrity is essential for TGN export of Cab45 clients

VAPA and VAPB proteins play an essential role in maintaining ER-TGN MCS integrity

by tethering ER and TGN. Deleting these proteins from cells leads to a reduction of contacts

between the organelles (Lev, 2010; Phillips and Voeltz, 2016). To examine the role of the integrity

of the ER-TGN contact sites on trafficking of Cab45-clients, we performed RUSH experiments,

and we analyzed the trafficking of COMP-EGFP in VAPA/VAPB depleted HeLa cells. To this

end, HeLa cells were transfected with non-targeting siRNA as control or siRNAs targeting VAPA

and VAPB. After 24 hours, HeLa cell lines were transfected with RUSH-constructs expressing

COMP-EGFP. RUSH experiments were performed, and samples were fixed after different

timepoints of biotin addition. Immunofluorescence images captured at 30, 40, and 60 minutes after

biotin addition showed that compared to control cells, VAPA/VAPB siRNA treated cells exhibited

a significant delay in the formation of COMP-EGFP containing post-Golgi vesicles (**Figure 2A**

[30, 40, 60 min]). Consistent with this result, quantification of post-Golgi vesicles after different

time points from a randomly selected population of control siRNA or VAPA/VAPB siRNA treated

cells revealed a decrease in the amount of TGN derived vesicles (**Figure 2B**).

We confirmed the phenotype of delayed export of Cab45 clients seen in VAPA and VAPB

knockdown with RUSH LyzC or COMP, respectively, in control versus VAPA/VAPB-dKO HeLa

cell lines (Dong et al., 2016) (see immunofluorescence **Supplementary S2A, S2B [0, 20 min]**),

and quantification (**Supplementary S2C, S2D**). In parallel, we measured time-dependent

intracellular decrease of COMP-EGFP by FACS in control versus VAPA/VAPB deficient cells.

The average intensity values in the GFP channel from VAPA and VAPB deficient cells were 1.5-

fold higher than in control cells at the 60 and 120 minutes after biotin addition, indicating

accumulation of COMP-EGFP in these cells (**Figure 2C, Supplementary S1A, S1C**).

To demonstrate that the secretion defect is caused by impaired TGN export we performed

live-cell imaging of the exiting EGFP-tagged Cab45 clients in the presence of the GALNT1-BFP

TGN marker. We observed time-dependent changes of LyzC-EGFP intensity within the TGN ROI

and a 1.6-fold difference in span values in control versus VAPA and VAPB depleted cells. In

addition, we observed a prolonged residence of LyzC in the TGN of HeLa cell lines transfected

with siRNA to VAPA and VAPB compared control cells (**Figure 2D, Supplementary S1F**). The

efficiency of knockdowns of VAPA and VAPB proteins were analyzed by western blotting

(**Figure 2E**).

To investigate that this phenotype is specific for certain secretory soluble proteins and does

not affect other proteins that are transported by bulk flow secretion, we measured the time-

dependent intracellular decrease of COMP-EGFP and EQ-sol-GFP by FACS in control versus

VAPA/VAPB deficient HeLa cells as well as 2-APB treated cells. These data indicated that the

secretion of EQ-sol, non-specific bulk flow secretion marker, representing soluble non-toxic

version of equinatoxin II-based reporter EQ-SM (Deng et al., 2016), is unaffected in VAPA and

VAPB-depleted HeLa line compared to WT cell line (**Suppl. S1A-C**).

Together these data showed that the TGN export of the Cab45 clients COMP and LyzC

from TGN requires intact MCSs between ER and TGN (**Figure 2**) and IP3R-dependent Ca^{2+}

release from the ER (**Figure 1**). These data also suggested that ER-TGN MCS could serve as

potential sites for Ca^{2+} transfer.

Targeting Twitch based FRET sensors to ER-TGN MCS

Our next goal was to directly test potential Ca^{2+} flow at ER-TGN MCSs. We hypothesized

that these MCSs would be potential structures that could serve as hotspots for Ca^{2+} transfer

between the organelles. To evaluate possible Ca^{2+} flows in these sites, we used Förster resonance

energy transfer (FRET)-based Ca^{2+} biosensors called Twitch. These sensors contain a minimal

Ca^{2+} binding moiety derived from the C-terminal domain of troponin C incorporated between

mCerulean3 and cpVenuscd (Thestrup et al., 2014). To target the sensors to ER-TGN MCS in

living cells, we introduced an amino acid sequence coding the N-terminal region of OSBP1 that

includes a PH and an FFAT motif (**Figure 3A**). As the N-terminal disordered domain of OSBP

seems to be crucial for active lipid transport in the ER-TGN MCS (Jamecna et al., 2019), we

generated sensors containing disordered domain (N-PH-FFAT) as well as sensor without that

domain (PH-FFAT) (**Supplementary S3A, S3B**). To obtain stable and equal protein expression

levels, we constructed stable HeLa cell lines expressing the MCS-Twitch sensors under a

doxycycline-inducible promoter. We confirmed the expression of the respective sensor after

doxycycline induction by immunofluorescence and live-cell imaging (**Supplementary S3B**). To

investigate the correct localization of the sensors at the TGN, cells were fixed and stained with

antibodies recognizing against GM130 or TGN46. The immunofluorescence microscopy data

confirmed that the MCS-Twitch sensors expressed in HeLa cells correctly localize

(**Supplementary S3C, S3D**).

The Twitch2b-MCS FRET sensor detects Ca^{2+} flows

To determine the range of Ca^{2+} signals detectable at the MCS at steady-state (non-treated

cells incubated at 37°C), we constructed four MCS-Twitch sensors with different Ca^{2+} affinities

(depicted in **Supplementary Tables 1, 2**). For quantitative FRET measurements of Ca^{2+}, we

calculated the FRET index value in the cell line expressing the respective sensor as an

approximation of the FRET/molecule (Grashoff et al., 2010; Kumar et al., 2016). The acquisition

of the normalized FRET index values was obtained by measuring FRET intensity, subtracting the

background noise and the bleed-through for the two fluorophores, and normalizing to FRET

acceptor intensity. Measuring FRET indexes from the MCS-Twitch sensors at steady revealed that

compared to MCS-Twitch9x, MCS-Twitch7x, MCS-Twitch8x, the Twitch2b sensor showed the

highest FRET index (**Supplementary S3E**).

To test if the Twitch2b sensor was responsive to Ca^{2+} perturbations in the cells, we

performed a series of control experiments. First, we determined the FRET values in HeLa cells

expressing the respective MCS-Twitch sensor by live-cell fluorescence microscopy in cells treated

with DMSO (control) or the Ca^{2+} chelating agent BAPTA-AM. As expected, chelating

intracellular Ca^{2+} ions with BAPTA-AM, showed a significant decrease in the FRET index of

MCS-Twitch2b (**Figure 3E, F**). The FRET indices of MCS-Twitch2b upon treating the cells with

BAPTA were similar to the baseline levels of the low Ca^{2+} affinity MCS-Twitch7x, 8x and 9x

sensors (**Figure C, D; Supplementary S3E**). Therefore, Twitch2b was selected for all further

experiments.

To quantify the range of free Ca^{2+} concentrations, we calibrated the Twitch2b sensor. To

calibrate Twitch2b in live cells, we used HeLa cells stably expressing CYTO-Twitch2b. We

applied a reciprocal dilution of buffers containing increasing ratios of Ca-EGTA/K2-EGTA

concentrations, and the free Ca^{2+} ion concentrations were calculated as described in Materials and

Methods (**Supplementary S4A, S4B**). The FRET indexes obtained from Twitch2b at different

ratios of Ca-EGTA/K2-EGTA in the calibration experiment allowed us to build a calibration curve

and which demonstrated a strong correlation between the FRET index value and the concentration

of free Ca^{2+} (calibration plot shown in **Figure 3B**). Thus, we developed a powerful tool to measure

Ca^{2+} levels in the ER-TGN-MCS. The calibration experiment allowed us to calculate the Ca^{2+}

levels at the MCS based on FRET indices.

To measure increased intracellular Ca^{2+} concentrations on the FRET signals from MCS-

Twitch2b sensors, we utilized active and passive means of increasing cytosolic Ca^{2+} levels.

Treatment of cells with ionomycin, that raises the intracellular Ca^{2+} level caused a significant

increase in the FRET index (**Figure 3G, H**). Several signaling pathways, including cell surface

receptors, are known to utilize Ca^{2+} ions as secondary messengers for downstream signaling

(Carafoli, 2002; Dickenson and Hill, 1994; Thillaiappan et al., 2017). Activation of Histamine

receptors (H1-receptor) at the plasma membrane causes the activation of PLC, which in turn

elevates intracellular Ca^{2+} through an IP3-dependent mechanism. To test if MCS sensor detects

these signals, we performed the FRET measurements in HeLa cells expressing PH-FFAT-

Twitch2b incubated with either DMSO (control) or after treatment with histamine. (**Figure 3I, J**).

The data clearly showed an increase in the FRET index– revealing a profound, physiological link

between MCS Ca^{2+} levels and the signaling receptor IP3.

Because we did not observe difference in the FRET index values between PH-FFAT-

Twitch and N-PH-FFAT-Twitch sensors (**Supplementary 3E, 3F, 3G, 3H**), we decided to

perform further experiments by using the PH-FFAT targeting motif.

The data indicates that release of Ca^{2+} caused by IP3R stimulation leads to an increase in

Ca^{2+} levels in MCS and our sensor precisely measures these changes.

TGN protein abundance and Ca^{2+} flux at MCS are coupled

Our former work has shown that TGN Ca^{2+} influx is necessary for the TGN exit of

secretory proteins (Crevenna et al., 2016; Deng et al., 2018; Kienzle et al., 2014; von Blume et

al., 2011). Therefore, we hypothesized that there must be a correlation between cargo influx into

the TGN and Ca^{2+} flow in the MCS. To test if Ca^{2+} in ER-TGN MCSs is influenced by protein

abundance, we treated HeLa cells expressing MCS-Twitch 2b with Cycloheximide (CHX), that

blocks protein synthesis. We incubated cells for 1, 2 and 4 hours and analyzed FRET signals.

Analyzing the FRET data showed a time-dependent decrease in the FRET index of cells treated

with CHX. To further demonstrate that this is correlated with cargo abundance in the TGN, we

incubated HeLa cells expressing MCS-Twitch 2b at 20°C to arrest secretory proteins in the TGN

(Ladinsky et al., 2002; Matlin and Simons, 1983). Notably, FRET values were significantly

decreased in cells incubated at 20°C (**Figure 4A, B**). Duration of the 20°C blocks for more than

1-hour did not affect the average FRET index values (**Figure 4C**). More importantly, incubation

of cells at 37°C after the 20°C-block resulted in a complete recovery of the FRET index (**Figure**

4B, C). To validate that the changes in the FRET indices were not due to other factors, cells were

incubated at 37°C and 20°C in the presence of ionomycin (**Figure 4E**). Independent of the

incubation temperature, ionomycin treatment led to a recovery of FRET indexes demonstrating

that the effects are specific to Ca^{2+}. These data supported that the Ca^{2+} flux in the MCS is

stimulated by the presence of newly synthesized protein. Furthermore, we showed that this is

specific to the cargo abundance in the TGN.

In the current study, we quantify the abundance of Ca^{2+} flux in the ER-TGN MCS, which

has remained unknown. We also demonstrate that cargo entering the TGN elicits a Ca^{2+} release in

the MCS that is required for sorting Cab45 clients into a TGN-derived carrier (**Figure 5A**).

Our previous work has shown that SPCA1, the only Ca^{2+} ATPase in the TGN, is required

for secretory cargo sorting in the TGN (von Blume et al., 2011). Interestingly, these studies also

exemplified that the Ca^{2+} binding protein Cab45 oligomerizes and forms a scaffold for collecting

its clients, including LyzC and COMP (Crevenna et al., 2016; von Blume et al., 2012).

Furthermore, sphingomyelin produced by SMS1 promotes SPCA activity and Cab45/client export

from the TGN (Deng et al., 2018). However, the source of Ca^{2+} that drives the process has

remained unknown (Ramazanov et al., 2021). In the current work, we demonstrate that the TGN

export COMP and LyzC, require the integrity of ER-TGN MCS and IP3R-dependent Ca^{2+} fluxes

in the MCS, suggesting that Ca^{2+} transfer between these organelles is the primary source for

SPCA1.

The autoregulation of the secretory pathway by cellular signaling remains mysterious. However,

central principles have been uncovered in recent years (Di Martino et al., 2019). The signaling

pathway that stimulates the Ca^{2+} release through the contact site remains unknown. Our work

elucidates that it is directly linked to IP3 production and activation of IP3R in close proximity to

ER-TGN contact sites. Although current work provides an understanding of upstream signaling

events that regulate of sorting of soluble proteins at TGN and depend on the activity of SPCA1,

the possibility that LyzC and COMP themselves could serve as a stimulus for the activation of

such pathways remains to be investigated.

Materials and methods

DNA techniques and plasmid construction

Restriction enzymes for molecular biology were obtained from New England Biolabs. PCRs were performed with a Phusion Polymerase (Thermofisher) and a Mastercycler Nexus (Eppendorf). All plasmids used in this study bear ampicillin resistance for selection in *Escherichia coli* and are listed in Table 1, where the transgenes and inserts are described. The DNA sequences encoding PH-FFAT/N-PH-FFAT domains of OSBP1 fused with Twitch sensors were integrated into donor plasmid of the transposon-based piggyBac system for stable transgene-expressing cell lines generation (Li et al., 2013). The piggyBac backbone vector (PB-T-PAF) and PB-RN and PBase were a gift of James Rini, University of Toronto, Ontario, Canada (Li et al., 2013). In brief, to generate PB-T-PAF-PHFFAT-Twitch constructs the PB-T-PAF vector was linearized with NheI and NotI-HF restriction enzymes (NEB). PH-FFAT/N-PH-FFAT sequences (5'-NheI/3'-AscI) were amplified by PCR from pLJM1-FLAG-GFP-OSBP plasmid (Addgene#134659). The sequences encoding Twitch (Twitch2b/Twitch7x/Twitch8x/Twitch9x) Calcium sensors were obtained from plasmids that have been generously provided from Oliver Griesbeck Lab and were amplified by adding corresponding restriction sites (5'-AscI/3'-NotI). All fragments were ligated in PB-T-PAF backbone. All cloning experiments were conducted using Phusion High-Fidelity Polymerase and T4 ligase (Thermo Fisher Scientific) according to the manufacturer's instructions. VAPA-Twitch2b and CYTO-Twitch2b constructs were generated by similar strategies. The sequences encoding Twitch2b for these constructs were amplified by adding corresponding restriction sites (5'-NheI/3'-AscI). The VAPA fragment was amplified from plasmid coding full length VAPA protein and was gifted from Pietro De Camilli with addition of AscI and NotI restriction sites. CYTO-Twitch2b construct was generated from VAPA-Twitch2b by replacing

VAPA sequence by short stop-codon containing fragment, annealed by using following sequences

(5'CGCGCCAGAGGAGTTTTAAGC3' and 5'GGCCGCTTAAAACTCCTCTGG3'). pLenti-

LyzC-EGFP for secretion assay lines was generated using gateway cloning reaction by amplifying

LyzC-EGFP from plpcx-LyzC-EGFP and cloning into pDONR221 using BP cloning reaction and

then subsequently using LR cloning reaction into the destination vector to generate the desired

construct. The correct sequence of all constructs was confirmed by DNA sequencing using the

SmartSeq Kit from Eurofins Genomics or KECK sequencing (Yale University).

Cell culture and generation of stable cell lines expressing Calcium sensor constructs.

Cell lines were maintained in DMEM media (Gibco) containing 10% FBS (Sigma

12306C-500ML) at 37°C and 5% CO_2. For transfection cells were plated in antibiotic-free media

24 hours before the procedure. DNA transfections were performed by using Lipofectamine 2000

reagent according to the manufacturer's protocol. After 8 hours the media was replaced. Transgene

expression was estimated at 48 hours after transfection. To generate cell lines stably expressing

the mentioned above transgenes HeLa lines were used. In brief, HeLa cells at 70% confluency

were transfected with PB-T-PAF (with corresponding transgene), PB-RN, and PBase (total DNA

1.5 µg; at ratio 8:1:1) using Lipofectamine 2000 in OptiMEM-I media. Cells were selected for 48

hours with 2 µg/ml puromycin dihydrochloride (Sigma-Aldrich) and for 7 days with 400 µg/ml

G418 disulfate salt (A1720-5G, Sigma-Aldrich). To induce transgene expression cells were

incubated in presence of doxycycline (J63805 Alfa Aesar, USA) (1ug/mL) for 24 hours. The

generated lines were sorted on BD FACS Aria to exclude resistant cells without transgene

expression.

HeLa cell lines stably expressing LyzC-EGFP were generated using lentiviral transduction

containing pLenti-LyzC-EGFP construct with followed Blasticidin 8 ug/mL (InvivoGen)

selection for 48 hours. siRNA transfections were performed using Lipofectamine RNAiMAX

according to standard protocol. VAPA/VAPB-dKO lines were generated by Pietro De Camilli Lab

and published previously (Dong et al., 2016).

381 RUSH cargo sorting assay and fluorescent microscopy

RUSH assay was performed as described previously (Boncompain et al., 2012; Deng et al.,

2018). Studied cell lines were cultured in 6-well plates (Cat#353046, Corning) on glass slides (Cat.

#72290-04, EMS) and transfected with RUSH-COMP-EGFP and RUSH-LyzC-EGFP constructs

using Lipofectamine 2000 according to standard protocol. At 24 hours post transfection, cells were

incubated with 40 µM d-Biotin (Sigma) ingrown media for different time points (20, 30, 40, 60,

and 90 min or without d-Biotin (control). For IF slides, cells were washed once with PBS, fixed in

4% PFA (Electron Microscopy Sciences) in PBS for 10 min, and mounted on 12mm coverslips

(Electron Microscopy Sciences) using ProLong Gold (Thermo Fisher Scientific). Nuclear

chromatin was stained by short incubation in 2.5uM DAPI (Biolegend) solution. Acquisition of

either GFP was performed by using a Delta Vision system by imaging z-stacks with a step size of

0.2 µm.

For quantification of vesicles, we empirically measured the sizes of objects between 4 and

20 pixels using the Analyze Particles function in ImageJ, which detects vesicular structures but

omits larger structures such as the Golgi. While small-fragmented and isolated Golgi structures

could be detected in error, such structures are rare. Furthermore, only vesicles of cells expressing

the RUSH construct were counted. The Fiji macro count_fixed_vesicles_V1.3 (M. Pakdel)

including the Particle Analyzer plug-in by Fiji was used to determine the number of vesicles (Deng

et al., 2018). Kruskal–Wallis one-way analysis of variance was used for comparisons in RUSH

experiments.

For immunostaining, cells were cultured in six wells on glass slides and fixed for 10 min

with 4% paraformaldehyde. After washing with PBS, cells were permeabilized for 5 min in 0.2%

Triton-X 100 and 0.5% SDS in 4% BSA. After washing with PBS, cells permeabilized with Triton-

X 100 were blocked with 4% BSA for 1 h. Cells were incubated with primary followed by

corresponding secondary antibody for 1 h at room temperature in blocking buffer in the dark.

Slides were washed three times with PBS after incubation with antibody. Glass slides were

mounted with ProLong Gold (Thermo Fisher Scientific). Antibodies to TGN46 (AHP500G,

Biorad) and GM130 (610822, BD) were used at dilution 1:200.

For FACS analysis cells were fixed with 4% PFA for 10 min. After fixation washed with

ice-cold DPBS and dissociated using trypsin-EDTA solution. Cells were washed with DPRS three

times by centrifugation at 200g for 5 min each. FACS analysis was performed using BD LSRII

machine. 10000 cells were analyzed for each sample. FACS data was analyzed using FlowJo

10.5.3 for Windows.

RUSH protein trafficking analysis

HeLa cells were seeded on Mattek dishes (p35gc-1.5-14-c, Mattek) and cultured at 37°C

with 5% CO_2. The next day or the day before imaging, cells were transfected by Lipofectamine

2000 (Invitrogen) with two plasmids expressing GalT-BFP (a trans-Golgi maker from James

Rothman lab) and RUSH COMP or RUSH Lysozyme C (see above for details of the plasmids).

After five to eight hours, cells were washed and incubated with DMEM with 10% FBS overnight.

Before imaging, cells were washed and briefly incubated with 37°C-warmed DMEM

supplemented with 10% FBS and HEPES buffer (Gibco, 21063029). Cells were then staged on the

microscope as described below.

Live-cell imaging was performed using a spinning disk confocal microscope CSUXfw-06p-01 (Yokogawa) on Nikon eclipse Ti2 (LWD NA=0.52) microscope stand with a motorized stage with stage top Piezo. sCMOS camera Photometrics Prim 95B and CFI Plan Apo Lambda 60x oil objective were used. Also, the Oko Lab temperature control system was set to 37°C and the fluorescence (405 nm and 488 nm) was induced by using an Agilent laser combiner. Images were acquired using Nikon Elements. Fluorescence images were taken every three minutes after adding biotin (see above for the methods for RUSH experiments). At each time point, the Nikon Elements stitched 6x5 fields to generate relatively large field images that could accommodate at least 10 distinguished secretion events through 90 or 120 minutes.

The time-course images generated above were imported to Fiji (ImageJ). Images containing individual cells with distinguishable Golgi and RUSH signals were cropped as stored separately. The Golgi masks were generated by using GalT-BFP signals (405 nm excitation) and ImageJ Auto-threshold and were imported as ROIs. The Golgi ROIs were used to measure mean RUSH intensity in the Golgi at each time point. The mean RUSH intensities were subtracted by background signals. Then, the maximum mean value of RUSH during the time course of the single-cell images was used to divide the RUSH intensities, thus normalizing the maximal RUSH intensities in the Golgi marker set to 1. The time course images for each single cell were analyzed separately and combined to generate the mean RUSH intensity in each time point and its standard deviation, as described in the Figures. Three independently cultured cells were analyzed, and their images were combined to generate the final data.

Secretion assay of LyzC-EGFP

For the secretion assay, 5×10^5 cells stably expressing LyzC-EGFP were plated in a 6 well

plate. After 24 hours cells were pretreated with 70μM 2-APB and DMSO. After that incubated in

grown medium containing 2-APB and DMSO for 1 hour. Cells and media were collected

separately. Cells were lysed using RIPA buffer. Collected grown media was incubated overnight

with GFP-trap beads at 4°C. On the next day the beads were washed 4 times with DPBS and then

protein was eluted from the beads by boiling it in 2X Laemmli SDS sample buffer (Biorad). Cell

lysates and IP fractions were analyzed by western blotting.

Live cell imaging and FRET analysis

For live cell imaging, DMEM without phenol red containing 4.5 mg ml – glucose, 25 mM

HEPES, 2 mM glutamine (Life Technologies) supplemented with 10 % FBS was used. Cells were

seeded on Glass bottom dishes (D35-14-1.5-N, Cellvis) on density 5×10^4 per dish. Next day to

induce transgene expression doxycycline at final concentration 1ug/mL was added. After 24 hours

of doxycycline was cancelled and cells were incubated in imaging media for additional 24 hours.

ImageJ (National Institutes of Health) was used for basic image processing. All analyses were

done using custom-written software (MATLAB R2014a; MathWorks). To manipulate

intracellular Ca^{2+} levels as well as inducing Ca^{2+} flux following drugs at corresponding

concentrations were used: calcium ionomycin (I3909-1ML, Sigma) at 1μM; 2-APB (100065-

100MG, Millipore) at 70μM, Histamine (H7125-1G, Sigma) 1 μM solution, BAPTA-AM

(126150-97-8, Millipore) at 25 μM final concentration. To inhibit protein synthesis, cells were

treated with cycloheximide at concentration 10ug/mL (DSC81040-5, Dot Scientific Inc).

FRET sensor calibration experiment.

Free Ca^{2+} calibration solutions were made by using Calcium Calibration Buffer Kit #1

(Cat. No. C3008MP, Biotium) according to standard protocol. To perform FRET calibration

469 experiment HeLa cell line stably expressing CYTO-Twitch2b was used. $1x10^5$ cells were plated

470 into Glass bottom dishes (D35-14-1.5-N, Cellvis) and doxycycline was added to the media at final

471 concentration 1 ug/mL. After 24 hours for each plate growth media was washed twice with 10mM

472 EGTA, 100mM KCl, 10mM MOPS (pH7.2). To chelate residual Ca^{2+} cells were incubated in

473 10mM EGTA, 100mM KCl, 10mM MOPS (pH7.2) with 1 μM ionomycin for 20 min at RT. 2ml

474 of stock solutions with free Ca^{2+} concentrations ranging from 0 μM to 39 μ μM were added to

475 cells and FRET indexes were measured for each condition.

476 **FRET imaging and analysis**

477  These analyses were done essentially as previously described (Kumar et al., 2016). High

478 resolution live FRET imaging was performed on Nikon Eclipse Ti widefiled microscope equipped

479 with a cooled charged-coupled device Cool SNAP HQ2 camera, using a ×100, 1.49 NA oil

480 objective at 37 °C. Images were acquired using Micromanager software. Three sequential images

481 with 500 ms exposure time were acquired with the following filter combinations: donor (Teal)

482 channel with 460/20 (excitation filter-ex), T455lp (dichroic mirror-di) and 500/22 (emission filter-

483 em); FRET channel with 460/20 (ex), T455lp (di) and 535/30 (em); and acceptor (Venus) channel

484 with 492/18 (ex), T515lp (di) and 535/30 (em) filter combinations. All filters and dichroic were

485 purchased from Chroma Technology. For data analysis, donor leakage was determined from HeLa

486 cells transiently transfected with Vinculin-Teal, whereas acceptor cross excitation was obtained

487 from Vinculin-Venus transfected cells. For all the calculations, respective background subtraction,

488 illumination gradient, and pixel shift correction were performed followed by three-point

489 smoothening. The slope of pixel-wise donor or acceptor channel intensity versus FRET channel

490 intensity gives leakage (x) or cross-excitation (y) fraction, respectively. FRET map and pixel-wise

FRET index for the sensors were determined from FRETindex =

[FRETchannel−x(Donorchannel)− y(Acceptorchannel)]/[Acceptorchannel]

ImageJ (National Institutes of Health) was used for basic image thresholding. Mean FRET

index per cell was calculated for each region within the mask. Student's t test was performed

between the two groups to calculate statistical significance and p-value. At least $P < 0.05$ was

considered significant. All analyses were done using custom-written software (MATLABR2020b;

MathWorks) (Kumar et al., 2016).

siRNA delivery and western blotting

Knockdown of VAPA and VAPB proteins was performed by using siRNA (VAPA: 5'-

AACTAATGGAAGAGTGTAAAA-3'; VAPB: 5'-AAGAAGGTTATGGAAGAATGT-

3')(Wakana et al., 2021). Non-targeting siRNA was purchased from Qiagen (Catalog No. –

1027281) and used as reference negative control. Knockdown of VAPA and VAPB proteins was

achieved by combined transfection (siVAPA and siVAPB) by using LipofectRNAiMAX

according to standard protocol. Knockdown efficiency was confirmed by Western blotting of cell

lysates in radioimmunoprecipitation assay (RIPA) buffer (50 mM Tris-HCl, pH 7.4 [American

Bioanalytical], 150 mM NaCl [American Bioanalytical], 1% Triton X100 [American

Bioanalytical], 1% sodium deoxycholate [Sigma-Aldrich], and 0.1% SDS [American

Bioanalytical] in milliQ water); protease and phosphatase inhibitor (Thermo Fisher Scientific) was

added just before extraction. Cell lysate was resolved using SDS-PAGE and transferred to the

nitrocellulose membrane (Biorad Laboratories) using a transfer system (Trans-Blot Turbo; Bio-

Rad Laboratories). The membrane was blocked using 5% skimmed milk (American Bioanalytical)

in TBS with 0.1% Tween 20 (TBST) for 1 hour and incubated with the following primary

antibodies diluted in 5% milk with TBST overnight at 4°C: Anti-β-actin (1:5000, A5441-.2ML

Sigma); Anti-VAPA (1:1000, SAB1402460-100G, Sigma); Anti-hVAPB (1:2000, MAB58551

R&D systems); anti-GFP (1:1000, 11814460001, Roche). Membrane was washed 3 times with 5%

milk TBST and incubated with HRP-conjugated secondary antibodies (1:5000, 32230/32260;

Invitrogen). Data was visualized using chemiluminescence detection on ChemiDoc Touch (Bio-

Rad Laboratories)**.**

Graphical data and image design

Graphs were plotted in GraphPad Prism version 9.2.0 for Mac, GraphPad Software, San Diego,

California USA. Images were compiled using Adobe Illustrator 2022 (Adobe Inc. (2022). Adobe

Illustrator. Retrieved from https://adobe.com/products/illustrator). Schemes were designed using

Biorender software (biorender.com).

Acknowledgments

We thank Dr. Pietro de Camilli for providing VAPA/VAPB-dKO cell lines and the Burd lab for discussions and support. Julia von Blume is funded by a Yale-start up grant and by the National Institute of General Medical Sciences of the NIH under the award number GM134083-01, an Administrative Supplement 3R01GM134083-03S1 and a Project and Feasibility award from Yale Diabetic Research Center (GR112420).

Author Contributions

B.R.Ramazanov and J.von Blume designed the experiments. B.R.Ramazanov carried out the experiments and analyzed the data. A.Kumar and M.Schwartz provided support in FRET data analysis and composed the software code for data analysis. A. Parchure and Y.Kim performed and analyzed secretion assays. O.Griesbeck provided Twitch sensors encoding plasmids. M.Schwartz provided the equipment. B.R.Ramazanov and J.von Blume wrote the paper.

Declaration of Interest

The authors declare no competing financial interests.

Abbreviations

546   ER - Endoplasmic reticulum

547   FRET - Fluorescence resonance energy transfer

548   nm - nanometer

549   TGN - Trans Golgi Network

References

Barlowe, C.K., and E.A. Miller. 2013. Secretory protein biogenesis and traffic in the early secretory pathway. *Genetics*. 193:383-410.

Berridge, M.J., M.D. Bootman, and H.L. Roderick. 2003. Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol*. 4:517-529.

Boncompain, G., S. Divoux, N. Gareil, H. de Forges, A. Lescure, L. Latreche, V. Mercanti, F. Jollivet, G. Raposo, and F. Perez. 2012. Synchronization of secretory protein traffic in populations of cells. *Nat Methods*. 9:493-U120.

Carafoli, E. 2002. Calcium signaling: A tale for all seasons. *Proceedings of the National Academy of Sciences*. 99:1115-1122.

Crevenna, A.H., B. Blank, A. Maiser, D. Emin, J. Prescher, G. Beck, C. Kienzle, K. Bartnik, B. Habermann, M. Pakdel, H. Leonhardt, D.C. Lamb, and J. von Blume. 2016. Secretory cargo sorting by Ca2+-dependent Cab45 oligomerization at the trans-Golgi network. *J Cell Biol*. 213:305-314.

De Matteis, M.A., and A. Luini. 2008. Exiting the Golgi complex. *Nat Rev Mol Cell Biol*. 9:273-284.

Deng, Y., M. Pakdel, B. Blank, E.L. Sundberg, C.G. Burd, and J. von Blume. 2018. Activity of the SPCA1 Calcium Pump Couples Sphingomyelin Synthesis to Sorting of Secretory Proteins in the Trans-Golgi Network. *Dev Cell*. 47:464-478 e468.

Deng, Y., F.E. Rivera-Molina, D.K. Toomre, and C.G. Burd. 2016. Sphingomyelin is sorted at the trans Golgi network into a distinct class of secretory vesicle. *Proc Natl Acad Sci U S A*. 113:6677-6682.

Di Martino, R., L. Sticco, and A. Luini. 2019. Regulation of cargo export and sorting at the trans-Golgi network. *FEBS Lett*. 593:2306-2318.

Dickenson, J.M., and S.J. Hill. 1994. Selective potentiation of histamine H1-receptor stimulated calcium responses by 1,4-dithiothreitol in DDT1MF-2 cells. *Biochem Pharmacol*. 48:1721-1728.

Dong, R., Y. Saheki, S. Swarup, L. Lucast, J.W. Harper, and P. De Camilli. 2016. Endosome-ER Contacts Control Actin Nucleation and Retromer Function through VAP-Dependent Regulation of PI4P. *Cell*. 166:408-423.

Dunlop, M.H., A.M. Ernst, L.K. Schroeder, D.K. Toomre, G. Lavieu, and J.E. Rothman. 2017. Land-locked mammalian Golgi reveals cargo transport between stable cisternae. *Nat Commun*. 8.

Dunphy, W.G., and J.E. Rothman. 1985. Compartmental organization of the Golgi stack. *Cell*. 42:13-21.

Ford, C., A. Parchure, J. von Blume, and C.G. Burd. 2021. Cargo sorting at the trans-Golgi network at a glance. *J Cell Sci*. 134.

Gillon, A.D., C.F. Latham, and E.A. Miller. 2012. Vesicle-mediated ER export of proteins and lipids. *Biochim Biophys Acta*. 1821:1040-1049.

Glick, B.S., T. Elston, and G. Oster. 1997. A cisternal maturation mechanism can explain the asymmetry of the Golgi stack. *FEBS Lett*. 414:177-181.

Glick, B.S., and A. Luini. 2011. Models for Golgi traffic: a critical assessment. *Cold Spring Harb Perspect Biol*. 3:a005215.

Glick, B.S., and A. Nakano. 2009. Membrane Traffic Within the Golgi Apparatus. *Annual Review*
of Cell and Developmental Biology. 25:113-132.
Grashoff, C., B.D. Hoffman, M.D. Brenner, R. Zhou, M. Parsons, M.T. Yang, M.A. McLean, S.G.
Sligar, C.S. Chen, T. Ha, and M.A. Schwartz. 2010. Measuring mechanical tension across
vinculin reveals regulation of focal adhesion dynamics. *Nature*. 466:263-266.
Guo, Y.S., D.W. Sirkis, and R. Schekman. 2014. Protein Sorting at the trans-Golgi Network. *Annu*
Rev Cell Dev Bi. 30:169-206.
Kawano, M., K. Kumagai, M. Nishijima, and K. Hanada. 2006. Efficient trafficking of ceramide
from the endoplasmic reticulum to the Golgi apparatus requires a VAMP-associated
protein-interacting FFAT motif of CERT. *J Biol Chem*. 281:30279-30288.
Kelly, R.B. 1985. Pathways of Protein Secretion in Eukaryotes. *Science*. 230:25-32.
Kienzle, C., N. Basnet, A.H. Crevenna, G. Beck, B. Habermann, N. Mizuno, and J. von Blume.
2014. Cofilin recruits F-actin to SPCA1 and promotes Ca2+-mediated secretory cargo
sorting. *J Cell Biol*. 206:635-654.
Kienzle, C., and J. von Blume. 2014. Secretory cargo sorting at the trans-Golgi network. *Trends*
Cell Biol. 24:584-593.
Kumagai, K., and K. Hanada. 2019. Structure, functions and regulation of CERT, a lipid-transfer
protein for the delivery of ceramide at the ER-Golgi membrane contact sites. *Febs*
Letters. 593:2366-2377.
Kumar, A., M. Ouyang, K. Van den Dries, E.J. McGhee, K. Tanaka, M.D. Anderson, A. Groisman,
B.T. Goult, K.I. Anderson, and M.A. Schwartz. 2016. Talin tension sensor reveals novel
features of focal adhesion force transmission and mechanosensitivity. *J Cell Biol*.
213:371-383.
Ladinsky, M.S., C.C. Wu, S. McIntosh, J.R. McIntosh, and K.E. Howell. 2002. Structure of the
Golgi and distribution of reporter molecules at 20 degrees C reveals the complexity of
the exit compartments. *Mol Biol Cell*. 13:2810-2825.
Lebreton, S., S. Paladino, D. Liu, M. Nitti, J. von Blume, P. Pinton, and C. Zurzolo. 2021. Calcium
levels in the Golgi complex regulate clustering and apical sorting of GPI-APs in polarized
epithelial cells. *Proc Natl Acad Sci U S A*. 118.
Lehto, M., and V.M. Olkkonen. 2003. The OSBP-related proteins: a novel protein family involved
in vesicle transport, cellular lipid metabolism, and cell signalling. *Biochim Biophys Acta*.
1631:1-11.
Lev, S. 2010. Non-vesicular lipid transport by lipid-transfer proteins and beyond. *Nat Rev Mol*
Cell Biol. 11:739-750.
Li, Z.J., I.P. Michael, D.X. Zhou, A. Nagy, and J.M. Rini. 2013. Simple piggyBac transposon-based
mammalian cell expression system for inducible protein production. *P Natl Acad Sci*
USA. 110:5004-5009.
Lissandron, V., P. Podini, P. Pizzo, and T. Pozzan. 2010. Unique characteristics of Ca2+
homeostasis of the trans-Golgi compartment. *Proc Natl Acad Sci U S A*. 107:9198-9203.
Lujan, P., and F. Campelo. 2021. Should I stay or should I go? Golgi membrane spatial
organization for protein sorting and retention. *Arch Biochem Biophys*. 707.
Maruyama, T., T. Kanaji, S. Nakade, T. Kanno, and K. Mikoshiba. 1997. 2APB, 2-
aminoethoxydiphenyl borate, a membrane-penetrable modulator of Ins(1,4,5)P3-
induced Ca2+ release. *J Biochem*. 122:498-505.

Masone, M.C., V. Morra, and R. Venditti. 2019. Illuminating the membrane contact sites
between the endoplasmic reticulum and the trans-Golgi network. *FEBS Lett*. 593:3135-
3148.
Matlin, K.S., and K. Simons. 1983. Reduced temperature prevents transfer of a membrane
glycoprotein to the cell surface but does not prevent terminal glycosylation. *Cell*.
34:233-243.
Mesmin, B., J. Bigay, J. Polidori, D. Jamecna, S. Lacas-Gervais, and B. Antonny. 2017. Sterol
transfer, PI4P consumption, and control of membrane lipid order by endogenous OSBP.
Embo J. 36:3156-3174.
Mesmin, B., J. Bigay, J.M. von Filseck, S. Lacas-Gervais, G. Drin, and B. Antonny. 2013. A Four-
Step Cycle Driven by PI(4)P Hydrolysis Directs Sterol/PI(4)P Exchange by the ER-Golgi
Tether OSBP. *Cell*. 155:830-843.
Missiaen, L., L. Dode, J. Vanoevelen, L. Raeymaekers, and F. Wuytack. 2007. Calcium in the Golgi
apparatus. *Cell Calcium*. 41:405-416.
Mostov, K.E., and M.H. Cardone. 1995. Regulation of protein traffic in polarized epithelial cells.
Bioessays. 17:129-138.
Nakano, A., and A. Luini. 2010. Passage through the Golgi. *Curr Opin Cell Biol*. 22:471-478.
Pakdel, M., and J. von Blume. 2018. Exploring new routes for secretory protein export from the
trans-Golgi network. *Mol Biol Cell*. 29:235-240.
Pfeffer, S.R., and J.E. Rothman. 1987. Biosynthetic protein transport and sorting by the
endoplasmic reticulum and Golgi. *Annu Rev Biochem*. 56:829-852.
Phillips, M.J., and G.K. Voeltz. 2016. Structure and function of ER membrane contact sites with
other organelles. *Nat Rev Mol Cell Biol*. 17:69-82.
Pizzo, P., V. Lissandron, P. Capitanio, and T. Pozzan. 2011. Ca2+ signalling in the Golgi
apparatus. *Cell Calcium*. 50:184-192.
Pizzo, P., V. Lissandron, and T. Pozzan. 2010. The trans-golgi compartment: A new distinct
intracellular Ca store. *Commun Integr Biol*. 3:462-464.
Ramazanov, B.R., M.L. Tran, and J. von Blume. 2021. Sending out molecules from the TGN. *Curr
Opin Cell Biol*. 71:55-62.
Rizzuto, R., M. Brini, M. Murgia, and T. Pozzan. 1993. Microdomains with High Ca2+ Close to
Ip(3)-Sensitive Channels That Are Sensed by Neighboring Mitochondria. *Science*.
262:744-747.
Rizzuto, R., P. Pinton, W. Carrington, F.S. Fay, K.E. Fogarty, L.M. Lifshitz, R.A. Tuft, and T. Pozzan.
1998. Close contacts with the endoplasmic reticulum as determinants of mitochondrial
Ca2+ responses. *Science*. 280:1763-1766.
Scherer, P.E., G.Z. Lederkremer, S. Williams, M. Fogliano, G. Baldini, and H.F. Lodish. 1996.
Cab45, a novel (Ca2+)-binding protein localized to the Golgi lumen. *J Cell Biol*. 133:257-
268.
Sepulveda, M.R., D. Marcos, M. Berrocal, L. Raeymaekers, A.M. Mata, and F. Wuytack. 2008.
Activity and localization of the secretory pathway Ca2+-ATPase isoform 1 (SPCA1) in
different areas of the mouse brain during postnatal development. *Mol Cell Neurosci*.
38:461-473.
Stalder, D., and D.C. Gershlick. 2020. Direct trafficking pathways from the Golgi apparatus to
the plasma membrane. *Semin Cell Dev Biol*. 107:112-125.

Tang, B.L. 2001. Protein trafficking mechanisms associated with neurite outgrowth and
polarized sorting in neurons. *J Neurochem*. 79:923-930.
Thillaiappan, N.B., A.P. Chavda, S.C. Tovey, D.L. Prole, and C.W. Taylor. 2017. Ca(2+) signals
initiate at immobile IP3 receptors adjacent to ER-plasma membrane junctions. *Nat*
Commun. 8:1505.
Uhlen, M., L. Fagerberg, B.M. Hallstrom, C. Lindskog, P. Oksvold, A. Mardinoglu, A. Sivertsson, C.
Kampf, E. Sjostedt, A. Asplund, I. Olsson, K. Edlund, E. Lundberg, S. Navani, C.A.
Szigyarto, J. Odeberg, D. Djureinovic, J.O. Takanen, S. Hober, T. Alm, P.H. Edqvist, H.
Berling, H. Tegel, J. Mulder, J. Rockberg, P. Nilsson, J.M. Schwenk, M. Hamsten, K. von
Feilitzen, M. Forsberg, L. Persson, F. Johansson, M. Zwahlen, G. von Heijne, J. Nielsen,
and F. Ponten. 2015. Proteomics. Tissue-based map of the human proteome. *Science*.
347:1260419.
Venditti, R., M.C. Masone, and M.A. De Matteis. 2020. ER-Golgi membrane contact sites.
Biochem Soc Trans. 48:187-197.
Venditti, R., M.C. Masone, L.R. Rega, G. Di Tullio, M. Santoro, E. Polishchuk, I.C. Serrano, V.M.
Olkkonen, A. Harada, D.L. Medina, R. La Montagna, and M.A. De Matteis. 2019. The
activity of Sac1 across ER-TGN contact sites requires the four-phosphate-adaptor-
protein-1. *J Cell Biol*. 218:783-797.
von Blume, J., A.M. Alleaume, G. Cantero-Recasens, A. Curwin, A. Carreras-Sureda, T.
Zimmermann, J. van Galen, Y. Wakana, M.A. Valverde, and V. Malhotra. 2011.
ADF/cofilin regulates secretory cargo sorting at the TGN via the Ca2+ ATPase SPCA1. *Dev*
Cell. 20:652-662.
von Blume, J., A.M. Alleaume, C. Kienzle, A. Carreras-Sureda, M. Valverde, and V. Malhotra.
2012. Cab45 is required for Ca(2+)-dependent secretory cargo sorting at the trans-Golgi
network. *J Cell Biol*. 199:1057-1066.
Wakana, Y., K. Hayashi, T. Nemoto, C. Watanabe, M. Taoka, J. Angulo-Capel, M.F. Garcia-Parajo,
H. Kumata, T. Umemura, H. Inoue, K. Arasaki, F. Campelo, and M. Tagaya. 2021. The ER
cholesterol sensor SCAP promotes CARTS biogenesis at ER-Golgi membrane contact
sites. *J Cell Biol*. 220.
Wong, A.K.C., P. Capitanio, V. Lissandron, M. Bortolozzi, T. Pozzan, and P. Pizzo. 2013.
Heterogeneity of Ca-2 handling among and within Golgi compartments. *J Mol Cell Biol*.
5:266-276.
Zanetti, G., S. Prinz, S. Daum, A. Meister, R. Schekman, K. Bacia, and J.A. Briggs. 2013. The
structure of the COPII transport-vesicle coat assembled on membranes. *Elife*. 2:e00951.
716

Main Figures

Main Figure legends

Figure 1. Inhibition of IP3R activity delays TGN export of Cab45 clients
(A) Schematic representation of the Retention Using Streptavidin Hooks (RUSH) assay with fluorescent tagged client molecules. (B) Representative immunofluorescence images of the RUSH experiments showing COMP-EGFP transport in HeLa lines treated with DMSO and 70 nM 2-APB. HeLa cells were transfected with KDEL-IRES-SBP-COMP-EGFP and fixed at 0, 20, 40, and 60 min after the addition of biotin. Z-stack images (d = 0.2 µm) were analyzed. The arrowheads indicate cytoplasmic vesicles. Scale bars, 10 µm. (C) The numbers of COMP budding vesicles from RUSH experiments with KDEL-IRES-SBP-COMP-EGFP in HeLa lines treated with DMSO and 2-APB were quantified. The cytoplasmic vesicles were counted at each time point by analyzing z-stack images (d = 0.2 µm). Scatter dot plot represents the means ± SD of at least three independent experiments (n > 30 cells per condition). Statistical test, Kruskal–Wallis. (D) Plot representing normalized average fluorescence intensity of COMP-EGFP in cells by FACS at 0, 60 and 120 minutes of RUSH experiments. (E) Plot representing normalized fluorescence intensity of LyzC-EGFP within TGN (ROI was defined by GALNT1 area) in cells treated with DMSO and 2-APB. (F) Western blot showing LyzC-GFP in Hela cells treated with DMSO and 2-APB in cell lysates and in secreted medium (top) and β-actin as a loading control.

Figure 2. Depletion of VAPA and VAPB delays TGN export of Cab45 clients
(A) Representative immunofluorescence images of RUSH experiments showing COMP-GFP transport in HeLa lines treated with siRNA against VAPA and VAPB proteins. Cells were transfected with KDEL-IRES-SBP-COMP-EGFP and fixed at 0, 20, 40, and 60 min after the addition of biotin. Z-stack images (d = 0.2 µm) were analyzed. Scale bars, 10 µm. (B) The numbers of COMP budding vesicles were quantified. The cytoplasmic vesicles were counted at each time point by analyzing z-stack images (d = 0.2 µm). Scatter dot plot represents the means ± SD of at least three independent experiments (n > 30 cells per condition). Statistical test, Kruskal–Wallis. (C) Plot representing normalized average fluorescence intensity of COMP-EGFP in cells by FACS at 0, 60 and 120 minutes of RUSH experiments in HeLa WT and VAPA/VAPB-dKO lines (D) Plot representing normalized fluorescence intensity of LyzC-EGFP within TGN (ROI was defined by Galnt1 area) in cells treated with non-targeting (control) siRNA and siRNA targeting VAPA and VAPB. (E) Expression of VAPA and VAPB proteins in Hela WT line transfected with control (non-targeting) siRNA, Hela line transfected with siVAPA/siVAPB and Hela VAPA/VAPB-dKO line. β-actin was used as a loading control.

Figure 3. MCS targeting FRET sensor responses to changes of Ca^{2+} concentration at ER-TGN MCS.
(A) Schematic representation of OSBP and Twitch2b Calcium Sensor targeting Endoplasmic Reticulum-Golgi membrane contact sites. PH domain of OSBP binds to PI4P on the outer leaflet of the TGN membrane and FFAT domain interacts with VAPA on the ER membrane. (B) Calibration curve for Twitch2b sensor performed used buffer with increasing concentration of free Ca^{2+} ions demonstrating correlation between FRET indices and free Ca^{2+}. (See Materials and Methods for a detailed description). (C) Pseudocolor map of FRET index and dot plot (D) representing FRET indices for PH-FFAT-Twitch2b and PH-FFAT-Twitch9x within Golgi ROI of

live cells and at steady state (***p<0.05). (E) Pseudocolor map of FRET index and dot plot (F)
representing FRET indices for PH-FFAT-Twitch2b at steady state and treated with 25uM BAPTA-
AM for 20 min (***p<0.05). (G) Pseudocolor map of FRET index and dot plot (H) representing
FRET indices for PH-FFAT-Twitch2b at steady state and treated with 1uM ionomycin for 20 min
(***p<0.05). (I) Pseudocolor map of FRET index and dot plot (J) representing FRET indices for
PH-FFAT-Twitch2b at steady state and treated with 1uM Histamine for 2 min (***p<0.05).
Pseudocolor bar FRET index value in range 0-0.4.
770
Figure 4. Ca^{2+} flux at ER-TGN MCS is coupled to protein trafficking
(A) Schematic representation of experiment and effect on protein trafficking in live cells during
20°C block and cycloheximide (CHX) treatment. (B) Pseudocolor maps of FRET index for PH-
FFAT-Twitch2b within Golgi ROI of live cells at described above conditions. Pseudocolor bar
FRET index value in range 0-0.4. (C) Dot plot representing FRET index for PH-FFAT-Twitch2b
at steady state, after 20°C block for 1 and 2 hours and after 10 min recovery at 37°C (***p<0.05).
(D) Dot plot representing FRET index for PH-FFAT-Twitch2b at steady state, treated with CHX
for 1, 2, 4 hours and after 1h 20°C block. (***p<0.05), (E) Dot plot representing normalized FRET
index for PH-FFAT-Twitch2b at steady state, after 1h 20°C block and effect of ionomycin at 37°C
and 20°C conditions (***p<0.05).
781
782 **Figure 5A. The model depicting the role of contact sites and IP3R-dependent release of**
783 **Calcium ions for SPCA1 dependent sorting at TGN.**
784
785 # Supplementary data
786
787 **Supplementary Figure legends**
788
Supplementary 1. RUSH experiments using Cab45 clients demonstrate that delay in protein
trafficking takes place at TGN
(A) Representative histograms plotting fluorescence intensity distribution of COMP-GFP in cells
at 30, 60 and 120 minutes (shown in red) during RUSH experiment compared to timepoint "0"
(shown in green overlay) in HeLa WT cells line treated with DMSO (upper row), HeLa WT cells
treated with 2-APB (middle row), and HeLa VAPA/VAPB-dKO line (bottom row). 10.000 cells
analyzed for each condition. (B) Graph depicting mean normalized fluorescence intensity (F.I.) of
RUSH LyzC-EGFP in BFP-GALNT1 (Golgi marker) region of interest (ROI) from and after the
time when the maximal LyzC-EGFP F.I. is reached in the ROI (peak Golgi F.I.). The decrease in
LyzC-EGFP F.I. represents the Golgi exit in HeLa line treated with DMSO (black) versus 2-APB
treated (red). At least 10 cells from independent experiments for each condition were analyzed
with nonlinear regression (exponential function) indicated in the plot using GraphPad Prism9 built-
in regression. (C) Graph depicting mean normalized fluorescence intensity (F.I.) of RUSH LyzC-
EGFP in BFP-GALNT1 (Golgi marker) region of interest (ROI) from and after the time when the
maximal LyzC-EGFP F.I. is reached in the ROI (peak Golgi F.I.). The decrease in LyzC-EGFP
F.I. represents the Golgi exit in HeLa line transfected with control siRNA (black) versus Hela line
transfected with siRNA to VAPA and VAPB (red). At least 10 cells from independent experiments
for each condition were analyzed with nonlinear regression (exponential function) indicated in the
plot using GraphPad Prism9 built-in regression.

Supplementary 2. VAPA/VAPB dKO HeLa lines demonstrate delay in TGN export of Cab45 clients in RUSH experiments

(A) Representative immunofluorescence images of RUSH experiments showing LyzC-GFP transport in HeLa WT and VAPA/B-dKO cell lines. Cells were transfected with KDEL-IRES-SBP-LyzC-EGFP and fixed at 0, 20, 40, and 60 min after the addition of biotin. Z-stack images (d = 0.2 μm) were analyzed. The arrowheads indicate cytoplasmic vesicles. Scale bars, 10 μm. (B) Representative immunofluorescence images of RUSH experiments showing COMP-GFP transport in HeLa WT and VAPA/B-dKO cell lines. Cells were transfected with KDEL-IRES-SBP-COMP-EGFP and fixed at 0, 20, 40, and 60 min after the addition of biotin. Z-stack images (d = 0.2 μm) were analyzed. The arrowheads indicate cytoplasmic vesicles. Scale bars, 10 μm. (C) The numbers of LyzC budding vesicles were quantified. The cytoplasmic vesicles were counted at each time point by analyzing z-stack images (d = 0.2 μm). Scatter dot plot represents the means ± SD of at least three independent experiments (n > 30 cells per condition). Statistical test, Kruskal–Wallis. (D) The numbers of COMP budding vesicles were quantified. The cytoplasmic vesicles were counted at each time point by analyzing z-stack images (d = 0.2 μm). Scatter dot plot represents the means ± SD of at least three independent experiments (n > 30 cells per condition). Statistical test, Kruskal–Wallis.

Supplementary 3. MCS targeting FRET sensors localize with TGN markers.

(A) Schematic representation of the domain structure of OSBP (top construct) and Twitch sensors fused proteins targeting ER-Golgi contact sites (N-PH-FFAT-Twitch and PH-FFAT-Twitch) as well as sensors targeting ER (fused with VAPA protein) and cytoplasmic version (two bottom constructs) used in this study. (B) IF images of HeLa lines stably expressing Twitch2b sensor targeting cytoplasm, ER and ER-TGN contact sites. (C) IF images of HeLa line expressing Twitch2b sensor targeting ER-TGN contact sites and its localization compared to TGN46 (TGN marker). (D) IF images of HeLa line expressing Twitch2b sensor targeting ER-TGN contact sites and its localization compared to GM130 (cis Golgi marker). (E) Dot plot representing FRET index for PH-FFAT-Twitch2b and PH-FFAT-Twitch7x, PH-FFAT-Twitch8x and PH-FFAT-Twitch9x within Golgi ROI of live cells and at steady state (***p<0.05); (F) Dot plot representing FRET index for PH-FFAT-Twitch2b and N-PH-FFAT-Twotch2b in cells at steady state. (G) Dot plot representing FRET index for N-PH-FFAT-Twitch2b cells at steady state and treated with 1uM ionomycin for 20 min (***p<0.05). (H) Dot plot representing FRET index for N-PH-FFAT-Twitch2b cells at steady state and treated with 1uM Histamine for 2 min(***p<0.05).

Supplementary 4. Calibration procedures of Twitch2b sensor and fluorescence channels for FRET experiments

(A) Schematic representation of the calibration experiment of Twich2b calcium sensor using Ca-EGTA buffers in live cells (Please see Materials and Methods section for detailed description). (B) Pseudocolor heatmap of FRET index for CYTO-Twitch2b during calibration experiment. Pseudocolor bar FRET index value in range 0-0.4. (C) Calibration of fluorescence channels for Twitch sensor using HeLa lines expressing PH-FFAT-mCerulean3 (only FRET donor), HeLa lines expressing PH-FFAT-cpVenus (only FRET acceptor)) and PH-FFAT-Twitch2b (FRET pair with calcium binding domain) sensor.

854 **Table legends**

856 Supplementary Table1. Twitch sensors and their affinity to Ca^{2+} ions.

FRET Sensor	Calcium affinity of TnC (KD)
Twitch 2b	200 nM
Twitch 7x	51.5 uM
Twitch 8x	139 uM
Twitch 9x	174 uM

860 Supplementary Table2. Twitch sensors and aminoacidic sequences of calcium binding domains.

FRET Sensor	Amino acid sequence of Calcium binding domain of Twitch sensors
Twitch 2b	MQVADASEEELSECFRIFDFDGN_GFIDREEFGDIIRLTGE QLTDEDVDEIFGDSDTDKN_GRIDFDEFLKMVENVQPIYP
Twitch 7x	MQVADASEEELSECFRIFDFDGNDGFIDREEFGDIIRLTGE QLTDEDVDEIFGDSDTDKN_GRIDFDEFLKMVENVQPIYP
Twitch 8x	MQVADASEEELSECFRIFDFDGN_GFIDREEFGDIIRLTGE QLTDEDVDEIFGDSDTDKNSGRIDFDEFLKMVENVQPIYP
Twitch 9x	MQ__DASEEELSECFRIFDKDGNDGFIDREEFGDIIRLTGE QLTDEDVDEIFGDSDTDKN_GRIDFDEFLKMVENVQPIYP

Figure 1. Inhibition of IP3R activity delays TGN export of Cab45 clients
(A) Schematic representation of the Retention Using Streptavidin Hooks (RUSH) assay with fluorescent tagged client molecules. (B) Representative immunofluorescence images of the RUSH experiments showing COMP-EGFP transport in HeLa lines treated with DMSO and 70 nM 2-APB. HeLa cells were transfected with KDEL-IRES-SBP-COMP-EGFP and fixed at 0, 20, 40, and 60 min after the addition of biotin. Z-stack images (d = 0.2 μm) were analyzed. The arrowheads indicate cytoplasmic vesicles. Scale bars, 10 μm. (C) The numbers of COMP budding vesicles from RUSH experiments with KDEL-IRES-SBP-COMP-EGFP in DMSO and 2-APB were quantified. The cytoplasmic vesicles were counted at each time point by analyzing z-stack images (d = 0.2 μm). Scatter dot plot represents the means ± SD of at least three independent experiments (n > 30 cells per condition). Statistical test, Kruskal–Wallis. (D) Plot representing normalized average fluorescence intensity (F.I.) of COMP-EGFP in cells by FACS at 0, 60 and 120 minutes of RUSH experiments, ***p<0.001 (E) Plot representing normalized fluorescence intensity of LyzC-EGFP within TGN (ROI was defined by GALNT1 area) in cells treated with DMSO and 2-APB. (F) Western blot showing LyzC-GFP in Hela cells treated with DMSO and 2-APB in cell lysates and in secreted medium (top) and β-actin as a loading control.

Figure 2. Depletion of VAPA and VAPB delays TGN export of Cab45 clients
(A) Representative immunofluorescence images of RUSH experiments showing COMP-GFP transport in HeLa lines treated with siRNA against VAPA and VAPB proteins. Cells were transfected with KDEL-IRES-SBP-COMP-EGFP and fixed at 0, 20, 40, and 60 min after the addition of biotin. Z-stack images (d = 0.2 μm) were analyzed. Scale bars, 10 μm. (B) The numbers of COMP budding vesicles were quantified. The cytoplasmic vesicles were counted at each time point by analyzing z-stack images (d = 0.2 μm). Scatter dot plot represents the means ± SD of at least three independent experiments (n > 30 cells per condition). Statistical test, Kruskal–Wallis. (C) Plot representing normalized average fluorescence intensity (F.I.) of COMP-EGFP in cells by FACS at 0, 60 and 120 minutes of RUSH experiments in HeLa WT and VAPA/VAPB-dKO lines, ***p<0.001; (D) Plot representing normalized fluorescence intensity of LyzC-EGFP within TGN (ROI was defined by Galnt1 area) in cells treated with non-targeting (control) siRNA and siRNA targeting VAPA and VAPB. (E) Expression of VAPA and VAPB proteins in Hela WT line transfected with control (non-targeting) siRNA, Hela line transfected with siVAPA/siVAPB and Hela VAPA/VAPB-dKO line. β-actin was used as a loading control.

Figure 3. MCS targeting FRET sensor responses to changes of Ca2+ concentration at ER-TGN MCS.
(A) Schematic representation of OSBP and Twitch2b Calcium Sensor targeting Endoplasmic Reticulum-Golgi membrane contact sites. PH domain of OSBP binds to PI4P on the outer leaflet of the TGN membrane and FFAT domain interacts with VAPA on the ER membrane. (B) Calibration curve for Twitch2b sensor performed used buffer with increasing concentration of free Ca2+ ions demonstrating correlation between FRET indices and free Ca2+. (See Materials and Methods for a detailed description). (C) Pseudocolor map of FRET index and dot plot (D) representing FRET indices for PH-FFAT-Twitch2b and PH-FFAT-Twitch9x within Golgi ROI of live cells and at steady state (***p<0.05). (E) Pseudocolor map of FRET index and dot plot (F) representing FRET indices for PH-FFAT-Twitch2b at steady state and treated with 25uM BAPTA-AM for 20 min (***p<0.05). (G) Pseudocolor map of FRET index and dot plot (H) representing FRET indices for PH-FFAT-Twitch2b at steady state and treated with 1uM ionomycin for 20 min (***p<0.05). (I) Pseudocolor map of FRET index and dot plot (J) representing FRET indices for PH-FFAT-Twitch2b at steady state and treated with 1uM Histamine for 2 min (***p<0.05). Pseudocolor bar FRET index value in range 0-0.4.

Figure 4. Ca2+ flux at ER-TGN MCS is coupled to protein trafficking
(A) Schematic representation of experiment and effect on protein trafficking in live cells during 20°C block and cycloheximide (CHX) treatment. (B) Pseudocolor maps of FRET index for PH-FFAT-Twitch2b within Golgi ROI of live cells at described above conditions. Pseudocolor bar FRET index value in range 0-0.4. (C) Dot plot representing FRET index for PH-FFAT-Twitch2b at steady state, after 20°C block for 1 and 2 hours and after 10 min recovery at 37°C (***p<0.05). (D) Dot plot representing FRET index for PH-FFAT-Twitch2b at steady state, treated with CHX for 1, 2, 4 hours and after 1h 20°C block. (***p<0.05), (E) Dot plot representing normalized FRET index for PH-FFAT-Twitch2b at steady state, after 1h 20°C block and effect of ionomycin at 37°C and 20°C conditions (***p<0.05).

A

Figure 5A. The model depicting the role of contact sites and IP3R-dependent release of Calcium ions for SPCA1 dependent sorting at TGN.