FER-LIKE IRON DEFICIENCY-INDUCED TRANSCRIPTION FACTOR (FIT)
accumulates in homo- and heterodimeric complexes in dynamic and inducible
nuclear condensates associated with speckle components

Short title: FIT localizes in condensates

Ksenia Trofimov[1], Regina Gratz[1,§], Rumen Ivanov[1], Yvonne Stahl[2,3], Petra Bauer[1,3,*],
Tzvetina Brumbarova[1,*]

[1] Institute of Botany, Heinrich-Heine-University, 40225 Düsseldorf, Germany
[2] Institute for Developmental Genetics, Heinrich-Heine-University, 40225 Düsseldorf,
Germany
[3] Cluster of Excellence on Plant Science (CEPLAS), Heinrich-Heine-University, 40225
Düsseldorf, Germany
* Shared corresponding authors: Tzvetina Brumbarova (Tzvetina.Brumbarova@hhu.de),
Petra Bauer (Petra.Bauer@hhu.de)
§ Present address: Department of Forest Genetics and Plant Physiology, Swedish
University of Agricultural Sciences, 90183 Umeå, Sweden

The author responsible for distribution of materials integral to the findings presented in
this article in accordance with the policy described in the Instructions for Authors
(https://academic.oup.com/plcell/pages/General-Instructions) is: Petra Bauer
(Petra.Bauer@hhu.de).

Highlights
• FIT undergoes light-induced, reversible condensation and localizes to nuclear
bodies (NBs), likely via liquid-liquid phase separation
• Functionally relevant Ser271/272 defines an intrinsically disordered region and
influences NB formation dynamics
• NBs are preferential sites for FIT dimerization with FIT and bHLH039, dependent
on Ser271/272
• FIT NBs colocalize with NB markers related to splicing and light signaling
• Light conditions inducing NBs are linked with active FIT, in agreement with
elevated FIT target gene expression in roots

Keywords

bHLH039, condensates, FIT, FRET, intrinsically disordered region, blue light, liquid-

liquid phase separation, nuclear body, speckle, SR45

Abbreviations

34	bHLH	basic helix-loop-helix
35	bHLH039	BASIC HELIX-LOOP-HELIX039
36	C	mCherry
37	FIT	FER-LIKE IRON DEFICIENCY-INDUCED TRANSCRIPTION FACTOR
38	FLIM	fluorescence lifetime imaging microscopy
39	FRAP	fluorescence recovery after photobleaching
40	FRET	fluorescence resonance energy transfer
41	G	GFP
42	GFP	GREEN FLUORESCENT PROTEIN
43	IDR	intrinsically disordered region
44	LLPS	liquid-liquid phase separation
45	mCherry	monomeric Cherry
46	mRFP	monomeric RED FLUORESCENT PROTEIN
47	NB	nuclear body
48	NP	nucleoplasm
49	PB	photobody
50	R	mRFP
51	TF	transcription factor

Abstract

Some nuclear proteins undergo condensation, but the functional importance remains often unclear. The basic helix-loop-helix (bHLH) FER-LIKE IRON DEFICIENCY-INDUCED TRANSCRIPTION FACTOR (FIT) integrates internal and external signals to control iron acquisition and growth. The previously described C-terminal residues Ser271/272 allow FIT to form active complexes with subgroup Ib bHLH factors such as bHLH039. FIT has lower nuclear mobility than mutant FITmSS271AA. Here, we show that FIT undergoes a light-inducible subnuclear partitioning into nuclear condensates that we termed FIT nuclear bodies (NBs). FIT NB characteristics were examined using a standardized FIT NB analysis procedure coupled with different types of quantitative and qualitative microscopy-based approaches. FIT condensates were reversible and likely formed by liquid-liquid phase separation. FIT accumulated preferentially in FIT NBs versus nucleoplasm when engaged in protein complexes with itself and with bHLH039. FITmSS271AA, instead, localized to NBs with different dynamics. FIT colocalized with splicing and light signaling NB markers. The light conditions inducing NBs were linked with active FIT and elevated FIT target gene expression in roots. Hence, the inducible highly dynamic FIT condensates link active transcription factor complexes with posttranscriptional regulation processes and light. Inducible FIT nuclear condensates may affect nuclear mobility and integrate environmental and Fe nutrition signals.

Introduction

As sessile organisms, plants must adjust to an ever-changing environment. Readout of environmental cues and rapid acclimation are necessary to ensure the plant's vitality. Accordingly, plants control micronutrient uptake. Overaccumulation causes toxicity but lack of a micronutrient leads to deficiency symptoms. Even though iron is one of the most abundant elements in the soil, its bioavailability as micronutrient is limited in most soils, rendering iron uptake a challenge for plants (Römheld and Marschner, 1986; Wedepohl, 1995).

An essential regulatory protein needed for iron acquisition is the basic helix-loop-helix (bHLH) transcription factor (TF) FER-LIKE IRON DEFICIENCY-INDUCED TRANSCRIPTION FACTOR (FIT; Colangelo and Guerinot, 2004; Jakoby et al., 2004; Yuan et al., 2005; Bauer et al., 2007). FIT is activated upon iron deficiency downstream of a cascade of bHLH TFs (Zhang et al., 2015; Li et al., 2016; Liang et al., 2017; Kim et al., 2019; Gao et al., 2020) and of a calcium-sensing protein kinase able to target phosphorylation site Ser271/272 of FIT (Gratz et al., 2019). FIT alone is not sufficient to upregulate iron acquisition, while it is active in a heterodimeric complex together with a member of the bHLH subgroup Ib such as bHLH039 (Yuan et al., 2008; Wang et al., 2013). Furthermore, FIT action is regulated through protein-protein contacts with multiple key players of hormonal and abiotic stress signaling pathways (Lingam et al., 2011; Le et al., 2016; Wild et al., 2016; Cui et al., 2018; Gratz et al., 2019, 2020). Thus, FIT behaves as a regulatory hub in root cells that perceives external and internal cues to adjust iron acquisition with growth (Schwarz and Bauer, 2020; Kanwar et al., 2021).

The subcellular localization of the FIT-bHLH039 module is remarkable. bHLH039 alone is inactive and present mainly close to the plasma membrane in cytoplasmic foci, while bHLH039 together with FIT localizes in the nucleus (Trofimov et al., 2019). FIT is predominately localized in the nucleus but not as mobile compared to mutant FITmSS271AA, that is a less active mutant form of FIT (Gratz et al., 2019). Subcellular partitioning of proteins involved in nutrient uptake has until now not been enough in the focus of research to understand the significance of the differing subcellular localization patterns.

One prominent type of subnuclear partitioning is conferred by biomolecular condensates, or nuclear bodies (NBs). NBs are membrane-less, nuclear sub-compartments, which can be of stable or dynamic nature. To form condensates, proteins

need to have particular features that enable protein interactions and compaction in three-
dimensional space. IDRs are flexible protein regions that allow conformational changes,
and thus various interactions, leading to the required multivalency of a protein for
condensate formation (Tarczewska and Greb-Markiewicz, 2019; Emenecker et al., 2020).
As Arabidopsis TFs are enriched in IDRs (Strader et al., 2022) it is not unlikely that the
resulting multivalency in TFs drives condensation and results in microenvironments for
interaction, probably more often than so far studied. IDRs are particularly characteristic in
bHLH TFs in vertebrates and invertebrates (Tarczewska and Greb-Markiewicz, 2019),
suggesting that this feature may also be relevant for the bHLH TFs of plants. One
possibility for condensates to form is to undergo liquid-liquid phase separation (LLPS). In
this process, a solution is demixed into two or more phases (Emenecker et al., 2020). This
mechanism has been examined in simplified *in vitro* systems, but the involvement of
different cell components renders the mechanism more complex *in vivo* (Fang et al., 2019;
Riback et al., 2020; Zhu et al., 2021).

NBs comprise an immense variety of types, and plants and animals share several
of them, e.g. the nucleolus, Cajal bodies, and speckles. The nucleolus is involved in
transcription of ribosomal DNA, processing of ribosomal RNA, and ribosome biogenesis
(Kalinina et al., 2018; Lafontaine et al., 2021). Nucleoli share components and function
with Cajal bodies, which are e.g. ribonucleoproteins and RNA processing (Love et al.,
2017; Trinkle-Mulcahy and Sleeman, 2017). Speckles are known spliceosomal sites
(Reddy et al., 2012; Galganski et al., 2017). Plant-specific NBs are photobodies (PBs),
which are triggered by light, temperature, and circadian clock (Pardi and Nusinow, 2021).
PBs harbor regulatory complexes of the photomorphogenic responses, including
photoreceptors like phytochromes (phy) and bHLH TFs belonging to the
PHYTOCHROME INTERACTING FACTORs (PIFs; Pardi and Nusinow, 2021). Another
trigger for inducible and reversible condensate formation is temperature (Jung et al., 2020;
Zhu et al., 2021).

NBs may act as hubs integrating environmental signals (Emenecker et al., 2020;
Meyer, 2020). Especially PBs may combine external cues, such as light, as an input signal
to steer developmental processes (Kaiserli et al., 2015; Meyer, 2020; Pardi and Nusinow,
2021). It is proposed that the formation of NBs could be an ancient mechanism for spatial
organization within the nucleus (Emenecker et al., 2020). As more evidence on

condensation in plants arises, this topic remains barely examined in the scope of plant
nutrition.

The motivation for our study was to elucidate the mechanism behind subcellular
distribution and nuclear mobility of FIT. We had found an interesting hint that FIT may
undergo light-inducible and reversible nuclear condensation, when we detected FIT
nuclear bodies (NBs). We developed a standardized FIT NB analysis procedure and
applied it to characterize quantitative and qualitative aspects of the dynamic NB formation
using different microscopy-based techniques. Thereby, we were able to link FIT NB
formation with the activity status of FIT to form functional protein complexes. We found
that splicing and light signaling were also associated with FIT NBs. Thus, this study lays
ground for FIT NBs being regulatory hubs steering nutritional signalling, and associating
functional significance to FIT protein condensate formation.

Results
FIT localizes to NBs in light-inducible and dynamic manner likely as a result of LLPS
The TF FIT has a dynamic mobility and capacity to form TF complexes inside plant
cells (Gratz et al., 2019; Trofimov et al., 2019). To explore possible mechanisms for
dynamic FIT subcellular localization, we performed a microscopic study on FIT-GFP
protein localization in the root epidermis of the root differentiation zone of 5-d-old iron-
deficient seedlings of *Arabidopsis thaliana* (Arabidopsis), where FIT is active and iron
acquisition occurs (FIT_{pro}:FIT-GFP, this study; $35S_{pro}$:FIT-GFP, Gratz et al., 2019). At first
microscopic inspection, FIT-GFP was evenly distributed within the nucleus (t= 0 in
Figure 1A, FIT_{pro}:FIT-GFP; t= 0 in **Supplemental Figure S1A**, $35S_{pro}$:FIT-GFP), as
expected (Gratz et al., 2019; Trofimov et al., 2019). We report and characterize here a
previously unknown localization behavior of FIT that we found to be coupled with its
activity status. We observed, when whole seedlings were exposed to 488 nm laser light
for several minutes or when they had been shifted from a white light growth chamber into
one with blue light for a time of up to two hours, that FIT became re-localized at the
subnuclear level. After a lag time, FIT-GFP was present in one to four discrete spots per
nucleus, visible after 40 min earliest, sometimes taking up to 120 min to appear (t= 90
minutes in **Figure 1A**, FIT_{pro}:FIT-GFP; t= 40 min in **Supplemental Figure S1A**,
$35S_{pro}$:FIT-GFP). Hence, FIT-GFP was re-localized in light-inducible manner at the
subnuclear level, irrespective of either of the two promoters used. During the imaging

process or an additional hour of darkness, these spots disappeared (**Figure 1A**, t= 120
min), indicating that the formation of these spots was reversible. The observation of FIT
nuclear spots in the root epidermis of the root differentiation zone was very interesting,
suggesting that these might perhaps be NBs containing FIT. However, further inspection
of the nuclear spots in root cells in this differentiating root zone was hampered by several
difficulties, namely the small size and low accessibility of the nucleus, and especially
considering the long lag time for detecting the nuclear spots. These factors made it
impossible for us to apply quantitative fluorescence microscopy techniques to draw
validated conclusions on the nature, dynamics, and functional significance of nuclear
spots in root epidermis cells of the root differentiation zone of iron-deficient seedlings.

We previously employed a reliable plant cell assay for studying FIT functionality,
and we adapted it to study the characteristics of FIT NBs (Gratz et al., 2019, 2020;
Trofimov et al., 2019). Using this assay, FIT-GFP that was transiently expressed in
Nicotiana benthamiana (*N. benthamiana*) leaf epidermis cells under a β-estradiol-
inducible promoter to control protein expression showed a very similar re-localization of
FIT-GFP into nuclear spots as observed in the Arabidopsis root epidermis, again triggered
by treatment with a 488 nm laser light stimulus. Differences were, however, the duration
of the lag time needed to observe this phenomenon, and the number of nuclear spots. As
in Arabidopsis, FIT-GFP localized initially in uniform manner to the entire nucleus (t=0) of
N. benthamiana leaf epidermis cells. A short duration of 1 min 488 nm laser light excitation
induced the formation of FIT-GFP signals in discrete spots inside the nucleus after a lag
time of only five minutes (t=5; **Figure 1B and Supplemental Movie S1A**). The nuclear
FIT spots were systematically initiated, and nearly all nuclei in the imaged leaf disk
showed numerous spots. A similar laser light excitation procedure was previously found
to elicit PB formation of cryptochrome2 (CRY2) in Arabidopsis protoplasts and HEK293T
cells (Wang et al., 2021). We deduced that the spots of FIT-GFP signal were NBs. FIT NB
formation was not dependent on the fluorescent tag, as it was similar for FIT-mCherry
when co-excited with 488 nm laser light (**Figure 1C**). Another TF and interactor of FIT,
ZINC FINGER OF ARABIDOPSIS THALIANA12-GFP (ZAT12-GFP; Le et al., 2016), did
not localize to NBs under the same imaging conditions (**Figure 1D and Supplemental**
Movie S1B). Therefore, we concluded that FIT localization to NBs was a specificity of FIT
and that formation of FIT NBs was not artificially caused by fluorescent tags or the imaging
setup. Importantly, the *N. benthamiana* epidermis expression system was suited to control

the parameters for light-induced triggering of FIT NBs and their quantitative analysis by

fluorescence microscopy. We then developed a standardized experimental procedure for

qualitative and quantitative FIT NB analysis in *N. benthamiana* (hereafter named

'standardized FIT NB analysis procedure'; **Supplemental Figure S1B**).

Liquid-liquid phase separation (LLPS) is a possible way for condensate formation,

and liquid-like features are quantifiable by mobility and shape analysis within condensates

(Shin et al., 2017; Wang et al., 2021). We used the standardized FIT NB analysis

procedure to examine whether this could also be a mechanism underlying the FIT NB

formation. Mobility of FIT NBs was tested with the fluorescence recovery after

photobleaching (FRAP) approach (Bancaud et al., 2010; Trofimov et al., 2019) by

recording the recovery of the fluorescence intensity over time in a bleached NB

(**Figure 1E-G**). According to relative fluorescence intensity the fluorescence signal

recovered to a high extent with FIT NBs (**Figure 1F**), and the calculated mobile fraction

of the NB protein was on average 80% (**Figure 1G**). Shape analysis of FIT NBs showed

that the NBs reached a high circularity score (**Figure 1H**). According to Wang et al. (2021),

fluorescence recovery and circularity scores as the ones measured for FIT NBs reflect

high mobility and circular shape. Thus, FIT NBs behave in a liquid-like manner suggesting

that LLPS might be the mechanism leading to FIT NB formation.

In summary, FIT-GFP forms light-inducible FIT NBs in roots and upon transient

expression in *N. bethamiana* leaf epidermis cells. The developed standardized FIT NB

analysis procedure was well suited for investigating dynamic properties of light-induced

FIT NBs and characterizing them as the likely result of LLPS. Because of these properties,

it is justified to term them 'FIT NBs'. We hypothesized that NB formation is a feature of the

FIT protein that provides regulatory specificity, and we subsequently investigated this

hypothesis using the developed standardized FIT NB analysis procedure in all subsequent

assays below.

FIT forms homodimeric complexes preferentially in NBs, dependent on Ser271/272

Next, we asked which properties of the FIT protein enable NB formation. Residue

Ser271/272 is important for the homo- and heterodimerization capacity of FIT (Gratz et

al., 2019). We therefore asked whether this site has an influence on FIT NB formation,

and we compared the ability for NB formation of phospho-mutated FIT with that of wild-

type FIT-GFP protein. The phospho-mimicking FITmS272E protein did not show

significant changes in its behavior compared to wild-type FIT in the work of Gratz et al. (2019). However, the FITmSS271AA-GFP mutant was less able to mediate induction of iron deficiency responses and its properties differed from wild-type FIT-GFP (Gratz et al., 2019). Hence, we focused on comparing the nuclear NB-forming characteristics between wild-type FIT-GFP and FITmSS271AA-GFP mutant.

FITmSS271AA-GFP also localized to NBs, however with different dynamics. The formation of FITmSS271AA NBs was delayed in time (**Figure 2A**; t=15 instead of t=5). While FIT-GFP NB formation started in the first minutes after excitation (**Supplemental Movie S1A**), FITmSS271AA-GFP NB formation occurred earliest 10 min after excitation (**Supplemental Movie S1C**). In addition, NB number and size of FITmSS271AA-GFP were decreased in comparison to the ones from wild-type FIT-GFP (**Figure 2, B and C**). Hence, the dynamics of NB formation were dependent on Ser271/272.

The process of condensation is facilitated when proteins possess IDRs, since, importantly, IDRs may engage in numerous interactions in space due to rapid conformational changes (Tarczewska and Greb-Markiewicz, 2019; Emenecker et al., 2020). The three-dimensional conformation of wild-type FIT had predicted stretches of intrinsic disorganization, peaking before and at the basic region of the bHLH domain, and two in the C-terminal part, one of them around the Ser271/272 site (termed IDR$^{Ser271/272}$; **Supplemental Figure S2A**). In contrast, in the FITmSS271AA mutant this C-terminal region was no longer classified as IDR (**Supplemental Figure S2B**). This underlined the significance of the Ser271/272 site, not only for interaction but also for FIT NB formation.

We then tested whether FIT homodimerization was preferentially associated with NB formation. For that, we investigated whether FIT-GFP shows a differentiated homodimerization strength, first, inside the NBs versus the nucleoplasm (NP), and second, as wild-type FIT versus the mutant FITmSS271AA-GFP protein by performing anisotropy (or homo-FRET) measurements. Energy transfer between the same kind of fluorescently tagged proteins leads to depolarization of the emitted light (Stahl et al., 2013; Weidtkamp-Peters et al., 2022). Fluorescence anisotropy (FA) describes this depolarization and gives hints on the dimerization and oligomerization status of a protein as the FA value decreases (**Figure 3A**). We measured FA before (t=0) and after NB formation (t=5 for FIT and t=15 for FITmSS271AA), and analyzed the homodimerization strength for the whole nucleus, the NBs, and the residual NP (**Figure 3B-D**). Free GFP

and GFP-GFP constructs were used as references for monomers and dimers (**Figure 3C**
and D).

Whole nucleus FA values were lower at t=5 than at t=0 for FIT-GFP. Additionally,
FA values were lower within the NBs compared to the NP (**Figure 3C**). Compared to wild-
type FIT-GFP, FA values were not reduced for mutant FITmSS271AA-GFP at t=15
compared to t=0. Also, the FA values did not differ between NBs and NP for the mutant
protein (**Figure 3D**). This indicated the presence of homodimeric FIT complexes in NBs.

In summary, wild-type FIT had better capacities to localize to NBs than mutant
FITmSS271AA, presumably due its IDR$^{Ser271/272}$ at the C-terminus. NBs were nuclear sites
in which FIT formed preferentially homodimeric protein complexes.

FIT-bHLH039 interaction complexes preferentially accumulate in FIT NBs

FIT engages in protein-protein interactions with bHLH039 to steer iron uptake
target gene induction in the nucleus, while mutant FITmSS271AA protein is less active in
interacting with bHLH039 (Gratz et al., 2019). Hence, we tested whether FIT also interacts
with bHLH039 preferentially inside NBs and whether mutant FITmSS271AA differs in this
ability from wild-type FIT protein. bHLH039 alone does not localize inside the nucleus but
requires FIT for nuclear localization (Trofimov et al., 2019), so that bHLH039 was not used
alone to test its subnuclear localization.

Upon co-expression, FIT-GFP and bHLH039-mCherry colocalized fully in NBs that
resembled the previously described FIT NBs. In the beginning, both proteins were
uniformly distributed within the nucleus (t=0), and later became localized in NBs (t=5;
Figure 4A).

We then examined the heterodimerization strength of FIT-GFP and bHLH039-
mCherry, and FITmSS271AA-GFP and bHLH039-mCherry by FRET-fluorescence
lifetime imaging microscopy (FRET-FLIM) measurements. In case of protein interaction
(close proximity, ≤10 nm), energy transfer between a fluorescently tagged donor and a
fluorescently tagged acceptor decreases the fluorescence lifetime of the donor
(**Figure 4B;** Borst and Visser, 2010; Weidtkamp-Peters and Stahl, 2017). We quantified
the fluorescence lifetime of FIT-GFP and FITmSS271AA-GFP respective of
heterodimerization before (t=0) and after NB formation (t=5 for FIT and t=15 for
FITmSS271AA) in the whole nucleus, in NBs, and in the NP (**Figure 4C-E**). FIT-GFP and
FITmSS271AA-GFP (donor only) served as negative controls.

Fluorescence lifetime was decreased for the pair FIT-GFP and bHLH039-mCherry
at t=5 within NBs compared to all other measured areas (**Figure 4D**). In contrast to that,
the fluorescence lifetime decreased for the pair FITmSS271AA-GFP and bHLH039-
mCherry at t=15 was not different between NBs and NP (**Figure 4E**). This indicated that
heterodimeric complexes accumulated preferentially in FIT NBs.

In summary, heterodimerization of FIT with bHLH039 was spatially concentrated in
NBs versus the remaining nuclear space and was less prominent for FITmSS271AA.
Hence, the capacity of FIT to form an active TF complex was coupled with its presence in
NBs. The occurrence of FIT homo- and heterodimerization preferentially in NBs suggests
that FIT protein interaction may drive condensation. We therefore concluded that FIT NBs
may be sites with active TF complexes for iron deficiency response regulation.

FIT NBs colocalize with speckle components

Numerous NB types are known, and they are associated with particular proteins
that are indicative of the NB type. To further understand the identity, dynamics, and
function of FIT NBs, we co-expressed FIT-GFP with seven different NB markers from The
Plant Nuclear Marker collection (NASC) and observed NB formation and protein
colocalization before (t=0) and after FIT NB formation (t=5). In cases where we detected
a colocalization with FIT-GFP, we analyzed the localization of NB markers also in the
single expression at t=0 and at t=5 after the 488 nm excitation, to detect potentially
different patterns in single and co-expression.

All seven NB markers were expressed together with FIT-GFP, and according to the
resulting extent of colocalization we subdivided them into three different types. The first
type (type I) did not colocalize with FIT-GFP neither at t=0 nor at t=5. This was the case
for the Cajal body markers coilin-mRFP and U2 snRNP-specific protein U2B"-mRFP
(**Supplemental Figure S3**; Lorković et al., 2004; Collier et al., 2006). Coilin-mRFP
localized into a NB within and around the nucleolus (**Supplemental Figure S3A**). The
NBs of U2B"-mRFP were also close to the nucleolus (**Supplemental Figure S3B**).
Hence, FIT-GFP was not associated with Cajal bodies.

The second type (type II) of NB markers were partially colocalized with FIT-GFP.
This included the speckle components ARGININE/SERINE-RICH45-mRFP (SR45) and
the serine/arginine-rich matrix protein SRm102-mRFP. SR45 is involved in splicing and
alternative splicing and is part of the spliceosome in speckles (Ali et al., 2003), and was

recently found to be involved in splicing of iron homeostasis genes (Fanara et al., 2022).
SRm102 is a speckle component (Kim et al., 2016). SR45-mRFP localized barely in the
NP but inside few and very large NBs that remained constant at t=0 and t=5. FIT-GFP did
not colocalize in those NBs at t=0, however, it colocalized with the large SR45-mRFP NBs
at t=5 (**Figure 5A**). FIT-GFP also localized in typical FIT NBs in the residual NP at t=5
(**Figure 5A**). SRm102-mRFP showed low expression in the NP and stronger expression
in a few NBs that also remained constant at t=0 and t=5. FIT-GFP colocalized with
SRm102-mRFP in only few instances at t=5, but not t=0, while most FIT NBs did not
colocalize with SRm102-mRFP NBs (**Figure 5B**). Both SR45-mRFP and SRm102-mRFP
had the same localization pattern at t=0 and t=5, irrespective of FIT-GFP co-expression
or 488 nm excitation (**Supplemental Figure S4**). These type II NB markers seemed to
recruit FIT-GFP into NBs after 488 nm excitation that were present (pre-existing) before
FIT-GFP NB formation, while FIT-GFP localized additionally in separate FIT NBs. Hence,
FIT became associated with splicing components and speckles upon the light trigger.

A third type (type III) of three NBs markers, namely UAP56H2-mRFP, P15H1-
mRFP, and PININ-mRFP, were fully colocalized with FIT-GFP. Until now, these NB
marker proteins are not well described in plants. UAP56H2 is a RNA helicase, which is
involved in mRNA export (Kammel et al., 2013). P15H1 was found as a putative
Arabidopsis orthologue of an exon junction complex component in humans (Pendle et al.,
2005), while PININ has a redundant role to its paralogue apoptotic chromatin
condensation inducer in the nucleus (ACINUS) in alternative splicing (Bi et al., 2021).
UAP56H2-mRFP and P15H1-mRFP did not localize in NBs and were not responsive to
the 488 nm excitation when expressed alone or together with FIT-GFP at t=0 (**Figure 6,
A and B and Supplemental Figure S4, C and D**). When co-expressed with FIT-GFP and
following the 488 nm excitation, at t=5, the two NB markers adopted the FIT NB pattern
and colocalized with FIT-GFP in FIT NBs (**Figure 6, A and B**). PININ-mRFP was also
uniformly distributed in the nucleus at t=0 like FIT-GFP and fully colocalized with FIT NBs
at t=5 (**Figure 6C**). But curiously, PININ-mRFP showed a very different localization in the
single expression. Predominately, it localized to a very large NB besides several small
NBs with no expression in the NP at t=0 and at t=5 (**Supplemental Figure S4E**). Thus,
FIT-GFP recruited these type III NB marker and speckle proteins fully into FIT NBs. Since
type III NB markers are also potentially involved in splicing and mRNA export from the
nucleus, these same functions may be relevant in FIT NBs.

Taken together, the colocalization studies underlined the dynamic behavior of
inducible FIT NB formation. FIT NBs had a speckle function, in which on the one hand FIT
was recruited itself into pre-existing splicing-related NBs (SR45-mRFP and SRm102-
mRFP, type II), while on the other hand it also recruited speckle-localized proteins into
FIT NBs (UAP56H2-mRFP, P15H1-mRFP, and PININ-mRFP, type III).

PB components influence FIT NB localization and formation
PBs are plant-specific condensates which harbor various light signaling
components (Kircher et al., 2002; Bauer et al., 2004). Among them are the bHLH TFs of
the PIF family. As key regulators of photomorphogenesis, they integrate light signals in
various developmental and physiological response pathways (Leivar and Monte, 2014;
Pham et al., 2018). Indeed, PIF4 may control iron responses in Arabidopsis based on
computational analysis of iron deficiency response gene expression networks
(Brumbarova and Ivanov, 2019) and both PIF proteins intersect with blue light signaling
(Ni et al., 1998; Oh et al., 2004; Pedmale et al., 2016). We tested in the same manner as
described above for NB markers, whether FIT NBs coincide with two of the described PB
markers, PIF3-mCherry and PIF4-mCherry (Van Buskirk et al., 2014; Qiu et al., 2019,
2021).

We detected distinct localization patterns for PIF3-mCherry and PIF4-mCherry. At
t=0, PIF3-mCherry was predominantly localized in a single large PB (**Figure 7A**). In
general, localization of single expressed PIF3-mCherry remained unchanged at t=0 and
t=15 (**Supplemental Figure S5A**). Upon co-expression, FIT-GFP was initially not present
in PIF3-mCherry PB at t=0. After 488 nm excitation and at t=5, FIT NBs were still not
visible. Instead, FIT-GFP accumulated and finally colocalized with the large PIF3-mCherry
PB at t=15, while the typical FIT NBs did not appear (**Figure 7A**).

PIF4-mCherry localized in two different patterns, and both differed substantially
from that of PIF3-mCherry. In the one pattern at t=0, PIF4-mCherry was not localized to
any PBs, but instead was uniformly distributed in the NP as was the case for FIT-GFP.
Such a pattern was also seen at t=15, and then neither PIF4-mCherry nor FIT-GFP were
localized in any PBs/NBs (**Figure 7B**). In the other pattern, PIF4-mCherry and FIT-GFP
localized in multiple PBs at t=0 and t=15, whereas none of them corresponded
morphologically to the typical FIT NBs (**Figure 7C**). The same two localization patterns

were also found for PIF4-mCherry in the single expression, whereby 488 nm excitation
did not alter PIF4-mCherry localization (**Supplemental Figure S5, B and C**).

Hence, FIT was able to localize to PBs when co-expressed with PIF3 and PIF4,
raising the possibility that FIT is a key regulator to cross-connect iron acquisition regulation
and light signaling pathways.

Blue light has a promoting effect on iron acquisition responses downstream of FIT

Previous studies have shown that iron uptake by FIT is diurnally and circadian
clock-regulated (Vert et al., 2003; Santi and Schmidt, 2009; Salomé et al., 2013). Knowing
that FIT NB formation was light-dependent as well as promoting the interaction of the FIT-
bHLH039 complex, and that FIT NBs were colocalizing with light signaling and PB
components, we reasoned that plants exposed to the FIT NB-forming light cues may show
differential FIT-dependent iron uptake responses with respect to control plants exposed
to the regular light conditions. To test this, we grew wild-type Arabidopsis plants for 5 d
under iron sufficient and iron deficient conditions under white light and exposed the plants
additionally for 1 d to blue light. As a control for light quality, we also included exposure to
red and far-red light and darkness besides the regular white light. We measured molecular
iron uptake responses known to be under the control of FIT-bHLH039 in roots (Gratz et
al., 2019).

Iron reductase activity is increased in roots when FIT and bHLH039 are activated
in response to iron deficiency in our growth system (Gratz et al., 2019). Iron reductase
activity was higher under iron deficient conditions compared to iron sufficient conditions
under all white light controls, as expected (**Figure 8A-D**). Interestingly, when seedlings
were exposed to blue light for a day, they induced iron reductase activity in the iron
deficient versus iron sufficient condition more than in the white light control (**Figure 8A**).
On the other hand, exposure to red light did not change the iron reductase activity
compared to the white light control (**Figure 8B**), while exposure to far-red light or darkness
abolished the induction (**Figure 8, C and D**). Hence, only exposure to blue light had an
extra promoting effect on iron uptake in the seedlings compared with the other tested light
qualities.

Increased iron acquisition under iron deficiency and enhanced iron reductase
activity require FERRIC REDUCTASE-OXIDASE2 (FRO2) protein (Yi and Guerinot, 1996;
Robinson et al., 1999), and *FRO2* to be induced, along with *IRON REGULATED*

TRANSPORTER1 (*IRT1*; Eide et al., 1996; Korshunova et al., 1999). *FRO2* and *IRT1* are

target genes of FIT-bHLH039 in our growth system (Gratz et al., 2019, 2020). As a readout

of the activity of the FIT-bHLH039 complex, we examined gene regulation of *FRO2* and

IRT1 as well as *FIT* and *BHLH039* under the same light conditions as described above.

Previously, it was shown that the expression of *IRT1* and *FRO2* was downregulated under

iron deficient conditions in darkness compared to iron deficient conditions in white light,

while the opposite was true for *FIT* (Santi and Schmidt, 2009).

Under white light conditions, gene expression for all four genes was tested in four

independent samples. In 15 out of the 16 cases, gene expression of *FIT*, *BHLH039*, *FRO2*

and *IRT1* was significantly higher under iron deficient than sufficient conditions, as

expected, and in only one case for *FIT*, it was similar and not significantly enhanced

(**Figure 9A-P, with exception of E**, *FIT*). Under blue light, *FIT* and *BHLH039* expression

was induced upon iron deficiency in the same manner as in white light (**Figure 9, A and

B**), whereas *FRO2* and *IRT1* were induced to a higher level in response to iron deficiency

in blue light versus white light (**Figure 9A-D**). Again, plant responses were different upon

exposure to other light conditions. *FIT* and *IRT1* were similarly induced, while *BHLH039*

and *FRO2* were even less induced by iron deficiency under red light than white light

(**Figure 9E-H**). In both, the far-red light and darkness situations, *FIT* was induced under

iron deficiency versus deficiency, while on the other side, *BHLH039*, *FRO2* and *IRT1* were

not induced at all in these light conditions (**Figure 9I-P**). *FIT* was similarly induced under

far-red light and white light (**Figure 9I**), whereas in darkness it was even induced to higher

level (**Figure 9M**), as previously reported (Santi and Schmidt, 2009).

To sum up, the FIT-bHLH039-dependent target gene expression is similarly light

quality-dependent as the FIT-bHLH039-dependent iron reductase activity. Hence, very

interestingly, blue light cues which lead to FIT NB formation, promote FIT-bHLH039-

dependent iron uptake responses, fitting with the above finding that FIT NBs are linked

with active TF complexes in condensates.

Discussion

In this study, we uncovered a previously unknown phenomenon, the blue light-

induced and reversible accumulation of FIT condensates in FIT NBs. LLPS was most

likely the underlying mechanism for this highly dynamic process. FIT NBs were enriched

in active FIT TF complexes that are required for iron deficiency gene regulation. FIT

associated with speckles and PBs in a highly dynamic fashion, indicating a function in
transcriptional and post-transcriptional control. Blue light exposure resulted in enhanced
molecular FIT-bHLH039-dependent iron acquisition responses. Based on these data, we
propose that FIT NBs are dynamic microenvironments with active FIT TF complexes that
possibly are hubs to enhance and cross-connect transcriptional iron deficiency gene
expression with post-transcriptional regulation and light signaling.

A standardized procedure for FIT NB induction was crucial to delineate the
characteristics of FIT NBs in reliable manner

A major aim of this study was to characterize the nature and potential function of
light-induced FIT NBs. To be able to apply the quantitative microscopy-based techniques,
we needed to control the appearance of NBs in reliable manner. This was clearly a
limitation for inspection of root epidermis cells of the root differentiation zone in iron-
deficient plants in which FIT-controlled iron uptake takes place. Not every root epidermis
cell showed NBs and only few FIT NBs were detectable after a delay of 40 min to 120 min.
Since condensation depleted FIT protein in the nucleoplasm, the remaining low FIT
protein concentration can be the reason why FIT NBs remained few in number in the
Arabidopsis root cells. The *N. benthamiana* protein expression system did not present
these limitations and high-quality measurement data were obtained for all experimental
series. Furthermore, this expression system is a well-established and widely utilized
system in plant biology (Martin et al., 2009; Bleckmann et al., 2010; Leonelli et al., 2016;
Burkart et al., 2022). The developed standardized assay generated reliable and accurate
data for statistical analysis and quantification to conclude about FIT NB characteristics.

Condensation likely explains the reduced mobility of FIT-GFP versus
FITmSS271AA-GFP seen in a previous study (Gratz et al., 2019). In liquid state,
condensates are still more mobile than in the solid one. The condensate formation was
reversible, speaking in favor of a liquid state of the condensates. According to FRAP data,
FIT NBs maintained a dynamic exchange of FIT protein with the surrounding NP. FIT NBs
were also mostly of circular shape. Circular condensates appear as droplets, in contrast
to solid-like condensates that are irregularly shaped (Shin et al., 2017). These
characteristics speak in favor of liquid-like features, suggesting that LLPS underlies FIT
NB formation. A similar situation was described for CRY2 PBs, which were also reversible
and of circular shape with mobile protein inside PBs (Wang et al., 2021). bHLH039 was

found accumulated in cytoplasmic foci at the cell periphery (Trofimov et al., 2019). In such
foci, bHLH039 was immobile, and we suspect it was in a non-functional state in the
absence of FIT. This underlines the understanding that liquid condensates such as FIT
NBs are dynamic microenvironments, whereas immobile condensates point rather
towards a solid and pathological state (Shin et al., 2017).

In conclusion, the properties of liquid condensation along with the findings that it
occurred irrespective of the fluorescence protein tag preferentially with wild-type FIT, but
with different dynamics for the mutant FITmSS271AA and not at all for ZAT12, allowed us
to coin the term of 'FIT NBs'.

499   IDR$^{Ser271/272}$ was crucial for interaction and NB formation of FIT

FIT NBs were hotspots for FIT interaction, allowing to assume that they are
integrated in the iron deficiency response as interaction hubs. FIT formed homodimers
and heterodimers with bHLH039 preferentially in NBs compared with the NP. These
abilities distinguished wild-type FIT and mutant FITmSS271AA. According to these
findings FITmSS271AA was less successful in interacting within NBs, indicating that wild-
type FIT is a multivalent protein and IDR$^{Ser271/272}$ is important for that. bHLH proteins
interact with other proteins via the helix-loop-helix interface, which may certainly also be
the case for FIT. Our study supports previous reports that FIT protein interaction via its C-
terminus is relevant (Lingam et al., 2011; Le et al., 2016; Gratz et al., 2019). The property
of being able to interact via the HLH and via the C-terminal domain allows FIT to be
multivalent. It could not be distinguished whether FIT homodimers were a prerequisite for
the localization of bHLH039 in NBs or whether FIT-bHLH039 complexes also initiated NBs
on their own.

The predicted C-terminal FIT IDR$^{Ser271/272}$ was relevant for NB formation capacity.
IDRs are often required for protein interactions of hub proteins since the flexible IDRs
adapt to interactions with multiple protein partners and are therefore crucial for
multivalency (Tarczewska and Greb-Markiewicz, 2019; Emenecker et al., 2020; Salladini
et al., 2020). Besides, evidence exists that the amino acid composition of IDRs is crucial
for condensation (Powers et al., 2019; Emenecker et al., 2021; Huang et al., 2022). Very
interestingly, posttranslational modification in form of phosphorylation within IDRs is
suggested to be a mechanism to regulate condensate formation (Owen and Shewmaker,
2019). Ser271/272 is targeted by a FIT-interacting protein kinase that was shown to affect

FIT activity *in vivo* and FIT phosphorylation *in vitro* (Gratz et al., 2019). Hence,
phosphorylation of Ser271/272 might perhaps be a trigger for NB formation *in vivo*.

524 Formation of FIT NB could happen *de novo* but also associate with pre-existing
525 condensates in the nucleus

FIT may have formed FIT NBs as entirely newly formed structures upon the light
trigger. But it is also possible that FIT joined pre-existing NBs, which then became the
structures we termed FIT NBs. Partial or full colocalization of FIT-GFP with NB and PB
markers revealed the remarkably high and intriguing dynamic nature of FIT NBs and
suggests that both possibilities are plausible. FIT NBs are light-triggered, and this speaks
in favor of pre-existing NBs. Since FIT does not possess light-responsive domains, it is
most likely that a light-responsive protein must be inducing FIT NB formation. The basic
leucine zipper TF ELONGATED HYPOCOTYL5 (HY5) could be a good candidate, since
HY5 is a mobile protein involved in iron acquisition in tomato (Gao et al., 2021; Guo et al.,
2021) and acts positively on iron uptake and gene expression of *FIT*, *FRO2*, and *IRT1* in
Arabidopsis (Mankotia et al., 2023). Possibly activation and condensation involve not only
the studied NB and PB markers but also potentially signaling proteins or further scaffold
proteins that are part of the multivalent protein complexes in FIT NBs. On the other hand,
FIT-GFP accumulated not only in FIT NBs but also in the pre-existing NBs with type II NB
markers (SR45 and SRm102) after the FIT NB induction procedure. In this respect, type II
markers were similar to PIF3 and PIF4. FIT-GFP was recruited to pre-existing PBs and
again only after the light trigger. Interestingly, typical FIT NB formation did not occur in the
presence of PB markers, indicating that they must have had a strong effect on recruiting
FIT. Overall, the dynamics of FIT colocalization with type II NB and PB markers suggest
that these condensates dictated FIT condensation in their own pre-existing NBs/PBs. This
recruiting process could be navigated via protein-protein interaction since this is the
driving force of condensation (Kaiserli et al., 2015; Emenecker et al., 2020).

Speaking in favor of a *de novo* FIT NB formation is the localization with type III NB
markers. The three fully colocalizing type III NB markers (UAP56H2, P15H1 and PININ)
accumulated only in FIT NBs upon co-expression with FIT and mostly not on their own.
The same was true for bHLH039, that joined FIT in FIT NBs, showing that FIT not only
facilitated bHLH039 nuclear localization (Trofimov et al., 2019) but also condensation.
Interestingly, FIT was able to change PININ nuclear localization. In single expression,

PININ was localized to a major large NB, but in colocalization with FIT it joined the typical FIT NBs. This suggests that FIT dictates bHLH039 and type III NB markers and highlights that FIT is also able to set the tone for NB formation. Hence, FIT can recruit other proteins into NBs, and it is possible that FIT forms its own NBs. Protein-protein interaction could underly this recruitment, as evident for bHLH039 (Kaiserli et al., 2015; Emenecker et al., 2020). Ultimately, as a high diversification of condensates exists, a combination of newly formed NBs and localization to pre-existing NBs cannot be ruled out. Given the variety of proteins localizing in condensates, effort in isolating FIT NBs and identification of proteins within FIT NBs is necessary to further uncover the driving forces of FIT NB formation.

FIT NBs might have a transcriptional and post-transcriptional function

Since the type II and III markers are splicing components, the colocalization studies suggest that FIT NBs are speckles. On the one hand, the speckle nature coincides well with the dynamic nature of FIT NBs. Like FIT NBs, speckles are highly dynamic. They are forming around transcriptionally active sites in the interchromatin regions recruiting several protein functions like mRNA synthesis, maturation, splicing and export (Reddy et al., 2012; Galganski et al., 2017). The type II speckle component SR45, for instance, was shown to be a highly mobile protein in speckles and required phosphorylation for proper speckle localization (Ali et al., 2003; Reddy et al., 2012). These processes fit well to the described FIT NB attributes. On the other hand, speckle components are also linked with epigenetic mechanisms (Mikulski et al., 2022). The characterization of FIT NBs as speckles is interesting because regulation of splicing and epigenetic regulation is associated with iron deficiency gene expression. Genes were spliced incorrectly in a *sr45-1* null mutant Arabidopsis line, and gene expression of *FIT* and FIT target genes was increased in *sr45-1* seedlings, showing that an interplay between SR45 and the iron uptake machinery exists (Fanara et al., 2022). Alternative splicing was detected for FIT targets and the *BHLH* subgroup Ib genes in iron-deficient versus iron-sufficient conditions (Li et al., 2013). Hence, FIT NBs may regulate iron uptake gene expression at posttranscriptional level. Notably, PININ (type III), together with ACINUS, were shown to stabilize SR45 (type II) in plants (Bi et al., 2021). Further, UAP56H2, P15H1, and PININ (type III) are connected to SR45 and SRm102 (type II) in mammalian cells as all being part of the exon junction complex and interacting with each other (Lin et al., 2004; Pendle et al., 2005). This is an interesting parallel, as it suggests that type II and type III marker

localization is conserved across kingdoms, underlying the ancient nature of condensates.

Indeed, SR45 and PININ located to a very large NB in non-induced cells. This opens the

possibility that the two proteins might localize to the same speckle, as also might FIT.

Taken together, the observations confirm the high diversification and complexity of FIT

NBs and speckles (Lorković et al., 2008) and it is tempting to speculate that FIT might

regulate splicing and alternative splicing of its target genes by recruiting speckle

components.

FIT is itself a direct target of FIT and the FIT-bHLH039 complex (Wang et al., 2007;

Naranjo-Arcos et al., 2017), and perhaps FIT NB speckles appear at the *FIT* transcription

site. Indeed, coupling of transcription with splicing or alternative splicing is an established

idea in the mammalian field, and evidence for co-transcriptional splicing in plants is also

recently rising (Nojima et al., 2015; Zhu et al., 2018; Chaudhary et al., 2019). Mediator

complex condensation was shown to drive transcriptional control (Boija et al., 2018) and

interestingly, FIT was also shown to interact with Mediator complexes, directly and

indirectly (Yang et al., 2014; Zhang et al., 2014). Besides, other studies suggest TF

condensation to be involved in transcriptional regulation (Kaiserli et al., 2015; Huang et

al., 2022). Possibly, the basic region of FIT and bHLH039 might be accessible for DNA

binding either within or outside of NBs to regulate target genes (Boija et al., 2018; Brodsky

et al., 2020). In further studies, it will be interesting to analyze whether DNA and mRNA

FIT targets are present inside FIT NBs and whether FIT may also interact directly with

other speckle components.

The interesting finding that blue light exposure not only caused the formation of FIT

NBs, that we could link with active TF complexes, but also promoted FIT-bHLH039-

dependent iron mobilization responses can be interpreted as follows: FIT NBs may indeed

serve the rapid rearrangement of TFs to enhance target gene expression and subsequent

physiological responses. Changes of the light spectrum over the day could serve as a cue

for a high photosynthetic period, such a high blue light component, and induce stronger

iron uptake. Due to photosynthesis, chloroplasts and photosynthetically active leaves may

serve as a major sink for iron. Under far-red light, which is as shade signal, and in

darkness, photosynthetic activity is lowered. In turn, lower amounts of iron may be

acquired. Regulation of FIT protein abundance occurs rather on the posttranslational level

than on transcriptional (Lingam et al., 2011; Meiser et al., 2011; Gratz et al., 2019, 2020).

On the other hand, *FIT* is itself a target of FIT-bHLH039 (Wang et al. 2007). The fact that

FIT gene expression is differently regulated from *IRT1* and *FRO2* depending on light

quality is intriguing. Perhaps FIT protein is readily available during the day. The formation

and dissociation of NBs could be a fast way to adjust iron uptake and specifically uncouple

IRT1 and *FRO2* from FIT regulation and therefore adapt even better to environmental

changes. Reversibility of FIT condensation might rapidly switch off the process, possibly

by phosphorylation triggers leading to inactivation of FIT (Gratz et al., 2020). Reversibility

of condensates was shown in the context of light, temperature, or stresses, once the

stimulus was removed (Jung et al., 2020; Wang et al., 2021; Zhu et al., 2021; Chen et al.,

2022; Wang et al., 2022). This supports the hypothesis that plant condensates are hubs

for fast adaptation to external cues.

As we focused to characterize the phenomenon, the physiological integration and

regulation of the induction of FIT NB formation can be subject of future studies. The rapid

speed by which FIT NB appeared within 5 min in *N. benthamiana* leaf cells speaks in favor

of protein rearrangement rather than protein synthesis. The long duration of FIT NB

formation after blue light induction in Arabidopsis roots suggests that signal transduction

was more complex and possibly involved intracellular or even cell-to-cell and long-

distance leaf-to-root signaling. In how far a long-distance signal or a signaling cascade

triggered by light is involved in FIT NB formation in roots remains to be investigated, but

CRY1/CRY2 and HY5 are promising candidates for further studies, as their involvement

in iron deficiency response regulation has been highlighted (Gao et al., 2021; Guo et al.,

2021; Mankotia et al., 2023). Since Arabidopsis germinates in the light, it is well

conceivable that the FIT NB-forming property is part of a seedling establishment

procedure during early phases when roots can be exposed in the light. In order to undergo

phase separation, a certain protein concentration must be reached (Bracha et al., 2018).

Since FIT protein is subject of proteasomal turnover in roots, FIT NB formation may

depend on FIT protein interaction partners in roots that need to be activated (Lingam et

al., 2011; Meiser et al., 2011).

In summary, FIT localizes to dynamic and reversible NBs. FIT NBs contain active

TF complexes for iron acquisition gene expression and are speckles that link

transcriptional with post-transcriptional regulation. The appearance of FIT NBs is inducible

by blue light, a condition that promotes iron mobilization responses, and light regulated

PB components are connected with FIT NBs and vice versa (**Figure 10**). It will be

interesting in the future to test hormonal and environmental triggers that may stabilize FIT

protein prior to examining the initiation of FIT NBs in root physiological situations and to

investigate the effects on transcriptional and posttranscriptional regulation of FIT targets.

Materials and methods

Plant material and growth conditions

Arabidopsis thaliana experiments were conducted with the Col-0 accession. Arabidopsis line FIT$_{pro}$:FIT-GFP used for imaging was obtained as follows: First, the FIT$_{pro}$:FIT-GFP construct was obtained in a plant gene expression binary vector via GreenGate cloning (Lampropoulos et al., 2013). *FIT* promoter sequence of 2012 bp upstream of the CDS was cloned into pGGA000 entry vector using pFIT GG fw and pFIT GG rv primers with respective overhang sequences (**Supplemental Table S1**). FIT CDS (954 bp, without stop codon) was cloned into pGGC000 entry vector using cFIT GG fw and cFIT GG rv primers with respective overhang sequences (**Supplemental Table S1**). The final construct was created via the GreenGate reaction with the pGGZ001 destination vector, containing the entry vectors pGGA000 FIT promoter, pGGB003 *B-dummy* for N-tag, pGGA000 FIT CDS, pGGD001 *linker*-GFP, pGGE009 *UBQ10* terminator, and pGGF001 $p^dMAS^e:BastaR^f:t^gMAS$, and verified by sequencing. The plasmid was transferred to *Rhizobium radiobacter* pSOUP strain (GV3103). Transgenic plants (in Col-0 wild-type background) were obtained by floral dip transformation (Clough and Bent, 1998) and subsequent Basta selection. *Arabidopsis thaliana* 2x35S$_{pro}$:FIT-GFP/*fit-3* line used for imaging was previously described in Gratz et al. (2019).

Arabidopsis thaliana FIT$_{pro}$:FIT-GFP seedlings were used for localization studies. Seeds were sterilized and grown upright on Hoagland medium plates (macronutrients: 1.5 mM Ca(NO$_3$)$_2$ · 4H$_2$O, 0.5 mM KH$_2$PO$_4$, 1.25 mM KNO$_3$, 0.75 mM MgSO$_4$ · 7H$_2$O; micronutrients: 0.075 µM (NH$_4$)6Mo$_7$O$_{24}$ · 4H$_2$O, 1.5 µM CuSO$_4$ · 5H$_2$O, 50 µM H$_3$BO$_3$, 50 µM KCl, 10 µM MnSO$_4$ · H$_2$O, 2 µM ZnSO$_4$ · 7H$_2$O; 1.4 % (w/v) plant agar, 1 % (w/v) sucrose, pH 5.8) with no iron supply for 5 d under long day conditions (16 h light/8 h dark) at 21°C in a plant chamber (CLF Plant Climatics) under white light (120 µmol m^{-2} s^{-1}).

For iron reductase activity assays and gene expression analysis, *Arabidopsis thaliana* seedlings were sterilized and grown upright on Hoagland medium plates with either no iron supply or with 50 µM FeNaEDTA supply for 5 d under long day conditions (16 h light/8 h dark) at 21°C in a plant chamber (CLF Plant Climatics) under white light (120 µmol m^{-2} s^{-1}). For different light treatments plates were transferred to LED light chambers (CLF Plant Climatics) for 1 d under continuous blue light (70 µmol m^{-2} s^{-1}), continuous red light (55 µmol m^{-2} s^{-1}), continuous far-red light (65 µmol m^{-2} s^{-1}) or darkness. White light was used as control (125 µmol m^{-2} s^{-1}).

687   *Nicotiana benthamiana* plants were grown in the greenhouse facility for approx. 4

688 weeks under long day conditions (16 h light/8 h dark) for use in transient protein

689 expression experiments.

Root iron reductase activity assay

6-d-old seedlings were grown as described in the text. Root iron reductase activity

was determined as described in Le *et al.* (2016). Plants were washed with 0.1 M

$Ca(NO_3)_2 \cdot 4H_2O$ and subsequently incubated for 1 h in darkness at room temperature in

the iron reductase solution (300 µM Ferrozine, 100 µM FeNaEDTA). The absorbance of

the ferrozine-Fe^{2+} complex was measured at 562 nm and used to calculate the root iron

reductase activity normalized to root fresh weight.

Gene expression analysis by RT-qPCR

6-d-old seedlings were grown as described in the text. The reverse transcription-

real time-quantitative polymerase chain reaction (RT-qPCR) procedure and analysis is

described in Abdallah and Bauer (2016). Briefly, plant material was harvested, using three

biological replicates, and shock frozen. RNA was extracted (peqGOLD Plant RNA Kit,

PEQLAB). Subsequently, an amount of 0.5 µg of total RNA was used for cDNA synthesis

(Thermo Scientific). Diluted cDNA, SYBR green master mix (Thermo Scientific) and

respective primers (**Supplemental Table S1**) were pipetted in a 96 well plate and qPCR

was performed with two technical duplicates with the CFX96 Touch™ Real-Time PCR

Detection System (Bio-Rad). qPCR data were analyzed, and quantification was obtained

according to the mass standard curve analysis procedure. Normalization was performed

using the reference gene *ELONGATION FACTOR1A* (*EFc*). Statistical analysis was

performed with absolute normalized gene expression levels.

Microscopy of *Arabidopsis thaliana* seedlings

Protein localization studies in roots of 5-d-old seedlings of the *Arabidopsis thaliana*

line FIT$_{pro}$:FIT-GFP and 35S$_{pro}$:FIT-GFP were performed with the widefield microscope

ELYRA PS (Zeiss) equipped with a EMCCD camera. To induce FIT NB formation, whole

seedlings were exposed to 488 nm laser light for several minutes. GFP was excited with

a 488 nm laser and detected with a BP 495-575 + LP 750 beam splitter. Images were

acquired with the C-Apochromat 63x/1.2 W Korr M27 (Zeiss) objective, pixel dwell time of

1.6 µs and frame size of 512x512. Pictures were processed with the manufacturer's
software ZEN lite (Zeiss).

Generation of fluorescent constructs

All constructs used in this study are listed in **Supplemental Table S2**. Generation
of fluorescent translational C-terminal fusion of PIF3 and PIF4 with mCherry was
performed with Gateway Cloning. CDS of *PIF3* was amplified with the PIF3 GW fw and
PIF3 GW rv primers (**Supplemental Table S1**), and CDS of *PIF4* was amplified with the
PIF4 GW fw and PIF4 GW rv primers (**Supplemental Table S1**), and introduced into the
entry vector pDONR207 via the BP reaction (Life Technologies) and subsequently into
the inducible pABind 35S$_{pro}$:mCherry destination vector (Bleckmann et al., 2010) via the
LR reaction (Life Technologies). Finally, *Rhizobium radiobacter* was transformed with the
constructs for transient transformation of *Nicotiana benthamiana* leaf epidermal cells.

Transient transformation of *Nicotiana benthamiana* leaf epidermal cells

Transient protein expression was performed in *Nicotiana benthamiana* leaf
epidermal cells according to Bleckmann et al. (2010). This was performed for localization
studies, FRAP measurements, anisotropy (homo-FRET) measurements, FRET-FLIM
measurements, and NB quantification. Cultures of *Rhizobium radiobacter* containing the
construct of interest (**Supplemental Table S2**) were incubated overnight and cell were
pelleted and dissolved in AS medium (250 µM acetosyringone (in DMSO), 5 % (w/v)
sucrose, 0.01 % (v/v) silwet, 0.01 % (w/v) glucose). An OD$_{600nm}$ of 0.4 was set for all
constructs. A *Rhizobium radiobacter* strain containing the silencing repressor p19 vector
(Shamloul et al., 2014) was used additionally for bHLH039-mCherry to enhance
expression. After 1 h incubation on ice the suspension was infiltrated with a syringe into
the abaxial side of the leaf. *Nicotiana benthamiana* plants were kept under long day
conditions (16 h light/8 h dark) in the laboratory after infiltration. Imaging was performed
2-3 d after infiltration. Expression of constructs with an inducible 35S promoter was
induced 16 h prior to imaging with β-estradiol (20 µM β-estradiol (in DMSO), 0.1 % (v/v)
Tween 20).

Confocal microscopy

For localization studies a confocal laser scanning microscope LSM780 (Zeiss) was
used. Imaging was controlled by the ZEN 2.3 SP1 FP3 (Black) (Zeiss) software. GFP was

excited with a 488 nm laser and detected in the range of 491-553 nm. mCherry and mRFP

were excited with a 561 nm laser and detected in the range of 562-626 nm. Fluorophore

crosstalk was minimized by splitting of the excitation tracks and reduction of emission

spectrum overlap. Images were acquired with the C-Apochromat 40x/1.20 W Korr M27

(Zeiss) objective, zoom factor of 8, pinhole set to 1,00 AU, pixel dwell time of 1.27 µs and

frame size of 1.024x1.024. Z-stacks for quantification were taken with the same settings,

except with pixel dwell time of 0.79 µs and frame size of 512x512. Pictures were

processed with the manufacturer's software ZEN lite (Zeiss).

Standardized FIT NB analysis procedure

Following *Nicotiana benthamiana* leaf infiltration with *Rhizobium radiobacter*, FIT-

GFP protein expression was induced after 2-3 d by β-estradiol, as described above. 16 h

later, a leaf disc was excised and FIT-GFP fluorescence signals were recorded (t=0). The

leaf disc was excited with 488 nm laser light for 1 min. 5 min later, FIT-GFP accumulation

in FIT NBs was observed (t= 5 min). See **Supplemental Figure S1**. This procedure was

modified by using different time points for NB analysis and different constructs

(**Supplemental Table S2**) and co-expression as indicated in the text. Imaging was

performed at the respective wavelengths for detection of GFP and mRFP/mCherry.

FRAP measurements

FRAP measurements (Bancaud et al., 2010; Trofimov et al., 2019) were performed

at the confocal laser scanning microscope LSM780 (Zeiss). Imaging was controlled by the

ZEN 2.3 SP1 FP3 (Black) (Zeiss) software. GFP was excited with a 488 nm laser and

detected in the range of 491-553 nm. Images were acquired with the C-Apochromat

40x/1.20 W Korr M27 (Zeiss) objective, zoom factor of 8, pinhole set to 2,43 AU, pixel

dwell time of 1.0 µs, frame size of 256x256, and 300 frames. After 20 frames, a NB was

bleached with 50 iterations and 100% 488 nm laser power. Fluorescence intensity was

recorded for the bleached NB (ROI), a non-bleached region equal in size to the NB (BG)

as well as for the total image (Tot). Values were calculated and processed in Excel

(Microsoft Corporation). Background subtraction and normalization to calculate the

relative fluorescence intensity was performed as follows: [(ROI(t)-BG(t)/Tot(T)-

BG(t))*(Tot(t0)-BG(t0)/ROI(t0)-BG(t0))]. The mobile fraction was calculated as follows:

$[(F_{end}-F_{post})/(F_{pre}-F_{post})*100]$. F_{pre} marks the average of the 20 values before bleaching,

F_{post} marks the value right after the bleaching, and F_{end} marks the average of the 280

values after the bleaching. Pictures were processed with the manufacturer's software ZEN
lite (Zeiss).

Anisotropy (homo-FRET) measurements

Anisotropy measurements (Stahl et al., 2013; Weidtkamp-Peters et al., 2022) were
performed at the confocal laser scanning microscope LSM780 (Zeiss) equipped with a
polarization beam splitter, bandpass filter (520/35), and a single-photon counting device
HydraHarp (PicoQuant) with avalanche photo diodes (τ-SPADs). Emission was detected
in parallel and perpendicular orientation. Rhodamine 110 was used to determine the G
factor to correct for the differential parallel and perpendicular detector sensitivity.
Calibration of the system was performed for every experiment and measurements were
conducted in darkness. Free GFP and GFP-GFP were used as references for mono- and
dimerization, respectively. GFP was excited with a linearly polarized pulsed (32 MHz)
485 nm laser and 0.05-1 µW output power. Measurements were recorded with a C-
Apochromat 40x/1.20 W Korr M27 (Zeiss) objective, zoom factor of 8, pixel dwell time of
12.5 µs, objective frame size of 256x256, and 40 frames. Measurements were controlled
with the manufacturer's ZEN 2.3 SP1 FP3 (Black) (Zeiss) software and SymPhoTime 64
(PicoQuant) software. SymPhoTime 64 (PicoQuant) software was used for analysis in the
respective regions of interest (whole nucleus, NB, NP) and to generate color-coded FA
value images. Minimal photon count was set to 200.

FRET-FLIM measurements

FRET-FLIM measurements (Borst and Visser, 2010; Weidtkamp-Peters and Stahl,
2017) were taken at the confocal laser scanning microscope FV3000 (Olympus) equipped
with a multi-photon counting device MultiHarp 150 (PicoQuant) with avalanche photo
diodes (τ-SPADs) and bandpass filter (520/35). Erythrosine B (quenched in saturated
potassium iodide) was used to record the Instrument Response Function to correct for the
time between laser pulse and detection. Calibration of the system was performed for every
experiment and measurements were conducted in darkness. FIT-GFP and
FITmSS271AA-GFP were used as negative controls (donor only), FIT-GFP or
FITmSS271AA-GFP (donor) and bHLH039-mCherry (acceptor) as FRET pair. GFP was
excited with a linearly polarized pulsed (32 MHz) 485 nm laser and 0.01-0.1 µW output
power. Measurements were recorded with a UPLSAPO 60XW (Olympus) objective, zoom
factor of 8, pixel dwell time of 12.5 µs, objective frame size of 256x256, and 60 frames.

Measurements were controlled with the manufacturer's FV31S-SW (Olympus) software
and SymPhoTime 64 (PicoQuant) software. SymPhoTime 64 (PicoQuant) software was
used for analysis in the respective regions of interest (whole nucleus, NB, NP) and to
generate color-coded fluorescence lifetime value images. Number of parameters for the
fit depended on the region of interest.

Circularity quantification

Circularity quantification was performed with the software ImageJ (National
Institutes of Health). Full intensity projection images were generated from Z-stacks in the
ZEN lite (Zeiss) software and exported as TIFF (no compression, all dimensions). Images
were duplicated in ImageJ and converted to RGB and 8-bit. Correct scale was set (in μm)
under 'Analyze' - 'Set Scale'. Threshold for the intensity limit (areas below that limit were
not considered for quantification) was set under 'Image' - 'Adjust' - 'Threshold' and was
set manually for every image. To separate the nuclear bodies better, 'Process' - 'Binary' -
'Watershed' was used. Parameters that should be quantified were selected under
'Analyze' - 'Set Measurements'. To perform the analysis, 'Analyze' - 'Analyze Particles'
was selected. Calculated values were further processed in Excel (Microsoft Corporation).

Nuclear body quantification

Nuclear body quantification was performed with the software ImageJ (National
Institutes of Health) and additional plugin '3D Object Counter'. Z-stacks were exported
from the ZEN lite (Zeiss) software as TIFF (no compression, all dimensions) first. In
ImageJ, Z-stacks were converted to RGB and 8-bit. Correct scale was set (in μm) under
'Properties'. Parameters that should be quantified were selected under 'Plugins' - '3D
Object Counter' - 'Set 3D Measurements'. To perform the analysis, 'Plugins' - '3D Object
Counter' - '3D object counter' was selected. Threshold for the intensity limit (areas below
that limit were not considered for quantification) was set manually for every z-stack.
Calculated values were further processed in Excel (Microsoft Corporation). Only size
between 0,01-15 μm³ was considered.

Protein domain prediction

IDRs in FIT/FITmSS271AA were predicted with the tool PONDR-VLXT
(www.pondr.com, Molecular Kinetics, Inc.). According to the sequence of the protein, a
PONDR score was determined for each amino acid. A score above 0.5 indicates intrinsic

disorder. The bHLH domain of FIT was predicted with InterPro (www.ebi.ac.uk/interpro,

EMBL-EBI).

Statistical analysis

Line and bar diagrams represent the mean and standard deviation. Box plots show

25-75 percentile with min-max whiskers, mean as small square and median as line.

Graphs and statistical analysis were created and performed with OriginPro (OriginLab

Corporation). Data was tested for normal distribution with the Shapiro-Wilk test. Statistical

significance of data with normal distribution was tested by one-way Anova with Tukey

post-hoc test. Statistical significance of data with non-normal distribution was tested by

Mann-Whitney test. Different letters indicate statistically significant differences ($P < 0.05$).

Illustrations were created with BioRender.com.

Accession numbers

Sequence data from this article can be found in the EMBL/GenBank data libraries

under accession numbers*: bHLH039* (AT3G56980), *COILIN* (AT1G13030), *FIT*

(AT2G28160), *FRO2* (AT1G01580), *IRT1* (AT4G19690), *P15H1* (AT1G11570), *PIF3*

(AT1G09530), *PIF4* (AT2G43010), *PININ* (AT1G15200), *SR45* (AT1G16610), *SRm102*

(AT2G29210), *U2B"* (AT2G30260), *UAP56H2* (AT5G11170), and *ZAT12* (AT5G59820).

Figures

Figure 1. FIT accumulated in nuclear condensates, termed FIT nuclear bodies (NBs) in a light-inducible manner, most likely following liquid-liquid phase separation (LLPS).

Figure 2. The FIT C-terminal Ser271/272 site was important for the capacity of FIT to localize to NBs.

Figure 3. FIT was present in homodimeric protein complexes in NBs, dependent on Ser271/272 site.

Figure 4. FIT was present in heterodimeric protein complexes with bHLH039 in NBs, dependent on Ser271/272 site.

Figure 5. Two NB markers and splicing components were present in NBs (designated type II), revealing the dynamics of FIT to accumulate in NBs.

Figure 6. Three NB markers and speckle components became localized in FIT NBs and colocalized fully with FIT (designated type III), suggesting that FIT NBs have speckle function.

Figure 7. FIT colocalized with photobody (PB) markers in distinct PBs.

Figure 8. Root iron reductase activity is promoted under blue light.

Figure 9. Induction of FIT target gene expression is enhanced under blue light.

Figure 10. Schematic summary models illustrating the dynamics of FIT NB formation, suggesting that FIT NBs are related to transcriptional and posttranscriptional regulation in speckles.

Supplemental Data

Supplemental Figure S1. FIT NBs induced by blue light and a standardized FIT NB analysis procedure was developed to analyze the characteristics and dynamics of FIT NBs. (Supports Figure 1)

Supplemental Figure S2. An intrinsically disordered region, IDR$^{Ser271/272}$, is present in the FIT C-terminus and disrupted in the FITmSS271AA mutant. (Supports Figure 2, 3, and 4)

Supplemental Figure S3. FIT NBs did not colocalize with Cajal body components (designated type I). (Supports Figure 5 and 6)

Supplemental Figure S4. Type II and III NB markers are similarly localized upon single expression as upon co-expression with FIT, except PININ. (Supports Figure 5 and 6)

Supplemental Figure S5. PB markers are similarly localized upon single expression and upon co-expression with FIT. (Supports Figure 7)

Supplemental Table S1. List of primers used in this study.

Supplemental Table S2. List of vectors used in this study.

Supplemental Movie S1. Light induction triggers the formation of NBs with FIT and
FITmSS271AA with different dynamics. (Supports Figure 1 and 2)

Acknowledgements

We thank Elke Wieneke and Monique Eutebach (Institute of Botany, Heinrich-
Heine-University, Düsseldorf, Germany) for excellent technical assistance. We
acknowledge the contribution of the student Daniel Torno. We thank Rebecca C. Burkart
(Institute of Developmental Genetics, Heinrich-Heine-University, Düsseldorf, Germany),
Sebastian Hänsch and Stefanie Weidtkamp-Peters (Center for Advanced Imaging,
Heinrich-Heine-University, Düsseldorf, Germany) for help and advice with microscopy and
data analysis. K.T. is an associate member of the DFG-funded International Graduate
School for Plant Science iGRAD*plant* (IRTG 2466 "Network, exchange, and training
program to understand plant resource allocation"), Düsseldorf, Germany.

Funding

This work was funded by Deutsche Forschungsgemeinschaft (DFG, German
Research Foundation) under GRK F020512056 (NEXT*plant*), SFB 1208 project B05, and
Germany´s Excellence Strategy – EXC-2048/1 – project ID 390686111. Funding for
instrumentation: Zeiss ELYRA PS: DFG- INST 208/613-1 FUGG; Zeiss LSM780 + 4-
channel FLIM extension (Picoquant): DFG- INST 208/551-1 FUGG; Olympus FV3000
Confocal Laser Scanning Microscope with 4-channel FLIM extension (PicoQuant
rapidFLIM): DFG- INST 1358/44-1 FUGB.

K.T., R.I., Y.S., P.B. and T.B. designed the research; K.T. performed research;
K.T., R.I., Y.S., and T.B. analyzed data; R.G. contributed key materials; K.T. wrote the
paper; P.B. acquired funding; all authors reviewed and edited the article.

References

Abdallah, H. Ben and Bauer, P. (2016). Quantitative reverse transcription-qPCR-based

919   gene expression analysis in plants. Methods Mol. Biol. **479**: 9–24.

Ali, G.S., Golovkin, M., and Reddy, A.S.N. (2003). Nuclear localization and in vivo

921   dynamics of a plant-specific serine/arginine-rich protein. Plant J. **36**: 883–893.

Bancaud, A., Huet, S., Rabut, G., and Ellenberg, J. (2010). Fluorescence Perturbation

923   Techniques to Study Mobility and Molecular Dynamics of Proteins in Live Cells:

924   FRAP, Photoactivation, Photoconversion, and FLIP. Cold Spring Harb. Protoc.:

925   1303–1325.

Bauer, D., Viczían, A., Kircher, S., Nobis, T., Nitschke, R., Kunkel, T., Panigrahi,

927   **K.C.S., Ádám, É., Fejes, E., Schäfer, E., and Nagy, F.** (2004). Constitutive

928   Photomorphogenesis 1 and Multiple Photoreceptors Control Degradation of

929   Phytochrome Interacting Factor 3, a Transcription Factor Required for Light

930   Signaling in Arabidopsis. Plant Cell **16**: 1433–1445.

Bauer, P., Ling, H.Q., and Guerinot, M. Lou (2007). FIT, the FER-LIKE IRON

932   DEFICIENCY INDUCED TRANSCRIPTION FACTOR in Arabidopsis. Plant Physiol.

933   Biochem. **45**: 260–261.

Bi, Y. et al. (2021). Arabidopsis ACINUS is O-glycosylated and regulates transcription

935   and alternative splicing of regulators of reproductive transitions. Nat. Commun. **12**:

936   1–13.

Bleckmann, A., Weidtkamp-Peters, S., Seidel, C.A.M., and Simon, R. (2010). Stem

938   Cell Signaling in Arabidopsis Requires CRN to Localize CLV2 to the Plasma

939   Membrane. Plant Physiol. **152**: 166–176.

Boija, A. et al. (2018). Transcription Factors Activate Genes through the Phase-

941   Separation Capacity of Their Activation Domains. Cell **175**: 1842–1855.

Borst, J.W. and Visser, A.J.W.G. (2010). Fluorescence lifetime imaging microscopy in

943   life sciences. Meas. Sci. Technol. **21**: 1–21.

Bracha, D., Walls, M.T., Wei, M.T., Zhu, L., Kurian, M., Avalos, J.L., Toettcher, J.E.,

945   **and Brangwynne, C.P.** (2018). Mapping Local and Global Liquid Phase Behavior

946   in Living Cells Using Photo-Oligomerizable Seeds. Cell **175**: 1467–1480.

Brumbarova, T. and Ivanov, R. (2019). The Nutrient Response Transcriptional

948   Regulome of Arabidopsis. iScience **19**: 358–368.

Burkart, R.C., Strotmann, V.I., Kirschner, G.K., Akinci, A., Czempik, L., Dolata, A.,

950   **Maizel, A., Weidtkamp-Peters, S., and Stahl, Y.** (2022). PLETHORA-WOX5

951   interaction and subnuclear localization control Arabidopsis root stem cell

952   maintenance. EMBO Rep. **23**: 1–18.

953   **Van Buskirk, E.K., Reddy, A.K., Nagatani, A., and Chen, M.** (2014). Photobody

954   Localization of Phytochrome B Is Tightly Correlated with Prolonged and Light-

955   Dependent Inhibition of Hypocotyl Elongation in the Dark. Plant Physiol. **165**: 595–

956   607.

957   **Chen, D., Lyu, M., Kou, X., Li, J., Yang, Z., Gao, L., Li, Y., Fan, L., Shi, H., and**

958   **Zhong, S.** (2022). Integration of light and temperature sensing by liquid-liquid

959   phase separation of phytochrome B. Mol. Cell **82**: 1–15.

960   **Clough, S.J. and Bent, A.F.** (1998). Floral dip: a simplified method for Agrobacterium-

961   mediated transformation of Arabidopsis thaliana. Plant J. **16**: 735–743.

962   **Colangelo, E.P. and Guerinot, M. Lou** (2004). The Essential Basic Helix-Loop-Helix

963   Protein FIT1 Is Required for the Iron Deficiency Response. Plant Cell **16**: 3400–

964   3412.

965   **Collier, S., Pendle, A., Boudonck, K., van Rij, T., Dolan, L., and Shaw, P.** (2006). A

966   Distant Coilin Homologue Is Required for the Formation of Cajal Bodies in

967   Arabidopsis. Mol. Biol. Cell **17**: 2942–2951.

968   **Cui, Y., Chen, C.L., Cui, M., Zhou, W.J., Wu, H.L., and Ling, H.Q.** (2018). Four IVa

969   bHLH Transcription Factors Are Novel Interactors of FIT and Mediate JA Inhibition

970   of Iron Uptake in Arabidopsis. Mol. Plant **11**: 1166–1183.

971   **Eide, D., Broderius, M., Fett, J., and Guerinot, M. Lou** (1996). A novel iron-regulated

972   metal transporter from plants identified by functional expression in yeast. Proc. Natl.

973   Acad. Sci. **93**: 5624–5628.

974   **Emenecker, R.J., Holehouse, A.S., and Strader, L.C.** (2020). Emerging Roles for

975   Phase Separation in Plants. Dev. Cell **55**: 69–83.

976   **Emenecker, R.J., Holehouse, A.S., and Strader, L.C.** (2021). Sequence determinants

977   of in cell condensate morphology, dynamics, and oligomerization as measured by

978   number and brightness analysis. Cell Commun. Signal. **19**: 1–15.

979   **Fanara, S., Schloesser, M., Hanikenne, M., and Motte, P.** (2022). Altered metal

980   distribution in the sr45-1 Arabidopsis mutant causes developmental defects. Plant J.

981   **110**: 1332–1352.

982   **Fang, X., Wang, L., Ishikawa, R., Li, Y., Fiedler, M., Liu, F., Calder, G., Rowan, B.,**

983  **Weigel, D., Li, P., and Dean, C.** (2019). Arabidopsis FLL2 promotes liquid–liquid

984  phase separation of polyadenylation complexes. Nature **569**: 265–269.

985 **Galganski, L., Urbanek, M.O., and Krzyzosiak, W.J.** (2017). Nuclear speckles:

986  Molecular organization, biological function and role in disease. Nucleic Acids Res.

987  **45**: 10350–10368.

988 **Gao, F., Robe, K., Bettembourg, M., Navarro, N., Rofidal, V., Santoni, V., Gaymard,**

989  **F., Vignols, F., Roschzttardtz, H., Izquierdo, E., and Dubos, C.** (2020). The

990  Transcription Factor bHLH121 Interacts with bHLH105 (ILR3) and Its Closest

991  Homologs to Regulate Iron Homeostasis in Arabidopsis. Plant Cell **32**: 508–524.

992 **Gao, Y.-Q., Bu, L.-H., Han, M.-L., Wang, Y.-L., Li, Z.-Y., Liu, H.-T., and Chao, D.-Y.**

993  (2021). Long-distance blue light signalling regulates phosphate deficiency-induced

994  primary root growth inhibition. Mol. Plant **14**: 1–15.

995 **Gratz, R. et al.** (2019). CIPK11-Dependent Phosphorylation Modulates FIT Activity to

996  Promote Arabidopsis Iron Acquisition in Response to Calcium Signaling. Dev. Cell

997  **48**: 726–740.

998 **Gratz, R., Brumbarova, T., Ivanov, R., Trofimov, K., Tünnermann, L., Ochoa-**

999  **Fernandez, R., Blomeier, T., Meiser, J., Weidtkamp-Peters, S., Zurbriggen, M.,**

1000  **and Bauer, P.** (2020). Phospho-mutant activity assays provide evidence for

1001  alternative phospho-regulation pathways of the transcription factor FER-LIKE IRON

1002  DEFICIENCY-INDUCED TRANSCRIPTION FACTOR. New Phytol. **225**: 250–267.

1003 **Guo, Z., Xu, J., Wang, Y., Hu, C., Shi, K., Zhou, J., Xia, X., Zhou, Y., Foyer, C.H., and**

1004  **Yu, J.** (2021). The phyB-dependent induction of HY5 promotes iron uptake by

1005  systemically activating FER expression. EMBO Rep. **22**: 1–15.

1006 **Huang, X., Xiao, N., Zou, Y., Xie, Y., Tang, L., Zhang, Y., Yu, Y., Li, Y., and Xu, C.**

1007  (2022). Heterotypic transcriptional condensates formed by prion-like paralogous

1008  proteins canalize flowering transition in tomato. Genome Biol. **23**: 1–21.

1009 **Jakoby, M., Wang, H.Y., Reidt, W., Weisshaar, B., and Bauer, P.** (2004). FRU

1010  (BHLH029) is required for induction of iron mobilization genes in Arabidopsis

1011  thaliana. FEBS Lett. **577**: 528–534.

1012 **Jung, J.-H. et al.** (2020). A prion-like domain in ELF3 functions as a thermosensor in

1013  Arabidopsis. Nature **585**: 256–260.

1014 **Kaiserli, E., Páldi, K., O'Donnell, L., Batalov, O., Pedmale, U. V., Nusinow, D.A.,**

1015  **Kay, S.A., and Chory, J.** (2015). Integration of Light and Photoperiodic Signaling in

Transcriptional Nuclear Foci. Dev. Cell **35**: 311–321.

Kalinina, N.O., Makarova, S., Makhotenko, A., Love, A.J., and Taliansky, M. (2018).

The Multiple Functions of the Nucleolus in Plant Development, Disease and Stress

Responses. Front. Plant Sci. **9**: 1–19.

Kammel, C., Thomaier, M., Sørensen, B.B., Schubert, T., Längst, G., Grasser, M.,

and Grasser, K.D. (2013). Arabidopsis DEAD-Box RNA Helicase UAP56 Interacts

with Both RNA and DNA as well as with mRNA Export Factors. PLoS One **8**: 1–12.

Kanwar, P., Baby, D., and Bauer, P. (2021). Interconnection of iron and osmotic stress

signalling in plants: is FIT a regulatory hub to cross-connect abscisic acid

responses? Plant Biol. **23**: 31–38.

Kim, D.W., Jeon, S.J., Hwang, S.M., Hong, J.C., and Bahk, J.D. (2016). The C3H-

type zinc finger protein GDS1/C3H42 is a nuclear-speckle-localized protein that is

essential for normal growth and development in Arabidopsis. Plant Sci. **250**: 141–

153.

Kim, S.A., Lacroix, I.S., Gerber, S.A., and Guerinot, M. Lou (2019). The iron

deficiency response in Arabidopsis thaliana requires the phosphorylated

transcription factor URI. Proc. Natl. Acad. Sci. **116**: 24933–24942.

Kircher, S., Gil, P., Kozma-Bognár, L., Fejes, E., Speth, V., Husselstein-Muller, T.,

Bauer, D., Ádám, É., Schäfer, E., and Nagy, F. (2002). Nucleocytoplasmic

Partitioning of the Plant Photoreceptors Phytochrome A, B, C, D, and E Is

Regulated Differentially by Light and Exhibits a Diurnal Rhythm. Plant Cell **14**:

1541–1555.

Korshunova, Y.O., Eide, D., Clark, W.G., Guerinot, M. Lou, and Himadri, B. (1999).

The IRT1 protein from Arabidopsis thaliana is a metal transporter with a broad

substrate range. Plant Mol. Biol. **40**: 37–44.

Lafontaine, D.L.J., Riback, J.A., Bascetin, R., and Brangwynne, C.P. (2021). The

nucleolus as a multiphase liquid condensate. Nat. Rev. Mol. Cell Biol. **22**: 165–182.

Lampropoulos, A., Sutikovic, Z., Wenzl, C., Maegele, I., Lohmann, J.U., and Forner,

J. (2013). GreenGate - A Novel, Versatile, and Efficient Cloning System for Plant

Transgenesis. PLoS One **8**: 1–15.

Le, C.T.T., Brumbarova, T., Ivanov, R., Stoof, C., Weber, E., Mohrbacher, J., Fink-

Straube, C., and Bauer, P. (2016). ZINC FINGER OF ARABIDOPSIS

THALIANA12 (ZAT12) Interacts with FER-LIKE IRON DEFICIENCY-INDUCED

1049  TRANSCRIPTION FACTOR (FIT) Linking Iron Deficiency and Oxidative Stress

1050  Responses. Plant Physiol. **170**: 540–557.

1051 **Leivar, P. and Monte, E.** (2014). PIFs: Systems Integrators in Plant Development. Plant

1052  Cell **26**: 56–78.

1053 **Leonelli, L., Erickson, E., Lyska, D., and Niyogi, K.K.** (2016). Transient expression in

1054  Nicotiana benthamiana for rapid functional analysis of genes involved in non-

1055  photochemical quenching and carotenoid biosynthesis. Plant J. **88**: 375–386.

1056 **Li, W., Lin, W.-D., Ray, P., Lan, P., and Schmidt, W.** (2013). Genome-Wide Detection

1057  of Condition-Sensitive Alternative Splicing in Arabidopsis Roots. Plant Physiol. **162**:

1058  1750–1763.

1059 **Li, X., Zhang, H., Ai, Q., Liang, G., and Yu, D.** (2016). Two bHLH Transcription

1060  Factors, bHLH34 and bHLH104, Regulate Iron Homeostasis in Arabidopsis

1061  thaliana. Plant Physiol. **170**: 2478–2493.

1062 **Liang, G., Zhang, H., Li, X., Ai, Q., and Yu, D.** (2017). bHLH transcription factor

1063  bHLH115 regulates iron homeostasis in Arabidopsis thaliana. J. Exp. Bot. **68**:

1064  1743–1755.

1065 **Lin, C.L., Leu, S., Lu, M.C., and Ouyang, P.** (2004). Over-expression of SR-cyclophilin,

1066  an interaction partner of nuclear pinin, releases SR family splicing factors from

1067  nuclear speckles. Biochem. Biophys. Res. Commun. **321**: 638–647.

1068 **Lingam, S., Mohrbacher, J., Brumbarova, T., Potuschak, T., Fink-Straube, C.,**

1069  **Blondet, E., Genschik, P., and Bauer, P.** (2011). Interaction between the bHLH

1070  Transcription Factor FIT and ETHYLENE INSENSITIVE3/ETHYLENE

1071  INSENSITIVE3-LIKE1 Reveals Molecular Linkage between the Regulation of Iron

1072  Acquisition and Ethylene Signaling in Arabidopsis. Plant Cell **23**: 1815–1829.

1073 **Lorković, Z.J., Hilscher, J., and Barta, A.** (2008). Co-localisation studies of

1074  Arabidopsis SR splicing factors reveal different types of speckles in plant cell nuclei.

1075  Exp. Cell Res. **314**: 3175–3186.

1076 **Lorković, Z.J., Hilscher, J., and Barta, A.** (2004). Use of Fluorescent Protein Tags to

1077  Study Nuclear Organization of the Spliceosomal Machinery in Transiently

1078  Transformed Living Plant Cells. Mol. Biol. Cell **15**: 3233–3243.

1079 **Love, A.J., Yu, C., Petukhova, N. V., Kalinina, N.O., Chen, J., and Taliansky, M.E.**

1080  (2017). Cajal bodies and their role in plant stress and disease responses. RNA Biol.

1081  **14**: 779–790.

Mankotia, S., Singh, D., Monika, K., Kalra, M., Meena, H., Meena, V., Yadav, R.K.,
Pandey, A.K., and Satbhai, S.B. (2023). ELONGATED HYPOCOTYL 5 regulates
BRUTUS and affects iron acquisition and homeostasis in Arabidopsis thaliana.
Plant J.: 1–18.

Martin, K., Kopperud, K., Chakrabarty, R., Banerjee, R., Brooks, R., and Goodin,
M.M. (2009). Transient expression in Nicotiana benthamiana fluorescent marker
lines provides enhanced definition of protein localization, movement and
interactions in planta. Plant J. **59**: 150–162.

Meiser, J., Lingam, S., and Bauer, P. (2011). Posttranslational Regulation of the Iron
Deficiency Basic Helix-Loop-Helix Transcription Factor FIT Is Affected by Iron and
Nitric Oxide. Plant Physiol. **157**: 2154–2166.

Meyer, H.M. (2020). In search of function: nuclear bodies and their possible roles as
plant environmental sensors. Curr. Opin. Plant Biol. **58**: 33–40.

Mikulski, P., Wolff, P., Lu, T., Nielsen, M., Echevarria, E.F., Zhu, D., Questa, J.I.,
Saalbach, G., Martins, C., and Dean, C. (2022). VAL1 acts as an assembly
platform co-ordinating co-transcriptional repression and chromatin regulation at
Arabidopsis FLC. Nat. Commun. **13**: 1–12.

Naranjo-Arcos, M.A., Maurer, F., Meiser, J., Pateyron, S., Fink-Straube, C., and
Bauer, P. (2017). Dissection of iron signaling and iron accumulation by
overexpression of subgroup Ib bHLH039 protein. Sci. Rep. **7**: 1–12.

Ni, M., Tepperman, J.M., and Quail, P.H. (1998). PIF3, a Phytochrome-Interacting
Factor Necessary for Normal Photoinduced Signal Transduction, Is a Novel Basic
Helix-Loop-Helix Protein. Cell **95**: 657–667.

Oh, E., Kim, J., Park, E., Kim, J.-I., Kang, C., and Choi, G. (2004). PIL5, a
Phytochrome-Interacting Basic Helix-Loop-Helix Protein, Is a Key Negative
Regulator of Seed Germination in Arabidopsis thaliana. Plant Cell **16**: 3045–3058.

Owen, I. and Shewmaker, F. (2019). The Role of Post-Translational Modifications in
the Phase Transitions of Intrinsically Disordered Proteins. Int. J. Mol. Sci. **20**: 1–14.

Pardi, S.A. and Nusinow, D.A. (2021). Out of the Dark and Into the Light: A New View
of Phytochrome Photobodies. Front. Plant Sci. **12**: 1–15.

Pedmale, U. V., Huang, S.-S.C., Zander, M., Cole, B.J., Hetzel, J., Ljung, K., Reis,
P.A.B., Sridevi, P., Nito, K., Nery, J.R., Ecker, J.R., and Chory, J. (2016).
Cryptochromes Interact Directly with PIFs to Control Plant Growth in Limiting Blue

Light. Cell **164**: 233–245.

Pendle, A.F., Cark, G.P., Boon, R., Lewandowska, D., Lam, Y.W., Andersen, J.,

Mann, M., Lamond, A.I., Brown, J.W.S., and Shaw, P.J. (2005). Proteomic

Analysis of the Arabidopsis Nucleolus Suggests Novel Nucleolar Functions. Mol.

Biol. Cell **16**: 260–269.

Pham, V.N., Kathare, P.K., and Huq, E. (2018). Phytochromes and Phytochrome

Interacting Factors. Plant Physiol. **176**: 1025–1038.

Powers, S.K. et al. (2019). Nucleo-cytoplasmic Partitioning of ARF Proteins Controls

Auxin Responses in Arabidopsis thaliana. Mol. Cell **76**: 177–190.

Qiu, Y., Li, M., Kim, R.J.A., Moore, C.M., and Chen, M. (2019). Daytime temperature is

sensed by phytochrome B in Arabidopsis through a transcriptional activator

HEMERA. Nat. Commun. **10**: 1–13.

Qiu, Y., Pasoreck, E.K., Yoo, C.Y., He, J., Wang, H., Bajracharya, A., Li, M., Larsen,

H.D., Cheung, S., and Chen, M. (2021). RCB initiates Arabidopsis

thermomorphogenesis by stabilizing the thermoregulator PIF4 in the daytime. Nat.

Commun. **12**: 1–13.

Reddy, A.S.N., Day, I.S., Gohring, J., and Barta, A. (2012). Localization and Dynamics

of Nuclear Speckles in Plants. Plant Physiol. **158**: 67–77.

Riback, J.A., Zhu, L., Ferrolino, M.C., Tolbert, M., Mitrea, D.M., Sanders, D.W., Wei,

M.-T., Kriwacki, R.W., and Brangwynne, C.P. (2020). Composition-dependent

thermodynamics of intracellular phase separation. Nature **581**: 209–214.

Robinson, N.J., Procter, C.M., Connolly, E.L., and Guerinot, M. Lou (1999). A ferric-

chelate reductase for iron uptake from soils. Nature **397**: 694–697.

Römheld, V. and Marschner, H. (1986). Mobilization of iron in the rhizosphere of

different plant species. Adv. Plant Nutr. **2**: 155–204.

Salladini, E., Jørgensen, M.L.M., Theisen, F.F., and Skriver, K. (2020). Intrinsic

Disorder in Plant Transcription Factor Systems: Functional Implications. Int. J. Mol.

Sci. **21**: 1–35.

Salomé, P.A., Oliva, M., Weigel, D., and Krämer, U. (2013). Circadian clock

adjustment to plant iron status depends on chloroplast and phytochrome function.

EMBO J. **32**: 511–523.

Santi, S. and Schmidt, W. (2009). Dissecting iron deficiency-induced proton extrusion

in Arabidopsis roots. New Phytol. **183**: 1072–1084.

Schwarz, B. and Bauer, P. (2020). FIT, a regulatory hub for iron deficiency and stress
signaling in roots, and FIT-dependent and -independent gene signatures. J. Exp.
Bot. **71**: 1694–1705.

Shamloul, M., Trusa, J., Mett, V., and Yusibov, V. (2014). Optimization and Utilization
of Agrobacterium-mediated Transient Protein Production in Nicotiana. J. Vis. Exp.
86: 1–13.

Shin, Y., Berry, J., Pannucci, N., Haataja, M.P., Toettcher, J.E., and Brangwynne,
C.P. (2017). Spatiotemporal Control of Intracellular Phase Transitions Using Light-
Activated optoDroplets. Cell **168**: 159–171.

Stahl, Y. et al. (2013). Moderation of Arabidopsis root Stemness by CLAVATA1 and
ARABIDOPSIS CRINKLY4 Receptor Kinase Complexes. Curr. Biol. **23**: 362–371.

Strader, L., Weijers, D., and Wagner, D. (2022). Plant transcription factors - being in
the right place with the right company. Curr. Opin. Plant Biol. **65**: 1–8.

Tarczewska, A. and Greb-Markiewicz, B. (2019). The Significance of the Intrinsically
Disordered Regions for the Functions of the bHLH Transcription Factors. Int. J. Mol.
Sci. **20**: 1–20.

Trinkle-Mulcahy, L. and Sleeman, J.E. (2017). The Cajal body and the nucleolus: "In a
relationship" or "It's complicated"? RNA Biol. **14**: 739–751.

Trofimov, K., Ivanov, R., Eutebach, M., Acaroglu, B., Mohr, I., Bauer, P., and
Brumbarova, T. (2019). Mobility and localization of the iron deficiency-induced
transcription factor bHLH039 change in the presence of FIT. Plant Direct **3**: 1–11.

Vert, G.A., Briat, J.F., and Curie, C. (2003). Dual Regulation of the Arabidopsis High-
Affinity Root Iron Uptake System by Local and Long-Distance Signals. Plant
Physiol. **132**: 796–804.

Wang, B., Zhang, H., Huai, J., Peng, F., Wu, J., Lin, R., and Fang, X. (2022).
Condensation of SEUSS promotes hyperosmotic stress tolerance in Arabidopsis.
Nat. Chem. Biol. **18**: 1361–1369.

Wang, H.Y., Klatte, M., Jakoby, M., Bäumlein, H., Weisshaar, B., and Bauer, P.
(2007). Iron deficiency-mediated stress regulation of four subgroup Ib BHLH genes
in Arabidopsis thaliana. Planta **226**: 897–908.

Wang, N., Cui, Y., Liu, Y., Fan, H., Du, J., Huang, Z., Yuan, Y., Wu, H., and Ling,
H.Q. (2013). Requirement and Functional Redundancy of Ib Subgroup bHLH
Proteins for Iron Deficiency Responses and Uptake in Arabidopsis thaliana. Mol.

Plant **6**: 503–513.

Wang, X., Jiang, B., Gu, L., Chen, Y., Mora, M., Zhu, M., Noory, E., Wang, Q., and Lin, C. (2021). A photoregulatory mechanism of the circadian clock in Arabidopsis. Nat. Plants **7**: 1397–1408.

Wedepohl, K.H. (1995). The composition of the continental crust. Geochim. Cosmochim. Acta **59**: 1217–1232.

Weidtkamp-Peters, S., Rehwald, S., and Stahl, Y. (2022). Homo-FRET Imaging to Study Protein–Protein Interaction and Complex Formation in Plants. In Plant Synthetic Biology: Methods and Protocols, M.D. Zurbriggen, ed (Springer US: New York, NY), pp. 197–208.

Weidtkamp-Peters, S. and Stahl, Y. (2017). The Use of FRET/FLIM to Study Proteins Interacting with Plant Receptor Kinases. In Plant Receptor Kinases: Methods and Protocols, R.B. Aalen, ed (Springer New York: New York, NY), pp. 163–175.

Wild, M., Davière, J.M., Regnault, T., Sakvarelidze-Achard, L., Carrera, E., Lopez Diaz, I., Cayrel, A., Dubeaux, G., Vert, G., and Achard, P. (2016). Tissue-Specific Regulation of Gibberellin Signaling Fine-Tunes Arabidopsis Iron-Deficiency Responses. Dev. Cell **37**: 190–200.

Yang, Y., Ou, B., Zhang, J., Si, W., Gu, H., Qin, G., and Qu, L.-J. (2014). The Arabidopsis Mediator subunit MED16 regulates iron homeostasis by associating with EIN3/EIL1 through subunit MED25. Plant J. **77**: 838–851.

Yi, Y. and Guerinot, M. Lou (1996). Genetic evidence that induction of root Fe(III) chelate reductase activity is necessary for iron uptake under iron deficiency. Plant J. **10**: 835–844.

Yuan, Y., Wu, H., Wang, N., Li, J., Zhao, W., Du, J., Wang, D., and Ling, H.Q. (2008). FIT interacts with AtbHLH38 and AtbHLH39 in regulating iron uptake gene expression for iron homeostasis in Arabidopsis. Cell Res. **18**: 385–397.

Yuan, Y.X., Zhang, J., Wang, D.W., and Ling, H.Q. (2005). AtbHLH29 of Arabidopsis thaliana is a functional ortholog of tomato FER involved in controlling iron acquisition in strategy I plants. Cell Res. **15**: 613–621.

Zhang, J., Liu, B., Li, M., Feng, D., Jin, H., Wang, P., Liu, J., Xiong, F., Wang, J., and Wang, H. (2015). The bHLH Transcription Factor bHLH104 Interacts with IAA-LEUCINE RESISTANT3 and Modulates Iron Homeostasis in Arabidopsis. Plant Cell **27**: 787–805.

1214 **Zhang, Y., Wu, H., Wang, N., Fan, H., Chen, C., Cui, Y., Liu, H., and Ling, H.-Q.**
1215   (2014). Mediator subunit 16 functions in the regulation of iron uptake gene
1216   expression in Arabidopsis. New Phytol. **203**: 770–783.
1217 **Zhu, P., Lister, C., and Dean, C.** (2021). Cold-induced Arabidopsis FRIGIDA nuclear
1218   condensates for FLC repression. Nature **599**: 657–661.

1219
1220
1221 **Figures and figure legends**
1222

Figure 1: FIT accumulated in nuclear condensates, termed FIT nuclear bodies (NBs) in a light-inducible manner, most likely following liquid-liquid phase separation (LLPS).

A, Induction of FIT NBs in Arabidopsis root epidermis cells of the root differentiation zone. Left, light microscopy overview image of a 5-d-old Arabidopsis seedling (proFIT:FIT-GFP) grown under iron deficiency. Right, nuclear localization of FIT-GFP in the root epidermis cells as indicated in the overview image, at t=0, t=90, and t=120 min. FIT-GFP signals were evenly distributed in the nucleus at t=0 min, and after induction by excitation with 488 nm laser NB formation accumulated in NBs at t=90 min, then disappeared as shown at t=120 min. Note that root epidermis cells developed few NBs, sometimes taking up to two hours to appear. Three independent experiments with three plants were conducted. In the indicated region of interest, approximately a quarter of the root epidermis cells showed NBs. A representative image from one nucleus is shown. B-D, Fluorescence protein analysis in transiently transformed *N. benthamiana* leaf epidermis cells. Confocal images of B, FIT-GFP, C, FIT-mCherry, and D, ZAT12-GFP at t=0 and t=5 min. At t=0 min, FIT-GFP and FIT-mCherry showed an even distribution within the nucleus. Following a 488 nm laser excitation, numerous NBs were clearly visible in all examined transformed cells at t=5 min. These NBs were termed FIT NBs. Under the same imaging conditions, ZAT12-GFP did not show NB formation. According to these results, a standardized FIT NB analysis procedure was set up (**Supplemental Figure S1B**). See also **Supplemental Movie S1A-C**. Representative images from two to three independent experiments. E-G, FRAP measurements to test for liquid-like behavior of FIT NBs, using the standardized FIT NB analysis procedure in transiently transformed *N. benthamiana* leaf epidermis cells. E, Representative images of the fluorescent signal during a FRAP experiment, taken before bleaching (0 s) and recovery of fluorescence at three time points after bleaching from 3 s to 45 s within the circled region of a NB. F, Line diagram representing the relative fluorescence during a FRAP measurement for 10 NBs, showing a high fluorescence recovery rate of FIT-GFP within NBs. Dark green line, mean value; light green filled area, variation. G, Box plot diagram representing the mobile fraction of FIT-GFP calculated based on the relative fluorescence recovery in F. The diagram indicates high mobility of FIT. The mean was calculated from 10 NBs from 10 nuclei from a transformed plant. Three independent experiments were conducted, one representative result is shown. H, Box plot diagram representing quantification of the FIT NB shape with

the software ImageJ (National Institutes of Health), indicating that FIT NBs have circular
shape. Mobility and circularity characteristics indicate that FIT NBs are most likely liquid
condensates that are the result of LLPS. The mean was calculated from all NBs visible in
15 nuclei from a transformed plant. Two independent experiments were conducted, one
representative result is shown.
Box plots show 25-75 percentile with min-max whiskers, mean as small square and
median as line. Scale bars of nuclei images, 2 µm; scale bar full seedling, 1 mm.
Arrowheads indicate NBs. G = GFP; C = mCherry.

Figure 2: The FIT C-terminal Ser271/272 site was important for the capacity of FIT to localize to NBs.

A, Confocal images of nuclear localization of FITmSS271AA-GFP at t=0 and t=15 min. FITmSS271AA-GFP accumulated in NBs, but NB formation required a longer time compared to FIT-GFP. See also **Supplemental Movie S1, A and C**. Two independent experiments. Representative images from one nucleus. B, Bar diagrams showing in B, number of NBs, and in C, the sizes of NBs with FIT-GFP and FITmSS271AA-GFP at t=5/15 min. NB number and size were determined with the software ImageJ (National Institute of Health). FIT-GFP accumulated in more and larger NBs than FITmSS271AA-GFP. See **Supplemental Movie S1, A and C**. FITmSS271AA-GFP lacks IDR$^{Ser271/272}$. This IDR may be relevant for FIT NB formation (**Supplemental Figure S2**). In B and C, bar diagrams represent the mean and standard deviation for a quantification of 15 nuclei from a transformed plant (n = 15). Two experiments were conducted, one representative result is shown.

Statistical analysis was performed with the Mann-Whitney test. Different letters indicate statistically significant differences (P < 0.05). Scale bar: 2 µm. Arrowheads indicate NBs. G = GFP. Analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure.

Figure 3: FIT was present in homodimeric protein complexes in NBs, dependent on Ser271/272 site.

Anisotropy (or homo-FRET) measurements of FIT-GFP and FITmSS271AA-GFP to determine homodimerization strength. A, Schematic illustration of the anisotropy principle. Energy transfer between the same kind of fluorescently tagged proteins leads to depolarization of the emitted light. Extent of the depolarization gives a hint on dimerization and oligomerization of a protein as the fluorescence anisotropy (FA) value decreases. B, Representative images showing color-coded FA values of FIT-GFP and FITmSS271AA at t=0 and t=5/15 min. C-D, Box plots representing quantification of FA values. FA was measured at t=0 within the whole nucleus and at t =5/15 min within the whole nucleus, in NBs and in residual NP. Free GFP and GFP-GFP served as references for mono- and dimerization. FA values for C, FIT-GFP, and D, FITmSS271AA-GFP. In C and D, FA values were calculated from 10-15 nuclei from a transformed plant (n = 10-15). Two experiments were conducted, one representative result is shown. C and D show the same free GFP and GFP-GFP references because both measurements were performed on the same day. FA values decreased for FIT-GFP, but not FITmSS271AA-GFP, in the whole nucleus (compare t=0 with t=5/15 min). FA values were also lowered in NBs versus NP in the case of FIT-GFP but not FITmSS271AA-GFP (compare t=5/15 min of NBs and NP). This indicates stronger homodimerization of FIT than FITmSS271AA-GFP in the whole nucleus and in NBs. IDR$^{Ser271/272}$ may therefore be relevant for FIT NB formation and FIT homodimerization (**Supplemental Figure S2**).

Box plots show 25-75 percentile with min-max whiskers, mean as small square and median as line. Statistical analysis was performed with one-way ANOVA and Tukey post-hoc test. Different letters indicate statistically significant differences ($P < 0.05$). Scale bar: 2 µm. Arrowheads indicate NBs. G = GFP. Fluorescence protein analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure.

Figure 4: FIT was present in heterodimeric protein complexes with bHLH039 in NBs, dependent on Ser271/272 site.

A, Confocal images with colocalization of FIT-GFP and bHLH039-mCherry in the nucleus. Both proteins were evenly distributed within the nucleus at t=0 and colocalized fully in FIT NBs at t=5 min. Two independent experiments with two plants each. Representative images from one nucleus. B-E, FRET-FLIM measurements to determine heterodimerization strength of FIT and FITmSS271AA with bHLH039, respectively. FIT-GFP and FITmSS271AA-GFP (donor only) served as negative controls. B, Schematic illustration of the FRET-FLIM principle. Energy transfer occurs between two different fluorophores. One fluorophore acts as the donor and the other as the acceptor of the energy. In case of interaction (close proximity, ≤10 nm) the fluorescence lifetime of the donor decreases. C, Representative images showing color-coded fluorescence lifetime values of FIT-GFP and FITmSS271AA-GFP co-expressed with bHLH039-mCherry at t=0 and t=5/15 min. D-E, Box plots diagrams representing FRET-FLIM measurements at t=0 within the whole nucleus and at t=5/15 min within the whole nucleus, inside NBs and in residual NP. Lifetime values represent measurements of 10 nuclei from a transformed plant (n = 10). Two experiments were conducted, one representative result is shown. Fluorescence lifetime was reduced for the pair of FIT-GFP and bHLH039-mCherry in NBs versus NP at t=5 min, indicating protein interaction preferentially inside NBs. Fluorescence lifetime values were not significantly different for the pair FITmSS271AA-GFP and bHLH039-mCherry in this same comparison at t=15 min, indicating that this pair did not preferentially interact in NBs. IDR[Ser271/272] may therefore be relevant for FIT NB formation, and FIT homo- and heterodimerization (**Supplemental Figure S2**).

Box plots show 25-75 percentile with min-max whiskers, mean as small square and median as line. Statistical analysis was performed with one-way ANOVA and Tukey post-hoc test. Different letters indicate statistically significant differences ($P < 0.05$). Scale bar: 2 µm. Arrowheads indicate NBs. G = GFP; C = mCherry. Fluorescence protein analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure.

Figure 5: Two NB markers and splicing components were present in NBs (designated type II), revealing the dynamics of FIT to accumulate in NBs.

Confocal images showing localization of FIT-GFP and NB markers (type II) upon co-expression in the nucleus at t=0 and t=5 min. Co-expression of FIT-GFP with A, SR45-mRFP, and B, SRm102-mRFP. Type II NB markers localized inside NBs at t=0 and t=5 min. Similar localization patterns were observed upon single expression, showing that SR45 and SRm102 are present in distinct NB types (compare with **Supplemental Figure S4, A and B**). FIT-GFP colocalized with type II markers in their distinct NBs at t=5 min, but not t=0. FIT-GFP additionally localized in FIT NBs at t=5 min. Type II markers were not present in FIT NBs, while FIT-GFP became recruited into the distinct type II NBs upon the light trigger. Hence, FIT NBs could be associated with speckle components.

Scale bar: 2 µm. Filled arrowheads indicate colocalization in NBs, empty arrowheads indicate no colocalization in NBs. G = GFP; R = mRFP. Fluorescence protein analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure. Representative images from two to five independent experiments are shown. For data with type I markers (no colocalization) and type III markers (full colocalization) see **Supplemental Figure S3** and **Figure 6**.

Figure 6: Three NB markers and speckle components became localized in FIT NBs

and colocalized fully with FIT (designated type III), suggesting that FIT NBs have

speckle function.

Confocal images showing localization of FIT-GFP and NB markers (type III) upon

co-expression in the nucleus at t=0 and t=5 min. Co-expression of FIT-GFP with A,

UAP56H2-mRFP, B, P15H1-mRFP, and C, PININ-mRFP. All three type III NB markers

were homogeneously distributed and colocalized with FIT-GFP in the nucleus at t=0, while

they colocalized with FIT-GFP in FIT NBs at t=5 min. UAP56H2-mRFP and P15H1-mRFP

showed homogeneous localization in the single expression at both t=0 and t=5 min

(compare with **Supplemental Figure S4, C and D**), while PININ-mRFP localized mainly

in one large and several small NBs upon single expression at t=0 and t=5 min (compare

with **Supplemental Figure S4E**). Hence, these three markers adopted the localization of

FIT-GFP upon co-expression and suggest that FIT NBs have a speckle function.

Scale bar: 2 µm. Arrowheads indicate colocalization within NBs. G = GFP; R =

mRFP. Fluorescence protein analysis was conducted in transiently transformed

N. benthamiana leaf epidermis cells, following the standardized FIT NB analysis

procedure. Representative images from four to seven independent experiments are

shown. For data with type I markers (no colocalization) and type II markers (partial

colocalization) see **Supplemental Figure S** and **Figure 5**.

Figure 7: FIT colocalized with photobody (PB) markers in distinct PBs.

Confocal images showing localization of FIT-GFP and PB markers upon co-expression in the nucleus at t=0 and t=15 min. Co-expression of FIT-GFP with A, PIF3-mCherry, and B and C, PIF4-mCherry, in B, showing a typical pattern with absence of NBs (ca. 50% of nuclei), in C, showing a typical pattern with presence of NBs (ca. 50% of cells). When FIT-GFP was co-expressed with PB markers, FIT NBs did not appear at t=5 min, but instead, FIT-GFP colocalized with PB markers in PBs at t=15 min. A, PIF3-mCherry localized predominantly to a single large PB at t=0 and t=15 min. FIT-GFP colocalized with PIF3-mCherry in this single large PB at t=15 min. B, PIF4-mCherry and FIT-GFP were both homogeneously distributed in the nucleoplasm at t=0 and t=15 min. In C, FIT-GFP colocalized with PIF4-mCherry in PBs at t=0 and t=15 min. The same localization patterns were found for PIF3-mCherry and PIF4-mCherry upon single expression (compare with **Supplemental Figure S5**). Hence, FIT-GFP was recruited to the two distinct types of PIF3 and PIF4 PBs, whereas PIF3 and PIF4 were not recruited to FIT NBs. This suggests that FIT NBs are affected by the presence of PIF3- and PIF4-containing PBs and a connection to light signaling exists.

Scale bar: 2 µm. Arrowheads indicate colocalization in PBs. G = GFP; C = mCherry. Fluorescence protein analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure. Representative images from four to six independent experiments are shown.

Figure 8. Root iron reductase activity is promoted under blue light.

Root iron reductase activity assay on 6-d-old Arabidopsis seedlings grown under white light for 5 d under iron deficient and iron sufficient conditions and then exposed for 1 d to blue, red, far-red light or darkness, or in parallel as control to white light. A, Induction of iron reductase activity upon iron deficiency versus sufficiency was higher under blue light than compared to the white light control. B, Exposure to red light did not change the iron reductase activity compared to the white light control. C-D, Plants exposed to far-red light and darkness did not show induction of iron reductase activity under iron deficient conditions compared to the iron deficient conditions.

Two experiments were conducted, one representative result is shown. Bar diagrams represent the mean and standard deviation of four replicates with four seedlings each (n=4). Statistical analysis was performed with one-way ANOVA and Tukey post-hoc test. Different letters indicate statistically significant differences ($P < 0.05$).

Figure 9. Induction of FIT target gene expression is enhanced under blue light.

Gene expression analysis of iron deficiency genes *FIT*, *BHLH039*, *FRO2*, and *IRT1* on 6-d-old Arabidopsis seedlings grown under white light for 5 d under iron deficient and iron sufficient conditions and then exposed for 1 d to blue, red, far-red light or darkness, or in parallel as control to white light. A-B, *FIT* and *bHLH039* gene induction in response to low iron supply did not change after blue light exposure compared to the white light control. C-D, *FRO2* and *IRT1* gene expression increased after blue light exposure in iron deficient conditions compared to respective iron sufficient condition and white light control. E, *FIT* gene expression did not change after red light exposure compared to the white light control. F-G, *BHLH039* and *FRO2* gene expression decreased after red light exposure in iron deficient conditions compared to respective iron sufficient condition and white light control. H, *IRT1* gene expression did not change after red light exposure compared to the white light control. I, *FIT* gene expression did not change after far-red light exposure compared to the white light control. J-L, *BHLH039*, *FRO2*, and *IRT1* gene expression decreased after far-red light exposure in iron deficient conditions compared to white light deficient condition, comparable to the respective iron sufficient condition and iron sufficient white light control. M, *FIT* gene expression was higher in darkness under iron deficient conditions compared to respective iron sufficient condition and white light control. N-P, *BHLH039*, *FRO2*, and *IRT1* gene expression decreased after being exposed to darkness in iron deficient conditions compared to white light deficient condition, comparable to the respective iron sufficient condition and iron sufficient white light control.

Two experiments were conducted, one representative result is shown. Bar diagrams represent the mean and standard deviation of three replicates with twelve seedlings and two technical replicates each (n=4). Statistical analysis was performed with one-way ANOVA and Tukey post-hoc test. Different letters indicate statistically significant differences (P < 0.05).

Figure 10: Summary model illustrating the effect of blue light on FIT NB formation and iron uptake, suggesting that FIT NBs are related to transcriptional and posttranscriptional regulation in speckles and that blue light has a promoting effect on iron uptake.

FIT accumulates in FIT NBs upon induction with blue light. FIT NBs are reversible, dynamic and of circular shape and may undergo LLPS. FIT homodimers and FIT-bHLH039 heterodimers are present in FIT NBs. FIT-bHLH039 is an active TF complex for upregulating the expression of iron acquisition genes in roots (Gratz et al., 2019). Hence, FIT NBs are subnuclear sites related to transcriptional regulation and because of their colocalization with speckle components, also to speckles. Blue light enhances root iron reductase activity and gene expression of *FRO2* and *IRT1*, all of which are downstream of FIT-bHLH039. In summary, FIT NBs are blue light-inducible subnuclear sites where active TF complexes of FIT and bHLH039 accumulate, linking transcriptional and posttranscriptional regulation in speckles with a promoting blue light effect on iron uptake.

Figure 1

Figure 1: FIT accumulated in nuclear condensates, termed FIT nuclear bodies (NBs) in a light-inducible manner, most likely following liquid-liquid phase separation (LLPS).

A, Induction of FIT NBs in Arabidopsis root epidermis cells of the root differentiation zone. Left, light microscopy overview image of a 5-d-old Arabidopsis seedling (proFIT:FIT-GFP) grown under iron deficiency. Right, nuclear localization of FIT-GFP in the root epidermis cells as indicated in the overview image, at t=0, t=90, and t=120 min. FIT-GFP signals were evenly distributed in the nucleus at t=0 min, and after induction by excitation with 488 nm laser NB formation accumulated in NBs at t=90 min, but also disappeared as shown at t=120 min. Note that root epidermis cells developed few NBs, sometimes taking up to two hours to appear. Three independent experiments with three plants were conducted. In the indicated region of interest, approximately a quarter of the root epidermis cells showed NBs. A representative image from one nucleus is shown. B-D, Fluorescence protein analysis in transiently transformed *N. benthamiana* leaf epidermis cells. Confocal images of B, FIT-GFP, C, FIT-mCherry, and D, ZAT12-GFP at t=0 and t=5 min. At t=0 min, FIT-GFP and FIT-mCherry showed an even distribution within the nucleus. Following a 488 nm laser excitation, numerous NBs were clearly visible in all examined transformed cells at t=5 min. These NBs were termed FIT NBs. Under the same imaging conditions, ZAT12-GFP did not show NB formation. According to these results, a standardized FIT NB analysis procedure was set up (**Supplemental Figure S1B**). See also **Supplemental Movie S1A-C**. Representative images from two to three independent experiments. E-G, FRAP measurements to test for liquid-like behavior of FIT NBs, using the standardized FIT NB analysis procedure in transiently transformed *N. benthamiana* leaf epidermis cells. E, Representative images of the fluorescent signal during a FRAP experiment, taken before bleaching (0 s) and recovery of fluorescence at three time points after bleaching from 3 s to 45 s within the circled region of a NB. F, Line diagram representing the relative fluorescence during a FRAP measurement for 10 NBs, showing a high fluorescence recovery rate of FIT-GFP within NBs. Dark green line, mean value; light green filled area, variation. G, Box plot diagram representing the mobile fraction of FIT-GFP calculated based on the relative fluorescence recovery in F. The diagram indicates high mobility of FIT. The mean was calculated from 10 NBs from 10 nuclei from a transformed plant. Three independent experiments were conducted, one representative result is shown. H, Box plot diagram representing quantification of the FIT NB shape with the software ImageJ (National Institutes of Health), indicating that FIT NBs have circular shape. Mobility and circularity characteristics indicate that FIT NBs are most likely liquid condensates that are the result of LLPS. The mean was calculated from all NBs visible in 15 nuclei from a transformed plant. Two independent experiments were conducted, one representative result is shown.

Box plots show 25-75 percentile with min-max whiskers, mean as small square and median as line. Scale bars of nuclei images, 2 µm; scale bar full seedling, 1 mm. Arrowheads indicate NBs. G = GFP; C = mCherry.

Figure 2

Figure 2: The FIT C-terminal Ser271/272 site was important for the capacity of FIT to localize to NBs.

A, Confocal images of nuclear localization of FITmSS271AA-GFP at t=0 and t=15 min. FITmSS271AA-GFP accumulated in NBs, but NB formation required a longer time compared to FIT-GFP. See also **Supplemental Movie S1, A and C**. Two independent experiments. Representative images from one nucleus. B, Bar diagrams showing in B, number of NBs, and in C, the sizes of NBs with FIT-GFP and FITmSS271AA-GFP at t=5/15 min. NB number and size were determined with the software ImageJ (National Institute of Health). FIT-GFP accumulated in more and larger NBs than FITmSS271AA-GFP. See **Supplemental Movie S1, A and C**. FITmSS271AA-GFP lacks IDR[Ser271/272]. This IDR may be relevant for FIT NB formation (**Supplemental Figure S2**). In B and C, bar diagrams represent the mean and standard deviation for a quantification of 15 nuclei from a transformed plant (n = 15). Two experiments were conducted, one representative result is shown.

Statistical analysis was performed with the Mann-Whitney test. Different letters indicate statistically significant differences ($P < 0.05$). Scale bar: 2 µm. Arrowheads indicate NBs. G = GFP. Analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure.

Figure 3

Figure 3: FIT was present in homodimeric protein complexes in NBs, dependent on Ser271/272 site.

Anisotropy (or homo-FRET) measurements of FIT-GFP and FITmSS271AA-GFP to determine homodimerization strength. A, Schematic illustration of the anisotropy principle. Energy transfer between the same kind of fluorescently tagged proteins leads to depolarization of the emitted light. Extent of the depolarization gives a hint on dimerization and oligomerization of a protein as the fluorescence anisotropy (FA) value decreases. B, Representative images showing color-coded FA values of FIT-GFP and FITmSS271AA at t=0 and t=5/15 min. C-D, Box plots representing quantification of FA values. FA was measured at t=0 within the whole nucleus and at t =5/15 min within the whole nucleus, in NBs and in residual NP. Free GFP and GFP-GFP served as references for mono- and dimerization. FA values for C, FIT-GFP, and D, FITmSS271AA-GFP. In C and D, FA values were calculated from 10-15 nuclei from a transformed plant (n = 10-15). Two experiments were conducted, one representative result is shown. C and D show the same free GFP and GFP-GFP references because both measurements were performed on the same day. FA values decreased for FIT-GFP, but not FITmSS271AA-GFP, in the whole nucleus (compare t=0 with t=5/15 min). FA values were also lowered in NBs versus NP in the case of FIT-GFP but not FITmSS271AA-GFP (compare t=5/15 min of NBs and NP). This indicates stronger homodimerization of FIT than FITmSS271AA-GFP in the whole nucleus and in NBs. IDR$^{Ser271/272}$ may therefore be relevant for FIT NB formation and FIT homodimerization (**Supplemental Figure S2**).

Box plots show 25-75 percentile with min-max whiskers, mean as small square and median as line. Statistical analysis was performed with one-way ANOVA and Tukey post-hoc test. Different letters indicate statistically significant differences ($P < 0.05$). Scale bar: 2 μm. Arrowheads indicate NBs. G = GFP. Fluorescence protein analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure.

Figure 4

Figure 4: FIT was present in heterodimeric protein complexes with bHLH039 in NBs, dependent on Ser271/272 site.

A, Confocal images with colocalization of FIT-GFP and bHLH039-mCherry in the nucleus. Both proteins were evenly distributed within the nucleus at t=0 and colocalized fully in FIT NBs at t=5 min. Two independent experiments with two plants each. Representative images from one nucleus. B-E, FRET-FLIM measurements to determine heterodimerization strength of FIT and FITmSS271AA with bHLH039, respectively. FIT-GFP and FITmSS271AA-GFP (donor only) served as negative controls. B, Schematic illustration of the FRET-FLIM principle. Energy transfer occurs between two different fluorophores. One fluorophore acts as the donor and the other as the acceptor of the energy. In case of interaction (close proximity, ≤10 nm) the fluorescence lifetime of the donor decreases. C, Representative images showing color-coded fluorescence lifetime values of FIT-GFP and FITmSS271AA-GFP co-expressed with bHLH039-mCherry at t=0 and t=5/15 min. D-E, Box plots diagrams representing FRET-FLIM measurements at t=0 within the whole nucleus and at t=5/15 min within the whole nucleus, inside NBs and in residual NP. Lifetime values represent measurements of 10 nuclei from a transformed plant (n = 10). Two experiments were conducted, one representative result is shown. Fluorescence lifetime was reduced for the pair of FIT-GFP and bHLH039-mCherry in NBs versus NP at t=5 min, indicating protein interaction preferentially inside NBs. Fluorescence lifetime values were not significantly different for the pair FITmSS271AA-GFP and bHLH039-mCherry in this same comparison at t=15 min, indicating that this pair did not preferentially interact in NBs. IDR[Ser271/272] may therefore be relevant for FIT NB formation, and FIT homo- and heterodimerization (**Supplemental Figure S2**).

Box plots show 25-75 percentile with min-max whiskers, mean as small square and median as line. Statistical analysis was performed with one-way ANOVA and Tukey post-hoc test. Different letters indicate statistically significant differences ($P < 0.05$). Scale bar: 2 µm. Arrowheads indicate NBs. G = GFP; C = mCherry. Fluorescence protein analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure.

Figure 5

Figure 5: Two NB markers and splicing components were present in NBs (designated type II), revealing the dynamics of FIT to accumulate in NBs.

Confocal images showing localization of FIT-GFP and NB markers (type II) upon co-expression in the nucleus at t=0 and t=5 min. Co-expression of FIT-GFP with A, SR45-mRFP, and B, SRm102-mRFP. Type II NB markers localized inside NBs at t=0 and t=5 min. Similar localization patterns were observed upon single expression, showing that SR45 and SRm102 are present in distinct NB types (compare with **Supplemental Figure S4, A and B**). FIT-GFP colocalized with type II markers in their distinct NBs at t=5 min, but not t=0. FIT-GFP additionally localized in FIT NBs at t=5 min. Type II markers were not present in FIT NBs, while FIT-GFP became recruited into the distinct type II NBs upon the light trigger. Hence, FIT NBs could be associated with speckle components.

Scale bar: 2 μm. Filled arrowheads indicate colocalization in NBs, empty arrowheads indicate no colocalization in NBs. G = GFP; R = mRFP. Fluorescence protein analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure. Representative images from two to five independent experiments are shown. For data with type I markers (no colocalization) and type III markers (full colocalization) see **Supplemental Figure S3** and **Figure 6**.

Figure 6

Figure 6: Three NB markers and speckle components became localized in FIT NBs and colocalized fully with FIT (designated type III), suggesting that FIT NBs have speckle function.

Confocal images showing localization of FIT-GFP and NB markers (type III) upon co-expression in the nucleus at t=0 and t=5 min. Co-expression of FIT-GFP with A, UAP56H2-mRFP, B, P15H1-mRFP, and C, PININ-mRFP. All three type III NB markers were homogeneously distributed and colocalized with FIT-GFP in the nucleus at t=0, while they colocalized with FIT-GFP in FIT NBs at t=5 min. UAP56H2-mRFP and P15H1-mRFP showed homogeneous localization in the single expression at both t=0 and t=5 min (compare with **Supplemental Figure S4, C and D**), while PININ-mRFP localized mainly in one large and several small NBs upon single expression at t=0 and t=5 min (compare with **Supplemental Figure S4E**). Hence, these three markers adopted the localization of FIT-GFP upon co-expression and suggest that FIT NBs have a speckle function.

Scale bar: 2 μm. Arrowheads indicate colocalization within NBs. G = GFP; R = mRFP. Fluorescence protein analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure. Representative images from four to seven independent experiments are shown. For data with type I markers (no colocalization) and type II markers (partial colocalization) see **Supplemental Figure S** and **Figure 5**.

Figure 7

Figure 7: FIT colocalized with photobody (PB) markers in distinct PBs.

Confocal images showing localization of FIT-GFP and PB markers upon co-expression in the nucleus at t=0 and t=15 min. Co-expression of FIT-GFP with A, PIF3-mCherry, and B and C, PIF4-mCherry, in B, showing a typical pattern with absence of NBs (ca. 50% of nuclei), in C, showing a typical pattern with presence of NBs (ca. 50% of cells). When FIT-GFP was co-expressed with PB markers, FIT NBs did not appear at t=5 min, but instead, FIT-GFP colocalized with PB markers in PBs at t=15 min. A, PIF3-mCherry localized predominantly to a single large PB at t=0 and t=15 min. FIT-GFP colocalized with PIF3-mCherry in this single large PB at t=15 min. B, PIF4-mCherry and FIT-GFP were both homogeneously distributed in the nucleoplasm at t=0 and t=15 min. In C, FIT-GFP colocalized with PIF4-mCherry in PBs at t=0 and t=15 min. The same localization patterns were found for PIF3-mCherry and PIF4-mCherry upon single expression (compare with **Supplemental Figure S5**). Hence, FIT-GFP was recruited to the two distinct types of PIF3 and PIF4 PBs, whereas PIF3 and PIF4 were not recruited to FIT NBs. This suggests that FIT NBs are affected by the presence of PIF3- and PIF4-containing PBs and a connection to light signaling exists.

Scale bar: 2 μm. Arrowheads indicate colocalization in PBs. G = GFP; C = mCherry. Fluorescence protein analysis was conducted in transiently transformed *N. benthamiana* leaf epidermis cells, following the standardized FIT NB analysis procedure. Representative images from four to six independent experiments are shown.

Figure 8

Figure 8. Root iron reductase activity is promoted under blue light.

Root iron reductase activity assay on 6-d-old Arabidopsis seedlings grown under white light for 5 d under iron deficient and iron sufficient conditions and then exposed for 1 d to blue, red, far-red light or darkness, or in parallel as control to white light. A, Induction of iron reductase activity upon iron deficiency versus sufficiency was higher under blue light than compared to the white light control. B, Exposure to red light did not change the iron reductase activity compared to the white light control. C-D, Plants exposed to far-red light and darkness did not show induction of iron reductase activity under iron deficient conditions compared to the iron deficient conditions.

Two experiments were conducted, one representative result is shown. Bar diagrams represent the mean and standard deviation of four replicates with four seedlings each (n=4). Statistical analysis was performed with one-way ANOVA and Tukey post-hoc test. Different letters indicate statistically significant differences ($P < 0.05$).

Figure 9

Figure 9. Induction of FIT target gene expression is enhanced under blue light.

Gene expression analysis of iron deficiency genes *FIT*, *BHLH039*, *FRO2*, and *IRT1* on 6-d-old Arabidopsis seedlings grown under white light for 5 d under iron deficient and iron sufficient conditions and then exposed for 1 d to blue, red, far-red light or darkness, or in parallel as control to white light. A-B, *FIT* and *bHLH039* gene induction in response to low iron supply did not change after blue light exposure compared to the white light control. C-D, *FRO2* and *IRT1* gene expression increased after blue light exposure in iron deficient conditions compared to respective iron sufficient condition and white light control. E, *FIT* gene expression did not change after red light exposure compared to the white light control. F-G, *BHLH039* and *FRO2* gene expression decreased after red light exposure in iron deficient conditions compared to respective iron sufficient condition and white light control. H, *IRT1* gene expression did not change after red light exposure compared to the white light control. I, *FIT* gene expression did not change after far-red light exposure compared to the white light control. J-L, *BHLH039*, *FRO2*, and *IRT1* gene expression decreased after far-red light exposure in iron deficient conditions compared to white light deficient condition, comparable to the respective iron sufficient condition and iron sufficient white light control. M, *FIT* gene expression was higher in darkness under iron deficient conditions compared to respective iron sufficient condition and white light control. N-P, *BHLH039*, *FRO2*, and *IRT1* gene expression decreased after being exposed to darkness in iron deficient conditions compared to white light deficient condition, comparable to the respective iron sufficient condition and iron sufficient white light control.

Two experiments were conducted, one representative result is shown. Bar diagrams represent the mean and standard deviation of three replicates with twelve seedlings and two technical replicates each (n=4). Statistical analysis was performed with one-way ANOVA and Tukey post-hoc test. Different letters indicate statistically significant differences ($P < 0.05$).

Figure 10

-Fe

FRO2

IRT1

Fe^{3+}

Fe^{2+}

FIT-bHLH039
heterodimer

FIT-FIT
homodimer

Speckle
component

DNA

RNA

Figure 10: Summary model illustrating the effect of blue light on FIT NB formation and iron uptake, suggesting that FIT NBs are related to transcriptional and posttranscriptional regulation in speckles and that blue light has a promoting effect on iron uptake.

FIT accumulates in FIT NBs upon induction with blue light. FIT NBs are reversible, dynamic and of circular shape and may undergo LLPS. FIT homodimers and FIT-bHLH039 heterodimers are present in FIT NBs. FIT-bHLH039 is an active TF complex for upregulating the expression of iron acquisition genes in roots (Gratz et al., 2019). Hence, FIT NBs are subnuclear sites related to transcriptional regulation and because of their colocalization with speckle components, also to speckles. Blue light enhances root iron reductase activity and gene expression of *FRO2* and *IRT1*, all of which are downstream of FIT-bHLH039. In summary, FIT NBs are blue light-inducible subnuclear sites where active TF complexes of FIT and bHLH039 accumulate, linking transcriptional and posttranscriptional regulation in speckles with a promoting blue light effect on iron uptake.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

FRO2

IRT1

Fe^{3+}

Fe^{2+}

-Fe

FIT-bHLH039
heterodimer

FIT-FIT
homodimer

Speckle
component

DNA

RNA