pGG-PIP: A GreenGate (GG) entry vector collection with Plant Immune system Promoters

(PIP)

Jacob Calabria[1,2,a], Madlen I. Rast-Somssich[2,5,b,#], Liu Wang[1,3,6,c,#], Hsiang-Wen Chen[1,3,d],

Michelle Watt[2,e], Alexander Idnurm[4,f], Staffan Persson[3,6,g], and Marc Somssich[1,7,h,*]

[1] Plant-Fusarium Interactions Research Team, School of BioSciences, University of Melbourne, Parkville, Australia

[2] Crop Root Physiology Lab, School of BioSciences, University of Melbourne, Parkville, Australia

[3] Plant Cell Biology Lab, School of BioSciences, University of Melbourne, Parkville, Australia

[4] Mycology Laboratory, School of BioSciences, University of Melbourne, Parkville, Australia

[5] Current address: Institute for Molecular Physiology, Heinrich Heine University, Düsseldorf, Germany

[6] Current address: Copenhagen Plant Science Center, University of Copenhagen, Frederiksberg C, Denmark

[7] Current address: Max Planck Institute for Plant Breeding Research, Cologne, Germany

Twitter: [a] @jacobcalabria, [b] @MSomssich, [c] @Liu_Wang1, [d] @HsiangWen_Chen, [e] @MelbourneRoots, [f]

@AlexanderIdnurm, [g] @perssonlab1, [h] @somssichm

[#] These two authors contributed equally

[*] For correspondence: marc.somssich@unimelb.edu.au

Abstract

The regulatory sequences controlling the expression of a gene (i.e., the promoter) are essential to

properly understand a gene's function. From their use in mutant complementation assays, to

studying their responsiveness to different stimuli via transcriptional reporter lines or using them

as proxy for the activation of certain pathways, assays using promoter sequences are valuable tools

for insight into the genetic architecture underlying plant life. The GreenGate (GG) system is a

plant-specific variant of the Golden Gate assembly method, a modular cloning system that allows

the hierarchical assembly of individual donor DNA fragments into one expression clone via a

single reaction step. Here, we present a collection of 75 GG entry vectors carrying putative

regulatory sequences for *Arabidopsis thaliana* genes involved in many different pathways of the

plant immune system, designated Plant Immune system Promoters (PIP). This pGG-PIP entry

vector set enables the rapid assembly of expression vectors to be used for transcriptional reporters

of plant immune system components, mutant complementation assays when coupled with coding

sequences, mis-expression experiments for genes of interest, or the targeted use of CRISPR/Cas9

genome editing. We used pGG-PIP vectors to create fluorescent transcriptional reporters in *A.*

thaliana and demonstrated the potential of these reporters to image the responsiveness of specific

plant immunity genes to infection and colonization by the fungal pathogen *Fusarium oxysporum*.

Using the PLANT ELICITOR PEPTIDE (PEP) pathway as an example, we show that several

components of this pathway are locally activated in response to colonization by the fungus.

Keywords

36 Golden Gate; GreenGate; plant immunity; *Arabidopsis thaliana*; plant-microbe interactions;

37 promoters; transcriptional reporters; PLANT ELICITOR PEPTIDE; synthetic biology; *Fusarium*

38 *oxysporum*

Introduction

Since the development of molecular cloning in the early 1970s, the isolation of genes and

promoters, and subsequent transgenesis of model organisms, has become standard practice in the

life sciences (Somssich, 2022). The development of more advanced cloning methods, particularly

the recombination-based Gateway technology in the year 2000, made the creation of expression

clones ready for transformation ever easier (Hartley *et al.*, 2000). However, with the rise of the

synthetic biology field, it is now no longer sufficient for these methods to facilitate the cloning of

individual DNA fragments. To recreate entire pathways and gene circuits in plants and other model

organisms, larger DNA constructs need to be readily assembled from individual components, and

these distinct building blocks need to be compatible to allow the flexibility to mix and match

different promoters, coding sequences, protein tags, terminators, and resistance genes for selection

of transgenic lines (Meng and Ellis, 2020). This requirement was met with the new modular

cloning techniques which use recombination-based hierarchical assembly of multiple donor-

modules (each containing, for example, a promoter of choice, tag of choice, gene of interest, etc.)

into one ordered expression clone to be used for transgenesis (Fig. 1) (Bird *et al.*, 2022). Among

the developed modular cloning methods, the Golden Gate system has emerged as the most widely

utilized version, and in 2013 Lampropoulos *et al.* developed the plant-specific GreenGate variant

of the Golden Gate technique (Engler *et al.*, 2008, 2009; Weber *et al.*, 2011; Lampropoulos *et al.*,

2013). The GreenGate toolkit provides users with a wide range of entry vectors that serve as donors

for standard promoters (e.g. CaMV35S, UBQ10), protein tags (e.g. GFP, NLS, HDEL),

terminators (e.g. CaMV35S, UBQ10) and plant resistance cassettes (e.g. BastaR, HygR, KanR),

to build basic gene expression modules for plants.

Figure 1: Overview of the modular Golden/GreenGate cloning principle
The promoter, N- and C-terminal tags, the gene-of-interest coding sequence (CDS), a terminator, and a plant resistance cassette are all cloned into individual entry clones (A-F), which are then combined with an 'empty' destination clone (Z) in the Golden Gate assembly mix. During the assembly reaction, the *Bsa*I restriction enzyme cuts the different fragments from their respective entry clones. These fragments subsequently self-align via complementary four base pair overhangs (indicated by the thin colored borders on each entry vector; complementary overhangs that will align have the same color), and the T4 DNA-ligase fuses the individual fragments in the final expression clone (Z) that can be used to transform plants.

Additional kits have been developed and added to the GreenGate toolbox since 2013, providing

entry vector sets to build plasmids for CRISPR/Cas9-guided genome editing (Wu *et al.*, 2018),

CRISPR/Cas9-guided tissue-specific gene knockout (Decaestecker *et al.*, 2019) and inducible and

cell-type specific gene expression (Schürholz *et al.*, 2018). In addition, a toolbox of fluorescent

proteins suitable for work in plants (Denay *et al.*, 2019), along with the necessary clones for *in*

planta proximity ligation assays using the biotin ligase (Goslin *et al.*, 2021), and destination

vectors with an already integrated plasma membrane-marker (Kümpers *et al.*, 2022) have been

designed. The development and availability of these various toolboxes, all compatible with each

other, demonstrate the usefulness of such modular systems to enable researchers to quickly adopt

new technologies, providing the flexibility and versatility to combine and recombine their existing

vectors (i.e., modules) with new vectors from all other kits.

In our work we use a microscopy-based live-imaging approach to monitor the *Arabidopsis*

thaliana's defense responses to infection and colonization by the pathogenic fungus *Fusarium*

oxysporum strain *Fo*5176 (*Fo*5176) on an individual cell level (Calabria *et al.*, 2022; Wang *et al.*,

2022*a*). Fluorescent transcriptional reporters are a good tool for such studies, as activation of

certain pathways is typically associated with transcriptional upregulation (Ngou *et al.*, 2021).

However, most studies investigating the transcriptional responsiveness of certain pathways to

pathogens tend to use transcriptomic analyses of whole tissues, organs, or seedlings, therefore

losing all spatial resolution and differences between cell groups and even individual cells. The

responses of only the few cells that could be the major contributors to a host-microbe outcome are

also typically lost in such experiments, as they are below the threshold of background noise. A

microscopy-based approach that allows for the spatial resolution and observation of responses in

individual cells therefore closes this knowledge gap. One reason why such an approach is not more

common is that a prerequisite for this work is the availability of fluorescent reporter lines for all

the different plant immune pathways to be investigated. To this end, we have cloned the putative

regulatory sequences (i.e., promoters) of 75 *A. thaliana* genes, representing many of the major

branches of the plant immune system (Fig. 2 and Table 1). We have done so using the GreenGate

cloning system, therefore creating a GreenGate plant promoter entry vector set that is compatible

with all other GreenGate-based toolkits, and that we believe will be a valuable tool for the scientific

community.

As a proof-of-principle, we selected a number of these promoters to create fluorescent

transcriptional reporter lines for genes involved in the PLANT ELICITOR PEPTIDE (PEP)-

pathway. The PEP pathway is involved in the immune response to several plant pathogens,

including bacteria, fungi and oomycetes (Saijo *et al.*, 2018). The PEP1-6 peptides function as

phytosulfokines. They are expressed as part of the plant's pattern-triggered immunity (PTI)

response and trigger cell autonomous and non-autonomous defense responses when they are

perceived by the plasma membrane localized receptors PEPR1 and PEPR2 (Yamaguchi *et al.*,

2006, 2010; Saijo *et al.*, 2018). For their function, they require the intracellular kinase BIK1, which

activates the NADPH oxidase RBOHD to trigger a ROS-burst, and also induces defense gene

expression (Liu *et al.*, 2013; Kadota *et al.*, 2014; Saijo *et al.*, 2018; Jing *et al.*, 2020). A role for

the PEP pathway in the defense against fungal disease has been hypothesized, based on large-scale

transcriptomic data. Plants carrying mutations in *MEDIATOR18* and *20* show increased resistance

to infection by *F. oxysporum* strain *Fo*5176, and mRNA sequencing has demonstrated that *PEP1*

is among the genes no longer induced after infection of these mutant plants (Fallath *et al.*, 2017;

Wang *et al.*, 2022*a*). Similarly, *PEPR1* and *2* were among the genes suppressed by the endophytic

F. oxysporum strain *Fo*47, indicating that the PEP pathway normally acts to restrict fungal

colonization (Guo *et al.*, 2021; Wang *et al.*, 2022*a*). However, no detailed data is available. Here,

we show that colonization of the *A. thaliana* vasculature is accompanied by a highly localized

Figure 2: Overview of the different immune pathways represented in the pGG-PIP collection
The red numbers correspond to the pGG-PIP plasmid numbers in column one of Table 1.

activation of this pathway via upregulation of genes encoding the peptide ligand PEP1, as well as the cognate PEP RECEPTORs 1 and 2, the downstream signaling intermediate BIK1 and the NADPH oxidase RBOHD.

Results

Selection and cloning of the promoters

By using these putative promoter sequences for the construction of transcriptional reporter lines, our main aim is to obtain information about the specific localized activation of different immune pathways in response to infection by a pathogen, in our case *Fo*5176. Once we have identified pathways involved in the plant's defense against this strain, we plan to further extend these experiments based on what is known about the function of these pathways. Thus, it is important to select promoters from many different pathways of the plant immune system, but also from pathways that are particularly well understood, since this existing knowledge will guide our future work. Some of the best studied immune pathways in the plant are the pattern-recognition flagellin (flg) and elongation factor thermos unstable (EF-Tu) pathways, the phytohormones jasmonic acid (JA), salicylic acid (SA) and ethylene (ET), and the transcriptional activity of several WRKY transcription factors. We therefore selected representative gene promoters from such well understood pathways, but also additional candidates such as MAP and calcium-dependent protein kinases (MPKs, CPKs), genes involved in production and signaling via reactive oxygen species (ROS), and genes involved in effector-triggered immunity (ETI) responses via the EDS1-PAD4 module. Figure 2 and Table 1 provide a complete overview of the 75 promoters included in the pGG-PIP collection. The numbers in Fig. 2 correspond to the pGG-PIP plasmid number in the first column of Table 1.

For the construction of the pGG-PIP plasmids, we considered the entire stretch of DNA from the 3´ upstream neighboring gene's coding sequence to the ATG of the gene of interest as the gene's putative regulatory sequences, and thus its 'promoter'. There are a few exceptions to this rule, where we used a defined stretch of DNA that has previously been described to complement a mutant (e.g., *FMO1*, *RPS4* (Wirthmueller *et al.*, 2007; Joglekar *et al.*, 2018)). Where present, this may include untranslated regions (UTRs), pseudogenes, or transposons, since these elements could indeed affect expression of a gene *in planta*. For the recombination-based cloning, the *Bsa*I recognition site and standard four base pair GreenGate A overhangs (5´-ACCT and TTGT-3´) were added via primers during amplification of the promoter sequences (see Supplementary Table 1 for a list of primer sequences). If the promoter sequence contained an internal *Bsa*I restriction site (GGTCTC), we mutated a single base, usually a G to a C via scar-free *Bsa*I cloning, i.e., we

inserted the base mutation by making it part of the four base pair overhang used for the scar-free

cloning (see Table 1 for the exact sequence edits made, and Supplementary Table 1 for primer

sequences). The promoters were then transferred into the GreenGate promoter entry vector

pGGA000 via the four base pair A overhangs in a Golden Gate assembly reaction.

GG-PIP plasmid	Gene	Bases	*Bsa*I site edits	Protein function	Full gene name	Gene code
pGG-PIP01	*PEP1*	1907		Activate defense genes (Yamada *et al.*, 2016)	*PLANT ELICITOR PEPTIDE 1*	*AT5G64900*
pGG-PIP02	*PEP2*	769		Activate defense genes (Yamada *et al.*, 2016)	*PLANT ELICITOR PEPTIDE 2*	*AT5G64890*
pGG-PIP03	*PEP3*	1694		Activate defense genes (Yamada *et al.*, 2016)	*PLANT ELICITOR PEPTIDE 3*	*AT5G64905*
pGG-PIP04	*RALF23*	2027	1458 C -> G	Phytosulfokine sensed by FER (Stegmann *et al.*, 2017)	*ARABIDOPSIS RAPID ALKALINIZATION FACTOR 23*	*AT3G16570*
pGG-PIP05	*SCOOP12*	2900		Phytosulfokine sensed by MIK2/BAK1 (Hou *et al.*, 2021; Rhodes *et al.*, 2021)	*PRECURSOR OF SERINE-RICH ENDOGENOUS PEPTIDE 12*	*AT5G44585*
pGG-PIP06	*BAK1*	1731		Co-receptor for several defense & development pathways (Greenwood and Williams, 2022)	*BRI1-ASSOCIATED RECEPTOR KINASE*	*AT4G33430*
pGG-PIP07	*CERK1*	493		Chitin-receptor (Cao *et al.*, 2014)	*CHITIN ELICITOR RECEPTOR KINASE 1*	*AT3G21630*
pGG-PIP08	*CIPP1*	3003		CERK1 co-receptor (Liu *et al.*, 2018)	*CERK-1 INTERACTING PROTEIN PHOSPHATASE 1*	*AT1G34750*
pGG-PIP09	*CORK1*	3376		DAMP-receptor (Tseng *et al.*, 2022)	*CELLOOLIGOMER RECEPTOR KINASE 1*	*AT1G56145*
pGG-PIP10	*EFR*	2376		Receptor for EF-Tu (Couto and Zipfel, 2016)	*EF-TU RECEPTOR*	*AT5G20480*
pGG-PIP11	*FER*	1243		Co-receptor for several defense & development pathways (Duan *et al.*, 2022)	*FERONIA*	*AT3G51550*
pGG-PIP12	*FLS2*	2913		Receptor for bacterial flagellin (Couto and Zipfel, 2016)	*FLAGELLIN-SENSITIVE 2*	*AT5G46330*
pGG-PIP13	*LYK5*	1560	1215 G->C	CERK1 co-receptor (Cao *et al.*, 2014)	*LYSM-CONTAINING RECEPTOR-LIKE KINASE 5*	*AT2G33580*
pGG-PIP14	*LYM1*	991		Fungal MAMP receptor (Zipfel and Oldroyd, 2017)	*LYSM DOMAIN GPI-ANCHORED PROTEIN 1 PRECURSOR*	*AT1G21880*
pGG-PIP15	*MIK2*	2513		SCOOP peptide receptor (Hou *et al.*, 2021; Rhodes *et al.*, 2021)	*MALE DISCOVERER 1-INTERACTING RECEPTOR-LIKE KINASE 2*	*AT4G08850*
pGG-PIP16	*PEPR1*	931	180 G->A	Receptor for PEP1-6 (Yamada *et al.*, 2016)	*PEP1 RECEPTOR 1*	*AT1G73080*
pGG-PIP17	*PEPR2*	1676		Receptor for PEP1 & 2 (Yamada *et al.*, 2016)	*PEP1 RECEPTOR 2*	*AT1G17750*
pGG-PIP18	*RFO1*	850		Cell wall-associated kinase (Huerta *et al.*, 2023)	*RESISTANCE TO FUSARIUM OXYSPORUM 1*	*AT1G79670*
pGG-PIP19	*RLP26*	822		Co-receptor for PRR receptors (Wu *et al.*, 2016)	*RECEPTOR LIKE PROTEIN 26*	*AT2G33050*
pGG-PIP20	*RLP29*	2790		Co-receptor for PRR receptors (Wu *et al.*, 2016)	*RECEPTOR LIKE PROTEIN 29*	*AT2G42800*
pGG-PIP21	*SOBIR1*	1163		Co-receptor for several defense pathways (Cho *et al.*, 2022)	*SUPPRESSOR OF BIR 1 / EVERSHED*	*AT2G31880*
pGG-PIP22	*BIK1*	2668		Defense signaling (Gonçalves Dias *et al.*, 2022)	*BOTRYTIS-INDUCED KINASE1*	*AT2G39660*
pGG-PIP23	*XLG2*	1256		G protein involved in immunity (Petutschnig *et al.*, 2022)	*EXTRA-LARGE GTP-BINDING PROTEIN 2*	*AT4G34390*
pGG-PIP24	*MPK3*	654		Activates immune response (Tsuda and Somssich, 2015)	*MITOGEN-ACTIVATED PROTEIN KINASE 3*	*AT3G45640*
pGG-PIP25	*MPK4*	1025		Activates immune response (Tsuda and Somssich, 2015)	*MITOGEN-ACTIVATED PROTEIN KINASE 4*	*AT4G01370*

pGG-PIP26	*MPK6*	811		Activates immune response (Tsuda and Somssich, 2015)	*MITOGEN-ACTIVATED PROTEIN KINASE 6*	*AT2G43790*
pGG-PIP27	*CPK5*	1983		Involved in calcium-dependent stress responses (Gao and He, 2013)	*CALMODULIN-DOMAIN PROTEIN KINASE 5*	*AT4G35310*
pGG-PIP28	*CPK29*	826	325 G->C	Involved in calcium-dependent stress responses (Patil and Senthil-Kumar, 2020)	*CALCIUM-DEPENDENT PROTEIN KINASE 29*	*AT1G76040*
pGG-PIP29	*GLR2.5*	2488	1488, 1489 G->C	Involved in defense-development balancing (Birkenbihl *et al.*, 2017*a*)	*GLUTAMATE RECEPTOR 2.5*	*AT5G11210*
pGG-PIP30	*GLR2.7*	4000		Involved in defense-development balancing (Birkenbihl *et al.*, 2017*a*)	*GLUTAMATE RECEPTOR 2.7*	*AT2G29120*
pGG-PIP31	*RBOHD*	2309		Produces ROS-burst (Kadota *et al.*, 2014)	*RESPIRATORY BURST OXIDASE HOMOLOGUE D*	*AT5G47910*
pGG-PIP32	*RBOHF*	4119		Produces ROS-burst (Morales *et al.*, 2016)	*RESPIRATORY BURST OXIDASE HOMOLOG F*	*AT1G64060*
pGG-PIP33	*EX1*	989		Produces ROS from chloroplasts (Dogra *et al.*, 2022)	*EXECUTER 1*	*AT4G33630*
pGG-PIP34	*HPCA1*	2944		ROS-sensor (Sun and Zhang, 2021)	*HP-INDUCED Ca2+ INCREASES 1*	*AT5G49760*
pGG-PIP35	*RCD1*	3137	2888 G -> C	Modulator of root-to-shoot ROS-signaling (Jin *et al.*, 2022)	*RADICAL-INDUCED CELL DEATH1*	*AT1G32230*
pGG-PIP36	*DORN1*	2325		eATP DAMP-receptor (Sun and Zhang, 2021)	*DOES NOT RESPOND TO NUCLEOTIDES 1*	*AT5G60300*
pGG-PIP37	*LECRK-VI.2*	1183		NAD(P)-receptor (Sun and Zhang, 2021)	*L-TYPE LECTIN RECEPTOR KINASE-VI.2*	*AT5G01540*
pGG-PIP38	*BPS1*	2142		Regulator of root-to-shoot communication (Lee *et al.*, 2016)	*BYPASS 1*	*AT1G01550*
pGG-PIP39	*WRKY11*	2026	240 C->G	Regulates defense gene expression (Journot-Catalino *et al.*, 2006)	*WRKY DNA-BINDING PROTEIN 11*	*AT4G31550*
pGG-PIP40	*WRKY17*	4565	4325 G->C	Regulates defense gene expression (Birkenbihl *et al.*, 2018)	*WRKY DNA-BINDING PROTEIN 17*	*AT2G24570*
pGG-PIP41	*WRKY33*	1665		Regulates defense gene expression (Birkenbihl *et al.*, 2018)	*WRKY DNA-BINDING PROTEIN 33*	*AT4G23810*
pGG-PIP42	*WRKY40*	4158		Regulates defense gene expression (Birkenbihl *et al.*, 2018)	*WRKY DNA-BINDING PROTEIN 40*	*AT1G80840*
pGG-PIP43	*WRKY53*	2515		Regulates defense gene expression (Birkenbihl *et al.*, 2017*b*)	*WRKY DNA-BINDING PROTEIN 53*	*AT2G38470*
pGG-PIP44	*WRKY70*	4099		Regulates defense gene expression (Journot-Catalino *et al.*, 2006)	*WRKY DNA-BINDING PROTEIN 70*	*AT3G56400*
pGG-PIP45	*ELI-3*	524		Elicitor-response gene (Tanaka *et al.*, 2018)	*CINNAMYL-ALCOHOL DEHYDROGENASE 7 / ELICITOR-ACTIVATED GENE3*	*AT4G37980*
pGG-PIP46	*FRK1*	1271		Defense gene (Birkenbihl *et al.*, 2017*b*)	*FLG22-INDUCED RECEPTOR-LIKE KINASE 1*	*AT2G19190*
pGG-PIP47	*MLO6*	3042		Defense gene (Acevedo-Garcia *et al.*, 2017)	*MILDEW RESISTANCE LOCUS O 6*	*AT1G61560*
pGG-PIP48	*PER5*	2007	558 C->G	Marker for activated immune signalling (Chuberre *et al.*, 2018)	*PEROXIDASE 5*	*AT1G14550*
pGG-PIP49	*PLP1*	2490		Pathogen-induced (Yang *et al.*, 2007)	*PATATIN-LIKE PROTEIN 1*	*AT4G37070*
pGG-PIP50	*ACS2*	2914	150 C->G	ET biosynthesis (Wang *et al.*, 2022*c*)	*1-AMINO-CYCLOPROPANE-1-CARBOXYLATE SYNTHASE 2*	*AT1G01480*
pGG-PIP51	*AOS*	2093	2088 G->C ;	JA biosynthesis (Yang *et al.*, 2019)	*ALLENE OXIDE SYNTHASE*	*AT5G42650*

			1434 G->C			
pGG-PIP52	*ARR5*	2229	639 C -> G	Cytokinin signaling (Lee *et al.*, 2016)	*ARABIDOPSIS RESPONSE REGULATOR 5*	*AT3G48100*
pGG-PIP53	*EDS16*	2975	1579 G->C ; 1049 C->G	SA biosynthesis (Calabria *et al.*, 2022)	*ENHANCED DISEASE SUSCEPTIBILITY TO ERYSIPHE ORONTII 16*	*AT1G74710*
pGG-PIP54	*ERF1*	2682		ET & JA response regulator (Wang *et al.*, 2022*c*)	*ETHYLENE RESPONSE FACTOR 1*	*AT3G23240*
pGG-PIP55	*PDF1.2*	1540		ET & JA-induced defense gene (Yang *et al.*, 2019)	*PLANT DEFENSIN 1.2*	*AT5G44420*
pGG-PIP56	*PR1*	2343		SA-induced defense gene (Yang *et al.*, 2019)	*PATHOGENESIS-RELATED GENE 1*	*AT2G14610*
pGG-PIP57	*VSP2*	1471		JA-induced defense gene (Yang *et al.*, 2019)	*VEGETATIVE STORAGE PROTEIN 2*	*AT5G24770*
pGG-PIP58	*FMO1*	1726		Systemic acquired resistance marker (Joglekar *et al.*, 2018)	*FLAVIN-DEPENDENT MONOOXYGENASE 1*	*AT1G19250*
pGG-PIP59	*MYB72*	4619		Involved in induced systemic resistance	*ARABIDOPSIS THALIANA MYB DOMAIN PROTEIN 72*	*AT1G56160*
pGG-PIP60	*CSLD2*	1963		Marker for cell death, possibly via ET-signaling (Salguero-Linares *et al.*, 2022)	*CELLULOSE-SYNTHASE LIKE D2*	*AT5G16910*
pGG-PIP61	*HRM1*	2238		Marker for cell death (Salguero-Linares *et al.*, 2022)	*HYPERSENSITIVE RESPONSE MARKER 1*	*AT5G17760*
pGG-PIP62	*EDS1*	1419	848 A->T ; 1269 G->C	Involved in ETI-signaling (Jia *et al.*, 2022)	*ENHANCED DISEASE SUSCEPTIBILITY 1*	*AT3G48090*
pGG-PIP63	*PAD4*	1596		Involved in ETI-signaling (Jia *et al.*, 2022)	*PHYTOALEXIN DEFICIENT 4*	*AT3G52430*
pGG-PIP64	*RPS4*	511		NLR effector receptor (Jia *et al.*, 2022)	*RESISTANT TO P. SYRINGAE 4*	*AT5G45250*
pGG-PIP65	*ATG8A*	1408		Autophagy marker (Yang *et al.*, 2018)	*AUTOPHAGY-RELATED 8A*	*AT4G21980*
pGG-PIP66	*TET8*	2068		Targets vesicular transport to infection sites (Cai *et al.*, 2018)	*TETRASPANIN8*	*AT2G23810*
pGG-PIP67	*SULTR4;1*	2087		Sulfate transporter (Wang *et al.*, 2022*b*)	*SULFATE TRANSPORTER 4.1*	*AT5G13550*
pGG-PIP68	*SULTR4;2*	712		Sulfate transporter (Wang *et al.*, 2022*b*)	*SULFATE TRANSPORTER 4.2*	*AT3G12520*
pGG-PIP69	*CAD5*	1134		Lignin biosynthesis (Kim *et al.*, 2020)	*CINNAMYL ALCOHOL DEHYDROGENASE 5*	*AT4G34230*
pGG-PIP70	*GPAT5*	1587		Suberin biosynthesis (Andersen *et al.*, 2015)	*GLYCEROL-3-PHOSPHATE SN-2-ACYLTRANSFERASE 5*	*AT3G11430*
pGG-PIP71	*MYB15*	2050		Lignin biosynthesis in response to pathogens (Kim *et al.*, 2020)	*ARABIDOPSIS THALIANA MYB DOMAIN PROTEIN 15*	*AT3G23250*
pGG-PIP72	*PMR4*	1954		PAMP-induced defense, callose-deposition (Blümke *et al.*, 2013)	*POWDERY MILDEW RESISTANT 4*	*AT4G03550*
pGG-PIP73	*FH16*	1158		Involved in microtubule/actin-network reorganisation (Wang *et al.*, 2013)	*FORMIN HOMOLOG 16*	*AT5G07770*
pGG-PIP74	*NET4A*	1718		Modulates the vacuole (Kaiser *et al.*, 2019)	*NETWORKED 4A*	*AT5G58320*
pGG-PIP75	*TUB6*	3349	2358 A->G	Component of microtubules (Liu *et al.*, 2019)	*BETA-6 TUBULIN*	*AT5G12250*

Table 1: List of the 75 pGG-PIP promoter entry vectors in the set described in this paper.
The pGG-PIP numbers correspond to the red numbers in Fig. 2.

Using pGG-PIP vectors to create transcriptional reporters for the PEP pathway

To test the usefulness of these promoters to map local individual cell responses upon attack by a

microbe, we created transcriptional reporter lines for genes involved in the PEP pathway. We used

a nuclear localized tandem of mTurqoise2 (mT2) fluorophores as a reporter, since mT2 has been shown to be very bright and photostable in plant cells (Goedhart *et al.*, 2012; Long *et al.*, 2018; Denay *et al.*, 2019). The nuclear localization enhances the brightness even further due to molecular crowding, and furthermore helps to identify individual cells.

We first probed our reporters for the PEP1 and PEP2 peptides (pGG-PIP01 and 02), as well as the PEPR1 and PEPR2 receptors (pGG-PIP16 and 17), assessing their local responsiveness to colonization by *Fo*5176. Under control uninfected conditions, *PEP1* was weakly expressed in inner tissue cells of the root differentiation zone (DZ) (Fig. S1A-D). No expression was detectable in the root tip, the meristematic or elongation zones (MZ and EZ). *PEP2* was robustly expressed in the DZ, but in contrast to *PEP1*, its expression was stronger in the outer tissues (Fig. S1G-H). In addition, it was also expressed in the root tip, around the MZ and early EZ (Fig. S1E-F). *PEPR1* was generally expressed at a very weak level. In the root tip, starting with the EZ, expression was limited to the vasculature, while from the young DZ (or root hair zone) onward, expression was restricted to the outer tissues (Fig. S2A-D). *PEPR2* was expressed only in the DZ, and only in the vasculature, but expression was much stronger compared to *PEPR1* (Fig. S2E-H). Thus, in vascular cells, *PEPR1* and *PEPR2* complemented each other under uninfected conditions. Following colonization of the vasculature by *Fo*5176, *PEP1*, as well as both *PEPR*s, showed a strong upregulation in the cells adjacent to the colonization site (Fig. 3). Interestingly, this

Figure 3: Local transcriptional responses in *PEP1*, *PEP2*, *PEPR1* and *PEPR2* expression to colonization by *Fo*5176
Bright field and fluorescence (**A, C, E, G, I**) and fluorescence only (**B, D, F, H, J**) images of colonized roots expressing the *PEP1* (A, B), *PEP2* (C, D), *PEPR1* (E, F) or *PEPR2* (G-J) reporters. *Fo*5176 is shown in magenta, the reporters in yellow. The purple bars indicate the fungal colonization front. All images were recorded with comparable imaging settings, except for G and H, which are the same as I and J, just with reduced exposure time. Expression of the *PEPR2* reporter was so strong (I, J), that we reduced the exposure time to visualize individual cells in G and H.

upregulation was limited to the cells of the vasculature, the exact tissue that is targeted and
colonized by *Fo*5176, showing how precisely the plant targets its response. Importantly, the
induction of *PEPR2* was much stronger than the induction of *PEP1* and *PEPR1* (Fig. 3G-J).
Interestingly, *PEP2* also showed an induction next to the colonized tissue, but its expression
appeared to be limited to the outer tissues, the epidermis and cortex, rather than the colonized
vasculature (Fig. 3C, D). The strong induction of the *PEPs* and *PEPRs* remained visible for
roughly 10 to 12 cells, and then tapered out, eventually returning to the expression pattern of the
uninfected control.

Since it was shown that the ability of the PEPs and PEPRs to induce any signaling downstream of
the receptors is strictly dependent on the activity of the cytoplasmic kinase BIK1, we also
investigated the responsiveness of a *BIK1* (pGG-PIP22) transcriptional reporter (Liu *et al.*, 2013;
Yamada *et al.*, 2016). Under uninfected control conditions, *BIK1* is expressed in the MZ and DZ,
with stronger expression in the outer tissues, and only weak expression in the vascular tissue of
the MZ, somewhat resembling the expression pattern of *PEP2* (compare Fig. S3A-D and S1E-H).
Following colonization of the root by *Fo*5176, this pattern changed significantly, with *BIK1* now
expressed specifically in the cells immediately bordering the colonized tissue, with a clearly visible
expression maximum in the vasculature (Fig. 4A, B). Hence, the expression maxima of *BIK1*,
PEP1, *PEPR1* and *PEPR2* all overlapped in a small group of vascular cells closest to the colonized
tissue. *BIK1* however, also showed expression in the endodermis and cortex, albeit at a much lower
level than in the vasculature, while *PEP1* and *PEPRs* expression were restricted to the vasculature
(Fig. 3 and Fig. 4A, B).

One of the downstream outputs of PEP-activated BIK1 is the activation of the NADPH oxidase
RBOHD to trigger an apoplastic ROS-burst (Holmes *et al.*, 2018; Jing *et al.*, 2020). We therefore
investigated if *RBOHD* (pGG-PIP31) is also activated in the cells expressing *PEP1*, the *PEPRs*
and *BIK1*. Under uninfected control conditions, *RBOHD* is expressed in all cells and tissues of the
DZ, but not in the root tip (MZ, EZ) (Fig. S3E-H). Upon infection by *Fo*5176, expression is
activated in the cells next to the colonization site, even if they are still part of the EZ, which
normally does not express *RBOHD* (Fig. 4C, D). The pattern we observed very much resembled

Figure 4: Transcriptional responses of *BIK1* and *RBOHD* expression to colonization by *Fo*5176
Bright field and fluorescence (**A, C**), fluorescence only (**B, D**), or bright field only (**E**) images of a colonized root expressing the *BIK1* (A, B), or *RBOHD* (C, D, E) reporters. The *Fo*5176 is shown in magenta, the reporters in yellow. The purple bars indicate the fungal colonization front. The white bracket in E indicates the area with 'root browning'.

the pattern of *BIK1* expression in response to colonization, with a maximum in a small group of vascular cells next to the colonization site (Fig. 4). However, in contrast to the expression of *BIK1* and the *PEPs* and *PEPRs*, upregulation of *RBOHD* appeared to be more spatially restricted to the cells closest to the colonization site. *RBOHD* expression quickly returned to control levels behind this small group of cells, while the activation of the other markers appeared to taper out more gradually, across a longer stretch of cells.

While imaging the fungal infected reporter lines, we also regularly observed a discoloration of the root in the colonized tissue (white bracket in Fig. 4E). This has previously been described as 'root browning', and has been observed in response to infection by *F. oxysporum* or treatment of the plant with SERINE-RICH ENDOGENOUS PEPTIDEs (Tintor *et al.*, 2020; Hou *et al.*, 2021). While it remains unclear what the exact cause for this discoloration is, redox/oxidative stress induced by ROS is one possibility. In our assay, the browning does not completely overlap with

the expression maximum of *RBOHD*, but as the colonization of the fungus progresses upward
through the vasculature, the *RBOHD* expression also extends along adjacent to the colonization
front. Therefore, the area showing root browning would be the region that last expressed *RBOHD*
at high levels, and thus could have been under redox/oxidative stress, but this requires further
examination.

Discussion

In this paper we present a GreenGate-based entry vector set with 75 plant immune system
promoters. In combination with the basic GreenGate kit from Lampropoulos et al. (2013) this is
already sufficient to build simple tools, such as transcriptional reporter lines, that can then be
probed for their responsiveness to a pathogen, elicitor, or other stimulant. Furthermore, owing to
the universal compatibility of these entry vectors with all other GreenGate-based systems, this
vector set can be used to screen for interactors of proteins in their native expression domain using
the proximity ligation kit from Goslin *et al.* (2021), or for tissue-specific gene knockouts by
combining them with the CRISPR-TSKO kit from Decaestecker *et al.* (2019), to name just two
possibilities.

As a straightforward proof-of-principle, we used some of the pGG-PIPs to create fluorescent
reporter lines for the PEP pathway and demonstrate how these can be used to gain insights into
spatial immunity – the highly localized activation of an immune pathway. Using these lines, we
showed that *PEP1*, and to some degree *PEP2*, were responsive to infection and colonization by
the vascular pathogen *Fo*5176. The danger signal PEP1 was upregulated in a specific group of
vascular cells immediately adjacent to the fungal colonization site, as were the two receptors
PEPR1 and *2*. We thus hypothesize that activation of the PEP pathway is part of the plant's
immune response to infection and colonization by *Fo*5176. Importantly, we could show the highly
targeted activation of this pathway in just a couple of vascular cells immediately adjacent to a
colonization site. Such a local response has rarely been shown *in vivo* and *in planta* before, and is
typically overlooked in large-scale transcriptomic analyses.

Since the PEPRs are unlikely to serve as pattern-recognition receptors (PRRs) to sense the
pathogen, it is more likely that the PEPRs are part of a larger complex containing the PRR co-
receptor BAK1, as well as other PRRs, such as MIK2 or FER, since the interaction between BAK1

Figure 5: Model of the PEP pathway in response to *Fo*5176

Alkalinization of the apoplast via *Fo*5176-derived alkalinization factors induces interaction of the PEPRs with their PRR co-receptors. Intracellularly the receptors phosphorylate and activate BIK1. BIK1 activates expression of PEP1, thereby amplifying the PEP signal and creating a positive feedback loop. BIK1 also activates other defense genes, possibly via a MAPK cascade, and activates RBOHD to produce an apoplastic ROS-burst, which is damaging to the pathogen, and functions as cell-cell communication signal.

and the PEPRs has been previously shown, and all of these PRRs have been implicated to function in the defense against *Fo*5176 (Yamada *et al.*, 2016; Wang *et al.*, 2022*a*) (Fig. 5). Possibly, these co-receptors are all co-localized in plasma membrane nanodomains to facilitate the rapid assembly of larger receptor platforms (Somssich *et al.*, 2015; McKenna *et al.*, 2019; Somssich, 2020; Gronnier *et al.*, 2022). Indeed, rather than functioning as a PRR sensing the pathogen directly, work from a recent publication suggests that the PEPRs act as extracellular pH sensors (Liu *et al.*, 2022). Alkalinization of the apoplast is a hallmark of activated immunity, and Liu *et al.*, (2022) show that the PEPRs specifically interact with their co-receptor BAK1 under alkaline conditions, while they only show a weak affinity for interaction under acidic conditions (Felix *et al.*, 1993; Liu *et al.*, 2022). Interestingly, *F. oxysporum* uses functional homologs of plant RAPID ALKALINIZATION FACTORs to induce alkalinization of the apoplast, as this also stimulates infectious growth of the fungus (Masachis *et al.*, 2016; Wang *et al.*, 2022*a*). Thus, the PEPRs pathway may act as pH sensors to counteract this action by the fungus, improving the plant's defense under these alkaline conditions. Extracellular binding of the PEPs to their receptors generally leads to the activation of the PEPRs intracellular kinase domain and trans-phosphorylation of the interacting cytoplasmic kinase BIK1 (Liu *et al.*, 2013). Indeed, we could show that our *BIK1* reporter is also activated in the same pattern as the *PEP1* and the *PEPR* reporters. BIK1 activation then facilitates downstream intracellular signaling (Yamada *et al.*,

2016). These downstream signaling events eventually result in the transcriptional activation of
defense genes, but also auto-activation of the *PEP* genes (Fig. 5). Since the PEPs do not have an
identifiable secretion signal that would direct them into the secretory pathway, it is assumed that
they are released into the apoplast indirectly when a pathogen breaches the plasma membrane and
causes local membrane disintegration (Fig. 5) (Yamaguchi and Huffaker, 2011; Yamada *et al.*,
2016). Once in the apoplast, they are then perceived by the PEPRs, creating a positive feedback-
loop of PEP-signaling and an auto-amplification of the plant's defense response (Fig. 5). A second
intracellular pathway activated via BIK1 leads to activation of the NADPH oxidase RBOHD (Fig.
5) (Kadota *et al.*, 2014; Holmes *et al.*, 2018; Jing *et al.*, 2020). This pathway was proposed to be
primarily driven by PEP1 and the PEPR2 (Jing *et al.*, 2020) and indeed, *PEP1* and *PEPR2* are also
the two reporters showing the strongest response to colonization by *Fo*5176, while *PEP2* was
primarily upregulated in the cells around the vasculature. *PEPR1* was also activated in the same
pattern as *PEP1*, *PEPR2*, and *BIK1*, but at a much lower level compared to *PEPR2*. Further, we
could also confirm that *RBOHD* expression is activated in these same cells, indicating that this
BIK1-RBOHD-dependent pathway for ROS-release is also potentially involved in the defense
against *Fo*5176 (Fig. 5). The activation of RBOHD, and the resulting ROS-burst, would aid in the
plant's defense against *Fo*5176, with ROS being directly harmful to the pathogen. Additionally,
ROS is known to also function as a signal in cell-cell communication to prime the surrounding
tissue for impeding attack (Waszczak *et al.*, 2018). We believe that the local 'root browning' that
we observed in the infected roots is likely the result of this apoplastic ROS-burst.

A role for the PEP-pathway in the defense against *F. oxysporum* has previously been hypothesized
on the basis of transcriptomic data showing up- or downregulation of some of the components in
response to infection (Fallath *et al.*, 2017; Guo *et al.*, 2021; Wang *et al.*, 2022*a*). By investigating
the responsiveness of our transcriptional reporters for PEP1 and 2, as well as PEPR1 and 2, we
added spatial resolution to the available transcriptomic data. We could show that PEP1 and the
PEPRs act in a clearly defined group of vascular cells immediately next to the fungal colonization
site, and this localized response overlaps with the activation of downstream signaling factors and
a local ROS-burst. *PEP2* was also induced in response to colonization, but it appeared to be
upregulated in the outer tissues, not the infected vasculature. While this may indicate a role for
PEP2 in priming these neighboring tissues, the fact that neither of the two *PEPRs* were expressed
there, makes it unclear how PEP2 would be perceived in these cells. Importantly, when Fallath *et*

al. (2017) investigated the transcriptional response of *med* mutants to infection by *Fo*5176, they

also only noted a deregulation of *PEP1* in these mutants, but not *PEP2*, indicating that *PEP1* is

indeed the major PEP signal acting in response to colonization by *Fo*5176.

Finally, we believe that the pGG-PIP plant immune promoter resource introduced in this article

will be valuable and helpful to the community for various research approaches. We will make the

75-plasmid collection available via AddGene for quick and simple distribution to the community.

We aim to provide further detailed 'spatial immunity' information by using these pGG-PIPs in our

future work.

Material & Methods

Cloning of the pGG-PIPs and transcriptional reporter constructs

The pGG-PIP entry vectors are based on the pGGA000 vector from the original GreenGate kit

(Lampropoulos *et al.*, 2013). The different promoters were amplified from total cellular *A. thaliana*

(natural accession Columbia) DNA, extracted from rosette leaves with the Qiagen DNeasy Plant

Kit, and using the primers in supplementary table 1. For error-free amplification, the Phusion high-

fidelity DNA polymerase with proofreading (New England Biolabs) was used. The fragments were

then transferred into pGGA000 in a Golden Gate assembly reaction using the NEBridge Golden

Gate Assembly Kit (BsaI-HF v2) (New England Biolabs). To clone the transcriptional reporter

lines, we used the destination vector pGGZ003, as well as the donor vectors pGGD007 (*linker-*

NLS (Nuclear Localization Signal)), pGGE009 (*UBQ10* terminator), and pGGF005

(*pUBQ10::HygR:tOCS*) from the original GreenGate kit from Lampropoulos *et al.* (2013). pGGB-

mT2 (*mTurquoise2*) and pGGC-mT2 (*mTurquoise2*) were created by cloning the *mT2* coding

sequence from an AddGene-derived template into pGGB000 and pGGC000 from the original

GreenGate kit (Goedhart *et al.*, 2012; Lampropoulos *et al.*, 2013). In combination with our pGG-

PIPs, this yielded the *pPIP::mT2-mT2-NLS:tUBQ10:pUBQ10::HygR:tOCS* reporters.

Plant growth and transformation

A. thaliana Columbia plants (Somssich, 2018) were grown at 16-hour light conditions, with 21 °C

during the light hours, and 18 °C during the dark hours. Light intensity was 120 mmol m^{-2}s^{-1}, and

humidity was 70%. Plant transformation was done via the *Agrobacterium tumefaciens* host strain

GV3101 *pMP90 pSoup* carrying one of the plasmids in table 1 using the floral dip method (Holsters

et al., 1980; Koncz and Schell, 1986; Clough and Bent, 1998; Hellens *et al.*, 2000; Somssich,

2019). Individual resistant bacteria colonies were selected after two days of growth at 28° C on

YT plates (Miller, 1972) supplemented with 50 µg/ml rifampicin, 25 µg/ml gentamycin, 5 µg/ml

tetracyclin, and 100 µg/ml spectinomycin, then grown overnight in 250 ml of liquid 2×YT medium

with 50 µg/ml rifampicin, 25 µg/ml gentamycin, and 100 µg/ml spectinomycin, shaking at 200

rpm. Cells were harvested by centrifugation at 3200 g for 20 min, and resuspended in 300 ml of a

5% sucrose solution with 0.08% Silwet L-77. Plants were then dipped into the solution for

approximately 30 seconds with gentle agitation and laid out into a tray covered with cling wrap

overnight to maintain humidity. The next day they were returned to upright, and the procedure was

repeated after seven days. Once the siliques had ripened, the seeds were harvested and dried for at

least two weeks. For the selection of positive transformants, seeds were surface-sterilized using

75% ethanol with 0.1% Triton X-100 on a rotating incubator for at least two hours, after which the

seeds were decanted onto filter paper and the ethanol left to evaporate. Once the seeds were dry,

they were sprinkled onto a plate of half-strength basal Murashige & Skoog (MS) medium with

vitamins and 30 µg/ml hygromycin B, wrapped in aluminum foil, and stratified at 4 °C for three

days (Murashige and Skoog, 1962). They were then placed into a growth cabinet for 10 to 14 days,

after which healthy looking seedlings were transferred to half-strength MS plates without the

hygromycin, and grown for another two weeks, at which stage they were transferred to soil.

Fungal growth and transformation

The *F. oxysporum* f. sp. *conglutinans* strain 5176 was collected in 1971 from white cabbage

(*Brassica oleracea* var. *capitata* (L.)) in Australia (Wang *et al.*, 2022a). It has since been

maintained by the Brisbane Pathology (BRIP) Plant Pathology Herbarium in Queensland,

Australia under the accession number BRIP 5176 a. It was first used as a model pathogen for *A.*

thaliana by (Campbell *et al.*, 2003). To generate a *Fo*5176 line expressing cytoplasmic tdTomato

(tdT), an expression clone was generated by cloning the *tdT* coding sequence (amplified from an

AddGene-derived plasmid (Shaner *et al.*, 2004)) into the *Bgl*II cloning site of plasmid pLAU2

using Gibson assembly (Idnurm *et al.*, 2017). This places the gene under the control of a

constitutive promoter from the *actin* gene of ascomycete *Leptosphaeria maculans*. After

replication in *E. coli*, this plasmid pMAI32 was electroporated into *A. tumefaciens* strain EHA105

with selection on LB medium with kanamycin (50 µg/ml). *Fo*5176 was routinely cultured on

potato dextrose agar (PDA) plates. To generate spores for transformation, five plugs of about 5

mm diameter were inoculated into 50 ml half strength potato dextrose broth and cultured at 150

rpm for five days at room temperature, then filtered through miracloth, centrifuged at 3000 g for

5 min, and resuspended in sterile H_2O. The spores were then transformed with the *A. tumefaciens*

strain that had been cultured overnight in LB broth with kanamycin, using standard methods

(Idnurm *et al.*, 2017). That is, fungal spores and bacteria were mixed on induction medium plates

and cocultured for three days. Selection for transformants used an overlay of PDA supplemented

with hygromycin and cefotaxime (50 µg/ml and 100 µg/ml, respectively). Transformants that grew

through the overlaid medium, were subcultured onto PDA containing hygromycin and cefotaxime

and allowed to produce conidia, which were separated with a metal loop to generate strains derived

from a single conidium.

Plant-fungus co-cultivation and infection

To obtain fungal spores, *Fo*5176 was grown for at least seven days on PDA plates at room

temperature, at which time five pieces of roughly 1 mm^2 size were cut from the plate and dropped

into a flask containing 50 ml yeast nitrogen base (YNB) medium with 1% sucrose. The liquid

cultures were incubated for four days at room temperature with shaking at 120 rpm. The solution

was then filtered through miracloth, the spores harvested by centrifugation at 3000 rcf for 10 min

and resuspended in 25 ml sterile MilliQ water. For the co-cultivation of plant and fungus, *A.*

thaliana seedlings were grown on a vertical petri dish containing half-strength basal MS medium

with vitamins for 11 days and then transferred to a horizontal petri dish with a 2-3 cm strip of half-

strength basal MS medium with vitamins at the top end, while the rest of the plate was filled

roughly 2-3 mm high with liquid quarter-strength basal MS medium with vitamins (this is a setup

with slight modifications as described in (Tintor *et al.*, 2020)). The seedlings were placed onto the

thin MS medium strip at the top end, with the root in the liquid medium. Fungal spores were then

added to the liquid medium. The plates were covered with aluminum foil up until the leaves of the

plant, and then placed into a growth chamber.

Microscopy

376   We imaged the infection and the progression of colonization on a Leica M205 FA

377   stereomicroscope. Infection could usually be observed on day three after spored addition, and at

378   day 5 there was robust colonization. We usually imaged daily from day 3 to 11 dpi. For the

379   fluorescence coming from the plant 2xmT2 reporters, we used the Leica ET CFP (ET436/20x

380   ET480/40m) filter, and for the fungal tdT reporter the Leica ET mCHER (ET560/40x ET630/75m)

filter. We use 80× magnification for the images used in this paper. The settings for the imaging (illumination strength, exposure time, gain, etc.) were kept constant for all imaging sessions, to allow for at least semi-quantitative imaging, and to make the images at least relatively comparable. The images were recorded using the Leica Application Suite software, and processed using Fiji Is Just ImageJ (FIJI) and the GNU Image Manipulation Program (GIMP) (Schindelin *et al.*, 2012).

Data availability

We have donated the pGG-PIP vector collection to AddGene (Deposit-ID: 82532, Catalog-#: 196739-196813).

Acknowledgements

This work was funded by the Australian Research Council (grant no. DE200101560) and was further supported by a seed grant from the Melbourne University Botany Foundation. LW was supported by the China Scholarship Council. H-WC was supported by a Graduate Research Scholarship from the University of Melbourne. Imaging was done on instruments maintained by the Biological Optical Microscopy Platform (BOMP) and the BioSciences Microscopy Unit at the University of Melbourne. The authors would like to thank Dr. Imre E. Somssich for critical reading of the manuscript.

References

Acevedo-Garcia J, Gruner K, Reinstädler A, *et al.* 2017. The powdery mildew-resistant Arabidopsis mlo2 mlo6 mlo12 triple mutant displays altered infection phenotypes with diverse types of phytopathogens. Scientific reports **7**, 9319.

Andersen TG, Barberon M, Geldner N. 2015. Suberization-the second life of an endodermal cell. Current Opinion in Plant Biology **28**, 9–15.

Bird JE, Marles-Wright J, Giachino A. 2022. A User's Guide to Golden Gate Cloning Methods and Standards. ACS Synthetic Biology.

Birkenbihl RP, Kracher B, Ross A, Kramer K, Finkemeier I, Somssich IE. 2018. Principles and characteristics of the Arabidopsis WRKY regulatory network during early MAMP-triggered immunity. The Plant Journal.

Birkenbihl RP, Kracher B, Somssich IE. 2017*a*. Induced Genome-Wide Binding of Three

Arabidopsis WRKY Transcription Factors during Early MAMP-Triggered Immunity. The Plant

cell **29**, 20–38.

Birkenbihl RP, Liu S, Somssich IE. 2017*b*. Transcriptional events defining plant immune

responses. Current Opinion in Plant Biology **38**, 1–9.

Blümke A, Somerville SC, Voigt CA. 2013. Transient expression of the Arabidopsis thaliana

callose synthase PMR4 increases penetration resistance to powdery mildew in barley. Advances

in Bioscience and Biotechnology **04**, 810–813.

Cai Q, Qiao L, Wang M, He B, Lin F, Palmquist J, Huang S-D, Jin H. 2018. Plants send

small RNAs in extracellular vesicles to fungal pathogen to silence virulence genes. Science **360**,

1126–1129.

Calabria J, Wang L, Rast-Somssich MI, Chen H-W, Watt M, Persson S, Idnurm A,

Somssich M. 2022. Spatially distinct phytohormone responses of individual Arabidopsis thaliana

root cells to infection and colonization by Fusarium oxysporum. bioRxiv, 521292.

Campbell EJ, Schenk PM, Kazan K, Penninckx IAMA, Anderson JP, Maclean DJ,

Cammue BPA, Ebert PR, Manners JM. 2003. Pathogen-responsive expression of a putative

ATP-binding cassette transporter gene conferring resistance to the diterpenoid sclareol is

regulated by multiple defense signaling pathways in Arabidopsis. Plant physiology **133**, 1272–

84.

Cao Y, Liang Y, Tanaka K, Nguyen CT, Jedrzejczak RP, Joachimiak A, Stacey G. 2014.

The kinase LYK5 is a major chitin receptor in Arabidopsis and forms a chitin-induced complex

with related kinase CERK1. eLife **3**, 1–19.

Cho H, Lee J, Oh E. 2022. Leucine-Rich Repeat Receptor-Like Proteins in Plants: Structure,

Function, and Signaling. Journal of Plant Biology.

Chuberre C, Plancot B, Driouich A, Moore JP, Bardor M, Gügi B, Vicré M. 2018. Plant

Immunity Is Compartmentalized and Specialized in Roots. Frontiers in Plant Science **9**, 1–13.

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated

transformation of Arabidopsis thaliana. The Plant Journal **16**, 735–43.

Couto D, Zipfel C. 2016. Regulation of pattern recognition receptor signalling in plants. Nature

reviews. Immunology **16**, 537–552.

Decaestecker W, Buono RA, Pfeiffer ML, Vangheluwe N, Jourquin J, Karimi M, Van

Isterdael G, Beeckman T, Nowack MK, Jacobs TB. 2019. CRISPR-TSKO: A Technique for

Efficient Mutagenesis in Specific Cell Types, Tissues, or Organs in Arabidopsis. The Plant Cell

31, 2868–2887.

Denay G, Blümke P, Hänsch S, Weidtkamp-Peters S, Simon R. 2019. Over the rainbow: A

practical guide for fluorescent protein selection in plant FRET experiments. Plant Direct **3**,

e00189.

Dogra V, Singh RM, Li M, Li M, Singh S, Kim C. 2022. EXECUTER2 modulates the

EXECUTER1 signalosome through its singlet oxygen-dependent oxidation. Molecular Plant **15**,

438–453.

Duan Z, Liu W, Li K, Duan W, Zhu S, Xing J, Chen T, Luo X. 2022. Regulation of immune

complex formation and signalling by FERONIA, a busy goddess in plant–microbe interactions.

Molecular Plant Pathology, 1–6.

Engler C, Gruetzner R, Kandzia R, Marillonnet S. 2009. Golden Gate Shuffling: A One-Pot

DNA Shuffling Method Based on Type IIs Restriction Enzymes (J Peccoud, Ed.). PLOS ONE **4**,

e5553.

Engler C, Kandzia R, Marillonnet S. 2008. A One Pot, One Step, Precision Cloning Method

with High Throughput Capability (HA El-Shemy, Ed.). PLOS ONE **3**, e3647.

Fallath T, Kidd BN, Stiller J, Davoine C, Björklund S, Manners JM, Kazan K, Schenk PM.

2017. MEDIATOR18 and MEDIATOR20 confer susceptibility to Fusarium oxysporum in

Arabidopsis thaliana. PLOS ONE **12**, e0176022.

Felix G, Regenass M, Boller T. 1993. Specific perception of subnanomolar concentrations of

chitin fragments by tomato cells: induction of extracellular alkalinization, changes in protein

phosphorylation, and establishment of a refractory state. The Plant Journal **4**, 307–316.

Gao X, He P. 2013. Nuclear dynamics of Arabidopsis calcium-dependent protein kinases in

effector-triggered immunity. Plant signaling & behavior **8**, e23868.

Goedhart J, von Stetten D, Noirclerc-Savoye M, Lelimousin M, Joosen L, Hink MA, van

Weeren L, Gadella Jr. TWJ, Royant A. 2012. Structure-guided evolution of cyan fluorescent

proteins towards a quantum yield of 93%. Nature communications **3**, 751.

Gonçalves Dias M, Soleimani F, Monaghan J. 2022. Activation and turnover of the plant

immune signaling kinase BIK1: a fine balance. Essays in Biochemistry, 1–12.

Goslin K, Finocchio A, Wellmer F. 2021. A Golden Gate-based Plasmid Library for the Rapid

Assembly of Biotin Ligase Constructs for Proximity Labelling. bioRxiv, 464533.

Greenwood JR, Williams SJ. 2022. Guarding the central regulator of extracellular perception in

plants – A job for two. Cell Host & Microbe **30**, 1657–1659.

Gronnier J, Franck CM, Stegmann M, *et al.* 2022. Regulation of immune receptor kinase

plasma membrane nanoscale organization by a plant peptide hormone and its receptors. eLife **11**,

212233.

Guo L, Yu H, Wang B, *et al.* 2021. Metatranscriptomic Comparison of Endophytic and

Pathogenic Fusarium –Arabidopsis Interactions Reveals Plant Transcriptional Plasticity.

Molecular Plant-Microbe Interactions **34**, 1071–1083.

Hartley JL, Temple GF, Brasch MA. 2000. DNA Cloning Using In Vitro Site-Specific

Recombination. Genome Research **10**, 1788–1795.

Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM. 2000. pGreen: a versatile

and flexible binary Ti vector for Agrobacterium-mediated plant transformation. Plant molecular

biology **42**, 819–32.

Holmes DR, Grubb LE, Monaghan J. 2018. The jasmonate receptor COI1 is required for

AtPep1-induced immune responses in Arabidopsis thaliana. BMC Research Notes **11**, 555.

Holsters M, Silva B, van Vliet F, *et al.* 1980. The functional organization of the nopaline A.

tumefaciens plasmid pTiC58. Plasmid **3**, 212–230.

Hou S, Liu D, Huang S, *et al.* 2021. The Arabidopsis MIK2 receptor elicits immunity by

sensing a conserved signature from phytocytokines and microbes. Nature Communications **12**,

5494.

Huerta AI, Sancho-Andrés G, Montesinos JC, *et al.* 2023. The WAK-like protein RFO1 acts
as a sensor of the pectin methylation status in Arabidopsis cell walls to modulate root growth and
defense. Molecular Plant **2**, 33–47.

Idnurm A, Urquhart AS, Vummadi DR, Chang S, Van de Wouw AP, López-Ruiz FJ. 2017.
Spontaneous and CRISPR/Cas9-induced mutation of the osmosensor histidine kinase of the
canola pathogen Leptosphaeria maculans. Fungal Biology and Biotechnology **4**, 12.

Jia A, Huang S, Song W, *et al.* 2022. TIR-catalyzed ADP-ribosylation reactions produce
signaling molecules for plant immunity. Science **377**.

Jin T, Wu H, Deng Z, Cai T, Li J, Liu Z, Waterhouse PM, White RG, Liang D. 2022.
Control of root-to-shoot long-distance flow by a key ROS-regulating factor in Arabidopsis.
Plant, Cell & Environment, 0–2.

Jing Y, Shen N, Zheng X, Fu A, Zhao F, Lan W, Luan S. 2020. Danger-Associated Peptide
Regulates Root Immune Responses and Root Growth by Affecting ROS Formation in
Arabidopsis. International Journal of Molecular Sciences **21**, 4590.

Joglekar S, Suliman M, Bartsch M, Halder V, Maintz J, Bautor J, Zeier J, Parker JE,
Kombrink E. 2018. Chemical Activation of EDS1/PAD4 Signaling Leading to Pathogen
Resistance in Arabidopsis. Plant & Cell Physiology **59**, 1592–1607.

Journot-Catalino N, Somssich IE, Roby D, Kroj T. 2006. The transcription factors WRKY11
and WRKY17 act as negative regulators of basal resistance in Arabidopsis thaliana. The Plant
cell **18**, 3289–3302.

Kadota Y, Sklenar J, Derbyshire P, *et al.* 2014. Direct regulation of the NADPH oxidase
RBOHD by the PRR-associated kinase BIK1 during plant immunity. Molecular cell **54**, 43–55.

Kaiser S, Eisa A, Kleine-Vehn J, Scheuring D. 2019. NET4 Modulates the Compactness of
Vacuoles in Arabidopsis thaliana. International Journal of Molecular Sciences **20**, 4752.

Kim SH, Lam PY, Lee M-H, Jeon HS, Tobimatsu Y, Park OK. 2020. The Arabidopsis R2R3
MYB Transcription Factor MYB15 Is a Key Regulator of Lignin Biosynthesis in Effector-
Triggered Immunity. Frontiers in Plant Science **11**, 1–10.

Koncz C, Schell J. 1986. The promoter of TL-DNA gene 5 controls the tissue-specific
expression of chimaeric genes carried by a novel type of Agrobacterium binary vector.
Molecular & General Genetics **204**, 383–396.

Kümpers BMC, Han J, Vaughan-Hirsch J, *et al.* 2022. Dual expression and anatomy lines
allow simultaneous visualization of gene expression and anatomy. Plant Physiology **188**, 56–69.

Lampropoulos A, Sutikovic Z, Wenzl C, Maegele I, Lohmann JU, Forner J. 2013.
GreenGate - a novel, versatile, and efficient cloning system for plant transgenesis. PLOS ONE **8**,
e83043.

Lee D-K, Parrott DL, Adhikari E, Fraser N, Sieburth LE. 2016. The Mobile bypass Signal
Arrests Shoot Growth by Disrupting Shoot Apical Meristem Maintenance, Cytokinin Signaling,
and WUS Transcription Factor Expression. Plant Physiology **171**, 2178–2190.

Liu J, Liu B, Chen S, *et al.* 2018. A Tyrosine Phosphorylation Cycle Regulates Fungal
Activation of a Plant Receptor Ser/Thr Kinase. Cell Host & Microbe **23**, 241-253.e6.

Liu L, Song W, Huang S, *et al.* 2022. Extracellular pH sensing by plant cell-surface peptide-
receptor complexes. Cell, 1–15.

Liu W, Wang C, Wang G, Ma Y, Tian J, Yu Y, Dong L, Kong Z. 2019. Towards a better
recording of microtubule cytoskeletal spatial organization and dynamics in plant cells. Journal of
Integrative Plant Biology **61**, 388–393.

Liu Z, Wu Y, Yang F, Zhang Y, Chen S, Xie Q, Tian X, Zhou J-M. 2013. BIK1 interacts
with PEPRs to mediate ethylene-induced immunity. Proceedings of the National Academy of
Sciences of the United States of America **110**, 6205–6210.

Long Y, Stahl Y, Weidtkamp-Peters S, Smet W, Du Y, Gadella Jr. TWJ, Goedhart J,
Scheres B, Blilou I. 2018. Optimizing FRET-FLIM Labeling Conditions to Detect Nuclear
Protein Interactions at Native Expression Levels in Living Arabidopsis Roots. Frontiers in Plant
Science **9**, 1–13.

Masachis S, Segorbe D, Turrà D, *et al.* 2016. A fungal pathogen secretes plant alkalinizing
peptides to increase infection. Nature Microbiology **1**, 16043.

McKenna JF, Rolfe DJ, Webb SED, Tolmie AF, Botchway SW, Martin-Fernandez ML,

Hawes C, Runions J. 2019. The cell wall regulates dynamics and size of plasma-membrane

nanodomains in Arabidopsis. Proceedings of the National Academy of Sciences of the United

States of America, 201819077.

Meng F, Ellis T. 2020. The second decade of synthetic biology: 2010–2020. Nature

Communications **11**, 5174.

Miller JH. 1972. Formulas and recipes. Experiments in molecular genetics. Cold Spring Harbor,

New York: Cold Spring Harbor Laboratory, 433.

Morales J, Kadota Y, Zipfel C, Molina A, Torres MÁ. 2016. The Arabidopsis NADPH

oxidases RbohD and RbohF display differential expression patterns and contributions during

plant immunity. Journal of experimental botany **67**, 1663–76.

Murashige T, Skoog F. 1962. A Revised Medium for Rapid Growth and Bio Assays with

Tobaoco Tissue Cultures. Physiologia Plantarum **15**, 473–497.

Ngou BPM, Ahn H, Ding P, Jones JDG. 2021. Mutual potentiation of plant immunity by cell-

surface and intracellular receptors. Nature **592**, 110–115.

Patil M, Senthil-Kumar M. 2020. Role of Plant Kinases in Combined Stress. Protein Kinases

and Stress Signaling in Plants. Wiley, 445–458.

Petutschnig E, Anders J, Stolze M, *et al.* 2022. EXTRA LARGE G-PROTEIN2 mediates cell

death and hyperimmunity in the chitin elicitor receptor kinase 1-4 mutant. Plant Physiology **5**.

Rhodes J, Yang H, Moussu S, Boutrot F, Santiago J, Zipfel C. 2021. Perception of a

divergent family of phytocytokines by the Arabidopsis receptor kinase MIK2. Nature

Communications **12**, 705.

Saijo Y, Loo EP, Yasuda S. 2018. Pattern recognition receptors and signaling in plant-microbe

interactions. The Plant Journal **93**, 592–613.

Salguero-Linares J, Serrano I, Ruiz-Solani N, Salas-Gómez M, Phukan UJ, González VM,

Bernardo-Faura M, Valls M, Rengel D, Coll NS. 2022. Robust transcriptional indicators of

immune cell death revealed by spatiotemporal transcriptome analyses. Molecular Plant **15**,

1059–1075.

Schindelin J, Arganda-Carreras I, Frise E, *et al.* 2012. Fiji: an open-source platform for
biological-image analysis. Nature methods **9**, 676–82.

Schürholz A-K, Lopez-Salmeron V, Li Z, *et al.* 2018. A Comprehensive Toolkit for Inducible,
Cell Type-Specific Gene Expression in Arabidopsis. Plant Physiology, pp.00463.2018.

Shaner NC, Campbell RE, Steinbach PA, Giepmans BNG, Palmer AE, Tsien RY. 2004.
Improved monomeric red, orange and yellow fluorescent proteins derived from Discosoma sp.
red fluorescent protein. Nature biotechnology **22**, 1567–72.

Somssich M. 2018. A short history of Arabidopsis thaliana (L.) Heynh. Columbia-0. PeerJ
Preprints **e26931v3**, 1–7.

Somssich M. 2019. A short history of plant transformation. PeerJ Preprints, 1–28.

Somssich M. 2020. FERONIA is the new BAK1, and membrane nanodomains are the new thing
- but the flg22/FLS2/BAK1 complex continues to serve as a platform for new discoveries.
preLights, 23746.

Somssich M. 2022. The Dawn of Plant Molecular Biology: How Three Key Methodologies
Paved the Way. Current Protocols **2**, 1–19.

Somssich M, Ma Q, Weidtkamp-Peters S, Stahl Y, Felekyan S, Bleckmann A, Seidel CAM,
Simon R. 2015. Real-time dynamics of peptide ligand-dependent receptor complex formation in
planta. Science Signaling **8**, 1–9.

Stegmann M, Monaghan J, Smakowska-Luzan E, Rovenich H, Lehner A, Holton NJ,
Belkhadir Y, Zipfel C. 2017. The receptor kinase FER is a RALF-regulated scaffold controlling
plant immune signaling. Science **355**, 287–289.

Sun T, Zhang Y. 2021. Short- and long-distance signaling in plant defense. The Plant Journal
105, 505–517.

Tanaka T, Ikeda A, Shiojiri K, *et al.* 2018. Identification of a Hexenal Reductase That
Modulates the Composition of Green Leaf Volatiles. Plant Physiology **178**, 552–564.

Tintor N, Paauw M, Rep M, Takken FLW. 2020. The root-invading pathogen Fusarium

oxysporum targets pattern-triggered immunity using both cytoplasmic and apoplastic effectors.

New Phytologist **227**, 1479–1492.

Tseng Y, Scholz SS, Fliegmann J, Krüger T, Gandhi A, Furch ACU, Kniemeyer O,

Brakhage AA, Oelmüller R. 2022. CORK1, A LRR-Malectin Receptor Kinase, Is Required for

Cellooligomer-Induced Responses in Arabidopsis thaliana. Cells **11**, 2960.

Tsuda K, Somssich IE. 2015. Transcriptional networks in plant immunity. New phytologist

206, 932–47.

Wang L, Calabria J, Chen H-W, Somssich M. 2022*a*. The Arabidopsis thaliana – Fusarium

oxysporum strain 5176 pathosystem: an overview (M Höfte, Ed.). Journal of Experimental

Botany, erac263.

Wang W, Liu J, Mishra B, Mukhtar MS, McDowell JM. 2022*b*. Sparking a sulfur war

between plants and pathogens. Trends in Plant Science, 1–13.

Wang X, Meng H, Tang Y, Zhang Y, He Y, Zhou J, Meng X. 2022*c*. Phosphorylation of an

ethylene response factor by MPK3/MPK6 mediates negative feedback regulation of pathogen-

induced ethylene biosynthesis in Arabidopsis. Journal of Genetics and Genomics **49**, 810–822.

Wang J, Zhang Y, Wu J, Meng L, Ren H. 2013. AtFH16, an Arabidopsis type II formin, binds

and bundles both microfilaments and microtubules, and preferentially binds to microtubules.

Journal of integrative plant biology **55**, 1002–15.

Waszczak C, Carmody M, Kangasjärvi J. 2018. Reactive Oxygen Species in Plant Signaling.

Annual Review of Plant Biology **69**, 209–236.

Weber E, Engler C, Gruetzner R, Werner S, Marillonnet S. 2011. A Modular Cloning

System for Standardized Assembly of Multigene Constructs (J Peccoud, Ed.). PLOS ONE **6**,

e16765.

Wirthmueller L, Zhang Y, Jones JDG, Parker JE. 2007. Nuclear Accumulation of the

Arabidopsis Immune Receptor RPS4 Is Necessary for Triggering EDS1-Dependent Defense.

Current Biology **17**, 2023–2029.

Wu J, Liu Z, Zhang Z, *et al.* 2016. Transcriptional regulation of receptor-like protein genes by

environmental stresses and hormones and their overexpression activities in Arabidopsis thaliana.

Journal of Experimental Botany **67**, 3339–3351.

Wu R, Lucke M, Jang Y, Zhu W, Symeonidi E, Wang C, Fitz J, Xi W, Schwab R, Weigel

D. 2018. An efficient CRISPR vector toolbox for engineering large deletions in Arabidopsis

thaliana. Plant Methods **14**, 65.

Yamada K, Yamashita-Yamada M, Hirase T, Fujiwara T, Tsuda K, Hiruma K, Saijo Y.

2016. Danger peptide receptor signaling in plants ensures basal immunity upon pathogen-

induced depletion of BAK1. The EMBO journal **35**, 46–61.

Yamaguchi Y, Huffaker A. 2011. Endogenous peptide elicitors in higher plants. Current

Opinion in Plant Biology **14**, 351–357.

Yamaguchi Y, Huffaker A, Bryan AC, Tax FE, Ryan CA. 2010. PEPR2 Is a Second Receptor

for the Pep1 and Pep2 Peptides and Contributes to Defense Responses in Arabidopsis. The Plant

Cell **22**, 508–522.

Yamaguchi Y, Pearce G, Ryan CA. 2006. The cell surface leucine-rich repeat receptor for

AtPep1, an endogenous peptide elicitor in Arabidopsis, is functional in transgenic tobacco cells.

Proceedings of the National Academy of Sciences of the United States of America **103**, 10104–9.

Yang W, Devaiah SP, Pan X, Isaac G, Welti R, Wang X. 2007. AtPLAI Is an Acyl Hydrolase

Involved in Basal Jasmonic Acid Production and Arabidopsis Resistance to Botrytis cinerea.

Journal of Biological Chemistry **282**, 18116–18128.

Yang J, Duan G, Li C, Liu L, Han G, Zhang Y, Wang C. 2019. The Crosstalks Between

Jasmonic Acid and Other Plant Hormone Signaling Highlight the Involvement of Jasmonic Acid

as a Core Component in Plant Response to Biotic and Abiotic Stresses. Frontiers in Plant Science

10, 1–12.

Yang F, Kimberlin AN, Elowsky CG, Liu Y, Gonzalez-Solis A, Cahoon EB, Alfano JR.

2018. A Plant Immune Receptor Degraded by Selective Autophagy. Molecular Plant, 1–39.

Zipfel C, Oldroyd GED. 2017. Plant signalling in symbiosis and immunity. Nature **543**, 328–

336.

Supplementary information:

Supplementary table 1: List of primers used to clone the promoters

MAS-GG-ACS2p-F AACAGGTCTCAACCTCGGTCGATGTAAATGGATTAAATTTTATA
MAS-GG-ACS2p-R AACAGGTCTCATGTTGCTGTGTCAATTCTCACTTCTTTG
MAS-GG-ACS2pm-F AACAGGTCTCAGGTGTCCTCAAGGTTTCTGTTTCAAC
MAS-GG-ACS2pm-R AACAGGTCTCACACCAAATAAAGTTGATGTGGGGTC
MAS-GG-AOSp-F AACAGGTCTCAACCTACACTTAGACACCCCAATATTTTAGATTT
MAS-GG-AOSp-R AACAGGTCTCATGTTCTATTCGAAACAGTGGCGAGT
MAS-GG-AOSpm1-F AACAGGTCTCAGCCAATATTTTAGATTTTACTTTAAAGAAAT
MAS-GG-AOSpm1-R AACAGGTCTCATGGCGTCTCTAAGTGTTTTTTTTT
MAS-GG-AOSpm2-F AACAGGTCTCAGACACCTAAGTATTTTCTTTCCAACAA
MAS-GG-AOSpm2-R AACAGGTCTCATGTCCTTCTCTTTGAATAAACCTCA
MAS-GG-ARR5p-F AACAGGTCTCAACCTATATGATTTTTTCAAAAGAAAACACCATTTAGT
MAS-GG-ARR5p-R AACAGGTCTCATGTTATCAAGAAGAGTAGGATCGTGACTCGT
MAS-GG-ARR5pm-F AACAGGTCTCAGATTAGGATTATTCTTTATAGAATGTTTTGGTGC
MAS-GG-ARR5pm-R AACAGGTCTCAAATCGTCTCGGTTTTTACCTCTCAAATAGTAT
MAS-GG-ATG8Ap-F AACAGGTCTCAACCTTTAGCAGTGCTTAGTGAGCTTAAATTATAGTT
MAS-GG-ATG8Ap-R AACAGGTCTCATGTTAATTAATAAACTCGATCGTCTGCTAGATCG
MAS-GG-BAK1p-F AACAGGTCTCAACCTGTTTTTTGGAAACAGAGAGAAACTCA
MAS-GG-BAK1p-R AACAGGTCTCATGTTATCCTCAAGAGATTAAAAACAAACCCTA
MAS-GG-BIK1p-F AACAGGTCTCAACCTCGTTCCCAAATCTCGGTCAATTG
MAS-GG-BIK1p-R AACAGGTCTCATGTTCAAAGCTAAGAACAGATTCGTTTTCTTCTT
MAS-GG-BIK1pm-F AACAGGTCTCAGTACATTCATTTTGATTGGGATTTATCTTTTTA
MAS-GG-BIK1pm-R AACAGGTCTCAGTACAGACCAAAACAGTGGACTTGGG
MAS-GG-BPS1p-F AACAGGTCTCAACCTAAAAAGAGAGTCACTTGGGTTAAGTGATTT
MAS-GG-BPS1p-R AACAGGTCTCATGTTCTGATAAGTTTCAACTGAAAAAAACAAGAA
MAS-GG-CAD5p-F AACAGGTCTCAACCTGACTTGGGTTCAGTTAAAAATCTCA
MAS-GG-CAD5p-R AACAGGTCTCATGTTTTTGATGATTCTTTCTTCTTTCTTATC
MAS-GG-CERK1p-F AACAGGTCTCAACCTGTATGAAGAAGGGTAACAATTCAACTCTAA
MAS-GG-CERK1p-R AACAGGTCTCATGTTTGAAGCTTCCTTAGATTCCCCAGAGGAAGGGTGTCTGTT
MAS-GG-CIPP1p-F AACAGGTCTCAACCTCAAACCAGACATACTTGCAGCTTCTC
MAS-GG-CIPP1p-R AACAGGTCTCATGTTGAGAGTCAGATGTTCCACAAAAATCTACTC
MAS-GG-CIPP1pm-F AACAGGTCTCAGTTTGCAATCTGTACTAGAATAATATCGTGG
MAS-GG-CIPP1pm-R AACAGGTCTCAAAACGGATCTATATAACATATACGTATGCAATTAA
MAS-GG-CORK1p-F AACAGGTCTCAACCTGTACACTTATTAACATATATTTTTAATTTTGTG
MAS-GG-CORK1p-R AACAGGTCTCATGTTCGTCGACGACCAAAGATGTGAGA
MAS-GG-CORK1pm-F AACAGGTCTCAAGCCTGTTATTCTTAGTATTGCTTTCTATTTAAG
MAS-GG-CORK1pm-R AACAGGTCTCAGGCTTTAAACAATATAAAGCCCACAAG
MAS-GG-CPK29p-F AACAGGTCTCAACCTAATGAGGTAATGGAGGTTATCTTATCGA
MAS-GG-CPK29p-R AACAGGTCTCATGTTGTGAGCAAAGTAGATCGGTCTTCGA
MAS-GG-CPK29pm-F AACAGGTCTCAAGACACCCCCTCCAAGGGGCT
MAS-GG-CPK29pm-R AACAGGTCTCAGTCTCTAGGTTCTCCTATTCCTTTGCAA
MAS-GG-CPK5pro-F AACAGGTCTCAACCTACCATGTGACTACGACAACTACTGG
MAS-GG-CPK5pro-R AACAGGTCTCATGTTGAAACAATGGGAATTACCAAATCC
MAS-GG-CSLD2p-F AACAGGTCTCAACCTGACGACCAAGACTAGAGTTTTGGTTCG
MAS-GG-CSLD2p-R AACAGGTCTCATGTTAGTTAGGATCTAACTTGGCAGATCCCT
MAS-GG-DORN1p-F AACAGGTCTCAACCTGATGTAAAATTTGAAGCTTGAAGATGAAC
MAS-GG-DORN1p-R AACAGGTCTCATGTTGCAGATGATGAATCAGAGAGTCTGG
MAS-GG-EDS16p-F AACAGGTCTCAACCTGACTGCAGAGATCAATTTTCTTTTATTTTAT
MAS-GG-EDS16p-R AACAGGTCTCATGTTGCAGAAATTCGTAAAGTGTTTCTTGA
MAS-GG-EDS16pm1-F AACAGGTCTCACACCGCGTCCAACATTTTAAAACA
MAS-GG-EDS16pm1-R AACAGGTCTCAGGTGTCAGACTCTCAGCTGAACATAATT
MAS-GG-EDS16pm2-F AACAGGTCTCATCTGAAAGAGCCTAAGTGGGTTTCC
MAS-GG-EDS16pm2-R AACAGGTCTCACAGACCAGTTTTTATCATTTAAAAAATATTGTTA

MAS-GG-EDS1p-F AACAGGTCTCAACCTGTTTATCAGATTCCACGTACGATATGTTCTT
MAS-GG-EDS1p-R AACAGGTCTCATGTTGATCTATATCTATTCTCTTTTCTTTAGTGGACTTTC
MAS-GG-EDS1pm1-F AACAGGTCTCACGTGACCAAATCTGAAAACCCAAGT
MAS-GG-EDS1pm1-R AACAGGTCTCACACGTATAGAAGAAATCTACTACTTTAACTCTGTT
MAS-GG-EDS1pm2-F AACAGGTCTCACTCTCATGATGGGGTATTTTGGGTAAC
MAS-GG-EDS1pm2-R AACAGGTCTCAAGAGCATTTCAATGCAAAAATGGGT
MAS-GG-EFRp-F AACAGGTCTCAACCTATCTAGACGATTAAGTAATTGAGCATGTAAAAG
MAS-GG-EFRp-R AACAGGTCTCATGTTGTCGATTATAAAAAGATAAAAGAAAGGTTCTTCT
MAS-GG-ELI3p-F AACAGGTCTCAACCTTAAAGTCGATGTTCTATATGTATTCAAAATAAT
MAS-GG-ELI3p-R AACAGGTCTCATGTTATGGATAAATAATAAGCGAATGGGA
MAS-GG-ERF1p-F AACAGGTCTCAACCTCTCTCCCAATTGATATTTTTGTTATTTCT
MAS-GG-ERF1p-R AACAGGTCTCATGTTGTAGAAAAAATACTCTGTTTCTTGACTACTCTGT
MAS-GG-EX1p-F AACAGGTCTCAACCTCCGCCGTCTTAAGTGGAATTTGG
MAS-GG-EX1p-R AACAGGTCTCATGTTCGCCGGAGAGATGTGAGAGCG
MAS-GG-FERp-F AACAGGTCTCAACCTAGAAAAGTTAAGAGTGGGAACTGGGA
MAS-GG-FERp-R AACAGGTCTCATGTTCGATCAAGAGCACTTCTCCGG
MAS-GG-FH16p-F AACAGGTCTCAACCTCTGACCATAGTGTGGTAACTCAGATATTTT
MAS-GG-FH16p-R AACAGGTCTCATGTTGGATCAGGAACCAAACAGATGATT
MAS-GG-FLS2p-F AACAGGTCTCAACCTTTTTTGTTAGAGAGATTTTTGTTGTTTTTGTT
MAS-GG-FLS2p-R AACAGGTCTCATGTTGGTTTAGACTTTAGAAGAGTTGAAATTGTGG
MAS-GG-FMO1p-F AACAGGTCTCAACCTCGGAAAAATCCTTCGTCAATGTGTG
MAS-GG-FMO1p-R AACAGGTCTCATGTTCTGAGAGAGGTTATGCTAGAGAGAAGAGAG
MAS-GG-FRK1p-F AACAGGTCTCAACCTAAATTAAACGCCTTTTTATCAACAAC
MAS-GG-FRK1p-R AACAGGTCTCATGTTACTTAATTGAGCTGCTTTCTCTGG
MAS-GG-FRK1pm-F AACAGGTCTCACGTCTCTTTACATTTGTGATGTGGT
MAS-GG-FRK1pm-R AACAGGTCTCAGACGAACACTGATATAAAAAATCTCACA
MAS-GG-GLR25p-F AACAGGTCTCAACCTGATAAGAAGTGATTCAGCTGGGGTTT
MAS-GG-GLR25p-R AACAGGTCTCATGTTATTGATAGCTCGCAAGCTCAAATCTG
MAS-GG-GLR25pm-F AACAGGTCTCAACCTCTCTCACCCTGGTCGCAAA
MAS-GG-GLR25pm-R AACAGGTCTCAAGGTTTTCTCTTTTCAACGTACAATATTTACATGA
MAS-GG-GLR27p-F AACAGGTCTCAACCTCACACACTGGTCACTTAATGGTTTATTTGA
MAS-GG-GLR27p-R AACAGGTCTCATGTTCCAGATTGAGGAACTTTATGTCATTCTTTAAC
MAS-GG-GLR27pm-F AACAGGTCTCAACTCAAAATGATGCGTATCATTAATTCTATAGTTT
MAS-GG-GLR27pm-R AACAGGTCTCAGAGTTTGAAATTTTTAACAAAAGTCTTAAGTTATATCAG
MAS-GG-GPAT5p-F AACAGGTCTCAACCTAAAAAGCGTTTTAATTAGAGAGATTTTTGC
MAS-GG-GPAT5p-R AACAGGTCTCATGTTCTTTTGTTTTTTGCTCGAATATTATTTT
MAS-GG-HPCA1p-F AACAGGTCTCAACCTAAACATAAGAGAAAACGCAAGTTGATGA
MAS-GG-HPCA1p-R AACAGGTCTCATGTTCTTCAAACCCAAAAAGAACCTCTTATCA
MAS-GG-HRMp-F AACAGGTCTCAACCTCTGAATATCTTCTTTTGGTTGCTCTGA
MAS-GG-HRMp-R AACAGGTCTCATGTTCGTAATATCTCTCTGTTTTTGCTCTGTTTT
MAS-GG-LECRKVI2p-F AACAGGTCTCAACCTACAAAGTCAATTACTTCGAGTTTTTTTCTG
MAS-GG-LECRKVI2p-R AACAGGTCTCATGTTGGGTGAGCGAAGTAAAGAAGGAGATA
MAS-GG-LYK5p-F AACAGGTCTCAACCTATTTTCTGTTAAGTTTGAACATTTGGTTGTAA
MAS-GG-LYK5p-R AACAGGTCTCATGTTTTGTGGTGTTCTGATCTGAAGAGG
MAS-GG-LYK5pm-F AACAGGTCTCACAACAGACCAAGACCATCTTTATGTCC
MAS-GG-LYK5pm-R AACAGGTCTCAGTTGGATCTCATGTGAAAGAGACACATT
MAS-GG-LYM1p-F AACAGGTCTCAACCTATATCATCGGTAAGTCACTAGACTATTGAACG
MAS-GG-LYM1p-R AACAGGTCTCATGTTTTGTGTTTAGGGTTTTACGAAATTCAA
MAS-GG-MIK2p-F AACAGGTCTCAACCTGTAAATAACGTTGAACTCGCGG
MAS-GG-MIK2p-R AACAGGTCTCATGTTACAGTTGCAGATTATCTCTCTACGGTC
MAS-GG-MLO6p-F AACAGGTCTCAACCTAAATATACATTTGGTTGACATGTTTCTCATT
MAS-GG-MLO6p-R AACAGGTCTCATGTTAGAACTCACAGAACAGTTCCAAGCAAA
MAS-GG-MPK3p-F AACAGGTCTCAACCTAAAAAAATTCTGATCGAAAATAGCTTAC
MAS-GG-MPK3p-R AACAGGTCTCATGTTCTCTCTCAATTGATCAAAGTCGA
MAS-GG-MPK4p-F AACAGGTCTCAACCTGACTTGTTTGTGAATATAGAGGAAACATGTAATTAT
MAS-GG-MPK4p-R AACAGGTCTCATGTTCGGAGCAAAATTCCTCACAACAACG

MAS-GG-MPK6p-F AACAGGTCTCAACCTAACACAAGAGAAGAGATTTATTGCTTC
MAS-GG-MPK6p-R AACAGGTCTCATGTTGACCGGTAAAGATGAAAGCTTTT
MAS-GG-MYB15p-F AACAGGTCTCAACCTGATGAATTTGAATAAACTAAACAAAATT
MAS-GG-MYB15p-R AACAGGTCTCATGTTCTCTTTGATTTGTGATTGCTGATAAA
MAS-MYB15m-F AGAGGACCATGGACACCTGAAGAAGATCAAATCTTT
MAS-GG-MYB72p-F AACAGGTCTCAACCTACACGATCTCTTTTGAGATTTAAGAAG
MAS-GG-MYB72p-R AACAGGTCTCATGTTCTTATTACACTACTTTCTTCTCTATAGCTACC
MAS-GG-MYB72pm1-F AACAGGTCTCACGTTTTAAAACTTTACCTTATGTCCAATCTCT
MAS-GG-MYB72pm1-R AACAGGTCTCAAACGTGACGTAGCATGTGTGGGTC
MAS-GG-MYB72pm2-F AACAGGTCTCAGCTCTCTCTCTACGAGTGAAGTGCCT
MAS-GG-MYB72pm2-R AACAGGTCTCAGAGCCAAAAGCATGGAACGTACG
MAS-GG-NET4Ap-F AACAGGTCTCAACCTTTAATCCTCTTCTCGTACATCACAT
MAS-GG-NET4Ap-R AACAGGTCTCATGTTGGCTGCAAAAATCAATGGACC
MAS-GG-PAD4p-F AACAGGTCTCAACCTAATTAGGGTTTTATCAGATTAAAGAGATTTACTGATT
MAS-GG-PAD4p-R AACAGGTCTCATGTTGATTGGATATCGAGTAGAGAGTTGCAGA
MAS-GG-PDF12p-F AACAGGTCTCAACCTTCTACCAAAAATCTTTGGTGCTTGATC
MAS-GG-PDF12p-R AACAGGTCTCATGTTGATGATTATTACTATTTTGTTTTCAATGTATAGA
MAS-GG-PEP1p-F AACAGGTCTCAACCTGAAGTCAAAAATTGAGTCGAAAAATC
MAS-GG-PEP1p-R AACAGGTCTCATGTTGAGATCTGATAAGACAGAGGAAAACTT
MAS-GG-PEP2p-F AACAGGTCTCAACCTTGAAGCTCTTGTGAATAGAGAAGAGA
MAS-GG-PEP2p-R AACAGGTCTCATGTTGAAATCCAATAGTTTGGTGAGTTATC
MAS-GG-PEP3p-F AACAGGTCTCAACCTGCACTTTAAGTTACATTGTTTAGTCTAATTATT
MAS-GG-PEP3p-R AACAGGTCTCATGTTCGTTGACTTCTTAATCTTTTTTTGGGAA
MAS-GG-PEPR1p-F AACAGGTCTCAACCTAGAGAAGGAAAACAACCATGTATTCCAG
MAS-GG-PEPR1p-R AACAGGTCTCATGTTCTGAGTTTAAAGATCGAGAAACATGCAG
MAS-GG-PEPR1pm-F AACAGGTCTCAGAAACCAAACATCTCGTCATAAAAAAC
MAS-GG-PEPR1pm-R AACAGGTCTCATTTCTCTGTATACCAACGATTGTGAGA
MAS-GG-PEPR2p-F AACAGGTCTCAACCTAGTTTGAGATGGAGTTGCATTGTG
MAS-GG-PEPR2p-R AACAGGTCTCATGTTGAGATTAGAGCTCAAGAGACTGAAATAT
MAS-GG-PER5p-F AACAGGTCTCAACCTCAGTGCGTAGTAGTGAGTTTTCTTCA
MAS-GG-PER5p-R AACAGGTCTCATGTTATTTGTAGATCTCACTTGGTATATATTTCGTAC
MAS-GG-PER5pm-F AACAGGTCTCAGACGAATATATATAATTAGCTACTAAATTAAATT
MAS-GG-PER5pm-R AACAGGTCTCACGTCTCAGAACGAGTGAATGATTC
MAS-GG-PLP1p-F AACAGGTCTCAACCTCTGATCATCTAGCCTCTTCCC
MAS-GG-PLP1p-R AACAGGTCTCATGTTAATAGTTGATCGATCTTCTTTTGAGTTAA
MAS-GG-PMR4p-F AACAGGTCTCAACCTGCTCGATGTCGATTTGAGACGTAGT
MAS-GG-PMR4p-R AACAGGTCTCATGTTAGTAGCATGTGGTAGATCTTAGAAATTTCTCG
MAS-GG-PR1pm-F AACAGGTCTCACTCCCTCCATATAAAAAAGTTTGATTTTATAG
MAS-GG-PR1pm-R AACAGGTCTCAGGAGAATCATTTTATAAGTTAAAACAAGCTTG
MAS-GG-PR1pro-F AACAGGTCTCAACCTATATATAACGATCATTGATTAGTATATATACATATTG
MAS-GG-PR1pro-R AACAGGTCTCATGTTTTCTAAGTTGATAATGGTTATTGTTGT
MAS-GG-RALF23p-F AACAGGTCTCAACCTGGTGATTCCGGTTCCGACG
MAS-GG-RALF23p-R AACAGGTCTCATGTTTCTTCTGTACACTGTAGCTTTAGCTCTCTC
MAS-GG-RALF23pm-F AACAGGTCTCAGAGCACTCATAATTGTACAAAATAAAAGTAAATG
MAS-GG-RALF23pm-R AACAGGTCTCAGCTCTCCTTCCATGATTTGAGACTATTTC
MAS-GG-RBOHDp-F AACAGGTCTCAACCTGACTTGTTAAATTGCTCTCTTAGTCTTA
MAS-GG-RBOHDp-R AACAGGTCTCATGTTCGAATTCGAGAAACCAAAAAGATC
MAS-GG-RBHOFp-F AACAGGTCTCAACCTACCGGTTGAAAATAAGAGTGGTGGA
MAS-GG-RBOHFp-R AACAGGTCTCATGTTAGATCCAAAGTCGGAATTCAAAGAGTT
MAS-GG-RBOHFpm-F AACAGGTCTCATGCAGAAGATAGTGAAGATAGTTGCAGAA
MAS-GG-RBOHFpm-R AACAGGTCTCATGCAACTTTTATAGTTTTCGAACGAAAGTA
MAS-GG-RCD1p-F AACAGGTCTCAACCTGGAGGAGCAGATTGGACACCGT
MAS-GG-RCD1p-R AACAGGTCTCATGTTCTATATATTAACAATACTAAACCTATAACCTTGATAG
MAS-GG-RCD1pm-F AACAGGTCTCAATACGTCTCATATAGTTATGCTGATTCTTTCTTG
MAS-GG-RCD1pm-R AACAGGTCTCAGTATGATCCTGTAATATCATTCCTTCACAAAA
MAS-GG-RFO1p-F AACAGGTCTCAACCTATATTAACCATGCATGCAAACAAA

MAS-GG-RFO1p-R AACAGGTCTCATGTTTTTTTTTCTCTAATGACTTTTATGTATG
MAS-GG-RLP26p-F AACAGGTCTCAACCTGATTAAAGGATTGATCGGTAAACAAC
MAS-GG-RLP26p-R AACAGGTCTCATGTTGGTGTTTGTGATTGAACCAACAAGT
MAS-GG-RLP29p-F AACAGGTCTCAACCTCCAGCAAAAAGCTTCTTCTACTCAA
MAS-GG-RLP29p-R AACAGGTCTCATGTTAGGTTTTGGTGTAAGAGAGAGGAAAGA
MAS-GG-RPS4p-F AACAGGTCTCAACCTCGAGAACCTTGGCGAACTTGTCA
MAS-GG-RPS4p-R AACAGGTCTCATGTTGGCCCAAAAGCTTTTTCCCGGT
MAS-GG-SCOOP12p-F AACAGGTCTCAACCTAATAGGTTTCGAGTACTGTATTGATGTTTAACTG
MAS-GG-SCOOP12p-R AACAGGTCTCATGTTCTCGATCTTTATTTTTTTCTCGAGTTTAGA
MAS-GG-SOBIR1p-F AACAGGTCTCAACCTTTTCGATTTTTCTAATCTCACAGCTGTTC
MAS-GG-SOBIR1p-R AACAGGTCTCATGTTTAATTAGAGAAAGTTTCTTCTTGTGGATGTT
MAS-GG-SULTR41p-F AACAGGTCTCAACCTATGATCCATCACACGCCTGCCT
MAS-GG-SULTR41p-R AACAGGTCTCATGTTGATGGCTCTTGCGCACGCTTGG
MAS-GG-SULTR42p-F AACAGGTCTCAACCTGTAGCTTCCACGCCCTTGCCTAA
MAS-GG-SULTR42p-R AACAGGTCTCATGTTCGGAATTGGTGGGATAGAGAAGAAT
MAS-GG-TET8p-F AACAGGTCTCAACCTCGGATGTATCAAAGGTAAAAATATC
MAS-GG-TET8p-R AACAGGTCTCATGTTGGTTTAGATTCAGAGAGAAAGATTG
MAS-GG-TUB6p-F AACAGGTCTCAACCTATTTAGAGGGTGTTATTGGTTTGTG
MAS-GG-TUB6p-R AACAGGTCTCATGTTCTTCTATTTTATCTGAAATCAACATTACA
MAS-GG-TUB6pm1-F AACAGGTCTCATAACAAAAAGTTATGAATATTCACAGACATA
MAS-GG-TUB6pm1-R AACAGGTCTCAGTTATGGTTAACCGAGGATGAGC
MAS-GG-TUB6pm2-F AACAGGTCTCAGAGGCCATTTTTTTTTCCCGT
MAS-GG-TUB6pm2-R AACAGGTCTCACCTCATTGCGTATGACAATGCG
MAS-GG-VSP2p-F AACAGGTCTCAACCTTCTCTCTGGTTATATTTTGTTGCTGCTT
MAS-GG-VSP2p-R AACAGGTCTCATGTTGTTTTTTATGGTATGGTTTATTGTTTAGTTTGTG
MAS-GG-WAKL22p-F AACAGGTCTCAACCTATATTAACCATGCATGCAAACAAA
MAS-GG-WAKL22p-R AACAGGTCTCATGTTTTTTTTTTCTCTAATGACTTTTATGTATG
MAS-GG-WRKY11p-F AACAGGTCTCAACCTTAGTTCCAAAACCGCATTGACAT
MAS-GG-WRKY11p-R AACAGGTCTCATGTTGATGATTTCTTGGTCTGAGGATTTT
MAS-GG-WRKY11pm-F AACAGGTCTCAGACGAAACTGTTGATTGCTTTATTCC
MAS-GG-WRKY11pm-R AACAGGTCTCACGTCTCCTCAAAGTTCGAGGTTACT
MAS-GG-WRKY17p-F AACAGGTCTCAACCTGTCTCGCAGAGGTTATTTATCTACTTGGTT
MAS-GG-WRKY17p-R AACAGGTCTCATGTTGATGAGAAACCAGAGGAGAAACTTGAAG
MAS-GG-WRKY17pm1-F AACAGGTCTCAGGTCAACGATTCCCATGTCGCTAA
MAS-GG-WRKY17pm1-R AACAGGTCTCAGACCTAACCGACTAATATATATGATTGTCGTG
MAS-GG-WRKY17pm2-F AACAGGTCTCAAAGCAGACCAAACTTTGATTACTTTATTCCATA
MAS-GG-WRKY17pm2-R AACAGGTCTCAGCTTGAGTTGTGAGATATGTAGGGTCTTCTT
MAS-GG-WRKY33p-F AACAGGTCTCAACCTCGCTGCTTTTTCGAGATAGATAG
MAS-GG-WRKY33p-R AACAGGTCTCATGTTACGAAAAATGGAAGTTTGTTTTATAA
MAS-GG-WRKY40p-F AACAGGTCTCAACCTTGTGTATAACTATTATGCAGCCTTTTTCAA
MAS-GG-WRKY40p-R AACAGGTCTCATGTTGTAAATATATGTAGGATGAATCTTCGATATGGGT
MAS-GG-WRKY40pm-F AACAGGTCTCATACAAGAATAGGTACAGTCCTGGTTTGTG
MAS-GG-WRKY40pm-R AACAGGTCTCATGTAATTGTGAATAATAAAATCTTAATTCAGAT
MAS-GG-WRKY53p-F AACAGGTCTCAACCTATCTTGTGAGCTGATTCAAAGATTTC
MAS-GG-WRKY53p-R AACAGGTCTCATGTTTTAGTATATGATTCCCAAAATAGATTTTTT
MAS-GG-WRKY70p-F AACAGGTCTCAACCTCATTGTAGATATGATATATGAAGCTTCCCC
MAS-GG-WRKY70p-R AACAGGTCTCATGTTGTTAGTTTTGAGGAAGTTTTTGGTGAG
MAS-GG-WRKY70pm-F AACAGGTCTCAGTATCTCGCATATTAACTTAGGCTAGAGAGC
MAS-GG-WRKY70pm-R AACAGGTCTCAATACTATGATAAACCAGTTGGTTCTGTAGCG
MAS-GG-XLG2p-F AACAGGTCTCAACCTGAGTGGAGGAGCATAGTGTGATTATTTAC
MAS-GG-XLG2p-R AACAGGTCTCATGTTCTTCTTACCCAATCAAGCACACATACAA

Figure S1: Expression pattern of *PEP1* and *PEP2*
A-D shows the expression of *PEP1* (yellow). Robust expression is found in the inner root tissues of the mature DZ (A, B). From the young to mature DZ, expression gradually becomes stronger (C, D). No expression in the root tip, MZ or EZ. **E-H** shows the expression of *PEP2* (yellow). Expression is found in the EZ of the tip, then disappears in the young DZ (E, F), and returns to all tissues in the mature DZ (G, H). A, C, E, G are bright field images plus fluorescence, B, D, F, H are fluorescence only.

Figure S2: Expression pattern of *PEPR1* and *PEPR2*

A-D shows the expression of *PEPR1* (yellow). Weak expression is found in vasculature of the MZ, EZ and young DZ (A, B). In the mature DZ, weak expression is found in all tissues (C, D). **E-H** shows the expression of *PEPR2* (yellow). Strong expression is found in the vasculature of the DZ, starting in the root hair zone (E, F) and becoming stronger in the mature DZ (G, H). A, C, E, G are bright field plus fluorescence, B, D, F, H are fluorescence only.

Figure S3: Expression pattern of *BIK1* and *RBOHD*
A-D shows the expression of *BIK1* (yellow). Expression is found in the root tip around the meristem, including the root cap and part of the EZ (A, B). Expression appears stronger in the outer tissues compared to the vasculature. Further up the root, expression is robust in the mature DZ, but still stronger in the outer tissues (C, D). **E-H** shows the expression of *RBOHD* (yellow). *RBOHD* is expressed in all cells and tissues from the young DZ onwards (E, F). Expression is strongest in differentiated tissue (G, H), A, C, E, G are bright field plus fluorescence, B, D, F, H are fluorescence only.