Identification of quiescent FOXC2+ spermatogonial stem cells in adult mammals

Zhipeng Wang[1,3], Cheng Jin[1,3], Pengyu Li[1], Yiran Li[1], Jielin Tang[1], Zhixin Yu[1], Tao Jiao[1],
Jinhuan Ou[1], Han Wang[1], Dingfeng Zou[1], Mengzhen Li[1], Xinyu Mang[1], Jun Liu[1], Yan Lu[1],
Kai Li[1], Ning Zhang[2], Jia Yu[1], Shiying Miao[1], Linfang Wang[1] and Wei Song[1*]

[1] Department of Biochemistry and Molecular Biology, State Key Laboratory of Medical
Molecular Biology, Institute of Basic Medical Sciences Chinese Academy of Medical
Sciences, School of Basic Medicine Peking Union Medical College; Beijing, 100005, China.
[2] MRC Protein Phosphorylation and Ubiquitylation Unit, School of Life Sciences, University
of Dundee; Dundee, DD1 5EH, UK.
[3] These authors contributed equally: Zhipeng Wang, Cheng Jin.
* Correspondence: Wei Song

Email: songwei@ibms.pumc.edu.cn

Competing Interest Statement: The authors declare that they have no competing interests.

Classification: BIOLOGICAL SCIENCES and Developmental Biology

Keywords: FOXC2; spermatogonial stem cells; quiescence

Abstract

In adult mammals, spermatogenesis embodies the complex developmental process from spermatogonial stem cells (SSCs) to spermatozoa. At the top of this developmental hierarchy lie a series of SSC subpopulations. Their individual identities as well as the relationships with each other, however, remain largely elusive. Using single-cell analysis and lineage tracing, we discovered both in mice and humans the quiescent adult SSC subpopulation marked specifically by forkhead box protein C2 (FOXC2). All spermatogenic progenies can be derived from FOXC2[+] SSCs and the ablation of FOXC2[+] SSCs led to the depletion of the undifferentiated spermatogonia pool. During germline regeneration, FOXC2[+] SSCs were activated and able to completely restore the process. Germ cell specific *Foxc2* knockout resulted in an accelerated exhaustion of SSCs and eventually led to male infertility. Furthermore, FOXC2 prompts the expressions of negative regulators of cell cycle thereby ensures the SSCs reside in quiescence. Thus, this work proposes that the quiescent FOXC2[+] SSCs are essential for maintaining the homeostasis and regeneration of spermatogenesis in adult mammals.

Introduction

Through spermatogenesis, spermatozoa are generated from spermatogenic cells that are originated from spermatogonial stem cells (SSCs). It is critical for this process to be continuous and successful that SSCs are maintained in a homeostatic balance between self-renewal and differentiation (S. Sharma et al., 2019). The SSCs, which belong to a subgroup of undifferentiated spermatogonia (uSPG), exhibit significant heterogeneity and dynamic characteristics. In recent years, great insights into SSC behaviors and regulations have been provided by a body of pioneer works, especially with recent advances in single-cell gene-expression profiling, highlighting great heterogeneity of SSC and focusing on characterizing the nature of SSC states. Within the population of uSPG, a number of genes relatively higher expressed in primitive subfractions have been identified and well investigated, i.e., *Gfra1*, *ID4*, *Ret*, *Eomes*, *Pax7*, *Nanos2*, *Shisa6*, *T*, *Pdx1*, *Lhx1*, *Egr2* and *Plvap* (Aloisio et al., 2014; Guo et al., 2004; Hara et al., 2014; Helsel et al., 2017; Jijiwa et al., 2008; La et al., 2018; Nakagawa et al., 2021; Oatley et al., 2007; Sada et al., 2009; M. Sharma et al., 2019; Tokue et al., 2017). Particularly, *Gfra1*, *ID4*, *Eomes*, *Pax7*, *Nanos2*, and *Plvap* are further validated through lineage tracing experiment, which is considered to be a reliable method to study the development of stem cells. However, some essential and primitive sub-populations remain undiscovered, and the identification of which is of great significance for elucidating the developmental process of SSC renewal and its behavior in testis.

Adult stem cells (ASCs), as the undifferentiated primitive cells that can be found in nearly all types of tissues in mammals, are characteristic for a unique quiescent status reflected by both reversible cell cycle arrest and specific metabolic alterations (van Velthoven & Rando, 2019). The role of the quiescent stem cells is to avoid premature exhaustion and to allow the long-term maintenance of a functional stem cell pool (Cheung & Rando, 2013). The adult SSCs appear to share this characteristic, as revealed in recent single-cell RNA-sequencing (scRNA-seq) analysis in humans and mice, being largely non-proliferative while capable of reciprocating between the quiescent and activated status during homeostasis and regeneration (Guo et al., 2018; Hermann et al., 2018; Suzuki et al., 2021; Tan & Wilkinson, 2019; Wang et al., 2018). This notion has also been significantly supported by the discoveries of important regulators such as USF1 (Faisal et al., 2019) and DNMT3L (Liao et al., 2014) as well as the pathways including the PI3K/MAPK and mTORC1 signalings (Suzuki et al., 2021). In addition, it is generally believed that cells in a quiescent state are supposed to be more resilient to genotoxic insults, thus theoretically the

quiescent SSCs can restore spermatogenesis upon such disturbance. However, by large, much
remain unknown as to the characteristics and regulation of the quiescence state of adult SSCs.
Thus, more important insights can be obtained through searching the quiescent SSC population
and defining the essential characteristics of this population as the foundation of successful
spermatogenesis.
Forkhead box protein C2 (FOXC2), a member of the forkhead/winged helix transcription factor
family, plays essential roles in the development of various tissues in mice (Bahuau et al., 2002;
Kume et al., 2001; Motojima et al., 2016; Sasman et al., 2012). Recently, FOXC2 has been found
to be expressed in A_s and A_{pr} spermatogonia of mouse testes and is required for SSC
maintenance (Wei et al., 2018). However, the role of FOXC2 in the development of adult SSCs
in vivo has yet to be explored. In this study, we identified a subpopulation of adult SSCs specifically
expressed by FOXC2. In adult mice, FOXC2[+] SSCs gave rise to all spermatogenic progenitor cells
that can complete the full spermatogenesis. Upon the loss of FOXC2[+] SSCs, the undifferentiated
spermatogonia pool was exhausted, eventually leading to defective spermatogenesis. Specifically,
FOXC2 is required for maintaining SSC quiescence by promoting the expression of negative
regulators of cell cycle. Moreover, the FOXC2[+] population endured the chemical insult with busulfan
and effectively restored spermatogenesis. Overall, we propose that FOXC2[+] represents the
quiescent state of the SSCs in adult mammals that is crucial for the homeostasis and regeneration
of SSCs.

Results

Identification of FOXC2$^+$ SSCs from the adult uSPGs. We performed single-cell RNA-seq (10x genomics) of the uSPG from adult mice testes marked by THY1, a widely recognized surface marker for uSPG with self-renewing and transplantable state (Hammoud et al., 2014; Kubota et al., 2004), to dissect the heterogeneity and developmental trajectory (Figure 1A, Figure 1—figure supplement 1A, B). Among 5 distinct clusters identified, Cluster1 was characterized by the high expression of stemness markers whereas other clusters were featured by progenitor or differentiating spermatogonia (dSPG) markers (Figure1B, Figure 1—figure supplement 1C, D). Primarily mapped to the extreme early point of the developmental trajectory, Cluster1 cells appeared quiescent and likely represented the primitive state of uSPG populations (Figure1B, Figure 1—figure supplement 1E-G). The top10 differentially expressed genes (DEGs) associated with Cluster1 are featured by SSC markers such as *Mcam* (Kanatsu-Shinohara et al., 2012), *Gfra1* (Hara et al., 2014), *Tcl1* and *Egr2* (Hermann et al., 2018; La et al., 2018) (Figure1C, Figure 1—figure supplement 2A, Figure 1—source data 1) in addition to six others expressed in different stages of germ cells and/or somatic cells, in which only FOXC2 was exclusively localized in the nucleus of a subgroup of ZBTB16$^+$ uSPG in mice (Buaas et al., 2004; Costoya et al., 2004) (Figure 1D, Figure 1—figure supplement 2B). More specifically, FOXC2 displayed differential expressions among various subtypes of uSPG, being more specific in A$_s$ (59.9%) than other subtypes including A$_{pr}$ (5.2%), A$_{pr-1}$ (4.1%), A$_{al4-1}$ (1.83%), A$_{al8-1}$ (1.5%), and A$_{al16-1}$ (1.67%) (Figure 1E). There was only a small fraction (5.1%) was active in proliferation as indicated by MKI67 (Figure 1F), suggesting that FOXC2$^+$ cells are primarily quiescent. Additionally, when examining the SSC markers validated previously by lineage tracing (La & Hobbs, 2019), we found that FOXC2 displays a higher level of co-localization with GFRA1 and EOMES than PAX7 and NEUROG3 (Nakagawa et al., 2007), indicating the FOXC2$^+$ cells contain but differ from the known SSC subsets (Figure 1G).

We next analyzed the expression of FOXC2 in adult human testis using the published scRNA-seq dataset (GSE112013) (Guo et al., 2018). As expected, FOXC2 was also specifically expressed in the human SSCs, most of which were MKI67$^-$ (Figure 1H, Figure 1—figure supplement 2C). Pseudotime analysis showed that the FOXC2$^+$ cells located at the start of the developmental trajectory with a proportion of about 90% that were MKI67$^-$ (Figure 1I). Immunofluorescence staining confirmed that FOXC2$^+$ cells were a subset of ZBTB16$^+$ spermatogonia in adult human testis, and most of them were MKI67$^-$ (Figure 1J), representing a quiescent subpopulation of SSCs in human testis(Clermont, 1966a, 1966b, 1969; Ehmcke & Schlatt, 2006; Ehmcke et al., 2006). These results suggest that FOXC2 is similarly expressed in the SSCs of adult human and mouse testis and may possess a conserved function.

FOXC2$^+$ SSCs sufficiently initiate and sustain spermatogenesis. We generated *Foxc2$^{iCreERT2/+}$;Rosa26$^{LSL-T/G/LSL-T/G}$* mice in which FOXC2$^+$ cells were specifically labeled with GFP to enable the progeny tracing after tamoxifen treatment (Figure 2—figure supplement 1) (Wang et al., 2015). Tamoxifen was introduced at 2-month of age, after which the FOXC2-expressing lineage (GFP$^+$) was tracked at d3 (day3), w1 (week1), w2, w4, w6, m4 (month4), m7, and m12 respectively (Figure 2A). At d3, the tracked cells were both GFP$^+$ and FOXC2$^+$ (Figure 2B) and constituted 0.027% of the total testicular cells as indicated by the fluorescence-activated cell sorting (FACS) analysis (Figure 2C). GFP$^+$ cells and THY1$^+$ cells sorted by FACS were then transplanted into testes of recipient mice pre-treated with busulfan, respectively. Two months after transplantation, FOXC2$^+$ cells generated 3.6 times greater number of colonies than the THY1$^+$ control (Figure 2D,

E), indicating that the FOXC2$^+$ cells possess higher stemness as convinced by stronger transplantable viability.

At w1, all GFP$^+$ cells were identified as uSPGs, encompassing A$_s$, A$_{pr}$, and A$_{al-4}$ (Figure 2F$_a$). Specifically, FOXC2$^+$ A$_s$ gave rise to 3 types of A$_{pr}$, i.e., FOXC2$^+$/FOXC2$^+$, FOXC2$^+$/FOXC2$^-$, and FOXC2$^-$/FOXC2$^-$ (Figure 2F$_{c1, b, c2, d2}$), which then either produced FOXC2$^+$ or FOXC2$^-$ A$_s$ through symmetric or asymmetric division (Figure 2F$_{c3, d1, f1}$), or developed into A$_{al}$ with no more than one FOXC2$^+$ cell in the chains (Figure 2F$_{e, f2}$). These results confirm that FOXC2$^+$ cells are capable of self-renewal to sustain the population as well as replenishing the uSPG pool by producing downstream progenies, thereby serving as SSCs. In the following 2-6 weeks, GFP$^+$ colonies further expanded and produced GFP$^+$ sperms in the epididymis, from which healthy GFP$^+$ offspring were given birth by C57BL/6J female recipients (Figure 2G). The GFP$^+$ colonies constituted 83.67%, 90.48%, 96.78%, 98.55%, and 99.31% of the total length of the seminiferous tubules at w6, m2, m4, m7, and m12 respectively (Figure 2H, I). All offspring were GFP$^+$ from m4 onwards (Figure 2J). Additionally, the EOMES$^+$, GFRA1$^+$ and PAX7$^+$ cells were all GFP$^+$ at w2, further confirming these progenies were derived from the FOXC2$^+$ cells (Figure 2K). Overall, FOXC2$^+$ SSCs can produce all subtypes of uSPG, thus initiating spermatogenesis in adult mice.

Specific ablation of the FOXC2$^+$ SSCs results in the depletion of uSPG pool. We then prepared *Foxc2$^{iCreERT2/+}$;Rosa26$^{LSL-DTA/+}$* mice to investigate the physiological requirement of FOXC2$^+$ SSCs in spermatogenesis (Wang et al., 2015). FOXC2$^+$ population in 2-month-old mice was specifically ablated with tamoxifen-induced diphtheria toxin (DTA). The testes of these mice were examined at d3, d7, and d14 post tamoxifen induction (Figure 3A). Gradual loss of weight in testes coincided with the reduction in the size of testes in all the mice while body weight was maintained (Figure 3B, C). Specifically, at d3, there were no detectable FOXC2$^+$ cells in addition to the decrease in the number of GFRA1$^+$, LIN28A$^+$ (Zheng et al., 2009) and ZBTB16$^+$ uSPG at the basement membrane of seminiferous tubules; at d14, all GFRA1$^+$, LIN28A$^+$ and ZBTB16$^+$ uSPG disappeared while vacuoles formed at the basement membrane with remaining spermatocytes and spermatids in the seminiferous lumen (Figure 3D–F, Figure 3—figure supplement 1). Meanwhile, the expression of DDX4 (Toyooka et al., 2000) and DAZL (Li et al., 2019) as germ cell markers was significantly reduced along with nearly undetectable expression of uSPG markers such as ZBTB16, LIN28A, GFRA1, RET, and NEUROG3 (Nakagawa et al., 2007) (Figure 3G). These results indicate an uSPG exhaustion as the result of the FOXC2$^+$ SSCs ablation, therefore supporting the critical role played by FOXC2$^+$ population in spermatogenesis.

FOXC2$^+$ SSCs are resistant to busulfan and indispensable for germline regeneration. Next, we examined the regenerative viability of FOXC2$^+$ SSCs. At d20 post busulfan treatment (20mg/kg), FOXC2$^+$ cells constituted the majority of uSPGs (Figure 4A). Following a sharp decrease in cell number in the first five days, ZBTB16$^+$ and GFRA1$^+$ cells began to recover from d25 while the number of FOXC2$^+$ cells remained stable (Figure 4B), indicating that this population is insensitive to busulfan. We then checked changes in the proportion of MKI67$^+$ cells in FOXC2$^+$ population after busulfan treatment (Figure 4C, D). At d30, the MKI67$^+$ proportion rose to 15.92%, indicating a higher level of proliferation, albeit the total cell number stayed static (Figure 4B, D), thereby becoming the driving force in restoring spermatogenesis. Up to d120, the MKI67$^+$ proportion had settled gradually back to the pre-treatment level, accompanied by the full recovery of spermatogenesis (Figure 4D). Further details of this process were revealed during lineage tracing (Figure 4E). Three days after tamoxifen induction, the 2-month-old *Foxc2$^{iCreERT2/+}$;Rosa26$^{LSL-T/G/LSL-T/G}$* mice were treated with busulfan. Consistent with the results

above, at d20, the survived uSPGs were predominantly GFP$^+$ (Figure 4F). Over 68.5% of the total
length of the seminiferous tubules were GFP$^+$ at m2, and this proportion rose to 95.43%, 98.41%,
and 99.27% at m4, m7, and m12 respectively (Figure 4G, H), which was comparable to the
proportion by tamoxifen induction alone (Figure 2I). From m4 onwards, nearly all germ cells,
spermatids, and their offspring were GFP$^+$ (Figure 4G, I). Together, these results confirm that
FOXC2$^+$ SSCs are indispensable for germline regeneration that is central to spermatogenesis
recovery from interruptions.

FOXC2 is essential for SSC maintenance in adult mice. We then focused on dissecting the role
of FOXC2 in the SSC maintenance using $Foxc2^{f/-};Ddx4^{Cre/+}$ mice (Gallardo et al., 2007) (Figure 5A).
No significant difference was observed in the expressions of various uSPG markers, including
ZBTB16 and LIN28A, between $Foxc2^{f/-};Ddx4^{Cre/+}$ and $Foxc2^{f/+}$ mice at the age of 1 week (Figure
5—figure supplement 1A). However, adult $Foxc2^{f/-};Ddx4^{Cre/+}$ mice displayed clear testis weight loss
without significant body weight loss (Figure 5B, C). Moreover, in these mice, we observed severe
degeneration of seminiferous tubules, reduced number of spermatids in the epididymis, and
decreased size of the uSPG population with age (Figure 5D–G) but without apparent signs of
apoptosis (Figure 5—figure supplement 1B). The 6-month-old $Foxc2^{f/-};Ddx4^{Cre/+}$ mice were infertile,
in which over 95% seminiferous tubules were Sertoli-only with hardly detectable expressions of
DAZL, DDX4, LIN28A, and ZBTB16 (Figure 5D–F, H). Therefore, FOXC2 is essential for
maintaining the SSC homeostasis and normal spermatogenesis in adult mice.

FOXC2 maintains the quiescent state of SSCs through negatively regulating the cell cycle.
We collected THY1$^+$ uSPGs from 4-month-old $Foxc2^{f/+}$ and $Foxc2^{f/-};Ddx4^{Cre/+}$ mice and compared
their transcriptome signatures revealed from scRNA-seq (Figure 6A). The pseudotime analysis
identified Cluster1, which represented the FOXC2-expressing SSCs in $Foxc2^{f/+}$ mice corresponding
to the FOXC2-deleting SSCs in the $Foxc2^{f/-};Ddx4^{Cre/+}$ mice, was specifically assigned to the
extremely early stage of the development trajectory in respective samples, which was validated by
the expression of corresponding markers (Figure 6B, Figure 6—figure supplement 1A, B).
Aggregated analysis of the overall uSPG populations showed that cells derived from
$Foxc2^{f/-};Ddx4^{Cre/+}$ mice were specifically associated with the late stage of the development
trajectory, as opposed to $Foxc2^{f/+}$ mice where nearly all the cells derived were concentrated at the
early stage of development (Figure 6C, Figure 6—figure supplement 1C). This implies that the loss
of FOXC2 prompts the SSCs to progress into a more differentiated stage. Further analysis of the
cells in Cluster1 revealed two distinct subclusters, i.e., Subclusters0 and Subclusters1 (Figure 6—
figure supplement 2A). Formed primarily by the Cluster1 cells derived from $Foxc2^{f/+}$ mice,
Subclusters0 was featured by stemness markers, while Subcluster1, representing the majority of
Cluster1 cells from $Foxc2^{f/-};Ddx4^{Cre/+}$ mice, was featured by progenitor markers (Figure 6—figure
supplement 2B, C). Consistently, pseudotime analysis showed that Cluster1 cells from $Foxc2^{f/+}$
mice projected a forward stage of the developmental trajectory indicated by stemness markers,
whereas Cluster1 cells from $Foxc2^{f/-};Ddx4^{Cre/+}$ mice were associated with a later stage of the
developmental trajectory (Figure6D, Figure 6—figure supplement 2D, E). More specifically, less
number of cells were found at the starting state1 in Cluster1 from $Foxc2^{f/-};Ddx4^{Cre/+}$ mice than in
$Foxc2^{f/+}$ mice, with rather more cells in the developmental progression (from state1 to state5),
especially at the advanced state5 (Figure 6E). Thus, FOXC2 deletion caused defective SSC
maintenance and committed the SSCs to a differentiation destiny. Further, there were 932 genes
down-regulated in Cluster1 cells derived from $Foxc2^{f/-};Ddx4^{Cre/+}$ mice in comparison to $Foxc2^{f/+}$ mice
(Figure 6F, Figure 6—source data 1), which were functionally associated with both stem cell

population maintenance and mitotic cell cycle (Figure 6G). Consistently, the GSEA analysis
revealed a more progressive cell cycle in Cluster1 upon *Foxc2*-knockout (Figure 6H), confirming
the role of FOXC2 in regulating the cell cycle of the SSCs.
We then performed Cleavage Under Targets and Tagmentation (CUT&Tag) sequencing to
explore the underlying mechanism (Kaya-Okur et al., 2020; Kaya-Okur et al., 2019), for which GFP$^+$
SSCs from *Foxc2$^{iCreERT2/+}$;Rosa26$^{LSL-T/G/LSL-T/G}$* mice 3 days after tamoxifen induction, representing
the FOXC2$^+$ SSCs, were isolated for CUT&Tag sequencing (Figure 7A). Specific peaks enriched
in the promoter region of 3629 genes (Figure 7B, C; Figure 7—source data 1) showed functional
enrichment in biological processes such as DNA repair and mitotic cell cycle regulation (Figure
7D). By overlapping with the 932 genes down-regulated in Cluster1 cells from *Foxc2$^{f/-}$;Ddx4$^{Cre/+}$*
mice, we obtained 306 genes as the candidates subjective to the regulation by FOXC2 (Figure 7E;
Figure 7—source data 1). Further, Gene Ontology (GO) enrichment analysis of these genes
highlighted a distinctive functional cluster (11 genes) focusing on the negative regulation of cell
cycle (Figure 7F; Figure 7—source data 1) (Bahar et al., 2002; Bakker et al., 2004; Gaudet et al.,
2011; Georgescu et al., 2014; Knudson & Korsmeyer, 1997; Nitiss, 2009; Raman et al., 2007; van
Oosten et al., 2005; Weaver et al., 2003; Yi et al., 2012). More specifically, significant peaks
enrichment at the promoter region were observed for these candidate genes (Figure 7G).
Meanwhile, as predicted using the JASPAR Scan function (binding potential >0.8), there showed
strong binding potential of FOXC2 towards these candidate genes (Figure 7I) via the binding motif
of FOXC2 (Figure 7H), which was further confirmed by the result from the CUT&Tag qPCR (Figure
7J). Overall results imply that FOXC2 may function as a gatekeeper that ensures the quiescent
state of SSCs by impeding cell cycle progression.
248
249 **Discussion**
250
In this work, a comprehensive analysis of uSPG populations with scRNA-seq and the following
lineage tracing study by whole-mount immunofluorescence assay led to the identification of
FOXC2$^+$ SSCs as a quiescent SSC subpopulation in adult mice. Further investigation through
functionality analysis confirmed FOXC2 is essential for SSC self-renewal and stemness, thereby is
required for maintaining the SSC population that is critical for continuous spermatogenesis.
Importantly, our data demonstrated that the colonies formed by FOXC2$^+$ cells constituted nearly
the total length of the seminiferous tubules (99.31%), implying that the FOXC2$^+$ SSCs can support
the complete spermatogenesis in adult mice.
GFRA1$^+$ A$_{pr}$ and A$_{al}$ cells are found to break randomly and a portion of them can return to the
stem cell state (Hara et al., 2014). Interestingly, our findings showed FOXC2 appeared in one of
the A$_{pr}$ or A$_{al}$ cells at times, therefore raising a possibility that the subset of GFRA1$^+$ cells that return
to stem cell state after intercellular bridge break, maybe FOXC2$^+$ due to different cell cycle state. If
so, based on both findings, GFRA1$^+$FOXC2$^+$ could represent a quiescent state whereas
GFRA1$^+$FOXC2$^-$ is proliferate active, which certainly requires further validation possibly through
multiple lineage tracing and live imaging.
We observed that the FOXC2$^+$ SSCs were almost all in a non-proliferative state (~94.9%), and
further revealed that FOXC2 functioned in the negative regulation of cell cycle progression, thus
confirming that FOXC2-expressing SSCs are quiescent SSC population in adult mice. The finding
that FOXC2 inhibited cell cycle and differentiation of SSCs in testis is consistent with that reported
in other tissues (Davis et al., 2004; Sabine et al., 2015). In general, the quiescent state is a
protective mechanism for stem cell storage and prevents stem cells from damage or depletion
under genotoxic stresses (Arai & Suda, 2007; Relaix & Zammit, 2012; Rodgers et al., 2014; S.

Sharma et al., 2019). In our work, after the busulfan treatment, the quantity of FOXC2$^+$ cells
remained stable and the survived uSPGs were predominantly FOXC2$^+$, indicating its insensitivity
to cytotoxic agents. However, the proportion of MKI67$^+$FOXC2$^+$ cells increased by 15.92% after 30
days of the busulfan treatment and decreased back to the pre-treatment level (5.08%) at 120 days,
implying that the quiescent FOXC2$^+$ cells are able to transform into the proliferative FOXC2$^+$ cells
to replenish the SSC pool to maintain the SSC homeostasis and normal spermatogenesis. We
further confirmed by lineage tracing analysis that FOXC2-expressing cells were the only remaining
SSC population and were responsible for germline regeneration after the busulfan treatment,
indicating that FOXC2$^+$ SSCs represent a functionally important stem cell population with
regenerative ability. In the future, more insights into the unique regulation of SSCs can be drawn
from studying and comparing the transition between the quiescent and proliferative states in
FOXC2$^+$ and other SSC subpopulations.
According to our findings, we propose a model for the maintenance of the FOXC2$^+$ SSC
subpopulation (Figure 7K). Under physiological conditions, FOXC2$^+$ A$_s$ cells (including
FOXC2$^+$GFRA1$^+$, FOXC2$^+$EOMES$^+$ cells, etc.) constitute the SSC pool, of which only a small
proportion (~5.1%) cells are proliferative while the majority remains quiescent (Figure 7Ka). This
population can divide symmetrically or asymmetrically into different A$_{pr}$ and A$_{al}$ (Figure 7Kb). Then
FOXC2$^+$ cells (Figure 7Kb) may break from the syncytial and return to A$_s$ state (Figure 7Ka) to
maintain the stable number of the SSC pool. FOXC2$^-$ progenies, derived from the FOXC2$^+$
population, form progenitors (Figure 7Kc) to support spermatogenesis. However, it requires
continuous supply from the FOXC2$^+$ population and is subject to exhaustion when the supply is
disrupted. In the context of regeneration conditions, the FOXC2$^+$MKI67$^-$ cells can survive and set
out the recovery process (Figure 7Kd). At the early stage, increasing proportion of FOXC2$^+$MKI67$^-$
cells is committed to transforming into proliferative FOXC2$^+$MKI67$^+$ cells, strengthening the supply
to the progenitors (Figure 7Ke). At the late recovery stage, MKI67$^+$/MKI67$^-$ ratio returns to the
physiological level in FOXC2$^+$ population (Figure 7Ka), leaving the total number of FOXC2$^+$ cells
stable, therefore maintaining the SSC homeostasis. However, it is necessary to perform more
investigation to further improve and modify this model to gain a complete understanding of the
connections between different SSC subpopulations in the testes of adult mice.
Based on our observation, FOXC2 seems nonessential for the transformation from gonocytes to
SSCs in infant mice, in contrast to its requirement for adult spermatogenesis. A recent study
showed that FOXC2$^+$ subpopulation in the postnatal mouse testis (<5 weeks) appeared more active
in proliferation than the adult counterpart (Wei et al., 2018). Such differential functionality might
reflect the difference in the physical nature of spermatogenesis between developmental stages.
For example, the maturity of spermatogenesis is still under development during the juvenile period
with a focus on establishing and expanding the SSC pool. Therefore, it would be interesting to
explore differences in individual functional contexts as well as the underlying regulatory
mechanisms. Meanwhile, FOXC2, highly conserved between mice and humans with 94% identity
in amino acid sequence (Miura et al., 1997), is also expressed in a subset of human adult SSCs,
raising the possibility of an evolutionarily conserved mechanism governing SSC homeostasis in
humans. Further work following this direction might be of great clinical significance specifically to
patients who suffer from infertility. Moreover, the developmental correlation between FOXC2$^+$ SSCs
and other SSC subpopulations proposed previously should be revealed via biological methods such
as multiple lineage tracing and live imaging. Collectively, our work here provides new insights into
the investigation of adult SSCs and serves as a reference for studying the homeostasis and
regeneration of other stem-cell systems.
319

320 **Materials and methods**
321

Mice
Animal experiments were approved by the Committee on Animal Care of the Institute of Basic
Medical Sciences, Chinese Academy of Medical Sciences and Peking Union Medical College. The
tab of animal experimental ethical inspection (No: ACUC-A01-2018-017) has been provided in
supplementary file. The 8-week-old C57BL/6J wild-type mice were used for magnetic-activated cell
sorting. The *Rosa26^{LSL-T/G/LSL-T/G}* mice (stock no. 007676), *Ddx4^{Cre/+}* mice (stock no. 000692) and
EGFP^{Tg/+} mice (stock no. 021930) were bought from the Jackson Laboratory. The *Foxc2^{iCreERt2}* mice
and the *Foxc2^{flox/flox}* (*Foxc2^{f/f}*) mice were constructed and bought from the Biocytogen. The
Rosa26^{LSL-DTA/+} mice were bought from GemPharmatech. All mice were housed and bred under
specific pathogen-free conditions (temperature: 22-26°C, humidity: 40-55%, 12-h light/dark cycle)
in the animal facility at the Institute of Basic Medical Sciences. DNA was isolated from the tails, and
the genotypes of the mice were checked using PCR with specific primers (Materials and methods—
source data 1). All mice were randomly assigned to experiments and no statistical methods were
used to predetermine sample size. The person performing the experiments did not know the sample
identity until after data analysis. No data were excluded from analyses and the data displayed
included a minimum of three independent experiments and a minimum of three biological replicates
for each independent experiment. The 8-week-old C57BL/6J WT mice were treated with busulfan
(40 mg/kg) and used as recipient mice 1 month later.
340

Magnetic-activated cell sorting (MACS)
The testes from 8-week-old C57BL/6J wild-type mice or 4-month-old *Foxc2^{f/+}* and *Foxc2^{f/-};Ddx4^{Cre/+}*
mice (n=4) were minced and digested in the collagenase type IV (1 mg/mL, Sigma) and DNase I
(500μg/mL, Sigma) at 37°C for 15 min. The cell suspension was pipetted up and down once every
5 minutes and the digestion process was stopped with DMEM (containing 10% FBS). The cell
suspension was filtered through a 40-μm nylon mesh, and after centrifugation, the cells were
resuspended in 8 mL PBS. The 15 mL conical centrifuge tubes were slowly overlayed with 2 mL of
70% Percoll solution, 2 mL of 30% Percoll solution, and then 2 mL of testicular cell suspension and
centrifuge at 600 × g for 10 min at 4 °C without using the centrifuge brake. After centrifugation, the
cells at the interface between the 70% and the 30% Percoll solution were carefully removed into
the new conical centrifuge tubes, washed with PBS, and then centrifuge at 600 × g for 10 min at
4 °C. After centrifugation, the cells were resuspended in 360μL MACS buffer, added with 40μL of
magnetic microbeads conjugated with anti-Thy-1 antibody (Miltenyi Biotec 130–049–101, Auburn,
CA), and mixed well. Incubate the cell suspension containing Thy-1 microbeads for 20 min at 4 °C.
Mix gently by tapping every 10 min. Add 20 mL of MACS buffer to the tube to dilute Thy-1
microbeads and centrifuge at 300 ×g for 10 min at 4 °C. Remove the supernatant completely and
resuspend in 2 mL of MACS buffer. Place the separation columns (MS Column; Miltenyi Biotec
130-042-201) in the magnetic field of the mini MACS Separation Unit (Miltenyi Biotec 130-142-102)
and rinse with 0.5 mL of MACS buffer. Apply the cell suspension to the columns (500μL/ column).
After the cell suspension has passed through the column and the column reservoir is empty, wash
the column with 0.5 mL of MACS buffer three times. Remove the column from the MACS Separation
Unit and elute the magnetically retained cells slowly into a 50 mL conical centrifuge tube with 1 mL
of MACS buffer using the plunger supplied with the column. Centrifuge the tube containing the cells
at 600 × g for 10 min at 4 °C and resuspend the cell pellet with 10 mL of MACS buffer for rinsing.
Repeat this step once. After the final rinsing step, resuspend cells in 0.04% BSA and count the cell
number.

Single-cell RNA-seq

The MACS-sorted THY1$^+$ cells were used for loading onto the Chromium Single Cell 3′ Chip kit v2 (10x Genomics, PN-120236) according to the instructions. Cell capturing and library preparation was performed following the kit instructions of the Chromium Single Cell 3' v2 Library and Gel Bead Kit (10x Genomics, PN-120237). In brief, 5000 cells were targeted for capture, and after cDNA synthesis, 10-12 cycles were used for library amplification. The libraries were then size-selected, pooled, and sequenced on a Novaseq 6000 (Illumina). Shallow sequencing was performed to access the library quality and to adjust the subsequent sequencing depth based on the capture rate and the detected unique molecular indices (UMI).

Single-cell RNA-seq data processing

Raw sequencing reads were processed using the Cell Ranger v.3.0.1 pipeline of the 10x Genomics platform. In brief, reads from each sample were demultiplexed and aligned to the mouse mm10 genome, and UMI counts were quantified for each gene per cell to generate a gene-barcode matrix. Default parameters were used. The UMI counts were analyzed using the Seurat R Package (Stuart et al., 2019) (v.3.0.1) following the Seurat pipeline. Cells with more than 200 detected genes or less than 10% mitochondria reads were retained. Genes not detected in at least 10 cells were removed from subsequent analysis. The resulting matrix was normalized, and the most variable genes were found using Seurat's default settings, then the matrix was scaled with regression against the mitochondria reads. The top 2000 variable genes were used to perform PCA, and Jackstraw was performed using Seurat's default settings. Variation in the cells was visualized by UMAP for the top principal components. Cell types were determined using marker genes identified from the literature (Kowalczyk et al., 2015). We used the Seurat function CellCycleScoring to determine the cell cycle phase, as this program determines the relative expression of a large set of G2-M and S-phase genes. After removing the undefined cells, the spermatogonia were used for trajectory analysis, and the single-cell pseudotime trajectory was constructed with the Monocle 2 package (v2.12.0) (Qiu, Hill, et al., 2017; Qiu, Mao, et al., 2017; Trapnell et al., 2014) according to the provided documentation. The Monocle function clusterCells was used to detect cell clusters between clusters. The Seurat function FindAllMarkers with default settings was used to find DEGs upregulated in each cluster compared to the other cells. The top200 DEGs of cluster1 were used for ordering cells, and the discriminative dimensionality reduction with trees (DDRTree) method was used to reduce the data to two dimensions. The dynamic expression patterns with the spermatogonial developmental trajectory of specific genes were visualized using the Monocle function plot_genes_in_pseudotime and plot_pseudotime_heatmap. The procession data of the adult human single-cell dataset was downloaded from Gene Expression Omnibus (GEO): GSE112013 (Guo et al., 2018) and the UMI counts were analyzed using the Seurat R Package (v.3.0.1) following the Seurat pipeline with the same parameters and functions as mentioned previously. According to the known markers, the germ cells characterized was used for trajectory analysis, and the single-cell pseudotime trajectory was constructed with the Monocle 2 package (v2.12.0) as mentioned previously.

CUT&Tag sequencing and analysis

CUT&Tag assay was performed using CUT&Tag 2.0 High-Sensitivity Kit (Novoprotein scientific Inc., Cat# N259-YH01). The detailed procedures were described in (Kaya-Okur et al., 2019; Wang et al., 2021). In brief, cells were harvested by trypsin and enriched by ConA-magnetic beads. 10,000 cells were re-suspended in 100 mL Dig-wash Buffer (20 mM HEPES pH 7.5; 150 mM NaCl;

0.5 mM Spermidine; 13 Protease inhibitor cocktail; 0.05% Digitonin) containing 2 mM EDTA and a
1:100 dilution of primary FOXC2 antibody. The primary antibody was incubated overnight at 4°C.
Beads were washed in Dig-wash Buffer 3 times and incubated with secondary antibody for 1 hour
at a dilution of 1:200. After incubation, the beads were washed 3 times in Dig-Hisalt Buffer (0.05%
Digitonin, 20 mM HEPES, pH 7.5, 300 mM NaCl, 0.5 mM Spermidine, 13 Protease inhibitor
cocktail). Cells were incubated with proteinA-Tn5 transposome at 25°C for 1 hour and washed 3
times in Dig-Hisalt buffer to remove unbound proteinA-Tn5. Next, cells were re-suspended in
100mL Tagmentation buffer (10 mM MgCl2 in Dig-Hisalt Buffer) and incubated at 37°C for 1 hour.
The tagmentation was terminated by adding 2.25 mL of 0.5 M EDTA, 2.75 mL of 10% SDS and 0.5
mL of 20 mg/mL Proteinase K at 55°C for 1 hour. The DNA fragments were extracted by phenol
chloroform and used for sequencing on an Illumina HiSeq instrument (Illumina NovaSeq 6000) to
generate 2 × 150-bp paired-end reads following the manufacturer's instructions.
Raw reads were analyzed by removing low-quality or adaptor sequences using Trim_galore
(v0.5.0) and cleaned reads were mapped to the reference genome mm10 using Bowtie2 (v2.2.5).
We used MACS2 (v2.1.2) to call peaks found in different groups. Homer (v4.11.1) de novo motif
discovery tool was used for finding the binding motifs of Foxc2 with the findMotifsGenome.pl
command. The binding potential of candidate target genes at the binding motif was predicated
using the JASPAR Scan function (binding potential >0.8). The peaks filtered by fold change more
than 5 and transcription start site (TSS) less than 3000 bp were annotated by R package Chip
Seeker for gene category analysis. R package Cluster profiler was used for gene function
annotation such as KEGG and GO analysis.
435
Enrichment analyses
Gene Ontology (GO) and KEGG pathway enrichment analyses were conducted using the
ClusterProfiler package (v3.12.0) (Yu et al., 2012) and the ClueGO app (v2.5.7) in Cytoscape
(v3.8.1) with default settings and a p-value cut-off of 0.05. GSEA enrichment analysis was
assessed using the GSEA (v4.0.2) algorithm with MSigDB (v7.0) with default settings. The signaling
pathways enriched by niche-derived paracrine factors and undifferentiated SPG-derived
membrane proteins in the DEGs of the four samples were characterized. Then for each niche cell
type, the niche-derived signaling pathways in all four samples were crossed with the SSC-derived
signaling pathways to identify the candidate signaling pathways pivotal to SSCs maintenance.
445
446 **Transplantation assay**
447 The 8-week-old C57BL/6J WT mice were treated with busulfan (40 mg/kg) and used as recipient
448 mice 1 month later. SSCs were transplanted into the testis of recipient mice (1 × 10³ cells/testis),
449 and two months after transplantation, the testes were harvested and observed under a
450 fluorescence microscope.
451
Fluorescence-activated cell sorting (FACS)
Single-cell suspensions were generated from testes or *in vitro* cultured SSCs. FACS was performed
using an SH800 machine (Sony Biotechnology) to isolate the GFP⁺ cells. Briefly, the GFP⁺ gating
area was based on the point of the fluorescence intensity axis where cells were considered as
being GFP⁺, set based on the background fluorescence intensity of a non-transgenic control testis
cell population.
458
Immunofluorescence

Mouse testes were fixed in 4% Paraformaldehyde (PFA) at 4°C overnight, dehydrated, embedded
in paraffin, and cut into 5-μm thick sections. The rehydrated mouse or human testis sections were
subjected to antigen retrieval, blocked in 5% BSA with 0.1% Triton X-100, and incubated with
primary antibody (Materials and methods—source data 1) at 4°C overnight, including the germ cell
marker DDX4, undifferentiated spermatogonia markers ZBTB16, LIN28A, ECAD (Tokuda et al.,
2007), GFRA1, EOMES, PAX7, progenitor marker NEUROG3, and spermatocyte marker SYCP3
(Yuan et al., 2000). After three 5-min washes in PBS, the sections were incubated with secondary
antibodies (Materials and methods—source data 1) and DAPI (Sigma) at 37°C for 1 hour. After
three 5-min washes in PBS, coverslips were then mounted on glass slides using anti-quencher
fluorescence decay (Solarbio). Images were captured using a Zeiss 780 laser-scanning confocal
microscope. Whole-mount immunofluorescence of seminiferous tubules was performed as
previously described (Di Persio et al., 2017). Briefly, seminiferous tubules were disentangled from
testicular biopsies and immediately fixed in 4% PFA at 4°C for 12 hours. After fixation, the
seminiferous tubules were permeabilized with 0.5% Triton X-100 in PBS and treated with 5% BSA
in PBS overnight at 4°C. After three 30-min washes, the seminiferous tubules were incubated with
primary antibody (Materials and methods—source data 1) overnight at 4°C. After three 30-min
washes, the seminiferous tubules were incubated with species-specific secondary antibodies and
DAPI at 4°C for 12 hours. After three 30-min washes, the seminiferous tubules were mounted on
slides with anti-quencher fluorescence decay (Solarbio) and observed with a Zeiss 780 laser-
scanning confocal microscope.
480
RNA isolation and quantitative RT-PCR analysis
Total RNA was extracted from the testes or cultured cells using the RNeasy kit (Qiagen), reverse-
transcribed using RevertAid First Strand cDNA Synthesis kit (Thermo), and processed for qRT-
PCR using PowerUp SYBR Green Master Mix (Applied Biosystems) and a LightCycler 480 system
(Roche) with gene-specific primers (Materials and methods—source data 1). Reactions were run
in triplicate and the mRNA levels were normalized to Gapdh and quantified using the delta-delta Ct
method. The values shown are mean ± s.e.m. from three biological replicates.
488
Tamoxifen inducible
According to a previous report for activation of iCre (M. Sharma et al., 2019), the mice were fed
with TD.130859 (TAM diet) for three days. The food was formulated for 400 mg tamoxifen citrate
per kg diet, which would provide ~40 mg tamoxifen per kg body weight per day.
Analyses of cyst length. The cyst length was obtained according to the previous report (Nakagawa
et al., 2010). Briefly, to determine the cyst length, after immunofluorescence staining with anti-E-
CAD antibody, the whole mount seminiferous tubule specimens were observed under a
fluorescence microscope. The E-CAD staining coupled with staining for FOXC2 enabled us to
reliably identify syncytial cysts of FOXC2$^+$ cells.
498
499 **Analyses of cell density**
500   The cell density was counted according to a previous report (Tegelenbosch & de Rooij, 1993).
501   Briefly, the densities of the ZBTB16$^+$, GFRA1$^+$, LIN28A$^+$, or FOXC2$^+$ cells were measured on the
502   seminiferous tubules with whole-mount staining, the numbers of which per 1000 Sertoli cells were
503   determined.
504
505 **Sperm counts**

506 Total sperm counts were obtained according to the previous report (Roy et al., 2007). Briefly,
507 epididymal caput and cauda were minced and incubated in prewarmed M16 medium (Sigma-
508 Aldrich) at 37°C in air containing 5% CO_2 for 30 min to allow the sperm to swim out. Then, the
509 sperm were diluted in water and counted using a hemocytometer.
510

Histology, evaluation of degenerating tubules
Testes of WT and mutant mice were fixed with PFA fixative and processed for paraffin-embedded
section preparation (5 µm thick) and hematoxylin and eosin staining, according to standard
procedures. The percentage of degenerating seminiferous tubules was calculated based on the
cross-sections of seminiferous tubules (n > 200) that appeared on one transverse section for each
testis. In normal (WT) mouse testes, four generations of germ cells, each synchronously
progressing through spermatogenesis, form cellular associations of fixed composition (called
seminiferous epithelial stages). In the testes of $Foxc2^{f/-};Ddx4^{Cre/+}$ mice, a few tubule cross-sections
lacked one or more out of the four germ cell layers, which was defined as "degenerative tubules"
in this study.
521

Statistical analysis
All statistical analyses were performed using GraphPad Prism (v7.0). All experiments were
repeated at least three times, and data for evaluated parameters are reported as mean ± s.e.m.
The p-values were obtained using two-tailed unpaired Student's t-tests or one-way ANOVA
followed by Tukey test (ns represents p-value > 0.05, * represents p-value < 0.05, ** represents p-
value < 0.01, *** represents p-value < 0.001, and **** represents p-value < 0.0001).
528

Acknowledgments
This work was supported by the National Key Research and Development Program of China grant
(2022YFA0806302, 2018YFC1003500, 2019YFA0801800), CAMS Innovation Fund for Medical
Sciences (CIFMS, 2021-I2M-1-019, 2017-I2M-3-009), National Natural Science Foundation of
China grant (92268111, 31970794, 32000586, 31725013, 32200646), and State Key Laboratory
Special fund from the Ministry of Science grant (2060204).

Materials and methods—source data 1. Excel spreadsheet with the primers and antibodies
used in this study.
538

Data Availability. All data are available in the main text or supplementary materials. The scRNA-
seq and CUT&Tag sequencing data produced in this study have been uploaded to the GEO
(https://www.ncbi.nlm.nih.gov/geo) with accession codes GSE183163, GSE180729, and
GSE180926. The expression of FOXC2 in adult human testis used the scRNA-seq dataset
GSE112013 from previously published "The Human Testis Cell Atlas via Single-cell RNA-seq" by
Jingtao Guo (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE112013) (Guo et al., 2018).
All of the R packages were available online and the code was used according to respective R
packages documentation as described in the Methods. The MSigDB (v.7.0) used in this study is
available at https://www.gsea-msigdb.org/gsea/msigdb.
548

Figures

Figure 1 with 2 supplements

Identification of the FOXC2⁺ SSCs in adult mouse and human testis. (**A**) Schematic illustration
of the single-cell analysis workflow. (**B**) t-SNE plot and developmental trajectory of all uSPG,
colored by cluster. (**C**) Heatmap of the Top10 DEGs in Cluster1. (**D**) Immunostaining for ZBTB16
(red), FOXC2 (green), and DAPI (blue) in testicular paraffin sections from wild-type adult C57 mice.
Scale bar, 50 μm; C57, C57BL/6J. (**E**) The proportion of FOXC2⁺ cells in different uSPG subtypes.
(**F**) Immunostainings for MKI67 (red), FOXC2 (green), and DAPI (blue) in adult mice testis and the
proportion of MKI67⁺ cells in FOXC2⁺ population (n=10). Scale bar, 50 μm; values, mean ± s.e.m.;
p-values were obtained using two-tailed t-tests (****p-value < 0.0001). (**G**) The co-expression
proportion between the FOXC2 and differential known SSCs makers (n=10). (**H**) t-SNE plot of germ
cells in adult human testis (GSE112013), colored by germ cell type. Feature plot showing the
expression patterns of FOXC2 and MKI67 in human germ cells. (**I**) The developmental trajectory of
the human germ cells, colored by germ cell type, FOXC2 expression cells (red), or MKI67
expression cells (red). (**J**) Immunostaining for ZBTB16/MKI67 (red), FOXC2 (green), and DAPI
(blue) in testicular paraffin sections from adult humans.

567 **Figure 1—figure supplement 1**

568
Validation and characterization of the MACS-sorted THY1⁺ uSPG from wild-type adult C57
mice. (**A**) Immunostainings of DAPI (blue), THY1 (green), and ZBTB16 (red) in the MACS-sorted
THY1⁺ cells (n=5). Scale bar, 50 μm. (**B**) Quantitative RT-PCR analysis of uSPG and dSPG
markers expressed in the MACS-sorted THY1⁺ cells (n=3). Values, mean ± s.e.m.; p-values were
obtained using two-tailed t-tests (ns > 0.05, *p-value < 0.05, **p-value < 0.01, ***p-value < 0.001,
****p-value < 0.0001). (**C**) Feature plots showing the expression pattern of classic SPG markers
(stemness and differentiation). (**D**) Heatmap showing the expression pattern of markers for SPG in
different clusters. (**E**) Expression pattern dynamics of the SPG markers with pseudotime
progression. (**F**) Heatmap showing the expression pattern of markers for cell cycle phase in
different clusters. (**G**) The developmental trajectory of the overall SPG, colored by pseudotime.
579
580
581

Figure 1—figure supplement 2

Expression of top10 DEGs of Cluster1 in Figure 1B and classic SSC and SPG markers in
adult human germ cells. (**A**) Feature plots and violin plots of the Top10 DEGs of Cluster1. (**B**)
Immunostainings for LIN28A (red), DAPI (blue), and newly-found markers (green) in testicular
paraffin sections from adult mice. Scale bar, 50 μm. (**C**) Feature plots showing the expression
pattern of classic SSCs and SPG markers in adult human germ cells.

Figure 1—source data 1. Excel spreadsheet with the list of the top30 differentially expressed
genes of different clusters.

Figure 2 with 1 supplement

Lineage tracing and functional validation of FOXC2⁺ SSCs in *Foxc2^{iCreERT2/+};Rosa26^{LSL-T/G/LSL-
^{T/G}* mice.** (**A**) Schematic illustration of the lineage tracing workflow for FOXC2⁺ cells. (**B**)
Immunostainings for DAPI (blue) and FOXC2 (red) at day 3 post TAM induction. Scale bar, 50 μm;
d, day. (**C**) FACS analysis of GFP⁺ populations derived from *Rosa26^{LSL-T/G/LSL-T/G}* or

Foxc2$^{iCreERT2/+}$;Rosa26$^{LSL-T/G/LSL-T/G}$ mice at day 3 post TAM induction. (**D**, **E**) The recipient mice
testes (**D**) and colony numbers (**E**) 2 months after transplantation (n=10) of the FACS-sorted GFP$^+$
cells from the *Foxc2$^{iCreERT2/+}$;Rosa26$^{LSL-T/G/LSL-T/G}$* mice 3 days after TAM diet and the FACS-sorted
THY1$^+$ cells from adult mice. Scale bar, 1 mm; values, mean ± s.e.m.; p-values were obtained using
two-tailed t-tests (****p-value < 0.0001). (**F**) Immunostaining for DAPI (blue), ZBTB16/FOXC2 (red),
and GFP (green) at week 1 post TAM induction (scale bar, 50 µm). (**G**) Seminiferous tubules of
Foxc2$^{iCreERT2/+}$;Rosa26$^{LSL-T/G/LSL-T/G}$ mice 2, 4, and 6 weeks post TAM induction. Scale bar, 50 µm.
(**H**) Testes (scale bar, 1 mm), seminiferous tubules, and epididymis (scale bar, 50 µm) at month 4,
7, and 12 post TAM induction in *Foxc2$^{iCreERT2/+}$;Rosa26$^{LSL-T/G/LSL-T/G}$* mice. (**I**, **J**) The GFP$^+$ patches
(**I**) and progeny (**J**) population dynamics (n=10). Values, mean ± s.e.m. (**K**) Immunostainings for
DAPI (blue), EOMES (red), GFRA1 (red), or PAX7 (red) in GFP$^+$ population at week 2 post TAM
induction. Scale bar, 50 µm.
611
612

613   **Figure 2—figure supplement 1**

614
615   **Construction of the *Foxc2^{iCreERt2}* mice.**

Figure 3 with 1 supplement

Specific ablation of FOXC2⁺ SSCs and phenotypic validation in *Foxc2^iCreERT2/+*;*Rosa26^LSL-DTA/+*
mice. (**A**) Schematic illustration of the lineage tracing workflow for FOXC2⁺ cells. (**B-D**) Phenotypic
validation of the *Rosa26^LSL-DTA/+* and *Foxc2^iCreERT2/+*;*Rosa26^LSL-DTA/+* mice (n=5) for testes size (**B**),
testis weight and body weight (**C**), and HE-staining of the testes (**D**). Scale bars in (**B**), 1 mm; in
(**D**), 50 μm; d, day; values were mean ± s.e.m.; p-values were obtained using two-tailed t-tests
(ns > 0.05, *p-value < 0.05, **p-value < 0.01). (**E**) ZBTB16⁺, GFRA1⁺, LIN28A⁺, and FOXC2⁺ SPG
populations dynamics. Values, mean ± s.e.m. (n=10); p-values were obtained using one-way
ANOVA followed by Tukey test (ns > 0.05, *p-value < 0.05, **p-value < 0.01, ****p-value < 0.0001).
(**F**) Immunostainings for DAPI (blue), SYCP3 (green), and LIN28A (magenta) at day 14 post TAM
induction. d, day; scale bar, 50 μm. (**G**) Quantitative RT-PCR analysis of SPG markers expression
in the testes of the *Rosa26^LSL-DTA/+* and *Foxc2^iCreERT2/+*;*Rosa26^LSL-DTA/+* mice (n=3). Values, mean ±
s.e.m.; p-values were obtained using two-tailed t-tests (***p-value < 0.001, ****p-value < 0.0001).

632 **Figure 3—figure supplement 1**

633
634 **Depletion of uSPG pool in *Foxc2iCreERT2/+;Rosa26LSL-DTA/+* mice 14 days after specific ablation**
635 **of FOXC2+ SSCs.** Immunostainings for DAPI (blue), DDX4 (white), ZBTB16 (green), and FOXC2
636 (magenta) at day 3 and day 14 post TAM induction in *Foxc2iCreERT2/+;Rosa26LSL-DTA/+* mice (scale
637 bar, 50 μm).
638

Figure 4

640
FOXC2+ SSCs are critical for germline regeneration. (**A**) Co-immunostaining of FOXC2 (green)
with ZBTB16 (red) in seminiferous tubules of the adult testes at day 20 post busulfan treatment.
Scale bar, 50 μm. (**B**) ZBTB16+, GFRA1+, and FOXC2+ population dynamics after busulfan
treatment (20 mg/kg, n=10). (**C**) Co-immunostaining of FOXC2 (green) with MKI67 (red) in
seminiferous tubules of the adult testes at day 20 post busulfan treatment. Scale bar, 50 μm. (**D**)
MKI67+FOXC2+ proportions in relation to the whole FOXC2+ population at different time points after
busulfan treatment (n=4). (**E**) Schematic illustration for lineage tracing of FOXC2+ cell after busulfan
treatment. (**F**) Lineage tracing of the GFP+ cells at day 20 and month 2 after busulfan treatment
(scale bar, 50 μm). (**G**) The testes (scale bar, 1 mm), seminiferous tubules, and epididymis (scale
bar, 50 μm) at month 4, 7, and 12 post TAM induction and busulfan injection. m, month. (**H**, **I**) The
proportion dynamics of GFP patches (**H**) and GFP+ progenies (**I**). Values, mean ± s.e.m. (n=10).
w, week; m, month.

 Figure 5 with 1 supplement

654
655 **Spermatogenesis exhaustion in the adult *Foxc2^f/-;Ddx4^Cre/+* mice.** (**A**) Construction of the
656 *Foxc2^f/-;Ddx4^Cre/+* mice. (**B**) The testes size of the *Foxc2^f/-;Ddx4^Cre/+* mice. Scale bar, 1mm; M,
657 month. (**C**) Body weight and testis weight of the *Foxc2^f/-;Ddx4^Cre/+* mice at different age (n=5). M,
658 month; values, mean ± s.e.m.; p-values were obtained using two-tailed t-tests (ns > 0.05, ****p-
659 value < 0.0001). (**D**) HE-staining of the testis and epididymis. Scale bar, 50 μm; M, month. (**E**)

Estimation of degenerative tubules and sperm counts in cauda epididymis of the *Foxc2$^{f/+}$* and *Foxc2$^{f/-}$;Ddx4$^{Cre/+}$* mice with age (n=5). Values, mean ± s.e.m.; p-values were obtained using two-tailed t-tests (**p-value < 0.01, ****p-value < 0.0001). (**F**) Immunostainings for DAPI (blue), ZBTB16 (green), FOXC2 (magenta), and DDX4 (white) in the seminiferous tubules of the *Foxc2$^{f/+}$* and *Foxc2$^{f/-}$;Ddx4$^{Cre/+}$* mice. Scale bar, 50 μm. (**G**) Estimation of ZBTB16$^+$ uSPG number in the *Foxc2$^{f/+}$* and *Foxc2$^{f/-}$;Ddx4$^{Cre/+}$* mice with age (n=5). Values, mean ± s.e.m.; p-values were obtained using two-tailed t-tests (ns > 0.05, ***p-value < 0.001, ****p-value < 0.0001). (**H**) Quantitative RT-PCR analysis of the uSPG and germ cell markers expressed in the testis of the *Foxc2$^{f/+}$* and *Foxc2$^{f/-}$;Ddx4$^{Cre/+}$* mice (n=3). M, month; values, mean ± s.e.m.; p-values were obtained using two-tailed t-tests (**p-value < 0.01, ***p-value < 0.001, ****p-value < 0.0001).

672 **Figure 5—figure supplement 1**

A

B

673
Phenotypic validation of the *Foxc2^{f/-};Ddx4^{Cre/+}* mice. (**A**) Immunostainings for DAPI (blue),
ZBTB16 (green), LIN28A (magenta), and TRA98 (white) in seminiferous tubules of 1-week-old
Foxc2^{f/+} and *Foxc2^{f/-};Ddx4^{Cre/+}* mice. Scale bar, 50 μm. (**B**) Immunostainings for DAPI (blue),
LIN28A (green), and Cleaved-CASP3 (red) in seminiferous of the *Foxc2^{f/+}* and *Foxc2^{f/-};Ddx4^{Cre/+}*
mice (2-month-old). M, month; scale bar, 50 μm.

Figure 6 with 2 supplements

680
scRNA-seq analysis of THY1⁺ uSPG in *Foxc2^{f/+}* and *Foxc2^{f/-};Ddx4^{Cre/+}* mice. (**A**) Schematic
illustration of the scRNA-seq workflow. (**B**) t-SNE plot and developmental trajectory of uSPG from
Foxc2^{f/+} and *Foxc2^{f/-};Ddx4^{Cre/+}* mice respectively, colored by cluster. (**C**) Developmental trajectories
of uSPG from *Foxc2^{f/+}* and *Foxc2^{f/-};Ddx4^{Cre/+}* mice, colored by sample or derivation. (**D**)
Developmental trajectories of the cells in Cluster1 from *Foxc2^{f/+}* (CON) and *Foxc2^{f/-};Ddx4^{Cre/+}* (KO)
mice, colored by derivation or developmental state. (**E**) The Cluster1 cells proportion of each state
in CON and KO mice. (**F**) Heatmap showing the DEGs in the Cluster1 cells from the
Foxc2^{f/-};Ddx4^{Cre/+} mice compared with the *Foxc2^{f/+}* mice. (**G**) Top GO terms enrichment by the
down-regulated DEGs in KO mice. (**H**) Gene set enrichment analysis (GSEA) of the Cluster1 cells
(*Foxc2^{f/-};Ddx4^{Cre/+}* v.s. *Foxc2^{f/+}* mice). NOM, nominal; FDR, false discovery rate; ES, enrichment
score; NES, normalized enrichment score.

Figure 6—figure supplement 1

scRNA-seq analysis of THY1⁺ uSPG in adult *Foxc2^{f/+}* and *Foxc2^{f/-};Ddx4^{Cre/+}* mice. (**A**) Feature plots of classic SPG markers for uSPG in adult *Foxc2^{f/+}* or *Foxc2^{f/-};Ddx4^{Cre/+}* mice. (**B**) Expression dynamics of SPG markers with pseudotime progression for uSPG from *Foxc2^{f/+}* or *Foxc2^{f/-};Ddx4^{Cre/+}* mice respectively. (**C**) Expression dynamics of SPG markers with pseudotime progression for overall uSPG from *Foxc2^{f/+}* and *Foxc2^{f/-};Ddx4^{Cre/+}* mice.

699
700

701 **Figure 6—figure supplement 2**

702
Re-cluster and developmental trajectory analysis of cells in Cluster1 derived from adult
***Foxc2^{f/+}* and *Foxc2^{f/-};Ddx4^{Cre/+}* mice.** (**A**) The t-SNE plot of the Cluster1 cells aggregated from the
Foxc2^{f/+} and *Foxc2^{f/-};Ddx4^{Cre/+}* mice colored by sample or subcluster. (**B**) The cell proportion of each
sample in each subcluster. (**C**) Feature plots of SPG markers expression. (**D**) Developmental
trajectory of the aggregated Cluster1 cells colored by subcluster or pseudotime. (**E**) Expression
dynamics of SPG markers with pseudotime progression.
709
710 **Figure 6—source data 1**. Excel spreadsheet with the list of the differentially expressed genes
711 found by scRNA-seq and enriched Gene Ontology terms.
712
713

714 **Figure 7**

715
FOXC2 is essential for sustaining the quiescent state of SSCs via regulating cell cycle. (**A**)
Workflow schematic illustration of the CUT&Tag_FOXC2 analysis on the FACS-sorted FOXC2+
cells. (**B**) Pie chart for CUT&Tag_FOXC2 peaks genome distribution. (**C**) Profiling of
CUT&Tag_FOXC2 peaks in proximity to transcriptional starting site (TSS). The distance to TSS
within 1000 was highlighted in the purple box. (**D**) Top GO terms enrichment by genes annotated
by CUT&Tag_FOXC2 peaks. (**E**) Venn diagram of FOXC2 target genes defined by overlapping the
CUT&Tag sequencing and scRNA-seq datasets. (**F**) GO terms enrichment by the FOXC2 target
genes related to cell cycle regulation. (**G**) Chromatin landscapes of CUT&Tag_FOXC2 peaks of
the candidates associated with negative cell cycle regulation. (**H**) The DNA-binding motif for FOXC2
(predicted with HOMER). (**I**) The cell cycle-related candidates possessing high binding potential
(>0.8, predicted with JASPAR SCAN). (**J**) CUT&Tag-qPCR validation of the cell cycle arrest
regulatory genes. (n=3). Values, mean ± s.e.m.; p-values were obtained using two-tailed t-tests
(****p-value < 0.0001). (**K**) The model for the maintenance of the FOXC2+ SSC subpopulation in
adult testis.

730
731 **Figure 7—source data 1**. Excel spreadsheet with the list of the differentially expressed genes
732 found by CUT&Tag sequencing and Gene Ontology terms of the 306 crossed candidates.
733

References

Aloisio, G. M., Nakada, Y., Saatcioglu, H. D., Peña, C. G., Baker, M. D., Tarnawa, E. D., Mukherjee,
J., Manjunath, H., Bugde, A., Sengupta, A. L., Amatruda, J. F., Cuevas, I., Hamra, F. K., &
Castrillon, D. H. (2014). PAX7 expression defines germline stem cells in the adult testis. *J*
Clin Invest, *124*(9), 3929-3944. https://doi.org/10.1172/jci75943
Arai, F., & Suda, T. (2007). Maintenance of quiescent hematopoietic stem cells in the
osteoblastic niche. *Ann N Y Acad Sci*, *1106*, 41-53.
https://doi.org/10.1196/annals.1392.005
Bahar, R., J, O. W., Kawamura, K., Seimiya, M., Wang, Y., Hatano, M., Okada, S., Tokuhisa, T.,
Watanabe, T., & Tagawa, M. (2002). Growth retardation, polyploidy, and
multinucleation induced by Clast3, a novel cell cycle-regulated protein. *J Biol Chem*,
277(42), 40012-40019. https://doi.org/10.1074/jbc.M205345200
Bahuau, M., Houdayer, C., Tredano, M., Soupre, V., Couderc, R., & Vazquez, M. P. (2002). FOXC2
truncating mutation in distichiasis, lymphedema, and cleft palate. *Clin Genet*, *62*(6), 470-
473. https://doi.org/10.1034/j.1399-0004.2002.620608.x
Bakker, W. J., Blázquez-Domingo, M., Kolbus, A., Besooyen, J., Steinlein, P., Beug, H., Coffer, P. J.,
Löwenberg, B., von Lindern, M., & van Dijk, T. B. (2004). FoxO3a regulates erythroid
differentiation and induces BTG1, an activator of protein arginine methyl transferase 1. *J*
Cell Biol, *164*(2), 175-184. https://doi.org/10.1083/jcb.200307056
Buaas, F. W., Kirsh, A. L., Sharma, M., McLean, D. J., Morris, J. L., Griswold, M. D., de Rooij, D. G.,
& Braun, R. E. (2004). Plzf is required in adult male germ cells for stem cell self-renewal.
Nat Genet, *36*(6), 647-652. https://doi.org/10.1038/ng1366
Cheung, T. H., & Rando, T. A. (2013). Molecular regulation of stem cell quiescence. *Nat Rev Mol*
Cell Biol, *14*(6), 329-340. https://doi.org/10.1038/nrm3591
Clermont, Y. (1966a). Renewal of spermatogonia in man. *Am J Anat*, *118*(2), 509-524.
https://doi.org/10.1002/aja.1001180211
Clermont, Y. (1966b). Spermatogenesis in man. A study of the spermatogonial population. *Fertil*
Steril, *17*(6), 705-721.
Clermont, Y. (1969). Two classes of spermatogonial stem cells in the monkey (Cercopithecus
aethiops). *Am J Anat*, *126*(1), 57-71. https://doi.org/10.1002/aja.1001260106
Costoya, J. A., Hobbs, R. M., Barna, M., Cattoretti, G., Manova, K., Sukhwani, M., Orwig, K. E.,
Wolgemuth, D. J., & Pandolfi, P. P. (2004). Essential role of Plzf in maintenance of
spermatogonial stem cells. *Nat Genet*, *36*(6), 653-659. https://doi.org/10.1038/ng1367
Davis, K. E., Moldes, M., & Farmer, S. R. (2004). The forkhead transcription factor FoxC2 inhibits
white adipocyte differentiation. *J Biol Chem*, *279*(41), 42453-42461.
https://doi.org/10.1074/jbc.M402197200
Di Persio, S., Saracino, R., Fera, S., Muciaccia, B., Esposito, V., Boitani, C., Berloco, B. P., Nudo, F.,
Spadetta, G., Stefanini, M., de Rooij, D. G., & Vicini, E. (2017). Spermatogonial kinetics in
humans. *Development*, *144*(19), 3430-3439. https://doi.org/10.1242/dev.150284
Ehmcke, J., & Schlatt, S. (2006). A revised model for spermatogonial expansion in man: lessons
from non-human primates. *Reproduction*, *132*(5), 673-680.
https://doi.org/10.1530/rep.1.01081
Ehmcke, J., Wistuba, J., & Schlatt, S. (2006). Spermatogonial stem cells: questions, models and
perspectives. *Hum Reprod Update*, *12*(3), 275-282.
https://doi.org/10.1093/humupd/dmk001

Faisal, I., Cisneros-Montalvo, S., Hamer, G., Tuominen, M. M., Laurila, P. P., Tumiati, M.,
Jauhiainen, M., Kotaja, N., Toppari, J., Mäkelä, J. A., & Kauppi, L. (2019). Transcription
Factor USF1 Is Required for Maintenance of Germline Stem Cells in Male Mice.
Endocrinology, *160*(5), 1119-1136. https://doi.org/10.1210/en.2018-01088
Gallardo, T., Shirley, L., John, G. B., & Castrillon, D. H. (2007). Generation of a germ cell-specific
mouse transgenic Cre line, Vasa-Cre. *Genesis*, *45*(6), 413-417.
https://doi.org/10.1002/dvg.20310
Gaudet, P., Livstone, M. S., Lewis, S. E., & Thomas, P. D. (2011). Phylogenetic-based propagation
of functional annotations within the Gene Ontology consortium. *Brief Bioinform*, *12*(5),
449-462. https://doi.org/10.1093/bib/bbr042
Georgescu, M. M., Cote, G., Agarwal, N. K., & White, C. L., 3rd. (2014). NHERF1/EBP50 controls
morphogenesis of 3D colonic glands by stabilizing PTEN and ezrin-radixin-moesin
proteins at the apical membrane. *Neoplasia*, *16*(4), 365-374.e361-362.
https://doi.org/10.1016/j.neo.2014.04.004
Guo, J., Grow, E. J., Mlcochova, H., Maher, G. J., Lindskog, C., Nie, X., Guo, Y., Takei, Y., Yun, J.,
Cai, L., Kim, R., Carrell, D. T., Goriely, A., Hotaling, J. M., & Cairns, B. R. (2018). The adult
human testis transcriptional cell atlas. *Cell Res*, *28*(12), 1141-1157.
https://doi.org/10.1038/s41422-018-0099-2
Guo, R., Yu, Z., Guan, J., Ge, Y., Ma, J., Li, S., Wang, S., Xue, S., & Han, D. (2004). Stage-specific
and tissue-specific expression characteristics of differentially expressed genes during
mouse spermatogenesis. *Mol Reprod Dev*, *67*(3), 264-272.
https://doi.org/10.1002/mrd.20026
Hammoud, S. S., Low, D. H., Yi, C., Carrell, D. T., Guccione, E., & Cairns, B. R. (2014). Chromatin
and transcription transitions of mammalian adult germline stem cells and
spermatogenesis. *Cell Stem Cell*, *15*(2), 239-253.
https://doi.org/10.1016/j.stem.2014.04.006
Hara, K., Nakagawa, T., Enomoto, H., Suzuki, M., Yamamoto, M., Simons, B. D., & Yoshida, S.
(2014). Mouse spermatogenic stem cells continually interconvert between equipotent
singly isolated and syncytial states. *Cell Stem Cell*, *14*(5), 658-672.
https://doi.org/10.1016/j.stem.2014.01.019
Helsel, A. R., Yang, Q. E., Oatley, M. J., Lord, T., Sablitzky, F., & Oatley, J. M. (2017). ID4 levels
dictate the stem cell state in mouse spermatogonia. *Development*, *144*(4), 624-634.
https://doi.org/10.1242/dev.146928
Hermann, B. P., Cheng, K., Singh, A., Roa-De La Cruz, L., Mutoji, K. N., Chen, I. C., Gildersleeve,
H., Lehle, J. D., Mayo, M., Westernströer, B., Law, N. C., Oatley, M. J., Velte, E. K.,
Niedenberger, B. A., Fritze, D., Silber, S., Geyer, C. B., Oatley, J. M., & McCarrey, J. R.
(2018). The Mammalian Spermatogenesis Single-Cell Transcriptome, from
Spermatogonial Stem Cells to Spermatids. *Cell Rep*, *25*(6), 1650-1667.e1658.
https://doi.org/10.1016/j.celrep.2018.10.026
Jijiwa, M., Kawai, K., Fukihara, J., Nakamura, A., Hasegawa, M., Suzuki, C., Sato, T., Enomoto, A.,
Asai, N., Murakumo, Y., & Takahashi, M. (2008). GDNF-mediated signaling via RET
tyrosine 1062 is essential for maintenance of spermatogonial stem cells. *Genes Cells*,
13(4), 365-374. https://doi.org/10.1111/j.1365-2443.2008.01171.x
Kanatsu-Shinohara, M., Morimoto, H., & Shinohara, T. (2012). Enrichment of mouse
spermatogonial stem cells by melanoma cell adhesion molecule expression. *Biol Reprod*,
87(6), 139. https://doi.org/10.1095/biolreprod.112.103861

Kaya-Okur, H. S., Janssens, D. H., Henikoff, J. G., Ahmad, K., & Henikoff, S. (2020). Efficient low-
cost chromatin profiling with CUT&Tag. *Nat Protoc*, *15*(10), 3264-3283.
https://doi.org/10.1038/s41596-020-0373-x

Kaya-Okur, H. S., Wu, S. J., Codomo, C. A., Pledger, E. S., Bryson, T. D., Henikoff, J. G., Ahmad, K.,
& Henikoff, S. (2019). CUT&Tag for efficient epigenomic profiling of small samples and
single cells. *Nat Commun*, *10*(1), 1930. https://doi.org/10.1038/s41467-019-09982-5

Knudson, C. M., & Korsmeyer, S. J. (1997). Bcl-2 and Bax function independently to regulate cell
death. *Nat Genet*, *16*(4), 358-363. https://doi.org/10.1038/ng0897-358

Kowalczyk, M. S., Tirosh, I., Heckl, D., Rao, T. N., Dixit, A., Haas, B. J., Schneider, R. K., Wagers, A.
J., Ebert, B. L., & Regev, A. (2015). Single-cell RNA-seq reveals changes in cell cycle and
differentiation programs upon aging of hematopoietic stem cells. *Genome Res*, *25*(12),
1860-1872. https://doi.org/10.1101/gr.192237.115

Kubota, H., Avarbock, M. R., & Brinster, R. L. (2004). Culture conditions and single growth factors
affect fate determination of mouse spermatogonial stem cells. *Biol Reprod*, *71*(3), 722-
731. https://doi.org/10.1095/biolreprod.104.029207

Kume, T., Jiang, H., Topczewska, J. M., & Hogan, B. L. (2001). The murine winged helix
transcription factors, Foxc1 and Foxc2, are both required for cardiovascular
development and somitogenesis. *Genes Dev*, *15*(18), 2470-2482.
https://doi.org/10.1101/gad.907301

La, H. M., & Hobbs, R. M. (2019). Mechanisms regulating mammalian spermatogenesis and
fertility recovery following germ cell depletion. *Cell Mol Life Sci*, *76*(20), 4071-4102.
https://doi.org/10.1007/s00018-019-03201-6

La, H. M., Mäkelä, J. A., Chan, A. L., Rossello, F. J., Nefzger, C. M., Legrand, J. M. D., De Seram,
M., Polo, J. M., & Hobbs, R. M. (2018). Identification of dynamic undifferentiated cell
states within the male germline. *Nat Commun*, *9*(1), 2819.
https://doi.org/10.1038/s41467-018-04827-z

Li, H., Liang, Z., Yang, J., Wang, D., Wang, H., Zhu, M., Geng, B., & Xu, E. Y. (2019). DAZL is a
master translational regulator of murine spermatogenesis. *Natl Sci Rev*, *6*(3), 455-468.
https://doi.org/10.1093/nsr/nwy163

Liao, H. F., Chen, W. S., Chen, Y. H., Kao, T. H., Tseng, Y. T., Lee, C. Y., Chiu, Y. C., Lee, P. L., Lin, Q.
J., Ching, Y. H., Hata, K., Cheng, W. T., Tsai, M. H., Sasaki, H., Ho, H. N., Wu, S. C., Huang,
Y. H., Yen, P., & Lin, S. P. (2014). DNMT3L promotes quiescence in postnatal
spermatogonial progenitor cells. *Development*, *141*(12), 2402-2413.
https://doi.org/10.1242/dev.105130

Miura, N., Iida, K., Kakinuma, H., Yang, X. L., & Sugiyama, T. (1997). Isolation of the mouse (MFH-
1) and human (FKHL 14) mesenchyme fork head-1 genes reveals conservation of their
gene and protein structures. *Genomics*, *41*(3), 489-492.
https://doi.org/10.1006/geno.1997.4695

Motojima, M., Tanimoto, S., Ohtsuka, M., Matsusaka, T., Kume, T., & Abe, K. (2016).
Characterization of Kidney and Skeleton Phenotypes of Mice Double Heterozygous for
Foxc1 and Foxc2. *Cells Tissues Organs*, *201*(5), 380-389.
https://doi.org/10.1159/000445027

Nakagawa, T., Jörg, D. J., Watanabe, H., Mizuno, S., Han, S., Ikeda, T., Omatsu, Y., Nishimura, K.,
Fujita, M., Takahashi, S., Kondoh, G., Simons, B. D., Yoshida, S., & Nagasawa, T. (2021). A
multistate stem cell dynamics maintains homeostasis in mouse spermatogenesis. *Cell*
Rep, *37*(3), 109875. https://doi.org/10.1016/j.celrep.2021.109875

Nakagawa, T., Nabeshima, Y., & Yoshida, S. (2007). Functional identification of the actual and
potential stem cell compartments in mouse spermatogenesis. *Dev Cell*, *12*(2), 195-206.
https://doi.org/10.1016/j.devcel.2007.01.002
Nakagawa, T., Sharma, M., Nabeshima, Y., Braun, R. E., & Yoshida, S. (2010). Functional
hierarchy and reversibility within the murine spermatogenic stem cell compartment.
Science, *328*(5974), 62-67. https://doi.org/10.1126/science.1182868
Nitiss, J. L. (2009). DNA topoisomerase II and its growing repertoire of biological functions. *Nat*
Rev Cancer, *9*(5), 327-337. https://doi.org/10.1038/nrc2608
Oatley, J. M., Avarbock, M. R., & Brinster, R. L. (2007). Glial cell line-derived neurotrophic factor
regulation of genes essential for self-renewal of mouse spermatogonial stem cells is
dependent on Src family kinase signaling. *J Biol Chem*, *282*(35), 25842-25851.
https://doi.org/10.1074/jbc.M703474200
Qiu, X., Hill, A., Packer, J., Lin, D., Ma, Y. A., & Trapnell, C. (2017). Single-cell mRNA quantification
and differential analysis with Census. *Nat Methods*, *14*(3), 309-315.
https://doi.org/10.1038/nmeth.4150
Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H. A., & Trapnell, C. (2017). Reversed
graph embedding resolves complex single-cell trajectories. *Nat Methods*, *14*(10), 979-
982. https://doi.org/10.1038/nmeth.4402
Raman, M., Earnest, S., Zhang, K., Zhao, Y., & Cobb, M. H. (2007). TAO kinases mediate
activation of p38 in response to DNA damage. *Embo j*, *26*(8), 2005-2014.
https://doi.org/10.1038/sj.emboj.7601668
Relaix, F., & Zammit, P. S. (2012). Satellite cells are essential for skeletal muscle regeneration:
the cell on the edge returns centre stage. *Development*, *139*(16), 2845-2856.
https://doi.org/10.1242/dev.069088
Rodgers, J. T., King, K. Y., Brett, J. O., Cromie, M. J., Charville, G. W., Maguire, K. K., Brunson, C.,
Mastey, N., Liu, L., Tsai, C. R., Goodell, M. A., & Rando, T. A. (2014). mTORC1 controls
the adaptive transition of quiescent stem cells from G0 to G(Alert). *Nature*, *510*(7505),
393-396. https://doi.org/10.1038/nature13255
Roy, A., Lin, Y. N., Agno, J. E., DeMayo, F. J., & Matzuk, M. M. (2007). Absence of tektin 4 causes
asthenozoospermia and subfertility in male mice. *Faseb j*, *21*(4), 1013-1025.
https://doi.org/10.1096/fj.06-7035com
Sabine, A., Bovay, E., Demir, C. S., Kimura, W., Jaquet, M., Agalarov, Y., Zangger, N., Scallan, J. P.,
Graber, W., Gulpinar, E., Kwak, B. R., Mäkinen, T., Martinez-Corral, I., Ortega, S.,
Delorenzi, M., Kiefer, F., Davis, M. J., Djonov, V., Miura, N., & Petrova, T. V. (2015).
FOXC2 and fluid shear stress stabilize postnatal lymphatic vasculature. *J Clin Invest*,
125(10), 3861-3877. https://doi.org/10.1172/jci80454
Sada, A., Suzuki, A., Suzuki, H., & Saga, Y. (2009). The RNA-binding protein NANOS2 is required
to maintain murine spermatogonial stem cells. *Science*, *325*(5946), 1394-1398.
https://doi.org/10.1126/science.1172645
Sasman, A., Nassano-Miller, C., Shim, K. S., Koo, H. Y., Liu, T., Schultz, K. M., Millay, M., Nanano,
A., Kang, M., Suzuki, T., & Kume, T. (2012). Generation of conditional alleles for Foxc1
and Foxc2 in mice. *Genesis*, *50*(10), 766-774. https://doi.org/10.1002/dvg.22036
Sharma, M., Srivastava, A., Fairfield, H. E., Bergstrom, D., Flynn, W. F., & Braun, R. E. (2019).
Identification of EOMES-expressing spermatogonial stem cells and their regulation by
PLZF. *Elife*, *8*. https://doi.org/10.7554/eLife.43352

Sharma, S., Wistuba, J., Pock, T., Schlatt, S., & Neuhaus, N. (2019). Spermatogonial stem cells:
updates from specification to clinical relevance. *Hum Reprod Update*, *25*(3), 275-297.
https://doi.org/10.1093/humupd/dmz006
Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., 3rd, Hao, Y.,
Stoeckius, M., Smibert, P., & Satija, R. (2019). Comprehensive Integration of Single-Cell
Data. *Cell*, *177*(7), 1888-1902.e1821. https://doi.org/10.1016/j.cell.2019.05.031
Suzuki, S., McCarrey, J. R., & Hermann, B. P. (2021). An mTORC1-dependent switch orchestrates
the transition between mouse spermatogonial stem cells and clones of progenitor
spermatogonia. *Cell Rep*, *34*(7), 108752. https://doi.org/10.1016/j.celrep.2021.108752
Tan, K., & Wilkinson, M. F. (2019). Human Spermatogonial Stem Cells Scrutinized under the
Single-Cell Magnifying Glass. *Cell Stem Cell*, *24*(2), 201-203.
https://doi.org/10.1016/j.stem.2019.01.010
Tegelenbosch, R. A., & de Rooij, D. G. (1993). A quantitative study of spermatogonial
multiplication and stem cell renewal in the C3H/101 F1 hybrid mouse. *Mutat Res*,
290(2), 193-200. https://doi.org/10.1016/0027-5107(93)90159-d
Tokuda, M., Kadokawa, Y., Kurahashi, H., & Marunouchi, T. (2007). CDH1 is a specific marker for
undifferentiated spermatogonia in mouse testes. *Biol Reprod*, *76*(1), 130-141.
https://doi.org/10.1095/biolreprod.106.053181
Tokue, M., Ikami, K., Mizuno, S., Takagi, C., Miyagi, A., Takada, R., Noda, C., Kitadate, Y., Hara, K.,
Mizuguchi, H., Sato, T., Taketo, M. M., Sugiyama, F., Ogawa, T., Kobayashi, S., Ueno, N.,
Takahashi, S., Takada, S., & Yoshida, S. (2017). SHISA6 Confers Resistance to
Differentiation-Promoting Wnt/β-Catenin Signaling in Mouse Spermatogenic Stem Cells.
Stem Cell Reports, *8*(3), 561-575. https://doi.org/10.1016/j.stemcr.2017.01.006
Toyooka, Y., Tsunekawa, N., Takahashi, Y., Matsui, Y., Satoh, M., & Noce, T. (2000). Expression
and intracellular localization of mouse Vasa-homologue protein during germ cell
development. *Mech Dev*, *93*(1-2), 139-149. https://doi.org/10.1016/s0925-
4773(00)00283-5
Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N. J., Livak, K. J.,
Mikkelsen, T. S., & Rinn, J. L. (2014). The dynamics and regulators of cell fate decisions
are revealed by pseudotemporal ordering of single cells. *Nat Biotechnol*, *32*(4), 381-386.
https://doi.org/10.1038/nbt.2859
van Oosten, M., Stout, G. J., Backendorf, C., Rebel, H., de Wind, N., Darroudi, F., van Kranen, H.
J., de Gruijl, F. R., & Mullenders, L. H. (2005). Mismatch repair protein Msh2 contributes
to UVB-induced cell cycle arrest in epidermal and cultured mouse keratinocytes. *DNA*
Repair (Amst), *4*(1), 81-89. https://doi.org/10.1016/j.dnarep.2004.08.008
van Velthoven, C. T. J., & Rando, T. A. (2019). Stem Cell Quiescence: Dynamism, Restraint, and
Cellular Idling. *Cell Stem Cell*, *24*(2), 213-225.
https://doi.org/10.1016/j.stem.2019.01.001
Wang, D., Cai, C., Dong, X., Yu, Q. C., Zhang, X. O., Yang, L., & Zeng, Y. A. (2015). Identification of
multipotent mammary stem cells by protein C receptor expression. *Nature*, *517*(7532),
81-84. https://doi.org/10.1038/nature13851
Wang, M., Liu, X., Chang, G., Chen, Y., An, G., Yan, L., Gao, S., Xu, Y., Cui, Y., Dong, J., Chen, Y.,
Fan, X., Hu, Y., Song, K., Zhu, X., Gao, Y., Yao, Z., Bian, S., Hou, Y., . . . Qiao, J. (2018).
Single-Cell RNA Sequencing Analysis Reveals Sequential Cell Fate Transition during
Human Spermatogenesis. *Cell Stem Cell*, *23*(4), 599-614.e594.
https://doi.org/10.1016/j.stem.2018.08.007

Wang, X., Liu, C., Zhang, S., Yan, H., Zhang, L., Jiang, A., Liu, Y., Feng, Y., Li, D., Guo, Y., Hu, X., Lin,
Y., Bu, P., & Li, D. (2021). N(6)-methyladenosine modification of MALAT1 promotes
metastasis via reshaping nuclear speckles. *Dev Cell*, *56*(5), 702-715 e708.
https://doi.org/10.1016/j.devcel.2021.01.015
Weaver, B. A., Bonday, Z. Q., Putkey, F. R., Kops, G. J., Silk, A. D., & Cleveland, D. W. (2003).
Centromere-associated protein-E is essential for the mammalian mitotic checkpoint to
prevent aneuploidy due to single chromosome loss. *J Cell Biol*, *162*(4), 551-563.
https://doi.org/10.1083/jcb.200303167
Wei, C., Lin, H., & Cui, S. (2018). The Forkhead Transcription Factor FOXC2 Is Required for
Maintaining Murine Spermatogonial Stem Cells. *Stem Cells Dev*, *27*(9), 624-636.
https://doi.org/10.1089/scd.2017.0233
Yi, X., Hong, M., Gui, B., Chen, Z., Li, L., Xie, G., Liang, J., Wang, X., & Shang, Y. (2012). RNA
processing and modification protein, carbon catabolite repression 4 (Ccr4), arrests the
cell cycle through p21-dependent and p53-independent pathway. *J Biol Chem*, *287*(25),
21045-21057. https://doi.org/10.1074/jbc.M112.355321
Yuan, L., Liu, J. G., Zhao, J., Brundell, E., Daneholt, B., & Höög, C. (2000). The murine SCP3 gene is
required for synaptonemal complex assembly, chromosome synapsis, and male fertility.
Mol Cell, *5*(1), 73-83. https://doi.org/10.1016/s1097-2765(00)80404-9
Zheng, K., Wu, X., Kaestner, K. H., & Wang, P. J. (2009). The pluripotency factor LIN28 marks
undifferentiated spermatogonia in mouse. *BMC Dev Biol*, *9*, 38.
https://doi.org/10.1186/1471-213x-9-38