1 **Abundant pleiotropy across neuroimaging modalities identified through a multivariate**
2 **genome-wide association study**

3
4

5 E.P. Tissink[1*], A.A. Shadrin[2], D. van der Meer[2,3], N. Parker[2], G. Hindley[2,4], D. Roelfs[2], O. Frei[2], C.C.
6 Fan[5,6], M. Nagel[1], T. Nærland[7], M. Budisteanu[8,9], S. Djurovic[2,7,10], L. T. Westlye[2,7,11], M.P. van den
7 Heuvel[1,12], D. Posthuma[1,12], T. Kaufmann[2,13], A.M. Dale[6,14,15] & O.A. Andreassen[2,7*]

8
9

10 [*] Corresponding authors: e.p.tissink@vu.nl, ole.andreassen@medisin.uio.no

11
12

[1] Department of Complex Trait Genetics, Center for Neurogenomics and Cognitive Research, Vrije
Universiteit Amsterdam, Amsterdam Neuroscience, 1081 HV Amsterdam, The Netherlands
[2] NORMENT Centre, Division of Mental Health and Addiction, Oslo University Hospital and Institute
of Clinical Medicine, University of Oslo, Building 48, Oslo, Norway
[3] School of Mental Health and Neuroscience, Faculty of Health, Medicine and Life Sciences, Maastricht
University, The Netherlands
[4] Psychosis Studies, Institute of Psychiatry, Psychology and Neurosciences, King's College London, 16
De Crespigny Park, London SE5 8AB, United Kingdom
[5] Laureate Institute for Brain Research, Tulsa, Oklahoma, USA.
[6] Department of Radiology, University of California San Diego, La Jolla, CA 92037, USA
[7] K.G. Jebsen Centre for Neurodevelopmental disorders, Division of Paediatric Medicine, Institute of
Clinical Medicine, University of Oslo, Building 31, Oslo, Norway
[8] Prof. Dr. Alex Obregia Clinical Hospital of Psychiatry, Bucharest, Romania
[9] "Victor Babes" National Institute of Pathology, Bucharest, Romania
[10] Department of Medical Genetics, Oslo University Hospital, Oslo, Norway
[11] Department of Psychology, University of Oslo, Oslo, Norway
[12] Department of Child and Adolescent Psychology and Psychiatry, section Complex Trait Genetics,
Amsterdam Neuroscience, VU University Medical Centre, Amsterdam, The Netherlands
[13] Department of Psychiatry and Psychotherapy, Tübingen Center for Mental Health, University of
Tübingen, Germany
[14] Center for Multimodal Imaging and Genetics, University of California San Diego, La Jolla, CA
92037, USA
[15] Department of Neurosciences, University of California San Diego, La Jolla, CA 92037, USA

Abstract

Genetic pleiotropy is abundant across spatially distributed brain characteristics derived from one neuroimaging modality (e.g. structural, functional or diffusion MRI). A better understanding of pleiotropy across modalities could inform us on the integration of brain function, micro- and macrostructure. Here we show extensive genetic overlap across neuroimaging modalities at a locus and gene level in the UK Biobank (N=34,029) and ABCD Study (N=8,607). When jointly analysing phenotypes derived from structural, functional and diffusion MRI with the Multivariate Omnibus Statistical Test (MOSTest), we boost the discovery of loci and genes beyond previously identified effects for each modality individually. Cross-modality genes are involved in fundamental biological processes and predominantly expressed during prenatal brain development. We additionally boost genetic discovery and prediction of psychiatric disorders by conditioning independent GWAS on our multimodal multivariate GWAS. These findings shed light on the shared genetic mechanisms underlying variation in brain morphology, functional connectivity, and tissue composition.

Graphical abstract

Introduction

The brain is our most complex organ, rapidly integrating information from many different sources[1], with strong genetic influences[2]. Studying genetic influences through genome-wide association studies (GWAS) has shown that most loci and genes show association with multiple traits[3], a phenomenon known as "statistical pleiotropy" (e.g. one gene influences multiple phenotypes directly, or indirectly via a causal pathway or common factor)[4]. Recently, numerous loci and genes with pleiotropic effects across brain characteristics derived from a single neuroimaging modality (e.g. structural, functional or diffusion MRI) have been discovered[5–9]. Yet the majority of pleiotropic loci act across rather than within phenotype domains[3], indicating that the genes associated within these loci may show pleiotropic effects across neuroimaging modalities, but the extent of this is underexplored. Investigating how genes influence a wide variety of brain imaging phenotypes may shed light on the mechanisms underlying alterations in brain morphology, activity, connectivity, and tissue composition that often co-occur in heritable psychiatric disorders[10,11].

Complex traits are affected by thousands of genetic variants scattered throughout the genome which makes pleiotropy between them inevitable[12]. In presence of such abundant pleiotropy, multivariate GWAS models have greater statistical power than univariate GWAS models[13]. This is because the cumulative evidence from different phenotypes leads to more sensitive detection of genetic associations and reduces the burden of multiple testing[7]. Previous studies have investigated either structural MRI-derived (sub)cortical volumes, surface area, and thickness[5–7], functional MRI-derived brain connectivity[8], or diffusion MRI-derived brain tissue composition[9] in a multivariate GWAS framework. The identified loci and genes inform us about the biological signal that is picked up by MRI, such as linking genetic effects of diffusion MRI to synaptic pruning, neuroinflammation, and axonal growth[9], structural MRI to neurogenesis and cell differentiation[5], and functional MRI to mental health[14] and psychiatric disorders[8]. Yet the modality specificity of these links remains unclear. Moreover, many loci are currently not discoverable in unimodal analyses: a multimodal multivariate GWAS could identify those loci that can improve our understanding of the interplay of biological processes contributing to brain structural organisation on multiple scales and reveal underpinnings of the structure-function relationship from a genetics viewpoint.

A large body of neuroimaging research established associations between psychiatric disorders and brain structure[15–20], function[21,22] and diffusion metrics[23,24]. However, previous studies were not able to detect non-null genetic correlations[25–27], possibly due to mixed effect directions[28] and differential polygenicity between neuroimaging phenotypes and psychiatric disorders[29]. Alternative methods that leverage the shared genetic signal between brain morphology and psychiatric disorders have been more fruitful[28] allowing improved prediction of disease liability[30]. Given that alterations of brain morphology, activity, connectivity, and tissue composition often co-occur in heritable psychiatric disorders[10,11], using multimodal GWAS associations may potentially further aid in leveraging the shared genetic signal between neuroimaging traits and psychiatric conditions.

Here, we demonstrate evidence of extensive pleiotropy across neuroimaging modalities using

583 structural (sMRI), resting-state functional (fMRI) or diffusion (dMRI) MRI-derived phenotypes in

the UK Biobank and the Adolescent Brain Cognitive Development (ABCD) Study®. We do so by

performing unimodal and multimodal multivariate GWAS with the Multivariate Omnibus Statistical

Test (MOSTest), which was designed to boost statistical power by capitalizing on the distributed nature

of genetic influences across imaging-derived phenotypes[7]. Specifically, we 1) overlap the previously

described multivariate effects for each modality and 2) combine all MRI-derived phenotypes into one

multimodal multivariate GWAS, obtaining an additional boost for discovery of cross-modality

pleiotropic loci and genes. Next, we functionally annotate single-modality (identified in only one

unimodal analysis) and cross-modality (identified in ≥ 2 unimodal analyses or unique to multimodal

analysis) loci and genes to describe the biological signal across MRI modalities. Last, we improve locus

discovery of five major psychiatric disorders by using the multimodal, multivariate, genetic signal of

brain morphology, functional connectivity, and tissue composition and observe improved polygenic

prediction for depression, autism, ADHD, and bipolar disorder. Thereby the current study provides

insight into pleiotropic effects across neuroimaging modalities and their relevance for understanding

the neurobiology of the human brain and mental health conditions.

Results

Abundant pleiotropy of loci and genes across neuroimaging modalities

We used data from three previous studies that applied MOSTest on single neuroimaging modality

phenotypes: 172 sMRI-derived brain morphology measures (68 regional surface area and 68 regional

thickness of the cerebral cortex, and 36 volumes of subcortical structures)[7], 153 fMRI-derived BOLD

signal and connectivity measures (17 network variances and 136 network correlations)[8], and 276 dMRI-

derived brain tissue composition principal components (65 restricted isotropic diffusion PCs, 124

restricted directional diffusion PCs, 87 normalized free water diffusion PCs)[9]. We investigated white

British UK Biobank samples (as derived from both self-declared ethnic background and genetic

principal component analysis) with quality-controlled genotypes and neuroimaging available (N_{sMRI} =

34,029, N_{fMRI}=31,023, N_{dMRI} = 30,106). We also included the ABCD cohort which had identical sMRI-

and dMRI-derived measures and similar fMRI-derived measures (see Methods). The ABCD sample

has a heterogeneous and admixed ancestral background. We created a replication sample within ABCD

similar to our UKB discovery sample by selecting the subset of European individuals (N_{sMRI} = 4,794,

N_{fMRI} = 4,132, N_{dMRI} = 4,418) assigned based on genetic ancestry factor[31] as defined in Methods. All

quality controlled ABCD samples (N_{sMRI} = 8,607, N_{fMRI} = 7,277, N_{dMRI} = 7,853) served as an additional

(ad)mixed ancestral replication sample. Next to genetic ancestry, ABCD samples were used to test the

generalizability of our results across age.

Heritability estimates obtained in UK Biobank with linkage disequilibrium (LD) Score
Regression[32] ranged from median = 5.80% (IQR = 3.09%) for fMRI-derived network variances to
28.00% (IQR = 6.92%) for sMRI-derived subcortical volumes (Supplementary Fig. 1 and
Supplementary Data 1), with dMRI-derived measures ranking in-between (median = 13.70%, IQR =
9.63%). Genetic and phenotypic correlations were generally similar ($\rho(r_g, r_p) = 0.53$, $p = 2.2 \times 10^{-16}$), and
stronger within modalities than between modalities (Supplementary Fig. 2).

First, we combined all heritable (nominal $p < 0.05$) phenotypes derived from the same modality
in multivariate GWAS analyses using MOSTest (Manhattan plots in Supplementary Fig. 4). This
heritability filter was applied (as previously by Roelfs et al[8]) because including non-heritable
phenotypes into MOSTest analyses have been shown to reduce statistical power[7]. MOSTest estimates
the correlation between measures from univariate GWAS (on randomly permuted genotype data) and
sums the squared decorrelated z-values across univariate GWAS summary statistics (from the original
genotype data) to integrate the effects across the measures into a multivariate test statistic[7]. These
unimodal multivariate analyses identified 590, 42 and 512 genome-wide significant loci associated with
sMRI, fMRI and dMRI respectively ($p < 5 \times 10^{-8}/3$; Supplementary Data 2). These results are in line
with earlier research (Supplementary Results 1). The number of genome-wide significant lead SNPs
from UK Biobank that replicated at nominal significance ($p < 0.05$) in ABCD-based MOSTest summary
statistics differed across modalities (EUR: 24.46% sMRI, 8.70% fMRI, 23.63% dMRI), and were higher
for the larger sample with (ad)mixed ancestries (42.12% sMRI, 15.38% fMRI, 35.39% dMRI).
Replication rates after correcting for the number of lead SNPs tested are provided in Supplementary
Data 3. Applying MAGMA[33] gene-level analyses to unimodal multivariate summary statistics identified
1,620, 39 and 1,453 genome-wide significant genes ($p < ((0.05/18,877)/3 =) 8.83 \times 10^{-7}$) for sMRI, fMRI
and dMRI respectively (Supplementary Data 4).

When overlapping the identified loci ($p < 5 \times 10^{-8}/3$) and genes ($p < 8.83 \times 10^{-7}$) from each
modality (Methods), we observed 326 loci (36.18% of total) and 1,021 genes (41.79% of total)
associated with at least two out of three modalities (Fig. 1a). This indicates pleiotropy across
neuroimaging modalities both at the genome-wide significant locus and gene level. We replicated this
pattern of overlap between sMRI and dMRI genome-wide significant loci and genes from UK Biobank
in the ABCD cohort, though fMRI genome-wide significant loci did not overlap and fMRI genome-
wide significant genes were not identified in ABCD (Supplementary Fig. 5, Supplementary Data 5-6).
Replication patterns were similar across ancestries, with all of the sMRI and dMRI loci identified in the
European ABCD sample overlapping with the loci identified within the (ad)mixed ancestry ABCD
sample. Locus definition parameters (LD cut-off and window size) did not affect on the observed locus
overlap (Fig. 1a), as was apparent from sensitivity analyses presented in Supplementary Fig. 3.

Figure 1. **Abundant pleiotropy of loci and genes across neuroimaging modalities.** Overlap of genome-wide significant a) loci and b) genes observed across neuroimaging modalities in single-modality and joint multimodal analyses ($p < 5 \times 10^{-8}/3$). When sMRI, fMRI and dMRI-derived phenotypes are jointly analysed in MOSTest (multimodal analysis), a boost in discovery of pleiotropic loci and genes is observed (yellow). This pattern partially replicates in the ABCD cohort (Supplementary Fig. 5). For 10 example lead SNPs from different parts of a) the univariate *z*-scores of all phenotypes used in this study (sMRI-derived phenotypes in green, dMRI-derived phenotypes in red, fMRI-derived phenotypes in blue) are plotted in c) as the distance from the centre of each circular plot. The dashed black line corresponds to $p = 5 \times 10^{-8}$ ($z = 5.45$) and *z*-scores that pass this threshold are depicted larger. Positive effect direction is shown as a filled circle and negative effect direction as a white circle.

Multimodal GWAS boosts discovery of cross-modality loci and genes

We next investigated whether combining all sMRI, fMRI, and dMRI-derived measures in one multivariate analysis generated greater statistical power to identify significant pleiotropic loci and genes which show sub-threshold associations in each unimodal multivariate analysis (Manhattan plot in Supplementary Fig. 4). We therefore applied MOSTest across neuroimaging modalities, combining 583 phenotypes, identifying 794 genetic loci (Supplementary Data 2). The LD Score Regression intercept (1.04, SE=0.04) indicated that the inflation in these multimodal MOSTest p-values (lambdaGC = 2.83) was driven by polygenicity and not population stratification or cryptic relatedness. Replication rates at nominal significance were comparable to previous MOSTest studies: EUR 26.45%, 37.55% (ad)mixed ancestries (see Supplementary Data 3 for rates after correction for the number of lead SNPs). One-hundred-thirty-six (15.09%) of these loci did not overlap with any of the loci identified in the unimodal multivariate analyses, suggesting that MOSTest leveraged the shared genetic signal across imaging modalities to boost the discovery of pleiotropic loci. Gene-based GWAS from MAGMA showed that of the 2,242 genome-wide significant multimodal genes, 384 (15.72%) were not discovered for unimodal gene-based GWAS (Supplementary Data 4). We used the ABCD cohort to investigate the generalizability of our findings and observed a similar boost in multimodal discovery on a genome-wide significant locus and gene level (Supplementary Fig. 5, Supplementary Data 5-6). Additionally, 81.82% of multimodal loci identified in the European ABCD sample were overlapping with the loci identified in the (ad)mixed ancestry ABCD sample.

We examined the univariate associations underlying multivariate associations from different parts of the Venn diagram (Fig. 1a). Fig. 1c shows the univariate z-scores across sMRI-, fMRI-, and dMRI-derived phenotypes for ten lead SNPs representing different parts of the Venn diagram (Fig. 1a). Cluster map of univariate z-scores for all multimodal MOSTest lead SNPs is displayed in Supplementary Fig. 6. Alternatively, for every lead SNP identified in the MOSTest analyses presented in the Venn diagram (Fig. 1a) we extracted the minimum univariate p-value across all analysed phenotypes. A lead SNP with a relatively high minimum univariate p-value would indicate that the signal should have been highly distributed across other measures for the variant to become genome-wide significant in the multivariate analysis. Supplementary Fig. 7 shows that lead SNPs boosted by the multimodal analysis had relatively high minimum univariate p-values ($0.05 > p > 5 \times 10^{-5}$) more frequently (87%) than lead SNPs of loci identified in one unimodal analysis only (sMRI-modality 72%; fMRI-modality 25%; dMRI-modality 78%), indicating that the discovery of these lead SNPs is driven by pleiotropic signals across modalities.

Comparing characteristics of single- and cross-modality loci and genes

We investigated to what extent single-modality (identified in only one unimodal analysis) and cross-modality (identified in ≥ 2 unimodal analyses or boosted by the multimodal analysis) loci and genes (Supplementary Data 7-8) differ in their biological effects compared to other complex traits. To this

end, we annotated 251 sMRI-, 11 fMRI-, 177 dMRI-modality, and 462 cross-modality lead SNPs using

ANNOVAR[34] (Supplementary Data 9). We used the annotations of 43,492 (unique) lead SNPs derived

from 558 traits (with reasonable power, i.e. N > 50,000) across 24 trait domains from Watanabe et al.[3]

as a reference. Fig. 3b demonstrates that cross-modality and single-modality lead SNPs were very

similarly located in genomic regions as other complex traits[3,35]. Notably, cross-modality lead SNPs

showed unique enrichment of exonic variants (6.58%, OR = 2.17, 95% CI = 1.43-3.17, p = 2.91×10⁻⁴).

Watanabe et al. already demonstrated that the contribution of exonic SNPs increases from ~1% (SNPs

associated to 1 trait domain) to ~5% (associated to ≥10 trait domain) with increasing pleiotropy[3]. The

enrichment we observed therefore emphasizes the strong pleiotropic nature of the cross-modality lead

SNPs.

Next, we tested whether any difference in results from gene-set enrichment analyses with Gene

Ontology (GO) biological processes, cellular components and molecular functions could be observed

using the 599 sMRI-, 7 fMRI-, 432 dMRI-modality, or 1,405 cross-modality genes (Fig. 3a). We

identified four networks of gene-sets that consisted of both single-modality and cross-modality subsets

of genes and showed significant enrichment after Bonferroni correction (Supplementary Data 10).

These processes were mostly related to nervous system development and neuronal growth and

differentiation. However, most gene-sets were significantly enriched of cross-modality genes only and

highlight the genes' involvement in fundamental biological processes, such as cell cycle processes,

cellular structure (chromatin, cytoskeleton, cell junction), and vesicle transport.

To investigate potential differential temporal patterns of cortical gene expression between

single-modality and cross-modality genes, we investigated transcriptome data of post-mortem brain

tissue (N=56) representing males and females of multiple ethnicities across the life span (Fig. 3c)[36]. The

number of fMRI-modality genes in the data (n=5) was too low to generate a reliable expression pattern

(Supplementary Fig. 8). Prenatal gene expression was high for both sMRI-, dMRI-modality and cross-

modality genes compared to all genes in the data. In (early) childhood until adolescence, a differential

dMRI-modality expression pattern was not apparent (compared to all genes in the data), whereas sMRI-

modality and cross-modality genes were generally lower expressed. This matches a large body of

research showing that pre- and postnatal cortical transcriptomes differ largely and pronounced prenatal

expression matches the course of cortical development[37].

Given the predominant prenatal gene-expression pattern, we investigated whether single-

modality and cross-modality genes were enriched for any cell-type identified by Bhaduri et al.[38] using

single-cell RNA sequencing analyses of the fetal brain (Fig. 2d). We observed that cross-modality genes

were enriched across all cell-types, confirming their importance in fetal brain tissue and suggesting

their expression pattern is not cell-specific and more general in nature (Supplementary Data 11).

Figure 2. **Comparing properties of single-modality and cross-modality lead SNPs and genes.** a) Gene-ontology biological processes, molecular functions and cellular components that were (Bonferroni-corrected) significantly enriched for cross-modality, sMRI-modality and/or dMRI-modality genes (none of the GO terms tested showed enrichment for the 6 fMRI-modality genes). Node size reflects gene-set size, edges reflect pathway similarity scores (Methods). b) Functional consequences of single-modality and cross-modality lead SNPs as

annotated with ANNOVAR[34]. When the null hypothesis (OR=1) could be rejected after Bonferroni correction (p
$< 1.14 \times 10^{-3}$), the solid line indicates significant enrichment of the annotation. Annotations of 43,492 (unique)
lead SNPs derived from 558 traits across 24 trait domains from Watanabe et al. were used as reference for Fisher
Exact Test (Supplementary Data 9). ncRNA = non-coding RNA. c) Mean-normalized expression (y-axis) of
cross-modality and single-modality genes over developmental timepoints (x-axis; log10 scale). Gray shading
indicates 95% confidence intervals. The mean-normalized expression of fMRI-modality genes is displayed in
Supplementary Fig. 8, since the number of genes (n=5) was low and therefore created an unreliable pattern. d)
Cell-type enrichment analysis for fetal brain tissue from Bhaduri et al.[38]. Bonferroni corrected significant results
($p < (0.05 / 30 =) 1.67 \times 10^{-3}$) are indicated by an asterisk (*). ipc = intermediate progenitor cells, oligo =
oligodendrocytes.

Leveraging shared genetics with psychiatric disorders for discovery
Alterations in brain morphology, connectivity, and tissue composition often co-occur in heritable
psychiatric disorders[11], suggesting that our multimodal, multivariate genetic signal may have relevance
for genetic discovery and prediction of psychiatric disorders. It is possible to boost locus discovery and
polygenic prediction by re-ranking the test-statistics from a given GWAS based on a genetically related
secondary GWAS[39]. We therefore conditioned five major psychiatric disorder GWAS summary
statistics from the Psychiatric Genomics Consortium (Supplementary Data 12) on our multimodal
MOSTest summary statistics using the conditional false discovery rate approach (condFDR)[40] to
identify loci associated with schizophrenia (SCZ)[41], bipolar disorder (BD)[42], major depressive disorder
(MDD)[43,44], attention-deficit hyperactivity disorder (ADHD)[45], and autism spectrum disorder (ASD)[46].
The rationale behind condFDR is that, in the presence of cross-trait enrichment, a variant with strong
associations with both traits is more likely to represent a true association[30].
Compared to the number of genome-wide significant loci identified in the original GWAS (Fig. 3a), we
observed a 5-19 fold increase in locus yield in the condFDR summary statistics (Supplementary Data
13). Note that the locus yield from the original GWAS may be lower than the number presented in the
original articles, since we relied on publicly available data (e.g. excluding 23andMe) and exclude some
additional cohorts to prevent sample overlap (with e.g. UKB, see Supplementary Results 2). We
calculated the sign concordance of the condFDR lead SNPs using independent disorder GWAS
summary statistics[47–49] (Supplementary Data 12) and observed higher and more significant sign
concordance across all disorders compared to the lead SNPs identified in the original GWAS
(Supplementary Data 14). We positionally mapped condFDR loci to genes (528 genes for MDD, 2,416
for SCZ, 116 for ASD, 1,162 for BD, 122 for ADHD; Supplementary Data 15) and subsequently used
these genes as input for hypergeometric test-based tissue enrichment analyses, both as implemented in
FUMA. The SCZ and BD genes were significantly enriched in gene-sets with upregulated gene
expression in brain tissue, but not in other body tissues (Supplementary Data 16, Fig. 3c). This contrasts
with the sets of genes mapped from the original GWAS loci (60 genes for MDD, 498 for SCZ, 5 for

ASD, 186 for BD, 14 for ADHD; Supplementary Data 15), which showed no enrichment of upregulated gene expression in brain tissues. This illustrates that genetic overlap may be used to identify genetic variants that may play a role in psychiatric disorders with a specific relevance for brain traits.

Figure 3. **Leveraging multimodal analysis to boost discovery and polygenic prediction of psychiatric disorders**. a) Loci discovered in the publicly available summary statistics of each disorder GWAS ($p < 5 \times 10^{-8}$), and after conditioning the disorder on multimodal MOSTest summary statistics with conditional FDR (FDR < 0.05). b) Phenotypic variance explained for five psychiatric disorders explained by polygenic scores based on original disorder GWAS summary statistics (original PGS) and conditional disorder-multimodal GWAS summary statistics (pleioPGS). Note that PGS using top 10 and top 100 SNPs had too low R^2 values to be visible and are therefore not plotted, but can be found in Supplementary Data 17. Independent target samples were used for prediction (see Methods). A random sampling procedure and t-test were applied per disorder to test if the observed difference in R^2 was significant (Supplementary Fig. 9). c) Enrichment of disorder genes (positionally mapped from the original disorder GWAS loci *vs* conditional FDR loci) in tissue-specific gene-sets with upregulated expression (Supplementary Data 16). The red line indicates Bonferroni correction for the number of tests ($\alpha = 0.05/540 = 9.26 \times 10^{-5}$).

We followed-up by testing whether our disorder-multimodal condFDR summary statistics

could improve polygenic prediction of the disorders in independent samples using a pleiotropy-

informed polygenic scoring method[30] (pleioPGS; see Methods). For that purpose, we constructed

polygenic scores with PRSice2 in two independent clinical datasets (TOP and MoBa, see

Supplementary Data 12) based on the top 10-150,000 original disorder GWAS ordered SNPs and

condFDR-based ordered SNPs (Fig. 3b). We observed a 1.24-fold increase in variance explained ($R^2 =$

6.22%, $p = 2.28 \times 10^{-15}$) for BD in the TOP sample ($N_{case} = 463$; $N_{control} = 1,073$) using the pleioPGS with

100,000 SNPs, compared with the original GWAS-based PGS ($R^2 = 5.02\%$, $p = 2.28 \times 10^{-15}$). We

employed a random sampling procedure to show that this was a significant improvement ($t = 42.16$, p

$= 3.64 \times 10^{-101}$; Supplementary Fig. 9). Also for MDD ($N_{case} = 135$, $N_{control} = 1,073$), the pleioPGS ($R^2 =$

3.98%, $p = 2.16 \times 10^{-6}$) significantly outperformed ($t = 14.57$, $p = 7.43 \times 10^{-33}$) the original GWAS-based

PGS ($R^2 = 3.22\%$, $p = 1.92 \times 10^{-5}$). The phenotypic variance explained by PGS in ASD and ADHD was

low ($R^2 < 2.43\%$) when using either the original PGS or pleioPGS (Supplementary Data 17), though a

significant improvement of the pleioPGS over the original PGS could be observed for both ASD (ΔR^2

$= 0.99\%$, $t = 30.99$, $p = 4.16 \times 10^{-78}$) and ADHD ($\Delta R^2 = 0.28\%$, $t = 46.44$, $p = 9.68 \times 10^{-109}$). When

predicting SCZ in TOP ($N_{case} = 735$; $N_{control} = 1,073$), the original GWAS-based PGS explained

significantly ($p = 1.44 \times 10^{-34}$) more variance in SCZ liability.

Discussion

Our findings demonstrated that many loci and genes show pleiotropic effects across brain characteristics

derived from three distinct MRI modalities. We found evidence of extensive pleiotropy across

structural, functional and diffusion MRI from genetic overlap (loci and genes being associated with at

least two out of three modalities), and a boost in discovery of loci (n=136) and genes (n=384) when all

MRI-derived phenotypes are jointly analysed using MOSTest. The results in the ABCD cohort show

generalizability of structural and diffusion MRI pleiotropic loci from adulthood to childhood, and from

European ancestry to (ad)mixed ancestries. Genes with cross-modality pleiotropic effects were highly

prenatally expressed, though not cell-type specific and mostly enriched in gene-sets of general

biological functions. Moreover, we showed how these results can be leveraged to improve locus

discovery for and prediction of major psychiatric disorders.

The human brain is a highly complex and inter-connected structure for which "the whole is

more than the sum of its parts"[50]. This complexity emerges from tight interplay between different units

and processes, where disturbance of any part may change the state of the whole system. With this in

mind, pervasive effects of genetic variants on brain-related traits are inevitable, resulting in abundant

pleiotropy not only across phenotypically linked traits such as brain morphology measures[37], but also

in widespread genetic overlap between distinct aspects of brain functioning such as personality and

cognition[39]. Our results implicate that the multivariate genetic signals of structural, functional, or

diffusion MRI are not only composed of pleiotropic effects within modalities as previously shown[5–9],
but also of a component that is shared across brain traits measured with different MRI modalities. This
provides a new conceptual insight into the integration of human brain functional connectivity (fMRI),
microstructure (dMRI) and macrostructure (sMRI) and highlights the importance of characterizing
patterns of specificity and pleiotropy to improve our understanding of molecular neurobiological
mechanisms. It is expected from theoretical analysis that rare variants would follow this pattern of
pleiotropy under assumption of the infinitesimal model[51], which would be an interesting direction of
future research given the recent availability of multivariate tools for gene-based rare-variant association
studies[52,53].

We observed genetic overlap across all three modalities in our discovery sample from UK
Biobank. Even though the heritability and number of loci and genes for fMRI was lower than in dMRI
and sMRI, the proportions of genome-wide significant loci and genes that did and did not overlap with
other modalities was generally equal across modalities. The results in the ABCD cohort showed
generalizability of structural-diffusion MRI pleiotropy from old age to late childhood, and from
European ancestry to (ad)mixed ancestries. Note that more sophisticated methods for analyses with
admixed individuals are becoming available[54], that allow to model individuals' local ancestry. A lack
of power in the smaller ABCD cohort most likely limited the robust estimation of genetic associations
for the relatively low heritable functional MRI-derived phenotypes, which complicated examining the
generalizability of structural-functional-diffusion MRI pleiotropy beyond structural-diffusion MRI
pleiotropy. The necessary future samples size to uncover the heritability of neuroimaging-derived
phenotypes depends on genetic architecture and varies across modalities or the brain trait of interest[6].
For example, researchers expect that multivariate analyses of the prospective UK Biobank
neuroimaging sample of 100,000 individuals will explain 32% and 24% of the additive genetic variance
in cortical surface area and thickness respectively[5]. This highlights that both larger neuroimaging-
genetic datasets and novel statistical approaches are required to delineate the genetic background of
neuroimaging phenotypes.

Our enrichment analyses showed that the biological processes and molecular components only
enriched for cross-modality genes include more general functions such as cell cycle processes,
regulation of gene expression, cell junctions. That gene-sets implicated by cross-modality genes alone
are involved in fundamental biological processes is consistent with previous findings that genes
associated with multiple trait domains are more likely to be involved in general biological functions[3].
This is in line with our finding that, although cross-modality genes were highly expressed in fetal brain
tissue, those genes were not necessarily cell-specific – suggesting they may serve in cellular functions
taking place in all cell-types. Future research may explore this putative relationship between the degree
of pleiotropy and the specificity of biological functions. The functional enrichment of cross-modality
lead SNPs investigated here can hint towards the mechanisms through which pleiotropic effects could
be exerted. Cross-modality pleiotropic lead SNPs were enriched for exonic variants as found previously

for pleiotropic SNPs[3,12]. There is previous evidence to suggest that pleiotropy emerges from the

variants' effect on total expression of functional protein, for example by the selective exclusion of

missense exons from the gene transcript[55]. However, identification of the (unmeasured) causal SNPs

tagged by these multivariate GWAS lead SNPs is necessary in future studies to uncover the mechanisms

through which variants exerts their pleiotropic effect.

When conditioning GWAS summary statistics for five major psychiatric disorders on our

multimodal multivariate analysis, we discovered new loci including genes with increased brain-specific

gene expression. The validity of these boosted loci needs to be replicated in independent samples in

future research. Nevertheless, prioritizing these variants already showed enhanced polygenic prediction

for MDD, BD, ASD, and ADHD. This highlights that future diagnostic strategies targeting psychiatric

disorders may benefit from aligning the PGSs to a relevant endophenotype. This rationale is similar to

a previously published study that showed improved prediction of SCZ after filtering on genetic variants

expressed in the placenta[56]. Exploring which disorders benefit from either a more specific or broader

range of neuroimaging phenotypes could also be an intriguing area for future research.

Some limitations are worth noting when interpreting our results. First, despite our efforts to

harmonize the three sets of phenotypes to the greatest degree, the differential spatial granularity and

number of features across modalities can result in differential representation of certain brain regions or

brain characteristics in the multivariate signal. Second, our definition of single-modality loci and genes

is inherent to two factors: 1) the grouping of phenotypes in unimodal and multimodal analyses –

MOSTest may miss a small number of hits when genetic signal is sparse across the jointly analyzed

phenotypes[7], and 2) the currently available data limits our statistical power – loci that are now associated

with one modality could become genome-wide significant in another modality once sample sizes

increase. Third, one should keep in mind that the presence of statistical pleiotropy as indicated in this

study can include instances of pleiotropy where multiple traits are affected by one gene but different

causal SNPs[4] or one locus with distinct gene effects that are in linkage disequilibrium[3]. Fourth,

MOSTest does not provide effect directions due to its multivariate nature and requires the use of

individual level data. This restricted access to certain post GWAS analyses that require direction of

effect, but these limitations were outweighed by MOSTest's ability to boost identification of variants

with shared effects across phenotypes and handle hundreds of phenotypes with sample size differences

in a computationally efficient manner[7]. Fifth, the temporal gene-expression patterns of single-modality

and cross-modality genes are based on a *postmortem* dataset with a limited number of donors per

timepoint[36]. A study that was recently released compared gene-expression between *postmortem* and

living brain samples and found a significant difference in 80% of the genes[57]. Although living postnatal

gene-expression samples may become available in the future for validation of our findings, prenatal

gene-expression will remain to rely on *postmortem* brain tissue.

In conclusion, we identified extensive cross-modality pleiotropy and demonstrated that

combining different neuroimaging modalities in multivariate analysis substantially increases genetic

variant and gene discovery compared to multivariate analyses within single modalities. The results

presented improve our understanding of the biology implicated by single-modality and cross-modality

genetic effects, and provide insights into the mechanistic pathways linking common genetic variation,

brain structure and function, and psychiatric disorders.

Methods

Samples

UK Biobank

The primary analyses of this study were conducted using data from UK Biobank participants who

provided written informed consent. This population-based resource obtained ethical approval from

the National Research Ethics Service Committee North West–Haydock (reference 11/NW/0382) and

the current study was conducted under application number 27412. We included participants that passed

quality control for functional[8] or diffusion[9] MRI-derived phenotypes as described in previous

publications (N_{fMRI} = 39,951; N_{dMRI} = 31,306). We increased our sample size for participants with

structural MRI-derived phenotypes compared to the original MOSTest publication[7], since new data had

been released (N_{sMRI} = 42,068). For all three subsamples, we excluded participants based on relatedness

(kinship coefficient > 0.05 as estimated in PLINK), non-European ancestry (UKB field 22006), a

genotype missing rate > 10%, and bad scan quality as indicated by an (age- and sex-adjusted) Euler

number > 3 SDs lower than the scanner site mean. This resulted in the sample characteristics described

in Supplementary Data 18.

Adolescent Brain Cognitive Development (ABCD) Study

Baseline data from ABCD participants from release 3.0 [NIMH Data Archive (NDA) DOI:10.151.54/

1519007] were used for the replication efforts in this study. All children in this cohort assented before

participation and their parents or guardians provided written informed consent. The procedures were

approved by a central Institutional Review Board (IRB) at the University of California, San Diego, and,

in some cases, by individual site Institutional Review Boards. We included participants with data for

the structural or functional MRI-derived phenotypes of interest (N_{sMRI} = 11,760; N_{fMRI} = 11,801) or

quality controlled diffusion MRI-derived phenotypes as previously described[9] (N_{dMRI} = 11,904). For all

three subsamples, we excluded participants based on recommended criteria for either modality as

provided by ABCD (e.g. imgincl_t1w_include), relatedness (first cousin), a genotype missing rate >

10%, and bad scan quality as indicated by an (age- and sex-adjusted) Euler number < 3 SDs lower than

the scanner site mean. This resulted in a sample with (ad)mixed ancestries (Table 1), as described by

Fan et al. recently[58]. We additionally excluded participants with a genetic ancestry factor[31] of European

ancestry < 90% (as provided by ABCD and applied previously in Loughnan *et al*[59]) to create a

replication sample that matched the ancestry characteristics of the discovery sample more specifically

(Supplementary Data 18).

Genotype data

UK Biobank

UK Biobank samples were genotyped from whole blood either using the UK BiLEVE or the UK Biobank axiom array and subsequently quality controlled and imputed by the UK Biobank Team[60]. Additional quality control was performed in-house and included SNP filters on minor allele frequency (MAF > 0.1%), imputation information score (INFO > 0.5), Hardy-Weinberg equilibrium (HWE; $p < 1 \times 10^{-9}$) and missingness (< 10%). This resulted in 9,061,587 SNPs used for association testing. Ancestral principal components were computed within European samples by UK Biobank and used to control for population stratification.

ABCD

Release 3.0 genotype data from ABCD participants was obtained through the Affymetric NIDA SmokeScreen Array, using either saliva or whole blood based on higher successful calls, higher non-missingness, matched genetic sex and less excessive identity by state. Initial quality control was performed by ABCD based on calling signals and variant call rates and subsequently following pre-imputation RICOPILI (Rapid Imputation and Computational Pipeline). We complemented ABCD quality control after creating two subsamples (European and (ad)mixed ancestries as described above) by further filtering pre-imputed variants on call rates (<5% missingness), MAF > 0.01, passing the HWE test ($p < 1 \times 10^{-9}$) and heterozygosity rate (deviating >6SD from the mean value) in PLINK2[61]. A pruned set of SNPs (r^2=0.1) was used to estimate 20 genetic principal components within each subsample to use downstream as covariates in multivariate GWAS. Genetic data was phased and imputed using the TOPMed imputation server and only SNPs with high imputation quality were retained (INFO > 0.9).

Neuroimaging data

Structural MRI-derived phenotypes

Three previous publications have used MOSTest on sMRI-derived phenotypes. These included either region-of-interest (ROI)-based cortical thickness, surface area, and subcortical volume[7] or vertex-based cortical thickness, surface area[5], and sulcal depth[6]. Given that the aim of this study was to combine phenotypes derived from three modalities and MOSTest can currently analyse a few thousand of phenotypes simultaneously, we opted to use the ROI-based cortical thickness, surface area, and subcortical volume[7] phenotypes given their relative low dimensionality. Supplementary Data 1 contains all the regional morphology measures included in the current study and indicates which measures were analysed for the left and right hemisphere separately. The respective publication by Van Der Meer & Frei *et al*[7] describes how the sets of 36 regional subcortical volumes, 68 cortical thickness and 68 surface area, as well as estimated intracranial volume (for covariate use downstream), were extracted from T1-weighted MRI using FreeSurfer v5.3[62,63] in UK Biobank samples. A similar procedure was performed

by the ABCD Data Acquisition and Integration Core and were readily available. As the importance of

normally distributed phenotypes for MOSTest was demonstrated in the original publication[7], we applied

rank-based inverse-normal transformation to each measure.

Functional MRI-derived phenotypes

We used functional MRI (fMRI)-derived phenotypes as previously described by Roelfs *et al*[8]. The UK

Biobank resting-state fMRI scans were processed into 1,000 Schaefer parcels[64] and mapped onto 17

large-scale brain networks defined by Yeo & Krienen *et al*[65]. The averaged time series within each Yeo-

defined network were Pearson correlated and represented 136 brain connectivity measures next to the

17 network variances (Supplementary Data 1). Rank-based inverse-normal transformation was applied

to each measure.

The ABCD Data Acquisition and Integration Core provided similar, but not identical, resting-

state fMRI-derived phenotypes for replication purposes. Instead of the 17 Yeo & Krienen networks

based on 1,000 parcels, temporal variance in 333 Gordon-defined parcels and 66 average correlations

between 12 Gordon-defined networks were available[66]. We averaged the parcel variances belonging to

the same network to achieve comparability to our discovery phenotypes. Subsequently, we rank-based

inverse-normal transformed the 12 network variances and 66 network connectivity phenotypes.

Diffusion MRI-derived phenotypes

We used diffusion MRI-derived phenotypes from UK Biobank and ABCD based on a voxel-wise

restriction spectrum imaging (RSI) model (as in Fan *et al*[9]). In short, RSI estimates the signal volume

fractions of separable pools of water in the human brain (i.e. intracellular, extracellular, and unhindered

free water) and their corresponding spherical harmonic coefficients[67,68]. Three RSI features were used:

1) restricted isotropic diffusion (N0) is most sensitive to isotropically diffusing water in the restricted

compartment (within cell bodies), restricted directional diffusion (ND) is sensitive to anisotropically

diffusing water in the restricted compartment (within oriented structures such as axons and dendrites),

and 3) normalized free water diffusion (NF) is sensitive to cerebrospinal fluid or intravascular spaces[69].

Fan *et al*[9] calculated the principal components (PCs) across all voxels and extracted the first 5,000 PCs

explaining more than 70% of the total variance of each feature. Here, due to dimensionality constraints,

we reduced the number of PCs for each feature by estimating the "elbow" of each scree plot of

eigenvalues using the nScree function of the nFactors R package (Supplementary Fig. 10). This resulted

in the first 124 ND-PCs, 87 NF-PCs and 65 N0-PCs used in our multivariate GWAS.

Statistical analyses

SNP-based GWAS

We performed discovery and replication SNP-based GWAS in PLINK2[61] for every MRI-derived phenotype separately while controlling for sex, age, age^2, genotype array (UKB only), scanner, 20 genetic principal components, and modality specific covariates. The latter included Euler number, and total surface area, mean thickness or intracranial volume (sMRI), signal to noise ratio and motion (fMRI), and intracranial volume (dMRI). A linear regression model with additive allelic effects was fitted for each SNP. Subsequently, SNP-based heritability (h_{SNP}^2) was estimated for each phenotype using Linkage Disequilibrium Score Regression (LDSC)[32]. Univariate GWAS summary statistics from non-heritable phenotypes (nominal significance threshold $\frac{h_{SNP}^2}{h_{SNP_{SE}}^2} > 1.96$ as used by Roelfs *et al*[8]) were dropped from further multivariate analyses, since including them may reduce statistical power[7]. This led to the exclusion of 3 dMRI-derived NF-PCs and 14 fMRI-derived connectivity phenotypes GWAS summary statistics (Supplementary Data 1). Then, variant *z*-scores from univariate GWAS were combined in the MOSTest framework to construct multivariate *p*-values as described by Shadrin *et al*[5]. This approach selects a regularization parameter optimized to the maximum yield of genetic loci (Supplementary Data 19). The alpha level for SNPs reaching genome-wide significance in the multivariate GWAS was $\alpha = 5 \times 10^{-8}$.

Locus definition

We defined genome-wide significant loci from MOSTest and conditional FDR summary statistics following a protocol as implemented in FUMA. First, independent genome-wide significant SNPs ($p < 5 \times 10^{-8}/3$) were obtained by clumping ($r^2 < 0.6$) and SNPs in linkage disequilibrium (LD) with them ($r^2 \geq 0.6$) were defined as candidate SNPs. LD was estimated using reference genotypes, using 5,000 random participants from the UK Biobank sample for UKB-based summary statistics and 1000 Genomes Phase 3 EUR for the European as well as (ad)mixed ancestry ABCD-based summary statistics. Second, independent significant SNPs with $r^2 < 0.1$ were defined as lead SNPs and the minimum and maximum positional coordinates of the corresponding candidate variants defined the locus start and end position. Loci in <250kb proximity were merged into a single locus. We excluded loci with a single SNP as these are more likely to be false positives. To test the robustness of our findings we performed sensitivity analyses for the locus definition parameters LD cut-off and merging proximity window, leading to twelve scenarios with different locus definition settings (Supplementary Fig. 3).

Multivariate gene-based GWAS

We explored the overlap between modalities and multimodal MOSTest on a gene-level by applying a SNP-wise mean model for 18,877 genes with MAGMA (Multi-marker Analysis of GenoMic Annotation) v1.08[33] in FUMA. The SNP-based MOSTest summary statistics from sMRI, dMRI, fMRI

and multimodal served as input with default settings and the UKB European population was used as

reference. The alpha level for genes reaching genome-wide significance was adjusted from $\alpha = 0.05$ to

$\alpha = (0.05/18,877)/3 = 8.83 \times 10^{-7}$ according to Bonferroni correction for multiple testing.

Definition of single-modality and cross-modality loci and genes

Locus overlap between the three MOSTest summary statistics (sMRI, fMRI, dMRI) was defined as

physically overlapping genome-wide significant loci after clumping (see above). We used the

GenomicRanges R-package[70] to compare the chromosome and start and end base pair positions of all

loci between any pair of summary statistics. A locus was considered single-modality when it did not

overlap with any of the loci identified for other modalities. All loci that were found to overlap between

two or more modalities also overlapped with the multimodal loci, hence we decided to represent these

cross-modality loci with the association statistics of the multimodal locus' lead SNP(s) in downstream

analyses. All univariate statistics that laid the basis of these single-modality and cross-modality lead

SNPs were plotted in a heatmap with the hierarchical clustering algorithm as implemented in

seaborn.clustermap (https://seaborn.pydata.org/generated/seaborn.clustermap.html) applied to the

phenotypes using (Pearson's) correlation as a distance metric and method='average' clustering. The

sMRI, fMRI, dMRI and multimodal MOSTest genes that were found to be genome-wide significant in

MAGMA were compared to provide a similar overview. The overlapping patterns were then plotted

with the eulerr R-package. This procedure was repeated for both discovery (UKB) and replication

(ABCD) MOSTest summary statistics, to investigate whether a similar overlapping pattern could be

observed.

Comparison of single-modality and cross-modality lead SNPs

We were interested in potential differences between single-modality and cross-modality loci compared

to other complex traits. Therefore we selected the lead SNPs within the respective loci (see *Locus*

definition) and annotated them with ANNOVAR[34]. As reference, we used the annotations of 43,492

(unique) lead SNPs derived from 558 traits (with reasonable power, i.e. N > 50,000) across 24 trait

domains (Table 2 from Watanabe et al.[3]). Enrichment of the single-modality and cross-modality lead

SNPs in positional annotation categories was then tested using Fisher's Exact test. The alpha level for

significant enrichment was Bonferroni corrected ($\alpha = 0.05 / 44 = 1.14 \times 10^{-3}$).

Comparison of single-modality and cross-modality gene properties

In order to interpret the biological processes, cellular components or molecular functions our single-

modality and cross-modality genes are involved in, Gene Ontology (GO)[71,72] gene-sets were tested for

enrichment of genes in these four lists using hypergeometric testing as implemented in FUMA[73]. Protein

coding genes were used as background genes and Bonferroni correction was applied to adjust for the

multiple comparisons. The significant single-modality and cross-modality GO terms were visualized as

a graph using Cytoscape, EnrichmentMap[74] and AutoAnnotate[75] following the Nature Protocol

by Reimand & Isserlin *et al*[76]. Stringent pathway similarity scores (Jaccard and overlap combined

coefficient = 0.6 as used in Paczkowska, Barenboim *et al*[77]) were used as edges.

To visualize the temporal gene expression pattern of the sets of single-modality and cross-

modality genes, we made use of gene expression data derived from brain tissue from 56 donors[36]. This

dataset ranges from 5 weeks post conception to 82 years of age and we used the data as pre-processed

in Kang *et al*[36]. We selected the probe with the highest differential stability for each gene (n = 16,849).

A number of single-modality ($n_{sMRI-modality}$ = 111, $n_{dMRI-modality}$ = 83, $n_{fMRI-modality}$ = 2) and cross-modality

genes (n=300) were not available in the data. Given the relatively high homogeneity of expression

patterns across cortical brain samples[78], we subsequently averaged over 13 cortical regions, within

donor, and normalized the expression values, within probe, across donors, to a range between 0 and

100. Mean expression over time per set of genes was plotted with ggplot2 in R v4.0.3., with

geom_smooth(method="gam") using default settings.

Lastly, we explored cell-type enrichment of single-modality and cross-modality genes by

performing Fisher Exact Tests for each cell-type identified by Bhaduri et al.[38] using single-cell RNA

sequencing analyses of the fetal brain. The minimum number of genes overlapping was set to >2.

Bonferroni correction was applied for the number of tests performed (α = 0.05 / 30 = 1.67×10^{-3}).

Conditioning major psychiatric disorder GWAS on multimodal MOSTest

We explored whether the multimodal multivariate summary statistics could be leveraged to improve

locus discovery for major psychiatric disorders (schizophrenia[41], bipolar disorder[42], major depressive

disorder[43], attention-deficit hyperactivity disorder[45], and autism spectrum disorder[46]). For that purpose,

we applied Conditional False Discovery Rate (cFDR)[40] on Psychiatric Genomics Consortium GWAS

summary statistics (listed in Supplementary Data 12) by conditioning on multimodal MOSTest

summary statistics. We obtained disorder summary statistics excluding UK Biobank to prevent sample

overlap. In cFDR analyses, original *p*-values are replaced by FDR values that reflect the posterior

probability that a SNP is null for the disorder given that the *p*-values for both phenotypes are as small

or smaller as the observed *p*-values:

(1) $$FDR(p_{disorder}|p_{multimodal}) = \frac{\pi_0(p_{multimodal})p_{disorder}}{F(p_{disorder}|p_{multimodal})}$$

with F = the conditional empirical cumulative distribution function and $\pi_0(p_{multimodal})$ = the

conditional proportion of null SNPs for the disorder given that *p*-values for the multimodal phenotype

are as small or smaller. We subsequently defined genome-wide significant loci (see *Locus definition*)

for the original ($p < 5 \times 10^{-8}$) and conditioned (FDR < 0.05) summary statistics and compared the results.

Since the probability of replicating these original and conditioned loci at genome-wide

significance is low due to the limited statistical power of replication GWAS, we first tested for en masse

sign concordance of effect direction in independent summary statistics, as is in line with previous

literature[79,80]. All summary statistics that were used to look up lead SNPs and test for sign concordance
are listed in Supplementary Data 12. An exact binomial test was used to test the null hypothesis that
directions of effect were randomly distributed (*prob.*=0.5), given the total number of variants and the
number of variants with concordant effects.
627
Tissue enrichment of condFDR genes and original disorder genes
Using the genome-wide significant loci from the condFDR and original summary statistics, we used
FUMA[73] (https://fuma. ctglab.nl/) to positionally map SNPs to genes. The resulting gene lists were used
for hypergeometric test-based tissue enrichment analyses (as implemented in FUMA), to test the tissue-
specificity of the input genes for differentially expressed (upregulated) genes in each of the 54 tissues
available in GTEx (v8)[81]. Bonferroni correction was applied for the number of tests performed ($\alpha =$
0.05 / (54 tissues × 2 gene lists × 5 disorders) = 9.26×10^{-5}).
635
PleioPGS
We then computed the prediction power of the original and conditioned summary statistics by
constructing polygenic scores (PGS) in independent case-control samples for the five major psychiatric
disorders described above in independent samples. The TOP, BUPGEN and MoBa[82] samples are
described in the Supplementary Methods (and Supplementary Data 12). We applied two different set-
ups, both based on the C+T (clumping + thresholding) approach[83] using different strategies for ranking
SNPs: 1) original GWAS *p*-value-based ranking and original GWAS effect sizes (standard PGS); 2)
cFDR-based ranking (described above) and original GWAS effect sizes (pleioPGS as introduced by
Van der Meer *et al*[30]; https://github.com/precimed/pleiofdr). For these two setups PGS were calculated
across five sets of LD-independent SNPs (n_{SNP} = 1,000, 10,000, 50,000, 100,000, 150,000) using
PRSice-2 (v2.3.3)[84] with no additional clumping (--no-clump option). Sets of LD-independent SNPs
were obtained using plink v1.90b6.1[61] based on the setup-defined SNP ranking with --clump-kb 250, -
-clump-r2 0.1 parameters and in-sample LD estimates. In both setups the phenotypic variance explained
by the PGS (R^2) was estimated as the difference between the R^2 of the full regression model and the R^2
of the null model only including the covariates (age, sex and the first 10 genetic PCs), also known as
Nagelkerke's pseudo-R^2. Bonferroni correction was applied for the number of tests ($p < (0.05 / (7\ n_{SNP}$
cut-offs × 2 PRS models × 5 disorders) =) 7.14×10^{-4}.
We applied random sampling without replacement procedure and *t*-test per disorder to test if
the highest R^2 (MDD pleioPGS/SCZ original PGS with n_{SNP} = 50,000 SNPs; ADHD/ASD/BD pleioPGS
with n_{SNP} = 100,000 SNPs) significantly higher than the R^2 of the alternative PGS (MDD original
PGS/SCZ pleioPGS with n_{SNP} = 50,000 SNPs; ADHD/ASD/BD original PGS with n_{SNP} = 100,000
SNPs). For each trait and type of PGS (original PGS, pleioPGS), we performed 200 iterations randomly
sampling 2/3 of controls and 2/3 of corresponding disorder cases with replacement and applying
PRSice2 to estimate R^2. Obtained distributions of R^2 values were used to assess confidence intervals.

Data availability

Genome-wide summary statistics will be made publicly available via https://ctg.cncr.nl/software/summary_statistics/ and GWAS Catalog upon acceptance. The individual-level data that support the discovery findings of this study are available from UK Biobank but restrictions apply to the availability of these data, which were used under license no. 27412 for the current study. Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development^{SM} (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive (NDA). ABCD data used for replication in this study is registered under the NDA study register at https://doi.org/10.15154/1527969. Data from the Norwegian Mother, Father and Child Cohort Study and the Medical Birth Registry of Norway used in this study are managed by the national health register holders in Norway (Norwegian Institute of public health) and can be made available to researchers, provided approval from the Regional Committees for Medical and Health Research Ethics (REC), compliance with the EU General Data Protection Regulation (GDPR) and approval from the data owners.

Code availability

Code to obtain the multimodal results presented in this manuscript are available via https://github.com/EPTissink/MOSTest-multimodal. Code from previously published studies based on different modalities from which we used phenotypic data are available at https://github.com/cmig-research-group/RSIGWAS (dMRI), https://github.com/precimed/mostest (sMRI) and https://www.github.com/norment/open-science (fMRI).

Acknowledgements

E.P.T. has been supported by the Foundation "De Drie Lichten" and The Simons Foundation Fund in The Netherlands. This work was partly performed on the TSD (Tjeneste for Sensitive Data) facilities, owned by the University of Oslo, operated and developed by the TSD service group at the University of Oslo, IT Department (USIT) (tsd-drift@usit.uio.no). We gratefully acknowledge support from the The Netherlands Organization for Scientific Research (NWO Gravitation: BRAINSCAPES: A Roadmap from Neurogenetics to Neurobiology - Grant No. 024.004.012, and VICI 452-16-015), The European Research Council (Advanced Grant No ERC-2018-AdG GWAS2FUNC 834057, Consolidator Grant No 101001062), American National Institutes of Health (NS057198, EB000790, 1R01MH124839, R01MH120025, R01MH122688), The Research Council of Norway (RCN) (229129, 213837, 324252, 300309, 273291, 223273, 248980, 276082, 323961), The South-East Norway Regional Health Authority (2019-108, 2022-073), KG Jebsen Stiftelsen (SKGJ-MED-021), EEA (#EEA2018-0573) collaborative grant in ASD: "Improving quality of life for Autism Spectrum Disorders patients by promoting strategies for early diagnosis and preventive measures" and EEA-RO-

NO-Grant 2014-2021 (contract No 6/2019), and the European Union's Horizon 2020 research and
innovation programme (No 847776 and 964874 and 801133, Marie Sklodowska-Curie grant
agreement). We want to acknowledge the participants and investigators of the UK Biobank, ABCD
Study, TOP, BUPGEN, PGC and FinnGen studies. The Norwegian Mother, Father and Child Cohort
Study is supported by the Norwegian Ministry of Health and Care Services and the Ministry of
Education and Research. We are grateful to all the participating families in Norway who take part in
this on-going cohort study. We thank the Norwegian Institute of Public Health (NIPH) for generating
high-quality genomic data. This research is part of the HARVEST collaboration, supported by the
Research Council of Norway (#229624). The graphical abstract includes images obtained from Servier
Medical Art (smart.servier.com). Data used in the preparation of this article were obtained from the
Adolescent Brain Cognitive Development SM Study (ABCD Study®) (https://abcdstudy.org), held in
the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than
10,000 children aged 9-10 and follow them over 10 years into early adulthood. The ABCD Study is
supported by the National Institutes of Health and additional federal partners under award numbers:
U01DA041022, U01DA041028, U01DA041048, U01DA041089, U01DA041106, U01DA041117,
U01DA041120, U01DA041134, U01DA041148, U01DA041156, U01DA041174, U24DA041123,
and U24DA041147. A full list of supporters is available at https://abcdstudy.org/federal-partners/. A
listing of participating sites and a complete listing of the study investigators can be found at
https://abcdstudy.org/principal-investigators.html. ABCD Study consortium investigators designed and
implemented the study and/or provided data but did not necessarily participate in analysis or writing of
this report. This manuscript reflects the views of the authors and may not reflect the opinions or views
of the NIH or ABCD Study consortium investigators. The ABCD data repository grows and changes
over time. The ABCD data used in this came from [NIMH Data Archive Digital Object Identifier
(10.151.54/1519007)].

Conflicts of interest

E.P.T., A.A.S., D.v.d.M, N.P., G.H., D.R., O.F., C.C.F., M.N., T.N., M.B., S.D., L.T.W., M.P.v.d.H.,
D.P. and T.K. declare no conflicts of interest. Dr. Andreassen has received speaker's honorarium from
Lundbeck and Janssen, and is a consultant to coretechs.ai. Dr. Dale is a Founder of and holds equity in
CorTechs Labs, Inc, and serves on its Scientific Advisory Board. He is a member of the Scientific
Advisory Board of Human Longevity, Inc. and receives funding through research agreements with
General Electric Healthcare and Medtronic, Inc. The terms of these arrangements have been reviewed
and approved by UCSD in accordance with its conflict of interest policies.

Author contributions

E.P.T., A.A.S., O.F., D.v.d.M. and O.A.A. conceived of the study. E.P.T. performed the analyses with A.A.S. and D.v.d.M. Neuroimaging data were processed by D.R., T.K., C.C.F., A.M.D, D.v.d.M and E.P.T. Genotype data were processed by E.P.T. and D.v.d.M. All authors contributed with conceptual input on methods and/or interpretation of results. E.P.T. wrote the manuscript and all authors contributed to the final manuscript.

References

1. Tononi, G., Edelman, G. M. & Sporns, O. Complexity and coherency: Integrating information in the brain. *Trends Cogn. Sci.* **2**, 474–484 (1998).

2. Jansen, A. G., Mous, S. E., White, T., Posthuma, D. & Polderman, T. J. C. What Twin Studies Tell Us About the Heritability of Brain Development, Morphology, and Function: A Review. *Neuropsychol. Rev.* **25**, 27–46 (2015).

3. Watanabe, K. *et al.* A global overview of pleiotropy and genetic architecture in complex traits. *Nat. Genet.* **51**, 1339–1348 (2019).

4. Solovieff, N., Cotsapas, C., Lee, P. H., Purcell, S. M. & Smoller, J. W. Pleiotropy in complex traits: Challenges and strategies. *Nat. Rev. Genet.* **14**, 483–495 (2013).

5. Shadrin, A. A. *et al.* Vertex-wise multivariate genome-wide association study identifies 780 unique genetic loci associated with cortical morphology. *Neuroimage* **244**, (2021).

6. van der Meer, D. *et al.* The genetic architecture of human cortical folding. *Sci. Adv.* **7**, (2021).

7. van der Meer, D. *et al.* Understanding the genetic determinants of the brain with MOSTest. *Nat. Commun.* **11**, (2020).

8. Roelfs, D. *et al.* Genetic overlap between multivariate measures of human functional brain connectivity and psychiatric disorders. *medRxiv* (2022).

9. Fan, C. C. *et al.* Multivariate genome-wide association study on tissue-sensitive diffusion metrics highlights pathways that shape the human brain. *Nat. Commun.* **13**, 2423 (2022).

10. Cross-Disorder Group of the Psychiatric Genomics Consortium. Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric Disorders. *Cell* **179**, 1469–1482 (2019).

11. Liu, S. *et al.* Multimodal neuroimaging computing: a review of the applications in neuropsychiatric disorders. *Brain Informatics* **2**, 167–180 (2015).

12. Sivakumaran, S. *et al.* Abundant pleiotropy in human complex diseases and traits. *Am. J. Hum. Genet.* **89**, 607–618 (2011).

13. Zhu, W. & Zhang, H. Why do we test multiple traits in genetic association studies? *J. Korean Stat. Soc.* **38**, 1–10 (2009).

14. Roelfs, D. *et al.* Shared genetic architecture between mental health and the brain functional

connectome in the UK Biobank. *BMC Psychiatry* **23**, (2023).

15.   Boedhoe, P. S. W. *et al.* Subcortical brain volume, regional cortical thickness, and cortical
surface area across disorders: Findings from the ENIGMA ADHD, ASD, and OCD working
groups. *Am. J. Psychiatry* **177**, 834–843 (2020).

16.   Hibar, D. P. *et al.* Cortical abnormalities in bipolar disorder: An MRI analysis of 6503
individuals from the ENIGMA Bipolar Disorder Working Group. *Mol. Psychiatry* **23**, 932–942
(2018).

17.   van Erp, T. G. M. *et al.* Cortical Brain Abnormalities in 4474 Individuals With Schizophrenia
and 5098 Control Subjects via the Enhancing Neuro Imaging Genetics Through Meta Analysis
(ENIGMA) Consortium. *Biol. Psychiatry* **84**, 644–654 (2018).

18.   van Erp, T. G. M. *et al.* Subcortical brain volume abnormalities in 2028 individuals with
schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Mol. Psychiatry* **21**, 547–
553 (2016).

19.   Schmaal, L. *et al.* Cortical abnormalities in adults and adolescents with major depression based
on brain scans from 20 cohorts worldwide in the ENIGMA Major Depressive Disorder Working
Group. *Mol. Psychiatry* **22**, 900–909 (2017).

20.   Schmaal, L. *et al.* Subcortical brain alterations in major depressive disorder: findings from the
ENIGMA Major Depressive Disorder working group. *Mol. Psychiatry* **21**, 806–812 (2016).

21.   van den Heuvel, M. P. & Sporns, O. A cross-disorder connectome landscape of brain
dysconnectivity. *Nat. Rev. Neurosci.* (2019). doi:10.1038/s41583-019-0177-6

22.   Rudie, J. D. *et al.* Altered functional and structural brain network organization in autism.
NeuroImage Clin. **2**, 79–94 (2013).

23.   Daianu, M. *et al.* Disrupted rich club network in behavioral variant frontotemporal dementia and
early-onset Alzheimer's disease. *Hum. Brain Mapp.* **37**, 868–883 (2016).

24.   Kelly, S. *et al.* Widespread white matter microstructural differences in schizophrenia across
4322 individuals: Results from the ENIGMA Schizophrenia DTI Working Group. *Mol.*
Psychiatry **23**, 1261–1269 (2018).

25.   Franke, B. *et al.* Genetic influences on schizophrenia and subcortical brain volumes: Large-scale
proof of concept. *Nat. Neurosci.* **19**, 420–431 (2016).

26.   Tissink, E. *et al.* Genome-wide association study of cerebellar volume provides insights into
heritable mechanisms underlying brain development and mental health. *Commun. Biol.* **5**,
(2022).

27.   Ohi, K. *et al.* Genetic correlations between subcortical brain volumes and psychiatric disorders.
Br. J. Psychiatry **216**, 280–283 (2020).

28.   Cheng, W. *et al.* Shared genetic architecture between schizophrenia and subcortical brain
volumes implicates early neurodevelopmental processes and brain development in childhood.
Mol. Psychiatry (2022). doi:10.1038/s41380-022-01751-z

29.   Cheng, W. *et al.* Genetic Association Between Schizophrenia and Cortical Brain Surface Area
803   and Thickness. *JAMA Psychiatry* 1–11 (2021). doi:10.1001/jamapsychiatry.2021.1435

804   30.   van der Meer, D. *et al.* Boosting Schizophrenia Genetics by Utilizing Genetic Overlap With
805   Brain Morphology. *Biol. Psychiatry* **92**, 291–298 (2022).

806   31.   Raj, A., Stephens, M. & Pritchard, J. K. FastSTRUCTURE: Variational inference of population
807   structure in large SNP data sets. *Genetics* **197**, 573–589 (2014).

808   32.   Bulik-Sullivan, B. *et al.* LD score regression distinguishes confounding from polygenicity in
809   genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).

810   33.   de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: Generalized Gene-Set
811   Analysis of GWAS Data. *PLoS Comput. Biol.* **11**, 1–19 (2015).

812   34.   Wang, K., Li, M. & Hakonarson, H. ANNOVAR: Functional annotation of genetic variants from
813   high-throughput sequencing data. *Nucleic Acids Res.* **38**, 1–7 (2010).

814   35.   Schork, A. J. *et al.* All SNPs are not created equal: Genome-wide association studies reveal a
815   consistent pattern of enrichment among functionally annotated SNPs. *PLoS Genet.* **9**, (2013).

816   36.   Kang, H. J. *et al.* Spatio-temporal transcriptome of the human brain. *Nature* **478**, 483–489
817   (2011).

818   37.   van der Meer, D. & Kaufmann, T. Mapping the genetic architecture of cortical morphology
819   through neuroimaging: progress and perspectives. *Transl. Psychiatry* **12**, 447 (2022).

820   38.   Bhaduri, A. *et al.* An atlas of cortical arealization identifies dynamic molecular signatures.
821   *Nature* **598**, 200–204 (2021).

822   39.   Hindley, G. *et al.* Multivariate genetic analysis of personality and cognitive traits reveals
823   abundant pleiotropy. *Nat. Hum. Behav.* (2023).

824   40.   Andreassen, O. A. *et al.* Improved Detection of Common Variants Associated with
825   Schizophrenia and Bipolar Disorder Using Pleiotropy-Informed Conditional False Discovery
826   Rate. *PLoS Genet.* **9**, (2013).

827   41.   Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic biology in
828   schizophrenia. *Nature* **604**, 502–508 (2022).

829   42.   Mullins, N. *et al.* Genome-wide association study of over 40,000 bipolar disorder cases provides
830   new insights into the underlying biology. *Nat. Genet.* **53**, 817–829 (2021).

831   43.   Wray, N. R. *et al.* Genome-wide association analyses identify 44 risk variants and refine the
832   genetic architecture of major depression. *Nat. Genet.* **50**, 668–681 (2018).

833   44.   Levey, D. F. *et al.* Reproducible Genetic Risk Loci for Anxiety: Results From ~200,000
834   Participants in the Million Veteran Program. *Am. J. Psychiatry* **177**, 223–232 (2020).

835   45.   Demontis, D. *et al.* Discovery of the first genome-wide significant risk loci for attention
836   deficit/hyperactivity disorder. *Nat. Genet.* **51**, 63–75 (2019).

837   46.   Grove, J. *et al.* Identification of common genetic risk variants for autism spectrum disorder. *Nat.*
838   *Genet.* **51**, 431–444 (2019).

839   47.   Middeldorp, C. M. *et al.* A Genome-Wide Association Meta-Analysis of Attention-
840   Deficit/Hyperactivity Disorder Symptoms in Population-Based Pediatric Cohorts. *J. Am. Acad.*
841   *Child Adolesc. Psychiatry* **55**, 896–905 (2016).

842   48.   Kurki, M. I., Karjalainen, J., Palta, P., Sipilä, T. P. & Kristiansson, K. FinnGen: Unique genetic
843   insights from combining isolated population and national health register data. *medRxiv* (2022).

844   49.   Lam, M. *et al.* Comparative Genetic Architectures of Schizophrenia in East Asian and European
845   Populations. *Nat. Genet.* **29**, (2019).

846   50.   Sporns, O. The complex brain: connectivity, dynamics, information. *Trends Cogn. Sci.* **26**,
847   1066–1067 (2022).

848   51.   Turelli, M. Heritable genetic variation via mutation-selection balance: Lerch's zeta meets the
849   abdominal bristle. *Theor. Popul. Biol.* **25**, 138–193 (1984).

850   52.   Luo, L. *et al.* Multi-trait analysis of rare-variant association summary statistics using MTAR.
851   *Nat. Commun.* **11**, 1–11 (2020).

852   53.   Dutta, D., Scott, L., Boehnke, M. & Lee, S. Multi-SKAT: General framework to test for rare-
853   variant association with multiple phenotypes. *Genet. Epidemiol.* **43**, 4–23 (2019).

854   54.   Atkinson, E. G. *et al.* Tractor uses local ancestry to enable the inclusion of admixed individuals
855   in GWAS and to boost power. *Nat. Genet.* **53**, 195–204 (2021).

856   55.   Drivas, T. G., Wojno, A. P., Tucker, B. A., Stone, E. M. & Bennett, J. Basal exon skipping and
857   genetic pleiotropy: A predictive model of disease pathogenesis. *Sci. Transl. Med.* **7**, 1–8 (2015).

858   56.   Ursini, G. *et al.* Convergence of placenta biology and genetic risk for schizophrenia. *Nat. Med.*
859   **24**, 792–801 (2018).

860   57.   Liharska, L. E. *et al.* A study of gene expression in the living human brain Living Brain Project
861   cohort. *bioRxiv* (2023).

862   58.   Fan, C. C. *et al.* Genotype Data and Derived Genetic Instruments of Adolescent Brain Cognitive
863   Development Study® for Better Understanding of Human Brain Development. *Behav. Genet.*
864   159–168 (2023). doi:10.1007/s10519-023-10143-0

865   59.   Loughnan, R. J. *et al.* Gene-experience correlation during cognitive development: Evidence
866   from the Adolescent Brain Cognitive Development (ABCD) Study. *bioRxiv* (2021).

867   60.   Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature*
868   **562**, 203–209 (2018).

869   61.   Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based
870   Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

871   62.   Desikan, R. S. *et al.* An automated labeling system for subdividing the human cerebral cortex
872   on MRI scans into gyral based regions of interest. *Neuroimage* **31**, 968–980 (2006).

873   63.   Fischl, B. *et al.* Whole Brain Segmentation: Automated Labeling of Neuroanatomical Structures
874   in the Human Brain. *Neuron* **33**, 341–355 (2002).

64.   Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic

Functional Connectivity MRI. *Cereb. Cortex* **28**, 3095–3114 (2018).

65.   Yeo, B. T. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic
functional connectivity. *J. Neurophysiol.* **106**, 2322–45 (2011).

66.   Gordon, E. M. *et al.* Generation and Evaluation of a Cortical Area Parcellation from Resting-
State Correlations. *Cereb. Cortex* **26**, 288–303 (2016).

67.   White, N. S., Leergaard, T. B., D'Arceuil, H., Bjaalie, J. G. & Dale, A. M. Probing tissue
microstructure with restriction spectrum imaging: Histological and theoretical validation. *Hum.*
Brain Mapp. **34**, 327–346 (2013).

68.   White, N. S. *et al.* Improved conspicuity and delineation of high-grade primary and metastatic
brain tumors using 'restriction spectrum imaging': Quantitative comparison with high B-value
DWI and ADC. *Am. J. Neuroradiol.* **34**, 958–964 (2013).

69.   Palmer, C. E. *et al.* Microstructural development from 9 to 14 years: Evidence from the ABCD
Study. *Dev. Cogn. Neurosci.* **53**, 101044 (2022).

70.   Lawrence, M. *et al.* Software for Computing and Annotating Genomic Ranges. *PLoS Comput.*
Biol. **9**, 1–10 (2013).

71.   The Gene Ontology Consortium. Gene Ontology: tool for the unification of biology. *Nat. Genet.*
25, 25–29 (2000).

72.   The Gene Ontology Consortium. The Gene Ontology resource: Enriching a GOld mine. *Nucleic*
Acids Res. **49**, D325–D334 (2021).

73.   Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and
annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).

74.   Merico, D., Isserlin, R., Stueker, O., Emili, A. & Bader, G. D. Enrichment map: A network-
based method for gene-set enrichment visualization and interpretation. *PLoS One* **5**, (2010).

75.   Kucera, M., Isserlin, R., Arkhangorodsky, A. & Bader, G. D. AutoAnnotate: A Cytoscape app
for summarizing networks with semantic annotations. *F1000Research* **5**, 1–14 (2016).

76.   Reimand, J. *et al.* Pathway enrichment analysis and visualization of omics data using g:profiler,
GSEA, Cytoscape and EnrichmentMap. *Nat. Protoc.* **14**, (2019).

77.   Paczkowska, M. *et al.* Integrative pathway enrichment analysis of multivariate omics data. *Nat.*
Commun. **11**, 1–16 (2020).

78.   Hawrylycz, M. *et al.* Canonical genetic signatures of the adult human brain. *Nat. Neurosci.* **18**,
1832–1844 (2015).

79.   Bahrami, S. *et al.* Dissecting the shared genetic basis of migraine and mental disorders using
novel statistical tools. *Brain* **145**, 142–153 (2022).

80.   Torgersen, K. *et al.* Shared genetic loci between depression and cardiometabolic traits. *PLoS*
Genet. **18**, 1–25 (2022).

81.   The GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across human
tissues. *Science (80-.).* **369**, 1318–1330 (2020).

82.   Magnus, P. *et al.* Cohort Profile Update: The Norwegian Mother and Child Cohort Study
(MoBa). *Int. J. Epidemiol.* **45**, 382–388 (2016).

83.   Choi, S. W., Mak, T. S. H. & O'Reilly, P. F. Tutorial: a guide to performing polygenic risk score
analyses. *Nat. Protoc.* **15**, 2759–2772 (2020).

84.   Choi, S. W. & O'Reilly, P. F. PRSice-2: Polygenic Risk Score software for biobank-scale data.
Gigascience **8**, 1–6 (2019).