Neural distinctiveness and discriminability in regions of the core network representations support
associative unitization

Authors: Steinkrauss, A. C.[1], Carpenter, C. M.[1], Tarkenton M. K.[2], Overman, A.A.[2] & Dennis,
N.A.[1]

The Pennsylvania State University[1], Elon University[2]

Corresponding Author:

Nancy A. Dennis

Professor

Department of Psychology

The Pennsylvania State University

450 Moore Building

University Park, Pennsylvania 16802

nad12@psu.edu

Key words: unitization, associative memory, neural reinstatement, multivariate pattern analyses, neuroimaging

Abstract:

Previous work has suggested unitized pairs behave as a single unit and more critically, are processed neurally different than those of associative memories. The current works examines the neural differences between unitization and non-unitized memory using fMRI and multivoxel analyses. Specifically, we examined the differences across face-occupation pairings as a function of whether the pairing was viewed as a person performing the given job (unitized binding) or a person saying they knew someone who had a particular job (non-unitized binding). The results show that at encoding, the angular gyrus can discriminate between unitized and non-unitized target trials. Additionally, during encoding, the medial temporal lobe (hippocampus and perirhinal cortex), frontal parietal regions (angular gyrus and medial frontal gyrus) and visual regions (middle occipital cortex) exhibit distinct neural patterns to recollected unitized and non-unitized targets. Furthermore, the medial frontal gyrus and middle occipital cortex show greater neural similarity for recollected unitized trials than those of recollected non-unitized trials. We conclude that visually unitized pairs may enhance unitization in older adults due to greater similarity of trials within the same condition during the encoding process.

1. Introduction:

The ability to form and remember correct associations is essential for recollection in episodic memory. Despite this necessity, the ability to successfully recall associations can be a difficult task in older adults (see Old & Naveh-Benjamin, 2008 for mata-analysis) For example, associating names with faces or stores with addresses can be increasingly difficult as a person ages. The associative deficit hypothesis attributes older adults' poor associative memory performance to an age-related difficulty in binding together novel pieces of information (Naveh-Benjamin, 2000; Naveh-Benjamin et al., 2004, 2007). Both unitization and visual imagery have been explored as encoding strategies intended to enhance the binding process and thus support older adults' associative memory (Kondo et al., 2005; Tu & Diana, 2021). The current study seeks to elucidate the neural basis underlying such associative memory strategies in aging individuals. Specifically, we use multivariate pattern classification and neural distinctiveness analyses to investigate whether the advantage afforded by a visual imagery task, used to promote unitization at encoding, supports discriminability and distinctiveness in the neural patterns of activation throughout the associative memory network in older adults.

Unitization is an encoding process used to bind arbitrary items together into a meaningful unit (Ahmad & Hockley, 2014; Bastin et al., 2013; Delhaye et al., 2014; Giovanello et al., 2006; Parks & Yonelinas, 2015; Parra et al., 2009; Quamme et al., 2007). Specifically, the strategy of unitization aims to integrate items in one bound unit which would reduce the involvement of hippocampal-based recollection and increase reliance on perirhinal-based familiarity processing (Ford et al., 2010), implying unitization may behave more similarly to a single item (Graf & Schacter, 1989; J. Haxby et al., 2001; Yonelinas et al., 1999). Most notably, this strategy has examined word-word associations for compound words compared to unrelated word pairs

(Ahmad et al., 2015), wherein older adults exhibit enhanced associative recollection for the

unitized pairings. Unitization strategies have also been applied to unrelated pairings (Huffer et

al., 2022; Memel & Ryan, 2017). For example, asking younger adults to encode face pairs as a

"married couple" resulted in better associative memory than two faces encoded as two unrelated

individuals (Parks & Yonelinas, 2015). Additionally, Huffer and colleagues (2022), show that

unrelated, yet unitized object pairs result in better associative memory compared to a non-

unitized presentation of the object pairing in older adults. Importantly, unitization instructions

have been shown to improve associative memory in older adults (Kamp et al., 2018; Liu et al.,

2022; Overman & Stephens, 2013; Zheng et al., 2016) by increasing both recollection and

familiarity-based responding (e.g., Liu et al., 2022).

76   One often-utilized strategy for inducing unitization has been visual imagery. By

visualizing, for example, a color as a feature of an item, it is more likely that a person will create

a unitized image of a single object with integrated characteristics (Bastin et al., 2013; Overman

& Stephens, 2013; Parks & Yonelinas, 2015; Tu & Diana, 2021; Zheng et al., 2015). For

example, Bastin and colleagues (2013) behaviorally, found enhanced associative memory in both

older and younger adults using item-color pairings with unitized compared to non-unitized

instructions. Using the same paradigm, Zheng and colleagues (2015) found that age differences

were reduced in the unitization instruction group due to familiarity-based source recognition.

Finally, Tu and Diana (2021) found greater associative memory for younger adults, in all

unitized imagery conditions which included a word with color, size, or both color and size.

Similarly, face-word pairings that encouraged unitization via visualization and intra-item binding

(e.g., imagine a person was a skier) were better remembered than those that encouraged inter-

item binding (e.g., the person had simply interacted with a skier) (Overman & Stephens, 2013;

Parks & Yonelinas, 2015). It is unknown if older adults will have similar capabilities as young

adults to use a unitization-based strategy for an associative memory task, in which they are

susceptible to memory declines in. The current work aims to see how in older adults' visual

imagery when using unitization-based strategies can enhance associative memory both

behaviorally and neurally. Specifically, this visual imagery will look at face-occupation pairings

and imagine those faces as *doing* the job associated with it to then *unitize* the pairing.

Neurally, associative memory has been shown to rely on activity in a common set of

regions, including the hippocampus (HC), medial frontal gyrus (MFG), and angular gyrus (AG)

(for a meta-analysis see Benoit & Schacter, 2015; Thakral et al., 2017). Specifically, associative

memories elicit increases in blood-oxygen-level-dependent (BOLD) activation within the HC for

recollection-based associative binding across novel items (Dennis, Bowman, et al., 2014; Dennis,

Johnson, et al., 2014; Piekema et al., 2010; Staresina & Davachi, 2010; Yonelinas et al., 2001).

Additionally, encoding-based strategies like unitization has been shown to elicit BOLD

activation in the perirhinal cortex (PrC) (Haskins et al., 2008; Memel & Ryan, 2017; Staresina &

Davachi, 2010), presumably because item-item associations are encoded and retrieved in a

manner more similar to that of an item in memory. BOLD activation in the MFG has been found

when detecting whether incoming information is congruent with its encoded memory state

(Bader et al., 2014). The MFG is also involved in the retrieval of schemas and is engaged when

people attempt to establish connections across associative items (Buuren et al., 2014; Kroes &

Fernández, 2012; Tse et al., 2011). Additionally, schema retrieval was found during increased

BOLD activation in the AG which allows for the encoding and consolidation of new information

(Wagner et al., 2015) Wagner and colleagues (2015) found multi-voxel representations of

different schema components in the AG during retrieval of novel, but related, information. They

concluded that the AG recombines encoded schemas into one integrated memory. In other

words, the AG may guide the binding of information from encoding to retrieval by condensing

encoded information for better retrieval recollection (Binder et al., 2009; Shimamura, 2011;

Wagner et al., 2015). In addition to this core associative network, occipital regions have been

consistently activated in visual memory tasks (Dennis et al., 2019; D. C. Park et al., 2004).

Specifically, classification accuracy discriminated between visual imagery and stimulus-driven

perception in the middle occipital cortex (MOC), showing the MOC is associated to visual

imagery (Stokes et al., 2009) and is linked to word-dependent source memory (Park et al., 2013).

Park and colleagues (2013) found that for multivariate neural activity in the MOC,

contextualized words are processed more like pictures, aiding to the imagery operations of

memory processing (Park et al., 2013).

While past work has focused on BOLD differences within this associative network, more

recent work is emerging that places an emphasis on understanding memory processing in aging

using multivariate analyses. Related to the current work, multivoxel pattern analysis (MVPA) in

older adults, has been used to identify unique neural patterns associated with subtle differences

in stimulus properties, such as classifying true and false memories (Carpenter et al., 2021;

Chamberlain et al., 2022; Dennis et al., 2022), categories like faces and houses (Carp et al.,

2011; Johnson et al., 2015) and recollection- and familiarity-based responding vs. responding at

chance (Kafkas et al., 2017). Previous work from our own group has found that neural patterns

across the encoding network, including those within prefrontal and occipital cortex, can reliably

distinguish between different types of associative memory during encoding (Dennis et al., 2019;

Elbich et al., 2021). The fact that such subtle differences in neural processing are detected in

older adults speaks to the idea that even for similar sets of information, different encoding

instructions can result in uniquely represented associations in the encoding network. That is,

neural representation for similar information can differ as a function of how information is

processed at encoding.

138   The examination of neural patterns also gives insight on the distinctiveness of different

memory processes and how neural representations contribute to and support episodic memory

performance (Sommer & Sander, 2022). By examining the distinct neural patterns related to

specific memory behaviors, such as associative recollection, we can gain further insight into how

unique representations during encoding lead to successful memory. Similar representations

within an encoding condition indicate that trials within that condition share overlapping

representations of information, which is to be expected when representing several related items

from the same category (Kriegeskorte & Kievit, 2013; Sommer & Sander, 2022). However,

when information is more distinctly processed in memory, neural patterns will represent this

uniqueness, reflected in less overlapping representation within a category of stimuli and instead,

more distinct patterns of neural activity. In neuroimaging memory research, different types of

associations, such as unitized associations compared to non-unitized associations, rely on

different neural processes (LaRocque et al., 2013; Staresina & Davachi, 2010). Knowing how

memory is represented and how those representations are related to successful retrieval is

important for understanding how we might be able to improve memory processing older adults.

It is unknown how older adults' neural activation may differ in their ability to encode and

retrieve information when using a unitization-based strategy versus an associative memory

strategy. Specifically using neural distinctiveness analyses in to identify differences in the neural

patterns underlying recollection-related activity and underlying unitization for associative

memories at each memory stage. Our focus is on regions within the core associative memory

network.

The current study aims to examine whether a visual imagery encoding strategy, intended

to promote unitization, will result in improved recollection-based associative memory in older

adults. The current study also aims to determine whether use of such an encoding strategy

enhances the distinctiveness of memory representations during both encoding and retrieval. If the

condition promoting unitization is effective in enhancing associative memory in aging, as we

have seen in previous work implementing a strategy-based unitization manipulation (Bastin et

al., 2013; Delhaye et al., 2014; Diana et al., 2008; Huffer et al., 2022; Quamme et al., 2007), then

we expect older adults to have greater recollection in the unitized condition than in the

associative memory condition. Neurally, we expect unitized and non-unitized memory for targets

to show neural discriminability within the core network, including the HC, PrC, MFG, AG, and

MOC. Further, we predict that the unitized encoding strategy pairs would show greater

distinctiveness than non-unitized associative memory pairs when taking into account subsequent

recollection of the associative information (Kriegeskorte & Kievit, 2013; Simmonite & Polk,

2022; Sommer & Sander, 2022). Our approach was undertaken in order to assess how encoding

strategies are processed successfully during an associative memory task.

2. Materials and Methods:

2.1 Subjects:

25 older adults were recruited from the Centre County community as a part of a larger study

including both younger and older adults. The older adults received monetary compensation for

their participation. Participants were screened for history of psychiatric and neurological illness,

head injury, stroke, learning disability, medication that affects cognitive and physiological

function, and substance abuse. On the day of the study, all participants provided written

informed consent for a protocol approved by The Pennsylvania State University Institutional

Review Board. All participants were native English speakers or had learned English before the

age of 8, with normal or corrected-to-normal vision and were right-handed. All participants had

completed high school. All 25 older adults were included in all analyses (M_{age}= 70.08 years,

SD_{age}= 6.77 years, range = 60-84 years; 14 female, 11 male). Participants identified as white (n =

23), more than one race (n = 1), and preferred not to answer (n = 1), and were all well-educated

(M_{years} = 16.50, SD_{years} = 2.65). All older adults scored above a 27 on the Mini Mental State

Examination (MMSE; Folstein et al., 1983; M = 29.22, SD = 1.02) and below a 3 on the Geriatric

Depression Scale (GDS; Yesavage, 1998; M = 0.88, SD = 1.17).

2.2 Regions of Interest (ROIs)*:* Based upon previous work mentioned above, we restricted our

analysis to the angular gyrus (AG), hippocampus (HC), perirhinal cortex (PrC), medial frontal

gyrus (MFG), and middle occipital cortex (MOC) (Bader et al., 2014; Binder et al., 2009;

Dennis, Bowman, et al., 2014; Dennis, Johnson, et al., 2014; Haskins et al., 2008; Stokes et al.,

2009). The ROIs were defined anatomically and created using the human AAL Pickatlas through

SPM12 (Folstein et al., 1983) (See Fig. 1).

2.3 Stimuli & Procedure: The current design and stimuli were modified from Overman &

Stephens (2013) and previously reported in Ricupero and colleagues (2023) where a young adult

sample was analyzed. The experimental stimuli consisted of 144 black and white photographs of

faces (see Criss & Shiffrin, 2004 for standardization details) and 144 single-word occupations

(ex: pianist; welder), with a majority of the occupations taken from Yovel & Paller (2004) and

additional occupations added as needed. During encoding, participants were shown an image of a

face and a name tag stating "Hello, I'm [occupation]" ('doing' condition) or a speech bubble

stating "I know [occupation]" ('speaking' condition). Participants were asked to imagine the

face-occupation association and remember the pairings using one of two strategies designed to

either promote unitization ('doing') or simply to create an association ('speaking'). Specifically,

in the 'doing' condition participants were asked to imagine the pictured individual performing

actions related to the occupation. In the 'speaking' condition participants were asked to imagine

the pictured individual knowing someone else with the given occupation. During practice,

participants were asked to repeat the instructions back and explain what they should be imaging

during the two strategy tasks. Participants were prompted by an instruction screen as to which

strategy they should employ, with both encoding strategies used in each run. Participants were

also asked to indicate, using a four-point button box, how easy or how difficult it was to use the

given strategy for each unique face and job pairing (response options included: very difficult,

somewhat difficult, somewhat easy, very easy). Finally, the background screen color randomly

alternated between either yellow or blue, with each appearing 50% of the time (analyses related

to this color manipulation were not included in the current set of analyses).

During encoding trials, participants received a prompt for a duration of 2.5 seconds

indicating which of the strategies to use (i.e., speaking). Participants then viewed 9 trials

followed by a prompt to use the other strategy (i.e., doing) and then viewed 9 trials. This process

repeated itself twice per run. Eleven participants completed this version of task before it was

changed to alternate between instructions after 18 trials (N=14), and prompt lasting 5 seconds.

There were no repetitions of strategy in each run (i.e., participants saw one doing and one

speaking strategy block per run). This change was made in response to feedback from older

adults. There were no significant differences in any result between versions. During retrieval

trials, participants were presented with both target pairs and recombined lure pairs on a white

background. Most face-job pairs were recombined within the same condition[1]. Participants

responded during retrieval using a standard Remember-Know-New paradigm (Yonelinas, 2002).

They responded 'Remember' if they remembered specific details about the face-occupation pair,

'Know' if they believed they had seen the pairs together previously, but could not remember

specific details of the pair, and 'New' if they believed the pair was not presented together

previously. Each retrieval run included 36 trials, 24 targets and 12 recombined lures. The order

of runs included two encoding runs, followed by two retrieval runs, repeated twice (for a total of

four encoding and four retrieval runs). All encoding and retrieval trials were presented for 5

seconds. Participants were provided with practice prior to beginning the task to facilitate learning

the two encoding strategies. Participants were given a total of 8 trials at encoding in which they a

prompt indicating they should use one of the strategies, then viewed 4 trials, and then were

presented with a prompt to use the other strategy and viewed another 4 trials. Following

encoding practice, they also had 8 trials of retrieval practice with 6 of those trials being targets.

(See Fig. 2).

2.5 Image Acquisition

Structural and functional images were acquired using a Siemens 3-T scanner equipped

with a 20-channel head coil, parallel to the AC–PC plane. Structural images were acquired with a

2300-msec repetition time, a 2.28-msec echo time, a 256-mm field of view, 192 axial slices, and

a 1.0-mm slice thickness for each participant. Echoplanar functional images were acquired using

a descending acquisition, a 2500-msec repetition time, a 33-msec echo time, a 192-mm field of

---

[1] Due to an error in programming, a subset of lures were recombined between (rather than within) condition. All between condition lures were removed from behavioral and imaging analyses.

view, a 80° flip angle, and 64 axial slices with a 2.0-mm slice thickness resulting in 2.0-mm

isotroFpic voxels.

2.6 Anatomical Data Processing

A total of 1 T1-weighted (T1w) images were found within the input BIDS dataset. The

T1-weighted (T1w) image was corrected for intensity non-uniformity (INU) with

N4BiasFieldCorrection (Tustison et al., 2010), distributed with ANTs 2.2.0 (Avants et al., 2008,

RRID:SCR_004757), and used as T1w-reference throughout the workflow. The T1w-reference

was then skull-stripped with a Nipype implementation of the antsBrainExtraction.sh workflow

(from ANTs), using OASIS30ANTs as target template. Brain tissue segmentation of

cerebrospinal fluid (CSF), white-matter (WM) and gray-matter (GM) was performed on the

brain-extracted T1w using fast (FSL 5.0.9, RRID:SCR_002823, Zhang et al., 2001). Brain

surfaces were reconstructed using recon-all (FreeSurfer 6.0.1, RRID:SCR_001847, Dale et al.,

1999), and the brain mask estimated previously was refined with a custom variation of the

method to reconcile ANTs-derived and FreeSurfer-derived segmentations of the cortical gray-

matter of Mindboggle (RRID:SCR_002438, Klein et al., 2017). Volume-based spatial

normalization to one standard space (MNI152NLin2009cAsym) was performed through

nonlinear registration with antsRegistration (ANTs 2.2.0), using brain-extracted versions of both

T1w reference and the T1w template. The following template was selected for spatial

normalization: ICBM 152 Nonlinear Asymmetrical template version 2009c [Fonov et al., 2009),

RRID:SCR_008796; TemplateFlow ID: MNI152NLin2009cAsym],

2.7 Functional data preprocessing

For each of the 8 BOLD runs found per subject (across all tasks and sessions), the

following preprocessing was performed. First, a reference volume and its skull-stripped version

were generated using a custom methodology of fMRIPrep. Head-motion parameters with respect

to the BOLD reference (transformation matrices, and six corresponding rotation and translation

parameters) are estimated before any spatiotemporal filtering using mcflirt (FSL 5.0.9, Jenkinson

et al., 2002). BOLD runs were slice-time corrected using 3dTshift from AFNI 20160207 (Cox &

Hyde, 1997, RRID:SCR_005927). Susceptibility distortion correction (SDC) was omitted. The

BOLD reference was then co-registered to the T1w reference using bbregister (FreeSurfer)

which implements boundary-based registration (Greve & Fischl, 2009). Co-registration was

configured with six degrees of freedom. The BOLD time-series (including slice-timing

correction when applied) were resampled onto their original, native space by applying the

transforms to correct for head-motion. These resampled BOLD time-series will be referred to as

preprocessed BOLD in original space, or just preprocessed BOLD. The BOLD time-series were

resampled into standard space, generating a preprocessed BOLD run in

MNI152NLin2009cAsym space. First, a reference volume and its skull-stripped version were

generated using a custom methodology of fMRIPrep. Several confounding time-series were

calculated based on the preprocessed BOLD: framewise displacement (FD), DVARS and three

region-wise global signals. FD was computed using two formulations following Power (absolute

sum of relative motions, Power et al., 2014)) and Jenkinson (relative root mean square

displacement between affines, Jenkinson et al., 2002)). FD and DVARS are calculated for each

functional run, both using their implementations in Nipype (following the definitions by Power

et al., 2014). The three global signals are extracted within the CSF, the WM, and the whole-brain

masks. The head-motion estimates calculated in the correction step were also placed within the

corresponding confounds file. The confound time series derived from head motion estimates and

global signals were expanded with the inclusion of temporal derivatives and quadratic terms for

each (Satterthwaite et al., 2013). Frames that exceeded a threshold of 0.5 mm FD or 1.5

standardized DVARS were annotated as motion outliers. All resamplings can be performed with

a single interpolation step by composing all the pertinent transformations (i.e. head-motion

transform matrices, susceptibility distortion correction when available, and co-registrations to

anatomical and output spaces). Gridded (volumetric) resamplings were performed using

antsApplyTransforms (ANTs), configured with Lanczos interpolation to minimize the smoothing

effects of other kernels (Lanczos, 1964). Non-gridded (surface) resamplings were performed

using mri_vol2surf (FreeSurfer).

2.8 Multivariate Pattern Analyses:

All analyses were conducted in normalized, MNI space. The anatomical masks were

drawn from the AAL pick atlas and were co-registered to a subject's brain in MNI space. All

trials were modeled within each individual trial GLMs. To estimate neural activity associated

with individual trials, separate GLMs on unsmoothed data were estimated in SPM12 defining

one regressor for each trial at encoding and retrieval (172 total for each phase). An additional 6

nuisance regressors were included in each run corresponding to motion. Whole-brain parameter

maps were generated for each trial for encoding and retrieval for each participant. In any given

parameter map, the value in each voxel represents the regression coefficient for that trial's

regressor in multiple regression containing all other trials in the run and the motion parameters.

These beta parameter maps were concatenated across runs and submitted to CoSMoMVPA

toolbox (Oosterhof et al., 2016) for pattern classification (Mumford et al., 2012), and

representational similarity/distinctiveness (Haxby et al., 2001) analyses.

Neural Pattern Classification

Given our interest in determining how unitization and non-unitized associative memory

strategies are discriminated in each region in the associative memory network, classification

analyses were conducted to determine if a classifier was able to discriminate between unitization

or non-unitized memory target trials in our selected ROIs. Separate classification accuracies

were computed between the foregoing trial types at both encoding and retrieval using a support

vector machine (SVM) classifier with a linear kernel using all voxels within each ROI (Mumford

et al., 2012). Training and testing were performed following an n-fold cross-validation wherein

the data was split across the 4 runs. The classifier was trained on 3 runs and tested on the 4th,

repeating across different folds. Group-level results were generated from averaging across

validation folds from all possible train-data/test-data permutations from the individual participant

level. Finally, we tested whether a classifier was significantly able to discriminate neural patterns

above chance between the two target types using a one-tailed one-sample t-test for classification

accuracy within each ROI. All t-tests were corrected for multiple comparisons using Benjamini-

Hochberg corrections and any significance was confirmed with permutation testing (using

10,000 Monte-Carlo simulations). The MVPA was conducted only on target trials, absent of

behavior. Classification in the current study begets the idea of pattern distinctiveness compared

to chance. The current classification analysis was run to determine how discriminable target

trials were from one another. In order to determine how behavior was affected by these trial

types, we next conducted pattern distinctiveness analyses accounting for that behavior.

Neural Pattern Distinctiveness

Neural distinctiveness metrics examine how distinct neural patterns are from one another

in different conditions in order to determine if brain regions are able to discriminate between

different conditions or stimuli (Haxby et al., 2001; Kriegeskorte, 2008; LaRocque et al., 2013).

Pattern distinctiveness analyses were conducted in the current study to examine the

representation of stimuli associated with unitization and non-unitized memory on subsequently

recollected targets. Thus, trials will be unequal between conditions and participants. Recollected

targets are encoded trials that are subsequently responded to with the subjective response of

'Remember' during retrieval, indicating a detailed memory for that pair. For the current analyses,

the similarity of activation patterns across different trials of the same category was calculated

(within-category correlation, a measure of reliability; Simmonite & Polk, 2022) in each

participant for participants subsequently recollected trials (e.g., correlation between all beta

parameter maps for unitized subsequently recollected trials correlated individually to all beta

parameter maps for all other unitized subsequently recollected trials at encoding) and a between-

category similarity score for each participant's subsequently recollected trials (e.g., one beta

parameter map for a unitized subsequently recollected trial correlated individually to all beta

parameter maps for non-unitized subsequently recollected trials, done for all unitized trials

individually and then averaged) (Haxby et al., 2001). As described in Simmonite and Polk

(2022), increased within-category similarity indicates greater reliability and consistency of

neural patterns within a given condition. Next, an overall distinctiveness score was calculated by

taking the mean of the unitized-within and non-unitized-within similarities scores and

subtracting the between similarity score. This value was then compared to 0 using one-sample t-

tests, for each ROI, to determine if distinctiveness differed from chance. To further examine

whether the discrete distinctiveness was driven by within-category differences, the within-

category similarity for the unitized and non-unitized conditions were compared using paired t-

tests within each ROI. This process was repeated for both encoding and retrieval trials

separately. (See Fig. 3 for an example).

3. Results:

Behavioral Results

Overall hit rates across conditions were relatively high, (unitized M= 0.79 SD=0.11 and

non-unitized M=0.68, SD=0.15), as were correct rejection rates (unitized M=0.54, SD=0.21 and

non-unitized M=0.52, SD=0.16). A series of ANOVAs were run between the unitized and non-

unitized memory trials for recollected hits, familiarity hits, and correct rejections (CR), with

version of the task included as a covariate. With respect to recollected hit rates, there was a main

effect of condition, $F(1,23) = 61.45$, $p < .001$, $d = .728$, such that the unitized memory trials had

greater recollected hits (M=0.54 SD=0.20) compared to the non-unitized memory trials (M=0.41

SD=0.17). The main effect of version, $F(1,23) = 0.41$, $p = .53$, $d = .018$, and the interaction

between version and condition, $F(1,23) = 0.03$, $p = .88$, $d = .001$, were not significant. With

respect to familiarity hits, we found no main effects between unitized (M=0.25 SD=0.16) and

non-unitized (M=0.27 SD=0.13) memory trials with version as a covariate, condition, $F(1,23) =$

0.76, $p = .39$, $d = .032$, version, $F(1,23) = 0.12$, $p = .73$, $d = .005$, or a condition by version

interaction, $F(1,23) = 0.36$, $p = .55$, $d = .015$. Similarly, for CR, we found no main effects for

condition, $F(1,23) = 0.13$, $p = .72$, $d = .006$, version, $F(1,23) = 2.21$, $p = .15$, $d = .088$, or a

condition by version interaction, $F(1,23) = 0.04$, $p = .85$, $d = .002$.

Classification Results

To examine whether classifiers were able to significantly discriminate between our two

target conditions, two multivoxel pattern analyses were run. The first classification analysis

attempted to classify all unitized and non-unitized memory trial targets at encoding, and the

second to classify all unitized and non-unitized memory trial targets at retrieval. Comparing

classification of unitized and non-unitized memory trial targets at encoding against chance (50%)

showed that in the AG, the classifier accuracy was significantly above chance (AG: $M_{accuracy}$ =

0.540, $t(24)$ = 1.94, $p < .05$). To confirm significance, permutation testing was conducted, using

10,000 Monte-Carlo simulations performed within the AG. In doing so, the AG maintained

significance, p = .02. Furthermore, in addition to identifying regions showing signals for unitized

targets that were discriminable from those of non-unitized targets, we also tested whether there

were univariate activation differences in the AG. In the AG, the region showing above-chance

classification performance, we computed an ANOCVA with classification accuracy, including

mean univariate activation of unitized targets versus baseline and non-unitized targets versus

baseline as nuisance covariates. The two covariates did not show a significant difference for

target classification (p's > .7). No other ROIs were significant at encoding (HC: t(24)= -0.97, p =

.83; MFG: t(24)= 0.544, p = .30; MidOcc: t(24)= 1.304, p = .10; PrC: t(24)= 0.153, p = .44. At

retrieval, no regions of interest showed significant classification of unitized and non-unitized

memory against chance (AG: t(24)= 1.57, p = .06; HC: t(24)= -0.52, p = .70; MFG: t(24)= 1.53,

p = .07; MidOcc: t(24)= 0.57, p = .29; PrC: t(24)= 0.47, p = .32). See Table 1 for all means and

standard deviations.

Encoding Distinctiveness

In order to examine neural discriminability related to successful memory at encoding, a

neural distinctiveness calculation (Haxby et al., 2001) was conducted for unitized and non-

unitized memory subsequently recollected targets (within-category similarity minus between-

category similarity) for each encoding condition. At encoding, overall distinctiveness of

recollected targets, collapsed across condition, was significantly greater than 0 within all ROIs

[AG: $t(24) = 4.43, p < .001$; HC: $t(24) = 2.98, p < .01$; MFG: $t(24) = 5.49, p < .001$; MOC: $t(24)$

$= 4.79, p < .001$; PrC: $t(24) = 2.31, p < .05$]. The within-condition similarity scores associated

with both conditions were then compared to determine if one condition showed higher similarity

in a given region compared to the other condition. A direct comparison of within-condition

similarity between conditions found significant differences in the MFG and MOC [MFG: $t(24) =$

$2.72, p < .05$, 95% CI[0.01, 0.04]; MOC: $t(24) = 2.11, p < .05$, 95% CI[0.00, 0.09]] such that the

unitized condition exhibited greater within-condition similarity than the non-unitized condition

in the MFG ($M_{unit} = 0.09$ $SD_{unit} = 0.05$, $M_{assoc} = 0.07$ $SD_{assoc} = 0.04$) and MOC ($M_{unit} = 0.31$ SD_{unit}

$= 0.16$, $M_{assoc} = 0.27$ $SD_{assoc} = 0.12$). There were no other significant differences in any other

ROIs [HC: $t(24)=0.97$, p=.34, 95% CI[-0.02, 0.06]; PrC: $t(24)=0.53$, p=.60, 95% CI[-0.02, 0.04];

AG: $t(24)=0.13$, p=.90, 95% CI[-0.03, 0.04]] (See Fig. 4).

Retrieval Distinctiveness

In order to examine neural discriminability between two successful retrieval conditions, a

neural distinctiveness calculation was conducted for unitized and non-unitized memory

recollected targets (within category similarity minus between category similarity). At retrieval,

overall distinctiveness of subsequently recollected targets, collapsed across condition, was not

significantly greater than 0 within any ROIs [AG: $t(24)= -1.17$, p=.87; HC: $t(24)= .62$, p = .27;

MFG: $t(24)= -.02$, p = .51; MOC: $t(24)= -1.00$, p = .84; PrC: $t(24)= 1.06$, p = .15]. Since no

region of interest was significantly distinct above 0, these conditions were not compared to

determine if one condition showed higher similarity in a given region compared to the other

condition.

Exploratory Analysis

As an exploratory analysis, correlation coefficients between distinctiveness in all ROIs

were computed in order to examine cross-region relationships in distinctiveness measures during

encoding (see Supplemental Fig. 1 for correlations and scatterplot) (see Simmonite & Polk, 2022

for similar analyses). Across all participants, there was a significant positive relationship

between the AG and MFG [$r(23) = .64$, $p < .001$.], AG and PrC [$r(23) = .71$, $p < .001$.], MOC

and HC [$r(23) = .48$, $p < .05$.], HC and PrC [$r(23) = .40$, $p < .05$.], and MFG and MOC [$r(23) =$

$.47$, $p < .05$.]. The relationship between all other ROIs were not significant (all p's $> .05$: AG and

HC [$r(23) = .24$, $p = .26$], AG and MOC [$r(23) = .32$, $p = .12$], HC and MFG [$r(23) = .13$, $p =$

$.54$], MFG and PrC [$r(23) = .31$, $p = .14$], and MOC and PrC [$r(23) = .28$, $p = .17$.]). All

significant findings were confirmed using Benjamini-Hochberg (BH) corrections.

4. Discussion:

The goal of the current study was to examine the underlying neural mechanisms of

unitization, induced through the manipulation of encoding instructions that employed visual

imagery. During encoding, participants were given two strategies, one of which promoted

unitization of face-occupation pairs through intra-item binding and visualization, while the other,

non-unitized encoding strategy, used visualization to promote inter-item binding. Specifically, in

the unitized encoding strategy, participants saw a face-occupation pairing and were prompted to

imagine the person they were viewing as doing the occupation. In contrast, in the non-unitized

encoding strategy participants saw a face-occupation pairing and were prompted to imagine the

person they were viewing as knowing someone with the occupation. In alignment with previous

work using this paradigm with younger adults (Ricupero et al., 2023), and older adults (Overman

& Stephens, 2013), as well as similar studies in older adults, (e.g., Bastin et al., 2013; Delhaye et

al., 2014; Diana et al., 2008; Parks & Yonelinas, 2015; Quamme et al., 2007) we found that the

unitized condition resulted in greater recollected hits than the non-unitized condition in our

sample of older adults. Results support the idea that intra-item binding leads to enhanced

associative memory and highlights the ability of older adults to take advantage of such a strategy.

We hypothesized that across both memory phases, information processed within the

unitized encoding condition would show greater neural discriminability and neural

distinctiveness than information in the non-unitized encoding condition (Kriegeskorte & Kievit,

2013; Simmonite & Polk, 2022; Sommer & Sander, 2022). However, only within the AG at

encoding, did MVPA exhibit greater than chance classification between the two encoding

conditions. This is consistent with prior work in which AG activation is associated with the

presence of schemas during encoding and posited to aid participants to consolidate new

information (Binder et al., 2009; Shimamura, 2011; Wagner et al., 2015), including contributing

to the encoding of a schema into a bound memory (Wagner et al., 2015). In the current study,

classification within the AG may have reflected the difference in processes used to encode an

integrated schema of person and their occupation as opposed to the mere association amongst a

face and an occupation of a separate individual. This differs from previous aging work in our lab

(Dennis et al., 2019) in which it was found that neural patterns across the encoding network can

reliably distinguish between two different types of visual associations. One difference across

results may be that the current study used similar configurations for both conditions even though

the visual associations created by the encoding instructions differed. Specifically, our prior study

found that when encoding face-scene associations, regions within prefrontal and occipital cortex,

as well as the perirhinal cortex, reliably discriminated between item-item and item-context

presentations in older adults. While both the current and prior study controlled for the content of

information across associative encoding conditions, our previous study included a greater

configural difference across conditions, whereas a similar visual layout was used across

encoding conditions in the current study (see Fig. 1). Taken together, the results suggest that

encoding processing differences alone may not be enough to promote distinct neural patterns

during associative memory in aging within medial temporal regions including the hippocampus

and perirhinal cortex. Rather, it may be that in order to represent information uniquely across

different encoding conditions, there must be both configural differences and differences in visual

encoding between conditions. This aligns with computational modeling work that has suggested

a primary age difference in associative memory may be a decrease in the uniqueness of encoded

features that enable accurate discrimination of old versus new information (Stephens &

Overman, 2018). Further work should continue to explore how encoding strategy and configural

differences across encoding conditions supports such neural discriminability.

Similar to the concept of neural discriminability across encoding conditions, the

investigation of neural distinctiveness, related to behavioral outcomes (Haxby et al., 2001),

allows us to investigate whether neural patterns of successfully recollected unitized and non-

unitized memories are presented uniquely both within and across study conditions. In the current

study, subsequently recollected targets from both conditions exhibited above-chance

distinctiveness across all regions of interest (HC, PrC, MFG, AG, and MOC), indicating that

neural patterns associated with these associative pairs are distinct from one another within each

condition, during encoding. The fact that recollected trials were only distinct from one another at

encoding, and not retrieval, suggests that older adults utilize encoding based strategies and

schemas to later benefit their recollection for face-occupation pairings. An exploratory analysis

examining the correspondence of distinctiveness across regions found significant cross-region

relationships in distinctiveness for the two encoding conditions. Specifically, the HC and PrC

had a positive relationship between each other's distinctiveness indicating these two MTL

regions elicit similar activation during encoding across conditions for recollected target trials.

Additionally, the AG and MFG, AG and PrC, MOC and HC, and MFG and MOC were also

positively correlated to each other with regard to distinctiveness amongst recollected target trials

during encoding. Together the results indicate that, not only do MTL regions exhibit similar

distinct activation between one another, but so do prefrontal and visual regions as well. This

finding suggests that for older adults, processing within different regions in the brain are related

to each other when discriminating between recollected trials.

When directly comparing pattern similarity of the unitized condition to that of the non-

unitized condition, the reliability of neural activity of each condition was significantly different

from each other in the MFG and the MOC, such that the unitized condition exhibited greater

reliability (greater within-condition similarity of trials) than the non-unitized condition. This

suggests that when processing targets that were later correctly recollected, participants encoded

the trials across the unitized condition in a more similar manner than they did those within the

non-unitized condition. Since analyses were completed on targets that were later recollected,

greater reliability related to within-category activation may likely underscore less detailed-based

memory traces. Additionally, this suggests that the MFG and MOC may play a role in

distinguishing between associations that are encoded in a different manner. The MFG has been

shown to aid in the detection of congruent information from encoding to retrieval as well as to

aid in supporting the retrieval of schemas (Kroes & Fernández, 2012; Tse et al., 2011). Since

neural patterns in the MFG associated with the unitized condition were more similar to other

recollected trials in the same condition, it further supports the idea that the MFG contributes to

the encoding of integrated schemas, such as imagining a relationship between a person and their

occupation. Since the unitization condition in the current study asked older adults to use schemas

to bind the face-occupation pairing, the additional detail associated with the schemas may have

led to greater recollection in the unitized condition compared to the non-unitized condition.

However, further work such as a questionnaire to identify strategy usage may be necessary to

determine how these schemas affected memory performance.

In the unitized condition, it was emphasized that they should be imagining and

visualizing the face-occupation pairing especially to a greater extent than the non-unitized

condition. Greater unitized trial activation in the MOC supports the idea that older adults

engaged in the imagery strategy that they were instructed to use, since the MOC is crucial for

visual imagery (e.g., Pidgeon et al., 2016; Stokes et al., 2009). In this study, unitization was

designed to be implemented as a top-down mechanism, aiding in the binding of two unrelated

items (face-occupation pairing) into a single representation using stored schemas associated with

occupation actions. Invoking occupation schemas, combined with generating mental imagery of

the actions associated with the occupation, allowed participants to draw on pre-existing

information during associative encoding, in a manner similar to that used in previous work

(Bastin et al., 2013; Memel & Ryan, 2017). Bastin and colleagues (2013) asked participants to

form visual associations and found that older adults had an increased associative memory

compared to the non-unitized strategy. In a similar manner, Memel and Ryan (2017) used a

visual integration task to bind objects and contexts, in which they placed objects within a logical

scene in order to promote schema-related unitization during encoding. This presentation resulted

in better associative recollection and increased activation in the prefrontal cortex within their

older participants. They concluded visual integrated unitization may be mediated by recollective

based encoding processes for older adults (Memel & Ryan, 2017).

5. Limitations/Future Directions:

While the current findings extend our knowledge of the role of schemas during encoding for older adults, there were some limitations to the study. Specifically, we cannot be certain that the unitized condition is truly being bound as a single unit, only that behavioral differences across encoding conditions suggest stronger memory for the unitization-based strategy. The unitized condition can only be concluded as being different from the non-unitized associative condition, in a manner that is beneficial to older adults' memory performance. Future work should aim to identify whether these differences in encoding strategies are due to unitization of unrelated information into a singular bound unit. Additionally, further research is needed to determine if a visualization unitization strategy is the most effective strategy to achieve unitized binding. In addition, the benefit of unitization has often been associated with greater use of familiarity, particularly in older adults (Delhaye & Bastin, 2018; Diana, Ranganath & Yonelinas, 2008). However, recent work in the realm of neuroimaging has also indicated that recollection may also be important to mediating the effect of unitization on memory (Memel & Ryan, 2017). In the current task, familiarity-based hit counts were very low and thus we chose to focus solely on recollection-based hits. Future work should aim to investigate and disentangle the role of familiarity and recollection in unitization in terms of neural distinctiveness.

6. Conclusions

Overall, the results show that older adults are able to benefit from associative encoding instructions that emphasize unitized binding and visualization, exhibiting greater recollection of targets in the unitized compared to non-unitized condition. Contrary to our predictions regarding neural discriminability, only the AG was able to discriminate between unitized and non-unitized targets during encoding, suggesting that more contextual differences may be required for older

adults to recruit other associative memory regions (Dennis et al., 2019). In contrast, we found

that neural patterns related to unitized and non-unitized associations were distinct from one

another within the core associative network and visual imagery regions. Specifically, when

comparing the similarity of unitized versus non-unitized trials, the MOC and MFG exhibited

greater reliability at encoding within neural representations of later recollected unitized pairs

compared to non-unitized pairs. Overall, the results suggest that unitization may help older adults

bind new information together into single units when using visual imagery strategies during

encoding, which may be in part due to the ability of visual imagery regions to distinguish

between types of associative information.

Acknowledgements

We would like to thank Courtney Gerver, Harini Babu, Kayla McGraw, Valerie Goodwin, Chloe

Hultman and Min Sung Seo for helping with data collection and analysis. This paper was

supported by the National Institutes of Health under grant R15 AG052903 awarded to A.A.O and

N.A.D.

Availability of Data and Materials

The datasets generated and analyzed during the current study are available from the

corresponding author on reasonable request. None of the experiments were preregistered.

References

Ahmad, F. N., Fernandes, M., & Hockley, W. E. (2015). Improving associative memory in older adults

with unitization. *Aging, Neuropsychology, and Cognition*, *22*(4), 452–472.

https://doi.org/10.1080/13825585.2014.980216

Ahmad, F. N., & Hockley, W. E. (2014). The role of familiarity in associative recognition of unitized

compound word pairs. *Quarterly Journal of Experimental Psychology*, *67*(12), 2301–2324.

https://doi.org/10.1080/17470218.2014.923007

Allen, R. J., Vargha-Khadem, F., & Baddeley, A. D. (2014). Item-location binding in working memory: Is it

hippocampus-dependent? *Neuropsychologia*, *59*, 74–84.

https://doi.org/10.1016/j.neuropsychologia.2014.04.013

Avants, B., Epstein, C., Grossman, M., & Gee, J. (2008). Symmetric diffeomorphic image registration with

cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain.

Medical Image Analysis, *12*(1), 26–41. https://doi.org/10.1016/j.media.2007.06.004

Bader, R., Opitz, B., Reith, W., & Mecklinger, A. (2014). Is a novel conceptual unit more than the sum of

its parts?: FMRI evidence from an associative recognition memory study. *Neuropsychologia*, *61*,

123–134. https://doi.org/10.1016/j.neuropsychologia.2014.06.006

Bastin, C., Diana, R. A., Simon, J., Collette, F., Yonelinas, A. P., & Salmon, E. (2013). Associative memory

in aging: The effect of unitization on source memory. *Psychology and Aging*, *28*(1), 275–283.

https://doi.org/10.1037/a0031566

Benoit, R. G., & Schacter, D. L. (2015). Specifying the core network supporting episodic simulation and

episodic memory by activation likelihood estimation. *Neuropsychologia*, *75*, 450–457.

https://doi.org/10.1016/j.neuropsychologia.2015.06.034

Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where Is the Semantic System? A Critical

Review and Meta-Analysis of 120 Functional Neuroimaging Studies. *Cerebral Cortex*, *19*(12),

2767–2796. https://doi.org/10.1093/cercor/bhp055

Buuren, V. M., Kroes, M. C. W., Wagner, I. C., Genzel, L., Morris, R. G. M., & Fernandez, G. (2014). Initial

Investigation of the Effects of an Experimentally Learned Schema on Spatial Associative Memory

in Humans. *Journal of Neuroscience*, *34*(50), 16662–16670.

https://doi.org/10.1523/JNEUROSCI.2365-14.2014

Carp, J., Park, J., Polk, T. A., & Park, D. C. (2011). Age differences in neural distinctiveness revealed by

multi-voxel pattern analysis. *NeuroImage*, *56*(2), 736–743.

https://doi.org/10.1016/j.neuroimage.2010.04.267

Carpenter, A. C., Thakral, P. P., Preston, A. R., & Schacter, D. L. (2021). Reinstatement of item-specific

contextual details during retrieval supports recombination-related false memories. *NeuroImage*,

236, 118033. https://doi.org/10.1016/j.neuroimage.2021.118033

Chadwick, M. J., Anjum, R. S., Kumaran, D., Schacter, D. L., Spiers, H. J., & Hassabis, D. (2016). Semantic

representations in the temporal pole predict false memories. *Proceedings of the National*

Academy of Sciences, *113*(36), 10180–10185. https://doi.org/10.1073/pnas.1610686113

Chamberlain, J. D., Bowman, C. R., & Dennis, N. A. (2022). Age-related differences in encoding-retrieval

similarity and their relationship to false memory. *Neurobiology of Aging*, *113*, 15–27.

https://doi.org/10.1016/j.neurobiolaging.2022.01.011

Cox, R. W., & Hyde, J. S. (1997). Software tools for analysis and visualization of fMRI data. *NMR in*

Biomedicine, *10*(4–5), 171–178. https://doi.org/10.1002/(SICI)1099-

1492(199706/08)10:4/5<171::AID-NBM453>3.0.CO;2-L

Criss, A. H., & Shiffrin, R. M. (2004). Pairs do not suffer interference from other types of pairs or single

items in associative recognition. *Memory & Cognition*, *32*(8), 1284–1297.

https://doi.org/10.3758/BF03206319

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical Surface-Based Analysis. I. Segmentation and

surface reconstruction. *NeuroImage*, *9*(2), 179–194.

Delhaye, E., Salmon, E., & Bastin, C. (2014). *Exploration of unitization processes in episodic memory in*

Alzheimer's disease. In Front. Hum. Neurosci. Conference Abstract: Belgian Brain Council 2014

MODULATING THE BRAIN: FACTS, FICTION, FUTURE.

Dennis, N. A., Bowman, C. R., & Peterson, K. M. (2014). Age-related differences in the neural correlates

mediating false recollection. *Neurobiology of Aging*, *35*(2), 395–407.

https://doi.org/10.1016/j.neurobiolaging.2013.08.019

Dennis, N. A., Johnson, C. E., & Peterson, K. M. (2014). Neural correlates underlying true and false

associative memories. *Brain and Cognition*, *88*, 65–72.

https://doi.org/10.1016/j.bandc.2014.04.009

Dennis, N. A., Overman, A. A., Carpenter, C. M., & Gerver, C. R. (2022). Understanding associative false

memories in aging using multivariate analyses. *Aging, Neuropsychology, and Cognition*, *29*(3),

500–525. https://doi.org/10.1080/13825585.2022.2037500

Dennis, N. A., Overman, A. A., Gerver, C. R., McGraw, K. E., Rowley, M. A., & Salerno, J. M. (2019).

Different types of associative encoding evoke differential processing in both younger and older

adults: Evidence from univariate and multivariate analyses. *Neuropsychologia*, *135*, 107240.

https://doi.org/10.1016/j.neuropsychologia.2019.107240

Diana, R. A., Yonelinas, A. P., & Ranganath, C. (2008). The effects of unitization on familiarity-based

source memory: Testing a behavioral prediction derived from neuroimaging data. *Journal of*

Experimental Psychology: Learning, Memory, and Cognition, *34*(4), 730–740.

https://doi.org/10.1037/0278-7393.34.4.730

Elbich, D. B., Webb, C. E., & Dennis, N. A. (2021). The influence of item familiarization on neural

discriminability during associative memory encoding and retrieval. *Brain and Cognition*, *152*,

105760. https://doi.org/10.1016/j.bandc.2021.105760

Favila, S. E., Chanales, A. J. H., & Kuhl, B. A. (2016). Experience-dependent hippocampal pattern

differentiation prevents interference during subsequent learning. *Nature Communications*, *7*(1),

11066. https://doi.org/10.1038/ncomms11066

Folstein, M. F., Robins, L. N., & Helzer, J. E. (1983). The mini-mental state examination. *Archives of*

General Psychiatry, *40*(7), 812–812.

Fonov, V. S., Evans, A. C., McKinstry, R. C., Almli, C. R., & Collins. (2009). Unbiased nonlinear average age-

appropriate brain templates from birth to adulthood. *NeuroImage*, *47*, S102.

Ford, J. H., Verfaellie, M., & Giovanello, K. S. (2010). Neural correlates of familiarity-based associative

retrieval. *Neuropsychologia*, *48*(10), 3019–3025.

https://doi.org/10.1016/j.neuropsychologia.2010.06.010

Giovanello, K. S., Keane, M. M., & Verfaellie, M. (2006). The contribution of familiarity to associative

memory in amnesia. *Neuropsychologia*, *44*(10), 1859–1865.

https://doi.org/10.1016/j.neuropsychologia.2006.03.004

Graf, P., & Schacter, D. L. (1989). Unitization and Grouping Mediate Dissociations in Memory for New

Associations. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *15*(5), 930.

Greve, D. N., & Fischl, B. (2009). Accurate and robust brain image alignment using boundary-based

registration. *NeuroImage*, *48*(1), 63–72. https://doi.org/10.1016/j.neuroimage.2009.06.060

Haskins, A. L., Yonelinas, A. P., Quamme, J. R., & Ranganath, C. (2008). Perirhinal Cortex Supports

Encoding and Familiarity-Based Recognition of Novel Associations. *Neuron*, *59*(4), 554–560.

https://doi.org/10.1016/j.neuron.2008.07.035

Haxby, J., Gobbini, M., Furey, M., Ishai, A., Schouten, J., & Pietrini, P. (2001). Distributed and Overlapping

Representations of Faces and Objects in Ventral Temporal Cortex. *Science (New York, N.Y.)*, *293*,

2425–2430. https://doi.org/10.1126/science.1063736

Haxby, J. V., Gobbini, M. I., Furey, M. L., Ishai, A., Schouten, J. L., & Pietrini, P. (2001). Distributed and

overlapping representations of faces and objects in ventral temporal cortex. *Science*, *293*(5539),

2425–2430.

Huffer, V., Bader, R., & Mecklinger, A. (2022). Can the elderly take the action? – The influence of

unitization induced by action relationships on the associative memory deficit. *Neurobiology of*

Learning and Memory, *194*, 107655. https://doi.org/10.1016/j.nlm.2022.107655

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved Optimization for the Robust and

Accurate Linear Registration and Motion Correction of Brain Images. *NeuroImage*, *17*(2), 825–

841. https://doi.org/10.1006/nimg.2002.1132

Johnson, M. K., Kuhl, B. A., Mitchell, K. J., Ankudowich, E., & Durbin, K. A. (2015). Age-related differences

in the neural basis of the subjective vividness of memories: Evidence from multivoxel pattern

classification. *Cognitive, Affective, & Behavioral Neuroscience*, *15*(3), 644–661.

https://doi.org/10.3758/s13415-015-0352-9

Kafkas, A., Migo, E. M., Morris, R. G., Kopelman, M. D., Montaldi, D., & Mayes, A. R. (2017). Material

Specificity Drives Medial Temporal Lobe Familiarity But Not Hippocampal Recollection:

FAMILIARITY-BASED RECOGNITION IN THE MTL. *Hippocampus*, *27*(2), 194–209.

https://doi.org/10.1002/hipo.22683

Kamp, S.-M., Bader, R., & Mecklinger, A. (2018). Unitization of word pairs in young and older adults:

Encoding mechanisms and retrieval outcomes. *Psychology and Aging*, *33*(3), 497–511.

https://doi.org/10.1037/pag0000256

Klein, A., Ghosh, S. S., Bao, F. S., Giard, J., Häme, Y., Stavsky, E., Lee, N., Rossa, B., Reuter, M., Chaibub

Neto, E., & Keshavan, A. (2017). Mindboggling morphometry of human brains. *PLOS*

Computational Biology, *13*(2), e1005350. https://doi.org/10.1371/journal.pcbi.1005350

Kondo, Y., Suzuki, M., Mugikura, S., Abe, N., Takahashi, S., Iijima, T., & Fujii, T. (2005). Changes in brain

activation associated with use of a memory strategy: A functional MRI study. *NeuroImage*, *24*(4),

1154–1163. https://doi.org/10.1016/j.neuroimage.2004.10.033

Kriegeskorte, N. (2008). Representational similarity analysis – connecting the branches of systems

neuroscience. *Frontiers in Systems Neuroscience*. https://doi.org/10.3389/neuro.06.004.2008

Kriegeskorte, N., & Kievit, R. A. (2013). Representational geometry: Integrating cognition, computation,

and the brain. *Trends in Cognitive Sciences*, *17*(8), 401–412.

https://doi.org/10.1016/j.tics.2013.06.007

Kroes, M. C. W., & Fernández, G. (2012). Dynamic neural systems enable adaptive, flexible memories.

Neuroscience & Biobehavioral Reviews, *36*(7), 1646–1666.

https://doi.org/10.1016/j.neubiorev.2012.02.014

Lanczos, C. (1964). *Evaluation of noisy data*. *1*(1), 76–85.

LaRocque, K. F., Smith, M. E., Carr, V. A., Witthoft, N., Grill-Spector, K., & Wagner, A. D. (2013). Global

Similarity and Pattern Separation in the Human Medial Temporal Lobe Predict Subsequent

Memory. *Journal of Neuroscience*, *33*(13), 5466–5474.

https://doi.org/10.1523/JNEUROSCI.4293-12.2013

Liu, Z., Li, X., Li, X., Yuan, J., & Guo, C. (2022). From encoding to retrieval: Change in level of unitization

resolves debate about Unitization's effect on associative recognition. *Psychophysiology*, *e14161*.

Memel, M., & Ryan, L. (2017). Visual integration enhances associative memory equally for young and

older adults without reducing hippocampal encoding activation. *Neuropsychologia*, *100*, 195–

206. https://doi.org/10.1016/j.neuropsychologia.2017.04.031

Mumford, J. A., Turner, B. O., Ashby, F. G., & Poldrack, R. A. (2012). Deconvolving BOLD activation in

event-related designs for multivoxel pattern classification analyses. *NeuroImage*, *59*(3), 2636–

2643. https://doi.org/10.1016/j.neuroimage.2011.08.076

Naveh-Benjamin, M. (2000). Adult age differences in memory performance: Tests of an associative

deficit hypothesis. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *26*(5),

1170–1187. https://doi.org/10.1037/0278-7393.26.5.1170

Naveh-Benjamin, M., Brav, T. K., & Levy, O. (2007). The associative memory deficit of older adults: The

role of strategy utilization. *Psychology and Aging*, *22*(1), 202.

Naveh-Benjamin, M., Guez, J., Kilb, A., & Reedy, S. (2004). The Associative Memory Deficit of Older

Adults: Further Support Using Face-Name Associations. *Psychology and Aging*, *19*(3), 541–546.

https://doi.org/10.1037/0882-7974.19.3.541

Old, S. R., & Naveh-Benjamin, M. (2008). Differential effects of age on item and associative measures of

memory: A meta-analysis. *Psychology and Aging*, *23*(1), 104–118. https://doi.org/10.1037/0882-

7974.23.1.104

Oosterhof, N. N., Connolly, A. C., & Haxby, J. V. (2016). CoSMoMVPA: Multi-Modal Multivariate Pattern

Analysis of Neuroimaging Data in Matlab/GNU Octave. *Frontiers in Neuroinformatics*, *10*, 27.

https://doi.org/10.3389/fninf.2016.00027

Overman, A. A., & Stephens, J. D. W. (2013). Synergistic effects of encoding strategy and context

salience on associative memory in older adults. *Psychology and Aging*, *28*(3), 654–665.

https://doi.org/10.1037/a0031441

Park, D. C., Polk, T. A., Park, R., Minear, M., Savage, A., & Smith, M. R. (2004). Aging reduces neural

specialization in ventral visual cortex. *Proceedings of the National Academy of Sciences*, *101*(35),

13091–13095. https://doi.org/10.1073/pnas.0405148101

Park, H., Leal, F., Spann, C., & Abellanoza, C. (2013). The effect of object processing in content-

dependent source memory. *BMC Neuroscience*, *14*(1), 71. https://doi.org/10.1186/1471-2202-

14-71

Parks, C. M., & Yonelinas, A. P. (2015). The importance of unitization for familiarity-based learning.

Journal of Experimental Psychology: Learning, Memory, and Cognition, *41*(3), 881–903.

https://doi.org/10.1037/xlm0000068

Parra, M. A., Abrahams, S., Logie, R. H., & Sala, S. D. (2009). Age and binding within-dimension features

in visual short-term memory. *Neuroscience Letters*, *449*(1), 1–5.

https://doi.org/10.1016/j.neulet.2008.10.069

Pidgeon, L. M., Grealy, M., Duffy, A. H. B., Hay, L., McTeague, C., Vuletic, T., Coyle, D., & Gilbert, S. J.

(2016). Functional neuroimaging of visual creativity: A systematic review and meta-analysis.

Brain and Behavior, *6*(10), e00540. https://doi.org/10.1002/brb3.540

Piekema, C., Rijpkema, M., Fernández, G., & Kessels, R. P. C. (2010). Dissociating the Neural Correlates of

Intra-Item and Inter-Item Working-Memory Binding. *PLoS ONE*, *5*(4), e10214.

https://doi.org/10.1371/journal.pone.0010214

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2014). Methods

to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage*, *84*, 320–

341. https://doi.org/10.1016/j.neuroimage.2013.08.048

Price, A. R., Peelle, J. E., Bonner, M. F., Grossman, M., & Hamilton, R. H. (2016). Causal Evidence for a

Mechanism of Semantic Integration in the Angular Gyrus as Revealed by High-Definition

Transcranial Direct Current Stimulation. *The Journal of Neuroscience*, *36*(13), 3829–3838.

https://doi.org/10.1523/JNEUROSCI.3120-15.2016

Quamme, J. R., Yonelinas, A. P., & Norman, K. A. (2007). Effect of unitization on associative recognition

in amnesia. *Hippocampus*, *17*(3), 192–200.

Ricupero, S., Carpenter, C. M., Steinkrauss, A. C., Gerver, C. R., Chamberlain, J. D., Monkman, R. G.,

Overman, A. A., & Dennis, N. A. (2023). Neural distinctiveness and reinstatement of

hippocampal representations support unitization for associations. *Brain Research*, *1798*, 148143.

https://doi.org/10.1016/j.brainres.2022.148143

Satterthwaite, T. D., Elliott, M. A., Gerraty, R. T., Ruparel, K., Loughead, J., Calkins, M. E., Eickhoff, S. B.,

Hakonarson, H., Gur, R. C., Gur, R. E., & Wolf, D. H. (2013). An improved framework for

confound regression and filtering for control of motion artifact in the preprocessing of resting-

state functional connectivity data. *NeuroImage*, *64*, 240–256.

https://doi.org/10.1016/j.neuroimage.2012.08.052

Shimamura, A. P. (2011). Episodic retrieval and the cortical binding of relational activity. *Cognitive,*

Affective, & Behavioral Neuroscience, *11*(3), 277–291. https://doi.org/10.3758/s13415-011-

0031-4

Simmonite, M., & Polk, T. A. (2022). Age-related declines in neural distinctiveness correlate across brain

areas and result from both decreased reliability and increased confusability. *Aging,*

Neuropsychology, and Cognition, *29*(3), 483–499.

https://doi.org/10.1080/13825585.2021.1999383

Sommer, V. R., & Sander, M. C. (2022). Contributions of representational distinctiveness and stability to

memory performance and age differences. *Aging, Neuropsychology, and Cognition*, *29*(3), 443–

462. https://doi.org/10.1080/13825585.2021.2019184

Staresina, B. P., & Davachi, L. (2010). Object Unitization and Associative Memory Formation Are

Supported by Distinct Brain Regions. *Journal of Neuroscience*, *30*(29), 9890–9897.

https://doi.org/10.1523/JNEUROSCI.0826-10.2010

Stephens, J. D. W., & Overman, A. A. (2018). Modeling age differences in effects of pair repetition and

proactive interference using a single parameter. *Psychology and Aging*, *33*(1), 182–194.

https://doi.org/10.1037/pag0000195

Stokes, M., Thompson, R., Cusack, R., & Duncan, J. (2009). Top-Down Activation of Shape-Specific

Population Codes in Visual Cortex during Mental Imagery. *Journal of Neuroscience*, *29*(5), 1565–

1572. https://doi.org/10.1523/JNEUROSCI.4657-08.2009

Thakral, P. P., Wang, T. H., & Rugg, M. D. (2017). Decoding the content of recollection within the core

recollection network and beyond. *Cortex*, *91*, 101–113.

https://doi.org/10.1016/j.cortex.2016.12.011

Tse, D., Takeuchi, T., Kakeyama, M., Kajii, Y., Okuno, H., Tohyama, C., Bito, H., & Morris, R. G. M. (2011).

Schema-dependent gene activation and memory encoding in neocortex. *Science*, *333*(6044),

891–895.

Tu, H.-W., & Diana, R. A. (2021). The interaction of relational encoding and unitization: Effects on medial

temporal lobe processing during retrieval. *Behavioural Brain Research*, *396*, 112878.

https://doi.org/10.1016/j.bbr.2020.112878

Tustison, N. J., Avants, B. B., Cook, P. A., Zheng, Y., Egan, A., Yushkevich, P. A., & Gee. (2010). N4ITK:

improved N3 bias correction. *IEEE Transactions on Medical Imaging*, *29*(6), 1310–1320.

Voss, J. L., Baym, C. L., & Paller, K. A. (2008). Accurate forced-choice recognition without awareness of

memory retrieval. *Learning & Memory*, *15*(6), 454–459. https://doi.org/10.1101/lm.971208

Wagner, I. C., van Buuren, M., Kroes, M. C., Gutteling, T. P., van der Linden, M., Morris, R. G., &

Fernández, G. (2015). Schematic memory components converge within angular gyrus during

retrieval. *ELife*, *4*, e09668. https://doi.org/10.7554/eLife.09668

Yesavage. (1998). *Geriatric depression scale.* (4th ed., Vol. 24). Psychopharmacol Bull.

Yonelinas, A. P. (1997). Recognition memory ROCs for item and associative information: The

contribution of recollection and familiarity. *Memory & Cognition*, *25*(6), 747–763.

https://doi.org/10.3758/BF03211318

Yonelinas, A. P. (2002). The Nature of Recollection and Familiarity: A Review of 30 Years of Research.

Journal of Memory and Language, 46(3), 441–517. https://doi.org/10.1006/jmla.2002.2864

Yonelinas, A. P., Hopfinger, J. B., Buonocore, M. H., Kroll, N. E. A., & Baynes, K. (2001). Hippocampal,

parahippocampal and occipital-temporal contributions to associative and item recognition

memory: An fMRI study: *Neuroreport, 12*(2), 359–363. https://doi.org/10.1097/00001756-

200102120-00035

Yonelinas, A. P., Kroll, N. E. A., Dobbins, I. G., & Soltani, M. (1999). Recognition memory for faces: When

familiarity supports associative recognition judgments. *Psychonomic Bulletin & Review, 6*(4),

654–661. https://doi.org/10.3758/BF03212975

Yovel, G., & Paller, K. A. (2004). The neural basis of the butcher-on-the-bus phenomenon: When a face

seems familiar but is not remembered. *NeuroImage, 21*(2), 789–800.

https://doi.org/10.1016/j.neuroimage.2003.09.034

Zhang, Y., Brady, M., & Smith, S. (2001). Segmentation of brain MR images through a hidden Markov

random field model and the expectation-maximization algorithm. *IEEE Transactions on Medical*

Imaging, 20(1), 45–57.

Zheng, Z., Li, J., Xiao, F., Broster, L. S., & Jiang, Y. (2015). Electrophysiological evidence for the effects of

unitization on associative recognition memory in older adults. *Neurobiology of Learning and*

Memory, 121, 59–71. https://doi.org/10.1016/j.nlm.2015.03.006

Zheng, Z., Li, J., Xiao, F., Ren, W., & He, R. (2016). Unitization improves source memory in older adults:

An event-related potential study. *Neuropsychologia, 89*, 232–244.

https://doi.org/10.1016/j.neuropsychologia.2016.06.025

Figure legends

Figure 1. Regions of interest for multivariate classification. Regions defined from AAL pickatlas. Tan = MOC, Dark Red = AG, Blue = MFG, Green = HC, Light Red = PrC. Slice numbers: y = -6; z = 36; x = 32.

Figure 2. Based on Overman and Stephens 2013, Example stimuli for the unitized and associative memory conditions. Face and job pairings were presented on one of two background colors (yellow or blue). Both the encoding conditions contained blue and yellow trials. At retrieval, the pairings were presented on a white background color. (Note: smile faces are added to replace actual faces for biorxiv, actual images will be reported in the paper)

Figure 3. Based on Haxby et al. 2001, calculation of neural distinctiveness. Calculations were repeated twice; once for encoding trials and once for retrieval trials. Distinctiveness was calculated by taking the within-condition similarity correlations for all trials within a condition subtracted by the between-condition similarity correlation for all trials between the condition. (Note: smile faces are added to replace actual faces for biorxiv, actual images will be reported in the paper)

Figure 4. Within-similarity between the two encoding conditions in the AG, HC, MFG, MOC, and PrC. Neural within-similarity scores significantly different between doing versus speaking conditions indicated in a black box; p's < 0.05.

ROI	Encoding		Retrieval	
	Mean	Standard Deviation	Mean	Standard Deviation
AG	0.540	0.104	0.517	0.053
HC	0.479	0.109	0.493	0.064
MFG	0.509	0.084	0.513	0.044
MOC	0.526	0.098	0.508	0.069
PrC	0.502	0.072	0.505	0.058

Table 1. Means and standard deviations of each ROI for classification analyses during encoding and retrieval.

Supplemental Figure

Supplemental Figure 1. Correlations of neural distinctiveness scores between each AG, HC, MFG, MOC, and PrC during encoding. * indicates regions significantly correlated to each other (p's $< .05$)