1

CEP signaling coordinates plant immunity with nitrogen status

Jakub Rzemieniewski[1], Henriette Leicher[1], Hyun Kyung Lee[2], Caroline Broyart[2], Shahran Nayem[3], Christian Wiese[1,†], Julian Maroschek[1], Zeynep Camgöz[1], Vilde Olsson Lalun[4], Michael Anthony Djordjevic[5], A. Corina Vlot[3,‡], Ralph Hückelhoven[1], Julia Santiago[2] and Martin Stegmann[1*]

[1]Phytopathology, TUM School of Life Sciences, Technical University of Munich, Freising, Germany

[2]The Plant Signaling Mechanisms Laboratory, Department of Plant Molecular Biology, University of Lausanne, Lausanne, Switzerland

[3]Helmholtz Zentrum Munich, Institute of Biochemical Plant Pathology, Neuherberg, Germany

[4]Section for Genetics and Evolutionary Biology, Department of Biosciences, University of Oslo, Oslo, Norway

[5]Division of Plant Science, Australian National University, Acton, Canberra, Australia

†present address: Biotechnology of Natural Products, TUM School of Life Sciences, Technical University of Munich, Freising, Germany

‡present address: Chair of Crop Plant Genetics, Faculty of Life Sciences: Food, Nutrition and Health, University of Bayreuth, Kulmbach, Germany

*Corresponding author: Email: martin.stegmann@tum.de

Abstract

Plant endogenous signaling peptides shape growth, development and adaptations to biotic and abiotic stress. Here, we identified C-TERMINALLY ENCODED PEPTIDEs (CEPs) as novel immune-modulatory peptides (phytocytokines) in *Arabidopsis thaliana*. Our data reveals that CEPs induce immune outputs and are required to mount resistance against the leaf-infecting bacterial pathogen *Pseudomonas syringae* pv. *tomato*. We show that effective immunity requires CEP perception by tissue-specific CEP RECEPTOR 1 (CEPR1) and CEPR2. Moreover, we identified the related RECEPTOR-LIKE KINASE 7 (RLK7) as a novel CEP4-specific CEP receptor contributing to CEP-mediated immunity, suggesting a complex interplay of multiple CEP ligands and receptors in different tissues during biotic stress. CEPs have a known role in the regulation of root growth and systemic nitrogen (N)-demand signaling. We now provide evidence that CEPs and their receptors promote immunity in an N status-dependent manner, suggesting a previously unknown molecular crosstalk between plant nutrition and cell surface immunity. We propose that CEPs and their receptors are central regulators for the adaptation of biotic stress responses to plant-available resources.

Main Text

Receptor kinases (RKs) sense external and internal cues to control multiple aspects of plant physiology, ranging from growth and development to plant immunity and abiotic stress tolerance. RKs can serve as pattern recognition receptors (PRRs) to detect microbe-associated molecular patterns (MAMPs) and activate pattern-triggered immunity (PTI). An example is the *Arabidopsis thaliana* (hereafter Arabidopsis) leucine-rich repeat RK (LRR-RK) FLAGELLIN SENSITIVE 2 (FLS2), which forms a receptor complex with BRASSINOSTEROID INSENSITIVE 1-ASSOCIATED RK 1 (BAK1) upon perception of a 22 amino acid epitope derived from bacterial flagellin (flg22) to activate PTI (*1–4*). Plants also perceive endogenous peptides to regulate multiple aspects of plant physiology (*5*). Importantly, plants utilize specific endogenous peptides for controlling immunity. These peptides are referred to as phytocytokines and their expression or secretion can be modulated upon PTI activation. They also often additionally regulate aspects of plant growth and development (*6, 7*). Examples are GOLVEN2 (GLV2) peptides which are perceived by ROOT MERISTEM GROWTH FACTOR INSENSITIVE 3 (RGI3) to modulate PRR stability and RAPID ALKALINIZATION FACTORs (RALFs) that are sensed by the malectin RK (MLRK) FERONIA (FER) to control PRR nanoscale dynamics at the plasma membrane and MAMP-induced PRR-BAK1 complexes for PTI initiation (*8–11*). GLV2 also controls hypocotyl gravicurvature (*12*) and RALF perception by FER and other MLRKs affects several aspects of plant growth, development and reproduction, suggesting that endogenous peptides coordinate these processes with stress responses (*13–18*).

Immune-modulatory peptides are often transcriptionally upregulated in response to MAMP perception, including SERINE-RICH ENDOGENOUS PEPTIDES (SCOOPs) and SMALL PHYTOCYTOKINES REGULATING DEFENSE AND WATER LOSS (SCREWs)/CTNIPs (*19–21*). Yet, *GLV2* transcription is not induced by biotic stress (*8*) and promotes immunity, suggesting that immunity-dependent transcriptional regulation is not a prerequisite for phytocytokine function. Phytocytokines and other endogenous peptides further regulate a multitude of abiotic stress responses, including adaptation to high salinity, drought and nutrient deprivation, indicating that they can integrate multiple external and internal cues to safeguard plant health (*22–24*). Yet, how different peptide-mediated pathways are coordinated remains largely unknown.

Here, we identified C-TERMINALLY ENCODED PEPTIDES (CEPs) as novel phytocytokines in

Arabidopsis. CEPs are important for sucrose-dependent lateral root growth, root system architecture,

systemic nitrogen (N)-demand signaling and promotion of root nodulation, but a function in plant

immunity remained unknown (*22, 25–31*). We show that the unusual class I CEP peptide CEP4 induces

immune responses. We found that *CEP4* and other *CEPs* are expressed in shoots and perceived by

canonical CEP receptors CEPR1 and CEPR2 to mount effective cell surface immunity. *CEPR1* and

CEPR2 show tissue-specific expression patterns, suggesting CEP sensing in distinct tissues spatially

cooperates to control plant immunity. Yet, CEP4-induced responses also require the CEPR-related

RECEPTOR-LIKE KINASE 7 (RLK7), which we identified as a novel CEP4-specific CEP receptor with

wide-spread expression in leaves. Importantly, we now show that a short-term reduction in seedling N

levels promotes flg22-induced PTI responses in a CEP and CEP receptor-dependent manner,

suggesting that CEPs coordinate a previously unknown cross-talk between cell surface immunity and

plant nutrition.

Results

CEPs are novel phytocytokines

We sought to identify novel phytocytokines regulating growth and immunity in Arabidopsis and

screened publicly available transcription data of known growth-regulatory plant peptide families for

members with differential expression after elicitor treatment. With this approach, we recently identified

GLV2 as a novel phytocytokine modulating PRR stability through RGIs (*8*). We noticed that a specific

member of the CEP family, *CEP4*, showed differential expression upon flg22 treatment with a moderate

downregulation in an *asr3* mutant background, a transcriptional repressor of flg22-induced genes

(Supplementary Fig. 1A) (*32*). Using RT-qPCR, we also observed a mild flg22-induced *CEP4*

downregulation in Col-0 seedlings compared to the mock control. At 4 hours of mock treatment, *CEP4*

expression was lower compared to 0-hour mock samples, suggesting some degree of circadian rhythm-

related regulation (Supplementary Fig. 1B). The majority of *CEPs* show predominant expression in root

tissue (*33*). We compared root and shoot *CEP4* transcripts in seedlings upon flg22 treatment. We

detected much stronger *CEP4* expression in roots with a similar pattern of flg22-dependent modulation.

Interestingly, *CEP4* expression in shoots was mildly upregulated upon 1 h of flg22 treatment

(Supplementary Fig. 1C). Arabidopsis encodes for 12 class I CEPs and 3 class II CEPs, which can be

distinguished by sequence differences in their peptide domain. CEPs are produced from larger peptide

precursors that carry one to five predicted mature CEP domain in their sequence and an N-terminal

signal peptide for secretion (*33*). CEP4 is classified as a class I CEP but has an unusual structure

compared to other class I CEPs (Supplementary Fig. 1D). CEP4 only carries two proline residues in its

peptide domain, unlike several characterized typical class I CEPs such as CEP1 and CEP3 (*33*).

To test whether CEP4 may be involved in immunity, we generated constitutive overexpression lines

using a full-length CEP4 precursor sequence (*35S::CEP4*, Supplementary Fig. 2A) and noticed that

these lines showed increased resistance to infection by *Pseudomonas syringae* pv. *tomato (Pto)*

lacking the effector molecule coronatine (*Pto*COR-), which is routinely used to assess PTI-associated

disease resistance phenotypes (Fig. 1A) (*34, 35*). The same lines were also more susceptible to

infection by the wild-type *Pto* DC3000 strain (Supplementary Fig. 2B).

The majority of mature CEPs previously identified are 15mer peptides with an N-terminal aspartate (D),

a C-terminal Histidine (H) and hydroxylated prolines (*22, 33*). We synthesized a 16mer peptide with

both proline residues hydroxylated and the N- and C-terminal D and H residue, respectively,

DAFRHypTHQGHypSQGIGH, to test whether it triggered or modulated immune responses. CEP4

application activated dose-dependent PTI outputs, including the cellular influx of calcium ions in a Col-

0 line expressing the calcium reporter Aequorin (Col-0AEQ) (*36*), the activation of MITOGEN-

ACTIVATED PROTEIN KINASEs (MAPKs), ethylene production and expression of the PTI marker

gene *FLAGELLIN-INDUCED RECEPTOR KINASE 1* (*FRK1*) in Col-0 seedlings (Fig. 1B-E,

Supplementary Fig. 3A). CEP4-induced calcium influx was detectable in the low nanomolar range of

CEP4 concentration (Supplementary Fig. 3A) and MAPK phosphorylation was detected at

concentrations of 100 nM, yet the magnitude of response was weaker compared to flg22

(Supplementary Fig. 3B). *FRK1* transcript accumulation and calcium influx activated by flg22 treatment

was much stronger compared to CEP4 in whole seedlings (Supplementary Fig. 3C, D). However, CEP4

induced *FRK1* expression in a similar range as previously described elicitors, suggesting biological

relevance (*37*). Moreover, CEP4 triggered nuclear YFP fluorescence in the vasculature of *pFRK1::NLS-*

3xmVenus seedlings, suggesting some degree of tissue specificity of CEP4-induced immune outputs

(*38*) (Fig. 1F). Finally, CEP4 treatment resulted in seedling growth inhibition (SGI) and systemic

resistance to *Pto* DC3000 infection (Fig. 1G, H). We also tested a 15mer peptide lacking the N-terminal

D residue (AFRHypTHQGHypSQGIGH), which triggered a dose-dependent calcium influx with no

significant differences to the response elicited by the 16mer peptide (Supplementary Fig. 3A). We next

tested whether other class I CEPs can activate PTI responses. Indeed, CEP1 and one peptide derived

from CEP9 (CEP9.5), which carries five CEP domains in its precursor sequence (*39*), were able to

trigger calcium influx in seedlings but only at higher concentrations of 10 µM (Supplementary Fig. 3E).

However, CEP1 induced nuclear YFP fluorescence in the vasculature of *pFRK1::NLS-3xmVenus* lines

at similar concentrations as CEP4 (Supplementary Fig. 3F), suggesting that likely several CEPs can

trigger immune responses with CEP4 being a very potent family member.

To confirm the role of CEPs in plant immunity and because we anticipated genetic redundancy of

immune-regulatory CEPs, we generated a *cep6x* mutant by CRISPR-Cas9 in which *CEP4*, as well as

the five additional class I CEPs *CEP1-CEP3*, *CEP6* and *CEP9* were mutated to predictable loss of

function (CRISPR alleles *cep1.1*, *cep2.1*, *cep3.1*, *cep4.1*, *cep6.1* and *cep9.1*, Supplementary Fig. 2C).

The resulting *cep6x* mutant had no obvious morphological defects (Supplementary Fig. 2D) but showed

compromised resistance to Pto^{COR-} infection, confirming that CEPs are important for antibacterial

resistance (Fig. 1I). To overcome the impact of different tissue-specific CEPs that cooperate to mount

disease resistance, we complemented the Pto^{COR-} hypersusceptibility phenotype of *cep6x* using a full-

length *CEP4* driven by the constitutive 35S promoter (Fig. 1J, Supplementary Fig. 2E). The *cep6x*

mutant was also more susceptible to infection with the fully virulent wild-type *Pto* DC3000 strain, and

35S::CEP4 expression partially complemented this phenotype (Supplementary Fig. 4A). We also tested

a *cep5x* mutant (CRISPR alleles *cep1.2*, *cep2.2*, *cep3.2*, *cep6.2* and *cep9.2*, *CEP4* wild-type) which

showed an intermediate phenotype upon Pto^{COR-} infection (Fig. 4B, Supplementary Fig. 2C, D). This

indicates the contribution of multiple CEPs to mount robust resistance.

CEPR1 and CEPR2 are CEP4 receptors and central regulators of plant immunity

CEPs bind to CEPR1 and CEPR2, two LRR-RLKs from LRR subfamily XI (*22*, *40*). Genetically, CEPR1

is predominantly required for class I CEP perception during root growth-related responses (*25–28*, *41*,

42). We tested whether *CEPR1* and *CEPR2* are also involved in bacterial immunity. We did not observe

a Pto^{COR-} infection phenotype in *cepr1-3* and *cepr2-4* single mutants (Supplementary Fig. 5A, Fig. 2A).

We generated a *cepr1-3 cepr2-4 (cepr1-3/2-4)* mutant by genetic crossing and this double mutant was

more susceptible to Pto^{COR-} (Fig. 2A). We also generated a new *cepr1 cepr2* double mutant by CRISPR-

Cas9 in a Col-0^{AEQ} background (CRISPR alleles *cepr1.1*^{AEQ}, *cepr2.1*^{AEQ}, hereafter *cepr1/2*^{AEQ})

(Supplementary Fig. 5B). Similar to *cepr1-3/2-4*, *cepr1/2*[AEQ] was more susceptible to *Pto*[COR-] infection

(Fig. 2A), further confirming a role of *CEPR1* and *CEPR2* in antibacterial resistance. These data

suggest that *CEPR1* and *CEPR2* may control immunity redundantly.

We then tested whether CEP4 perception depends on CEPR1 and/or CEPR2. Using SGI as a readout,

we noticed that *cepr1-3* was insensitive and two *cepr2* alleles (*cepr2-3* and *cepr2-4*) were insensitive

and less sensitive to CEP4 treatment, respectively (Supplementary Fig. 5C). The *cepr1-3/2-4* mutant

and the previously reported No-0 *cepr1-1 cepr2-1* double mutant also did not respond to CEP4 in SGI

experiments (Fig. 2B, Supplementary Fig. 5D). Similarly, the *cepr1-3/2-4* double mutant was largely

insensitive to CEP4-induced systemic resistance (Fig. 2C). Of note, mock-treated *cepr1-3/2-4* mutants

do not show enhanced bacterial growth compared to Col-0, suggesting that CEPR1/CEPR2 primarily

regulate immunity to tissue invasion, an effect that is bypassed by syringe infiltration in this experimental

setup. Collectively, these data suggest that both CEPR1 and CEPR2 are involved in CEP4 perception.

We then tested whether CEP4 can directly bind to the ectodomain (ECD) of CEPR1 and/or CEPR2.

We expressed CEPR1[ECD] and CEPR2[ECD] in *Trichoplusia ni* Tnao38 cells and purified them for

quantitative binding experiments. We analyzed protein quality by Coomassie stain and size exclusion

chromatography (SEC, Supplementary Fig. 6A, B). Unfortunately, CEPR1[ECD] aggregated, as indicated

by the early elution of the bulk sample during SEC analysis (~10min, Supplementary Fig. 6A).

Nevertheless, we obtained good quality protein for CEPR2[ECD] with a single SEC elution peak at ~13min,

which we subsequently tested for quantitative binding to CEP4 using isothermal titration calorimetry

(ITC). CEP4, but not a scrambled control (CEP4[scr]), directly bound to CEPR2[ECD] with a K_D of 15.7 µM

(± 4.5 µM) (Fig. 2D-E, G, Supplementary Fig. 6C). CEP1, which was previously shown to bind to CEPR1

and CEPR2 (*22*) also bound to CEPR2[ECD] with a K_D of 9.3 µM (± 0.6 µM) (Fig. 2F-G, Supplementary

Fig. 6C). These data are in range with previously reported peptide-LRR-RK binding affinities obtained

by ITC and suggest that CEPR2 is a bona fide CEP4 receptor (*43–45*).

We were then interested to characterize a possible role of CEPR1 and CEPR2 for FLS2-mediated

signaling. We did not observe strong differences in flg22-induced ethylene accumulation *in cepr1-3/2-*

4 (Supplementary Fig. 7A). Of note, flg22-induced ethylene production was higher in *cepr1-3/2-4*, but

basal ethylene production was enhanced in this mutant background, making the result difficult to

interpret. This is interesting in light of a previous report showing that the *Medicago truncatula* CEP1-

CRA2 (CEPR1 orthologue) pathway negatively regulates ethylene signaling during root nodule

symbiosis (*46*). Interestingly though, flg22-induced *FRK1* and *PR1* expression were reduced in adult

cepr1-3 or *cepr2-4* plants, which was pronounced in *cepr1-3/2-4* (Fig. 2H). Moreover, *cepr1-3/2-4*

showed compromised flg22-induced resistance to *Pto* DC3000 infection (Fig. 2I). The flg22-induced

seedling growth inhibition was unaffected in *cepr1-3/2-4* (Supplementary Fig. 7B), suggesting that

CEPR1 and CEPR2 are selectively required for specific flg22-induced outputs associated with

antibacterial defense. We next tested whether CEP-CEPR1/2 signaling may be important for systemic

acquired resistance (SAR). To induce SAR, we used a *Pto* strain producing the effector avrRPM1 (*Pto*

avrRPM1), which is recognized by Col-0 RESISTANCE TO PSEUDOMONAS SYRINGAE 1, to activate

effector-triggered immunity and consequently SAR (*47*, *48*). Local inoculation of *Pto avrRPM1* and

subsequent infection of systemic tissue with virulent *Pto* revealed that *cepr1-3/2-4*, as well as *cep6x*,

were strongly compromised in *Pto* avrRPM1-triggered SAR (Fig. 2J). Collectively, these data show that

CEP-CEPR signaling is a central regulator of PTI and SAR in Arabidopsis.

RLK7 is a novel CEP4-specific CEP receptor

To further characterize the role of CEPR1 and CEPR2 in CEP4-induced signaling, we tested early

CEP4-triggered responses in *cepr1-3/2-4* and *cepr1/2*[AEQ]. To our surprise, we found that *cepr1/2*[AEQ]

did not show compromised calcium influx upon CEP4 treatment (Fig. 3A). CEP4-induced MAPK

activation was also unaffected in *cepr1-3/2-4* (Fig. 3B). Moreover, *cepr1-3* and *cepr2-4* were largely

unaffected in CEP4-induced *FRK1* expression and the response was also not abolished in *cepr1-3/2-*

4 (Supplementary Fig. 8A). These results suggested that other receptor(s) may be involved in CEP4

perception. CEPR1 and CEPR2 are phylogenetically close to IKU2, which is involved in seed size

regulation, and RLK7, which plays a role in controlling germination speed, lateral root formation and

salt stress adaptation (*24*, *40*, *49–51*). RLK7 also senses endogenous PAMP-INDUCED PEPTIDES

(PIPs) to regulate PTI and resistance to the fungal wilt pathogen *Fusarium oxysporum* and *Pto* (*52*, *53*).

We tested whether *iku2-4*, *rlk7-1* and *rlk7-3* are compromised for CEP4 perception. The *iku2-4* mutant

showed unaltered CEP4-induced ethylene accumulation, but this response was compromised in *rlk7-*

1 and *rlk7-3* (Supplementary Fig. 8B). Similarly, *rlk7-1* and *rlk7-3* showed strongly reduced CEP4-

induced MAPK activation (Fig. 3C). We also generated an *rlk7/iku2*[AEQ] mutant by CRISPR-Cas9 in a

Col-0[AEQ] background (CRISPR alleles *rlk7.1*[AEQ] and *iku2.1*[AEQ], Supplementary Fig. 8C). The

rlk7/iku2[AEQ] line showed compromised CEP4-induced calcium influx (Supplementary Fig. 8D). Two

additional *rlk7*[AEQ] single mutants generated by CRISPR-Cas9 (CRISPR alleles *rlk7.2*[AEQ], *rlk7.3*[AEQ]

Supplementary Fig. 8C) were also compromised in CEP4-triggered calcium influx (Fig. 3D). Yet,

residual CEP4 activity remained in *rlk7* mutants, both for CEP4-induced MAPK activation and calcium

influx (Fig. 3C, D). To resolve this, we generated a CRISPR *cepr1 cepr2 rlk7*[AEQ] (hereafter

cepr1/2/rlk7[AEQ]) triple mutant using the *cepr1/2*[AEQ] background (CRISPR allele *rlk7.4*[AEQ] in *cepr1/2*[AEQ],

Supplementary Fig. 5B, 8C), which showed abolishment of CEP4-induced calcium influx and MAPK

activation (Fig. 3E, F). This suggests that RLK7, CEPR1 and CEPR2 each participate in mounting a

full CEP4 response, with RLK7 playing a predominant genetic role. The *cepr1/2/rlk7*[AEQ] mutant also

showed abolished CEP4-induced resistance to *Pto* DC3000 infection (Fig. 3G). Both *rlk7*[AEQ] and

cepr1/2/rlk7[AEQ] were also insensitive to PIP1 in MAPK activation (Supplementary Fig. 8E), consistent

with RLK7's function as a PIP receptor (*52*). Interestingly, *rlk7-1* and *rlk7-3* were only moderately

affected in CEP4-induced SGI (Supplementary Fig. 8F), suggesting that certain CEP4 responses

require selective specificity for one of the three CEP receptors. Moreover, CEP1 and CEP9.5-induced

calcium influx was abolished in *cepr1/2*[AEQ] and unaltered *in rlk7/iku2*[AEQ] or *rlk7.2*[AEQ], respectively,

indicating that CEPR1/2 are required for the early responses triggered by these canonical class I CEPs,

which again show differential receptor requirements (Fig. 3H, Supplementary Fig. 8G).

We next used ITC to test whether CEP4 can bind to the ectodomain of RLK7 (RLK7[ECD]). Similar to

CEPR2[ECD], we obtained good quality proteins for RLK7[ECD] eluting with a main single peak at ~13min

during SEC analysis (Supplementary Fig. 6A, B). CEP4 bound to RLK7[ECD] with a K_D of 9 μM (± 4.9

μM) (Fig. 3I, L, Supplementary Fig. 6C), similar to CEPR2[ECD] (Fig. 2D, G). We also tested binding of

PIP1 to RLK7[ECD], a described RLK7 ligand (*24, 52*). RLK7[ECD] bound PIP1 with a higher affinity (K_D 500

nM ± 60 nM) (Fig. 3J, L, Supplementary Fig. 6C). Importantly, consistent with unaltered CEP1-induced

responses in *rlk7/iku2*[AEQ] (Fig. 3H), CEP1 did not bind to RLK7 (Fig. 3K, Supplementary Fig. 6C). These

data suggest that RLK7 can function as a CEP4-specific CEP receptor.

The *cepr1/2*[AEQ] mutants show compromised resistance to *Pto*[COR-] (Fig. 2A), similar to previously

published *rlk7* single mutants (*53*). Further mutation of *rlk7* in *cepr1/2*[AEQ] background did not

significantly enhance this phenotype (Supplementary Fig. 8H). When using the fully virulent *Pto*

DC3000 strain, *cepr1/2*[AEQ] was moderately more susceptible (Fig. 3M), whereas *rlk7* was not (*53*).

Interestingly, the *cepr1/2/rlk7*[AEQ] triple mutant showed significantly increased susceptibility to *Pto*

DC3000 compared to *cepr1/2*[AEQ] (Fig. 3M), indicating that all three CEP receptors mount full

antibacterial resistance.

CEP-CEP receptor signaling promotes local immunity against *Pto*

Class I *CEPs* show predominant expression in root tissue (*33*) (Supplementary Fig. 1C) and function

as systemic root-to-shoot transmitters of N starvation (*22*). We hypothesized that root-expressed *CEPs*

may regulate leaf immunity upon infection with *Pto*. Therefore, we performed reciprocal grafts between

Col-0 and *cep6x* and tested whether root or shoot expression of *CEPs* is required for immune

regulation. Surprisingly, we found that *CEP* mutation in the shoot conferred increased *Pto*COR-

susceptibility in *cep6x* (Fig. 4A). We used the hypovirulent *Pto*COR- strain in this experiment, as the

cep6x phenotype is stronger with this bacterial strain (Fig. 1I, Supplementary Fig. 4A). Consistently, we

detected weak *CEP1-CEP4*, *CEP6* and *CEP9* expression in leaf tissue with *CEP4* being slightly

upregulated upon *Pto* DC3000 infection (Fig. 4B), similar to the mild *CEP4* upregulation upon flg22

treatment in seedling shoots (Supplementary Fig. 1C). We similarly performed reciprocal grafting with

Col-0AEQ and *cepr1/2/rlk7*AEQ mutants, which demonstrated that shoot expressed CEP receptors are

required to confer enhanced *Pto* DC3000 resistance (Fig. 4C). *Pto* DC3000 was used because the

phenotype of *cepr1/2/rlk7*AEQ was more pronounced with this bacterial strain (Fig. 3M). These data

suggest that CEP function in the shoot is necessary for their immune-modulatory function, unlike the

root-to-shoot CEP mobility required for N-demand signaling (*22*).

Next we wanted to investigate spatial expression patterns of *CEP4* by generating a *pCEP4::NLS-*

3xmVenus line. Consistent with previous reports, *CEP4* did not show expression in the main root, but

in emerging lateral roots (Fig. 4D) (*33*). Despite weak shoot signals for *CEP4* expression obtained by

qPCR (Supplementary Fig. 1C), we found widespread expression of *CEP4* in seedling leaf tissue, but

not in the vasculature or stomatal guard cells (Fig. 4D). This data further supports a role for leaf-

expressed CEP4 in local responses.

We were next interested in resolving the spatial expression pattern of CEPR1, CEPR2 and RLK7 in

shoot tissue. Previous reports using promoter::β-GLUCURONIDASE lines suggested restricted *CEPR1*

expression to the vasculature and more widespread *CEPR2* promoter activity (*22*, *54*). Consistently, a

pCEPR1::NLS-3xmVenus revealed that *CEPR1* expression was specific to vasculature tissue, while

pCEPR2::NLS-3xmVenus signals were largely restricted to stomatal guard cells (Fig. 4E). The

pRLK7::Venus-H2B line showed widespread promoter activity, including the vasculature, mesophyll

and epidermal cells (Fig. 4E). *RLK7* showed a large overlap with *CEP4* expression in leaves,

suggesting that CEP4 and RLK7 can meet *in vivo* to function as a receptor-ligand pair. Although our

results point to CEPR1/CEPR2 and RLK7 being required for full CEP4 sensitivity (Fig. 3E, F), only

RLK7 shows an overlapping expression pattern with *CEP4*. Moreover, *CEP4* expression was absent in

the vasculature (Fig. 4D) but CEP4 induced *FRK1* expression in this tissue (Fig. 1F, Supplementary

Fig. 3F). This raises the possibility for CEP4 mobility between tissue layers and that the peptide may

exert its function in a combination of cell-autonomous and short-to-long distance signaling.

The restriction of expression of *CEPR1* and *CEPR2* in the vasculature and guard cells, respectively,

may also explain the minor contribution of these receptors to mount CEP4-induced early responses

upon elicitation in whole seedlings (Fig. 3A-F). For this reason, we generated *CEPR2-GFP* and *CEPR1-*

GFP overexpression lines to test whether the constitutive expression of these receptors can promote

CEP4-induced responses. Two *35S::CEPR2-GFP* lines overexpressed the receptor ~100-150 fold

compared to Col-0 and protein accumulation was detected by western blots (Supplementary Fig. 9A,

B). We only obtained one *35S::CEPR1-GFP* line with a ~10 fold overexpression relative to Col-0

(Supplementary Fig. 9C). However, we failed to detect CEPR1-GFP accumulation in this line,

suggesting that the protein may be unstable. Consistent with a function as a CEP4 receptor (Fig. 2D,

G), the overexpression of CEPR2-GFP enhanced the responsiveness of seedlings to CEP4 in MAPK

activation experiments (Fig. 4F). As expected from the lack of detectable protein, *CEPR1* transcript

overexpression did not alter CEP4-induced MAPK activation (Supplementary Figure 9D). However,

CEPR1 overexpression mildly promoted CEP3-induced MAPK activation (Supplementary Fig. 9D), in

line with previous reports of CEPR1 being the primary receptor for canonical class I CEPs (*22*, *27*, *30*,

31, *42*). This suggests that CEPR2 is a physiologically relevant CEP4 receptor.

CEP signaling promotes FLS2 signaling under reduced nitrogen supply

Unlike other immune-promoting phytocytokines, such as PIPs and SCOOPs, *CEP4* is transcriptionally

downregulated upon flg22 perception in whole seedling and roots (Supplementary Fig. 1A-C), and only

mildly upregulated upon flg22 treatment in seedling shoots or upon *Pto* infection in leaf tissue

(Supplementary Fig. 1C, Fig. 4B). We were thus interested to determine the biological relevance of

CEP-mediated control of cell surface immunity. First, we tested whether CEP treatment affects FLS2-

dependent signaling. CEP4 application could promote flg22-induced ethylene accumulation and

resistance induced by a low dose of flg22 (100 nM) (Fig. 5A, B). Yet, we did not observe a noticeable

defect in flg22-induced ethylene accumulation in *cep6x* and *cepr1/2/rlk7*[AEQ] (Supplementary Fig. 10A,

B). CEP4 did not induce ethylene production in *cepr1/2/rlk7*[AEQ], similar to abolished CEP4-triggered

calcium influx and MAPK activation in this mutant background (Supplementary Fig. 10B, Fig. 3E, F).

Unlike *cepr1-3/2-4* (Supplementary Fig. 7A), the triple receptor mutant did not show elevated basal

ethylene levels, raising the question whether RLK7 promotes ethylene accumulation in the absence of

CEPR1/CEPR2 (Supplementary Fig. 10B). Also, flg22-induced MAPK activation and/or *FRK1*

expression were unaltered in *cep6x* and *cepr1/2/rlk7*[AEQ] (Supplementary Fig. 10C-E).

As modulators of systemic N-demand signaling, several *CEPs* are transcriptionally upregulated in N-

starved roots, including *CEP1* (*22*, *55*, *56*). It is known that the N status of plants affects disease

resistance to different pathogens, but the underlying molecular mechanisms remain unknown (*57*, *58*).

We noticed that *CEP4* is transcriptionally upregulated upon placing seedlings in ½ MS containing 10%

N concentrations compared to the control medium (100% N ½ MS medium, 20 mM NO_3^-, 10 mM NH_4^+,

Fig. 5C). Importantly, N starvation promoted resistance to *Pto* (*59*), suggesting a connection between

N homeostasis and antibacterial resistance. We hypothesized that CEPs coordinate the plant's N status

with cell surface immunity. We transferred two-week-old seedlings for 24 h to ½ MS medium with

different N concentrations before challenging them with flg22. Ten percent N and 5% N medium

promoted flg22-induced MAPK activation, while 1% N did not (Supplementary Fig. 10F). This suggests

that different N concentrations modulate the ability to mount FLS2 signaling. The promotion of flg22-

induced MAPK activation upon 10% N treatment was compromised in *cep6x*, suggesting that CEPs

promote FLS2 signaling under reduced N conditions (Fig. 5D, E). We confirmed this phenotype by

measuring *FRK1* expression upon 10% N treatment and subsequent flg22 elicitation in wild-type and

cep6x (Fig. 5F). We next tested whether CEP receptors are required for the N-dependent regulation of

FLS2 signaling. Indeed, 10% N-promoted flg22-induced MAPK activation was reduced in

cepr1/2/rlk7[AEQ] (Fig. 5G, H).

We raised the question whether the impairment in FLS2 signaling under 10% N conditions in *cep6x*

and *cepr1/2/rlk7*[AEQ] mutants might be associated with defects in N homeostasis. Indeed, the No-0

cepr1/2 mutant has paler leaves, smaller rosette size and constitutive anthocyanin accumulation,

related to defects in nitrate uptake (*22*) but the *cep6x* shoot is indistinguishable from the wild-type plant

(Supplementary Fig. 2D). We grew *cep6x* and *cepr1/2/rlk7*[AEQ] seedlings in 100% N, 10% N and 1% N

medium. We noticed a mild reduction in seedling growth of all genotypes in 10% N medium, which was

more pronounced in 1% N medium after seven days (Supplementary Fig. 10G, H). The *cep6x* mutant

phenotype was indistinguishable from the WT (Supplementary Fig. 10G). The *cepr1/2/rlk7*[AEQ] seedlings

also showed similar growth at lower N concentrations compared to its WT control (Supplementary Fig.

10H). This suggests that 10% N-induced promotion of flg22-triggered responses is not a mere

consequence of deregulated metabolism in *cep6x* and *cepr1/2/rlk7*[AEQ]. Strongly reduced nitrate

concentrations enhance the expression of the high-affinity nitrate transporter *NRT2.1* (*60–62*). We

could not detect enhanced *NRT2.1* expression in WT under 10% N conditions, which remains higher

than nitrate concentrations previously tested for *NRT2.1* expression (2 mM nitrate vs 1 mM) (Fig. 5I)

(*61*). However, flg22 promoted *NRT2.1* transcript accumulation which was abrogated in *cep6x* (Fig. 5I).

Altogether, these results indicate a CEP-and CEP receptor-dependent connection between FLS2-

triggered PTI and the plant´s N supply, revealing a previously unknown mechanism of signaling cross-

talk between cell surface immunity and the plant`s nutritional status.

Discussion

This work shows that CEP perception in Arabidopsis is achieved by three partially cell type and tissue-

specific receptors that regulate plant immunity and likely coordinate biotic stress with plant nutritional

cues. CEP4 binds to both CEPR2 and RLK7, and CEP4 outputs show variable grades of dependency

on specific receptors. The canonical group I CEP1 and CEP9.5, however, exclusively depend on

CEPR1/2 and CEP1 does not bind to RLK7. The overlap of *CEP4* with *RLK7* expression in leaves, but

not with *CEPR1* and *CEPR2*, suggests a combination of cell autonomous and short-distance signaling

contributing to CEP-mediated immune modulation. Our work reveals an unexpected and previously

undescribed complexity of phytocytokine signaling. A challenge for the future will be to determine how

three CEP receptors with distinct expression patterns integrate responses between tissues and the

concerted action of multiple ligands. RLK7 also recognizes PIPs to regulate growth, salt stress and

immunity (*50, 52*) (Fig. 3J). RLK7 binds PIP1 with higher affinity and likely preferentially recognizes this

ligand. Members of the CLE peptide family also bind multiple receptors with variable affinities to

regulate epidermal cell patterning (*44, 63*). Dissecting spatiotemporal ligand availability and CEP4-PIP1

signaling specificity will be an important future task. RLK7 binds CEP4, but not CEP1, suggesting that

CEP4`s receptor specificities are unique among CEPs, likely caused by its distinct sequence

(Supplementary Fig. 1D). It will be interesting to compare the molecular mechanisms of PIP1/CEP4-

RLK7 and CEP1/CEP4-CEPR2/CEPR1 recognition and activation in future structure analyses. Plant

peptides may be versatile tools for the evolution of plant plasticity. High numbers of diverse ligands and

receptors may provide the plant with vast combinations for fine-tuned signal outputs during adaption to

environmental challenges.

Our data suggest that short-term reduction in N supplementation promotes *CEP* expression to promote

FLS2 activation. Other phytocytokines modulate immune signaling by regulating PRR abundance or

receptor complex formation and dynamics, but in most cases the mechanism remains unknown (*7–9,*

11). It will be interesting to reveal whether CEPs directly or indirectly modulate PRR signaling.

Accumulating evidence suggests a direct integration of nutrient homeostasis and PTI in plants.

Perception of flg22 by FLS2 induces PHT1.4 phosphorylation to inhibit phosphate (Pi) uptake and

promote root immunity (*64*). It remains unknown whether immune activation also regulates N transport

in root or shoot tissue. CEPs induce nitrate, Pi and sulfate uptake, suggesting CEP-

CEPR1/CEPR2/RLK7-dependent modulation of several transporter pathways (*65*). This raises the

question whether nutrient uptake directly contributes to CEP-mediated immune modulation. The main

source of inorganic N for plant utilization is nitrate, which also functions as a signaling molecule to

induce adaptive growth responses (*66*). Nitrate is sensed by the plasma membrane transceptor NRT1.1

and the nuclear transcriptional regulator NLP7 (*67, 68*). NRT1.1 and similar transporters are regulated

by phosphorylation to control transport activity, including NRT1.2 phosphorylation by CEPR2 (*67, 69–*

73), which we identified as a CEP4 receptor. It will be interesting to resolve whether and how cell

surface signaling and nitrogen sensing/transport directly or indirectly intersect. Since seedlings grown

under high N concentrations (as provided by ½ MS medium) show reduced flg22 responsiveness, it is

also possible that N saturation inhibits PTI by suppressing *CEP* expression and accumulation.

CEPR1/CEPR2/RLK7 and CEPs are widely conserved among angiosperms, including crop plants (*40*).

CEPs are important to promote nodulation in legumes and also regulate sucrose-dependent root growth

inhibition and fecundity (*28, 41, 74, 75*). This places CEPs as central integrators of biotic interactions

(symbiosis and pathogen defense) with plant nutrition, growth and development (*65, 76*). It will be

critical to understand whether N-dependent and CEP-mediated PTI modulation extends beyond

Arabidopsis, and to decipher how CEPs can promote immunity and control symbiosis in diverse

species. This will provide important insights for future crop improvement strategies that coordinate crop

nutrition with disease resistance.

Materials and methods

<u>Molecular cloning</u>

To generate *CEP4* overexpression lines, the coding sequence of *CEP4* (AT2G35612) was synthesized (Twist Bioscience, USA) with attB attachment sites for subsequent gateway cloning into pDONRZeo (Invitrogen, USA) and recombination with pB7WG2 (VIB, Ghent). To generate CRISPR-Cas9 mutants, appropriate target sites (two per gene of interest) were designed using the software tool chopchop (https://chopchop.cbu.uib.no/). Individual guide RNA constructs containing gene-specific target sites were synthesized (Twist Bioscience, USA) and subsequently stacked in a GoldenGate-adapted pUC18-based vector. To generate different order CRISPR *cep* mutants, *cepr1/2*[AEQ], *rlk7/iku2*[AEQ] and *rlk7*[AEQ] 12, 4, 4 and 2 target site-containing gRNA constructs were stacked, respectively (Supplementary Table 2). Together with FastRed-pRPS5::Cas9, higher-order gRNA stacks were subsequently cloned into pICSL4723 for in planta expression (*77*).

To generate the *pCEP4::NLS-3xmVenus*, *pCEPR1::NLS-3xmVenus* and *pCEPR2::NLS-3xmVenus* reporter constructs, 1000, 1696 and 2788 bp fragments upstream of the start codon, respectively, were amplified from genomic DNA and assembled together with the sequence coding for the nuclear localization signal of SV40 large T antigen followed by 3 consecutive mVenus YFP fluorophores (*78*) into a GoldenGate-modified pCB302 binary vector for plant expression. For *pRLK7* the 1957 bp promoter sequence upstream from the start codon was amplified with primers containing attB attachment sites for subsequent gateway cloning into pDONRZeo (Invitrogen, USA) and recombination with promotor::Venus (YFP)-H2B destination vector (*78*). For *CEPR1* (AT5G49660) and *CEPR2* (AT1G72180) overexpression lines, the coding sequence of both genes was amplified from cDNA with attB attachment sites for subsequent cloning into a pDONR223 (Invitrogen, USA) and recombination with pK7FWG2 (VIB Ghent, Belgium). All of the generated plant expression constructs were subsequently transformed into *Agrobacterium tumefaciens* strain GV3101 before floral dip transformation of Arabidopsis. All primers used for cloning are listed in (Supplementary Table 3).

<u>Plant material and growth conditions</u>

Arabidopsis Col-0, Col-0[AEQ] (*36*) and No-0 were used as wild types for experiments and generation of transgenic lines or CRISPR mutants. The *cepr1-3* (GK-467C01), *cepr2-3* (SALK_014533), *rlk7-1*

(SALK_056583), *rlk7-3* (SALK_120595), *iku2-4* (Salk_073260) and the novel *cepr2-4* allele (GK-

695D11) were obtained from NASC (UK) (*28*, *52*, *79*, *80*). The No-0 *cepr2-1xcepr2-1* was obtained from

RIKEN (Japan) (*22*). T-DNA insertion mutants were genotyped by PCR using T-DNA- and gene-specific

primers as listed in Supplementary Table 3. The lack of *CEPR2* transcript in *cepr2-4* (GK-695D11) was

determined by semi-quantitative PCR from cDNA using *cepr2-4* genotyping primers (Supplementary

Fig. 5A). The *cepr1-3/2-4* double mutant was obtained by genetic crossing. The *bak1-5/bkk1* mutant

was characterized previously (*81*). For visualizing tissue-specific *FRK1* expression, a *pFRK1::NLS-

3xmVenus* line was used (*82*). To isolate homozygous CRISPR mutants, pICSL4723 transformant T1

seeds showing red fluorescence were selected and grown on soil before genotyping with gene-specific

primers and Sanger sequencing (Supplementary Table 3). Mutants lacking the transgene were

identified by loss of fluorescence. To generate the *cep6x 35S::CEP4* lines, the same pB7WG2 CEP4

construct used for the generation of *CEP4* overexpression lines was utilized for floral dip transformation

of homozygous *cep6x*.

Plants for physiological assays involving mature plants were vernalized for 2-3 days in the dark at 4°C

and later grown in individual pots in environmentally controlled growth rooms (20-21°C, 55% relative

humidity, 8 h photoperiod). For seedling-based assays, seeds were sterilized using chlorine gas and

grown axenically on ½ Murashige and Skoog (MS) media supplemented with vitamins (Duchefa,

Netherlands), 1% sucrose, with or without 0.8% agarose at 22°C and a 16 h photoperiod unless stated

otherwise. For experiments using ½ MS medium with reduced N concentrations, modified MS salts

without nitrogen-containing compounds (Duchefa, Netherlands) were used and supplemented with

KNO_3/NH_4NO_3 to achieve 100% N (KNO_3 9.395 mM, NH_4NO_3 10.305 mM), 10% N (KNO_3 0.9395 mM,

NH_4NO_3 1.0305 mM), 5% (KNO_3 0.4698 mM, NH_4NO_3 0.5153 mM) and 1% (KNO_3 0.09395 mM,

NH_4NO_3 0.10305 mM) conditions. To keep the ionic strength equal in 10%, 5% and 1% N conditions,

media were supplemented with 90% (8.455 mM), 95% (8.925 mM) and 99% (9.301 mM) KCl,

respectively.

Grafting

Arabidopsis seedlings were grown vertically on ½ MS agar medium without sucrose in short-day

conditions seven days before grafting. Grafting was performed aseptically under a stereo microscope

as previously described (*83*). Vertically mounted plates with reciprocally grafted seedlings were

returned to short-day conditions for 10 days. Healthy seedlings were transferred to the soil.

<u>Imaging and microscopy</u>

Confocal laser-scanning microscopy was performed using a Leica TCS SP5 (Leica, Germany)

microscope (with Leica Application Suite X 3.7.4.23463) and all the pictures were taken with a 20x

water immersion objective. For the mVenus fluorophore, pictures were imaged with argon laser

excitation at 514nm and a detection window of 525-535 nm. Propidium iodide was visualized using

DPSS 561 laser emitting at 561 nm with a detection window of 575-590 nm. For imagining the

expression pattern of NLS-3xmVenus or Venus (YFP)-H2B under the control of different promoters

(*pCEP4*, *pCEPR1*, *pCEPR2* and *pRLK7*), vertically-grown seedlings were stained with propidium iodide

immediately before microscopic analysis. For imagining the *pFRK1::NLS-3xmVenus* reporter line, 12-

day-old seedlings were transferred to a 24-well-plate containing ddH$_2$O with or without (mock) peptides

in the indicated concentrations. For comparison of *FRK1* promoter activity, seedlings were analysed by

confocal microscopy using identical laser intensities and interval/number of slices for Z stack projection

16 h after treatment.

<u>Calcium influx assay</u>

Apoaequorin-expressing liquid-grown eight-day-old seedlings were transferred individually to a 96-well

plate containing 100 µl of 5 µM coelenterazine-h (PJK Biotech, Germany) and incubated in the dark

overnight. Luminescence was measured using a plate reader (Luminoskan Ascent 2.1, Thermo Fisher

Scientific, USA). Background luminescence was recorded by scanning each well 12 times at 10 s

intervals, before adding a 25 µl elicitor solution to the indicated final concentration. Luminescence was

recorded for 30 min at the same interval. The remaining aequorin was discharged using 2 M CaCl$_2$,

20% ethanol. The values for cytosolic Ca^{2+} concentrations ([Ca^{2+}]$_{cyt}$) were calculated as luminescence

counts per second relative to total luminescence counts remaining (L/L$_{max}$).

<u>Ethylene measurement</u>

Leaf discs (4 mm in diameter) from four-to-five-week-old soil-grown Arabidopsis were recovered

overnight in ddH$_2$O. Three leaf discs per sample were transferred to a glass vial containing 500 µl of

ddH$_2$O before adding ddH$_2$O (mock) or peptides to the indicated final concentration. Glass vials were

capped with a rubber lid and incubated under gentle agitation for 3.5 h. One mL of the vial headspace

was extracted with a syringe and injected into a Varian 3300 gas chromatograph (Varian, USA) to

measure ethylene.

<u>MAPK activation and western blot analysis</u>

Five-day old Arabidopsis seedlings growing on ½ MS agar plates, were transferred into a 24-well plate

containing liquid medium for seven days. 24 h before the experiment, seedlings were equilibrated in a

fresh ½ MS medium. For N reduction experiments, modified ½ MS containing 100% N, 10% N, 5% N

and 1 % N supplemented with KCl was used. MAPK activation was elicited by adding the peptides to

the indicated concentrations. Six seedlings per sample were harvested, frozen in liquid nitrogen and

homogenized using a tissue lyser (Qiagen, Germany). Proteins were extracted using a buffer containing

50 mM Tris-HCl (pH 7.5), 50 mM NaCl, 10% glycerol, 5 mM DTT, 1% protease inhibitor cocktail, 1 mM

phenylmethylsulfonyl fluoride (PMSF), 1% IGEPAL, 10 mM EGTA, 2 mM NaF, 2 mM Na$_3$VO$_4$, 2 mM

Na$_2$MoO$_4$, 15 mM ß-Glycerophosphate and 15 mM p-nitrophenylphosphate before analysis by SDS-

PAGE and western blot. Phosphorylated MAPKs were detected by α-p44/42 antibodies (Cell Signaling,

USA). To quantify the intensity of the specific bands, ImageJ software (version 1.53t) was used. Each

band was selected with the same-sized frame and the intensity peak was determined. The area under

each peak was calculated and normalized to Coomassie staining as a measure of relative band intensity

(RBI). The RBI of the wild-type genotypes at 100% N upon flg22 treatment was set to one.

To determine CEPR1 and CEPR2 protein levels in *35S::CEPR1-GFP* and *35S::CEPR2-GFP*

overexpression lines, seedlings were grown in ½ MS liquid medium for 12 days. Afterwards, harvested

seedlings were frozen in liquid nitrogen, homogenized using a tissue lyser (Qiagen, Germany) and the

proteins were isolated using an extraction buffer contenting 50 mM Tris-HCl (pH 7.5), 50 mM NaCl,

10% glycerol, 2 mM EDTA, 2 mM DTT, 1% protease inhibitor cocktail, 1 mM phenylmethylsulfonyl

fluoride (PMSF) and 1% IGEPAL. After SDS-PAGE and western blot, GFP-tagged proteins were

detected by α-GFP antibodies (ChromoTek).

<u>Seedling growth inhibition</u>

Arabidopsis seedlings were grown for five days on ½ MS agar plates before the transfer of individual

seedlings into each well of a 48-well plate containing liquid medium with or without elicitors in the

indicated concentration. After seven-day treatment, the fresh weight of individual seedlings was

measured.

Pathogen growth assay

Pseudomonas syringae pv. *tomato* (*Pto*) DC3000 and *Pto* lacking the effector molecular coronatine

*Pto*COR- were grown on King's B agar plates containing 50 µg/mL rifampicin and 50 µg/mL kanamycin at

28°C. After two-to-three days bacteria were resuspended in ddH_2O containing 0.04% Silwet L77 (Sigma

Aldrich, USA). The bacterial suspension was adjusted to an OD_{600} = 0.2 (10^8 cfu/mL) for *Pto*COR- or

OD_{600} = 0.02 (10^7 cfu/mL) for *Pto* DC3000 before spray inoculating four-to-five-week-old plants. For

peptide-induced resistance in local tissues, ddH_2O (mock), flg22 (100 nM) and/or CEP4 (1 µM) were

syringe-infiltrated into mature leaves. After 24 h, *Pto* DC3000 (OD_{600} = 0.0002, 10^5 cfu/mL) was syringe-

infiltrated into pre-treated leaves and incubated for three days before determining bacterial counts. For

CEP4-induced resistance in systemic tissues, CEP4 (1 or 5 µM) or ddH_2O (mock) were syringe-

infiltrated into the first two true leaves of young three-to-four-week-old Arabidopsis. After four days, *Pto*

DC3000 (OD_{600}=0.0002, 10^5 cfu/mL) was syringe-infiltrated into leaves three and four of the pre-treated

plants. Bacterial counts were determined four days after infection.

Systemic acquired resistance

SAR experiments were performed as previously described (*84*). Briefly, plants were cultivated in a

mixture of substrate (Floradur) and silica sand in a 5:1 ratio under short day (SD) conditions (10 h) in a

growth chamber at 22 °C /18 °C (day/night) with a light intensity of 100 µmol $m^{-2}s^{-1}$, and 70% relative

humidity (RH). SAR assays were performed using *Pto* DC3000 and *Pto* AvrRpm1. Bacteria were grown

on NYGA media (0.5% peptone, 0.3% yeast extract, 2% glycerol, 1.8% agar, 50 µg/mL kanamycin, 50

µg/mL Rifampicin) at 28° C before infiltration. Freshly grown *Pto* avrRpm1 was diluted in 10 mM $MgCl_2$

(to reach a final concentration of $1×10^6$ cfu/mL) and syringe-infiltrated in the first two true leaves of four-

and-a-half-week-old plants. Concurrently, 10 mM $MgCl_2$ was applied to a separate set of plants as the

mock control treatment. Three days after *Pto* avrRpm1 infiltration, plants were challenged in their 3rd

and 4th leaves with *Pto* DC3000 (1×10^5 cfu/mL). Bacterial titers were determined four days after *Pto*

DC3000 infection.

<u>Gene expression analysis</u>

For seedlings-based assays, 12-day-old liquid-grown seedlings were equilibrated in fresh medium for

24 h before treatment with the indicated peptides. For adult plants, four-to-five-week-old Arabidopsis

leaves were syringe-infiltrated with ddH$_2$O (mock), flg22 (1 µM) or *Pto* DC3000 (OD$_{600}$ = 0.001, 5×10^5

cfu/mL) and incubated for 24 h. All samples for RT-qPCR analysis were harvested at the indicated time

points, frozen in liquid nitrogen and homogenized using a tissue lyser (Qiagen, Germany). Total RNA

was isolated using TRIzol reagent (Roche, Switzerland) and purified using Direct-zol™ RNA Miniprep

Plus kit (Zymo Research, Germany). 2 µg of the total RNA was digested with DNase I and reverse

transcribed with oligo (dT)18 and Revert Aid reverse transcriptase. RT-qPCR experiments were

performed using Takyon™ Low ROX SYBR MasterMix (Eurogentec, Belgium) with the AriaMx Real-

Time PCR system (Agilent Technologies, USA). Expression levels of all tested genes were normalized

to the house-keeping gene *Ubiquitin 5* (*UBQ5*). Sequences of all primers used for RT-qPCR analysis

are found in Supplementary Table 3.

<u>Expression and purification of recombinant receptor ectodomains</u>

Spodoptera frugiperda codon-optimized synthetic genes (Invitrogen GeneArt), coding for Arabidopsis

CEPR1 (residues 23 to 592), CEPR2 (residues 32 to 620) and RLK7 (residues 29 to 608) were cloned

into a modified pFastBAC vector (Geneva Biotech) providing a 30K signal peptide (*85*), a C-terminal

TEV (tobacco etch virus protease) cleavable site and a StrepII-9xHis affinity tag. For protein expression,

Trichoplusia ni Tnao38 cells (*86*) were infected with CEPR1, CEPR2 or RLK7 virus with a multiplicity of

infection (MOI) of 3 and incubated one day at 28°C and two days at 21°C at 110 rounds per minute

(rpm). The secreted proteins were purified by Ni^{2+} (HisTrap excel, Cytiva, equilibrated in 25 mM KP$_i$ pH

7.8 and 500 mM NaCl) followed by Strep (Strep-Tactin Superflow high-capacity, IBA Lifesciences,

equilibrated in 25 mM Tris pH 8.0, 250 mM NaCl, 1 mM EDTA) affinity chromatography. All proteins

were incubated with TEV protease to remove the tags. Proteins were purified by SEC on a Superdex

200 Increase 10/300 GL column (Cytiva, USA) equilibrated in 20 mM citrate pH 5.0, 150 mM NaCl and

further concentrated using Amicon Ultra concentrators from Millipore (Merck, Germany) with a 30,000

Da molecular weight cut-off. Purity and structural integrity of the different proteins were assessed by

SDS-PAGE.

Analytical size-exclusion (SEC) chromatography

Analytical SEC experiments were performed using a Superdex 200 Increase 10/300 GL column (GE,

USA). The columns were pre-equilibrated in 20 mM citric acid pH 5, 150 mM NaCl. 150 μg of CEPR1,

CEPR2 and RLK7 were injected sequentially onto the column and eluted at 0.5 mL/min. Ultraviolet

absorbance (UV) at 280 nm was used to monitor the elution of the proteins. The peak fractions were

analyzed by SDS-PAGE followed by Coomassie blue staining.

Isothermal titration calorimetry (ITC)

Experiments were performed at 25°C on a MicroCal PEAQ-ITC (Malvern Instruments, UK) using a 200

μL standard cell and a 40 μL titration syringe. CEP1, CEP4, CEP4^{scr} and PIP1 peptides were dissolved

in the SEC buffer to match the receptor protein. A typical experiment consisted of injecting 1 μL of a

300 μM solution of the peptide into 30 μM CEPR2 or RLK7 solution in the cell at 150 s intervals. ITC

data were corrected for the heat of dilution by subtracting the mixing enthalpies for titrant solution

injections into protein-free ITC buffer. Experiments were done in duplicates and data were analyzed

using the MicroCal PEAQ-ITC Analysis Software provided by the manufacturer. The N values were

fitted to 1 in the analysis.

Synthetic peptides

The flg22 peptide was kindly provided by Dr. Justin Lee (IPB Halle). Other peptides were synthesized

by Pepmic (China) with at least 90% purity and dissolved in ddH$_2$O. Sequences of all the synthetic

peptides can be found in Supplementary Table 1.

Quantification and statistical analysis

Statistical analyses were performed using GraphPad Prism (Version 10.1.0). Sample size, p-values and

statistical methods employed are described in the respective figure legends.

Data and material availability

All data are available in the main text or the supplementary materials. All newly generated mutant lines are available upon request to M.S.

References

1. L. Gómez-Gómez, G. Felix, T. Boller, A single locus determines sensitivity to bacterial flagellin in Arabidopsis thaliana. *Plant J.* **18**, 277–84 (1999).

2. C. Zipfel, S. Robatzek, L. Navarro, E. J. Oakeley, J. D. G. Jones, G. Felix, T. Boller, Bacterial disease resistance in Arabidopsis through flagellin perception. *Nature.* **428**, 764–767 (2004).

3. A. Heese, D. R. Hann, S. Gimenez-Ibanez, A. M. E. Jones, K. He, J. Li, J. I. Schroeder, S. C. Peck, J. P. Rathjen, The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants. *Proc. Natl. Acad. Sci. USA.* **104**, 12217–22 (2007).

4. D. Chinchilla, C. Zipfel, S. Robatzek, B. Kemmerling, T. Nürnberger, J. D. G. Jones, G. Felix, T. Boller, A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. *Nature.* **448**, 497–500 (2007).

5. V. Olsson, L. Joos, S. Zhu, K. Gevaert, M. A. Butenko, I. De Smet, Look closely, the beautiful may be small: precursor-derived peptides in plants. *Annu. Rev. Plant Biol.* **70**, 1-1.34 (2019).

6. A. A. Gust, R. Pruitt, T. Nürnberger, Sensing danger: Key to activating plant immunity. *Trends Plant Sci.* **22**, 779–791 (2017).

7. J. Rzemieniewski, M. Stegmann, Regulation of pattern-triggered immunity and growth by phytocytokines. *Curr. Opin. Plant Biol.* **68**, 102230 (2022).

8. M. Stegmann, P. Zecua-Ramirez, C. Ludwig, H.-S. Lee, B. Peterson, Z. L. Nimchuk, Y. Belkhadir, R. Hückelhoven, RGI-GOLVEN signalling promotes cell surface immune receptor abundance to regulate plant immunity. *EMBO Rep.* **23**, e53281 (2022).

9. M. Stegmann, J. Monaghan, E. Smakowska-Luzan, H. Rovenich, A. Lehner, N. Holton, Y. Belkhadir, C. Zipfel, The receptor kinase FER is a RALF-regulated scaffold controlling plant immune signaling. *Science.* **355**, 287–289 (2017).

10. Y. Xiao, M. Stegmann, Z. Han, T. A. DeFalco, K. Parys, L. Xu, Y. Belkhadir, C. Zipfel, J. Chai, Mechanisms of RALF peptide perception by a heterotypic receptor complex. *Nature.* **572**, 270–274 (2019).

11. J. Gronnier, C. M. Franck, M. Stegmann, T. DeFalco, A. Abarca, M. Von Arx, K. Duenser, W.

Lin, Z. Yang, J. Kleine-Vehn, C. Ringli, C. Zipfel, Regulation of immune receptor kinases plasma

membrane nanoscale landscape by a plant peptide hormone and its receptors. *Elife.* **11**, e74162

(2022).

12.   R. Whitford, A. Fernandez, R. Tejos, A. C. Pérez, J. Kleine-Vehn, S. Vanneste, A. Drozdzecki,

J. Leitner, L. Abas, M. Aerts, K. Hoogewijs, P. Baster, R. De Groodt, Y. C. Lin, V. Storme, Y.

Van de Peer, T. Beeckman, A. Madder, B. Devreese, C. Luschnig, J. Friml, P. Hilson, GOLVEN

secretory peptides regulate auxin carrier turnover during plant gravitropic responses. *Dev. Cell.*

22, 678–685 (2012).

13.   M. R. Blackburn, M. Haruta, D. S. Moura, Twenty Years of Progress in Physiological and

Biochemical Investigation of RALF peptides. *Plant Physiol.* **182**, 1657–1666 (2020).

14.   S. Zhu, Q. Fu, F. Xu, H. Zheng, F. Yu, New paradigms in cell adaptation: decades of discoveries

on the CrRLK1L receptor kinase signalling network. *New Phytol.* **232**, 1168–1183 (2021).

15.   C. Liu, L. Shen, Y. Xiao, D. Vyshedsky, C. Peng, X. Sun, Z. Liu, L. Cheng, H. Zhang, Z. Han, J.

Chai, H. Wu, A. Y. Cheung, C. Li, Pollen PCP-B peptides unlock a stigma peptide – receptor

kinase gating mechanism for pollination. *Science.* **372**, 171–175 (2021).

16.   S. Zhong, L. Li, Z. Wang, Z. Ge, Q. Li, A. Bleckmann, J. Wang, Z. Song, Y. Shi, T. Liu, L. Li, H.

Zhou, Y. Wang, L. Zhang, H.-M. Wu, L. Lai, H. Gu, J. Dong, A. Y. Cheung, T. Dresselhaus, L.-

J. Qu, RALF peptide signaling controls the polytubey block in Arabidopsis. *Science.* **375**, 290–

296 (2022).

17.   Z. Lan, Z. Song, Z. Wang, L. Li, Y. Liu, S. Zhi, R. Wang, J. Wang, Q. Li, Antagonistic RALF

peptides control an intergeneric hybridization barrier on Brassicaceae stigmas. *Cell.* **186**, 1–15

(2023).

18.   J. Huang, L. Yang, L. Yang, X. Wu, X. Cui, L. Zhang, J. Hui, Y. Zhao, H. Yang, S. Liu, Q. Xu, M.

Pang, X. Guo, Y. Cao, Y. Chen, X. Ren, J. Lv, J. Yu, J. Ding, G. Xu, N. Wang, X. Wei, Q. Lin,

Y. Yuan, X. Zhang, C. Ma, C. Dai, P. Wang, Y. Wang, F. Cheng, W. Zeng, R. Palanivelu, X.

Zhang, A. Y. Cheung, Q. Duan, Stigma receptors control intraspecies and interspecies barriers

in Brassicaceae. *Nature.* **614**, 303–308 (2022).

19.   K. Gully, S. Pelletier, M.-C. C. Guillou, M. Ferrand, S. Aligon, I. Pokotylo, A. Perrin, E. Vergne,

M. Fagard, E. Ruelland, P. Grappin, E. Bucher, J.-P. P. Renou, S. Aubourg, The SCOOP12

peptide regulates defense response and root elongation in Arabidopsis thaliana. *J. Exp. Bot.* **70**,

1349–1365 (2019).

20.   J. Rhodes, A.-O. Roman, M. Bjornson, B. Brandt, P. Derbyshire, M. Wyler, M. Schmid, F. L. H.
Menke, J. Santiago, C. Zipfel, Perception of a conserved family of plant signalling peptides by
the receptor kinase HSL3. *Elife.* **11**, e74687 (2022).

21.   Z. Liu, S. Hou, O. Rodrigues, P. Wang, D. Luo, K. Nomura, C. Yin, H. Wang, W. Zhang, K. Zhu-
salzman, Phytocytokine signalling reopens stomata in plant immunity and water loss. *Nature.*
605, 332–339 (2022).

22.   R. Tabata, K. Sumida, T. Yoshii, K. Ohyama, H. Shinohara, Y. Matsubayashi, Perception of root-
derived peptides by shoot LRR-RKs mediates systemic N-demand signaling. *Science.* **346**, 343–
346 (2014).

23.   F. Takahashi, T. Suzuki, Y. Osakabe, S. Betsuyaku, Y. Kondo, N. Dohmae, H. Fukuda, K.
Yamaguchi-Shinozaki, K. Shinozaki, A small peptide modulates stomatal control via abscisic
acid in long-distance signalling. *Nature.* **556**, 235–238 (2018).

24.   H. Zhou, F. Xiao, Y. Zheng, G. Liu, Y. Zhuang, Z. Wang, Y. Zhang, J. He, C. Fu, H. Lin, PAMP-
INDUCED SECRETED PEPTIDE 3 modulates salt tolerance through RECEPTOR-LIKE
KINASE 7 in plants. *Plant Cell.* **34**, 927–944 (2021).

25.   K. Ohyama, M. Ogawa, Y. Matsubayashi, Identification of a biologically active, small, secreted
peptide in Arabidopsis by in silico gene screening, followed by LC-MS-based structure analysis.
Plant J. **55**, 152–160 (2008).

26.   M. Taleski, N. Imin, M. A. Djordjevic, CEP peptide hormones: Key players in orchestrating
nitrogen-demand signalling, root nodulation, and lateral root development. *J. Exp. Bot.* **69**,
1829–1836 (2018).

27.   C. Delay, K. Chapman, M. Taleski, Y. Wang, S. Tyagi, Y. Xiong, N. Imin, M. A. Djordjevic, CEP3
levels affect starvation-related growth responses of the primary root. *J. Exp. Bot.* **70**, 4763–4774
(2019).

28.   K. Chapman, M. Taleski, H. A. Ogilvie, N. Imin, M. A. Djordjevic, CEP-CEPR1 signalling inhibits
the sucrose-dependent enhancement of lateral root growth. *J. Exp. Bot.* **70**, 3955–3967 (2019).

29.   S. Roy, L. M. Müller, A rulebook for peptide control of legume – microbe endosymbioses. *Trends*
Plant Sci. **27**, 870–889 (2022).

30.   K. Chapman, M. Taleski, M. Frank, M. A. Djordjevic, CEP and cytokinin hormone signaling

intersect to promote shallow lateral root angles. *J. Exp. Bot.* **in press**, erad353 (2023).

31.   M. Taleski, K. Chapman, O. Novák, T. Schmülling, M. Frank, M. A. Djordjevic, CEP peptide and

cytokinin pathways converge on CEPD glutaredoxins to inhibit root growth. *Nat. Commun.* **13**,

1683 (2023).

32.   B. Li, S. Jiang, X. Yu, C. Cheng, S. Chen, Y. Cheng, J. S. Yuan, D. Jiang, P. He, L. Shan,

Phosphorylation of Trihelix Transcriptional Repressor ASR3 by MAP KINASE4 Negatively

Regulates Arabidopsis Immunity. *Plant Cell.* **27**, 839–856 (2015).

33.   I. Roberts, S. Smith, B. De Rybel, J. Van Den Broeke, W. Smet, S. De Cokere, M. Mispelaere,

I. De Smet, T. Beeckman, The CEP family in land plants: Evolutionary analyses, expression

studies, and role in Arabidopsis shoot development. *J. Exp. Bot.* **64**, 5371–5381 (2013).

34.   M. Melotto, W. Underwood, J. Koczan, K. Nomura, S. Y. He, Plant Stomata Function in Innate

Immunity against Bacterial Invasion. *Cell.* **126**, 969–980 (2006).

35.   V. Nekrasov, J. Li, M. Batoux, M. Roux, Z.-H. Chu, S. Lacombe, A. Rougon, P. Bittel, M. Kiss-

Papp, D. Chinchilla, H. P. van Esse, L. Jorda, B. Schwessinger, V. Nicaise, B. P. H. J. Thomma,

A. Molina, J. D. G. Jones, C. Zipfel, Control of the pattern-recognition receptor EFR by an ER

protein complex in plant immunity. *EMBO J.* **28**, 3428–38 (2009).

36.   M. R. Knight, A. K. Campbellt, S. M. Smith, A. J. Trewavas, Transgenic plant aequorin reports

the effects of touch and cold-shock and elicitors on cytoplasmic calcium. *Nature.* **314**, 524–526

(1991).

37.   L. Huang, Y. Yuan, C. Lewis, L. Dandurand, I. Zasada, L. Huang, Y. Yuan, C. Lewis, J. Kud, J.

C. Kuhl, A. Caplan, L. Dandurand, NILR1 perceives a nematode ascaroside triggering immune

signaling and resistance. *Curr. Biol.* **33**, 1–6 (2023).

38.   L. Poncini, I. Wyrsch, V. D. Tendon, T. Vorley, T. Boller, N. Geldner, J. P. Métraux, S. Lehmann,

In roots of Arabidopsis thaliana, the damage-associated molecular pattern AtPep1 is a stronger

elicitor of immune signalling than flg22 or the chitin heptamer. *PLoS One.* **12**, e0185808 (2017).

39.   C. Delay, N. Imin, M. A. Djordjevic, CEP genes regulate root and shoot development in response

to environmental cues and are specific to seed plants. *J. Exp. Bot.* **64**, 5383–5394 (2013).

40.   C. Furumizu, A. K. Krabberød, M. Hammerstad, R. M. Alling, M. Wildhagen, S. Sawa, R. B.

Aalen, The sequenced genomes of nonflowering land plants reveal the innovative evolutionary

history of peptide signaling. *Plant Cell.* **33**, 2915–2934 (2021).

41. M. Taleski, K. Chapman, N. Imin, M. A. Djordjevic, M. Groszmann, The peptide hormone
receptor CEPR1 functions in the reproductive tissue to control seed size and yield. *Plant Physiol.*
182, 620–636 (2020).

42. K. Chapman, A. Ivanovici, M. Taleski, C. J. Sturrock, J. L. P. Ng, N. A. Mohd-Radzman, F.
Frugier, M. J. Bennett, U. Mathesius, M. A. Djordjevic, CEP receptor signalling controls root
system architecture in Arabidopsis and Medicago. *New Phytol.* **226**, 1809–1821 (2020).

43. J. Rhodes, H. Yang, S. Moussu, F. Boutrot, J. Santiago, C. Zipfel, Perception of a divergent
family of phytocytokines by the Arabidopsis receptor kinase MIK2. *Nat. Commun.* **12**, 5494
(2021).

44. P. Qian, W. Song, T. Yokoo, A. Minobe, G. Wang, T. Ishida, S. Sawa, J. Chai, T. Kakimoto, The
CLE9/10 secretory peptide regulates stomatal and vascular development through distinct
receptors. *Nat. Plants.* **4**, 1071–1081 (2018).

45. S. Hou, D. Liu, S. Huang, D. Luo, Z. Liu, Q. Xiang, P. Wang, R. Mu, Z. Han, S. Chen, J. Chai,
L. Shan, P. He, The Arabidopsis MIK2 receptor elicits immunity by sensing a conserved
signature from phytocytokines and microbes. *Nat. Commun.* **12**, 5494 (2021).

46. F. Zhu, J. Deng, H. Chen, P. Liu, L. Zheng, Q. Ye, R. Li, M. Brault, J. Wen, F. Frugier, J. Dong,
T. Wang, A CEP Peptide Receptor-like Kinase Regulates Auxin Biosynthesis and Ethylene
Signaling to Coordinate Root Growth and Symbiotic Nodulation in Medicago truncatula. *Plant*
Cell. **32**, 2855–2877 (2020).

47. D. Mackey, B. F. Holt, A. Wiig, J. L. Dangl, RIN4 interacts with Pseudomonas syringae type III
effector molecules and is required for RPM1-mediated resistance in Arabidopsis. *Cell.* **108**, 743–
54 (2002).

48. H. H. Breitenbach, M. Wenig, F. Wittek, L. Jordá, A. M. Maldonado-Alconada, H. Sarioglu, T.
Colby, C. Knappe, M. Bichlmeier, E. Pabst, D. Mackey, J. E. Parker, A. Corina Vlot, Contrasting
roles of the apoplastic aspartyl protease APOPLASTIC, ENHANCED DISEASE
SUSCEPTIBILITY1-DEPENDENT1 and LEGUME LECTIN-LIKE PROTEIN1 in Arabidopsis
systemic acquired resistance. *Plant Physiol.* **165**, 791–809 (2014).

49. M. Luo, E. S. Dennis, F. Berger, W. J. Peacock, A. Chaudhury, MINISEED3 (MINI3), a WRKY
family gene, and HAIKU2 (IKU2), a leucine-rich repeat (LRR) KINASE gene, are regulators of
seed size in Arabidopsis. *Proc. Natl Acad. Sci. USA.* **102**, 17531–17536 (2005).

50. K. Toyokura, T. Goh, H. Shinohara, A. Shinoda, Y. Kondo, Y. Okamoto, T. Uehara, K. Fujimoto,
Y. Okushima, Y. Ikeyama, K. Nakajima, T. Mimura, M. Tasaka, Y. Matsubayashi, H. Fukaki,
Lateral inhibition by a peptide hormone-receptor cascade during Arabidopsis lateral root founder
cell formation. *Dev. Cell.* **48**, 64-75.e5 (2019).

51. D. Pitorre, C. Llauro, E. Jobet, J. Guilleminot, J. P. Brizard, M. Delseny, E. Lasserre, RLK7, a
leucine-rich repeat receptor-like kinase, is required for proper germination speed and tolerance
to oxidative stress in Arabidopsis thaliana. *Planta.* **232**, 1339–1353 (2010).

52. S. Hou, X. Wang, D. Chen, X. Yang, M. Wang, D. Turrà, A. Di Pietro, W. Zhang, The secreted
peptide PIP1 amplifies immunity through Receptor-Like Kinase 7. *PLoS Pathog.* **10**, e1004331
(2014).

53. S. Hou, H. Shen, H. Shao, PAMP-induced peptide 1 cooperates with salicylic acid to regulate
stomatal immunity in Arabidopsis thaliana. *Plant Signal. Behav.* **14**, 1666657 (2019).

54. Y. Wu, Q. Xun, Y. Guo, J. Zhang, K. Cheng, T. Shi, K. He, S. Hou, X. Gou, J. Li, Genome-Wide
Expression Pattern Analyses of the Arabidopsis Leucine-Rich Repeat Receptor-Like Kinases.
Mol. Plant. **9**, 289–300 (2016).

55. Y. Ohkubo, M. Tanaka, R. Tabata, M. Ogawa-Ohnishi, Y. Matsubayashi, Shoot-to-root mobile
polypeptides involved in systemic regulation of nitrogen acquisition. *Nat. Plants.* **3**, 17029
(2017).

56. R. Ota, Y. Ohkubo, Y. Matsubayashi, M. Ogawa-ohnishi, Shoot-to-root mobile CEPD-like 2
integrates shoot nitrogen status to systemically regulate nitrate uptake in Arabidopsis. *Nat.*
Commun. **11**, 641 (2020).

57. E. Hoffland, M. J. Jeger, M. L. van Beusichem, Effect of nitrogen supply rate on disease
resistance in tomato depends on the pathogen. *Plant Soil.* **218**, 239–247 (2000).

58. M. Fagard, A. Launay, G. Clément, J. Courtial, A. Dellagi, M. Farjad, A. Krapp, M. C. Soulié, C.
Masclaux-Daubresse, Nitrogen metabolism meets phytopathology. *J. Exp. Bot.* **65**, 5643–5656
(2014).

59. C. Verly, A. C. R. Djoman, M. Rigault, F. Giraud, L. Rajjou, M. E. Saint-Macary, A. Dellagi, Plant
defense stimulator mediated defense activation is affected by nitrate fertilization and
developmental stage in Arabidopsis thaliana. *Front. Plant Sci.* **11**, 583 (2020).

60. D. Zhuo, M. Okamoto, J. J. Vidmar, A. D. M. Glass, Regulation of a putative high-affinity nitrate

transporter (Nrt2;1At) in roots of Arabidopsis thaliana. *Plant J.* **17**, 563–568 (1999).

61.   T. Kiba, A. B. Feria-Bourrellier, F. Lafouge, L. Lezhneva, S. Boutet-Mercey, M. Orsel, V. Bréhaut,

A. Miller, F. Daniel-Vedele, H. Sakakibara, A. Krapp, S. Figure, S. Figure, The Arabidopsis

nitrate transporter NRT2.4 plays a double role in roots and shoots of nitrogen-starved plants.

Plant Cell. **24**, 245–258 (2012).

62.   Z. Luo, J. Wang, F. Li, Y. Lu, Z. Fang, M. Fu, K. S. K. S. Mysore, J. Wen, J. Gong, J. D. Murray,

F. Xie, The small peptide CEP1 and the NIN-like protein NLP1 regulate NRT2.1 to mediate root

nodule formation across nitrate concentrations. *Plant Cell.* **35**, 776–794 (2023).

63.   A. Roman, P. Jimenez-sandoval, S. Augustin, C. Broyart, L. A. Hothorn, J. Santiago, HSL1 and

BAM1/2 impact epidermal cell development by sensing distinct signaling peptides. *Nat.*

Commun. **13**, 876 (2022).

64.   J. Dindas, T. A. DeFalco, G. Yu, L. Zhang, P. David, M. Bjornson, M.-C. Thibaud, V. Custódio,

G. Castrillo, L. Nussaume, A. P. Macho, C. Zipfel, Direct inhibition of phosphate transport by

immune signaling in Arabidopsis. *Curr. Biol.* **32**, 488-495.e5 (2022).

65.   S. Roy, M. Griffiths, I. Torres-Jerez, B. Sanchez, E. Antonelli, D. Jain, N. Krom, S. Zhang, L. M.

York, W. R. Scheible, M. Udvardi, Application of Synthetic Peptide CEP1 Increases Nutrient

Uptake Rates Along Plant Roots. *Front. Plant Sci.* **12**, 793145 (2022).

66.   E. A. Vidal, J. M. Alvarez, V. Araus, E. Riveras, M. D. Brooks, G. Krouk, S. Ruffel, L. Lejay, N.

M. Crawford, G. M. Coruzzi, R. A. Gutiérrez, Nitrate in 2020: Thirty years from transport to

signaling networks. *Plant Cell.* **32**, 2094–2119 (2020).

67.   C. H. Ho, S. H. Lin, H. C. Hu, Y. F. Tsay, CHL1 Functions as a Nitrate Sensor in Plants. *Cell.*

138, 1184–1194 (2009).

68.   K. Liu, M. Liu, Z. Lin, Z. Wang, B. Chen, C. Liu, A. Guo, M. Konishi, S. Yanagisawa, G. Wagner,

J. Sheen, NIN-like protein 7 transcription factor is a plant nitrate sensor. *Science.* **377**, 1419–

1425 (2022).

69.   A. Jacquot, V. Chaput, A. Mauries, Z. Li, P. Tillard, C. Fizames, P. Bonillo, F. Bellegarde, E.

Laugier, V. Santoni, S. Hem, A. Martin, A. Gojon, W. Schulze, L. Lejay, NRT2.1 C-terminus

phosphorylation prevents root high affinity nitrate uptake activity in Arabidopsis thaliana. *New*

Phytol. **228**, 1038–1054 (2020).

70.   Y. Ohkubo, K. Kuwata, Y. Matsubayashi, A type 2C protein phosphatase activates high-affinity

806   nitrate uptake by dephosphorylating NRT2.1. *Nat. Plants.* **7**, 310–316 (2021).

807 71. X. Zou, M. Y. Liu, W. H. Wu, Y. Wang, Phosphorylation at Ser28 stabilizes the Arabidopsis

808   nitrate transporter NRT2.1 in response to nitrate limitation. *J. Integr. Plant Biol.* **62**, 865–876

809   (2019).

810 72. W. Wang, B. Hu, A. Li, C. Chu, NRT1.1s in plants: Functions beyond nitrate transport. *J. Exp.*

811   *Bot.* **71**, 4373–4379 (2020).

812 73. L. Zhang, Z. Yu, Y. Xu, M. Yu, Y. Ren, S. Zhang, G. Yang, J. Huang, K. Yan, C. Zheng, C. Wu,

813   Regulation of the stability and ABA import activity of NRT1.2/NPF4.6 by CEPR2-mediated

814   phosphorylation in Arabidopsis. *Mol. Plant.* **14**, 633–646 (2021).

815 74. C. Laffont, A. Ivanovici, P. Gautrat, M. Brault, M. A. Djordjevic, F. Frugier, The NIN transcription

816   factor coordinates CEP and CLE signaling peptides that regulate nodulation antagonistically.

817   *Nat. Commun.* **11**, 3167 (2020).

818 75. A. Ivanovici, C. Laffont, E. Larrainzar, N. Patel, C. S. Winning, H. C. Lee, N. Imin, F. Frugier, M.

819   A. Djordjevic, The Medicago SymCEP7 hormone increases nodule number via shoots without

820   compromising lateral root number. *Plant Physiol.* **191**, 2012–2026 (2023).

821 76. M. Taleski, M. Jin, K. Chapman, K. Taylor, C. Winning, M. Frank, N. Imin, M. A. Djordjevic, CEP

822   hormones at the nexus of nutrient acquisition and allocation, root development, and plant-

823   microbe interactions. *J. Exp. Bot.* **in press**, erad444 (2023).

824 77. B. Castel, L. Tomlinson, F. Locci, Y. Yang, J. D. G. Jones, Optimization of T-DNA architecture

825   for Cas9-mediated mutagenesis in Arabidopsis. *PLoS One.* **14**, e0204778 (2019).

826 78. M. Somssich, A. Bleckmann, R. Simon, Shared and distinct functions of the pseudokinase

827   CORYNE (CRN) in shoot and root stem cell maintenance of Arabidopsis. *J. Exp. Bot.* **67**, 4901–

828   4915 (2016).

829 79. I. Dimitrov, F. E. Tax, Lateral root growth in Arabidopsis is controlled by short and long distance

830   signaling through the LRR RLKs XIP1/CEPR1 and CEPR2. *Plant Signal. Behav.* **13**, e1489667

831   (2018).

832 80. A. Fatihi, A. M. Zbierzak, P. Dörmann, Alterations in seed development gene expression affect

833   size and oil content of Arabidopsis seeds. *Plant Physiol.* **163**, 973–985 (2013).

834 81. M. Roux, B. Schwessinger, C. Albrecht, D. Chinchilla, A. Jones, N. Holton, F. G. Malinovsky, M.

835   Tör, S. de Vries, C. Zipfel, The Arabidopsis leucine-rich repeat receptor-like kinases

BAK1/SERK3 and BKK1/SERK4 are required for innate immunity to hemibiotrophic and biotrophic pathogens. *Plant Cell.* **23**, 2440–55 (2011).

82. F. Zhou, A. Emonet, V. Dénervaud Tendon, P. Marhavy, D. Wu, T. Lahaye, N. Geldner, Co-incidence of damage and microbial patterns controls localized immune responses in roots. *Cell.* **180**, 440–453 (2020).

83. C. W. Melnyk, Grafting with arabidopsis thaliana. *Methods Mol. Biol.* **1497**, 9–18 (2017).

84. M. Wenig, A. Ghirardo, J. H. Sales, E. S. Pabst, H. H. Breitenbach, F. Antritter, B. Weber, B. Lange, M. Lenk, R. K. Cameron, J. P. Schnitzler, A. C. Vlot, Systemic acquired resistance networks amplify airborne defense cues. *Nat. Commun.* **10**, 3813 (2019).

85. M. Futatsumori-Sugai, K. Tsumoto, Signal peptide design for improving recombinant protein secretion in the baculovirus expression vector system. *Biochem. Biophys. Res. Commun.* **391**, 931–935 (2010).

86. Y. Hashimoto, S. Zhang, S. Zhang, Y. R. Chen, G. W. Blissard, Correction: BTI-Tnao38, a new cell line derived from Trichoplusia ni, is permissive for AcMNPV infection and produces high levels of recombinant proteins. *BMC Biotechnol.* **12** (2012), doi:10.1186/1472-6750-12-12.

Author contributions

Conceptualization: MS; Investigation: JR, HL, HKL, CB, SN, CW, JM, ZC, VOL, MS; Funding

acquisition: MS, JA, RH, ACV, MAD; Project administration: MS; Supervision: MS, JR, ACV, JS, MAD,

RH; Writing – original draft: MS, JR; Writing – review & editing: HL, HKL, CB, SN, CW, JM, VOL, MAD,

ACV, RH, JS.

Competing interests

Authors declare that there are no competing interests.

Acknowledgments

We thank Stefanie Ranf for providing GoldenGate vectors for molecular cloning and Ulrich Hammes for

advice in grafting experiments. This work was funded by the following agencies: Deutsche

Forschungsgemeinschaft (DFG) grants STE2448/3-1 (MS and JR), STE2448/4-1 (MS and HL) and

SFB924 TP B06 (ACV), the Technical University of Munich (MS, JR, RH, ZC, CW and JM), the

Research Council of Norway, Grant 230849 (VOL), Australian Research Council, Grant DP200101885

(MAD), the University of Lausanne (CB), the Swiss National Science Foundation grants no.

310030_204526 and the European Research Council (ERC) grant agreement no. 716358 (JS, HKL).

Figures

Figure 1

Figure 1: CEPs induce immune responses and are important determinants of plant immunity. A)
Colony forming units (cfu) of Pto^{COR-} 3 days post inoculation (3 dpi) upon spray infection; n = 16 pooled
from four experiments ± SD (one-way ANOVA, Tukey post-hoc test; a-b, p<0.01; a/b-c p<0.0001). **B**)
Kinetics of cytosolic calcium concentrations ($[Ca^{2+}]_{cyt}$) in Col-0^{AEQ} seedlings upon mock (ddH$_2$O) or
CEP4 (1 µM) treatment; n = 8, ± SD. **C**) MAPK activation in Col-0 upon CEP4 (1 µM) or flg22 (100 nM)
treatment for the indicated time. Western blots were probed with α-p44/42. CBB = Coomassie brilliant
blue. **D**) Ethylene concentration in Col-0 leaf discs 3.5 h upon mock (ddH$_2$O) or CEP4 (1 µM) treatment;
n = 30 pooled from six experiments ± SD (two-tailed Student's t-test, **** p<0.0001). **E**) RT-qPCR of
FRK1 in seedlings upon CEP4 (1 µM) or mock (ddH$_2$O) treatment for 4 h. Housekeeping gene *UBQ5*;
n = 6 pooled from six experiments ± SD (two-tailed Student's t-test, ** p<0.01). **F**) NLS-3xmVenus signal
in p*FRK1::NLS-3xmVenus* upon mock (ddH$_2$O) or CEP4 (100 nM) treatment for 16 h. Cyan dotted line
indicates vasculature. PI = propidium iodide, scale bar = 100 µm. **G**) Relative fresh weight (as percent
of ½ MS medium control = % MS ctrl) of five-day-old seedlings treated with CEP4 (1 µM) for seven days
(n=16, ± SD, **** p<0.0001, two-tailed Student's t-test). **H**) cfu of *Pto* DC3000 (4 dpi) in distal leaves
upon local CEP4 (1 µM) or mock (ddH$_2$O) pre-treatment; n = 12 pooled from three experiments ± SD
(two-tailed Student's t-test, *** p=0.0001). **I**) cfu of Pto^{COR-} (3 dpi) upon spray infection. The *bak1-5/bkk1-*
1 mutant was used as a hypersusceptible control; n = 16 pooled from four experiments ± SD (one-way
ANOVA, Tukey post-hoc test, a-b p≤0.001). **J**) cfu of Pto^{COR-} (3dpi) upon spray infection; n = 4 ± SD
(one-way ANOVA, Tukey post-hoc test, a-b p<0.05). All experiments were repeated at least three times
in independent biological repeats with similar results.

Figure 2: CEPR1 and CEPR2 are CEP4 receptors and are important determinants of plant immunity. A) cfu of *Pto*^COR- (3 dpi) upon spray infection. The dotted line indicates different experiments; n = 12, 21 and 13 pooled from three, five and three experiments, respectively ± SD (one-way ANOVA, Dunnett's post-hoc test, a-b p<0.0001, two-tailed Student's t-test, A-B p<0.0001). **B)** Relative fresh weight of five-days-old seedlings treated with CEP4 (1 µM) for seven days; n = 35-36 pooled from three experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.0001). **C)** cfu of *Pto* DC3000 (4 dpi) in distal leaves upon local CEP4 (1 µM) or mock (ddH₂O) pre-treatment; n = 12 pooled from three experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.01). **D)** CEP4, **E)** CEP4^scr and **F)** CEP1 were titrated into a solution containing CEPR2^ECD in ITC cells. Top: raw data thermogram; bottom: fitted integrated ITC data curves. DP = differential power between reference and sample cell; ΔH = enthalpy change. **G)** ITC table summarizing CEPR2^ECD vs CEP4/CEP4^scr/CEP1 as means ± SD of two experiments. The dissociation constant (K_d) indicates receptor-ligand binding affinity. N indicates reaction stoichiometry (n = 1 for 1:1 interaction). **H)** RT-qPCR of *PR1* and *FRK1* in adult leaves after treatment with flg22 (1 µM) or mock (ddH₂O) for 24 h. Housekeeping gene *UBQ5*. Different symbols represent four independent experiments; n = 4 ± SD. **I)** cfu of *Pto* DC3000 (3 dpi) in leaves upon flg22 (1 µM) or ddH₂O pre-treatment; n = 12 pooled from three experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.0001). **J)** cfu of *Pto* DC3000 (4 dpi) in distal leaves upon local infection with *Pto avrRPM1* or 10 mM MgCl₂; n = 9 pooled from three independent experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.0001). Experiments in **A-C**, **H-J,** and **D-F** were repeated at least three times in independent biological repeats or in two independent technical repeats, respectively, with similar results.

Figure 3: RLK7 is an additional CEP4 receptor. A) ($[Ca^{2+}]_{cyt}$) kinetics in seedlings upon CEP4 (1 µM) treatment; n = 6 ± SD. **B-C)** MAPK activation upon CEP4 (1 µM) treatment for the indicated time. Western blots were probed with α-p44/42. CBB = Coomassie brilliant blue. **D-E)** $[Ca^{2+}]_{cyt}$ kinetics in seedlings upon CEP4 treatment (1 µM); n = 3, n = 4, respectively ± SD. *c1c2r7 = cepr1/2/rlk7*[AEQ] **F)** MAPK activation upon CEP4 (1 µM) treatment for the indicated time. Western blots were probed with α-p44/42. CBB = Coomassie brilliant blue. **G)** cfu of *Pto* DC3000 (4 dpi) in distal leaves upon local CEP4 (5 µM) or mock (ddH$_2$O) pre-treatment; n = 9 pooled from three experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.01). **H)** $[Ca^{2+}]_{cyt}$ kinetics in seedlings upon CEP1 (10 µM) treatment; n = 4, ± SD. **I)** CEP4, **J)** PIP1 and **K)** CEP1 were titrated into a solution containing CEPR2[ECD] in ITC cells. Top: raw data thermogram; bottom: fitted integrated ITC data curves. DP = differential power between reference and sample cell; ΔH = enthalpy change. **L)** ITC table summarizing RLK7[ECD] vs CEP4/PIP1/CEP1 as mean ± SD of two experiments. The dissociation constant (K$_d$) indicates receptor-ligand binding affinity. N indicates reaction stoichiometry (n = 1 for 1:1 interaction). **M)** cfu of *Pto* DC3000 (3 dpi) upon spray infection; n = 12 pooled from three independent experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.0021; a-c p<0.0001; b-c p<0.0209). Experiments in **A-H**, **M** and **I-K** were repeated at least two times in independent biological repeats or in two independent technical repeats, respectively, with similar results.

Figure 4: Shoot-expressed *CEPs* and *CEPR1*, *CEPR2*, *RLK7* receptors are required for basal immunity against *Pto*. A) Cfu of *Pto*COR- (3 dpi) upon spray infection of reciprocally grafted Col-0 (C) and *cep6x* (*6x*) plants. The genotype of the shoot scion is indicated with S, and the genotype of the rootstock is indicated with R; n = 17-19 pooled from four experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.0001). **B)** RT-qPCR analysis of *CEP* expression in mock (ddH₂O) and *Pto* DC3000-inoculated leaves 24 h post-treatment. Housekeeping gene *UBQ5*; n = 3, ± SD. **C)** cfu of *Pto*COR- (3 dpi) upon spray infection of reciprocally grafted Col-0AEQ (C) and *cepr1/2/rlk7*AEQ (*3x*) plants. The genotype of the shoot scion is indicated with S, and the genotype of the rootstock is indicated with R; n = 15-17 pooled from three experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.0001). **D)** pCEP4::NLS-3xmVenus signal in the indicated plant tissues. For imaging the lateral root emergence zone and the hydathode region, the maximum projection of Z-stacks for mVenus is merged with Z-stacked propidium iodide (PI) signal. To show the lack of mVenus signal in the vasculature, the maximum projection of Z-stacks for mVenus is merged with the same single section of PI, showing a single epidermal layer or vasculature (cyan-dotted line). **E)** NLS-3xmVenus or Venus-H2B signal in the leaves of indicated lines. The maximum projection of Z-stacks for mVenus is merged with Z-stacked PI signal. The cyan-dotted line represents vasculature; scale bar = 100 µm. **F)** MAPK activation upon CEP1 (1), CEP3 (3) and CEP4 (4) 1 µM treatment. Western blots were probed with α-p44/42. CBB = Coomassie brilliant blue. Different symbols in **B** represent independent experiments. Experiments were repeated at least two times in independent biological repeats with similar results, except **F**, which has been repeated once with identical results.

Figure 5: CEPs promote FLS2 signaling during reduced N conditions. A) Ethylene concentration
in Col-0 leaf discs 3.5 h after mock (ddH$_2$O), CEP4 (1 µM) and/or flg22 (500 nM) treatment; n = 30,
seven pooled experiments ± SD (two-tailed Student's t-test, ** p<0.05, *** p = 0.0001). **B)** cfu of *Pto*
DC3000 (3 dpi) in Col-0 leaves upon mock (ddH$_2$O), CEP4 (1 µM) and/or flg22 (100 nM) pre-treatment;
n = 4 ± SD (one-way ANOVA, Tukey post hoc, a-b p<0.05). **C)** RT-qPCR of *CEP4* after 24 h transfer of
Col-0 seedlings to 100% N or 10 % N medium. Housekeeping gene *UBQ5;* n = 3 ± SD. **D)** MAPK
activation upon flg22 (100 nM) treatment after 24 h transfer of seedlings to 100% N or 10% N medium.
Western blots were probed with α-p44/42. CBB = Coomassie brilliant blue. **E)** Quantification of
pMPK6/MPK3 band intensities normalized to the CBB band and relative to 100% N flg22 set as 1 using
ImageJ software; n = 4 biological replicates. **F)** RT-qPCR of *FRK1* upon flg22 (100 nM) treatment for 4
h after 24 h transfer of seedlings to 100% N or 10% N medium. Housekeeping gene *UBQ5*; n = 3 ± SD
(one-way ANOVA, Tukey post hoc, a-b p<0.01, b-c ≤ 0,001, a-c <0.0001). **F)** MAPK activation upon
flg22 (100 nM) treatment after 24 h transfer of seedlings to normal N or reduced N medium. Western
blots were probed with α-p44/42. CBB = Coomassie brilliant blue. **H)** Quantification of pMPK6/MPK3
band intensities normalized to the CBB band and relative to 100% N flg22 set as 1 using ImageJ
software.; n = 4 biological replicates. **I)** RT-qPCR of *NRT2.1* upon flg22 (100 nM) treatment of
seedlings for 4 h after 24 h transfer to 100% N or 10% N. Housekeeping gene *UBQ5*; n = 3 ± SD.
Different symbols in figures **C**, **E**, **F**, **H** and **I** represent independent experiments. All experiments were
repeated at least three times in independent biological repeats with similar results.

   Supplementary Figures

Supplementary Fig. 1: *CEP4* expression is differentially regulated after flg22 perception.
A) *CEP4* is downregulated after flg22 treatment in an *asr3* mutant background (*32*). Data were obtained using Genevestigator software and are based on the AT_mRNASeq_ARABI_GL-1 data set. All *CEP* genes for which RNAseq data was available are depicted **B)** *CEP4* expression in seedlings is weakly downregulated after flg22 treatment. Col-0 seedlings were treated with flg22 (100 nM) or mock (ddH$_2$O) for the indicated time, after which seedlings were harvested for RT-qPCR analysis of *CEP4* abundance. Housekeeping gene *UBQ5*; n = 3 ± SD. **C)** *CEP4* shows higher expression in the roots. Col-0 seedlings were treated with flg22 (100 nM) or mock (ddH$_2$O) for the indicated time, after which seedlings were harvested for RT-qPCR analysis of *CEP4* abundance. Housekeeping gene *UBQ5*; n = 3 ± SD. **D)** Alignment of group I CEP peptide domains from *Arabidopsis thaliana* marked with colors based on amino acid properties and conservation. The conserved proline (P) residues are indicated in yellow. Different symbols in **B** and **C** represent independent biological experiments.

Supplementary Fig. 2: Characterization of *35S::CEP4*, CRISPR *cep* mutant alleles and *cep6x 35S::CEP4*.

A) *CEP4* transcript levels in two independent *CEP4* overexpression lines, shown as fold induction compared to Col-0. Housekeeping gene *UBQ5*; n = 4 ± SD. **B)** Cfu of *Pto* DC3000 3 dpi upon spray infection; n = 16 pooled from four experiments ± SD (one-way ANOVA, Tukey post-hoc test; a-b/c p<0.05; b-c p<0.0001). **C)** Characterization of four independent *cep* mutant alleles. Schematic diagram of *CEP1*, *CEP2*, *CEP3*, *CEP4*, *CEP6* and *CEP9* gene structure and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. The locus number and the length of the coding sequence (CDS) are indicated above the scheme for each gene. CRISPR *cep6x* is mutated in *cep1/2/3/4/6/9* (alleles *cep1.1, cep2.1, cep3.1, cep4.1, cep6.1, cep9.1*), CRISPR *cep5x* is mutated in *cep1/2/3/6/9* with CEP4 wild-type (alleles *cep1.2, cep2.2, cep3.2, cep6.2, cep9.2*). The specific location and type of mutations for each gene are indicated in the schematics describing the mutants. The CEP domain is indicated in blue, and the two CRISPR target sites are indicated in purple. **D)** Pictures of 5-week-old plants of the indicated genotypes grown on soil; scale bar = 2 cm. **E)** *CEP4* transcript levels in two independent lines of *cep6x 35S::CEP4* shown as fold induction compared to Col-0; Housekeeping gene *UBQ5*; n = 4 ± SD. Different symbols in **A** and **E** represent independent biological repeats.

1010

Supplementary Fig. 3: CEP4 is the strongest inducer of PTI responses on the whole tissue level.

A) Both *CEP4* variants trigger influx of calcium ions. Col-0^{AEQ} seedlings were treated with the indicated concentrations of CEP4 and CEP4-15mer. ([Ca^{2+}]$_{cyt}$) was measured for 30 min. Shown is total calcium influx; n = 12 pooled from two experiments. **B)** MAPK activation in Col-0 upon treatment with indicated concentrations of CEP4 or flg22. Western blots were probed with α-p44/42. CBB = Coomassie brilliant blue. **C)** RT-qPCR of *FRK1* in seedlings upon mock (ddH$_2$O), CEP4 (1 µM) or flg22 (100 nM) treatment for 4 h, shown as a fold induction compared to Col-0. Different symbols represent independent biological repeats. Housekeeping gene *UBQ5*; n = 2 ± SD. **D)** Kinetics of cytosolic calcium concentrations ([Ca^{2+}]$_{cyt}$) in Col-0^{AEQ} seedlings upon CEP4 (100 nM) and flg22 (100nM) treatment; n = 12, ± SD. **E)** Some CEPs induce Ca^{2+} influx at higher concentrations. Col-0^{AEQ} seedlings were treated with the indicated concentrations of CEPs. ([Ca^{2+}]$_{cyt}$) was measured for 30 min. The average calcium influx triggered by individual CEPs is normalized to mock (ddH$_2$O). The dotted line represents different experiments. Statistical analysis was performed separately for each group. Statistical significance was compared to CEP4^{scr} of the indicated concentration; n = 6 (one-way ANOVA, Dunnett post hoc test, a-b p<0.0001, A-B p<0.001, A-C p<0.0001). **F)** CEP4 and CEP1 induce *FRK1* promoter activity in the vasculature. Representative images of NLS-3xmVenus signal in *pFRK1::NLS-3xmVenus* lines upon mock (ddH$_2$O), CEP1 (100 nM), CEP4 (100 nM) or flg22 (100 nM) treatment for 16 h. Maximum projection of Z-stack for mVenus merged with Z-stacked propidium iodide (PI) signal. Cyan dotted line indicates vasculature,scale bar = 100 µm. The experiment was repeated three times in independent biological repeats with similar results.

1032

Supplementary Fig. 4: CEPs are important for resistance against *Pto*.
A) *CEP4* overexpression does not fully rescue the CRISPR *cep6x* susceptibility to *Pto* DC3000. Cfu of
Pto DC3000 (3 dpi) upon spray infection; n = 16 pooled from four experiments ± SD (one-way ANOVA,
Tukey post-hoc test; a-b p<0.0001; a-c, b-c p<0.05; a/c-b p≤0.0001). **B)** Loss of *CEPs* increases
Arabidopsis susceptibility to *Pto*COR-. Cfu of *Pto*COR- (3 dpi) upon spray infection; n = 4 ± SD (one-way
ANOVA, Tukey post-hoc test; a-b p = 0.0015; a-bc/a-c p<0.0001; b-c p = 0.0041). All experiments were
repeated at least three times in independent biological repeats with similar results.

1040

1041

1042

Supplementary Fig. 5: Characterization of *cepr2-4* and *cepr1/2*[AEQ] mutants and CEP4-induced seedling growth inhibition.

A) Characterization of the *cepr2-4* mutant. The *CEPR2* transcript could not be detected in *cepr2-4* mutant in a semi-quantitative RT-PCR analysis. Housekeeping gene *UBQ5*. **B)** Schematic diagram of the *CEPR1* and *CEPR2* genomic sequence, structure, and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. The locus number and the length of the coding sequence (CDS) are indicated above the scheme for each gene. The CRISPR *cepr1/2*[AEQ] mutant was generated in Col-0[AEQ] background. CRISPR *cepr1*[AEQ] (in *cepr1/2*[AEQ]) has two 1 bp insertions, which lead to a frameshift mutation and an early stop codon. CRISPR *cepr2*[AEQ] (in *cepr1/2*[AEQ]) has a 3020 bp deletion between the start and stop codon. The specific location and type of mutations for each gene are indicated in the schematics describing the mutants. The two CRISPR target sites are indicated in purple. The T-DNA insertion site in *cepr2-4* is indicated with a black triangle. **C)** CEPR1 and CEPR2 are similarly required for CEP4-induced seedling growth inhibition. Relative fresh weight of five-day-old seedlings treated with CEP4 (1 µM) for seven days. The dotted line represents different experiments. Statistical analysis was performed separately for each group; n = 42-60 and n = 46-48 pooled from five and four experiments, respectively (one–way ANOVA, Tukey post-hoc test, a-b p<0.0001; A-B p<0.0001; A/B-C p<0.05). **D)** CEP4-induced seedling growth inhibition is abolished in *cepr1-1xcepr2-1*. Relative fresh weight of five-day-old seedlings treated with CEP4 (1 µM) for seven days; n=16 ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.01). Similar results were obtained in three independent biological repeats.

1062

1063

1064

1065

1066
1067
Supplementary Fig. 6: Quality control of recombinant CEPR1, CEPR2 and RLK7 ectodomains
and supplemental ITC data.
A) Analytical size-exclusion chromatography (SEC) experiments show CEPR1^{ECD} aggregates in
comparison to single peaks for CEPR2^{ECD} and RLK7^{ECD}. **B)** SDS-PAGE of the ECD peaks of the SEC
analysis in **A**. **C)** ITC thermograms of the second technical repeat performed for each ectodomain and
peptide analyzed in Fig. 2 (**D-F**) and Fig. 3 (**I-K**).
1074

1075

1076

Supplementary Fig. 7: CEPR1 and CEPR2 are dispensable for flg22-induced ethylene accumulation and growth inhibition.

A) Basal and flg22-triggered ethylene production is higher in *cepr1-3/2-4* mutants. Ethylene concentration in leaf discs upon mock (ddH$_2$O) or CEP4 (1 µM) for 3.5 h; n = 7 - 8 pooled from two experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b/c/d, c-b/d p<0.0001, b-d p<0.005). **B)** flg22-induced seedling growth inhibition is not affected in *cepr1-3/2-4*. Relative fresh weight of five-day-old seedlings treated with flg22 (100 nM) for seven days; n = 14 ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.0001). All experiments were performed at least twice in independent biological repeats with similar results.

1086

Supplementary Fig. 8: Identification of RLK7 as an additional CEP4 receptor.
A) RT-qPCR of *FRK1* in seedlings upon mock (ddH$_2$O) or CEP4 (1 µM) treatment for 4 h. Housekeeping gene *UBQ5*. Different symbols represent three independent biological repeats ± SD. **B)** CEP4-triggered ethylene production is reduced in *rlk7* mutants. Ethylene concentration in leaf discs upon mock (ddH$_2$O) or CEP4 (1 µM) treatment for 3.5 h; n = 15 pooled from three experiments ± SD (one-way ANOVA, Tukey post hoc, a-b p<0.01). **C)** Schematic diagram of the *RLK7* and *IKU2* genomic sequence, structure, and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. The locus number and the length of the coding sequence (CDS) are indicated above the scheme for each gene. There are four *rlk7* alleles generated in Col-0^{AEQ} background: *rlk7.1*^{AEQ} in *rlk7/iku2*^{AEQ} double mutant, two single *rlk7*^{AEQ} mutants: *rlk7.2*^{AEQ} and *rlk7.3*^{AEQ}, and *rlk7.4*^{AEQ} in CRISPR *cepr1/2/rlk7*^{AEQ}. CRISPR *iku2* (in *rlk7/iku2*^{AEQ}) has a 29 bp insertion, followed by a 17 bp deletion. Mutations in both genes lead to a frameshift mutation and an early stop codon. The specific location and type of mutations for each gene are indicated in the schematics describing the mutants. The two CRISPR target sites are indicated in purple. **D)** [Ca^{2+}]$_{cyt}$ kinetics in seedlings upon CEP4 treatment (1 µM); n = 6, ± SD. **E)** MAPK activation upon CEP4 (1 µM) treatment. Western blots were probed with α-p44/42. CBB = Coomassie brilliant blue. **F)** Relative fresh weight of five-day-old seedlings treated with CEP4 (1 µM) for seven days; n = 48 pooled from four experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b/c/bc p<0.0001, b-c p<0.05). **G)** [Ca^{2+}]$_{cyt}$ kinetics in seedlings upon CEP9.5 (10 µM) treatment; n = 6, ± SD. **H)** cfu of *Pto*$^{COR-}$ (3 dpi) upon spray infection; n = 8 pooled from two experiments ± SD (one-way ANOVA, Tukey post-hoc test; a-b, p<0.0120; a-bc p=0.0019). All experiments were performed at least twice in independent biological repeats with similar results.

1111
Supplementary Fig. 9: Characterization of *35S::CEPR1-GFP* and *35S::CEPR2-GFP*
overexpression lines.
A) *CEPR2* transcript levels in two independent *CEPR2-GFP* overexpression lines. *CEPR2* expression
was normalized to *UBQ5* and is shown as fold induction compared to Col-0. **B)** CEPR2-GFP protein
levels in wild-type Col-0 and *CEPR2* overexpression lines. Western blots were probed with α-GFP. CBB
= Coomassie brilliant blue. **C)** *CEPR1* transcript levels in the *CEPR1-GFP* overexpression line. *CEPR1*
expression was normalized to *UBQ5* and is shown as fold induction compared to Col-0. **D)** MAPK
activation in Col-0 and *35S::CEPR1-GFP* line upon CEP1 (1), CEP3 (3) and CEP4 (4) 1 µM treatment.
Western blots were probed with α-p44/42. CBB = Coomassie brilliant blue. Different symbols in **A** and
C represent two independent experiments. All experiments were performed at least twice in
independent biological repeats with similar results.

1123

Supplementary Fig. 10: Further characterization of CRISPR *cep* and CRISPR *cepr1/2/rlk7*[AEQ] mutants.

A) Flg22-triggered ethylene production is not impaired in *cep6x*. Ethylene concentration in leaf discs upon mock (ddH$_2$O) or flg22 (500 nM) treatment for 3.5 h; n = 21 pooled from four experiments ± SD (one-way ANOVA, Tukey post hoc, a-b p<0.05; a/b-c p<0.0001). **B)** CEP4-triggered ethylene production is abolished in *cepr1/2/rlk7*[AEQ]. Ethylene concentration in leaf discs upon mock (ddH$_2$O), CEP4 (1 µM) or flg22 (500 nM) treatment for 3.5 h; n = 10 pooled from two experiments ± SD (one-way ANOVA, Tukey post hoc, a-b p=0.0166; a/b/ab-c p<0.0001). *cep6x* **(C)** and *cepr1/2/rlk7*[AEQ] **(D)** are not affected in flg22-triggered MAPK activation under normal conditions. MAPK activation upon flg22 (100 nM) treatment. Western blots were probed with α-p44/42. CBB = Coomassie brilliant blue. **E)** *cep6x* is not affected in flg22-triggered expression of *FRK1* in seedlings. RT-qPCR of *FRK1* upon flg22 (100 nM) treatment for 4 h normalized to *UBQ5*. Different symbols represent different experiments; n = 3 ± SD (one-way ANOVA, Tukey post hoc, a-b p<0.0001). **F)** N availability modulates flg22-triggered MAPK activation. MAPK activation upon flg22 (100 nM) treatment after 24 h transfer of seedlings to 100% N, 10% N, 5% N and 1% N medium. Western blots were probed with α-p44/42. CBB = Coomassie brilliant blue. **G)** The *cep6x* mutant shows wild-type growth response to different medium N concentration. Relative fresh weight of five-day-old seedlings grown in 100% N, 10% N and 1% N medium for seven days; n = 40 pooled from five independent experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.001). **H)** The *cepr1/2/rlk7*[AEQ] shows slightly stronger growth response to different medium N concentrations. Relative fresh weight of five-day-old seedlings grown in 100% N, 10% N and 1% N medium for seven days; n = 39-40 pooled from five independent experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-cd p<0.0001, a-bc p=0.0002, a-d p<0.0001, ab-d, p<0.0001, bc-d p=0.0303).

1146

1147

Peptide name	Sequence
CEP1	DFRHypTNPGNSHypGVGH
CEP2	DFAHypTNPGDSHypGIRH
CEP3	TFRHypTEPGHSHypGIGH
CEP4^{scr}	TGQHypDHQRHypFAHIGGS
CEP4^{15mer}	AFRHypTHQGHypSQGIGH
CEP4^{16mer}	DAFRHypTHQGHypSQGIGH
CEP6.2	DFEPTTHypGHSHypGVGH
CEP9.1	DFVHypTSHypGNSHypGVGH
CEP9.2	DFAHypTSHypGHSHypGVGH
CEP9.3	DFAHypTSHypGNSHypGIGH
CEP9.4	DFAHypTTHypGNSHypGMGH
CEP9.5	DFKPTTHypGHSHypGVGH
PIP1	RLASGHypSPRGRGH
flg22	Ac-QRLSTGSRINSAKDDAAGLQIA

1148
1149 **Supplementary Table 1: Peptides used in this study.**

1150
1151 Hyp (in red) indicates hydroxylated proline residues.

1152

1153
1154

Target site	Sequence	Purpose
CEP1.1	TGGTCACATATACACGGCG	Generation of CRISPR *cep* mutants
CEP1.2	CAGGCGTTGGACACTCTAA	
CEP2.1	GTCCCGGTATCAGGCATCC	
CEP2.2	ATGTATGCCCTGCAAAAGT	
CEP3.1	TACGGAATGTCCAATACCG	
CEP3.2	TATCAACCCCGTGGCTCGG	
CEP4.1	CATTCCGACCAACCCACCA	
CEP4.2	TATTTTAAGGAGCACCTGG	
CEP6.1	ATGATCATCATTTCACAGT	
CEP6.2	TCGGGTACACTGATGATTT	
CEP9.1	CAAGATGACTTCAAGCCCA	
CEP9.2	ACACCAGGACATAGCCCCG	
CEPR1.1	AGAGAGACGAAGTACACGG	Generation of *cepr1/2*[AEQ] mutant
CEPR1.2	AGCTCATGAGGTATAACAC	
CEPR2.1	GACCTCCTACGTGGCAGCG	
CEPR2.2	CTCAGACCGAGTATGAGAG	
RLK7.1	AGATTGATCTCTCTCGTCG	Generation of *rlk7*[AEQ] and *rlk7/iku2*[AEQ] mutants
RLK7.2	GCGACGGCTGATTTCCCGG	
IKU2.1	GCTTGAGATACTTAGACCT	Generation of *iku2*[AEQ] mutant
IKU2.2	CTTGTTCGTCTACAAAACT	

1155 **Supplementary Table 2: CRISPR-Cas9 target sites.**

1156

Primers used for cloning		
Primer name	**Sequence (5'→3')**	**Purpose**
pCEPR1-BsaI-F	GCGCGCGGTCTCTCGAGTCAGTTTTTTT GGATTCAGATTTG	*pro*CEPR1 for pCEPR1::NLS-3xmVenus-tNOS
pCEPR1-BsaI-R	GCGCGCGGTCTCTCATTTTTCAGAGAAA GATCAAAAGTAACC	
pCEPR2-BsaI-F	GCGCGCGGTCTCTCGAGTTTTGTATACT CTTAAAAAAACTATG	*pro*CEPR2 for pCEPR2::NLS-3xmVenus-tNOS
pCEPR2-BsaI-R	GCGCGCGGTCTCTCATTTTATAGTGATT CCCAAGGG	
pCEP4-BpiI-F	TTGAAGACTTCGAGAATATAAACAAATA TAGGTAAC	*pro*CEP4 for pCEP4::NLS-3xmVenus-t35S
pCEP4-BpiI-R	TTGAAGACTTCATTCACTTGACAACTTTC TG	
pRLK7-AttB1	GGGGACAAGTTTGTACAAAAAAGCAGG CTTAGAGCTAAGACATATAACT	*pro*RLK7 for pRLK7-H2B-mVenus
pRLK7-AttB2	GGGGACCACTTTGTACAAGAAAGCTGG GTAGTCGGAGAGAAGAAGTGG	
CEPR1-AttB-F	GGGGACAAGTTTGTACAAAAAAGCAGG CTTTTATGCGTCTCAAAAATTTCCC	35S::CEPR1-GFP
CEPR1-AttB-R	GGGGACCACTTTGTACAAGAAAGCTGG GTCGAGTCTTGTTTGCGTGAG	
CEPR2-AttB-F	GGGGACAAGTTTGTACAAAAAAGCAGG CTATGTCGAGAAGACCAGACC	35S::CEPR2-GFP
CEPR2-AttB-R	GGGGACCACTTTGTACAAGAAAGCTGG GTCTACTGTAATCTTTCCAGTTGTGTC	
Primers used for RT-qPCR		
Primer name	**Sequence (5'→3')**	**Purpose**
qUBQ5-F	ACTCCTTCCTCAAACGCTGA	*UBQ5*
qUBQ5-R	CCAAGCCGAAGAAGATCAAG	
qFRK1-F	TGCAGCGCAAGGACTAGAG	*FRK1*
qFRK1-R	ATCTTCGCTTGGAGCTTCTC	
qPR1-F	CGGAGCTACGCAGAACAACT	*PR1*
qPR1-R	CAGACAAGTCACCGCTACCC	
qNRT2.1-F	AACAAGGGCTAACGTGGATG	*NRT2.1*
qNRT2.1-R	CTGCTTCTCCTGCTCATTCC	
qCEP1-F	AATGCTAAAGGGGTGTTTGG	*CEP1*
qCEP1-R	ACAAACCCACGACAAAGACA	
qCEP2-F	TGGTGACCATTTTGACCATC	*CEP2*
qCEP2-R	CCGACCATCTTTTTCGACTT	
qCEP3-F	GACCTACGGAACCTGGTCAT	*CEP3*
qCEP3-R	AAAAAGTCACCAGGCCAATC	
qCEP4-F	AGAATACAAAAGCAGCTCGAC	*CEP4*
qCEP4-R	TCCAATACCTTGACTAGGACC	
qCEP6-F	GTCGGGTAATCTAGACCTTCCTC	*CEP6*
qCEP6-R	CGGTTTTTCGCAACTGTCTCG	
qCEP9-F	AGCTGGATTTACAGATGATTTCG	*CEP9*
qCEP9-R	TGTCCCACACCAGGACTGT	
qCEPR1-F	ATGCGTACTCGTCCAAAGCA	*CEPR1*
qCEPR1-R	CTCCCCGAAACACGAATCCA	
qCEPR2-F	CTTCAAAAACCGCCTCGACG	*CEPR2*
qCEPR2-R	CCGATGACTTCGCCGGATAA	
Primers used for genotyping and sequencing		
Primer name	**Sequence (5'→3')**	**Purpose**
GG pUC18 Seq-F	GTATCACGAGGCCCTTTCGT	Inserts in GoldenGate pUC18
GG pUC18 Seq-R	TAATGAATCGGCCAACGC	
CEP1-CRISP-s1	CCTTGTTTTGTTTTACATATTCTT	CRISPR CEP1
CEP1-CRISP-s2	TCTGAATGAAAATACATGTAATTTTC	

CEP2-CRISP-s1	CATTCAATACTTCTCATATACAAAAC	CRISPR CEP2
CEP2-CRISP-s2	TCACCAGTTTTCTTTAGTCTTC	
CEP3-CRISP-s1	CATTTTCGCCTTCGACTA	CRISPR CEP3
CEP3-CRISP-s2	CAATCATATTTTCAACATGTAATCAC	
CEP4-CRISP-s1	GACATACGCTTTTAAAGGAAG	CRISPR CEP4
CEP4-CRISP-s2	GAAGAACTGTGTGCGATAAAG	
CEP6-CRISP-s1	CCTCTCTCGTATACACTAGAAAC	CRISPR CEP6
CEP6-CRISP-s2	TTAATTTAGCAAAAGCATTGAAG	
CEP9-CRISP-s1	ATATATACAAACTCCTAAACCTTGC	CRISPR CEP9
CEP9-CRISP-s2	CATATCGTAAACCAATGTATTAGA	
CEPR1-CRISP-s1	CGTAAGATGTGACGGTCAA	CRISPR CEPR1
CEPR1-CRISP-s2	GGTTGCCTTGCAGTACAA	
CEPR2-CRISP-s1	GAATTCCAGTGCATCGACC	CRISPR CEPR2
CEPR2-CRISP-s2	CTAAAACGTGGTAGGTGAATCAT	
RLK7-CRISP-s1	TCAACCACCGTCACCACTT	CRISPR RLK7
RLK7-CRISP-s2	TCCAAATTCCGAAGCTCCGT	
IKU2-CRISP-s1	GCATGTGAATTCGCCGGAAT	CRISPR IKU2
IKU2-CRISP-s2	CCGCAGCTCTGATAAATCACC	
pCEPR1-s1	CTAGTCACCTGCAGCTCG	pCEPR1
pCEPR1-s2	GTGTAGTTCACGTCGGATG	
pCEPR1-s3	CAAACGCTTCACTTATGTAATG	
pCEPR1-s4	GCATCTATCTGATTCTGATCGTG	
pCEPR1-s5	CATCCGACGTGAACTACAC	
pCEPR2-s1	ACGACAATCTGAGCTCCAC	pCEPR2
pCEPR2-s2	TGGGATTCGGAGCATAAGG	
pCEPR2-s3	TCCATCTAGAAACATTAACCGA	
pCEPR2-s4	CCTAGAACGTGGTCCAAG	
pCEPR2-s5	CTGCACTGTAATGAGTACC	
pCEPR2-s6	CGGTAATAGATAGTGAAATGG	
cepr1-3-LP	TAAATCAAGAATCCACTTCCATGC	*cepr1-3*
cepr1-3-RP	CCTAATGGTAACTTATGGGACGCT	
cepr2-4-LP	GGTGAGTTCGTTATCGCTGAG	*cepr2-4*
cepr2-4-RP	AATGTTGAATCGACCGTTGAG	
GABI_o8474	ATAATAACGCTGCGGACATCTACATTTT	T-DNA primer for GABI-kat
cepr2-3-LP	TCACAACTCTGTAACGCAACG	*cepr2-3*
cepr2-3-RP	AACTCGGAGTTTTGAAGGAGC	
rlk7-1-LP	CCGCCTCTCTCTCTCTCTC	*rlk7-1*
rlk7-1-RP	AAGCAGAGCTTTCATCTTCCC	
rlk7-3-LP	ACGATTTGATCGTCGTGCTAC	*rlk7-3*
rlk7-3-RP	TTACAACAACTCCTTGACCGG	
iku2-4-LP	TCTTTAAGAACCGCAGCTCTG	*iku2-4*
iku2-4-RP	GTTGTTTCGCCTACAATGACC	
LBb1.3	ATTTTGCCGATTTCGGAAC	T-DNA LB primer for *rlk7-1, rlk7-3, iku2-4*
LBb1	GCGTGGACCGCTTGCTGCAACT	T-DNA LB primer for *cepr2-3*

1157

1158 **Supplementary Table 3: Primers used in this study.**

1159

1160
1161

1162
1163