Phytocytokine signaling integrates cell surface immunity and nitrogen limitation

Jakub Rzemieniewski[1], Henriette Leicher[1], Hyun Kyung Lee[2], Caroline Broyart[2], Shahran Nayem[3], Christian Wiese[1,†], Julian Maroschek[1], Zeynep Camgöz[1], Vilde Olsson Lalun[4], Michael Anthony Djordjevic[5], A. Corina Vlot[3,‡], Ralph Hückelhoven[1], Julia Santiago[2] and Martin Stegmann[1*]

[1]Phytopathology, TUM School of Life Sciences, Technical University of Munich, Freising, Germany

[2]The Plant Signaling Mechanisms Laboratory, Department of Plant Molecular Biology, University of Lausanne, Lausanne, Switzerland

[3]Helmholtz Zentrum Munich, Institute of Biochemical Plant Pathology, Neuherberg, Germany

[4]Section for Genetics and Evolutionary Biology, Department of Biosciences, University of Oslo, Oslo, Norway

[5]Division of Plant Science, Australian National University, Acton, Canberra, Australia

†present address: Botany, TUM School of Life Sciences, Technical University of Munich, Freising, Germany

‡present address: Chair of Crop Plant Genetics, Faculty of Life Sciences: Food, Nutrition and Health, University of Bayreuth, Kulmbach, Germany

*Corresponding author: Email: martin.stegmann@tum.de

Abstract

Little is known about plant genetic and biochemical components that coordinate immune responses with growth and environmental cues. C-TERMINALLY ENCODED PEPTIDEs (CEPs) control plant development and nitrogen demand signaling. Here, we identified CEP4 as an immune-modulatory peptide (phytocytokine) in *Arabidopsis thaliana*. CEP4 and related CEPs are important regulators of resistance to plant pathogenic bacteria and are perceived by the tissue-specific receptor kinases CEP RECEPTOR 1 (CEPR1), CEPR2 and the phylogenetically related RECEPTOR-LIKE KINASE 7 (RLK7). CEP4 promotes flagellin-triggered responses and we provide evidence that CEPs modulate cell surface immunity upon N limitation. We propose that CEPs integrate biotic and abiotic stress-associated signals to safeguard plant health.

Introduction

Receptor kinases (RKs) sense external and internal cues to control multiple aspects of plant physiology. RKs can serve as pattern recognition receptors (PRRs) to detect microbe-associated molecular patterns (MAMPs) and activate pattern-triggered immunity (PTI). An example is the *Arabidopsis thaliana* (hereafter Arabidopsis) leucine-rich repeat RK (LRR-RK) FLAGELLIN SENSITIVE 2 (FLS2), which forms a receptor complex with BRASSINOSTEROID INSENSITIVE 1-ASSOCIATED RK 1 (BAK1) upon perception of a 22 amino acid epitope derived from bacterial flagellin (flg22) to activate PTI (*1*). Plants also perceive endogenous peptides that plants release during infection or MAMP recognition. These peptides are referred to as phytocytokines and are also perceived by RKs. They modulate immunity but often also plant growth and development (*2*, *3*). In many cases, phytocytokines are transcriptionally induced upon MAMP

perception, but this is not a prerequisite for their function, as previously shown for GLV2 (*4–7*). Phytocytokines and other endogenous peptides regulate several abiotic stress responses, including adaptation to high salinity, drought and nutrient deprivation, suggesting that they might integrate multiple external and internal cues to safeguard plant health (*8–10*). Yet, information is sparse on plant cell surface components with functions in more than one peptide-mediated pathway. C-TERMINALLY ENCODED PEPTIDES (CEPs) are important for sucrose-dependent lateral root growth, root system architecture, systemic nitrogen (N)-demand signaling and promotion of root nodulation, but a role in plant immunity remained unknown (*8*, *11–15*). Here, we studied CEPs as new phytocytokines in Arabidopsis and discovered a function in adaptation of immune responses to N availability.

Results

CEPs are novel phytocytokines

Screening of public transcript abundance data identified a specific member of the CEP family, *CEP4*, that showed differential flg22-induced expression (Fig. S1A) (*16*). RT-qPCR confirmed flg22-dependent downregulation of *CEP4* expression in Col-0 roots (Fig. S1B). CEPs are produced as peptide precursors and contain one to five CEP domains that require maturation for peptide release (*17*). To test for CEP4 involvement in immunity, we generated CEP4 precursor overexpression lines (*35S::CEP4*, Fig. S2A) which were more resistant to *Pseudomonas syringae* pv. *tomato (Pto)* lacking the chemical effector molecule coronatine (*Pto*[COR-]) (Fig. 1A). Arabidopsis encodes 12 class I CEPs including CEP4 which has a distinctive sequence (*18*). CEP4 only carries two proline residues in its peptide domain and the second proline is positioned uniquely compared to other classical class I CEPs with three prolines (e.g. CEP1 and CEP3) (Table S1). Most previously identified CEPs are 15mers with an N-terminal aspartate (D), a C-terminal histidine (H) and hydroxylated prolines (*8*). We tested whether a 16mer CEP4 with proline hydroxylation (Hyp), DAFRHypTHQGHypSQGIGH, triggers immune responses. CEP4 induced dose-dependent PTI signaling outputs, including cellular influx of calcium ions in Col-0 expressing the calcium reporter aequorin (Col-0[AEQ]), activation of MITOGEN-ACTIVATED PROTEIN KINASEs (MAPKs), ethylene production and expression of the PTI marker gene *FLAGELLIN-INDUCED RECEPTOR KINASE 1* (*FRK1*) (Fig. 1B-E, Fig. S3A). CEP4 also induced YFP fluorescence in the vasculature of *pFRK1::NLS-3xmVenus* seedlings (*19*) (Fig. 1F), seedling growth inhibition (SGI) and systemic resistance to *Pto* (Fig. 1G,H). The 15mer CEP4 (AFRHypTHQGHypSQGIGH) triggered dose-dependent calcium influx similar to the 16mer (Fig. S3A). CEP1 and CEP9.5 (Table S1) also triggered significant calcium influx in seedlings, but only at high concentrations (10 µM) (Fig. S3B). CEP1 further induced YFP fluorescence in the vasculature of *pFRK1::NLS-3xmVenus* lines (Fig. S3C). We confirmed a role for class I CEPs in plant immunity by generating a *cep6x* hextuple mutant by CRISPR-Cas9 in which we mutated CEP4, CEP1-CEP3, CEP6 and CEP9 (Fig. S2B,C). The *cep6x* mutant showed compromised resistance to both *Pto*[COR-] and *Pto* wild type, demonstrating that CEPs are important for antibacterial resistance (Fig. 1I, Fig. S2D). The *cep6x* susceptibility phenotype was complemented with *35S::CEP4* (Fig. 1J, Fig. S2E).

Class I CEPs predominantly express in root tissue (*17*) and function as systemic root-to-shoot transmitters of N starvation (*8*). We hypothesized that root-expressed *CEPs* regulate leaf immunity upon *Pto* infection. Reciprocal grafts between *cep6x* and Col-0 showed, however, that *CEP* mutation in the shoot conferred increased *Pto* susceptibility (Fig. S4A). We detected *CEP1-CEP4*, *CEP6* and *CEP9* expression in leaf tissue using RT-qPCR and imaging *pCEP4::NLS-3xmVENUS* reporter plants (Fig. S4B,C) with *CEP4* being slightly upregulated upon *Pto* infection (Fig. S4B). The *CEP4* promoter showed activity in seedling leaf tissue but not in the vasculature (Fig. S4D). Consistent with previous reports, *CEP4* expressed in emerging lateral roots (Fig. S4C) (*17*). Given that CEP4 induced *FRK1* expression in the vasculature (Fig. 1F, S3C), it is possible that CEP4 is mobile between different tissue layers of the leaf.

CEPR1 and CEPR2 are CEP4 receptors and central regulators of plant immunity

CEPs bind to CEPR1 and CEPR2, two LRR-RLKs from subfamily XI (*8*). We resolved expression patterns of *CEPR1* and *CEPR2* in leaf tissues using *pCEPR1::NLS-3xmVenus* and *pCEPR2::NLS-3xmVenus* lines. As expected, *CEPR1* expressed in vasculature tissue (Fig. 2A), as previously reported (*8*), whereas *CEPR2* was largely restricted to stomatal guard cells (Fig. 2A). We tested whether *CEPR1* and *CEPR2* control immunity. Single *cepr1-3* or *cepr2-4* mutants did not show a Pto^{COR-} infection phenotype (Fig. 2B, S5A,B). However, *cepr1-3 cepr2-4 (cepr1-3/2-4)* and a CRISPR-Cas9 *cepr1 cepr2* double mutant in Col-0^{AEQ} (*cepr1/2*^{AEQ}) were both more susceptible to Pto^{COR-} (Fig. 2B, S5A). These data suggest that *CEPR1* and *CEPR2* spatially complement each other to control Pto^{COR-} resistance.

We tested whether CEP4 perception depends on CEPR1 and/or CEPR2. Using SGI as a readout, *cepr1-3* was insensitive and two *cepr2* alleles (*cepr2-3* and *cepr2-4*) were insensitive and less sensitive to CEP4 treatment, respectively (Fig. S6A,B). The *cepr1-3/2-4* mutant and No-0 *cepr1-1 cepr2-1* double mutant did also not respond to CEP4 (Fig. 2C, Fig. S6C). Similarly, the *cepr1-3/2-4* mutant was insensitive to CEP4-induced systemic resistance (Fig. 2D). These data suggest that both CEPR1 and CEPR2 mediate CEP4 perception.

We then tested whether CEP4 binds to purified insect cell-expressed ectodomains (ECDs) of CEPR1 and/or CEPR2. We analyzed protein quality by Coomassie stain and size exclusion chromatography (SEC, Fig. S7A,B). Unfortunately, CEPR1^{ECD} aggregated, as indicated by the early elution of the bulk sample during SEC analysis (~8 mL corresponding to the column void volume, Fig. S7A). By contrast, isothermal titration calorimetry (ITC) showed that CEPR2^{ECD}, which eluted with a single SEC peak at ~13 mL (Fig. S7A,B), directly bound CEP4 with a K_D of 15.7 µM (± 4.5 µM) (Fig. 2E, H, Fig. S7C). A scrambled control (CEP4^{scr}) did not bind to CEPR2^{ECD} (Fig. 2F, H, Fig. S7C). CEP1, which was previously shown to bind to CEPR1 and CEPR2 (*8*), also bound to CEPR2^{ECD} with a K_D of 9.3 µM (± 0.6 µM) (Fig. 2G,H, Fig. S7C). These data are in range with previously reported peptide-LRR-RK binding affinities obtained by ITC and suggest that CEPR2 is a CEP4 receptor (*20–23*).

We tested whether *CEPR1* and *CEPR2* were important for FLS2-mediated PTI. We did not observe strong differences in flg22-induced ethylene accumulation in *cepr1-3/2-4* (Fig. S8A). Of note, flg22-induced ethylene production was higher in *cepr1-3/2-4*, but basal ethylene production was also enhanced, making

the result difficult to interpret. However, we found that flg22-induced *FRK1* and *PR1* expression was strongly reduced in *cepr1-3/2-4* and reduced in *cepr1-3* and *cepr2-4* (Fig. 2I). The flg22-induced seedling growth inhibition was not affected in *cepr1-3/2-4* (Fig. S8B), suggesting that CEPR1/CEPR2 are selectively required for specific flg22-induced outputs. We also tested whether CEP-CEPR1/2 signaling was important for systemic acquired resistance (SAR). Local inoculation of *Pto* avrRPM1 and subsequent infection of systemic tissue with virulent *Pto* (*24*) revealed that *cepr1-3/2-4* and *cep6x* were strongly compromised in *Pto* avrRPM1-triggered SAR (Fig. 2J). Collectively, these data show that CEP-CEPR signaling is a central regulator of plant immunity in Arabidopsis.

RLK7 is a novel CEP4-specific CEP receptor

To further characterize the role of CEPR1 and CEPR2 in CEP4 signaling, we tested early CEP4-induced responses in *cepr1-3/2-4* and *cepr1/2*[AEQ]. Surprisingly, we found that *cepr1/2*[AEQ] did not show compromised calcium influx upon CEP4 treatment (Fig. 3A). CEP4-induced MAPK activation was also unaffected in *cepr1-3/2-4* (Fig. 3B). Moreover, *cepr1-3* and *cepr2-4* showed mildly reduced CEP4-induced *FRK1* expression, which was more pronounced, but not abolished, in *cepr1-3/2-4* (Fig. 3C). These results suggested that other receptor(s) are involved in CEP4 perception. CEPR1 and CEPR2 are phylogenetically close to IKU2, which is involved in seed size regulation, and RLK7 which controls germination speed, lateral root formation and salt stress adaptation (*10*, *25–27*). RLK7 also senses endogenous PAMP-INDUCED PEPTIDES (PIPs) to regulate resistance to the fungal vascular pathogen *Fusarium oxysporum* and *Pto* (*28*, *29*). The *iku2-4* mutant showed unaltered CEP4-induced ethylene accumulation, but the response was strongly compromised in *rlk7-1* and *rlk7-3* (Fig. S9A). CEP4-induced ethylene production was also impaired in *cepr1-3/2-4*, but the results are difficult to interpret given the mutant´s high basal ethylene accumulation (Fig. S9B). Similarly, *rlk7-1* and *rlk7-3* showed strongly reduced CEP4-induced MAPK activation (Fig. 3D). We also generated a *rlk7/iku2*[AEQ] mutant (Fig. S5C) which showed strongly compromised CEP4-induced calcium influx (Fig. 3E). CEP4-induced *FRK1* expression was mildly affected in *rlk7-1* and *rlk7-3,* similar to *cepr1-3/2-4* (Fig. 3F,C). These data suggest that RLK7 also functions as a CEP4 receptor. Interestingly, *rlk7-1* and *rlk7-3* were only moderately affected in CEP4-induced SGI (Fig. 3G), suggesting that CEP4 activates distinct responses via specific receptors. Moreover, CEP1-induced calcium influx was abolished in *cepr1/2*[AEQ] and unaltered in *rlk7/ iku2*[AEQ], indicating that CEPR1/2 are required for canonical class I CEP-mediated early immune signaling responses (Fig. 3H).

We used ITC to test whether CEP4 binds to RLK7[ECD]. Similar to CEPR2[ECD], we obtained good quality proteins for RLK7[ECD] eluting with a main single SEC peak at ~13 mL (Fig. S7A,B). CEP4 bound to RLK7[ECD] with a K_D of 9 µM (± 4.9 µM) (Fig. 3I,L, Fig. S7), like CEPR2[ECD] (Fig. 2E,H). We also tested binding of PIP1, a defined RLK7 ligand, to RLK7[ECD] (*10*, *28*). RLK7[ECD] bound PIP1 with higher affinity (K_D 500 nM ± 60 nM) (Fig. 3J,L, S7C). Importantly, consistent with unaltered CEP1-induced responses in *rlk7/iku2*[AEQ] (Fig. 3H), CEP1 did not bind to RLK7 (Fig. 3K,L Fig. S7C). Collectively, these data suggest that RLK7 can also function as a CEP4-specific CEP receptor. By analyzing a *pRLK7::Venus-H2B*, we found wide-spread

expression of *RLK7* which overlapped with *CEP4* in aerial parts of seedlings (Fig. 3M, S4C). This suggests

that CEP4 and RLK7 meet *in vivo* to function as a receptor-ligand pair.

CEPs promote FLS2 signaling during N starvation

Class I *CEP* expression, including *CEP4*, is not strongly affected by flg22 treatment or *Pto* infection (Fig.

S1A,B, S4B). We were thus interested to determine the biological context of CEP-mediated control of

immunity. We tested whether CEP4 treatment also affects FLS2-dependent signaling. CEP4 promoted

flg22-induced ethylene accumulation and resistance induced by a low dose of flg22 (100 nM) (Fig. 4A,B).

Yet, we did not observe defects in flg22-induced MAPK activation and *FRK1* expression in *cep6x* (Fig.

4C,D). Several *CEPs* are transcriptionally upregulated in N-starved roots, including *CEP1* (*8*). Similarly,

CEP4 was transcriptionally upregulated upon N limitation of seedlings (Fig. 4E). The N status of plants

affects disease resistance to different pathogens, but the underlying molecular mechanisms remain

unknown (*30*). N starvation promoted resistance to *Pto*, suggesting a connection between N homeostasis

and antibacterial resistance (*31*). We thus hypothesized that CEPs coordinate the plant´s N status with cell

surface immunity. We N-deprived two-week-old seedlings for 24 h by transferring them to ½ MS medium

with low N content (10% N concentration of ½ MS medium, low N). Upon elicitation, wild type seedlings

showed enhanced flg22-triggered MAPK activation and *FRK1* expression (Fig 4F,G). This N status-

dependent promotion of flg22-triggered PTI was abolished in *cep6x.* These data suggest that CEPs

contribute to FLS2-dependent signaling under N limited conditions. Moreover, flg22 stimulated expression

of the high affinity *NITRATE TRANSPORTER 2.1 (NRT2.1)* (*32*) upon N limitation in a *CEP*-dependent

manner (Fig. 4H). Together, these results indicate a tight CEP-dependent connection between PTI and N

availability, highlighting signaling cross-talk between cell surface immunity and the plant`s nutritional status.

Discussion

This work reveals that CEP perception is achieved by three partially tissue-specific receptors that regulate

plant immunity and likely coordinate biotic stress with plant nutritional cues. The close overlap of *CEP4* and

RLK7 expression in leaves, but not with *CEPR1* and *CEPR2*, suggests a combination of cell autonomous

and short distance signaling contributing to CEP-mediated immune modulation. A challenge for the future

will be to determine how three CEP receptors with distinct expression patterns integrate responses between

tissues and the concerted action of diverse ligands. RLK7 also recognizes endogenous PIPs, likely with

higher affinity over CEP4 (*26*, *28*). Members of the CLE peptide family also bind multiple receptors with

variable binding affinities to regulate epidermal cell patterning (*22*, *33*). Dissecting spatiotemporal ligand

availability and CEP4-PIP1 signaling specificity will be an important future task. RLK7 does not bind to

CEP1, but CEP4, suggesting that CEP4`s receptor specificities are unique among CEPs, likely caused by

its distinct sequence. It will be interesting to compare the molecular mechanism of PIP1/CEP4-RLK7 and

CEP1/CEP4-CEPR2/CEPR1 recognition in future structure analyses. Plant peptides may be versatile tools

for the evolution of plant plasticity. Increasing ligand and receptor diversity may provide vast combinations
for fine-tuned signal outputs.

Our data suggest that N starvation facilitates CEP expression to promote flg22 responsiveness. Other
phytocytokines modulate immune signaling by regulating PRR abundance or receptor complex formation
and dynamics, but in most cases the mechanism remains unknown (*3*, *7*, *34*, *35*). It will be interesting to
reveal whether CEPs directly or indirectly modulate PRR signalling.

Recent evidence suggests a direct integration of nutrient homeostasis and PTI in plants. Perception of
flg22 induces PHT1.4 phosphorylation to inhibit phosphate (Pi) uptake and promote root immunity (*36*).
CEPs induce nitrate, Pi and sulfate uptake, suggesting CEP-CEPR1/CEPR2/RLK7-dependent transporter
modulation (*37*). NRT transport activity can be regulated by kinases, including phosphorylation of NRT1.2
by CEPR2, which we identified as a CEP4 receptor (*38*, *39*). Since N starvation promotes CEP-dependent
PTI, it is also possible that sufficient N dampens immune signaling via a yet unknown mechanism. This
may be important in ecological or agronomic contexts, when plants need to decide whether to invest into
defense or growth depending on nutrient availability.

CEP signaling also regulates sucrose-dependent root growth inhibition and fecundity (*14*, *40*), placing
CEPs as central integrators of biotic stress with plant nutrition, growth and development (*37*).
CEPR1/CEPR2/RLK7 and CEPs are widely conserved among angiosperms, including crop plants (*41*).
CEPs are important to promote nodulation in legumes (*42*, *43*). It will be critical to understand whether N-
dependent and CEP-mediated promotion of PTI extends beyond Arabidopsis and to decipher how CEPs
can both promote immunity and control symbiosis in diverse species. This will provide important insights
for future crop improvement strategies that coordinate crop nutrition with disease resistance.

Materials and methods

<u>Molecular cloning</u>
To generate *CEP4* overexpression lines, the coding sequence of *CEP4* (AT2G35612) was synthesized
(Twist Bioscience, USA) with attB attachment sites for subsequent gateway cloning into pDONRZeo
(Invitrogen, USA) and recombination with pB7WG2 (VIB, Ghent). To generate CRISPR-Cas9 mutants,
appropriate target sites (two per gene of interest) were designed using the software tool chopchop
(https://chopchop.cbu.uib.no/). Individual guide RNA constructs containing gene specific target sites were
synthesized (Twist Bioscience, USA) and subsequently stacked in a GoldenGate-adapted pUC18-based
vector (45). To generate *cep6x*, *cepr1/2*^AEQ and *rlk7/iku2*^AEQ, 12, 4 and 4 target site-containing gRNA
constructs were stacked, respectively (Table 2). Together with FastRed-pRPS5::Cas9, higher order gRNA
stacks were subsequently cloned into pICSL4723 for in planta expression (46).

To generate the *pCEP4::NLS-3xmVenus*, *pCEPR1::NLS-3xmVenus* and *pCEPR2::NLS-3xmVenus*
reporter constructs, 1000, 1696 and 2788 bp fragments upstream of the start codon, respectively, were
amplified from genomic DNA and assembled together with the sequence coding for the nuclear localization
signal of SV40 large T antigen followed by 3 consecutive mVenus YFP fluorophores (47) into a GoldenGate-

modified pCB302 binary vector for plant expression. For p*RLK7* the 1957 bp promoter sequence upstream from the start codon was amplified with primers containing attB attachment sites for subsequent gateway cloning into pDONRZeo (Invitrogen, USA) and recombination with promotor::Venus (YFP)-H2B destination vector (56). All of the generated plant expression constructs were subsequently transformed into *Agrobacterium tumefaciens* strain GV3101 before floral dip transformation of Arabidopsis. All primers used for cloning are listed in Table S3.

Plant material and growth conditions

Arabidopsis Col-0, Col-0^{AEQ} and No-0 were used as wild types for experiments and generation of transgenic lines or CRISPR mutants (48). The *cepr1-3* (GK-467C01), *cepr2-3* (SALK_014533), *rlk7-1* (SALK_056583), *rlk7-3* (SALK_120595), *iku2-4* (Salk_073260) and the novel *cepr2*-4 allele (GK-695D11) were obtained from NASC (UK) (14, 28, 49, 50) . The No-0 *cepr2-1xcepr2-1* was obtained from RIKEN (Japan) (8). T-DNA insertion mutants were genotyped by PCR using T-DNA- and gene-specific primers as listed in Table S3. The lack of *CEPR2* transcript in *cepr2-4* (GK-695D11) was determined by semi-quantitative PCR from cDNA using *cepr2-4* genotyping primers (Fig. S5B). The *cepr1-3/2-4* double mutant was obtained by genetic crossing. The *bak1-5/bkk1* mutant was characterized previously (51). For visualizing tissue-specific *FRK1* expression, a *pFRK1::NLS-3xmVenus* line was used (19). To isolate homozygous CRISPR mutants, pICSL4723 transformant T1 seeds showing red fluorescence were selected and grown on soil before genotyping with gene specific primers and Sanger sequencing (Table 3). Mutants lacking the transgene were identified by loss of fluorescence. To generate the *cep6x p35S::CEP4* lines, pB7WG2 CEP4 was used for floral dip transformation of homozygous *cep6x*.

Plants for physiological assays involving mature plants were vernalized for 2-3 days in the dark at 4°C and later grown in individual pots in environmentally controlled growth rooms (20-21°C, 55% relative humidity, 8 h photoperiod). For seedling-based assays, seeds were sterilized using chlorine gas and grown axenically on ½ Murashige and Skoog (MS) media supplemented with vitamins (Duchefa, Netherlands), 1% sucrose, with or without 0.8% agarose at 22°C and a 16 h photoperiod unless stated otherwise. For experiments using N depleted ½ MS medium, modified MS salts without nitrogen-containing compounds (Duchefa, Netherlands) were used and supplemented with KNO_3/NH_4NO_3 to achieve 100% N (KNO_3 9.395 mM, NH_4NO_3 10.305 mM) and 10% N (KNO_3 0.9395 mM, NH_4NO_3 1.0305 mM) conditions. To keep the ionic strength equal in 10% N conditions, the medium was supplemented with 90% KCl (8.455 mM).

Grafting

Arabidopsis seedlings were grown vertically on ½ MS agar medium without sucrose in short day conditions seven days before grafting. Grafting was performed aseptically under a stereo microscope as previously described (52). Vertically mounted plates with reciprocally grafted seedlings were returned to short-day conditions for 10 days. Healthy seedlings were transferred to soil.

Imaging and microscopy

Confocal laser-scanning microscopy was performed using a Leica TCS SP5 (Leica, Germany) microscope (with Leica Application Suite X 3.7.4.23463) and all the pictures were taken with a 20x water immersion objective. For the mVenus fluorophore, pictures were imaged with argon laser excitation at 514nm and a detection window of 525-535 nm. Propidium iodide was visualized using DPSS 561 laser emitting at 561 nm with a detection window of 575-590 nm. For imagining the expression pattern of NLS-3xmVenus or Venus (YFP)-H2B under the control of different promoters (*pCEP4*, *pCEPR1*, *pCEPR2* and *pRLK7*), vertically-grown seedlings were stained with propidium iodide immediately before microscopic analysis. For imagining the *pFRK1::NLS-3xmVenus* reporter line, 12-day old seedlings were transferred to a 24-well-plate containing ddH$_2$O with or without (mock) peptides in the indicated concentrations. For comparison of *FRK1* promoter activity, seedlings were analysed by confocal microscopy using identical laser intensities and interval/number of slices for Z stack projection 16 h after treatment.

Calcium influx assay

Apoaequorin-expressing liquid-grown eight-day-old seedlings were transferred individually to a 96-well plate containing 100 µl of 5 µM coelenterazine-h (PJK Biotech, Germany) and incubated in the dark overnight. Luminescence was measured using a plate reader (Luminoskan Ascent 2.1, Thermo Fisher Scientific, USA). Background luminescence was recorded by scanning each well 12 times at 10 s intervals, before adding a 25 µl elicitor solution to the indicated final concentration. Luminescence was recorded for 30 min at the same interval. The remaining aequorin was discharged using 2 M CaCl$_2$, 20% ethanol. The values for cytosolic Ca^{2+} concentrations ([Ca^{2+}]$_{cyt}$) were calculated as luminescence counts per second relative to total luminescence counts remaining (L/L$_{max}$).

Ethylene measurement

Leaf discs (4 mm in diameter) from four-to-five-week-old soil-grown Arabidopsis were recovered overnight in ddH$_2$O. Three leaf discs per sample were transferred to a glass vial containing 500 µl of ddH$_2$O before adding ddH$_2$O (mock) or peptides to the indicated final concentration. Glass vials were capped with a rubber lid and incubated under gentle agitation for 3.5 h. One mL of the vial headspace was extracted with a syringe and injected into a Varian 3300 gas chromatograph (Varian, USA) to measure ethylene.

MAPK activation

Five-day old Arabidopsis seedlings growing on ½ MS agar plates, were transferred into a 24-well plate containing liquid medium for seven days. 24 h before the experiment, seedlings were equilibrated in a fresh ½ MS medium. For N depletion experiments modified ½ MS containing 100% N or 10% N supplemented with 90% KCl was used. MAPK activation was elicited by adding the peptides to the indicated concentrations. Six seedlings per sample were harvested, frozen in liquid nitrogen and homogenized using a tissue lyser (Qiagen, Germany). Proteins were extracted using a buffer containing 50 mM Tris-HCl (pH 7.5), 50 mM NaCl, 10% glycerol, 5 mM DTT, 1% protease inhibitor cocktail, 1 mM phenylmethylsulfonyl

fluoride (PMSF), 1% IGEPAL, 10 mM EGTA, 2 mM NaF, 2 mM Na_3VO_4, 2 mM Na_2MoO_4, 15 mM ß-
Glycerophosphate and 15 mM p-nitrophenylphosphate before analysis by SDS-PAGE and western blot.
Phosphorylated MAPKs were detected by α-p44/42 antibodies (Cell Signaling, USA).

<u>Seedling growth inhibition</u>
Arabidopsis seedlings were grown for five days on ½ MS agar plates, before transfer of individual seedlings
into each well of a 48-well plate containing liquid medium with or without elicitors in the indicated
concentration. After seven-day treatment, fresh weight of individual seedlings was measured.

<u>Pathogen growth assay</u>
Pseudomonas syringae pv. *tomato* (*Pto)* DC3000 and *Pto* lacking the effector molecular coronatine *Pto*[COR-]
were grown on King's B agar plates containing 50 µg/mL rifampicin and 50 µg/mL kanamycin at 28ºC. After
two-to-three days bacteria were resuspended in ddH_2O containing 0.04% Silwet L77 (Sigma Aldrich, USA).
The bacterial suspension was adjusted to an OD_{600} = 0.2 (10^8 cfu/mL) for *Pto*[COR-] or OD_{600} = 0.02 (10^7
cfu/mL) for *Pto* DC3000 before spray inoculating four-to-five-week-old plants. For peptide-induced
resistance in local tissues, ddH_2O (mock), flg22 (100 nM) and/or CEP4 (1 µM) were syringe-infiltrated into
mature leaves. After 24 h, *Pto* DC3000 (OD_{600} = 0.0002, 10^5 cfu/mL) was syringe-infiltrated into pre-treated
leaves and incubated for three days before determining bacterial counts. For CEP4-induced resistance in
systemic tissues, CEP4 (1 µM) or ddH_2O (mock) were syringe-infiltrated into the first two true leaves of
young three-to-four-week-old Arabidopsis. After three days, *Pto* DC3000 (OD_{600}=0.0002, 10^5 cfu/mL) was
syringe-infiltrated into leaves three and four of the pre-treated plants. Bacterial counts were determined four
days after infection.

<u>Systemic acquired resistance</u>
SAR experiments were performed as previously described (53). Briefly, plants were cultivated in a mixture
of substrate (Floradur) and silica sand in a 5:1 ratio under short day (SD) conditions (10 h) in a growth
chamber at 22 °C /18 °C (day/night) with a light intensity of 100 µmol $m^{-2}s^{-1}$, and 70% relative humidity
(RH). SAR assays were performed using *Pto* DC3000 and *Pto* AvrRpm1. Bacteria were grown on NYGA
media (0.5% peptone, 0.3% yeast extract, 2% glycerol, 1.8% agar, 50 µg/mL kanamycin, 50 µg/mL
Rifampicin) at 28° C before infiltration. Freshly grown *Pto* avrRpm1 was diluted in 10 mM $MgCl_2$ (to reach
a final concentration of $1×10^6$ cfu/mL) and syringe-infiltrated in the first two true leaves of four-and-a-half-
week-old plants. Concurrently, 10 mM $MgCl_2$ was applied to a separate set of plants as the mock control
treatment. Three days after *Pto* avrRpm1 infiltration, plants were challenged in their 3rd and 4th leaves with
Pto DC3000 ($1×10^5$ cfu/mL). Bacterial titers were determined four days after *Pto* DC3000 infection.

<u>Gene expression analysis</u>
For seedlings-based assays, 12-day-old liquid-grown seedlings were equilibrated in fresh medium for 24 h
before treatment with the indicated peptides. For adult plants, four-to-five-week-old Arabidopsis leaves

were syringe-infiltrated with ddH$_2$O (mock), flg22 (1 µM) or *Pto* DC3000 (OD$_{600}$ = 0.001, 5x10^5 cfu/mL) and

incubated for 24 h. All samples for RT-qPCR analysis were harvested at the indicated time points, frozen

in liquid nitrogen and homogenized using a tissue lyser (Qiagen, Germany). Total RNA was isolated using

TRIzol reagent (Roche, Switzerland) and purified using Direct-zol™ RNA Miniprep Plus kit (Zymo

Research, Germany). 2 µg of the total RNA was digested with DNase I and reverse transcribed with oligo

(dT)18 and Revert Aid reverse transcriptase. RT-qPCR experiments were performed using Maxima SYBR

green mix (Thermo Fisher Scientific, USA) with the AriaMx Real-Time PCR system (Agilent Technologies,

USA). Expression levels of all tested genes were normalized to the house-keeping gene *Ubiquitin 5* (*UBQ5*).

Sequences of all primers used for RT-qPCR analysis are found in Table S3.

<u>Expression and purification of recombinant receptor ectodomains</u>

Spodoptera frugiperda codon-optimized synthetic genes (Invitrogen GeneArt), coding for Arabidopsis

CEPR1 (residues 23 to 592), CEPR2 (residues 32 to 620) and RLK7 (residues 29 to 608) were cloned into

a modified pFastBAC vector (Geneva Biotech) providing a 30K signal peptide (54), a C-terminal TEV

(tobacco etch virus protease) cleavable site and a StrepII-9xHis affinity tag. For protein expression,

Trichoplusia ni Tnao38 cells (55) were infected with CEPR1, CEPR2 or RLK7 virus with a multiplicity of

infection (MOI) of 3 and incubated one day at 28°C and two days at 21°C at 110 rounds per minute (rpm).

The secreted proteins were purified by Ni^{2+} (HisTrap excel, Cytiva, equilibrated in 25 mM KP$_i$ pH 7.8 and

500 mM NaCl) followed by Strep (Strep-Tactin Superflow high-capacity, IBA Lifesciences, equilibrated in

25 mM Tris pH 8.0, 250 mM NaCl, 1 mM EDTA) affinity chromatography. All proteins were incubated with

TEV protease to remove the tags. Proteins were purified by SEC on a Superdex 200 Increase 10/300 GL

column (Cytiva, USA) equilibrated in 20 mM citrate pH 5.0, 150 mM NaCl and further concentrated using

Amicon Ultra concentrators from Millipore (Merck, Germany) with a 30,000 Da molecular weight cut-off.

Purity and structural integrity of the different proteins were assessed by SDS-PAGE.

<u>Analytical size-exclusion (SEC) chromatography</u>

Analytical SEC experiments were performed using a Superdex 200 Increase 10/300 GL column (GE, USA).

The columns were pre-equilibrated in 20 mM citric acid pH 5, 150 mM NaCl. 150 µg of CEPR1, CEPR2 and

RLK7 were injected sequentially onto the column and eluted at 0.5 mL/min. Ultraviolet absorbance (UV) at

280 nm was used to monitor the elution of the proteins. The peak fractions were analyzed by SDS-PAGE

followed by Coomassie blue staining.

<u>Isothermal titration calorimetry (ITC)</u>

Experiments were performed at 25°C on a MicroCal PEAQ-ITC (Malvern Instruments, UK) using a 200 µL

standard cell and a 40 µL titration syringe. CEP1, CEP4, CEP4^{scr} and PIP1 peptides were dissolved in the

SEC buffer to match the receptor protein. A typical experiment consisted of injecting 1 µL of a 300 µM

solution of the peptide into 30 µM CEPR2 or RLK7 solution in the cell at 150 s intervals. ITC data were

corrected for the heat of dilution by subtracting the mixing enthalpies for titrant solution injections into

protein-free ITC buffer. Experiments were done in duplicates and data were analyzed using the MicroCal

PEAQ-ITC Analysis Software provided by the manufacturer. The N values were fitted to 1 in the analysis.

<u>Synthetic peptides</u>

The flg22 peptide was kindly provided by Dr. Justin Lee (IPB Halle). Other peptides were synthesized by

Pepmic (China) with at least 90% purity and dissolved in ddH$_2$O. Sequences of all the synthetic peptides

can be found in Table S1.

<u>Quantification and statistical analysis</u>

Statistical analyses were performed using GraphPad Prism (Version 9.4.1). Sample size, p-values and

statistical methods employed are described in the respective figure legends.

Data and material availability

All data are available in the main text or the supplementary materials. All newly generated mutant lines are

available upon request to M.S.

References

1. B. P. M. Ngou, P. Ding, J. D. G. Jones, Thirty years of resistance: Zig-zag through the plant
immune system. *Plant Cell*. **34**, 1447–1478 (2022).

2. A. A. Gust, R. Pruitt, T. Nürnberger, Sensing danger: Key to activating plant immunity. *Trends
Plant Sci.* **22**, 779–791 (2017).

3. J. Rzemieniewski, M. Stegmann, Regulation of pattern-triggered immunity and growth by
phytocytokines. *Curr. Opin. Plant Biol.* **68**, 102230 (2022).

4. K. Gully, S. Pelletier, M.-C. C. Guillou, M. Ferrand, S. Aligon, I. Pokotylo, A. Perrin, E. Vergne, M.
Fagard, E. Ruelland, P. Grappin, E. Bucher, J.-P. P. Renou, S. Aubourg, The SCOOP12 peptide
regulates defense response and root elongation in Arabidopsis thaliana. *J. Exp. Bot.* **70**, 1349–
1365 (2019).

5. J. Rhodes, A.-O. Roman, M. Bjornson, B. Brandt, P. Derbyshire, M. Wyler, M. Schmid, F. L. H.
Menke, J. Santiago, C. Zipfel, Perception of a conserved family of plant signalling peptides by the
receptor kinase HSL3. *Elife*. **11**, e74687 (2022).

6. Z. Liu, S. Hou, O. Rodrigues, P. Wang, D. Luo, K. Nomura, C. Yin, H. Wang, W. Zhang, K. Zhu-
salzman, Phytocytokine signalling reopens stomata in plant immunity and water loss. *Nature*. **605**,
332–339 (2022).

7. M. Stegmann, P. Zecua-Ramirez, C. Ludwig, H.-S. Lee, B. Peterson, Z. L. Nimchuk, Y. Belkhadir,
R. Hückelhoven, RGI-GOLVEN signalling promotes cell surface immune receptor abundance to
regulate plant immunity. *EMBO Rep.* **23**, e53281 (2022).

8. R. Tabata, K. Sumida, T. Yoshii, K. Ohyama, H. Shinohara, Y. Matsubayashi, Perception of root-
derived peptides by shoot LRR-RKs mediates systemic N-demand signaling. *Science*. **346**, 343–
346 (2014).

9. F. Takahashi, T. Suzuki, Y. Osakabe, S. Betsuyaku, Y. Kondo, N. Dohmae, H. Fukuda, K.
Yamaguchi-Shinozaki, K. Shinozaki, A small peptide modulates stomatal control via abscisic acid
in long-distance signalling. *Nature*. **556**, 235–238 (2018).

10.   H. Zhou, F. Xiao, Y. Zheng, G. Liu, Y. Zhuang, Z. Wang, Y. Zhang, J. He, C. Fu, H. Lin, PAMP-
INDUCED SECRETED PEPTIDE 3 modulates salt tolerance through RECEPTOR-LIKE KINASE 7
in plants. *Plant Cell*. **34**, 927–944 (2021).

11.   K. Ohyama, M. Ogawa, Y. Matsubayashi, Identification of a biologically active, small, secreted
peptide in Arabidopsis by in silico gene screening, followed by LC-MS-based structure analysis.
Plant J. **55**, 152–160 (2008).

12.   M. Taleski, N. Imin, M. A. Djordjevic, CEP peptide hormones: Key players in orchestrating
nitrogen-demand signalling, root nodulation, and lateral root development. *J. Exp. Bot.* **69**, 1829–
1836 (2018).

13.   C. Delay, K. Chapman, M. Taleski, Y. Wang, S. Tyagi, Y. Xiong, N. Imin, M. A. Djordjevic, CEP3
levels affect starvation-related growth responses of the primary root. *J. Exp. Bot.* **70**, 4763–4774
(2019).

14.   K. Chapman, M. Taleski, H. A. Ogilvie, N. Imin, M. A. Djordjevic, CEP-CEPR1 signalling inhibits
the sucrose-dependent enhancement of lateral root growth. *J. Exp. Bot.* **70**, 3955–3967 (2019).

15.   S. Roy, L. M. Müller, A rulebook for peptide control of legume – microbe endosymbioses. *Trends
Plant Sci.* **27**, 870–889 (2022).

16.   B. Li, S. Jiang, X. Yu, C. Cheng, S. Chen, Y. Cheng, J. S. Yuan, D. Jiang, P. He, L. Shan,
Phosphorylation of Trihelix Transcriptional Repressor ASR3 by MAP KINASE4 Negatively
Regulates Arabidopsis Immunity. *Plant Cell*. **27**, 839–856 (2015).

17.   I. Roberts, S. Smith, B. De Rybel, J. Van Den Broeke, W. Smet, S. De Cokere, M. Mispelaere, I.
De Smet, T. Beeckman, The CEP family in land plants: Evolutionary analyses, expression studies,
and role in Arabidopsis shoot development. *J. Exp. Bot.* **64**, 5371–5381 (2013).

18.   C. Delay, N. Imin, M. A. Djordjevic, CEP genes regulate root and shoot development in response
to environmental cues and are specific to seed plants. *J. Exp. Bot.* **64**, 5383–5394 (2013).

19.   L. Poncini, I. Wyrsch, V. D. Tendon, T. Vorley, T. Boller, N. Geldner, J. P. Métraux, S. Lehmann, In
roots of Arabidopsis thaliana, the damage-associated molecular pattern AtPep1 is a stronger
elicitor of immune signalling than flg22 or the chitin heptamer. *PLoS One*. **12**, e0185808 (2017).

20.   J. Santiago, B. Brandt, M. Wildhagen, U. Hohmann, L. A. Hothorn, M. A. Butenko, M. Hothorn,
Mechanistic insight into a peptide hormone signaling complex mediating floral organ abscission.
Elife. **5**, 1–19 (2016).

21.   J. Rhodes, H. Yang, S. Moussu, F. Boutrot, J. Santiago, C. Zipfel, Perception of a divergent family
of phytocytokines by by the Arabidopsis receptor kinase MIK2. *Nat. Commun.* **12**, 5494 (2021).

22.   P. Qian, W. Song, T. Yokoo, A. Minobe, G. Wang, T. Ishida, S. Sawa, J. Chai, T. Kakimoto, The
CLE9/10 secretory peptide regulates stomatal and vascular development through distinct
receptors. *Nat. Plants*. **4**, 1071–1081 (2018).

23.   S. Hou, D. Liu, S. Huang, D. Luo, Z. Liu, Q. Xiang, P. Wang, R. Mu, Z. Han, S. Chen, J. Chai, L.
Shan, P. He, The Arabidopsis MIK2 receptor elicits immunity by sensing a conserved signature
from phytocytokines and microbes. *Nat. Commun.* **12**, 5494 (2021).

24.   H. H. Breitenbach, M. Wenig, F. Wittek, L. Jordá, A. M. Maldonado-Alconada, H. Sarioglu, T.
Colby, C. Knappe, M. Bichlmeier, E. Pabst, D. Mackey, J. E. Parker, A. Corina Vlot, Contrasting
roles of the apoplastic aspartyl protease APOPLASTIC, ENHANCED DISEASE
SUSCEPTIBILITY1-DEPENDENT1 and LEGUME LECTIN-LIKE PROTEIN1 in Arabidopsis
systemic acquired resistance. *Plant Physiol*. **165**, 791–809 (2014).

25.   M. Luo, E. S. Dennis, F. Berger, W. J. Peacock, A. Chaudhury, MINISEED3 (MINI3), a WRKY
family gene, and HAIKU2 (IKU2), a leucine-rich repeat (LRR) KINASE gene, are regulators of
seed size in Arabidopsis. *Proc. Natl Acad. Sci. USA*. **102**, 17531–17536 (2005).

26.   K. Toyokura, T. Goh, H. Shinohara, A. Shinoda, Y. Kondo, Y. Okamoto, T. Uehara, K. Fujimoto, Y.
Okushima, Y. Ikeyama, K. Nakajima, T. Mimura, M. Tasaka, Y. Matsubayashi, H. Fukaki, Lateral
inhibition by a peptide hormone-receptor cascade during Arabidopsis lateral root founder cell

472    formation. *Dev. Cell*. **48**, 64-75.e5 (2019).

473 27. D. Pitorre, C. Llauro, E. Jobet, J. Guilleminot, J. P. Brizard, M. Delseny, E. Lasserre, RLK7, a
474    leucine-rich repeat receptor-like kinase, is required for proper germination speed and tolerance to
475    oxidative stress in Arabidopsis thaliana. *Planta*. **232**, 1339–1353 (2010).

476 28. S. Hou, X. Wang, D. Chen, X. Yang, M. Wang, D. Turrà, A. Di Pietro, W. Zhang, The secreted
477    peptide PIP1 amplifies immunity through Receptor-Like Kinase 7. *PLoS Pathog.* **10**, e1004331
478    (2014).

479 29. S. Hou, H. Shen, H. Shao, PAMP-induced peptide 1 cooperates with salicylic acid to regulate
480    stomatal immunity in Arabidopsis thaliana. *Plant Signal. Behav.* **14**, 1666657 (2019).

481 30. M. Fagard, A. Launay, G. Clément, J. Courtial, A. Dellagi, M. Farjad, A. Krapp, M. C. Soulié, C.
482    Masclaux-Daubresse, Nitrogen metabolism meets phytopathology. *J. Exp. Bot.* **65**, 5643–5656
483    (2014).

484 31. C. Verly, A. C. R. Djoman, M. Rigault, F. Giraud, L. Rajjou, M. E. Saint-Macary, A. Dellagi, Plant
485    defense stimulator mediated defense activation is affected by nitrate fertilization and
486    developmental stage in Arabidopsis thaliana. *Front. Plant Sci.* **11**, 583 (2020).

487 32. D. Zhuo, M. Okamoto, J. J. Vidmar, A. D. M. Glass, Regulation of a putative high-affinity nitrate
488    transporter (Nrt2;1At) in roots of Arabidopsis thaliana. *Plant J.* **17**, 563–568 (1999).

489 33. A. Roman, P. Jimenez-sandoval, S. Augustin, C. Broyart, L. A. Hothorn, J. Santiago, HSL1 and
490    BAM1/2 impact epidermal cell development by sensing distinct signaling peptides. *Nat. Commun.*
491    **13**, 876 (2022).

492 34. M. Stegmann, J. Monaghan, E. Smakowska-Luzan, H. Rovenich, A. Lehner, N. Holton, Y.
493    Belkhadir, C. Zipfel, The receptor kinase FER is a RALF-regulated scaffold controlling plant
494    immune signaling. *Science*. **355**, 287–289 (2017).

495 35. J. Gronnier, C. M. Franck, M. Stegmann, T. DeFalco, A. Abarca, M. Von Arx, K. Duenser, W. Lin,
496    Z. Yang, J. Kleine-Vehn, C. Ringli, C. Zipfel, Regulation of immune receptor kinases plasma
497    membrane nanoscale landscape by a plant peptide hormone and its receptors. *Elife*. **11**, e74162
498    (2022).

499 36. J. Dindas, T. A. DeFalco, G. Yu, L. Zhang, P. David, M. Bjornson, M.-C. Thibaud, V. Custódio, G.
500    Castrillo, L. Nussaume, A. P. Macho, C. Zipfel, Direct inhibition of phosphate transport by immune
501    signaling in Arabidopsis. *Curr. Biol.* **32**, 488-495.e5 (2022).

502 37. S. Roy, M. Griffiths, I. Torres-Jerez, B. Sanchez, E. Antonelli, D. Jain, N. Krom, S. Zhang, L. M.
503    York, W. R. Scheible, M. Udvardi, Application of Synthetic Peptide CEP1 Increases Nutrient
504    Uptake Rates Along Plant Roots. *Front. Plant Sci.* **12**, 793145 (2022).

505 38. A. Jacquot, V. Chaput, A. Mauries, Z. Li, P. Tillard, C. Fizames, P. Bonillo, F. Bellegarde, E.
506    Laugier, V. Santoni, S. Hem, A. Martin, A. Gojon, W. Schulze, L. Lejay, NRT2.1 C-terminus
507    phosphorylation prevents root high affinity nitrate uptake activity in Arabidopsis thaliana. *New*
508    *Phytol.* **228**, 1038–1054 (2020).

509 39. L. Zhang, Z. Yu, Y. Xu, M. Yu, Y. Ren, S. Zhang, G. Yang, J. Huang, K. Yan, C. Zheng, C. Wu,
510    Regulation of the stability and ABA import activity of NRT1.2/NPF4.6 by CEPR2-mediated
511    phosphorylation in Arabidopsis. *Mol. Plant*. **14**, 633–646 (2021).

512 40. M. Taleski, K. Chapman, N. Imin, M. A. Djordjevic, M. Groszmann, The peptide hormone receptor
513    CEPR1 functions in the reproductive tissue to control seed size and yield. *Plant Physiol.* **182**, 620–
514    636 (2020).

515 41. C. Furumizu, A. K. Krabberød, M. Hammerstad, R. M. Alling, M. Wildhagen, S. Sawa, R. B. Aalen,
516    The sequenced genomes of nonflowering land plants reveal the innovative evolutionary history of
517    peptide signaling. *Plant Cell*. **33**, 2915–2934 (2021).

518 42. C. Laffont, A. Ivanovici, P. Gautrat, M. Brault, M. A. Djordjevic, F. Frugier, The NIN transcription
519    factor coordinates CEP and CLE signaling peptides that regulate nodulation antagonistically. *Nat.*
520    *Commun.* **11**, 3167 (2020).

43.   Z. Luo, J. Wang, F. Li, Y. Lu, Z. Fang, M. Fu, K. S. Mysore, J. Wen, J. Gong, J. D. Murray, F. Xie,
The small peptide CEP1 and NIN-like protein NLP1 regulate NRT2.1 to mediate root nodule
formation across nitrate concentrations. *Plant Cell*. **in press** (2022).

44.   C. Engler, R. Kandzia, S. Marillonnet, A one pot, one step, precision cloning method with high
throughput capability. *PLoS One*. **3**, e3647 (2008).

45.   B. Castel, L. Tomlinson, F. Locci, Y. Yang, J. D. G. Jones, Optimization of T-DNA architecture for
Cas9-mediated mutagenesis in Arabidopsis. *PLoS One*. **14**, e0204778 (2019).

46.   J. E. M. Vermeer, D. von Wangenheim, M. Barberon, Y. Lee, E. H. K. Stelzer, A. Maizel, N.
Geldner, A spatial accommodation by neighboring cells is required for organ initiation in
Arabidopsis. *Science*. **343**, 178–183 (2014).

47.   M. Somssich, A. Bleckmann, R. Simon, Shared and distinct functions of the pseudokinase
CORYNE (CRN) in shoot and root stem cell maintenance of Arabidopsis. *J. Exp. Bot.* **67**, 4901–
4915 (2016).

48.   M. R. Knight, A. K. Campbellt, S. M. Smith, A. J. Trewavas, Transgenic plant aequorin reports the
effects of touch and cold-shock and elicitors on cytoplasmic calcium. *Nature*. **314**, 524–526
(1991).

49.   I. Dimitrov, F. E. Tax, Lateral root growth in Arabidopsis is controlled by short and long distance
signaling through the LRR RLKs XIP1/CEPR1 and CEPR2. *Plant Signal. Behav.* **13**, e1489667
(2018).

50.   A. Fatihi, A. M. Zbierzak, P. Dörmann, Alterations in seed development gene expression affect
size and oil content of Arabidopsis seeds. *Plant Physiol.* **163**, 973–985 (2013).

51.   M. Roux, B. Schwessinger, C. Albrecht, D. Chinchilla, A. Jones, N. Holton, F. G. Malinovsky, M.
Tör, S. de Vries, C. Zipfel, The Arabidopsis leucine-rich repeat receptor-like kinases BAK1/SERK3
and BKK1/SERK4 are required for innate immunity to hemibiotrophic and biotrophic pathogens.
Plant Cell. **23**, 2440–55 (2011).

52.   C. W. Melnyk, Grafting with arabidopsis thaliana. *Methods Mol. Biol.* **1497**, 9–18 (2017).

53.   M. Wenig, A. Ghirardo, J. H. Sales, E. S. Pabst, H. H. Breitenbach, F. Antritter, B. Weber, B.
Lange, M. Lenk, R. K. Cameron, J. P. Schnitzler, A. C. Vlot, Systemic acquired resistance
networks amplify airborne defense cues. *Nat. Commun.* **10**, 3183 (2019).

54.   M. Futatsumori-Sugai, K. Tsumoto, Signal peptide design for improving recombinant protein
secretion in the baculovirus expression vector system. *Biochem. Biophys. Res. Commun.* **391**,
931–935 (2010).

55.   Y. Hashimoto, S. Zhang, S. Zhang, Y. R. Chen, G. W. Blissard, Correction: BTI-Tnao38, a new
cell line derived from *Trichoplusia ni*, is permissive for AcMNPV infection and produces high levels
of recombinant proteins. *BMC Biotechnol.* **12** (2012).

Author contributions

Conceptualization: MS; Investigation: JR, HL, HKL, CB, SN, CW, JM, ZC, VOL, MS; Funding acquisition:

MS, JA, RH, ACV, MAD; Project administration: MS; Supervision: MS, JR, ACV, JS, MAD, RH; Writing –

original draft: MS, JR; Writing – review & editing: HL, HKL, CB, SN, CW, JM, VOL, MAD, ACV, RH, JS.

Competing interests

Authors declare that there are no competing interests.

Acknowledgments

We thank Stefanie Ranf for providing GoldenGate vectors for molecular cloning and Ulrich Hammes for

advice in grafting experiments. This work was funded by the following agencies: Deutsche

Forschungsgemeinschaft (DFG) grants STE2448/3-1 (MS and JR), STE2448/4-1 (MS and HL) and SFB924

TP B06 (ACV), the Technical University of Munich (MS, JR, RH, ZC, CW and JM), the Research Council

of Norway, Grant 230849 (VOL), Australian Research Council, Grant DP200101885 (MAD), the University

of Lausanne (CB), the Swiss National Science Foundation grants no. 310030_204526 and the European

Research Council (ERC) grant agreement no. 716358 (JS, HKL).

Figures

Figure 1: CEPs induce immune responses and are important determinants of plant immunity. A)
Colony forming units (cfu) of Pto^{COR-} 3 days post inoculation (3 dpi) upon spray infection; n = 16 pooled
from four experiments ± SD (one-way ANOVA, Tukey post-hoc test; a-b, p<0.01; a/b-c p<0.0001) **B)**
Kinetics of cytosolic calcium concentrations ($[Ca^{2+}]_{cyt}$) in Col-0^{AEQ} seedlings upon ddH$_2$O or CEP4 (1 µM)
treatment; n = 8, ± SD. **C)** MAPK activation in Col-0 upon CEP4 (1 µM) or flg22 (100 nM) treatment. Western
blots were probed with α-p44/42. CBB = Coomassie brilliant blue. **D)** Ethylene concentration in Col-0 leaf
discs 3.5 h upon mock (ddH$_2$O) or CEP4 (1 µM) treatment; n = 30 pooled from six experiments ± SD (two-
tailed Student's t-test, **** p<0.0001). **E)** RT-qPCR of *FRK1* in seedlings upon CEP4 (1 µM) or ddH$_2$O
treatment for 4 h; n = 6 pooled from six experiments ± SD (two-tailed Student's t-test, ** p<0.01).
Housekeeping gene *UBQ5*. **F)** NLS-3xmVenus signal in p*FRK1::NLS-3xmVenus* upon ddH$_2$O or CEP4 (1
µM) treatment for 16 h. Cyan dotted line indicates vasculature. BF = bright field, PI = propidium iodide,
scale bar = 100 µm. **G)** Relative fresh weight of five-day-old seedlings treated with CEP4 (1 µM) for seven
days (n=16, ± SD, **** p<0.0001, two-tailed Student's t-test). **H)** cfu of *Pto* DC3000 (4 dpi) in distal leaves
upon local CEP4 (1 µM) or ddH$_2$O pre-treatment; n = 12 pooled from three experiment ± SD (two-tailed
Student's t-test, *** p=0.0001). **I)** cfu of Pto^{COR-} (3 dpi) upon spray infection; n = 16 pooled from four
experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p≤0.001). **J)** cfu of Pto^{COR-} (3dpi) upon spray
infection; n = 4 ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.05). All experiments were repeated
at least three times in independent biological repeats with similar results.

594
595
Figure 2: CEPR1 and CEPR2 are tissue-specific CEP4 receptors and important determinants of plant
immunity. A) NLS-3xmVenus signal in the indicated lines. Cyan dotted line indicates vasculature. BF =
bright field, PI = propidium iodide, scale bar = 100 μm. **B)** cfu of Pto^{COR-} (3 dpi) upon spray infection. Dotted
line indicates different experiments; n = 12, 21 and 13 pooled from three, five and three experiments,
respectively ± SD (one-way ANOVA, Dunnett's post-hoc test, a-b p<0.0001, two-tailed Student's t-test, A-
B p<0.0001). **C)** Relative fresh weight of five-days-old seedlings treated with CEP4 (1 μM) for seven days;
n = 35-36 pooled from three experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.0001). **D)**
cfu of Pst DC3000 (4 dpi) in distal leaves upon local CEP4 (1 μM) or ddH$_2$O pre-treatment; n = 12 pooled
from three experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.01). **E)** CEP4, **F)** CEP4^{scr}
and **G)** CEP1 were titrated into a solution containing CEPR2^{ECD} in ITC cells. Top: raw data thermogram;
bottom: fitted integrated ITC data curves. **H)** ITC table summarizing CEPR2^{ECD} vs CEP4/CEP4^{scr}/CEP1 as
means ± SD of two experiments. Dissociation constant (K$_d$) indicates receptor-ligand binding affinity. N
indicates reaction stoichiometry (n = 1 for 1:1 interaction). **I)** RT-qPCR of $PR1$ and $FRK1$ in adult leaves
after treatment with flg22 (100 nM) or ddH$_2$O for 24h; n = 4 ± SD. Housekeeping gene $UBQ5$. Different
symbols represent four independent experiments. **J)** cfu of Pto DC3000 (4 dpi) in distal leaves upon local
infection with Pto $avrRPM1$ or 10 mM MgCl$_2$; n = 9 pooled from three independent experiments ± SD (one-
way ANOVA, Tukey post-hoc test, a-b p<0.0001). Experiments in **A-D**, **I**, **J** and **E-G** were repeated at least
three times in independent biological repeats or in two independent technical repeats, respectively, with
similar results.
615
616

617
618
Figure 3: RLK7 is an additional CEP4 receptor. A) ([Ca^{2+}]$_{cyt}$) kinetics in seedlings upon CEP4 (1 μM)
treatment; n = 6 ± SD. **B)** MAPK activation upon CEP4 (1 μM) treatment for 10 min. Western blots were
probed with α-p44/42. CBB = Coomassie brilliant blue. **C)** RT-qPCR of *FRK1* in seedlings upon CEP4 (1
μM) or ddH$_2$O treatment for 10 min; n = 3, ± SD. Housekeeping gene *UBQ5*. **D)** MAPK activation upon
CEP4 (1 μM) treatment for 10 min. Western blots were probed with α-p44/42. CBB = Coomassie brilliant
blue. **E)** [Ca^{2+}]$_{cyt}$ kinetics in seedlings upon CEP4 treatment (1 μM); n = 6, ± SD. **F)** RT-qPCR analysis of
FRK1 in seedlings upon CEP4 (1 μM) or ddH$_2$O treatment for 4 h; n = 3 ± SD. Housekeeping gene *UBQ5*.
G) Relative fresh weight of five-day-old seedlings treated with CEP4 (1 μM) for seven days; n = 48 pooled
from four experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b/c/bc p<0.0001, b-c p<0.05). **H)**
[Ca^{2+}]$_{cyt}$ kinetics in seedlings upon CEP1 (10 μM) treatment; n = 4, ± SD. **I)** CEP4, **J)** PIP1 and **K)** CEP1
(**K**) were titrated into a solution containing CEPR2^{ECD} in ITC cells. Top: raw data thermogram; bottom: fitted
integrated ITC data curves. **I)** ITC table summarizing RLK7^{ECD} vs CEP4/PIP1/CEP1 as mean ± SD of two
experiments. K$_d$ indicates receptor-ligand binding affinity. N indicates reaction stoichiometry. **M)** NLS-
3xmVenus signal in p*RLK7*::*Venus-H2B*. Cyan-dotted line represents vasculature. BF = bright field, PI =
propidium iodide, scale bar = 100 μm. Different symbols in **C** and **F** represent three independent
experiments. Experiments in **A-H**, **M** and **I-K** were repeated at least three times in independent biological
repeats or in two independent technical repeats, respectively, with similar results.
636

637
638
Figure 4: CEPs promote FLS2 signaling during N starvation. A) Ethylene concentration in Col-0 leaf
discs 3.5 h after CEP4 (1 µM) and/or flg22 (500 nM) treatment; n = 30, seven pooled experiments ± SD
(two-tailed Student's t-test, ** p<0.05, *** p = 0.0001). **B)** cfu of *Pst* DC3000 (3dpi) in Col-0 leaves upon
ddH$_2$O, CEP4 (1 µM) and/or flg22 (100 nM) pretreatment; n = 4 ± SD (one-way ANOVA, Tukey post hoc,
a-b p<0.05). **C)** MAPK activation upon flg22 (100 nM) treatment. Western blots were probed with α-p44/42.
CBB = Coomassie brilliant blue. **D)** RT-qPCR of *FRK1* upon flg22 (100 nM) treatment for 4 h; n = 3 ± SD.
Housekeeping gene *UBQ5* **E)** RT-qPCR of *CEP4* after 24 h transfer of Col-0 seedlings to normal N/low N
medium (NN/LN); n = 3 ± SD. Housekeeping gene *UBQ5*. **F)** MAPK activation upon flg22 (100 nM)
treatment after 24 h transfer of seedlings to NN or LN. Western blots were probed with α-p44/42. CBB =
Coomassie brilliant blue. **G)** RT-qPCR of *FRK1* upon flg22 (100 nM) treatment for 4h after 24 h transfer of
seedlings to NN or LN; n = 3 ± SD. Housekeeping gene *UBQ5*; n = 3 ± SD (one-way ANOVA, Tukey post
hoc, a-b p<0.01, b-c ≤ 0,001, a-c <0.0001). **H)** RT-qPCR of *NRT2.1* upon flg22 (100 nM) treatment of
seedlings for 4h after 24 h transfer to NN or LN; n = 3 ± SD. Housekeeping gene *UBQ5*. Different symbols
in figure **D-E** and **G-H** represent independent biological experiments. All experiments were repeated at least
three times in independent biological repeats with similar results.
654

Supplementary figures

Figure S1: CEP4 expression is differentially regulated after flg22 perception.
A) *CEP4* is downregulated after flg22 treatment. Data were obtained using Genevestigator software and are based on the AT_mRNASeq_ARABI_GL-1 data set. **B)** *CEP4* expression in roots is weakly downregulated after flg22 treatment. Col-0 seedlings were treated with flg22 (100 nM) for the indicated time after which roots were harvested for RT-qPCR analysis of *CEP4* abundance. Expression of *CEP4* was normalized to *UBQ5*. Different symbols represent three independent experiments ± SD.

Figure S2: Characterization of *35S::CEP4*, *cep6x* and *cep6x 35S::CEP4*.

A) *CEP4* transcript levels in two independent *CEP4* overexpression lines. Expression of *CEP4* was normalized to *UBQ5* and is shown as a fold induction compared to Col-0. **B)** Characterization of *cep6x*. Schematic diagram of *CEP1*, *CEP2*, *CEP3*, *CEP4*, *CEP6* and *CEP9* gene structure and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. CRISPR *cep1* has an 88 bp deletion including the mature peptide sequence which is indispensable for CEP activity. CRISPR *cep2* mutations show two 1 bp insertions. The first insertion causes a frameshift mutation and leads to an early stop codon before the mature peptide sequence. CRISPR *cep3* has a 20 bp deletion followed by a 1 bp insertion. The mutations lead to an early stop codon before the mature peptide sequence. CRISPR *cep4* has a 41 bp deletion which partially cuts out the coding sequence of the mature peptide. CRISPR *cep6* has a 1 bp insertion which leads to a frame-shift mutation and premature stop codon before the peptide sequence. CRISPR *cep9* includes a 1 bp mutation between CEP9.1 and CEP9.2 sequences which leads to an early stop codon, followed by a 60bp deletion starting in the coding sequence of CEP9.5. The specific location and type of mutations for each gene are indicated in the schematics describing the mutants. **C)** Pictures of 5-week-old plants of the indicated genotypes grown on soil; scale bar = 2 cm **D)** Loss of *CEPs* increases Arabidopsis susceptibility to *Pto* DC3000. Bacterial titer three days after leaf spray infection; n = 12 pooled from three experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b, p<0.005). **E)** *CEP4* transcript levels in two independent lines of *cep6x 35S::CEP4*. Expression of *CEP4* was normalized to *UBQ5* and is shown as a fold change to Col-0. Different symbols in **A** and **E** represent four independent biological repeats ± SD.

Figure S3: CEP4 is the strongest inducer of PTI responses on the whole tissue level.

A) Both *CEP4* variants trigger influx of calcium ions. Col-0^{AEQ} seedlings were treated with the indicated concentrations of CEP4 and CEP4-15mer. ($[Ca^{2+}]_{cyt}$) was measured for 30 min. Shown is $\log_{10}$ of total calcium influx; n = 12 pooled from two experiments. **B)** Some CEPs induce Ca^{2+} influx at higher concentrations. Col-0^{AEQ} seedlings were treated with the indicated concentrations of CEPs. ($[Ca^{2+}]_{cyt}$) was measured for 30 min. The average calcium influx triggered by individual CEPs is normalized to mock (ddH$_2$O). Dotted line represents different experiments. Statistical analysis was performed separately for each group. Statistical significance was compared to CEP4^{scr} of the indicated concentration; n = 6 (one-way ANOVA, Dunnett post hoc test, a-b p<0.0001, A-B p<0.001, A-C p<0.0001). All experiments were repeated at least three times with similar results. **C)** CEP4 and CEP1 induce *FRK1* promoter activity in the vasculature. Representative images of NLS-3xmVenus signal in *pFRK1::NLS-3xmVenus* lines upon ddH$_2$O, CEP1 (1 µM), CEP4 (1 µM) or flg22 (10 nM) treatment for 16 h. Maximum projection of Z-stack for mVenus merged with either Z-stacked PI signal or single section BF. Cyan dotted line indicates vasculature. BF = bright field, PI = propidium iodide, scale bar = 100 µm. All experiments were repeated three times in independent biological repeats with similar results.

Figure S4: Shoot-expressed *CEPs* are required for basal immunity against *Pto*.

A) Root-expressed *CEPs* are dispensable for resistance to *Pto*[COR-]. Cfu of *Pto*[COR-] upon spray infection of reciprocally grafted plants. The genotype of the shoot scion is indicated with S, the genotype of the rootstock is indicated with R; n = 17-19 pooled from four experiments ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.0001). **B)** *CEPs* are weakly expressed in the leaf tissue of adult plants. RT-qPCR analysis of *CEP* expression in ddH$_2$O and *Pto* DC3000-inoculated leaves 24 h post-treatment; n = 3, ± SD. Housekeeping gene *UBQ5*. **C)** Representative images of *CEP4* promoter activity. For imaging lateral root emergence zone and the hydathode region, maximum projection of Z-stacks for mVenus is merged with either Z-stacked PI signal or single section BF. To show lack of signal mVenus signal in the vasculature, maximum projection of Z-stacks for mVenus is merged with the same single section of PI or BF showing a single epidermal layer or vasculature. Cyan dotted line indicates vasculature. BF = bright field, PI = propidium iodide, scale bar = 100 µm. Different symbols in **B** represent independent experiments. All experiments were repeated at least three times in independent biological repeats with similar results.

Figure S5: Characterization of *cepr1/2*^{AEQ}, *cepr2-4* and *rlk7/iku2*^{AEQ}.

A) Schematic diagram of the *CEPR1* and *CEPR2* gene structure and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. CRISPR *cepr1* has two 1 bp insertions which lead to a frameshift mutation and an early stop codon. CRISPR *cepr2* has a 3020 bp deletion between the start and stop codon. T-DNA insertion site in *cepr2-4* is indicated. **B)** Characterization of the *cepr2-4* mutant. The *CEPR2* transcript could not be detected in *cepr2-4* mutant in a semi-quantitative RT-PCR analysis. Housekeeping gene *UBQ5*. **C)** Schematic diagram of the *RLK7* and *IKU2* gene structure and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. CRISPR *rlk7* mutant has a 1 bp insertion and CRISPR *iku2* has a 29 bp insertion, followed by a 17 bp deletion. Mutations in both genes lead to a frameshift mutation and an early stop codon. The specific location and type of mutations for each gene are indicated in the schematics describing the mutants.

731
Figure S6: CEPR1 and CEPR2 are similarly required for CEP4-induced seedling growth inhibition.
A) CEP4-induced seedling growth inhibition is abolished in *cepr1-3*. Relative fresh weight of five-day-old
seedlings treated with CEP4 (1 µM) for seven days; n = 42-60 pooled from five experiments ± SD (one-way
ANOVA, Tukey post-hoc test, a-b p<0.0001). **B)** CEP4-induced seedling growth inhibition is reduced and
abolished in *cepr2-3* and *cepr2-4,* respectively. Relative fresh weight of five-day-old seedlings treated with
CEP4 (1 µM) for seven days; n = 46-48 of four pooled experiments ± SD (one –way ANOVA, Tukey post-
hoc test, a-b, p<0.0001; c-a/b p<0.05). **C)** CEP4-induced seedling growth inhibition is abolished in *cepr1-*
1xcepr2-1. Relative fresh weight of five-day-old seedlings treated with CEP4 (1 µM) for seven days; n=16
± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.01). Similar results were obtained in three
independent biological repeats.
742

Figure S7: Quality control of recombinant CEPR1, CEPR2 and RLK7 ectodomains and supplemental ITC data.
A) Analytical size-exclusion chromatography (SEC) experiments show CEPR1^{ECD} aggregates in comparison to single peaks for CEPR2^{ECD} and RLK7^{ECD}. **B)** SDS-PAGE of the ECD peaks of the SEC analysis in **A**. **C)** ITC thermograms of the second technical repeat performed for each ectodomain and peptide analyzed in Fig. 2 (**E-G**) and Fig. 3 (**I-K**).

752
Figure S8: CEPR1 and CEPR2 are dispensable for flg22-induced ethylene accumulation and
growth inhibition.
A) Basal and flg22-triggered ethylene production is higher in *cepr1-3/2-4* mutants. Ethylene concentration
in leaf discs upon ddH_2O or CEP4 (1 µM) for 3.5 h; n = 7 - 8 pooled from two experiments ± SD (one-way
ANOVA, Tukey post-hoc test, a-b/c/d, c-b/d p<0.0001, b-d p<0.005). **B)** flg22-induced seedling growth
inhibition is not affected in *cepr1-3/2-4*. Relative fresh weight of five-day-old seedlings treated with flg22
(100 nM) for seven days; n = 14 ± SD (one-way ANOVA, Tukey post-hoc test, a-b p<0.0001). The
experiment was performed twice in independent biological repeats with similar results.
761

Figure S9: *rlk7* **and** *cepr1/2* **mutants show compromised CEP4-induced ethylene accumulation.**
A) CEP4-triggered ethylene production is reduced in *rlk7* mutants. Ethylene concentration in leaf discs upon ddH$_2$O or CEP4 (1 µM) treatment for 3.5 h; n = 15 pooled from three experiments ± SD (one-way ANOVA, Tukey post hoc, a-b p<0.01). **B)** CEP4-triggered ethylene production is compromised in *cepr1-3/2-4*. Ethylene concentration in leaf discs upon ddH$_2$O or CEP4 (1 µM) treatment for 3.5 h; n = 19 pooled from four independent experiments ± SD (one-way ANOVA, Tukey post hoc, a-b p<0.0001).

Peptide name	Sequence
CEP1	DFRHypTNPGNSHypGVGH
CEP2	DFAHypTNPGDSHypGIRH
CEP3	TFRHypTEPGHSHypGIGH
CEP4scr	TGQHypDHQRHypFAHIGGS
CEP415mer	AFRHypTHQGHypSQGIGH
CEP416mer	DAFRHypTHQGHypSQGIGH
CEP6.2	DFEPTTHypGHSHypGVGH
CEP9.1	DFVHypTSHypGNSHypGVGH
CEP9.2	DFAHypTSHypGHSHypGVGH
CEP9.3	DFAHypTSHypGNSHypGIGH
CEP9.4	DFAHypTTHypGNSHypGMGH
CEP9.5	DFKPTTHypGHSHypGVGH
PIP1	RLASGHypSPRGRGH
Flg22	Ac-QRLSTGSRINSAKDDAAGLQIA

770 **Table S1: Peptides used in this study**

Target site	Sequence	Purpose
CEP1.1	TGGTCACATATACACGGCG	
CEP1.2	CAGGCGTTGGACACTCTAA	
CEP2.1	GTCCCGGTATCAGGCATCC	
CEP2.2	ATGTATGCCCTGCAAAAGT	
CEP3.1	TACGGAATGTCCAATACCG	
CEP3.2	TATCAACCCCGTGGCTCGG	Generation of *cep6x* mutant
CEP4.1	CATTCCGACCAACCCACCA	
CEP4.2	TATTTTAAGGAGCACCTGG	
CEP6.1	ATGATCATCATTTCACAGT	
CEP6.2	TCGGGTACACTGATGATTT	
CEP9.1	CAAGATGACTTCAAGCCCA	
CEP9.2	ACACCAGGACATAGCCCCG	
CEPR1.1	AGAGAGACGAAGTACACGG	
CEPR1.2	AGCTCATGAGGTATAACAC	Generation of *cepr1/2*^{AEQ} mutant
CEPR2.1	GACCTCCTACGTGGCAGCG	
CEPR2.2	CTCAGACCGAGTATGAGAG	
RLK7.1	AGATTGATCTCTCTCGTCG	
RLK7.2	GCGACGGCTGATTTCCCGG	Generation of *rlk7/iku2*^{AEQ} mutant
IKU2.1	GCTTGAGATACTTAGACCT	
IKU2.2	CTTGTTCGTCTACAAAACT	

773 **Table S2: CRISPR-Cas9 target sites.**

Primers used for cloning		
Primer name	**Sequence (5'→3')**	**Purpose**
pCEPR1-BsaI-F	GCGCGCGGTCTCTCGAGTCAGTTTTTTTGGATTCAGATTTG	*pro*CEPR1 for pCEPR1::NLS-3xmVenus-tNOS
pCEPR1-BsaI-R	GCGCGCGGTCTCTCATTTTTCAGAGAAAGATCAAAAGTAACC	
pCEPR2-BsaI-F	GCGCGCGGTCTCTCGAGTTTTGTATACTCTTAAAAAAACTATG	*pro*CEPR2 for pCEPR2::NLS-3xmVenus-tNOS
pCEPR2-BsaI-R	GCGCGCGGTCTCTCATTTTATAGTGATTCCCAAGGG	
pCEP4-BpiI-F	TTGAAGACTTCGAGAATATAAACAAATATAGGTAAC	*pro*CEP4 for pCEP4::NLS-3xmVenus-t35S
pCEP4-BpiI-R	TTGAAGACTTCATTCACTTGACAACTTTCTG	
pRLK7-Attb1	GGGGACAAGTTTGTACAAAAAAGCAGGCTTAGAGCTAAGACATATAACT	*pro*RLK7 for pRLK7-H2B-mVenus
pRLK7-Attb2	GGGGACCACTTTGTACAAGAAAGCTGGGTAGTCGGAGAGAAGAAGTGG	
Primers used for RT-qPCR		
Primer name	**Sequence (5'→3')**	**Purpose**
qUBQ5-F	ACTCCTTCCTCAAACGCTGA	*UBQ5*
qUBQ5-R	CCAAGCCGAAGAAGATCAAG	
qFRK1-F	TGCAGCGCAAGGACTAGAG	*FRK1*
qFRK1-R	ATCTTCGCTTGGAGCTTCTC	
qPR1-F	CGGAGCTACGCAGAACAACT	*PR1*
qPR1-R	CAGACAAGTCACCGCTACCC	
qNRT2.1-F	AACAAGGGCTAACGTGGATG	*NRT2.1*
qNRT2.1-R	CTGCTTCTCCTGCTCATTCC	
qCEP1-F	AATGCTAAAGGGGTGTTTGG	*CEP1*
qCEP1-R	ACAAACCCACGACAAAGACA	
qCEP2-F	TGGTGACCATTTTGACCATC	*CEP2*
qCEP2-R	CCGACCATCTTTTTCGACTT	
qCEP3-F	GACCTACGGAACCTGGTCAT	*CEP3*
qCEP3-R	AAAAAGTCACCAGGCCAATC	
qCEP4-F	AGAATACAAAAGCAGCTCGAC	*CEP4*
qCEP4-R	TCCAATACCTTGACTAGGACC	
qCEP6-F	GTCGGGTAATCTAGACCTTCCTC	*CEP6*
qCEP6-R	CGGTTTTTCGCAACTGTCTCG	
qCEP9-F	AGCTGGATTTACAGATGATTTCG	*CEP9*
qCEP9-R	TGTCCCACACCAGGACTGT	
Primers used for genotyping and sequencing		
Primer name	**Sequence (5'→3')**	**Purpose**
GG pUC18 Seq-F	GTATCACGAGGCCCTTTCGT	Inserts in GoldenGate pUC18
GG pUC18 Seq-R	TAATGAATCGGCCAACGC	
CEP1-CRISP-s1	CCTTGTTTTGTTTTACATATTCTT	CRISPR CEP1
CEP1-CRISP-s2	TCTGAATGAAAATACATGTAATTTTC	
CEP2-CRISP-s1	CATTCAATACTTCTCATATACAAAC	CRISPR CEP2
CEP2-CRISP-s2	TCACCAGTTTTCTTTAGTCTTC	
CEP3-CRISP-s1	CATTTTCGCCTTCGACTA	CRISPR CEP3
CEP3-CRISP-s2	CAATCATATTTTCAACATGTAATCAC	
CEP4-CRISP-s1	GACATACGCTTTTAAAGGAAG	CRISPR CEP4
CEP4-CRISP-s2	GAAGAACTGTGTGCGATAAAG	
CEP6-CRISP-s1	CCTCTCTCGTATACACTAGAAAC	CRISPR CEP6
CEP6-CRISP-s2	TTAATTTAGCAAAAGCATTGAAG	
CEP9-CRISP-s1	ATATATACAAACTCCTAAACCTTGC	CRISPR CEP9

CEP9-CRISP-s2	CATATCGTAAACCAATGTATTAGA	
CEPR1-CRISP-s1	CGTAAGATGTGACGGTCAA	CRISPR CEPR1
CEPR1-CRISP-s2	GGTTGCCTTGCAGTACAA	
CEPR2-CRISP-s1	GAATTCCAGTGCATCGACC	CRISPR CEPR2
CEPR2-CRISP-s2	CTAAAACGTGGTAGGTGAATCAT	
RLK7-CRISP-s1	TCAAACCACCGTCACCACTT	CRISPR RLK7
RLK7-CRISP-s2	TCCAAATTCCGAAGCTCCGT	
IKU2-CRISP-s1	GCATGTGAATTCGCCGGAAT	CRISPR IKU2
IKU2-CRISP-s2	CCGCAGCTCTGATAAATCACC	
pCEPR1-s1	CTAGTCACCTGCAGCTCG	
pCEPR1-s2	GTGTAGTTCACGTCGGATG	
pCEPR1-s3	CAAACGCTTCACTTATGTAATG	pCEPR1
pCEPR1-s4	GCATCTATCTGATTCTGATCGTG	
pCEPR1-s5	CATCCGACGTGAACTACAC	
pCEPR2-s1	ACGACAATCTGAGCTCCAC	
pCEPR2-s2	TGGGATTCGGAGCATAAGG	
pCEPR2-s3	TCCATCTAGAAACATTAACCGA	pCEPR2
pCEPR2-s4	CCTAGAACGTGGTCCAAG	
pCEPR2-s5	CTGCACTGTAATGAGTACC	
pCEPR2-s6	CGGTAATAGATAGTGAAATGG	
cepr1-3-LP	TAAATCAAGAATCCACTTCCATGC	*cepr1-3*
cepr1-3-RP	CCTAATGGTAACTTATGGGACGCT	
cepr2-4-LP	GGTGAGTTCGTTATCGCTGAG	*cepr2-4*
cepr2-4-RP	AATGTTGAATCGACCGTTGAG	
GABI_o8474	ATAATAACGCTGCGGACATCTACATTTT	T-DNA primer for GABI-kat
cepr2-3-LP	TCACAACTCTGTAACGCAACG	*cepr2-3*
cepr2-3-RP	AACTCGGAGTTTTGAAGGAGC	
rlk7-1-LP	CCGCCTCTCTCTCTCTCTC	*rlk7-1*
rlk7-1-RP	AAGCAGAGCTTTCATCTTCCC	
rlk7-3-LP	ACGATTTGATCGTCGTGCTAC	*rlk7-3*
rlk7-3-RP	TTACAACAACTCCTTGACCGG	
iku2-4-LP	TCTTTAAGAACCGCAGCTCTG	*iku2-4*
iku2-4-RP	GTTGTTTCGCCTACAATGACC	
LBb1.3	ATTTTGCCGATTTCGGAAC	T-DNA LB primer for *rlk7-1, rlk7-3, iku2-4*
LBb1	GCGTGGACCGCTTGCTGCAACT	T-DNA LB primer for *cepr2-3*

776 **Table S3: Primers used in this study.**

778
779