1

XAP5 CIRCADIAN TIMEKEEPER regulates RNA splicing and the circadian clock via genetically separable pathways

Authors:

Hongtao Zhang (张弘韬)[1], Roderick W. Kumimoto[1], Shajahan Anver[1,2], Stacey L. Harmer[1*]

Author affiliations:

[1]Department of Plant Biology, College of Biological Sciences, University of California, Davis, CA 95616, USA

[2]Department of Genetics, Evolution and Environment, University College London, London, UK

[*]**Corresponding author:**

slharmer@ucdavis.edu

Running title:

Separable roles for *XCT* in splicing and the clock

One sentence summary:

XCT plays genetically separable roles in controlling the fidelity of 3' splice site selection during pre-mRNA splicing and the pace of circadian clock in Arabidopsis

Abstract

The circadian oscillator allows organisms to synchronize their cellular and physiological

activities with diurnal environmental changes. In plants, the circadian clock is primarily

composed of multiple transcriptional-translational feedback loops. Regulators of post-

transcriptional events, such as pre-mRNA splicing factors, are also involved in controlling the

pace of the clock. However, in most cases the underlying mechanisms remain unclear. We have

previously identified *XAP5 CIRCADIAN TIMEKEEPER* (*XCT*) as an *Arabidopsis thaliana*

circadian clock regulator with uncharacterized molecular functions. Here, we report that XCT

physically interacts with components of the spliceosome, including members of the Nineteen

Complex (NTC). PacBio Iso-Seq data show that *xct* mutants have transcriptome-wide pre-

mRNA splicing defects, predominantly aberrant 3' splice site selection. Expression of a genomic

copy of *XCT* fully rescues those splicing defects, demonstrating that functional *XCT* is important

for splicing. Dawn-expressed genes are significantly enriched among those aberrantly spliced in

xct mutants, suggesting that the splicing activity of *XCT* may be circadian regulated. Furthermore,

we show that loss of function mutations in *PRP19A* or *PRP19B*, two homologous core NTC

components, suppress the short circadian period phenotype of *xct-2*. However, we do not see

rescue of the splicing defects of core clock genes in *prp19 xct* mutants. Therefore, our results

suggest that *XCT* may regulate splicing and the clock function through genetically separable

pathways.

Introduction

Most eukaryotes have evolved an endogenous timekeeper known as the circadian clock, which

allows them to anticipate the daily fluctuating environmental conditions caused by the earth's

rotation (Harmer, 2009). Although the central oscillators of circadian clocks in diverse

eukaryotes lack conserved individual components, they share similar general architectures

(Nohales and Kay, 2016). In plants, the approximately 24-h periodicity of the clock is

maintained by a complex gene regulatory network consisting primarily of repressors and

activators of transcription. Those regulators, often referred to as core circadian clock genes,

regulate each other's expression and the expression of thousands of output genes (Creux and

Harmer, 2019).

Additionally, post-transcriptional and post-translational mechanisms such as alternative splicing

of precursor messenger RNAs (pre-mRNAs) provide critical regulation of clock function. It has

been suggested that changes in splicing of core circadian clock genes are important for

Arabidopsis in response to environmental stresses (James et al., 2012; Kwon et al., 2014). For

example, cold temperatures significantly suppress production of a splice variant of *CIRCADIAN*

CLOCK-ASSOCIATED1 (*CCA1*) in which intron four is retained. This incompletely spliced

isoform encodes a truncated protein which competitively inhibits the function of fully spliced

CCA1 (Seo et al., 2012). However, in most cases the effects of alternative splicing of pre-

mRNAs on circadian clock function remain unclear.

In contrast, the mechanisms underlying pre-mRNA splicing are increasingly well understood

(Wilkinson et al., 2020). There are two catalytic transesterification steps, which sequentially

remove first the 5' and then the 3' splice sites (5'SS and 3'SS) of introns from their adjacent

exons (Shi, 2017). This process is carried out by five small nuclear ribonucleoproteins (snRNPs)

and hundreds of non-snRNP splicing factor proteins which assemble on a pre-mRNA to form the

spliceosome complex (Wilkinson et al., 2020). One of these spliceosome complex components is

the Nineteen Complex (NTC, also known as PRP19 complex), named after PRECURSOR RNA

PROCESSING 19 (PRP19), a U-box E3 ubiquitin ligase that forms the core of the NTC

(Hatakeyama et al., 2001; Koncz et al., 2012). The NTC is associated with the spliceosome

during the two transesterification steps and helps facilitate conformational rearrangements and

promote splicing fidelity (Fig. 1; Hogg et al., 2010). The NTC is highly conserved across

eukaryotes. In Arabidopsis, multiple orthologs of yeast NTC proteins including two PRP19

homologs (PRP19A/MAC3A and PRP19B/MAC3B) have been shown to physically interact

with the spliceosome (Monaghan et al., 2009; Deng et al., 2016). Plants mutant for *prp19a*

prp19b or homologs of other NTC-associated proteins such as *pleiotropic regulatory locus1*

(*prl1*) and *snw/ski-interacting protein* (*skip*) have genome-wide intron retention defects (Jia et al.,

2017; Wang et al., 2012; Li et al., 2019). These data indicate that NTC components act as

evolutionarily conserved splicing factors in Arabidopsis.

Mutation of splicing factors can lead to disruption of circadian clock function (Shakhmantsir and

Sehgal, 2019). For example, loss-of-function alleles of NTC components, including *PRP19*,

PRL1 and *SKIP*, cause lengthening of circadian period (Wang et al., 2012; Feke et al., 2019; Li

et al., 2019). Aberrantly spliced mRNA variants of core circadian clock genes have been

detected in these splicing factor mutants (Sanchez et al., 2010; Wang et al., 2012; Jones et al.,

2012; Schlaen et al., 2015; Marshall et al., 2016; Li et al., 2019; Feke et al., 2019), suggesting

that changes in the pace of the clock might be caused by aberrant splicing of core clock genes. In

some cases, epistasis analysis suggests that this may be true (Marshall et al., 2016; Schlaen et al.,

2015). However, in other cases, genetic analysis has either not been performed or has revealed

additive interactions between the splicing factor and clock gene mutants. In addition, the levels

of mis-spliced mRNA variants of clock genes are usually only a small fraction of total transcripts

(Jones et al., 2012; Perez-Santángelo et al., 2014; Feke et al., 2019; Sanchez et al., 2010; Wang

et al., 2012). Thus, it is unclear whether splicing factors affect clock function solely by

controlling the splicing of core clock genes.

RNA splicing factors often function in multiple biological pathways: plants deficient for

components of the NTC have defects in immunity, microRNA biogenesis, DNA damage

response and transcriptional elongation (Koncz et al., 2012; Monaghan et al., 2009; Zhang et al.,

2013; Jia et al., 2017). Some splicing factors are known to carry out roles in nuclear processes

biochemically separable from their roles in splicing. In addition to its structural role in the

spliceosome, PRP19 also senses DNA damage and promotes DNA repair via its ubiquitin ligase

activity (Maréchal et al., 2014). Additionally, SKIP acts in two distinct complexes to regulate

splicing and the transcription of genes involved in flowering time control (Li et al., 2019).

Nonetheless, whether splicing factors might control circadian clock function via splicing-

independent activities has not been investigated.

We previously identified *XAP5 CIRCADIAN TIMEKEEPER* (*XCT*) as a novel regulator of the

Arabidopsis circadian clock (Martin-Tryon and Harmer, 2008). Like NTC components, XCT is

also well-conserved across eukaryotes. Homologs of XCT (also known as XAP5 proteins) share

a highly conserved C-terminal protein domain and are nuclear-localized (Martin-Tryon and

Harmer, 2008; Anver et al., 2014; Li et al., 2018; Lee et al., 2020). Previously, we found that

transgenic expression of Arabidopsis *XCT* fully rescued the slow-growth phenotype of fission

yeast mutant for *xap5* (Anver et al., 2014). Taken together, these data suggest that XCT

homologs might share similar molecular and cellular functions across kingdoms.

It has been reported that FAM50A, one of the two XCT orthologs in humans, physically

associates with the spliceosomal C complex and its mutants have defects in RNA splicing

(Bessonov et al., 2008; Lee et al., 2020). Similarly, a recent study in Arabidopsis also reported

the association of XCT with the spliceosome (Liu et al., 2022), implying that XAP5 proteins may

be evolutionarily conserved splicing factors. However, evidence suggests that XAP5 proteins

may also participate in fundamental biological processes other than splicing. Fission yeast and

Chlamydomonas XAP5 proteins associate with chromatin and directly regulate transcription

(Anver et al., 2014; Li et al., 2018). In addition to its role in the clock, Arabidopsis *XCT* has been

implicated in diverse processes including small RNA production, immune signaling, and DNA

damage responses (Fang et al., 2015; Xu et al., 2017; Kumimoto et al., 2021). Notably, NTC

components have also been reported to function in all these pathways (Jia et al., 2017; Maréchal

et al., 2014; Chanarat and Sträßer, 2013). Although pleiotropic defects are seen in *xct* mutants,

interconnections between these phenotypes and the molecular function of *XCT* have not been

extensively studied.

In the current work, we report that XCT physically interacts with NTC and other spliceosome-

associated proteins in Arabidopsis. We use long-read RNA sequencing to reveal that *XCT*

controls the fidelity of 3' splice site selection for hundreds of pre-mRNA splicing events,

probably by rejecting downstream suboptimal 3' splice sites. Intriguingly, circadian-regulated

genes that are aberrantly spliced in *xct* mutants are significantly enriched for peak expression at

subjective dawn. This implies that the splicing-related activity of *XCT* may be circadian

regulated. We also demonstrate that *PRP19A* and *PRP19B* display epistatic genetic interactions

with *XCT* in control of the circadian clock period length but not of the splicing of core clock

genes. This is consistent with our finding that although *xct-2* causes more severe splicing defects

than *xct-1*, *xct-1* mutants have a stronger circadian phenotype. Collectively, our results suggest

that XCT works close to NTC and may regulate pre-mRNA splicing and the circadian clock

function via distinct pathways.

Results

XCT physically interacts with the Nineteen Complex and other spliceosome-associated

proteins

To uncover the molecular functions of XCT, we carried out mass spectrometry (MS)

experiments to identify its possible interactors. We expressed epitope-tagged XCT protein under

the control of the native *XCT* promoter in *xct-2* mutant background (*xct-2 XCT*). This transgene

largely rescued the morphological and circadian clock defects of *xct-2* (Supplemental Fig. 1).

Next, we affinity purified tagged XCT from plant extracts and analyzed co-purifying proteins via

MS. In total, 26 proteins were significantly enriched in the XCT immunoprecipitation compared

to control immunoprecipitations (Welch's two sample *t*-test, $P < 0.05$; Supplemental Dataset 1).

Among those XCT-associated proteins, 15 (57.7%) were annotated as mRNA splicing-related,

including 11 core NTC or NTC-associated proteins (Table 1), consistent with the MS and co-

immunoprecipitation data from a recent report (Liu et al., 2022). Notably, studies in human and

yeast have revealed that NTC is physically associated with the spliceosomal complex throughout

the two catalytic transesterification steps (Fig. 1; Hogg et al., 2010). Taken together, those results

imply that *XCT* may act close to NTC and function in the catalytic steps during pre-mRNA

splicing. Interestingly, we also observed 8 (30.8%) chloroplast proteins enriched in the XCT

immunoprecipitation (Supplemental Dataset 1), which may be related to the delayed leaf

greening phenotypes observed in *xct-2* mutant plants (Martin-Tryon and Harmer, 2008).

***XCT* is required for the fidelity of 3' splice site selection during pre-mRNA splicing**

To investigate a possible role for *XCT* in RNA splicing, we performed PacBio Isoform

Sequencing (Iso-Seq) on wild-type Col-0, reduction-of-function allele *xct-1*, null allele *xct-2*, and

the complemented line *xct-2 XCT*. Additionally, we also sequenced *prl1-2*, which contains a T-

DNA insertion mutation *in PRL1*, an NTC member that has been demonstrated to control

genome-wide splicing efficiency (Jia et al., 2017). Since both transcript levels and RNA splicing

of a large proportion of the Arabidopsis transcriptome are circadian regulated (Romanowski et

al., 2020; Yang et al., 2020), the time of day at which plants are harvested has significant effects

on gene expression and splicing analysis (Hsu and Harmer, 2012). Therefore, we grew plants in

constant environmental conditions for ten days to desynchronize clocks of individual plants. To

further minimize any differences in subjective time of day between wild-type plants, the long-

circadian-period *prl1-2*, and the short-period *xct* mutants (Supplemental Fig. 2), we pooled

seedlings harvested at 2-hour intervals across a twenty-four hour period (Fig. 2A). For each

genotype, we sequenced three biological replicates and acquired an average of 784,832 full-

length transcript reads per genotype per replicate (Supplemental Dataset 2). With no alignment

needed, those reads were directly mapped to the TAIR 10 Arabidopsis reference genome and

then subjected to differential splicing analysis using the JunctionSeq R package (Hartley and

Mullikin, 2016).

Previous studies have demonstrated that intron retention is the most prevalent type of alternative

splicing event in Arabidopsis (Wang and Brendel, 2006; Filichkin et al., 2010). Indeed, plants

mutant for multiple NTC components have been reported to have global intron retention defects

(Jia et al., 2017; Meng et al., 2022). In our analysis of *prl1-2*, we found 5,730 out of 44,496

detected splicing events were significantly differentially enriched from Col-0 (Supplemental

Dataset 3). Notably, 99% of these events were decreases in known junctions (Supplemental Fig.

3), consistent with previous studies suggesting that loss of *PRL1* function mainly causes intron

retention without affecting splice site selection (Jia et al., 2017).

We next examined transcript composition in *xct* mutants. In the null allele *xct-2*, we detected

75,073 splicing events, of which 875 were significantly down-regulated and 532 were up-

regulated compared with Col-0 (FDR<0.05, Fig. 2B-C). Meanwhile, 209 and 337 splicing events

were significantly decreased or increased, respectively, in the partial loss-of-function mutant *xct-*

1. Comparing mis-regulated splicing between the two *xct* mutant alleles, 196 (93.8%) of the

down-regulated and 329 (97.6%) of the up-regulated events were shared. Thus, the two *xct*

alleles have similar splicing phenotypes but the severity of the defect is greater in the null allele

despite the stronger effect of *xct-1* on circadian period (Supplemental Fig. 1). Next, we analyzed

the splicing events in the complemented line *xct-2 XCT*. Only 16 out of 84,965 events were

significantly differentially enriched compared with Col-0, indicating that the splicing defects of

xct-2 were almost completely rescued by the restored *XCT* expression. Taken together, these

results demonstrate that *XCT* is important for transcriptome-wide pre-mRNA splicing.

To further characterize the major types of splicing defects caused by loss of *XCT* function, we

analyzed the splice sites for the differentially spliced junctions in *xct* mutants. Specifically, we

categorized *xct*-induced mis-splicing events into four classes based on whether the 5' and 3'

splice sites used were previously annotated (known) or not (novel). As expected, most down-

regulated splicing events displayed decreases in usage of junctions with known 5' and 3' splice

sites (Fig. 2D). Interestingly, 307 (90.8%) in *xct-1* and 462 (86.5%) in *xct-2* of up-regulated

splicing events involved increased usage of junctions with a known 5' splice site but a novel 3'

splice site (Fig. 2E). The decreases in abundance of junctions with known splice sites hence

reflects both intron retention events and novel 3' splice sites usage. Therefore, our results

demonstrate that in addition to controlling the efficient removal of introns, *XCT* is also

responsible for the fidelity of 3' splice site selection.

To investigate how *XCT* contributes to the fidelity of 3' splice site selection, we compared the

significantly up- or down-regulated 3' splice sites in *xct* mutants with all detected 3' splice sites

in Col-0. Intriguingly, most of the up-regulated novel 3' splice sites in *xct-1* and *xct-2* were

located less than 50 nucleotides downstream of the wild-type 3' splice sites (Fig. 3B;

Supplemental Fig. 4A). Studies in human and yeast have demonstrated that the sequences

preceding a 3' splice site, including the polypyrimidine tract (PPT) and the pyrimidine at the -3

position (Fig. 3A), are important for the strength of the 3' splice site (Horowitz, 2012). Therefore,

we further examined the 23-mer sequences flanking the 3' splice sites (from -20 to +3 position).

We found that the frequency of the canonical AG sequence at the 3' splice sites was not altered

in either *xct-1* or *xct-2* (Fig. 3C; Supplemental Fig. 4B-C). However, the average predicted

strength of the 875 down-regulated 3' splice sites in *xct-2* was significantly weaker than the

average of total detected 3' splice sites in Col-0 (Fig. 3D). Specifically, the percentage of

pyrimidine residues at the -3 position was significantly reduced in down-regulated 3' splice sites

compared to that found in all detected 3' splice sites (Fisher's exact test, $P < 0.001$, Fig. 3E),

suggesting that functional *XCT* is important for the removal of 3' splice sites with a suboptimal

sequence at the -3 position. Moreover, the 532 up-regulated 3' splice sites had an even lower 3'

splice site score than the down-regulated sites (Fig. 3D). The frequency of pyrimidines both at

the -3 position and throughout the PPT region was significantly lower in the up-regulated

junctions (Fig. 3E-F), showing that *xct-2* is less able to discriminate between 3' splice sites

during splicing. Taken together, our Iso-Seq data demonstrates that *XCT* controls the accuracy of

3' splice site selection, possibly by helping to recognize weaker 3' splice sites and reject sub-

optimal downstream sites.

The splicing defects of core clock genes are generally more severe in *xct-2* than *xct-1*

Since aberrant splicing of core clock genes may fully or partially contribute to the altered

circadian clock period phenotype in splicing factor mutants (Sanchez et al., 2010; Wang et al.,

2012; Jones et al., 2012; Schlaen et al., 2015; Marshall et al., 2016; Li et al., 2019; Feke et al.,

2019), we searched for core clock genes with aberrant splicing events in both the short-period

xct-1 and *xct-2* mutants. In total, our Iso-Seq data detected five aberrantly spliced core clock

genes, including *LHY*, *LNK2*, *PRR7*, *TOC1* and *TIC* (Fig. 4A). However, we noticed that the

ratio of aberrantly-spliced relative to fully-spliced transcripts for several clock genes are

significantly higher in *xct-2* than *xct-1*, even though the circadian period defect is more severe in

the latter (Supplemental Fig. 5A-D; Supplemental Fig. 1B). We confirmed this finding via semi-

qRT-PCR assay (Supplemental Fig. 5E-H). Thus, those results imply that there may be a

disconnect between the splicing and circadian clock defects in *xct-1* and *xct-2*.

Genes expressed at subjective dawn are enriched among those aberrantly spliced in *xct*

mutants

To more broadly explore possible links between the circadian clock and splicing phenotypes in

xct mutants, we looked for enrichment of circadian clock regulation within genes aberrantly

spliced in *xct*. Among the 5,602 previously reported clock-regulated genes (Romanowski et al.,

2020) that are detected in our Iso-Seq data, we identified 182 and 402 genes that are mis-spliced

in *xct-1* and *xct-2*, respectively (Fig. 4A; Supplemental Dataset 4). However, there is no

significant over-representation in either mutant (one-tailed Fisher's exact test, $P = 0.87$ for *xct-1*

and 0.99 for *xct-2*), suggesting that *XCT* does not preferentially affect the splicing of clock-

regulated genes.

Previous studies have identified transcripts that are differentially spliced at various times of day

(Romanowski et al., 2020; Yang et al., 2020). This suggests that splicing activity may be

circadian clock regulated. Therefore, to test whether *XCT* preferentially affects pre-mRNA

splicing at certain times of day, we examined the distribution of estimated peak expression times

of genes aberrantly spliced in *xct* mutants. We grouped all 5,602 clock-regulated genes into

twelve 2-hour intervals based on their estimated peak expression time (Fig. 4B). Genes that are

aberrantly spliced in *xct-2* are significantly enriched for peak expression between ZT22 and ZT2

compared with the other intervals of the day (Fig. 4C-D, Fisher's exact test, $P < 1e-4$). Similarly,

clock-regulated genes that are aberrantly spliced in *xct-1* also showed a dawn-enriched

expression pattern (Supplemental Fig. 6). Collectively, those results indicate that *XCT* activity

within the spliceosome may be regulated by the circadian clock.

To further investigate whether *XCT* preferentially affects splicing of transcripts produced at

dawn, we monitored the abundance of candidate splice variants via quantitative reverse

transcriptase-polymerase chain reaction (qRT-PCR) over a 24-hour light/dark period (LD)

followed by 48 hours in constant light (LL). We selected four circadian-clock-regulated genes

and one non-clock-regulated gene that are aberrantly spliced in *xct-2* and fully rescued in *xct-2*

XCT (Fig. 5A-E). Total transcript levels of clock-regulated genes displayed rhythmic abundance

with peak *LHY* and *LNK2* expression at dawn and peak *PRR7* and *TOC1* expression late in the

afternoon (Supplemental Fig. 7), as expected (Creux and Harmer, 2019). We found significantly

advanced phases of expression in *xct-2* but this mutation did not affect the overall abundance of

total transcripts for those genes. However, both the peak and average abundance of the aberrantly

spliced isoforms were significantly different in *xct-2* compared with Col-0 (Fig. 5F-J). This

serves as validation of the pooling strategy we used to generate our Iso-Seq libraries, where the

daily abundance of transcripts was averaged (Fig. 2A; Fig. 5A-E). Contrary to our expectations,

instead of preferentially accumulating at dawn, abundance of all splice variants in *xct-2* was

synchronized with total transcript levels (Fig. 5F-I; Supplemental Fig. 7). Likewise, the mis-

spliced isoform of the non-clock-regulated gene *SPPA* did not show a significant morning peak

(Fig. 5J). Meanwhile, the genomic *XCT* complemented line almost completely rescued the

splicing defects across the whole experimental period. Taken together, those results suggest that

although morning-expressed genes are preferentially enriched among *XCT* targets, *XCT* is

important for the accuracy of pre-mRNA splicing throughout the day under both LD and free-

running LL conditions.

Reduction of *PRP19* function rescues the circadian clock but not the splicing defects in *xct-*

2

Previous studies revealed that NTC components participate in regulation of circadian clock

function in Arabidopsis (Feke et al., 2019; Li et al., 2019). Here we examined circadian period in

mutants for several NTC components. We found that mutation in either of the two Arabidopsis

PRP19 homologs, *PRP19A* (*MAC3A*) or *PRP19B* (*MAC3B*), only caused minor lengthening of

circadian period. In contrast, the *prp19a-1 prp19b-1* double mutant had a significantly slower

circadian clock than Col-0 (Supplemental Fig. 8A), consistent with previously reported

redundant roles for *PRP19A* and *PRP19B* in circadian clock regulation (Feke et al., 2019).

Similarly, loss of function of other NTC members, including *PRL1*, *CDC5* and *SKIP*, also

lengthened the clock period by 1 to 3 hours (Supplemental Fig. 2B; Supplemental Fig. 8B-C),

similar to previous reports for *prl1-9* and *skip-1* mutants (Li et al., 2019).

Since *PRP19* co-purifies with *XCT* (Table 1) and they both control the pace of the circadian

clock (Supplemental Fig. 8A), we hypothesized they might function in the same pathway to

regulate the clock. To test this hypothesis, we introduced *prp19a-1* and *prp19b-1* mutant alleles,

which express greatly decreased and near-null levels of *PRP19A* and *PRP19B*, respectively

(Supplemental Fig. 9), into the null *xct-2* background and assayed their circadian clock

phenotypes. Interestingly, we found that neither the circadian period of *prp19a-1 xct-2* nor of

prp19b-1 xct-2 was significantly different from Col-0 (Fig. 6A; Supplemental Fig. 10),

indicating that both *prp19* mutants can fully suppress the short-period phenotype of *xct-2*.

Notably, *prp19a-1* and *prp19b-1* single mutants only have slightly longer circadian periods than

Col-0. This suggests that the rescued clock function in *prp19 xct* double mutants is not likely due

to additive genetic interactions. Thus, our data show that functional *PRP19A* and *PRP19B* are

both necessary for *XCT* in regulation of circadian period.

To further investigate the functional association between *XCT* and *PRP19*, we next asked

whether *prp19* mutants suppress other developmental and physiological defects observed in *xct-2*.

In contrast to the circadian period, the reduced rosette size in *xct-2* was not significantly restored

in either *prp19a-1 xct-2* or *prp19b-1 xct-2* (Supplemental Fig. 11), indicating that the effects of

XCT on rosette development and the circadian clock period are genetically separable. This is

consistent with our observation that *xct-1* and *xct-2* both have short circadian period phenotypes

but only *xct-2* is morphologically different from Col-0 (Supplemental Fig. 1).

Next, to determine whether suppression of the short clock period phenotype in *xct-2 prp19*

mutants is due to reversal of the *xct-2* splicing defects, we conducted qRT-PCR experiments to

examine the abundance of *xct*-induced aberrantly-spliced mRNA isoforms of core clock genes in

prp19a-1 xct-2 and *prp19b-1 xct-2*. Surprisingly, none of the aberrantly-spliced isoforms tested

showed significantly different abundance in *prp19a-1 xct-2* and *prp19b-1 xct-2* relative to *xct-2*

(Fig. 6B-F). Consistent with previous reports for functional redundancy between *PRP19A* and

PRP19B (Monaghan et al., 2009; Li et al., 2019), we noticed a statistically significant defect in

splicing of the third intron of *LNK2* in the *prp19a-1 prp19b-1* double mutant but not the *prp19a-*

1 or *prp19b-1* single mutants. We also checked the expression levels of fully-spliced or total

mRNA isoforms of the five core clock genes that are aberrantly-spliced in *xct-2*. In all cases,

addition of *prp19a-1* or *prp19b-1* to *xct-2* did not alter the abundance of functionally spliced

isoforms of those clock genes (Supplemental Fig. 12). Thus, similar to the developmental defects,

the splicing defects of core clock genes are genetically separable from the circadian clock

phenotype of *xct-2*.

Discussion

The accurate removal of introns from pre-mRNAs is an essential step of gene expression in all

eukaryotes. In this work, we report that *XCT* is a global regulator of pre-mRNA splicing in

Arabidopsis. Multiple lines of evidence suggest that orthologs of XCT share conserved

molecular and cellular functions across kingdoms. Previous mass spectrometry data showed that

FAM50A and FAM50B, two human homologs of XCT, physically associate with the affinity-

purified spliceosomal C complex (Bessonov et al., 2008; Bessonov et al., 2010). Loss of

FAM50A function induces transcriptome-wide pre-mRNA splicing defects in both human and

zebrafish (Lee et al., 2020). Similarly, a recent study in Arabidopsis also found that XCT

associates with spliceosomal proteins and regulates splicing (Liu et al., 2022). In this paper, we

show that XCT physically co-purifies with PRP19 and other NTC-associated proteins (Table 1).

Furthermore, our PacBio Iso-Seq data indicate that the efficiency and fidelity of pre-mRNA

splicing are negatively impacted in both the partial loss-of-function mutant *xct-1* and the null

mutant *xct-2* but are largely restored in *xct-2 XCT* (Fig. 2). Therefore, XCT orthologs likely play

evolutionarily conserved roles in pre-mRNA splicing. It would be interesting to apply structural

biology approaches to further investigate what conserved functional domains among XCT

homologs might contribute to their splicing activities.

Studies have shown that intron retention is the most prevalent type of alternative splicing event

in Arabidopsis (Wang and Brendel, 2006; Filichkin et al., 2010). Correspondingly, mutations in

many Arabidopsis splicing factors mainly cause intron retention (Supplemental Fig. 3; Schlaen et

al., 2015; Li et al., 2019). Here we identify *XCT* as an unusual splicing regulator that specifically

controls the fidelity of 3' splice site selection (Fig. 2D-E; Fig. 3). Biochemical studies of splicing

in yeast revealed that several DEAH-box ATPases are important for the selection of 3' splice

sites (Horowitz, 2012). For example, PRP22, a DEAH-box ATPases that co-purifies with XCT

(Table 1; Liu et al., 2022), represses usage of aberrant 3' splice sites and promotes spliceosome

scanning for downstream alternative 3' splice sites in yeast (Mayas et al., 2006; Semlow et al.,

2016). Studies in human showed that hPRP22, FAM50A and FAM50B, are all abundant in the

spliceosomal C complex but nearly absent in B and B* complexes (Bessonov et al., 2008;

Bessonov et al., 2010; Zhan et al., 2022). This implies that PRP22 and XCT may have

evolutionarily conserved interactions during the second transesterification reaction of splicing,

when 5' and 3' splice sites are joined (Fig. 1). Intriguingly, we detected decreased fidelity of 3'

splice site selection in both *xct-1* and *xct-2* mutants (Fig. 3; Supplementary Fig. 4), which is also

observed in yeast *prp22* mutants (Semlow et al., 2016). Therefore, our results imply that XCT

may work with PRP22 to control the fidelity of 3' splice site selection. Future research on the

biochemical functions of XCT using *in vitro* and *in vivo* systems could help reveal how it

controls splicing fidelity.

Previous microarray and RNAseq studies demonstrated that transcription and splicing of a large

proportion of Arabidopsis genes are under circadian clock regulation (Covington and Harmer,

2007; Hsu and Harmer, 2012; Romanowski et al., 2020; Yang et al., 2020). Consequently, timing

of sample collection is an important consideration in gene expression and splicing analysis,

especially when comparing genotypes with different circadian periods. Here we employed a

more efficient sampling method by harvesting plants at twelve evenly distributed time points

across a day and pooling them before analysis. Our pooling strategy enabled us to detect

transcripts of genes with the distribution of peak phases of expression mirroring that of all clock-

regulated genes (Fig. 4B; Romanowski et al., 2020). Overall, this strategy allowed us to identify

more clock-regulated differentially spliced transcripts in *xct-2* mutants than in a recent study in

which plants were collected at a single time point (Liu et al., 2022). For example, we report

novel differential splicing events in afternoon-expressed genes including *PRR7* and *TOC1* (Fig.

4A; Supplemental Fig. 7C-D). In fact, our time-course qPCR data suggest that abundance of the

aberrantly spliced isoforms of these clock-regulated genes fluctuates by over 99% depending on

time of sample collection (Fig. 5F-I). Thus, our results demonstrate the advantages of pooling

samples when conducting transcriptome-wide splicing analysis of clock-regulated genes.

Using this strategy, we were able to investigate whether *XCT* preferentially affects splicing at

certain times of day. Indeed, we found that dawn-expressed genes are significantly more likely to

be mis-spliced in *xct* mutants than those with other peak phases (Fig. 4C-D; Supplemental Fig. 6).

This suggests that *XCT* activity in or its association with the spliceosome may be clock-regulated.

An alternative explanation for the overrepresented mis-splicing of dawn-expressed genes could

be that *cis*-regulatory elements of those genes might preferentially recruit XCT to control their

splicing. This possibility is supported by previous studies showing that XCT orthologs in fission

yeast and *Chlamydomonas* are chromatin-associated and in the latter case recruit RNA

polymerase II (Pol II) to promoter regions (Anver et al., 2014; Li et al., 2018).

Alternative splicing regulates various biological processes, including the function of the

circadian clock (Hsu and Harmer, 2014; Nolte and Staiger, 2015). In Arabidopsis, such

regulation is supported by identification of splicing factor mutants that alter circadian clock

period length (Shakhmantsir and Sehgal, 2019). Some splicing factors, such as *GEMIN2* and

SICKLE, interact epistatically with one or more alternatively spliced clock genes in the control of

period length (Schlaen et al., 2015; Marshall et al., 2016), suggesting changes in the pace of the

clock are due to altered splicing of these clock genes. Whereas in other cases, the mis-regulated

circadian period can only be partially attributed to changes in levels of splicing variants of clock

genes (Sanchez et al., 2010; Wang et al., 2012). Additionally, in most splicing factor mutants,

only small fractions of total transcripts are aberrantly processed (Jones et al., 2012; Perez-

Santángelo et al., 2014; Feke et al., 2019; Li et al., 2019). Therefore, whether these small

changes in levels of aberrantly spliced isoforms could lead to significantly decreased levels of

functional proteins and thereby cause the observed circadian phenotypes remains unclear.

Here, we demonstrate that in *xct* mutants the circadian clock phenotype is genetically separable

from altered levels of aberrantly spliced mRNA variants of core clock gene. We found that loss

of *XCT* function accelerates the clock and causes aberrant splicing of five core clock genes

(Supplemental Fig. 1; Fig. 4A). Surprisingly, the clock phenotype but not the splicing defects of

the five clock genes in *xct-2* is suppressed by reduction of *PRP19A* or *PRP19B* function (Fig. 6).

Similarly, the clock in *xct-1* runs slightly faster than *xct-2* but the splicing defects, including the

aberrant splicing of *TOC1*, *TIC* and *CCA1*, are more severe in the latter (Fig. 2; Supplemental

Fig. 5). Collectively, these data demonstrate that the short period phenotypes and altered levels

of clock gene splicing variants are genetically separable in *xct* mutants.

How *XCT* regulates the pace of the circadian clock is still an outstanding question. Although we

argue that changes in levels of aberrantly spliced clock mRNAs are not responsible for the

accelerated clock in *xct* mutants, one possibility is that alterations in the overall kinetics of pre-

mRNA splicing may cause circadian period phenotypes. Indeed, pharmacological perturbations

of global transcription and translation efficiency can both lengthen the circadian period (de Melo

et al., 2021; Uehara et al., 2022). Those results suggest that changes in the kinetics of RNA

processing might cause the period phenotypes observed in plants mutated for some splicing

factors.

Alternatively, *XCT* may control the circadian clock function independent of its role in pre-

mRNA splicing. In Chlamydomonas, the XCT homolog XAP5 co-immunoprecipitates with

RNA Pol II (Li et al., 2018), suggesting that XCT orthologs may possess transcriptional

regulatory activities. In this paper, we show that NTC components physically and genetically

interact with XCT to regulate circadian clock period (Table 1; Fig. 6A). Besides splicing, another

well-characterized role of the NTC in gene expression is regulation of transcriptional elongation

(Chanarat and Sträßer, 2013). Studies in both yeast and Arabidopsis have revealed that multiple

NTC members, including PRP19, PRL1 and CDC5, physically associate with RNA Pol II and

participate in transcript elongation (Chanarat et al., 2011; Zhang et al., 2013; Zhang et al., 2014).

In addition, the NTC component SKIP interacts with Polymerase-Associated Factor 1 complex

to regulate transcription in a splicing-independent manner (Li et al., 2019). Interestingly, a recent

study showed that inhibition of transcriptional elongation by decreasing phosphorylation of the

RNA Pol II C-terminal domain lengthens circadian period in Arabidopsis (Uehara et al., 2022). It

is therefore possible that *XCT* affects circadian period by altering transcriptional elongation.

Yet another possibility is that *XCT* affects a cellular process independently of its role in gene

expression. For example, PRP19 is known to facilitate DNA repair by its E3 ubiquitin ligase

activity independent of its involvement in the spliceosomal complex (Maréchal et al., 2014;

Idrissou and Maréchal, 2022). Indeed, there is increasing evidence that many RNA binding

proteins directly participate in DNA double-strand break responses (Klaric et al., 2021). A recent

study showed that the NTC-associated protein MOS4-ASSOCIATED COMPLEX SUBUNIT

5A (MAC5A) physically interacts with the 26S proteasome and regulates its activities in

response to DNA damage (Meng et al., 2022). Intriguingly, we recently reported that loss of *XCT*

function also disturbs the DNA damage response pathway (Kumimoto et al., 2021). Thus, it is

possible that *XCT* works with *PRP19* in this pathway separately from its role in RNA processing.

Future research is required to understand the nature of the relationship of between XCT and the

NTC in the control of circadian clock function and other essential biological processes.

Materials and Methods

Plant materials and growth conditions

All *Arabidopsis thaliana* plants used in this study are Columbia-0 (Col-0) ecotype and contain a

CCR2::*LUC* reporter for circadian clock assays. The *xct-1*, *xct-2*, *xct-2* p*XCT*::g*XCT-YFP-HA*,

prp19a-1, *prp19b-1*, *prl1-2*, *cdc5-1* and *skip-1* genotypes have been previously described

(Martin-Tryon and Harmer, 2008; Monaghan et al., 2009; Zhang et al., 2014; Wang et al., 2012).

All the double mutants in this study were produced by crossing. Unless otherwise specified,

seeds were surface sterilized with chlorine gas for 3 hours (50 ml 100% bleach + 3ml 1M HCl)

and then plated on 1x Murashige and Skoog (MS) growth media containing 0.7% agar, pH 5.7.

After 3d stratification in dark at 4˚C, plates were transferred to 12-h light (cool white fluorescent

bulbs, 55 µmol m^{-2} s^{-1}) / 12-h dark cycles at 22˚C for a variable number of days depending on the

experiment.

Immunoprecipitations and mass spectrometry

Plants were grown in 12-h light / 12-h dark cycles and harvested on day 10 at ZT3 or ZT17.

Approximately 7.5g of vegetative tissue was flash frozen in liquid nitrogen and ground into fine

powder with a mortar and pestle. Ground tissue was resuspended in nuclei enrichment buffer (50

mM Tris pH7.5, 400 mM sucrose, 2.5 mM MgCl$_2$, 1 mM DTT, 1 mM PMSF, cOmplete Protease

Inhibitor cocktail (Roche)), the nuclear pellet was collected, resuspended in IP buffer (100mM

Tris pH7.5, 1mM EDTA, 75mM NaCl, 10% glycerol, 0.3% Triton X-100, 0.05% SDS, 10 mM

MG-132, 1 mM PMSF, , cOmplete Protease Inhibitor cocktail (Roche)) and pelleted again.

Nuclei were resuspended in IP buffer, disrupted via sonication, and adjusted to 1 mg/mL in IP

buffer. 1 mg of extract was incubated with µMACS MicroBeads conjugated to an anti-HA

monoclonal antibody (Miltenyi Biotec) and beads were then captured on µMACS M-colums.

Beads were washed 3x with ice cold IP buffer and 1x with ice cold TE buffer (10 mM Tris-HCl

pH 8.0, 0.1 mM EDTA). Proteins were eluted off the beads with elution buffer (Miltenyi Biotec).

Proteins were resolved using SDS polyacrylamide gel electrophoresis and gels were sent to the

Rutgers Biological Mass Spectrometry Facility, Robert Wood Johnson Medical School. Proteins

were eluted and subjected to mass spectrometry as previously described (Wei et al., 2020).

PacBio Iso-Seq and bioinformatic analysis

Arabidopsis seeds were surface sterilized in 70% ethanol prepared in 0.1% (v/v) Triton X-100

(Sigma) for 5 minutes and then in 100% Ethanol for 20 minutes. After air drying, seeds were

plated on 1x MS growth media containing 0.7% agar. After cold stratification for 3 days, plates

were transferred to constant light (55 µmol m^{-2} s^{-1}) and temperature (22˚C). Approximately 60

mg of whole seedlings of each genotype were collected every 2h over 24 hours on day 11 and

pooled as indicated in Fig. 2A. Three biological replicates were collected for each genotype.

Samples were flash-frozen in liquid nitrogen, and ground into fine powder using a beadbeater.

Total RNA was extracted using an RNeasy Plant Mini Kit (Qiagen) followed by purification

using the RNA Clean & Concentrator Kit (Zymo Research). Purified RNAs were quantified

using a Nanodrop (ThermoFisher) and quality control was performed by an Agilent Bioanalyzer

2100. All samples had a 260/230 nm ratio higher than 1.9, 260/280nm ratio between 2 and 2.15,

and RIN score higher than 8. PacBio Sequel II library preparation and RNA-sequencing (Iso-Seq)

was performed by UC Davis DNA Technologies & Expression Analysis Core

(https://dnatech.genomecenter.ucdavis.edu/pacbio-library-prep-sequencing/).

Raw reads generated by the PacBio Sequel II sequencer were imported into PacBio SMRT Link

v 8.0 for Circular Consensus Sequence (CCS) calling and demultiplexing. Next, poly(A) tails

and concatemers were removed using the refine command from isoseq3 package via Bioconda (v

3.10) with the option '--require-polya'. Then the fastq files containing full-length non-

concatemer reads were mapped to Arabidopsis TAIR 10 genome assembly using minimap2 (v

2.17) with the parameter '-ax splice -t 30 -uf --secondary=no -C5'. For differential splicing

analysis, counts of exonic regions and known/novel splice junctions were generated using

QoRTs software package (v 1.3.6) (Hartley and Mullikin, 2015) with the parameter '--stranded --

singleEnded --stranded_fr_secondstrand --keepMultiMapped'. To adapt to the long-read data,

the 'maxPhredScore' was set to 93 and the 'maxReadLength' was adjusted manually for each

library. To increase the power of detecting novel splice junctions, the '--minCount' threshold was

set to 5. The raw count data was then loaded to JunctionSeq R package (v 1.5.4) (Hartley and

Mullikin, 2016) to determine differential usage of exons or splice junctions relative to the overall

expression of the corresponding gene with a false discovery rate (FDR) of 0.05. Genes with at

least one differentially spliced exon/junction were compared to the total circadian-clock-

regulated genes (Romanowski et al., 2020) as described in Fig. 4.

RNA extraction and RT-PCR

For time-course qRT-PCR analysis, Arabidopsis seedlings were entrained in 12-h light / 12-h

dark cycles for 9 days before transferred to constant light and temperature (55 μmol m^{-2} s^{-1},

22˚C). Starting from day 9, approximately 15-20 whole seedlings of each genotype were

harvested every 4h over the 72-h period. For single time point gene expression and splicing

analysis, 10-d-old seedlings grown in 12-h light / 12-h dark cycles at 22˚C were harvested at

estimated peak time of expression for each studied gene. Collected tissue was immediately

frozen in liquid nitrogen and ground into fine powder using a beadbeater. Total RNA was

isolated using TRIzol reagent (Invitrogen) and quantified with a Nanodrop (ThermoFisher). 200

ng of total RNA was used for cDNA synthesis with an oligo(dT)18 primer and SuperScriptIII

(Invitrogen) reverse transcriptase. Diluted cDNAs were used as templates for qRT-PCR and

semi-qRT-PCR reactions. The qRT-PCR was carried out as previously described (Martin-Tryon

et al., 2007) using a Bio-Rad CFX96 thermocycler. Primers amplifying the aberrantly spliced or

total transcripts were tested by standard curve and melt curve assays. Relative expression (ΔΔCq)

values were normalized to the geometric mean of *PROTEIN PHOSPHATASE 2a* (*PP2A*) and

ISOPENTENYL DIPHOSPHATE ISOMERASE 2 (*IPP2*) expression levels. For semi-qRT-PCR,

splice-junction specific primer pairs were designed to amplify regions flanking the aberrantly

spliced introns of interest. The size and abundance of the resultant PCR products were then

analyzed and quantified by LabChip GX bioanalyzer (PerkinElmer). The expression analyses

represent two to three biological replicates. All primers used in this study are described in

Supplemental Dataset 5.

Circadian period analysis

After growing on MS plates under 12-h light / 12-h dark cycles for 6 days, Arabidopsis seedlings

were sprayed with 3 mM D-luciferin (Biosynth) prepared in 0.01% (v/v) Triton X-100 (Sigma)

and then transferred to a growth chamber with a constant 22˚C and constant light provided by red

and blue LED SnapLites (Quantum Devices, 35 μmol m^{-2} s^{-1} each) for imaging. Luciferase

activity was detected using a cooled CCD camera (DU434-BV [Andor Technology]).

Bioluminescence signals from the images were quantified using MetaMorph 7.7.1.0 software

(Molecular Devices). Subsequent circadian period estimation and rhythmicity analysis were

performed using Biological Rhythms Analysis Software System 3.0 (BRASS) by fitting the

bioluminescence data to a cosine wave through Fourier Fast Transform-Non-Linear Least

Squares (Plautz et al., 1997).

Rosette size measurement

Seeds on MS plates were germinated in a growth chamber with 12-h light / 12-h dark cycles at a

constant 22˚C. 15-day-old seedlings were transferred to soil and grown in 16-h light (cool-white

fluorescent bulbs, 75 µmol m^{-2} s^{-1}) / 8-h dark long day condition at a constant 22˚C. Rosette sizes

of 39-day-old plants were determined by the greatest distance between rosette leaves using a

digital caliper.

Accession Numbers

All the *Arabidopsis thaliana* genes studied in this paper can be found under the following

accession numbers:

XCT, AT2G21150

PRP19A, AT1G04510

PRP19B, AT2G33340

PRL1, AT4G15900

SKIP, AT1G77180

CDC5, AT1G09770

LHY, AT1G01060

LNK2, AT3G54500

TIC, AT3G22380

PRR7, AT5G02810

TOC1, AT5G61380

SPPA, AT1G73990

IPP2, AT3G02780

PP2A, AT1G69960

CCA1, AT2G46830

Data Availability

All our MS data were deposited and available at Center for Computational Mass Spectrometry

(CCMS, Dataset: MSV000090830). Our PacBio Iso-Seq data were deposited and available in

NCBI Gene Expression Omnibus (GEO, Accession number: GSE220902).

Funding

This work was supported by awards from the National Institutes of Health (R01 GM069418 to

SLH) and the United States Department of Agriculture-National Institute of Food and

Agriculture (CA-D-PLB-2259-H to SLH). HZ was supported by a fellowship from China

Scholarship Council (CSC #201806010204 to HZ).

Acknowledgments

We thank the Arabidopsis Biological Resource Center for providing seeds. We thank Haiyan

Zheng and the Biological Mass Spectrometry Facility of Robert Wood Johnson Medical School

and Rutgers for mass spectrometry analysis. They are supported by NIH Shared Instrumentation

Grant S10OD01640. We thank the DNA Technologies and Expression Analysis Core at the UC

Davis Genome Center for performing PacBio Iso-Seq experiment. They are supported by NIH

Shared Instrumentation Grant 1S10OD010786-01. We thank Julin Maloof for statistical advice,

as well as members of the Harmer, Maloof, and Shabek labs for helpful discussions.

Author Contributions

HZ, RWK and SLH designed the research; HZ, RWK, and SA performed the research; HZ,

RWK, and SLH analyzed data; HZ and SLH wrote the paper.

Figure Legends

Figure 1. Simplified overview of pre-mRNA splicing reactions. A schematic diagram

highlighting the two catalytic transesterification steps and the association between the NTC and

the spliceosomal complex during pre-mRNA splicing. Gray boxes and solid black lines represent

exons and introns, respectively. U1-U6 small nuclear ribonucleoproteins (snRNPs) are indicated

by yellow circles. The NTC is indicated by a green oval.

Figure 2. Transcriptome-wide analysis reveals *XCT* as a global pre-mRNA splicing

regulator. A, Experimental design and sampling method for the PacBio Iso-Seq experiment.

White and grey boxes represent subjective day and subjective night, respectively. Arabidopsis

seedlings were grown at constant light and temperature for 10 days before being harvested and

pooled. B and C, Numbers of differentially enriched splicing events represented by significantly

differentially down- (B) or up-regulated (C) splice junctions in *xct-1*, *xct-2* and *xct-2*

complemented with p*XCT*::g*XCT-YFP-HA* compared with Col-0 (false discovery rate < 0.05). D

and E, Frequency of different classes of 5' and 3' splice sites among down- (D) or up-regulated

(E) splice junctions in *xct-1* and *xct-2*. Known or novel splice sites were classified by comparing

to TAIR10 genome annotation.

Figure 3. *XCT* is required for the fidelity of 3' splice site selection during pre-mRNA

splicing. A, A schematic diagram showing the structure of a typical U2-type splice junction. Y:

pyrimidine. B, Distribution of the distance between each pair of novel 3' splice site and its

corresponding canonical 3' splice sites in *xct-2*. C, Pictograms showing the frequency of

nucleotides in the 23-mers sequences flanking the 3' splice sites in all expressed, down-regulated

and up-regulated splice junctions in *xct-2*. D, Maximum Entropy Model scores showing the

strength of 3' splice sites of all expressed, down-regulated and up-regulated junctions in *xct-2*. E,

Percentage of pyrimidines at the -3 position (i.e. the nucleotide preceding the AG at 3' splice

site) in *xct-2*. F, Counts of pyrimidines in the Y10 polypyrimidine tract upstream of the 3' splice

sites in *xct-2*. PPT, polypyrimidine tract. The lines in the boxplot represent the 75% quartile,

median and 25% quartile of the data, respectively. Statistical significance in (D) and (F) was

determined using linear regression model with junction class as a fixed effect and is shown in

lower case letters (Tukey's multiple comparison test, $P < 0.05$). Statistical significance in (E) was

determined by Fisher's exact test: **: $P < 0.01$; ***: $P < 0.001$.

Figure 4. Dawn-phased genes are significantly enriched among all the circadian-clock-

regulated genes that are aberrantly spliced in *xct-2*. A, Venn diagram showing the overlaps of

aberrantly spliced genes in *xct-1, xct-2*, and total circadian-clock-regulated genes (Romanowski

et al., 2020). The core circadian clock genes that are mis-spliced in both *xct* mutants are indicated

in red (non-clock-regulated) and blue (clock-regulated) fonts. Only genes considered as

'detected' in all three RNA-Seq datasets are included. B and C, Phases of estimated peak

expression of all circadian clock-regulated genes (Romanowski et al., 2020) that are detected (B)

or significantly aberrantly spliced (C) in *xct-2*. The white and gray backgrounds represent the

subjective day and subjective night, respectively. D, Distribution of *P*-values from Fisher's extact

tests calculating whether the ratio of the number of aberrantly spliced genes in (B) to total clock-

regulated genes in (A) is significantly higher than expected by chance in each 2-hour interval.

Figure 5. Time-course qRT-PCR experiments validate the role of *XCT* in regulating pre-

mRNA splicing of clock-regulated and non-clock-regulated genes. A - E, Sashimi plots

showing PacBio Iso-Seq reads mapped to *LHY* (A)*, LNK2* (B)*, PRR7* (C)*, TOC1* (D) and *SPPA*

(E) in Col-0 (teal)*, xct-2* (orange) and *xct-2 XCT* (purple). The red rectangles highlight the

aberrantly spliced exon-exon junctions that are examined by qRT-PCR in (F) - (J). F - J,

Normalized expression of the aberrantly spliced isoforms of *LHY* (F)*, LNK2* (G)*, PRR7* (H)*,

TOC1 (I) and *SPPA* (J) in Col-0*, xct-2* and *xct-2 XCT*. Samples were collected every four hours

over a 72-h window. Expression levels were examined by qRT-PCR using splice-junction-

specific primers and normalized to *PP2*A and *IPP2*. Data points represent mean ± se from three

independent biological replicates. For each isoform in each biological replicate, the normalized

expression levels were relative to the highest expression levels of their total transcripts in Col-0

across all time points. Teal lines, wild type Col-0; orange lines, *xct-2* mutants; purple lines, *xct-2*

XCT. Black background, dark period; white and gray background, light period during subjective

day and night, respectively.

Figure 6. Loss of *PRP19* function suppresses the short circadian clock period phenotype

but not the splicing defects of core clock genes in *xct-2*. A, Circadian periods of Col-0, *xct-2,*

prp19a-1, *prp19b-1*, *prp19a-1 prp19b-1*, *prp19a-1 xct-2* and *prp19b-1 xct-2* plants. B - F,

Normalized expression of the aberrantly spliced isoforms of *LHY, LNK2, TIC, PRR7* and *TOC1*

in Col-0, *xct-2, prp19a-1, prp19b-1, prp19a-1 prp19b-1, prp19a-1 xct-2* and *prp19b-1 xct-2*.

Samples were collected at the estimated peak expression time for each gene. Expression levels

were examined by qRT-PCR using splice-junction-specific primers and normalized to *PP2*A and

IPP2. Data points represent mean ± se from two independent biological replicates. Statistical

significance was determined using linear regression model with genotype as a fixed effect and is

shown in lower case letters (Tukey's multiple comparison test, $P < 0.05$).

Table 1. XCT physically associates with splicing-related proteins, especially the NTC

components, in Arabidopsis. Mass Spectrometry (MS) data showing that spliceosome-

associated proteins are significantly more enriched in XCT-YFP-HA-IP than control IP (fold

enrichment > 3; *P*-value < 0.05, Welch two sample t.test). The full list of proteins co-purified

with XCT are described in Supplemental Dataset 1. Only proteins detected in each biological

replicate and with 20 or more total peptides counts are shown.

Supplemental Materials

Supplemental Figure 1. *XCT* regulates plant development and the pace of the circadian

clock in *Arabidopsis thaliana.* A, Morphology of 52-day-old Col-0, *xct-1, xct-2* mutant plants

and *xct-2* complemented with p*XCT*::g*XCT-YFP-HA*. Bar = 10 mm. B, Circadian periods of

genotypes described in (A). The lines in the boxplot represent the 75% quartile, median and 25%

quartile of the data, respectively. Statistical significance was determined using linear regression

model with genotype as a fixed effect and is shown in lower case letters (Tukey's multiple

comparison test, $P < 0.05$).

Supplemental Figure 2. *XCT* and *PRL1* control the pace of the circadian clock in an

opposite manner. A, Morphology of 21-day-old Col-0, *xct-1, xct-2* and *prl1-2* mutant

Arabidopsis seedlings. Bar = 1 mm. B, Circadian clock periods of plant genotypes described in

(A). The lines in the boxplot represent the 75% quartile, median and 25% quartile of the data,

respectively. Statistical significance was determined using linear regression model with genotype

as a fixed effect and is shown in lower case letters (Tukey's multiple comparison test, $P < 0.05$).

Supplemental Figure 3. *prl1-2* **mutant mainly induces intron retention defects in**

Arabidopsis. Frequency of different classes of 5' and 3' splice sites of down- (teal) or up-

regulated (orange) junctions in *prl1-2* compared to wild type. Known or novel splice sites were

classified based on TAIR10 genome annotation.

Supplemental Figure 4. *xct-1* **mutant displays reduced fidelity of 3' splice site selection**

during pre-mRNA splicing. A, Distribution of the distances between each mis-selected novel 3'

splice site and its corresponding canonical 3' splice site in *xct-1*. B - C, Pictograms showing the

frequency of nucleotides in the 23-mers sequences flanking the 3' splice sites in splice junctions

that are down-regulated or up-regulated in *xct-1*.

Supplemental Figure 5. The splicing defects of core clock genes are more severe in *xct-2*

than *xct-1***.** A - D, Sashimi plots showing PacBio Iso-Seq reads mapped to *CCA1* (A), *LNK2* (B),

TOC1 (C) and *TIC* (D) in Col-0 (teal)*, xct-1* (yellow) and *xct-2* (orange). The red rectangles

highlight the aberrantly spliced exon-exon junctions that are examined by semi-qRT-PCR in (E)

- (H). E - H, Relative abundance of aberrantly-spliced to fully-spliced isoforms of *CCA1* (E)*,*

LNK2 (F)*, TOC1* (G) and *TIC* (H). Differentially spliced isoforms were amplified by semi-qRT-

PCR using primers flanking the examined regions, followed by separation and quantification

using a Lab Chip GX bioanalyzer. Statistical significance was determined using linear regression

models with genotype as a fixed effect and is shown in lower case letters (Tukey's multiple

comparison test, $P < 0.05$).

Supplemental Figure 6. Dawn-phased genes are significantly enriched among genes mis-

spliced in *xct-1***.** A - B, Phases of peak expression of all the circadian-clock-regulated genes

(Romanowski et al., 2020) that are detected (A) or aberrantly spliced (B) in *xct-1*. The white and

gray backgrounds represent the subjective day and subjective night, respectively. C, Distribution

of *P*-values from Fisher's exact test calculating whether the ratio of the number of mis-spliced

genes in (B) to all clock-regulated genes in (A) is significantly higher than expected by chance

for each 2-hour interval.

Supplemental Figure 7. Loss of *XCT* function does not affect overall abundance of

transcripts of core-circadian-clock genes that are aberrantly spliced in *xct* mutants. A - F,

Normalized expression of the total transcript levels of *LHY* (A)*, LNK2* (B), *PRR7* (C) and *TOC1*

(D) in Col-0*, xct-2* and *xct-2 XCT*. Samples were collected every four hours over a 72-h window.

Expression levels were examined by qRT-PCR using primers that detect abundance of all

transcripts and normalized to *PP2*A and *IPP2.* Data points represent mean ± se from three

independent biological replicates. For each isoform in each biological replicate, the normalized

expression levels were relative to the highest expression level in Col-0 across all time points.

Teal lines, wild type Col-0; orange lines, *xct-2* muants; purple lines, *xct-2* complemented with

p*XCT*::g*XCT-YFP-HA*. Black background, dark period; white and gray background, light period

during subjective day and night, respectively.

Supplemental Figure 8. Mutation of Arabidopsis NTC members lengthens circadian

period. A - C, Circadian periods of Col-0 and NTC mutant plants, including (A) *prp19a-1*,

prp19b-1, *prp19a-1 prp19b-1*, (B) *cdc5-1*, and (C) *skip-1*. The lines in the boxplot represent the

75% quartile, median and 25% quartile of the data, respectively. Statistical significance was

determined using linear regression model with genotype as a fixed effect and is shown in lower

case letters (Tukey's multiple comparison test, *P* < 0.05).

Supplemental Figure 9. *PRP19A* (*MAC3A*) and *PRP19B* (*MAC3B*) transcript levels are

highly reduced in *prp19a-1* and *prp19b-1* mutant backgrounds, respectively. A - B,

Normalized expression of (A) *PRP19A* and (B) *PRP19B* in Col-0, *xct-2*, *prp19a-1*, *prp19b-1*,

prp19a-1 prp19b-1, *prp19a-1 xct-2* and *prp19b-1 xct-2*. Expression levels were examined by

qRT-PCR and normalized to *PP2*A and *IPP2.* Data points represent mean ± se from two

independent biological replicates. Samples were collected at ZT0. Statistical significance was

determined using linear regression model with genotype as a fixed effect and is shown in lower

case letters (Tukey's multiple comparison test, *P* < 0.05).

Supplemental Figure 10. Loss of *PRP19* function suppresses the *xct-2* short circadian

period phenotype. An independent replicate of the experiment in Figure 6A showing the

circadian periods of Col-0, *xct-2*, *prp19a-1*, *prp19b-1*, *prp19a-1 prp19b-1*, *prp19a-1 xct-2* and

prp19b-1 xct-2 plants.

Supplemental Figure 11. *PRP19* and *XCT* additively regulate Arabidopsis rosette size. A,

Morphology of 39-day-old Col-0, *xct-2*, *prp19a-1*, *prp19b-1*, *prp19a-1 prp19b-1*, *prp19b-1 xct-2*

and *prp19a-1 xct-2* plants. Bar = 10 mm. B, Rosette diameter of 36-day-old plants grown under

long-day conditions at 22˚C (n = 5-11). Data are shown as mean ± se. Statistical significance was

determined using linear regression model with genotype as a fixed effect and is shown in lower

case letters (Tukey's multiple comparison test, $P < 0.05$).

Supplemental Figure 12. Levels of total transcripts of *XCT*-targeted core clock genes are

not altered in *prp19a-1 xct-2* or *prp19b-1 xct-2* compared to *xct-2*. A - F, Normalized

expression of total transcript levels (A, C, E, F) or specific normally spliced isoforms (B, D) of

LHY, LNK2, TIC, PRR7 and *TOC1* in Col-0, *xct-2*, *prp19a-1*, *prp19b-1*, *prp19a-1 prp19b-1*,

prp19a-1 xct-2 and *prp19b-1 xct-2*. Expression levels were examined by qRT-PCR using primers

that detect abundance of all transcripts or normally spliced isoforms and normalized to *PP2*A and

IPP2. Data points represent mean ± se from two independent biological replicates. Samples were

collected at the estimated peak expression time for each gene. Statistical significance was

determined using linear regression model with genotype as a fixed effect and is shown in lower

case letters (Tukey's multiple comparison test, $P < 0.05$).

Supplemental Dataset 1. A list of proteins co-purified with XCT-IP detected by MS.

Supplemental Dataset 2. Summary of full-length reads obtained from each PacBio Iso-Seq

library.

Supplemental Dataset 3. Lists of significant differential splicing events identified by

JunctionSeq.

Supplemental Dataset 4. Lists of total and aberrantly spliced circadian-clock-regulated

genes described in this study.

Supplemental Dataset 5. A list of primers used in this study.

References

Anver S, Roguev A, Zofall M, Krogan NJ, Grewal SIS, Harmer SL (2014) Yeast X-
chromosome-associated protein 5 (Xap5) functions with H2A.Z to suppress aberrant
transcripts. EMBO Rep **15**: 894–902

**Bessonov S, Anokhina M, Krasauskas A, Golas MM, Sander B, Will CL, Urlaub H, Stark
H, Lührmann R** (2010) Characterization of purified human Bact spliceosomal
complexes reveals compositional and morphological changes during spliceosome
activation and first step catalysis. RNA **16**: 2384–2403

Bessonov S, Anokhina M, Will CL, Urlaub H, Lührmann R (2008) Isolation of an active step
I spliceosome and composition of its RNP core. Nature **452**: 846–850

Chanarat S, Seizl M, Strässer K (2011) The Prp19 complex is a novel transcription elongation
factor required for TREX occupancy at transcribed genes. Genes Dev **25**: 1147–1158

Chanarat S, Sträßer K (2013) Splicing and beyond: the many faces of the Prp19 complex.
Biochim Biophys Acta **1833**: 2126–2134

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J **16**: 735–743

Covington MF, Harmer SL (2007) The circadian clock regulates auxin signaling and responses
in Arabidopsis. PLoS Biol **5**: e222

Creux N, Harmer S (2019) Circadian Rhythms in Plants. Cold Spring Harb Perspect Biol **11**:
a034611

Deng X, Lu T, Wang L, Gu L, Sun J, Kong X, Liu C, Cao X (2016) Recruitment of the
NineTeen Complex to the activated spliceosome requires AtPRMT5. Proc Natl Acad Sci
U S A **113**: 5447–5452

Fang X, Shi Y, Lu X, Chen Z, Qi Y (2015) CMA33/XCT Regulates Small RNA Production
through Modulating the Transcription of Dicer-Like Genes in Arabidopsis. Mol Plant **8**:
1227–1236

Feke A, Liu W, Hong J, Li M-W, Lee C-M, Zhou EK, Gendron JM (2019) Decoys provide a
scalable platform for the identification of plant E3 ubiquitin ligases that regulate
circadian function. Elife **8**: e44558

Filichkin SA, Priest HD, Givan SA, Shen R, Bryant DW, Fox SE, Wong W-K, Mockler TC
(2010) Genome-wide mapping of alternative splicing in Arabidopsis thaliana. Genome
Res **20**: 45–58

Harmer SL (2009) The circadian system in higher plants. Annu Rev Plant Biol **60**: 357–377

Hartley SW, Mullikin JC (2016) Detection and visualization of differential splicing in RNA-
Seq data with JunctionSeq. Nucleic Acids Res **44**: e127

Hartley SW, Mullikin JC (2015) QoRTs: a comprehensive toolset for quality control and data
processing of RNA-Seq experiments. BMC Bioinformatics **16**: 224

Hatakeyama S, Yada M, Matsumoto M, Ishida N, Nakayama KI (2001) U box proteins as a
new family of ubiquitin-protein ligases. J Biol Chem **276**: 33111–33120

Hogg R, McGrail JC, O'Keefe RT (2010) The function of the NineTeen Complex (NTC) in
regulating spliceosome conformations and fidelity during pre-mRNA splicing. Biochem
Soc Trans **38**: 1110–1115

Horowitz DS (2012) The mechanism of the second step of pre-mRNA splicing. Wiley
Interdiscip Rev RNA **3**: 331–350

Hsu PY, Harmer SL (2012) Circadian phase has profound effects on differential expression
analysis. PLoS One **7**: e49853

Hsu PY, Harmer SL (2014) Wheels within wheels: the plant circadian system. Trends Plant Sci
19: 240–249

Idrissou M, Maréchal A (2022) The PRP19 Ubiquitin Ligase, Standing at the Cross-Roads of
886   mRNA Processing and Genome Stability. Cancers (Basel) **14**: 878

887   **James AB, Syed NH, Bordage S, Marshall J, Nimmo GA, Jenkins GI, Herzyk P, Brown**
888   **JWS, Nimmo HG** (2012) Alternative splicing mediates responses of the Arabidopsis
889   circadian clock to temperature changes. Plant Cell **24**: 961–981

890   **Jia T, Zhang B, You C, Zhang Y, Zeng L, Li S, Johnson KCM, Yu B, Li X, Chen X** (2017)
891   The Arabidopsis MOS4-Associated Complex Promotes MicroRNA Biogenesis and
892   Precursor Messenger RNA Splicing. Plant Cell **29**: 2626–2643

893   **Jones MA, Williams BA, McNicol J, Simpson CG, Brown JWS, Harmer SL** (2012) Mutation
894   of Arabidopsis spliceosomal timekeeper locus1 causes circadian clock defects. Plant Cell
895   **24**: 4066–4082

896   **Klaric JA, Wüst S, Panier S** (2021) New Faces of old Friends: Emerging new Roles of RNA-
897   Binding Proteins in the DNA Double-Strand Break Response. Front Mol Biosci **8**:
898   668821

899   **Koncz C, Dejong F, Villacorta N, Szakonyi D, Koncz Z** (2012) The spliceosome-activating
900   complex: molecular mechanisms underlying the function of a pleiotropic regulator. Front
901   Plant Sci **3**: 9

902   **Kumimoto RW, Ellison CT, Toruño TY, Bak A, Zhang H, Casteel CL, Coaker G, Harmer**
903   **SL** (2021) XAP5 CIRCADIAN TIMEKEEPER Affects Both DNA Damage Responses
904   and Immune Signaling in Arabidopsis. Front Plant Sci **12**: 707923

905   **Kwon Y-J, Park M-J, Kim S-G, Baldwin IT, Park C-M** (2014) Alternative splicing and
906   nonsense-mediated decay of circadian clock genes under environmental stress conditions
907   in Arabidopsis. BMC Plant Biol **14**: 136

908   **Lee Y-R, Khan K, Armfield-Uhas K, Srikanth S, Thompson NA, Pardo M, Yu L, Norris**
909   **JW, Peng Y, Gripp KW, et al** (2020) Mutations in FAM50A suggest that Armfield
910   XLID syndrome is a spliceosomopathy. Nat Commun **11**: 3698

Li L, Tian G, Peng H, Meng D, Wang L, Hu X, Tian C, He M, Zhou J, Chen L, et al (2018)
New class of transcription factors controls flagellar assembly by recruiting RNA
polymerase II in Chlamydomonas. Proc Natl Acad Sci U S A **115**: 4435–4440

Li Y, Yang J, Shang X, Lv W, Xia C, Wang C, Feng J, Cao Y, He H, Li L, et al (2019) SKIP
regulates environmental fitness and floral transition by forming two distinct complexes in
Arabidopsis. New Phytol **224**: 321–335

Liu L, Li X, Yuan L, Zhang G, Gao H, Xu X, Zhao H (2022) XAP5 CIRCADIAN
TIMEKEEPER specifically modulates 3' splice site recognition and is important for
circadian clock regulation partly by alternative splicing of LHY and TIC. Plant Physiol
Biochem **172**: 151–157

Maréchal A, Li J-M, Ji XY, Wu C-S, Yazinski SA, Nguyen HD, Liu S, Jiménez AE, Jin J,
Zou L (2014) PRP19 transforms into a sensor of RPA-ssDNA after DNA damage and
drives ATR activation via a ubiquitin-mediated circuitry. Mol Cell **53**: 235–246

Marshall CM, Tartaglio V, Duarte M, Harmon FG (2016) The Arabidopsis sickle Mutant
Exhibits Altered Circadian Clock Responses to Cool Temperatures and Temperature-
Dependent Alternative Splicing. Plant Cell **28**: 2560–2575

Martin-Tryon EL, Harmer SL (2008) XAP5 CIRCADIAN TIMEKEEPER coordinates light
signals for proper timing of photomorphogenesis and the circadian clock in Arabidopsis.
Plant Cell **20**: 1244–1259

Martin-Tryon EL, Kreps JA, Harmer SL (2007) GIGANTEA acts in blue light signaling and
has biochemically separable roles in circadian clock and flowering time regulation. Plant
Physiol **143**: 473–486

Mayas RM, Maita H, Staley JP (2006) Exon ligation is proofread by the DExD/H-box ATPase
Prp22p. Nat Struct Mol Biol **13**: 482–490

de Melo JRF, Gutsch A, Caluwé TD, Leloup J-C, Gonze D, Hermans C, Webb AAR,
Verbruggen N (2021) Magnesium maintains the length of the circadian period in
Arabidopsis. Plant Physiol **185**: 519–532

Meng X, Wang Q, Hao R, Li X, Li M, Hu R, Du H, Hu Z, Yu B, Li S (2022) RNA-binding
protein MAC5A interacts with the 26S proteasome to regulate DNA damage response in
Arabidopsis. Plant Physiol kiac510

Monaghan J, Xu F, Gao M, Zhao Q, Palma K, Long C, Chen S, Zhang Y, Li X (2009) Two
Prp19-like U-box proteins in the MOS4-associated complex play redundant roles in plant
innate immunity. PLoS Pathog **5**: e1000526

Nohales MA, Kay SA (2016) Molecular mechanisms at the core of the plant circadian oscillator.
Nat Struct Mol Biol **23**: 1061–1069

Nolte C, Staiger D (2015) RNA around the clock - regulation at the RNA level in biological
timing. Front Plant Sci **6**: 311

Perez-Santángelo S, Mancini E, Francey LJ, Schlaen RG, Chernomoretz A, Hogenesch JB,
Yanovsky MJ (2014) Role for LSM genes in the regulation of circadian rhythms. Proc
Natl Acad Sci U S A **111**: 15166–15171

Plautz JD, Straume M, Stanewsky R, Jamison CF, Brandes C, Dowse HB, Hall JC, Kay SA
(1997) Quantitative analysis of Drosophila period gene transcription in living animals. J
Biol Rhythms **12**: 204–217

Romanowski A, Schlaen RG, Perez-Santangelo S, Mancini E, Yanovsky MJ (2020) Global
transcriptome analysis reveals circadian control of splicing events in Arabidopsis thaliana.
Plant J **103**: 889–902

Sanchez SE, Petrillo E, Beckwith EJ, Zhang X, Rugnone ML, Hernando CE, Cuevas JC,
Godoy Herz MA, Depetris-Chauvin A, Simpson CG, et al (2010) A methyl transferase
links the circadian clock to the regulation of alternative splicing. Nature **468**: 112–116

Schlaen RG, Mancini E, Sanchez SE, Perez-Santángelo S, Rugnone ML, Simpson CG,
Brown JWS, Zhang X, Chernomoretz A, Yanovsky MJ (2015) The spliceosome
assembly factor GEMIN2 attenuates the effects of temperature on alternative splicing and
circadian rhythms. Proc Natl Acad Sci U S A **112**: 9382–9387

Semlow DR, Blanco MR, Walter NG, Staley JP (2016) Spliceosomal DEAH-Box ATPases
Remodel Pre-mRNA to Activate Alternative Splice Sites. Cell **164**: 985–998

Seo PJ, Park M-J, Lim M-H, Kim S-G, Lee M, Baldwin IT, Park C-M (2012) A self-
regulatory circuit of CIRCADIAN CLOCK-ASSOCIATED1 underlies the circadian
clock regulation of temperature responses in Arabidopsis. Plant Cell **24**: 2427–2442

Shakhmantsir I, Sehgal A (2019) Splicing the Clock to Maintain and Entrain Circadian
Rhythms. J Biol Rhythms **34**: 584–595

Shi Y (2017) Mechanistic insights into precursor messenger RNA splicing by the spliceosome.
Nat Rev Mol Cell Biol **18**: 655–670

Uehara TN, Nonoyama T, Taki K, Kuwata K, Sato A, Fujimoto KJ, Hirota T, Matsuo H,
Maeda AE, Ono A, et al (2022) Phosphorylation of RNA Polymerase II by CDKC;2
Maintains the Arabidopsis Circadian Clock Period. Plant Cell Physiol **63**: 450–462

Wang B-B, Brendel V (2006) Genomewide comparative analysis of alternative splicing in
plants. Proc Natl Acad Sci U S A **103**: 7175–7180

Wang X, Wu F, Xie Q, Wang H, Wang Y, Yue Y, Gahura O, Ma S, Liu L, Cao Y, et al
(2012) SKIP is a component of the spliceosome linking alternative splicing and the
circadian clock in Arabidopsis. Plant Cell **24**: 3278–3295

Wei Y, Yee PP, Liu Z, Zhang L, Guo H, Zheng H, Anderson B, Gulley M, Li W (2020)
NEDD4L-mediated Merlin ubiquitination facilitates Hippo pathway activation. EMBO
Rep **21**: e50642

Wilkinson ME, Charenton C, Nagai K (2020) RNA Splicing by the Spliceosome. Annu Rev
Biochem **89**: 359–388

Xu Y-J, Lei Y, Li R, Zhang L-L, Zhao Z-X, Zhao J-H, Fan J, Li Y, Yang H, Shang J, et al
(2017) XAP5 CIRCADIAN TIMEKEEPER Positively Regulates RESISTANCE TO
POWDERY MILDEW8.1-Mediated Immunity in Arabidopsis. Front Plant Sci **8**: 2044

Yang Y, Li Y, Sancar A, Oztas O (2020) The circadian clock shapes the Arabidopsis
transcriptome by regulating alternative splicing and alternative polyadenylation. J Biol
Chem **295**: 7608–7619

Zhan X, Lu Y, Zhang X, Yan C, Shi Y (2022) Mechanism of exon ligation by human
spliceosome. Mol Cell **82**: 2769-2778.e4

Zhang S, Liu Y, Yu B (2014) PRL1, an RNA-binding protein, positively regulates the
accumulation of miRNAs and siRNAs in Arabidopsis. PLoS Genet **10**: e1004841

Zhang S, Xie M, Ren G, Yu B (2013) CDC5, a DNA binding protein, positively regulates
posttranscriptional processing and/or transcription of primary microRNA transcripts.
Proc Natl Acad Sci U S A **110**: 17588–17593

Table 1. XCT physically associates with splicing-related proteins, especially the NTC components, in Arabidopsis. Mass Spectrometry (MS) data showing that spliceosome-associated proteins are significantly more enriched in XCT-YFP-HA-IP than control IP (fold enrichment > 3; P-value < 0.05, Welch two sample t.test). The full list of proteins co-purified with XCT are described in Supplemental Dataset 1. Only proteins detected in each biological replicate and with 20 or more total peptides counts are shown.

AGI	*A. thaliana* gene name	*P*-value	fold enrichment	category	*S. cerevisiae* homolog	*H. sapiens* homolog
AT2G21150	XCT	6.21E-04	Inf			FAM50A
AT1G04510	PRP19A/MAC3A/PUB59	1.52E-02	4.2	NTC	Prp19p	PRPF19; PSO4; SNEV
AT2G33340	PRP19B/MAC3B/PUB60	2.17E-03	4.8	NTC		
AT4G15900	PRL1/MAC2	1.24E-02	7.7	NTC	Prp46p	PLRG1
AT1G77180	SKIP/MAC6	1.36E-03	4.9	NTC	Prp45p	SNW1
AT1G09770	CDC5/MAC1	4.06E-03	6.7	NTC	Cef1p	CDC5L
AT1G07360	MAC5A	2.15E-03	8.8	NTC	Ecm2p	RBM22
AT5G41770	CRNK1/MAC10	5.78E-03	7.7	NTC	Clf1p/Syf3	CRNKL1/SYF3
AT3G18790	ISY1/MAC8	1.26E-03	6.1	NTC	Isy1p	ISY1
AT2G38770	EMB2765/MAC7	1.36E-02	4.4	NTC	Cwf11p	AQR
AT5G28740		8.86E-03	4.6	NTC	Syf1p	XAB2
AT3G18165	MOS4	4.09E-03	4.5	NTC	Snt309p	BCAS2/SPF27
AT5G51280		1.65E-02	4.3	other splicing	Dbp2p	DDX41
AT2G47640		4.93E-03	4.3	other splicing	Smd2p	SNRPD2
AT3G26560		8.15E-03	24	other splicing	Prp22p	DHX8
AT1G09760	U2A'	3.51E-02	3.3	U2 complex	Lea1p	SNRPA1