Switching between external and internal attention

in hippocampal networks

Craig Poskanzer and Mariam Aly

Department of Psychology, Columbia University

Please address correspondence to:
Craig Poskanzer
crp2170@columbia.edu
Columbia University
Department of Psychology
406 Schermerhorn Hall
1190 Amsterdam Avenue MC 5501
New York, NY 10027

ORCiD:

Craig Poskanzer: https://orcid.org/0000-0002-4457-4941
Mariam Aly: https://orcid.org/0000-0003-4033-6134

Acknowledgements: We would like to thank Dr. Serra Favila, Dr. Katherine Duncan, and the Alyssano Group for helpful feedback on this project. We also thank Dr. Eren Günseli for providing the dataset that is analyzed in the current work. This work was funded by an NSF CAREER Award (BCS-1844241) to M.A.

Abstract

Everyday experience requires processing external signals from the world around us and internal information retrieved from memory. To do both, the brain must fluctuate between states that are optimized for external vs. internal attention. Here, we focus on the hippocampus as a region that may serve at the interface between these forms of attention, and ask how it switches between prioritizing sensory signals from the external world vs. internal signals related to memories and thoughts. Pharmacological, computational, and animal studies have identified input from the cholinergic basal forebrain as important for biasing the hippocampus towards processing external information, whereas complementary research has suggested the dorsal attention network (DAN) may aid in allocating attentional resources towards accessing internal information. We therefore tested the hypothesis that the basal forebrain and DAN drive the hippocampus towards external and internal attention, respectively. We used data from 29 human participants (17 female) who completed 2 attention tasks during fMRI. One task primarily required external attention ("explicitly instructed" task); the other required switching between external and internal attention ("memory-guided" task). We discovered that background connectivity between the basal forebrain and hippocampus was stronger during the explicitly instructed vs. memory-guided task. In contrast, DAN-hippocampus background connectivity was stronger during the memory-guided vs. explicitly instructed task. Finally, the strength of DAN-hippocampus connectivity correlated with performance on the memory-guided but not explicitly instructed task. Together, these results provide evidence that the hippocampus may switch between external and internal states by preferentially communicating with the basal forebrain and DAN.

Significance Statement

How does the brain balance the need to pay attention to internal thoughts and external sensations? We focused on the human hippocampus, a region that may serve at the interface between internal and external attention, and asked how it varies its functional connectivity based on attentional states. The hippocampus was more strongly coupled with the cholinergic basal forebrain during tasks that emphasized external vs. internal attention. This pattern flipped for connectivity between the hippocampus and dorsal attention network, which was higher for attention tasks that were guided by memory rather than external cues. Together, these findings elucidate how the hippocampus may coordinate switches between external and internal attention by modulating its communication with distinct networks in the brain.

Introduction

While navigating daily life, we must balance paying attention to sensory stimuli streaming in from the external world and paying attention to our internal thoughts and memories. To accomplish this, the brain must dynamically shift between states in which attention is externally vs. internally oriented (Chun et al., 2011; Verschooren et al., 2019; Li et al., 2022). This flexible reorienting of attention allows us to both acquire new information and leverage previously stored information, which is critical for interacting with our environment, making decisions, and planning future actions (Lepsien & Nobre, 2006).

What neural mechanisms allow the brain to efficiently switch between external and internal processing? Although a burgeoning line of research addresses this question (Honey et al., 2017; Verschooren et al., 2019; Li et al., 2022), much of this work focuses on attentional networks. Here, we test how attentional networks might differentially interact with the hippocampus to switch between internal and external states.

Although the majority of literature on the hippocampus explores its role in episodic and relational memory, recent work has demonstrated its involvement in external attention. For example, hippocampal activity patterns represent top-down attentional goals (Aly & Turk-Browne, 2016a, 2016b; Córdova et al., 2019). Moreover, individuals with medial temporal lobe damage are impaired on relational attention tasks that require spatial processing (Ruiz et al., 2020). The hippocampus is also involved in guiding external attention by leveraging information from memory (Hutchinson & Turk-Browne, 2012; Günseli & Aly, 2020). Thus, the hippocampus is ideally situated to coordinate internal and external attention. Indeed, computational and rodent work suggests that the hippocampus oscillates between attentional states that prioritize processing of either external or internal input (Honey et al., 2017; Tarder-Stoll, Jayakumar, et al., 2020). The peak vs. trough of local theta cycles in the hippocampus may increase the strength of external and internal input respectively (Hasselmo et al., 2002; Honey et al., 2017). To better understand these state switches, it is necessary to explore what drives this switching, and which brain areas bias the hippocampus towards external vs. internal attention.

One area that may drive hippocampal switching is the basal forebrain, which provides acetylcholine to the hippocampus (Newman et al., 2012). The basal forebrain promotes external attention through cholinergic projections across the cortex (Villano et al., 2017). Within the hippocampus, acetylcholine strengthens afferent input about the external world and suppresses recurrent connections that support internally oriented processing (Newman 2006; Tarder-Stoll, Jayakumar, et al., 2020; Decker & Duncan, 2020). At a behavioral level, acetylcholine supports encoding (Newman et al., 2012), and cholinergic agonists benefit performance on a hippocampally mediated external attention task (Ruiz, Thieu, & Aly, 2021). This combination of animal, computational, and human behavioral work provides compelling evidence that the basal forebrain may bias the hippocampus into an externally oriented state.

Conversely, the dorsal attention network (DAN) may bias the hippocampus toward an internally oriented state. The DAN, along with the hippocampus, is recruited when accessing long-term memories to direct attention (Stokes et al., 2012) and dorsal parietal damage (overlapping with the DAN) impairs episodic recall (Berryhill et al., 2007). More generally, dorsal parietal regions, overlapping with the DAN, may allocate attentional resources towards mnemonic processing (Cabeza, 2008; Cabeza et al., 2008; Wagner et al., 2005). This "attention to memory" (AtoM) hypothesis suggests that the DAN may interact with the hippocampus to access information from memory, and may thus be an important network for biasing the hippocampus towards internal attention.

Here, we evaluate this two-fold hypothesis in which the basal forebrain and DAN dynamically interact with the hippocampus to bias external and internal attentional states respectively. We leverage two attention tasks that vary in their requirements to switch between externally and internally oriented attention. We hypothesize that hippocampal connectivity with the basal forebrain and DAN will switch based on attentional goals, with higher hippocampus-basal forebrain connectivity during tasks that are primarily externally vs. internally oriented, and an inversion of this pattern for hippocampus-DAN connectivity.

Methods

To test these hypotheses, we analyzed a dataset (Günseli & Aly, 2020) in which participants performed 2 attention tasks, one in which they searched for rooms with the same spatial layout and one in which they searched for paintings with a similar artistic style. These 2 attentional states each took place in 2 conditions: memory-guided attention and explicitly instructed attention. These conditions were identical in terms of stimuli and motor demands; they differed only in whether participants selected their attentional state (art or room attention) based on their memory for images in the preceding trial ("memory-guided" condition) or were assigned an attentional state at the beginning of each trial ("explicitly instructed" condition). These attentional states and task conditions strongly modulate the hippocampus (Günseli & Aly, 2020; also see Aly & Turk-Browne, 2016a, 2016b; Ruiz et al., 2020) but differ in that only the memory-guided task requires switching between external attention and long-term memory retrieval. This allowed us to examine the hippocampal network mechanisms that vary between attention tasks that are relatively externally oriented and those that require switches between internal and external attention.

Participants

Participants, stimuli, and tasks were previously described in Günseli & Aly (2020) and are summarized again here. A total of 30 participants were recruited from the Columbia University

community. Participants provided written, informed consent and were paid $72 for their participation. As in Günseli & Aly (2020), we analyzed data from 29 participants, excluding 1 whose performance was three standard deviations below average on the memory-guided attention task. The included participants (17 female; 1 left-handed; all normal/corrected-to-normal vision) were aged 18-35 (M = 26; SD = 4.07) and had 13-21 years of education (M = 17.1; SD = 2.2). Study procedures were approved by the Columbia University Institutional Review Board.

Stimuli

A total of 141 images were created for the main fMRI task by combining a picture of a 3D-rendered room with an image of a painting. An initial 20 virtual rooms were rendered using Sweet Home 3D (sweethome3d.com). They were designed to have a unique combination of shape, furniture, and wall color. After capturing an image of each room, the viewpoint was rotated 30° (50% of rooms were rotated clockwise and 50% counterclockwise). Next, these rotated images were edited while preserving their spatial layout: the wall color was changed and all furniture was substituted with a different piece of the same category in the same position (e.g. substituting a couch with a different couch). A second image was then captured of this spatially consistent room, which we refer to as a "room match" to the original image.

20 painting images were initially selected from the Google Arts & Culture Project (https://artsandculture.google.com/). Each art image was then paired with an "art match" by selecting a second painting by the same artist. These 2 paintings were thus stylistically similar to each other, while containing unique features. To create a combined art/room image, a painting image was superimposed on a room image to appear as if it were hanging on one of the visible walls.

A set of images were then selected to be "stay" or "switch" cues. The presence of a "stay" cue in a given trial in the memory-guided condition would indicate that the participant should stay in the same attentional state (art or room) on the following trial; the presence of a "switch" cue would indicate that they should switch to the other attentional state on the following trial (**Figure 1A**). Stay/switch cues were created by selecting 2 rooms (1 stay cue and 1 switch cue) and 2 paintings (1 stay cue and 1 switch cue). Each room cue was paired with 3 selected paintings to form 3 room "stay" images and 3 room "switch" images; similarly, each painting cue was paired with 3 room images to form 3 painting "stay" images and 3 painting "switch" images. To create parity in the non-attended "background" of the stay/switch cues (i.e., "background" paintings for room stay/switch cues and "background" rooms for painting stay/swich cues), the same 3 paintings were used for both room "stay" and "switch" cues, and the same 3 rooms were used for both art "stay" and "switch" cues. This ensured that participants could not identify whether a given (art or room) cue was "stay" or "switch" based on the unattended "background" feature (room or art, respectively). The non-attended "backgrounds" of the stay/switch cues were also used to make

art/room images that were not "stay" or "switch" cues. In particular, each of the 3 room backgrounds for art "stay" and "switch" cues was also paired with 2 painting images that were not "stay" or "switch" cues; and each of the paintings paired with room "stay" and "switch" cues was also paired with 2 rooms that were not "stay" or "switch" cues. This ensured that participants could not use these "background" features to identify that a "stay" or "switch" cue had been presented. In sum, this procedure resulted in 12 images in which stay/switch cues were embedded; for all of these, the "background" features were not indicative of the cue being presented.

The complete set of 129 main-task images (excluding the 12 stay/switch cue images) used in the fMRI task was created by permuting the 43 room images (20 rooms in 2 perspectives each and 3 "background" rooms also used with art stay/switch cues) with 43 paintings (20 artists with 2 paintings each and 3 "background" paintings also used with room stay/switch cues) such that each painting was paired with multiple rooms and each room was paired with multiple paintings.

From the set of 129 main-task images, 20 images were selected as "base images" (**Figure 1B**). Each base image was included in a set with 6 other images to form a "base set." In addition to the base image, a base set included: a room match (another image containing a room with the same spatial layout as the room in the base image, shown from a different angle), an art match (an image containing a painting from the same artist as the painting in the base image), and 4 "distractor" images containing rooms and paintings that were neither art nor room matches to the base image, nor included any paintings or rooms used elsewhere in the base set. Base images could only be used in their own set, but art/room matches and distractors could also appear in other base sets. No single image in the base set could serve as **both** an art match and a room match to a single base image. For a single trial in the fMRI task, a base image was shown followed by 4 "search" images selected from the base set (art match, room match, 4 distractors) and pool of stay/switch cue images. "Stay" and "switch" cues on a given trial were always consistent with the attended dimension on that trial, i.e., art "stay" and "switch" cues only appeared on art attention trials, and room "stay" and "switch" cues only appeared on room attention trials. Each base set was used to create 5 trials for the explicitly instructed attention task and 5 trials for the memory-guided attention task; these trials differed in the images presented in the search set and the order in which they were presented. Art and room matches were independently likely to appear, such that an equal number of trials had both matches present in the search set, neither present, an art match only, or a room match only.

A second, separate set of stimuli was generated for a practice session, which occurred approximately 2 days before the scan. These practice stimuli consisted of 12 images with embedded stay/switch cues and 70 practice-task images. These 70 practice-task images were divided into 10 base sets consisting of a base image, art match, room match, and 4 distractors. Twelve novel images with embedded stay/switch cues were created using the same method described above. All practice stimuli were distinct from the fMRI task images — including the

practice "stay" and "switch" cues — so that participants could learn the task while still preserving the novelty of the fMRI task stimuli.

A third, separate set of 82 stimuli was created for a practice session on the day of the scan. This practice set contained 70 novel art/room images as well as 12 images with embedded stay/switch cues. Importantly, the stay/switch cues used in this practice session were the **same** art and room images that would subsequently be used in the fMRI task; however, the non-attended "backgrounds" paired with each cue (3 room images and 3 paintings) were selected from the set of images used to create the 70 novel art/room stimuli for this practice session. In this way, participants could practice using the stay/switch cues that would ultimately be used in fMRI, while decoupling this practice from the unattended "backgrounds" of the stay/switch cues that would be present in the fMRI task.

Task Overview

Participants performed an attention task on the combined art/room images described above (**Figure 1C**). On each trial, participants were first shown a base image, followed by a search set of 4 images. There were 2 potential attentional states: room attention and art attention. On a room attention trial, the participant paid attention to the spatial layout of the rooms; on an art attention trial, the participant paid attention to the artistic style of the paintings embedded in the rooms. Specifically, on a room attention trial, participants had to search the search set for a room match (an image showing a room with the same spatial layout as the base image, but from a different perspective); on an art attention trial, participants had to search the search set for an art match (an image showing a painting with a similar artistic style as the base image).

Figure 1: Task Overview. (A) Before entering the fMRI scanner, participants learned 2 "stay" cues (one painting, one room) and 2 "switch" cues (one painting, one room). These cues were embedded into the set of search images for the attention task in the fMRI scanner, and indicated how the participant should direct their attention in the following trial during the memory-guided condition. Stay cues instructed the participants to continue in the same attentional state for the following trial (i.e., art→art or room→room). Switch cues instructed the participants to change their attentional state for the following trial (i.e., art→room or room→art). Stay/switch cues were also embedded in the search images for the explicitly instructed condition but were not relevant for that task; stay/switch cues were only informative in the memory-guided condition. **(B)** For each trial of the attention task, participants saw a single "base image" of a virtual room with a painting hanging on a wall. Participants could either pay attention to the spatial layout of the room in the image *or* to the artistic style of the art on the wall. Depending on their attentional goal, participants were instructed to look for an image that either showed the same room as the base image from a different perspective (room trial) or a painting with a similar style as that in the base image (art trial). Shown are a sample base image and its room match and art match. **(C)** The attention task consisted of two conditions that occurred in different task runs: the explicitly instructed condition and the memory-guided condition. In each trial of the explicitly instructed condition, participants were randomly assigned either a room or art attentional state. Next, they were shown a base image and a series of search images. Finally, they were asked whether the search set contained an image that matched the room in the base image (same room from a different perspective, "room" probe) or the art in the base image (painting with a similar style, "art" probe). The memory-guided task had a similar structure, except that participants also had to attend to the stay or switch cue that was embedded in the search images. The stay or switch cue on a given trial was then used to select the attentional state at the start of the subsequent trial (e.g., a "stay" cue in a "room" trial, as shown above, should result in "room" being selected as the attentional state on the following trial). One third of trials did not contain a stay or switch cue; on these "no-cue" trials, participants were free to select either art or room on the subsequent trial. Room stay/switch cues were only presented on room trials and art stay/switch cues were only presented during art trials.

Both of these attention tasks (art and room) took place in 2 conditions: explicitly instructed attention and memory-guided attention. In the explicitly instructed condition, the participant was directed at the start of each trial to pay attention to either the room or the art with an explicit cue ("room" or "art"). Conversely, in the memory-guided condition, the participant searched for an embedded stay/switch cue — that could appear anywhere in the search set on a given trial — which indicates to them to either stay in the same attentional state in the following trial ("stay" cue), or to switch attentional states in the following trial ("switch" cue).

Critically, the stimuli were identical for the explicitly instructed and memory-guided conditions. Thus, the stay/switch cue images were also embedded in the search set for the explicitly instructed condition; they were just irrelevant to participants' attentional state. This allowed us to compare these conditions without a confound of different stimuli.

Participants completed 4 adjacent runs (25 trials each) of the explicitly instructed condition and an additional 4 adjacent runs (25 trials each) of the memory-guided condition. The order of the conditions was counterbalanced across participants.

Task Design

All stimuli and instructions were presented using Psychtoolbox for MATLAB. Each trial began with the presentation of an initiation screen. For the explicitly instructed task, the initiation screen read "Press any key with the left hand to start the trial". For the memory-guided condition, the initiation screen read "Press left index for Room, left middle for Art." The structure of the rest of the trial was identical across the explicitly instructed and memory-guided conditions.

After the initiation screen, participants were shown their attentional state for the trial: "ROOM" for trials in which they were supposed to search for a room match and "ART" for trials in which they were supposed to search for an art match. This attentional state screen was presented for either 1.5s, 2s, or 2.5s (randomly assigned across trials). In the explicitly instructed condition, this attentional state was randomly assigned; however, in the memory-guided condition, the attentional state was determined by the participant's key press during the initiation screen. On the first trial of a memory-guided run (25 trials), the participant was free to select whichever attentional state they preferred. On the following trials of the memory-guided run, participants used stay/switch cues embedded in trial N to select their attentional state in trial N+1. A "stay" cue indicated that the participant should maintain the same attentional state in the subsequent trial (e.g., if trial N were a room trial, a stay cue indicated that trial N+1 should also be a room trial). A "switch" cue indicated that the participant should choose the opposite attentional state on the subsequent trial (e.g., if trial N were a room trial, a "switch" cue indicated that trial N+1 should be an art trial). If a participant incorrectly selected their attentional state at the start of a memory-guided trial, the trial continued with whichever attentional state they selected. One-third

of trials did not contain a stay/switch cue, and participants were told to select either attentional state on the following trial if this occurred. Participants were instructed to attempt to balance their choice of art and room attentional states following "no-cue" trials, but on average, they selected room trials more often (as reported in Günseli & Aly, 2020: average number of selected room trials = 16.828, 95% CI [16.299, 17.356]; average number of selected art trials = 14.655, 95% CI [14.144, 15.166]; t(28) = 5.90, p < 0.00001, d = 1.10, 95% CI [1.418, 2.927]). No-cue trials were not analyzed further because our main hypotheses were about differences in the memory-guided and explicitly instructed conditions that arise due to the stay/switch cues in the search set; these cues, in the memory-guided task, may cause participants to switch from an externally oriented mode to an internally oriented one, to retrieve the meaning of the cue.

After the initiation screen and the attentional state screen, participants were shown a base image for 2s. Depending on the attentional state (art or room), participants could either pay attention to the spatial layout of the room in the base image or to the style of the art in the base image. Next, participants saw 4 search images (selected from the trial's base set and set of stay/switch cue images) each presented for 1.25s with a 0.1s inter-stimulus interval (ISI). Finally, 0.1s after the last search image, participants were shown a probe screen (2s) asking either "ART?" or "ROOM?" Participants responded "yes" or "no" to indicate if there was a match in the probed dimension using their right-hand index finger (match) or middle finger (no-match) on a button box. When presented with the probe "ART?" participants had to respond if any of the search images contained a painting that could have been painted by the same artist as the painting in the base image. After a "ROOM?" probe, participants responded whether or not one of the search images contained a room with the same spatial layout as the room in the base image. For 80% of trials across all runs, the probe matched the cue on the attentional state screen (e.g., if the cued attentional state was room, the probe read "ROOM?"); these were termed "valid" trials. However, 20% of trials across all runs were "invalid", meaning the probe did not match the attentional state cue (e.g., if the cued attentional state was room, the probe read "ART?"). Comparing performance on valid vs. invalid trials allowed us to determine whether attention conferred a performance benefit (Posner, 1980; Stokes et al., 2012), and whether this performance benefit was matched across the memory-guided and explicitly instructed conditions. (Behavioral performance — including performance on valid trials only and the valid vs. invalid difference — was not different between these conditions, as noted in Günseli & Aly, 2020.)

Between trials, a blank screen was presented for a variable time (the inter-trial interval, ITI). Twenty-five ITI lengths were generated (truncated exponential, lambda = 1.5, mean = 6.66s, T = 9s) and were randomly shuffled to be used after each of the 25 trials across all 8 runs of the task.

After each run, participants were shown the percent of responses to the art/room probe that they had answered correctly. In memory-guided runs, participants were also shown the percent of stay/switch cues that they correctly identified.

The order of trials within a run was structured such that each of the 20 base images was shown once every 20 trials, no two adjacent trials could have the same base image, and no image could appear multiple times in the same trial. Matches for the probed attentional state were present in 50% of trials; independently, matches for the non-probed attentional state were present in 50% of trials. Stay/switch cues were presented in ⅔ of trials in both the explicitly instructed and memory-guided conditions (though they were only informative in the memory-guided condition). Finally, within each condition (explicitly instructed and memory-guided), 80% of trials were valid. Valid trials were equally distributed across the art/room attentional states, presence of probed match (present/absent), presence of non-probed match (present/absent), and stay/switch cue presence (stay cue/switch cue/neither cue). It was not possible to perfectly equate the number of valid trials across each of these trial types for a given participant, but valid trials were balanced across these trial types across every 6 participants.

The design of the practice sessions mirrored the structure of the main experiment with the exception of the ITI lengths, which were randomly selected as either 2s or 2.5s. Additionally, for the practice sessions, participants were provided with feedback about their performance following their response to the art/room probe, and in the memory-guided condition, their selection of an attentional state.

Procedure

The study was completed in 3 phases: an initial practice session (approximately 2 days before the fMRI scan), a second practice session (just before the scan), and the fMRI task. The initial practice session gave the participants an opportunity to learn the task and practice the explicitly instructed and memory-guided conditions with an entirely separate set of stimuli from those used in the fMRI task. Similarly, the second practice session allowed the participants to practice the task using another (mostly) separate set of stimuli from those in the subsequent fMRI task; however, we included the same stay/switch cues as those in the fMRI task to ensure that they were learned prior to the start of fMRI.

Both practice sessions began with a run of 10 trials of the explicitly instructed condition. Participants repeated this run until they answered at least 65% of the validly cued art/room probes correctly. Following practice of the explicitly instructed condition, participants entered a cue learning phase, in which they were shown each of the stay/switch cues and their meanings 4 times in a randomly shuffled order. Each image was shown for a minimum of 1 second, along with its meaning (i.e., stay or switch), and the participant had to press a button to advance to the next image. Participants were then tested on the stay/switch cues: they were shown all stay/switch cues one at a time (without their meanings) in a shuffled order. The participant had to correctly identify each of the stay/switch cues with a button press (to indicate whether the cue meant "stay" or "switch") 5 times in a row, or else they would have to restart the cue learning sequence. Upon completion of the cue learning phase, participants performed a practice run of 10 trials of

the memory-guided condition. This run was repeated until participants met two criteria: 1) they answered at least 65% of the validly cued art/room probes correctly and 2) they used stay/switch cues to correctly identify the attentional state (art or room) for the following trial in at least 80% of the trials.

After the second practice session, participants entered the scanner. The fMRI task consisted of 8 runs of 25 trials each. The explicitly instructed and memory-guided conditions each contained 100 trials (4 contiguous runs), with condition order counterbalanced across participants. Before each condition, participants completed 5 practice trials of the upcoming condition (explicitly instructed or memory-guided); practice trials were repeated until participants achieved at least 65% accuracy on the art/room probes and, in the memory-guided practice, correctly identified the attentional state (art or room) from the stay/switch cues in 80% of the trials. After each run of the memory-guided condition, participants were shown all 4 stay/switch cues (2 art images and 2 room images) and their meanings as a reminder. If on any run of the memory-guided condition, a participant was not able to correctly select the attentional state (art or room) in at least 85% of the trials, they repeated the cue learning procedure from the previous practice sessions before advancing to the next run.

MRI Image Acquisition

All MRI scans were performed in a 3T Siemens Magnetom Prisma scanner using a 64-channel coil. Functional scans (8 task runs) used a multiband echo-planar imaging (EPI) sequence with the following settings: TR = 1.5s, echo time = 30ms, flip angle = 65°, acceleration factor = 3, voxel size = 2mm iso, phase encoding direction = posterior to anterior. 69 oblique axial slices (14° transverse to coronal) were collected in an interleaved order. Each participant also underwent a T1-weighted anatomical scan (resolution = 1mm iso) using a magnetization-prepared rapid acquisition gradient-echo sequence (MPRAGE). Field maps were collected for each participant using 69 oblique axial slices (resolution = 2mm iso).

Regions of Interest

We focused on 4 bilateral regions of interest (ROIs): the hippocampus, the dorsal attention network (DAN), the ventral attention network (VAN), and the basal forebrain (**Figure 2**). The hippocampus was defined using the Harvard-Oxford atlas in FSL thresholded at 0.5 (Jenkinson et al. 2012). Both the DAN and VAN were defined using the procedure outlined in Jimenez et al. (2016). The DAN was defined by using FSLMATHS to create 9mm radius spheres around coordinates in the bilateral lateral frontal cortex centered at [+/- 48, 8, 35], anterior intraparietal cortex centered at [+/- 30, -58, 45], and posterior intraparietal cortex centered at [+/-46, -43, 45]. The VAN was defined by creating 9mm radius spheres around coordinates in the bilateral temporoparietal junction centered at [+/- 53, -34, 21], anterior cingulate cortex centered at [+/- 3, 20, 36], and anterior insula centered at [+/-32, 22, 4]. Finally, the basal forebrain region was

defined using a probabilistic atlas from Zaborszky et al. (2021), thresholded at 0.5. All ROIs were binarized and registered to 1mm standard space.

Figure 2: Regions of Interest. Hippocampus (green), basal forebrain (red), dorsal attention network (DAN) (teal), and ventral attention network (VAN) (purple) are shown on a 1mm T1 template image in MNI152 standard space. All regions of interest were bilateral.

Preprocessing

Data were preprocessed using FEAT, FNIRT, and FSL tools (as reported in Günseli & Aly, 2020). The first 4 volumes of each functional run were removed to account for T1 equilibration; for 1 participant, only 1 volume was removed at the start of each functional run (because not enough volumes were collected at the start of each run to discard 4). Both brain extraction (BET; Smith, 2002) and motion correction (MCFLIRT; Jenkinson et al., 2002) were performed. Data were high-pass filtered (cut-off = 128s) and spatially smoothed using a 3mm full width at half maximum Gaussian kernel. Custom scripts were generated based on the FUGUE guide (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FUGUE/Guide) to preprocess field maps. This process consisted of using the skull-stripped average of 2 magnitude images along with the phase image to generate a field map image with the fsl_prepare_fieldmap command. Field map images and magnitude images were used in FEAT preprocessing. Finally, functional images were registered to the standard MNI152 1mm anatomical image using a nonlinear warp (10mm resolution and 12 degrees of freedom). This preprocessing procedure was used in both GLMs described below.

Univariate Activity

Data were modeled using a single-trial GLM following the procedure in Günseli & Aly (2020). Each trial was made up of 3 parts (**Figure 1C**): first, an *orienting period* spanning the time from the onset of the initiation screen to the offset of the attentional state screen; second, an *image period*, spanning the 7.4s epoch from the onset of the base image until the offset of the final search image; and finally, a *probe period* spanning the 2s duration of the art/room probe screen. Each image period was modeled separately (as a 7.4s epoch); 1 additional regressor was included to model all orienting periods (modeled as epochs of varying duration depending on participant response time (RT) during the initiation screen and the length of the attentional cue); and a final

regressor that modeled all probe periods (as 2s epochs). All regressors were convolved with a double-gamma hemodynamic response function as implemented in FEAT. To account for movement-related noise, 6 motion regressors were included in the model. FILM prewhitening was used to remove autocorrelation. Finally, each task run was modeled separately. All regressors of this model were registered to high-resolution 1mm MNI152 standard space.

The task included both valid trials (in which the cue and probe were the same) and invalid trials (in which participants were cued to attend to one dimension, e.g., art, but probed about the other, e.g., room). Invalid vs. valid probes lead to behavioral costs in response times and sensitivity (Aly & Turk-Browne, 2016a, b; Gunseli & Aly, 2020) and enhanced activity in the brain's reorienting network, overlapping with both the ventral and dorsal attention networks (Aly & Turk-Browne, 2016a; Corbetta & Shulman, 2002). For these reasons, analyses of univariate activity (as well as subsequent analyses of brain-behavior correlations) focused on valid trials.

We examined univariate activity (and background connectivity, see below) for the *image period*, because this part of the task differentially taxed internal vs. external attention in our 2 conditions: individuals had to shift between external attention and memory retrieval in the memory-guided condition (i.e., to identify and retrieve the meaning of a stay/switch cue) but could remain primarily in an externally oriented mode in the explicitly instructed condition. For each condition and region of interest, the beta values for a given image period were first averaged across all voxels, and then separately averaged across art trials and room trials across all 4 runs of the condition. Finally, the beta values were averaged across art and room trials resulting in 1 average parameter estimate per participant per condition, equally weighted across art and room attentional states. To correspond with subsequent analyses, univariate activity was only averaged across valid trials in which the participant correctly answered the subsequent art/room probe, and, for the memory-guided condition, correctly identified the attentional state in the following trial. The same pattern of results was found when using all trials with a stay/switch cue (i.e., valid and invalid trials, regardless of accuracy on the art/room probe and, in the memory-guided task, regardless of the accuracy of attentional state selection on the following trial), as well as all valid trials with a stay/switch cue (regardless of accuracy). Univariate parameter estimates were compared across the explicitly instructed and memory-guided conditions using a within-participant t-test.

Background Connectivity

Task-evoked fluctuations in brain activity can lead to spurious findings of functional connectivity between brain areas: if two brain regions are similarly modulated by a task, they can appear to be functionally coupled even if they are acting in isolation. For this reason, we examined functional connectivity with the "background connectivity" approach, in which correlations between brain regions are estimated after regressing out task-evoked activity (Al-Aidroos et al., 2012). For this approach, we used a separate GLM than that used for univariate activity. Rather than modeling

individual trials, the GLM was designed to capture common task-evoked activity across trials (Al-Aidroos et al., 2012). This GLM therefore had 13 regressors that captured attentional states (art or room), trial validity (valid or invalid), participant accuracy (correct or incorrect), and trial component (orienting period, image period, probe): 1) the image periods of all valid trials with a cued (and probed) art attentional state for which participants correctly answered the art/room probe; 2) the image periods of all valid trials with a cued (and probed) room attentional state for which participants correctly answered the art/room probe; 3) the image periods of all invalid trials with a cued art attentional state (probed room state) for which participants correctly answered the art/room probe; 4) the image periods of all invalid trials with a cued room attentional state (probed art state) for which participants correctly answered the art/room probe; 5) the image periods of all valid trials with a cued (and probed) art attentional state for which participants did not correctly answer the art/room probe; 6) the image periods of all valid trials with a cued (and probed) room attentional state for which participants did not correctly answer the art/room probe; 7) the image periods of all invalid trials with a cued art attentional state (probed room state) for which participants did not correctly answer the art/room probe; 8) the image periods of all invalid trials with a cued room attentional state (probed art state) for which participants did not correctly answer the art/room probe; 9) the probe period for all valid trials for which the participant correctly answered the art/room probe; 10) the probe periods for all invalid trials for which the participant correctly answered the art/room probe; 11) the probe period for all valid trials for which the participant did not correctly answer the art/room probe; 12) the probe periods for all invalid trials for which the participant did not correctly answer the art/room probe; and 13) the orienting period of all trials. In parallel with the procedure described above for univariate activity, this GLM also included 6 motion regressors and employed FILM prewhitening. All runs were modeled separately.

The residuals of this model were registered to high-resolution 1mm MNI152 standard space. For each region of interest, the residual values were averaged across voxels and across the timepoints (5 TRs) for each image period (with onset time shifted 6s / 4 TRs from the start of the image period, to account for hemodynamic lag) resulting in a single value for each ROI for each image period. For consistency with univariate analyses, the calculation of background connectivity used valid trials only, although analyses were also repeated to include invalid trials (because evoked activity was regressed out prior to conducting background connectivity analyses, trial validity should have minimal effects on background connectivity).

Additionally, analyses were restricted to valid trials in which the participant correctly answered the art/room probe, and in the memory-guided condition, correctly identified the attentional state on the following trial. This was to make sure that background connectivity was estimated across trials in which participants were in good attentional states, both in terms of external attention (identifying art/room matches and detecting stay/switch cues) and internal attention (retrieving the meaning of stay/switch cues).

For each task condition (explicitly instructed and memory-guided) and for each attentional state (room and art) the residuals for each ROI (averaged across voxels and image period timepoints) were concatenated across all aforementioned trials across all runs of the condition of interest. Background connectivity was calculated by taking the Pearson correlation of the resulting timecourses of the 2 relevant ROIs. Finally, for each condition (explicitly instructed and memory-guided) background connectivity across correct, valid art trials and across correct, valid room trials was averaged together to balance the contribution of the 2 attentional states. This procedure resulted in a single measure of background connectivity for each participant in each condition. Background connectivity was compared across the 2 conditions (explicitly instructed and memory-guided) and regions (basal forebrain-hippocampus and DAN-hippocampus) with a 2-by-2 repeated measures ANOVA on the Fisher-transformed r-values. Follow-up comparisons were conducted with a within-participant t-test.

The same pattern of results was found when using all trials that included stay/switch cues (i.e., whether valid or invalid, whether performance was correct or incorrect), as well as when background connectivity was calculated without averaging the timepoints within each trial.

Temporal Multivariate-Univariate Dependence

Background connectivity provides a measure of how strongly the activity of 2 brain areas is correlated over time, but relating this measure to behavior on any given trial requires a timepoint-by-timepoint measure of connectivity strength. To accomplish this, Tompary et al. (2018) developed temporal multivariate-univariate dependence (TMUD) analysis to quantify the contribution of each time point to the overall correlation between 2 brain areas, giving a measure of how connectivity fluctuates over time. We applied the TMUD procedure to background connectivity for each task condition (explicitly instructed and memory-guided) as estimated from valid trials for both correct and incorrect judgments on the art/room probe, so that we could predict behavioral performance. First, for each region, the mean of the residual timecourse was calculated and subtracted from each individual time point in the residuals. Each mean-centered timepoint was then squared; these squared values were then summed and the square root obtained to find the "root sum-of-squares", i.e., the square root of the sum of squared deviations from the mean. Then, each mean-centered timepoint was divided by the root sum-of-squares, resulting in a normalized timecourse of "background activity" for each region of interest. Finally, we obtained the element-wise product of 2 regions' normalized timecourses, which provides a measure of their connectivity strength for each trial. Indeed, the sum of these element-wise products is equal to the background connectivity between the 2 regions, offering an intuitive reason for why the element-wise product for each trial is a measure of connectivity strength: the larger the value for a given trial, the larger the contribution of that trial to overall connectivity.

Mixed Effects Logistic Regression

The relationship between brain activity and behavior was modeled using mixed-effects logistic regression. Trialwise measures of connectivity (TMUD values calculated according to the procedure above) between the DAN and the hippocampus, and between the basal forebrain and the hippocampus, were used to predict participants' accuracy (0 or 1) on the art/room probe for each valid trial. First, the TMUD values were z-scored; outliers were then removed on an individual participant basis using Rosner's test for outliers as implemented in R. The model also included the attentional state of the given trial (art or room, effect coded) as well as a random intercept for each participant.

We first modeled the explicitly instructed and memory-guided conditions in the same model (effect coding the conditions), and incorporated an interaction between condition and TMUD, as follows: Accuracy ~ TMUD + Condition + TMUD*Condition + Attentional State + (1|Participant). Separate models were run for DAN-hippocampus and basal forebrain-hippocampus.

We also ran follow-up models separately for the two conditions, as follows: Accuracy ~ TMUD + Attentional State + (1|Participant).

Finally, an additional model for each condition was also run using univariate activity in the DAN in place of TMUD values.

Results

Validation of Background Connectivity Approach

Prior to examining our hypotheses regarding background connectivity, we first sought to ensure that we effectively removed task-evoked activity with our GLM. To that end, we examined the residuals from the background connectivity GLM (described in **Methods**), averaging across time-points separately for art and room trials. Our prior studies have found robust differences in univariate activity across art and room trials in the hippocampus and across the brain more generally (Aly & Turk-Browne, 2016a, 2016b). We verified that we effectively removed this task-evoked activity in our background connectivity GLMs, for the trials of interest used in subsequent analyses (i.e., valid trials with correct responses for the art/room probe and, in the memory-guided condition, correct identification of the attentional state on the following trial). Averaging the residuals for these trials confirmed no difference between art and room attentional states in the hippocampus (memory-guided: $t(28) = 0.45$, $p = 0.66$; explicitly-instructed: $t(28) = 0.41$, $p = 0.68$), basal forebrain (memory-guided: $t(28) = -0.72$, $p = 0.48$; explicitly-instructed: $t(28) = -1.63$, $p = 0.11$), DAN (memory-guided: $t(28) = -0.22$, $p = 0.82$; explicitly-instructed: $t(28) = -1.28$, $p = 0.21$), or a control region of interest, the ventral attention network (VAN; memory-guided: $t(28) =$

0.31, p = 0.76; explicitly-instructed: t(28) = -0.57, p = 0.57). There was also no difference in the averaged residuals between the memory-guided and explicitly instructed conditions in any region of interest (hippocampus: t(28) = -1.59, p = 0.12; DAN: t(28) = -1.27, p = 0.21; basal forebrain: t(28) = -1.75, p = 0.09; VAN: t(28) = -0.83, p = 0.41). Thus, these residuals — to be examined in our background connectivity analyses — did not contain reliable information about attentional states or task conditions.

Background Connectivity

Our main analyses centered on changes in hippocampal network interactions across the explicitly instructed and memory-guided attention conditions. We hypothesized that we would find stronger interactions between the basal forebrain and the hippocampus during explicitly instructed vs. memory-guided attention. We expected the opposite effect for DAN and hippocampus, with higher connectivity during memory-guided vs. explicitly instructed attention.

To test these hypotheses, we measured "background connectivity" by calculating functional connectivity between the regions of interest after removing task-evoked activity (Al-Aidroos et al., 2012). We observed the hypothesized interaction between condition (memory-guided or explicitly instructed) and region pair (basal forebrain-hippocampus or DAN-hippocampus), $F(1,28) = 15.68$, p = 0.0005, generalized $\eta^2 = 0.06$ (**Figure 3**).

Figure 3: Background connectivity in regions of interest. Background connectivity was calculated between the hippocampus and the dorsal attention network (DAN) and between the hippocampus and basal forebrain for each attention condition (explicitly instructed and memory-guided). Solid circles represent the mean background connectivity across participants in each condition, and error bars represent the standard error of the within-participant difference between conditions (separately for hippocampus-DAN and hippocampus-basal forebrain connectivity). Background connectivity between the DAN and the hippocampus was significantly higher during the memory-guided vs. explicitly instructed condition. Conversely, background connectivity between the basal forebrain and hippocampus was significantly higher in the explicitly instructed vs. memory-guided condition. The X symbol represents a significant interaction between attention condition and inter-regional interactions. (* p < .05, ** p < .01, *** p < .001).

Given this interaction, we conducted follow-up t-tests to compare connectivity patterns across conditions. Consistent with our hypothesis, background connectivity between the basal forebrain and the hippocampus was significantly stronger during explicitly instructed vs. memory guided attention, t(28) = 2.78, p = 0.009, 95% CI [0.044, 0.286], Cohen's dz = 0.52 (**Figure 3**, red). Conversely, background connectivity between the DAN and hippocampus was significantly higher during memory-guided vs. explicitly instructed attention, t(28) = 2.14, p = 0.04, 95% CI [0.005, 0.208], Cohen's dz = 0.40 (**Figure 3**, teal).

We next tested whether the switches in hippocampal connectivity, reported above, were relatively selective to our networks of interest. To that end, we examined hippocampal connectivity with the ventral attention network (VAN). The VAN is associated with bottom-up attention and shows increased activity for salient stimuli (Corbetta and Shulman, 2002; Vossel et al., 2014). Although the VAN is centrally involved in attentional orienting, we did not expect this network to differentially interact with the hippocampus across the explicitly instructed and memory-guided attention conditions.

Figure 4: Background connectivity between hippocampus and ventral attention network (VAN). Solid circles represent the mean background connectivity across participants in each condition (explicitly instructed, memory guided), and error bars represent the standard error of the within-participant difference between conditions. Background connectivity between the VAN and the hippocampus was not significantly different across conditions.

The logic for this hypothesis was as follows. To keep stimuli identical across tasks, the stay/switch cues were embedded in both memory-guided and explicitly instructed trials; however, in the latter, they were irrelevant to the attentional state on the following trial. Because participants were extensively trained, prior to the fMRI task, to identify stay/switch cues, the stay/switch cues may have captured bottom-up attention to some extent in both the memory-guided and explicitly

instructed conditions. If so, then the VAN — involved in bottom-up attention capture, as noted above — may not show differential connectivity with the hippocampus across tasks.

We first tested whether the VAN shows differential univariate activity across conditions. As we expected, we observed no difference, t(28) = -0.19, p = 0.85, 95% CI [-4.69, 3.88], Cohen's dz = -0.04. Furthermore, as we hypothesized, background connectivity between the VAN and the hippocampus was not significantly different across the explicitly instructed and memory-guided conditions, t(28) = -0.92, p = 0.37, 95% CI [-.19, 0.07], Cohen's dz = -0.17 (**Figure 4**). Thus, switches in hippocampal connectivity across conditions did not extend more broadly to other attention networks in the brain.

Relationship between Background Connectivity and Behavior

We next sought to test whether the strength of background connectivity between the hippocampus and our regions of interest (DAN, basal forebrain) was related to performance on the attention task. To test this, we first used a TMUD analysis to generate trial-by-trial measures of background connectivity. We then used logistic regression to model the relationship between these trial-wise connectivity measures and participants' accuracy on the art/room probe. We also included attentional state (art or room) as a control regressor in the model, to account for differences in accuracy across these attention tasks. Thus, we predicted trial-wise accuracy based on background connectivity (with separate models for DAN-hippocampus and basal forebrain-hippocampus), condition (memory-guided, explicitly instructed), attentional state (art, room), and the interaction between background connectivity and condition.

We first examined the relationship between DAN-hippocampus background connectivity and behavior. We found a significant interaction between trial-wise background connectivity strength and attention condition (memory-guided or explicitly instructed), β = 0.32, SE = 0.14, p = 0.02 (**Figure 5A**). To probe this interaction, we conducted separate logistic regression models for the two conditions. There was a significant positive relationship between DAN-hippocampus background connectivity and art/room probe accuracy in the memory-guided condition, β = 0.26, SE = 0.10, p = 0.007 (**Figure 5B**). There was no relationship between DAN-hippocampus background connectivity and probe accuracy in the explicitly instructed condition, β = -0.06, SE = 0.10, p = 0.56 (**Figure 5C**).

Basal-forebrain hippocampal background connectivity strength did not predict accuracy on the art/room probe (β = -0.006, SE = 0.08, p = 0.94) nor was there an interaction between connectivity strength and condition (β = -0.08, SE = 0.11, p = 0.46). Thus, basal-forebrain hippocampus connectivity generally did not predict behavior, nor was there evidence for differential effects across the memory-guided (β = -0.09, SE = 0.07, p = 0.20) and explicitly instructed conditions (β = -0.01, SE = 0.08, p = 0.89) (**Figure 5D-F**).

Figure 5: Relationship between DAN-hippocampus (top) and basal forebrain-hippocampus (bottom) background connectivity and accuracy on the attention task. Top: (A) The parameter estimates of the full model are displayed as odds ratios; background connectivity = trial-wise z-scored DAN-hippocampus background connectivity; condition = memory-guided or explicitly instructed (effect coded); attentional state = art or room (effect coded). DAN-hippocampus background connectivity predicted accuracy on the attention task in the memory-guided condition **(B)** but not the explicitly instructed condition **(C)**. **Bottom:** The parameter estimates of the full model are depicted as odds ratios **(D)** with the same regressors as in **(A)** except with basal-forebrain hippocampus background connectivity. Basal forebrain-hippocampus background connectivity did not predict attention task performance in either the memory-guided **(E)** or explicitly instructed **(F)** conditions. Colored lines represent the fitted model for each participant and colored dots represent trial-wise connectivity for each participant separately for incorrect (0) and correct (1) trials. (* $p < .05$, ** $p < .01$, *** $p < .001$).

Analysis of DAN Univariate Activity

We found that DAN-hippocampal connectivity predicted behavioral performance on the art/room probe in the memory-guided condition. Are these results selective to DAN-hippocampal connectivity or does DAN activity on its own predict performance? It is possible, for example, that the DAN is more engaged in the memory-guided vs. explicitly instructed condition because the former requires more top-down control and multitasking. To test this, we first probed whether

task-evoked univariate DAN activity was higher during memory-guided attention than explicitly instructed attention; however, there was no difference between conditions, t(28) = 1.24, p = 0.23, 95% CI [-2.47, 10.00], Cohen's dz = 0.23 **(Figure 6A)**. Thus, the memory-guided condition may not have required more top-down attentional effort than the explicitly instructed condition, perhaps because participants were extensively trained on the task; indeed, this is consistent with the lack of a behavioral difference between conditions (as reported in Günseli & Aly, 2020).

Figure 6: Univariate activity in the DAN and its relationship to behavior. (A) Mean DAN activity was not significantly different across the explicitly instructed and memory-guided conditions. Circles represent the average parameter estimates across participants and error bars represent the standard error of the within-participant difference between conditions. **(B)** DAN activity did not predict accuracy on the attentional task in either condition. Colored lines represent the fitted model for each participant and colored dots represent trial-wise DAN activity for each participant separately for incorrect (0) and correct (1) trials.

Although there was no difference in DAN activity across conditions, it is possible that DAN activity may predict behavior in the memory-guided but not explicitly instructed task. DAN activity did not, however, predict art/room probe accuracy in either condition (memory-guided: β = -0.03 , SE = 0.06, p = 0.62, explicitly instructed: β = -0.03 , SE = 0.06, p = 0.62; **Figure 6B**). Thus, DAN-hippocampus background connectivity, but not DAN univariate activity, predicts behavioral performance in the memory-guided attention condition.

Together, these results show that the hippocampus switches connectivity between the basal forebrain and DAN in accordance with external vs. internal attentional goals, respectively. Furthermore, hippocampal coupling with the DAN is associated with better attentional performance in the memory-guided vs. explicitly instructed condition. These results were specific to our networks and measures of interest, and did not extend to hippocampal coupling with the VAN or to univariate DAN activity.

Discussion

Hippocampus switches connectivity during external vs. internal attention

Everyday experience requires individuals to constantly switch between attending to the external environment and their internal memories and thoughts. How is this attentional switching coordinated? We focused on hippocampal connectivity profiles with attention networks in the brain, given evidence that the hippocampus is involved in both internal and external attention (Eichenbaum, 2004; Aly & Turk-Browne, 2016a, 2016b; Honey et al., 2017; Córdova et al., 2019). We compared connectivity patterns during an explicitly instructed attention task, which primarily required external attention, and a memory-guided task, which required switching between internal and external attention. Consistent with our hypotheses, we found that the hippocampus showed stronger background connectivity with the basal forebrain during explicitly instructed vs. memory-guided attention. In contrast, the hippocampus exhibited stronger background connectivity with the dorsal attention network (DAN) during memory-guided vs. explicitly instructed attention. Importantly, these changes in hippocampal connectivity did not extend to the ventral attention network (VAN), a region we hypothesized would be involved in both tasks due to its role in bottom-up attention (Corbetta and Shulman, 2002; Vossel et al., 2014). Finally, hippocampal coupling with the DAN predicted attentional behavior on the memory-guided but not explicitly instructed task.

We propose that DAN-hippocampus background connectivity may allow for the efficient retrieval of associative memories in the memory-guided attention task, thus freeing up attentional resources to process external visual signals and perform accurately on art/room match detection. DAN-hippocampus connectivity may not have been as important in predicting task performance in the explicitly instructed condition because that task did not require balancing internal and external attention. Unexpectedly, however, there was no relationship between hippocampal-basal forebrain coupling and attentional behavior. This was surprising because the basal forebrain should bias the hippocampus toward an externally oriented state (Newman et al., 2012; Tarder-Stoll, Jayakumar, et al., 2020; Decker & Duncan, 2020), which should in turn improve externally oriented attentional behavior. We return to this null effect below when we discuss limitations.

Together, these results suggest that switches between internal and external attention may be coordinated at least in part by changes in hippocampal communication with different attention networks in the brain.

Relation to existing work

The relationship between the basal forebrain and the hippocampus has been a rich area of research within the computational modeling and rodent literatures (Hasselmo, 1995; Parent & Baxter, 2004; Newman et al., 2012; Honey et al., 2017). Within the hippocampus, acetylcholine from the basal forebrain strengthens afferent (external) synapses and suppresses reciprocal (internal) synapses (Hasselmo 2006; Tarder-Stoll, Jayakumar, et al., 2020; Decker & Duncan, 2020). Together, this modulation of external and internal connections can drive an externally oriented state in the hippocampus. Our findings critically expand on this work by providing empirical evidence in humans that functional connectivity between the basal forebrain and the hippocampus is strengthened when there are greater demands for external attention. Our results therefore suggest that top-down goals to attend to the external world may modulate coupling between the cholinergic basal forebrain and hippocampus, driving the hippocampus toward an externally oriented state.

Recent work by Li et al. (2022) explored external and internal states across the brain using background connectivity, the same approach used in the current study. Li et al. found that sustained background connectivity within the default mode network characterized an internally oriented state, whereas stable patterns of connectivity within the DAN/cognitive control network defined an externally oriented state. Moreover, the retrosplenial cortex flexibly switched between these two networks during internal vs. external attention. This work, along with other research (Spreng et al., 2010; Kim, 2015), emphasizes the role of the DAN as an externally oriented, task-positive network; however, Li et al. argue that this externally oriented state does not preclude situations in which the DAN is recruited for internal attention as well. Furthermore, they propose that the hippocampus (which did not emerge in their whole-brain analysis) may shift between externally and internally oriented modes (Li et al., 2022).

Indeed, in our study, we found that the hippocampus does switch its connectivity patterns in tasks that vary external and internal attention requirements; however, contrary to Li et al., we found that the DAN is more strongly coupled with the hippocampus during *internal* vs. external attention. Because the DAN is involved with both external attention (e.g., Li et al.) and internal attention (e.g., Stokes et al., 2012), it is possible that the recruitment of the DAN to help shift the hippocampus towards internal processing is dependent on the particular cognitive load and the demands of the given task. Future research can systematically compare DAN-hippocampus connectivity in various attention tasks to determine when and why the DAN is more important for internally vs. externally oriented attention.

Limitations

We found that DAN-hippocampus connectivity during the memory-guided condition (but not the explicitly instructed condition) predicted performance on art/room match detection. This null result for the explicitly instructed condition is intriguing because one might have instead predicted a *negative* relationship. Because internal attention demands were minimal for the

explicitly instructed condition, moments of increased DAN-hippocampus connectivity in this condition might instead reflect internally oriented distraction (e.g., mind wandering) which should then hamper performance. Future research may uncover such a negative relationship between DAN-hippocampus connectivity and performance on external attention tasks, perhaps in settings in which mind wandering might be particularly common.

In contrast, because internal attention demands were higher for the memory-guided condition, we expected DAN-hippocampal connectivity to benefit attentional performance — and indeed observed such an effect. This relationship may reflect the importance of DAN-hippocampus connectivity in retrieving memories of the stay/switch cues in the memory-guided condition, which in turn frees up cognitive resources for performing the primary art/room match detection task. If this is true, one might also predict that stronger connectivity between the DAN and hippocampus might be associated with better performance in selecting the correct attentional state in the memory-guided condition (based on the stay/switch cues). Our study was not ideally set up to test this particular hypothesis, however, for two reasons. First, accuracy in selecting attentional states in the memory-guided condition approached ceiling performance (94.9% for stay cues and 96.7% for switch cues, as reported in Günseli & Aly, 2020) This high performance was a result of overtraining, which was necessary to ensure matched art/room task performance across the memory-guided and explicitly instructed conditions. Second, because the attentional state selection occurred at the start of the subsequent trial, after an approximately 6.66s inter-trial interval (ITI), RTs may not be particularly sensitive measures — participants had extensive time to prepare a response during the ITI. For these reasons, neither accuracy nor RT were ideal measures for assessing the relationship between DAN-hippocampus connectivity and the ability to retrieve memories of the stay/switch cues.

Our study is also limited in that we found a surprising null effect. Although the basal forebrain showed increased connectivity with the hippocampus during explicitly instructed vs. memory-guided attention, as hypothesized, connectivity strength did not predict behavior in either condition. This was unexpected because, to the extent that such connectivity enhances the processing of external signals, stronger connectivity should predict better identification of art/room matches, perhaps in both the memory-guided and explicitly instructed conditions. One possible reason for this null effect is the difficulty of the art and room attention tasks; high hippocampus-basal forebrain connectivity may have put individuals in a strong externally oriented state, but they may have nevertheless failed to identify art/room matches because these tasks were relatively difficult. Future research can explore the impact of basal forebrain-hippocampus connectivity on behavioral measures of external attention in tasks in which poor performance is most likely to result from mind wandering rather than task difficulty.

Finally, this study was designed to compare memory-guided and explicitly instructed conditions rather than interactions between these conditions and attentional states (art or room). Future experiments can explore whether hippocampal coupling with the basal forebrain and DAN varies

both by the nature of attentional cueing (from memory or external signals) and by the content of attentional states.

Conclusion

We explored how the hippocampus may shift between external attention to the outside world and internal attention to retrieved memories. We found that the hippocampus shows stronger functional connectivity with the cholinergic basal forebrain during attention tasks that are guided by external cues rather than retrieved memories. Conversely, the hippocampus exhibited stronger connectivity with the dorsal attention network during memory-guided vs. externally cued attention, and this connectivity predicted trial-by-trial attentional behavior. Taken together, these results suggest that the basal forebrain and dorsal attention network may help the brain balance attention to the external and internal world by differentially communicating with the hippocampus — allowing us to efficiently guide behavior based on both sensory signals and memories.

References

Al-Aidroos, N., Said, C. P., & Turk-Browne, N. B. (2012). Top-down attention switches coupling between low-level and high-level areas of human visual cortex. *Proceedings of the National Academy of Sciences*, *109*(36), 14675-14680.

Aly, M., & Turk-Browne, N. B. (2016). Attention stabilizes representations in the human hippocampus. *Cerebral Cortex*, *26*(2), 783-796.

Aly, M., & Turk-Browne, N. B. (2016). Attention promotes episodic encoding by stabilizing hippocampal representations. *Proceedings of the National Academy of Sciences*, *113*(4), E420-E429.

Berryhill, M. E., Phuong, L., Picasso, L., Cabeza, R., & Olson, I. R. (2007). Parietal lobe and episodic memory: bilateral damage causes impaired free recall of autobiographical memory. Journal of Neuroscience, 27(52), 14415-14423.

Cabeza, R. (2008). Role of parietal regions in episodic memory retrieval: the dual attentional processes hypothesis. Neuropsychologia, 46(7), 1813-1827.

Cabeza, R., Ciaramelli, E., Olson, I. R., & Moscovitch, M. (2008). The parietal cortex and episodic memory: an attentional account. Nature reviews neuroscience, 9(8), 613-625.

Chun, M. M., Golomb, J. D., & Turk-Browne, N. B. (2011). A taxonomy of external and internal attention. Annual review of psychology, 62(1), 73-101.

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in the brain. *Nature reviews neuroscience*, *3*(3), 201-215.

Córdova, N. I., Turk-Browne, N. B., & Aly, M. (2019). Focusing on what matters: Modulation of the human hippocampus by relational attention. Hippocampus, 29(11), 1025-1037.

Decker, A. L., & Duncan, K. (2020). Acetylcholine and the complex interdependence of memory and attention. Current Opinion in Behavioral Sciences, 32, 21-28.

Eichenbaum, H. (2004). Hippocampus: cognitive processes and neural representations that underlie declarative memory. Neuron, 44(1), 109-120.

Günseli, E., & Aly, M. (2020). Preparation for upcoming attentional states in the hippocampus and medial prefrontal cortex. *Elife*, *9*, e53191.

Hasselmo, M. E. (1995). Neuromodulation and cortical function: modeling the physiological basis of behavior. Behavioural brain research, 67(1), 1-27.

Hasselmo, M. E. (2006). The role of acetylcholine in learning and memory. Current Opinion in Neurobiology, 16(6), 710–715.

Hasselmo, M. E., Bodelón, C., & Wyble, B. P. (2002). A proposed function for hippocampal theta rhythm: separate phases of encoding and retrieval enhance reversal of prior learning. Neural computation, 14(4), 793-817.

Honey, C. J., Newman, E. L., & Schapiro, A. C. (2017). Switching between internal and external modes: a multiscale learning principle. Network Neuroscience, 1(4), 339-356.

Hutchinson, J. B., & Turk-Browne, N. B. (2012). Memory-guided attention: Control from multiple memory systems. Trends in cognitive sciences, 16(12), 576-579.

Hutchinson, J. B., Uncapher, M. R., & Wagner, A. D. (2015). Increased functional connectivity between dorsal posterior parietal and ventral occipitotemporal cortex during uncertain memory decisions. Neurobiology of learning and memory, 117, 71-83.

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage*, *17*(2), 825-841.

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., & Smith, S. M. (2012). Fsl. *Neuroimage*, *62*(2), 782-790.

Jimenez, A. M., Lee, J., Wynn, J. K., Cohen, M. S., Engel, S. A., Glahn, D. C., ... & Green, M. F. (2016). Abnormal ventral and dorsal attention network activity during single and dual target detection in schizophrenia. *Frontiers in psychology*, *7*, 323.

Kim, H. (2015). Encoding and retrieval along the long axis of the hippocampus and their relationships with dorsal attention and default mode networks: the HERNET model. Hippocampus, 25(4), 500-510.

Lepsien, J., & Nobre, A. C. (2006). Cognitive control of attention in the human brain: Insights from orienting attention to mental representations. Brain research, 1105(1), 20-31.

Li, Y. P., Wang, Y., Turk-Browne, N. B., Kuhl, B. A., & Hutchinson, J. B. (2022). Perception and Memory Retrieval States are Reflected in Distributed Patterns of Background Functional Connectivity. bioRxiv.

Newman, E. L., Gupta, K., Climer, J. R., Monaghan, C. K., & Hasselmo, M. E. (2012). Cholinergic modulation of cognitive processing: insights drawn from computational models. Frontiers in behavioral neuroscience, 6, 24.

Parent, M. B., & Baxter, M. G. (2004). Septohippocampal acetylcholine: involved in but not necessary for learning and memory?. Learning & Memory, 11(1), 9-20.

Posner, M. I. (1980). Orienting of attention. Quarterly journal of experimental psychology, 32(1), 3-25.

Ruiz, N. A., Meager, M. R., Agarwal, S., & Aly, M. (2020). The medial temporal lobe is critical for spatial relational perception. *Journal of Cognitive Neuroscience*, *32*(9), 1780-1795.

Ruiz, N. A., Thieu, M. K., & Aly, M. (2021). Cholinergic modulation of hippocampally mediated attention and perception. Behavioral Neuroscience, 135(1), 51.

S.M. Smith. Fast robust automated brain extraction. Human Brain Mapping, 17(3):143-155, November 2002.

Spreng, R. N., Stevens, W. D., Chamberlain, J. P., Gilmore, A. W., & Schacter, D. L. (2010). Default network activity, coupled with the frontoparietal control network, supports goal-directed cognition. Neuroimage, 53(1), 303-317.

Stokes, M. G., Atherton, K., Patai, E. Z., & Nobre, A. C. (2012). Long-term memory prepares neural activity for perception. Proceedings of the National Academy of Sciences, 109(6), E360-E367.

Tarder-Stoll, H., Jayakumar, M., Dimsdale-Zucker, H. R., Günseli, E., & Aly, M. (2020). Dynamic internal states shape memory retrieval. Neuropsychologia, 138, 107328.

Tompary, A., Al-Aidroos, N., & Turk-Browne, N. B. (2018). Attending to what and where: Background connectivity integrates categorical and spatial attention. *Journal of Cognitive Neuroscience*, *30*(9), 1281-1297.

Verschooren, S., Liefooghe, B., Brass, M., & Pourtois, G. (2019). Attentional flexibility is imbalanced: Asymmetric cost for switches between external and internal attention. Journal of experimental psychology: human perception and performance, 45(10), 1399.

Verschooren, S., Schindler, S., De Raedt, R., & Pourtois, G. (2019). Switching attention from internal to external information processing: A review of the literature and empirical support of the resource sharing account. Psychonomic bulletin & review, 26(2), 468-490.

Villano, I., Messina, A., Valenzano, A., Moscatelli, F., Esposito, T., Monda, V., ... & Messina, G. (2017). Basal forebrain cholinergic system and orexin neurons: effects on attention. Frontiers in behavioral neuroscience, 11, 10.

Vossel, S., Geng, J. J., & Fink, G. R. (2014). Dorsal and ventral attention systems: distinct neural circuits but collaborative roles. The Neuroscientist, 20(2), 150-159.

Wagner, A. D., Shannon, B. J., Kahn, I., & Buckner, R. L. (2005). Parietal lobe contributions to episodic memory retrieval. Trends in cognitive sciences, 9(9), 445-453.

Záborszky, L., Gombkoto, P., Varsanyi, P., Gielow, M. R., Poe, G., Role, L. W., ... & Chiba, A. A. (2018). Specific basal forebrain–cortical cholinergic circuits coordinate cognitive operations. Journal of Neuroscience, 38(44), 9446-9458.

Zaborszky, L., Hoemke, L., Mohlberg, H., Schleicher, A., Amunts, K., & Zilles, K. (2021). *Probabilistic cytoarchitectonic map of Ch 123 (Basal Forebrain) (v4.3)* [Data set]. EBRAINS. DOI: 10.25493/19M1-B4B