Spatially distinct phytohormone responses of individual *Arabidopsis thaliana* root cells to infection and colonization by *Fusarium oxysporum*

Jacob Calabria[1,2,a], Liu Wang[1,3,b], Madlen I. Rast-Somssich[2,c], Hsiang-Wen Chen[1,3,d], Michelle Watt[2,e], Staffan Persson[3,4,f], Alexander Idnurm[5,g], and Marc Somssich[1,h,*]

[1] Plant-Fusarium Interactions Research Team, School of BioSciences, University of Melbourne, Parkville, Australia

[2] Crop Root Physiology Lab, School of BioSciences, University of Melbourne, Parkville, Australia

[3] Plant Cell Biology Lab, School of BioSciences, University of Melbourne, Parkville, Australia

[4] Current address: Copenhagen Plant Science Center, University of Copenhagen, Frederiksberg C, Denmark

[5] Mycology Laboratory, School of BioSciences, University of Melbourne, Parkville, Australia

Twitter: [a] @jacobcalabria, [b] @Liu_Wang1, [c] @MSomssich, [d] @HsiangWen_Chen, [e] @MelbourneRoots, [f] @perssonlab1, [g] @AlexanderIdnurm, [h] @somssichm

[*] For correspondence: marc.somssich@unimelb.edu.au

Graphical Abstract

- Colonization of the *A. thaliana* root tip by *F. oxysporum* strain *Fo*5176 leads to immediate cell death of the colonized and surrounding tissue.

- As the colonization front progresses through the vasculature, the cell death front moves along with it, through not only the vasculature, but also the surrounding tissues.

- WRKY70 positively regulates salicylic acid (SA) biosynthesis in cells immediately adjacent to the colonized tissue, inducing a hypersensitive response (HR), thereby killing off the cells deemed lost to the intruder, establishing the cell death front.

- Slightly further removed from the HR zone, WRKY11 induces jasmonate (JA) biosynthesis in cells of the vasculature to launch a defense response aimed at actively repelling the fungus.

Abstract

Jasmonic acid (JA), ethylene (ET) and salicylic acid (SA) are the three major phytohormones coordinating a plant's defense response to pathogenic attack. While JA and ET are assumed to primarily control the defense against necrotrophic pathogens, SA-induced defense responses target mainly biotrophic microbes, and can include drastic measures such as programmed cell death as part of the plant's hypersensitive response (HR). *Fusarium oxysporum* is a hemibiotrophic fungal pathogen of several plant species, including many important food crops, and the model plant species *Arabidopsis thaliana*. Colonization of the plant's root vascular tissue by the fungus eventually results in wilting and plant death. A general role for JA, ET and SA in combating infection and colonization of the plant by *F. oxysporum* has been demonstrated, but their distinct roles and modes of action have so far not been described. Here, using high resolution microscopy with fluorescent marker lines of *A. thaliana* roots infected with *F. oxysporum* we show that SA acts spatially separate from JA/ET, in a distinct set of root cells immediately neighboring the fungal colonization site. JA and ET, however, act together in a different, but also clearly defined set of cells, slightly removed from the colonization site. SA induces HR in the cells directly in contact with already colonized tissue, that presumably can no longer be rescued from infection, stopping the spread of colonization. In comparison, JA and ET act in a group of cells slightly removed from the HR site, initiating a defense response to actively resist the invader. These results show how the three phytohormones act together, but spatially and functionally separate from each other, to fight this hemibiotrophic pathogen. Such a high-resolution analysis to resolve the plant's immune response to pathogenic infection on an individual cell level and in intact tissue has so far been lacking.

Keywords

Arabidopsis thaliana, *Fusarium oxysporum*, *Fo*5176, salicylic acid, jasmonic acid, ethylene, plant immunity, spatial immunity, plant-microbe interactions

Introduction

Phytohormones play an important role in coordinating a plant's defense response to attack by microbes (Hou and Tsuda, 2022). A plant senses and recognizes a potential attacker via their microbe-associated molecular patterns (MAMPs); highly conserved structures on the outside of the microbe, such as flagellin in the bacterial flagellum, or chitin in the fungal cell wall (Wang *et al.*, 2022*b*). Since the root is continuously surrounded by microbes carrying MAMPs in the soil, recognition of a MAMP alone will not trigger the plant immune system. Instead, it is the co-incidence of MAMPs with cell damage caused by a pathogenic microbe that will launch a full immune response (Zhou *et al.*, 2020). MAMPs and damage-associated molecular patterns (DAMPs) are sensed by the plant outside the cell via plasma membrane localized PATTERN RECOGNITION RECEPTORS (PRRs), which then transduce the information about the attacker into the cell via an intracellular kinase domain (Lee *et al.*, 2021). Inside the cell, several defense pathways are then activated, including phytohormone-dependent signaling pathways (Lee *et al.*, 2021). Jasmonic acid (JA), ethylene (ET) and salicylic acid (SA) are the three main phytohormones involved in coordinating the plant's defense against pathogenic invaders (Hou and Tsuda, 2022). Among them, JA and ET mostly act in concert, while JA and SA appear to act in mutual antagonism. It is generally assumed that JA/ET control the plant's defense against necrotrophic pathogens, while SA is the regulator of pathways combating biotrophic pathogens (Hou and Tsuda, 2022). SA-mediated immunity can involve drastic measures such as the hypersensitive response (HR), a form of programmed cell death aimed at containing an infection by killing off the tissue surrounding the infection site in an attempt to prevent further spread of the microbe to other healthy cells and tissues (Pitsili *et al.*, 2020). This form of defense is not activated against necrotrophic pathogens, as these feed on dead plant material and would benefit from such a cell death response (McCombe *et al.*, 2022). Thus, the JA/ET-mediated response to necrotrophs instead focusses on active mechanisms that can prevent spread of the pathogen, for example, releasing reactive oxygen species (ROS) into the apoplast that can harm and kill the microbe, producing antimicrobial compounds, or changing the apoplastic pH (Masachis *et al.*, 2016; McCombe *et al.*, 2022).

Fusarium oxysporum, a soil-inhabiting ascomycete fungus, is a destructive pathogen to many plant species, including several major crop plants, such as the tomato (*Solanum lycopersicum*), cotton (*Gossypium barbadense*), and banana (*Musa acuminata*) (Takken and Rep, 2010; Gordon, 2017;

Dita *et al.*, 2018). *F. oxysporum* reproduces asexually and the resulting clonal lineages all represent

individual formae speciales (ff. spp.), that together form the *F. oxysporum* species complex

(Summerell, 2019). The different ff. spp. mostly share an 11 chromosome core genome and it is

their lineage-specific accessory chromosomes that account for the host specificity of each f. sp.,

and whether a f. sp. is pathogenic, non-compatible, or beneficial to a certain plant (Ma *et al.*, 2010;

Redkar *et al.*, 2022). *F. oxysporum* hyphae grow in the soil and are chemotactically guided toward

the root of their host plant (Wang *et al.*, 2022*a*). Upon reaching the plant, they attach and grow

along the root until they enter via natural openings, such as lateral root emergence sites or wounds

(Wang *et al.*, 2022*a*). Inside the root, the hyphae grow into the vasculature, infecting and

colonizing xylem vessel cells (van der Does *et al.*, 2008). Once inside the vasculature, the fungus

drains the plant of water and nutrients, and can spread to all parts of the plant, which begins to wilt

and eventually dies (Gordon, 2017). Based on these observations, *F. oxysporum* is considered a

hemibiotrophic fungus, that initially lives biotrophically within the host's vasculature before

eventually turning necrotrophic and killing the plant (Gordon, 2017). Research into the plant's

defense against *F. oxysporum* has originally focused on the crop plants threatened by Fusarium

wilt (e.g. Panama disease of bananas) and the *F. oxysporum* ff. spp. targeting them. However, more

recently, the *Arabidopsis thaliana – F. oxysporum* f. sp. *conglutinans* strain 5176 (*Fo*5176)

pathosystem has also been established to study this plant-fungus interaction with all the modern

molecular, cell biological, genetic and genomic tools available for this plant model system (Wang

et al., 2022*a*).

Early work on the *A. thaliana – Fo*5176 pathosystem has focused on creating several large-scale

transcriptomic datasets, as well as comprehensive infection studies of *A. thaliana* mutants, to

investigate the role of JA and SA in combating infection by *Fo*5176 (Wang *et al.*, 2022*a*). These

studies have established a role for the different hormones in the plant's defense against *Fo*5176,

but were also often inconclusive as to their exact roles (Wang *et al.*, 2022*a*). For example, draining

a plant of SA enhances disease severity, while treatment of *A. thaliana* leaves with SA led to

systemic acquired resistance (SAR) against *Fo*5176 in the treated leaves (Berrocal-Lobo and

Molina, 2004; Diener and Ausubel, 2005; Edgar *et al.*, 2006). However, no other than the treated

tissue showed SAR, and marker genes for activated SA-signaling were either unresponsive, or

even slightly downregulated in plants infected by *Fo*5176 (Edgar *et al.*, 2006; Kidd *et al.*, 2011;

Wang *et al.*, 2022*a*). Similarly, infection of the plant roots by *Fo*5176 resulted in upregulation of

JA-biosynthesis genes in shoot tissue, but not in root tissue, the actual site of infection, where
makers for JA-biosynthesis and signaling were either unresponsive, downregulated or only very
slightly induced (Kidd *et al.*, 2011; Chen *et al.*, 2014; Wang *et al.*, 2022*a*). Cole *et al.* (2014)
furthermore found that the absence of JA biosynthesis in the *allene oxide synthase* (*aos*) mutant
was inconsequential to both infection and disease progression, and the JA-receptor mutant
coronatine-insensitive 1 exhibited no disease symptoms upon infection (Thatcher *et al.*, 2009; Cole
et al., 2014). Thus, while it is assumed that both SA and JA are involved in the plant's defense
response against this hemibiotrophic fungus, the available transcriptomic and mutant data are often
inconclusive in this respect and therefore insufficient to substantiate this claim (Wang *et al.*,
2022*a*).

A possible explanation for the failure of such large-scale transcriptomic approaches to identify
novel and specific pathways involved in the defense against *Fo*5176, as well as the apparent
inconclusiveness in relation to known pathways, such as the phytohormones, may lie in the tissue
and timing used in the experiments. In contrast to other plants, such as tomato, where *F. oxysporum*
can infect the plant via several openings simultaneously, the fungus generally appears to infect *A.*
thaliana solely via the root tip, thereby limiting the number of infection sites and thus infected
tissue available to be subjected to transcriptomic analyses (Czymmek *et al.*, 2007; Wang *et al.*,
2022*a*; Martínez-Soto *et al.*, 2022). Thus, by using either whole seedlings or the whole root system
to extract RNA, the transcriptional changes in these few infected cells may be diluted too much by
the other uninfected cells. This would also explain why RNA sequencing did eventually detect a
slight activation of JA-responsive genes at six days post infection, when colonization has most
likely progressed up the main root, and in several lateral roots (Lyons *et al.*, 2015). Furthermore,
as the plant also senses the fungal hyphae in its rhizosphere via their MAMPs, a majority of root
cells will launch a general MAMP-response, which further dilutes the response of the cells
undergoing infection.

For our work, we have set up a microscopy-based live-imaging approach monitoring
transcriptional activity of different pathways *in planta*, over time and on an individual cell level.
We have previously published our pGG-PIP GreenGate entry vector collection with regulatory
sequences for plant immunity genes, including reporters for the different phytohormone pathways
(Calabria *et al.*, 2022). Using this approach, we detected local responses of genes involved in SA

and JA biosynthesis, signaling and regulation, targeted to specific tissues and to a very limited

number of root cells upon infection with *Fo*5176. Interestingly, we found that SA acts spatially

separate from JA/ET, in a distinct set of root cells while JA and ET act together in a different, but

also clearly defined set of cells.

Results

***Fo*5176 colonization of the root leads to cell death of the infected tissue**

To monitor the infection and colonization of *A. thaliana* roots by *Fo*5176, we cultivated 10-day

old seedlings in square petri dishes, added fungal spores and then imaged hyphal growth, infection

and colonization over the next 14 days. We used a transgenic *Fo*5176 line expressing a cytoplasmic

tdTomato fluorophore to visualize the fungus (Calabria *et al.*, 2022). One of the first things

noticeable when imaging infected plants was that the tissue surrounding the colonization site

appeared to undergo immediate cell death. When cells die, the plasma membrane and vacuole

collapse, leaving behind the cell wall material. As this material scatters light, the dead tissue

appears as bright white in the image (Fig. 1A). In fact, when the vasculature of a root tip was

colonized by *Fo*5176, the entire tip underwent cell death (Fig. 1A, B). By imaging a plant line

expressing a fluorescent marker for the plant plasma membrane (PM) (mNeonGreen(mNG)-

LTI6b), this became even more apparent, as the fluorescence signal from the PM outlined all cells

of the plant but disappeared completely from all cells around the colonized tissue (Fig. 1C, D). As

fungal colonization progressed through the vasculature as a 'colonization front', a 'cell death front'

was also observed that moved in line with it. In contrast though, the cell death front spanned the

entire circumference of the root, including the outer tissues, epidermis, cortex, and endodermis,

which are not colonized by the fungus. Because cell death was not limited to the colonized tissue,

we reasoned that cell death may be the result of a hypersensitive response (HR) initiated by the

plant, rather than caused directly by *Fo*5176.

Salicylic acid signaling is activated in the hypersensitive response zone

Since HR is often activated via an SA- and PATHOGENESIS-RELATED PROTEIN 1 (PR1)-

dependent pathway, we next investigated if SA biosynthesis and signaling via the antimicrobial

protein PR1 were indeed activated in the cells bordering the colonized tissue. We utilized reporters

for the key SA biosynthesis enzyme ENHANCED DISEASE SUSCEPTIBILITY 16 (EDS16; also

Figure 1: Colonization of *A. thaliana* roots by *Fo*5176 causes cell death
A, B: Bright field (A) and fluorescence image (B) of a colonized root tip. (A) Dead cells appear bright white. (B) Fluorescent fungal hyphae (magenta) can be seen in the tip and the vasculature. The colonization front (magenta bars to the left and right of the root) aligns with the cell death front. **C, D**: Overlay of bright field and fluorescence image (C) and fluorescence image only (D) of a root expressing the PM-marker mNG-LTI6b (yellow) and colonized by *Fo*5176 (magenta). Fluorescent hyphae (magenta) can be seen in the tip and the colonized vasculature. The fluorescence coming from the PM-marker (yellow) is lost at the colonization front (magenta bars).

named SALICYLIC ACID DEFICIENT 2 (SID2)) and PR1. In an uninfected control, *EDS16* was

expressed in the meristematic and elongation zones (MZ/EZ) of the root tip, as well as in older

differentiated cells, with no expression detectable in young differentiating cells (Fig. S1A, B). In

the tip, expression also appeared to be stronger in the outer layers, epidermis and cortex, compared

to the vasculature (Fig. S1A, B). *PR1* expression was only weakly detectable in the mature

differentiation zone (DZ; Fig. S1C, D). However, following infection and colonization by *Fo*5176,

EDS16 expression was upregulated in the root tip, and spatially targeted to a group of vascular

cells immediately at the colonization front of the fungus (Fig. 2A, B). *PR1* expression, which was

Figure 2: Expression of *EDS16* and *PR1* following colonization of the root by *Fo*5176
A, B: Overlay of bright field and fluorescence (A), and fluorescence-only (B) images of a root tip expressing the *EDS16* reporter (yellow) in response to colonization by *Fo*5176 (magenta). **C, D**: Overlay of bright field and fluorescence (C), and fluorescence-only (D) images of a root tip expressing the *PR1* reporter (yellow) in response to colonization by *Fo*5176 (magenta). The colonization front is marked by magenta bars.

not detectable in the tip of the uninfected control, was also upregulated in these same cells, the set of cells likely to subsequently initiate HR (Fig. 2C, D).

Jasmonic acid and ethylene act in a distinct group of cells close to the colonized tissue

We then investigated the responses of our JA and ET reporters. In the case of JA, we used regulatory sequences for *ALLENE OXIDE SYNTHASE* (*AOS*), a key enzyme in the JA biosynthesis pathway, *PLANT DEFENSIN 1.2* (*PDF1.2*), a defense gene activated as the result of active JA-signaling, and *ETHYLENE RESPONSE FACTOR 1* (*ERF1*), an activator of JA- and ET-responsive defense gene expression, as well as an integrator of the JA- and ET-signaling pathways (Wang *et al.*, 2022*a*). To monitor ET-biosynthesis, we used our reporter for 1-AMINOCYCLOPROPANE-

1-CARBOXYLIC ACID SYNTHASE 2 (ACS2), a key enzyme in the ET biosynthesis pathway.

We found that *AOS* expression was mostly absent in the uninfected control roots, with very weak

expression limited to sites of natural stress occurrence, such as lateral root emergence sites (Fig.

S2A, B). Similarly, *PDF1.2* was not expressed in the root; however, it was strongly expressed in

above ground tissue. *ERF1* was expressed in all differentiated tissues, but was absent from the root

tip, including the MZ and EZ (Fig. S2C-F). This contrasting expression pattern between *AOS* and

ERF1 most likely represents activation via the ET-, rather than the JA-pathway, since ET also acts

in several developmental pathways, while JA-signaling is mostly restricted to stress conditions.

Indeed, the marker for ET-biosynthesis, *ACS2*, was expressed throughout the root system in the

uninfected control, with the main root showing stronger expression than the lateral roots (Fig. S2G-

J). Furthermore, expression was absent from the epidermis in differentiated tissues, and the

expression was generally very weak between the EZ and early DZ. Expression in the MZ was

moderate and included the epidermal cells (Fig. S2G-J).

Following infection and colonization of the plant by *Fo*5176, *AOS* expression was robustly

induced in a distinct group of cells roughly four to seven cells away from the colonized tissue (Fig.

3A, B). The expression then slowly tapered out along the root toward the base. Interestingly,

expression was limited to cells of the vasculature, with no upregulation in the epidermis, cortex,

or endodermis. Thus, *AOS* expression and JA signaling are precisely targeted to the tissue that is

colonized by the fungus. Interestingly, *PDF1.2* was not responsive in the group of cells next to the

colonization site, despite this gene being routinely used as a general readout for JA-signaling

activity in PCR or transcriptomic assays (Fig. 3C, D). However, at later stages of infection, when

the fungus had progressed far enough along the vasculature to reach the DZ, either through the

main root or along a lateral root, *PDF1.2* was upregulated in the vasculature of these mature,

differentiated cells now threatened by the pathogen (Fig. S3A, B). Therefore, PDF1.2 possibly

only acts in mature, differentiated cells, while JA probably acts via a different pathway, such as

VEGETATIVE STORAGE PROTEIN 2 (VSP2), in the undifferentiated tissue of the root tip. We

next imaged *ERF1* and, like *AOS*, found it to be robustly upregulated in the same set of vascular

cells that showed the *AOS* expression maximum (Fig. 4E, F). However, in contrast to *AOS*, the

ERF1 expression pattern returned to normal immediately behind the distinct group of cells that

showed upregulation of *AOS*, i.e., it was moderately expressed in all cell types, not just the

vasculature (Fig. 4E, F). The marker for ET biosynthesis, *ACS2*, was also upregulated in a distinct

Figure 3: Expression of *AOS* and *PDF1.2* following colonization of the root by *Fo*5176
A, B: Overlay of bright field and fluorescence (A), and fluorescence-only (B) images of a root tip expressing the *AOS* reporter (yellow) in response to colonization by *Fo*5176 (magenta). **C, D**: Overlay of bright field and fluorescence (C), and fluorescence-only (D) images of a root tip expressing the *PDF1.2* reporter (yellow) in response to colonization by *Fo*5176 (magenta). The colonization front is marked by magenta bars.

group of cells next to the colonized tissue, with its maximum partially overlapping that of *AOS*

and *ERF1* (Fig. 4A-D). However, based on fluorescence intensity, *ACS2* expression appeared to

be induced much stronger than *AOS* or *ERF1* (Fig. 4A-D). Further, *ACS2* expression started closer

to the colonization front, as if ET biosynthesis was upregulated at the end of the HR-zone already,

and not just in the defense response zone as JA is. Following the expression maximum in the group

of cells that form the defense response zone, *ACS2* expression immediately dropped back to basal

levels, thereby resembling the pattern of *ERF1* not *AOS*. Expression remained restricted to the

central cylinder for a long stretch, however, and only expanded to all cells and tissues again in the

older parts of the root, close to the root's base and the root-to-shoot transition zone. Thus, while

SA and JA appear to be strictly separated spatially, ET may bridge the two zones.

Figure 4: Expression of *ACS2* and *ERF1* following colonization of the root by *Fo*5176
A, B: Overlay of bright field and fluorescence (A, C), and fluorescence-only (B, D) images of a root tip expressing the *ACS2* reporter (yellow) in response to colonization by *Fo*5176 (magenta). The images in C and D are the same as A and B with reduced exposure time to allow for the identification of individual cells. **E, F**: Overlay of bright field and fluorescence (E), and fluorescence-only (F) images of a root tip expressing the *ERF1* reporter (yellow) in response to colonization by *Fo*5176 (magenta). The colonization front is marked by magenta bars.

A WRKY11/WRKY70 module could coordinate the spatially distinct JA and SA responses

WRKY11 and WRKY70 act upstream of JA and SA signaling respectively, and act as integrators

of these two mutually antagonistic pathways (Journot-Catalino *et al.*, 2006). Therefore, the action

of these two transcription factors may also establish the two distinct zones of JA and SA activity

in the colonized root tip. WRKY11 is considered a negative regulator of basal plant resistance, but

a positive regulator of JA biosynthesis. WRKY70, on the other hand, is a positive regulator of SA,

but a negative regulator of JA. Furthermore, *WRKY70* itself is under negative regulation by

WRKY11 and JA-signaling (Journot-Catalino *et al.*, 2006). To test if this regulatory module is

also active in response to *Fo*5176 infection, we imaged our *WRKY11* and *WRKY70* reporter lines.

We found that under control, non-infected conditions *WRKY11* is expressed at a moderate level

throughout the plant in all tissues, as expected for a gene involved in negatively regulating basal

resistance (Fig. 5A, B). However, upon colonization of the vasculature by *Fo*5176, the pattern of

WRKY11 expression changed, now confined to the vasculature only, with a clear expression

maximum in the cells neighboring the colonized cells, overlapping the maxima of *AOS* and *ERF1*

Figure 5: Expression of *WRKY11* in an uninfected control and following colonization of the root by *Fo*5176
A, B: Overlay of bright field and fluorescence (A), and fluorescence-only (B) images of an uninfected control root expressing the *WRKY11* reporter (blue). These overview images were taken with 12x magnification. **C, D**: Overlay of bright field and fluorescence (C), and fluorescence-only (D) images of a root tip expressing the *WRKY11* reporter (yellow) in response to colonization by *Fo*5176 (magenta). The colonization front is marked by magenta bars.

(Fig. 5C, D). Expression was removed from the surrounding tissues (epidermis, cortex), possibly

to allow for the expression of basal resistance genes normally repressed by WRKY11. This

enhanced expression pattern persisted for a long stretch up the root, before it reverted to normal

levels in cells closer to the root base, far away from the colonization site. This long tail of

expression mirrored that of *AOS* expression but was differed from that of *ACS2* or *EDS16*. Thus,

transcriptional activation of JA biosynthesis and signaling indeed overlaps with the activation of

its regulator WRKY11.

We then imaged the *WRKY70* reporter to see if its proposed role as a positive regulator of SA-,

and negative regulator of JA-signaling is also reflected in its expression pattern in response to

Figure 6: Expression of *WRKY70* in an uninfected control and following colonization of the root by *Fo*5176
A, B: Overlay of bright field and fluorescence (A), and fluorescence-only (B) images of an uninfected control root expressing the *WRKY70* reporter (yellow). **C, D**: Overlay of bright field and fluorescence (C), and fluorescence-only (D) images of a root tip expressing the *WRKY70* reporter (yellow) in response to colonization by *Fo*5176 (magenta). The colonization front is marked by magenta bars. The yellow brace brackets indicate the zone in which *WRKY70* appears to be excluded from the vascular tissue.

colonization by *Fo*5176. Under control conditions, *WRKY70* was expressed in all cells and tissues

of the root (Fig. 6A, B). Following colonization by *Fo*5176, we observed some striking changes.

First, there was strong upregulation in the cells neighboring the colonized tissue, with a maximum

in cells directly in contact with the colonized cells, overlapping the maximum seen for SA

biosynthesis and signaling (Fig. 6C, D). Second, in the group of cells slightly removed from the

colonization site, which show the JA biosynthesis/signaling maximum, *WRKY70* expression was

excluded from the vasculature, the tissue that expresses JA markers and *WRKY11*, while still

expressed in the surrounding tissues (Fig. 6C, D, yellow bracket). Slightly further up the root,

WRKY70 expression was restored in the vasculature (Fig. 6C, D). Thus, WRKY70 may indeed be

responsible for promoting SA biosynthesis and signaling in the HR zone, while the mutual
antagonism of WRKY11 and WRKY70 could contribute to establishing these two distinct
response zones.

Discussion

With the work described here we aimed at clarifying some of the inconclusive existing large-scale
transcriptomic data concerning the roles of JA, ET and SA in regulating the plant's defense against
the fungal pathogen *Fo*5176. Consistent with our own microscopic observations, and previously
described by Cyzmmek et al. (2007), *Fo*5176 infects and colonizes *A. thaliana* almost exclusively
via the root tip. We therefore reasoned that previous bulk transcriptomic analyses using whole
seedlings, or the entire root system were insufficient to detect highly specific events occurring
within the limited amount of infected tissue. Furthermore, general MAMP responses, initiated by
most root cells in response to the plant sensing the fungal hyphae in its rhizosphere, or growing
outside along the root, would most likely drown out any specific PAMP responses launched by the
few cells under actual acute infection. This hypothesis was strengthened by our initial observation,
that the infected tissue undergoes immediate cell death (Fig. 1), thereby limiting the amount of
infected tissue available for transcriptomic analyses even further. By employing a microscopy-
based approach to live-image the plant's response to infection in real time and at individual cell
resolution, we were then indeed able to show that JA, ET and SA all contribute to the plant's
defense against *Fo*5176, albeit in clearly distinct groups of cells.

An SA signaling pathway establishes the HR zone next to the colonization site

We first noticed that colonization of the vasculature was accompanied by local cell death of the
entire infected root tip. As the fungus continued to move through the root, colonizing more cells,
it formed a clearly visible 'colonization front' in the vasculature. As this colonization front moved
forward, a 'cell death front' moved in line with the colonization front. The only difference between
these two fronts being that the cell death front ran along all tissues of the root, from epidermis to
epidermis, and not just in the vasculature, as is seen with the colonization front (Fig. 7). We
hypothesized that this cell death front was the result of the plant's own HR, rather than
necrotrophic behavior by the fungus, and therefore imaged our reporters for SA biosynthesis and
signaling, since SA is a known inducer of HR (Pitsili *et al.*, 2020). Indeed, both SA biosynthesis
and signaling, were upregulated precisely in a small group of cells directly in contact with the

Figure 7: Model showing the distinct activity zones of the phytohormones and transcription factors
*Fo*5176 infects an *A. thaliana* root tip via the meristem in the tip, where it colonizes the vasculature (purple cells = colonized zone). The tissue around this zone undergoes cell death (dark grey cells). As the colonization front (purple line) progresses in the vasculature, so does a cell death front (black line) across all tissues. In an apparent attempt to prevent the fungus from infecting more tissue, the plant triggers a cell death response in a group of cells immediately adjacent to the colonization front (yellow cells), the HR zone. HR is activated by WRKY70 via SA. Slightly removed from the HR and colonized zones, the plant launches an active defense response to combat the pathogen in the defense response zone (orange cells). This response is dependent on WRKY11 and JA/ET-biosynthesis and signaling. WRKY11/JA and WRKY70/SA are mutually antagonistic, thereby establishing the two spatially separate zones of distinct action.

already colonized and dead tissue. Our interpretation of this observation is that SA induces HR in the cells that will otherwise be colonized next by the pathogen, thereby trying to prevent the fungus from spreading further. In our model we designated this group of cells the 'HR-zone' (Fig. 7). JA, on the other hand, was upregulated in a distinct group of vascular cells slightly (4-7 cells) removed from the HR-zone. In this region, that we named the 'defense response zone', we think the plant launches an active defense response to combat the invader (Fig. 7). Such defense can include the activation of NADPH oxidases to produce ROS in the apoplast that may damage the fungus, an acidification of the apoplast via proton pumps, or the production of antimicrobial compounds (Wang *et al.*, 2022*a*). ET activity bridges the boundary between these two zones, but most likely contributes to the defense response zone with JA.

WRKY70 acts via SA to establish the HR zone, and WRKY11 acts via JA to establish a defense response zone'.

We then asked how this distinct spatial separation of the two hormonal responses can be achieved and investigated the role of a WRKY transcription factor module, consisting of WRKY11 and WRKY70 that have previously been described as controlling JA and SA signaling respectively

(Journot-Catalino *et al.*, 2006). In this regard, we not only found that *WRKY11*, a positive regulator

of JA, was indeed upregulated in the defense response zone, while *WRKY70*, a positive regulator

of SA, was upregulated in HR-zone, but we also observed that these two transcription factors

altered their expression patterns. In the uninfected state, *WRKY11* was expressed in all cells and

tissues of the root, in accordance with a proposed role as a negative regulator of basal defense gene

expression (Journot-Catalino *et al.*, 2006). This is consistent with the hypothesis that under regular

growth conditions, defense genes need to be downregulated/repressed, to allow for normal

development of the plant, as part of the defense-development trade-off. Following colonization of

the plant by *Fo*5176, *WRKY11* was robustly upregulated in the defense response zone, likely

activating JA biosynthesis and signaling. In contrast, *WRKY11* expression was no longer observed

in all other tissues. This would enable the release of basal defense genes from repression and prime

such tissue for a possible pathogen attack. Furthermore, WRKY11 would also contribute to the

spatial separation of the defense response and HR zones, since WRKY11 and JA jointly repress

SA via WRKY70. Consequently, *WRKY70* expression is excluded from the defense response zone

following colonization by *Fo*5176 and most likely excludes WRKY11 and JA from the HR zone.

Thus, the mutual exclusion of WRKY11 and WRKY70 appears to establish the separate defense

response and HR zones (Fig. 7).

ET-signaling and RBOHD-derived ROS may establish the boundary between HR and

defense response zone

Importantly, this WRKY-module regulates SA and JA, but has so far not been linked to the

regulation of ET. Interestingly, both JA and ET-induced defense signaling has been linked to the

PLANT ELICITOR PEPTIDE (PEP) pathway (Liu *et al.*, 2013; Tintor *et al.*, 2013; Holmes *et al.*,

2018). Liu *et al.* (2013) have shown that PEP1, via the PEP RECEPTORs 1 and 2 and

phosphorylation of the cytoplasmic BOTRYTIS-INDUCED KINASE 1 (BIK1), induces ethylene-

dependent defense genes, including *ERF1*. Holmes *et al.* (2018) demonstrated the requirement of

the JA-receptor COI1 for PEP1-induced growth inhibition (as an indirect readout for activated

immune signaling) and apoplastic ROS-burst via a member of the RBOH family of NADPH

oxidases. In our previous work we showed that all of these components, *PEP1*, *PEPR1* and *2*,

BIK1, and *RBOHD*, are also locally induced in the vascular cells neighboring the *Fo*5176-

colonized zone (Calabria *et al.*, 2022). Of these components, *PEP1* and *PEPR2* show the strongest

induction, with both not just induced in cells of the defense response zone, but also in a long stretch

covering the entire region from the HR zone and the defense response zone (Calabria *et al.*, 2022).

It is therefore a possibility that this PEP pathway activates ET biosynthesis and signaling. As

mentioned above, the observed expression maximum of the ET biosynthesis marker *ACS2* did

overlap with the JA-biosynthesis marker, but also appeared to partially cover the HR zone. Thus,

one speculation could be that ET signaling in the region between the HR and defense response

zones is required to form a boundary between these two zones, and possibly to prime the cells (Fig.

8). The co-incidence of ET and JA in the defense response zone would then launch the full-blown

defense response (Fig. 8). In this case, the PEP pathway would act as an amplifier of both, JA- and

ET-signaling (Liu *et al.*, 2013; Holmes *et al.*, 2018; Calabria *et al.*, 2022). *RBOHD* activation

appeared to be strongest in the HR and boundary zones, and this is also where we observed 'root

browning' (Calabria *et al.*, 2022). Thus, RBOHD may play a dual role in promoting cell death in

the HR zone and establishing the boundary. Indeed, it was previously shown that RBOHD may

play such a dual function, as RBOHD seemed to contribute to HR-induction in infected cells, while

suppressing HR in cells slightly removed from the infection site (Torres *et al.*, 2005; Pogány *et*

al., 2009; Lukan and Coll, 2022). In this model, first hypothesized by Torres *et al.,* (2005),

RBOHD suppresses SA-accumulation in cells that have been primed for attack, but are not under

Figure 8: Hypothetical model of how the phytohormones and the PEP pathway may establish the different zones and boundary

In the cells of the HR zone, PEP1, via PEPR2 and BIK1, activates RBOHD to produce ROS. In combination with an over-accumulation of SA via WRKY70- and cell damage-signaling, these two factors induce HR. In the boundary, RBOHD still produces ROS, but SA no longer accumulates to high levels, since WRKY70 and WRKY11 functions exclude each other from this zone, and there is also no cell damage to induce SA. ROS therefore inhibits HR via SA. In these cells, PEP1 also activates ET biosynthesis, and ROS and ET together prime these cells for active defense. In the cells of the defense response zone, WRKY11 can activate JA biosynthesis. JA and ET jointly activate active defense via ERF1.

acute attack, thereby preventing HR. In cells under attack, however, cell damage would cause an

elevation of SA-levels to the point where RBOHD can no longer counteract it, thus triggering HR

(Torres *et al.*, 2005; Pogány *et al.*, 2009; Lukan and Coll, 2022). In this second scenario, SA

accumulation and RBOHD-produced ROS therefore together result in HR. This model fits very

well with our observations from Calabria *et al.,* (2022) and the work described here, and we thus

speculate that this could be how the boundary between the HR and defense response zone is created

(Fig. 8).

Outlook and conclusion

Whilst the inclusion of the WRKY-module and the dual role of RBOHD lead to an elegant

hypothesis of how the separate zones can be established, further work is needed to validate these

observations. First, it will be important to image the phytohormone markers in *wrky* mutant plants,

to definitively establish that they act upstream of the hormone pathways. Further, it is not clear

how WRKY11 and WRKY70 can both be expressed in all cells and tissues under normal

conditions, if they are mutually exclusive. Similarly, under normal conditions, JA and SA are not

expressed in all tissues, despite the expression of the WRKY transcription factors. This could be a

matter of lower expression levels or the lack of additional co-activators/co-repressors, but so far,

we have no proof for this. Furthermore, while we found that the HR-zone spreads out across all

tissues of the root, we only found WRKY70 and SA upregulation in the vasculature, not in the

outer tissues. Thus, it is not clear what actually induces the HR in these cells. This could, however,

be a normal response to a dying vasculature. Should the vasculature undergo cell death, it is

conceivable that the loss of conductivity and structural integrity of the inner cylinder of the root

spontaneously induces cell death of the outer tissue, without the need of active SA-signaling.

Finally, *rbohd* mutants need to be investigated for boundary formation, and the inclusion of

RBOHF may help differentiate between ROS production involved in HR and boundary formation,

and ROS involved in active defense. It will be interesting to further investigate the different

pathways activated specifically in the defense response and HR zones.

We believe that the work reported here is an important step in uncovering very local spatial plant

responses to this fungus. It also highlights the need for further high-resolution analysis to resolve

the plant's immune response to pathogenic infection on an individual cell level within intact tissue,

that has been lacking so far. By building on these initial results, it should be possible to further

map the distinct spatial and temporal responses of other plant pathways to infection by *F.*
oxysporum, thereby creating a root atlas of the plant's immune system.

Material & Methods:

Cloning of *A. thaliana* reporter constructs

All expression clones for the different reporter lines were assembled as in Calabria *et al.* (2022),
and the putative promoter sequences are from our pGG-PIP vector collection (Calabria *et al.*,
2022). The expression clones are based on the pGGZ003 GreenGate destination vector backbone
from (Lampropoulos *et al.*, 2013). GreenGate modules for the Nuclear Localization Signal
(pGGD007), UBQ terminator (pGGE009) and hygromycin resistance cassette (pGGF005) were
also from the original GreenGate kit (Lampropoulos *et al.*, 2013). The B- and C-modules with
mTurquoise2 (amplified from an AddGene-derived template (Goedhart *et al.*, 2012)), as well as
the B-module with *mNeonGreen* (amplified from an AddGene-derived template (Shaner *et al.*,
2013)) and the C-module with *LTI6b* (amplified from total cellular *A. thaliana* Columbia DNA)
were created using the pGGB000, pGGC000 and pGGD000 entry clones from the original
GreenGate kit (Lampropoulos *et al.*, 2013). Expression clone assembly was done with the NEB
Golden Gate Assembly Kit (BsaI-HF® v2).

Plant growth and transformation

A. thaliana Columbia plants were grown until flowering stage at day/night conditions of 21 °C/18
°C, 16 h/8 h with a light intensity of 120 mmol m^{-2} s^{-1} and constant relative humidity of 70%
(Somssich, 2018). Plants were transformed using a slightly modified *Agrobacterium tumefaciens*-
mediated floral dip protocol (Clough and Bent, 1998; Somssich, 2019). *A. tumefaciens* strain
GV3101 *pMP90 pSoup* carrying the plasmids of interest were grown in 250 mL of 2×YT media
supplemented with 50 µg/ml rifampicin, 25 µg/ml gentamycin and 100 µg/ml spectinomycin for
16 h at 28 °C, shaking at 200 rpm (Miller, 1972; Holsters *et al.*, 1980; Koncz and Schell, 1986;
Hellens *et al.*, 2000). *A. tumefaciens* cultures were centrifuged at 3750 rpm for 20 min, and the
pellet resuspended in 300 mL of 5% sucrose solution with 0.08% Silwet L-77 (Clough and Bent,
1998). Plants were inverted and the flowers submerged in *A. tumefaciens* solution for 30 s with
agitation. Plants were briefly patted dry to remove excess *A. tumefaciens* before laying horizontal
in a plant tray covered with plastic film to maintain humidity at room temperature overnight. The

following day, plants were reverted upright and returned to the original growth conditions. The
floral dipping procedure was repeated after a 7-day interval and upon maturation of siliques, seeds
were harvested. Harvested seeds were dried for at least two weeks and then surface-sterilized by
washing them for 2-3 h in 75% ethanol with 0.1% Triton X-100 on an upright tube rotator. The
seeds were then decanted onto filter paper in a laminar flow cabinet to evaporate the ethanol. The
dry seeds were then sprinkled onto a petri dish with half-strength basal Murashige & Skoog (MS)
medium with vitamins supplemented with 30 µg/ml hygromycin (Murashige and Skoog, 1962).
Following a 3–5-day stratification period in darkness at 4 °C, the plates were placed into a growth
cabinet and grown for 10 to 14 days. After this, seedlings that survived selection were transferred
to fresh plates without antibiotics and grown until they were too big for the plates, at which point
they were transferred to soil until maturation of siliques.

Fungal growth and transformation

The *Fo*5176 line expressing the cytoplasmic tdTomato (tdT) was described in our previous paper,
Calabria *et al.* (2022). It was created using plasmid pMAI32, carrying a constitutive promoter from
the *actin* gene from ascomycete *Leptosphaeria maculans* (Idnurm *et al.*, 2017). The *tdT* coding
sequence was amplified from an AddGene-derived template and inserted into pMAI32 via Gibson
assembly (Shaner *et al.*, 2004). The resulting expression clone was used to transform *A.*
tumefaciens strain EHA105, and positive colonies were selected on LA plates supplemented with
50 µg/ml kanamycin. *Fo*5176 was cultured on potato dextrose agar (PDA) plates (Idnurm *et al.*,
2017). To obtain spores for transformation, a 50 ml half strength potato dextrose broth was
inoculated with five 5 mm plugs from the PDA plates. This culture was incubated at room
temperature while shaking at 150 rpm for five days, then filtering through miracloth, and the spores
harvested via centrifugation at 3000 rcf for 5 min. The spores were then resuspended in sterile
water. For the transformation, the *A. tumefaciens* strain was cultured overnight in LB
supplemented with kanamycin, and fungal spores and bacteria were then mixed on induction
medium plates and cocultured for three days (Idnurm *et al.*, 2017). For selection of positive
transformants, an overlay of PDA supplemented with 50 µg/ml hygromycin, and 100 µg/ml
cefotaxime was used. The transformants that grew through the overlay were subcultured on PDA
containing hygromycin and cefotaxime and allowed to produce conidia. The final strain was then
derived from a single conidium.

Plant-fungus co-cultivation and infection

Plant-fungus co-cultivation was done as in Calabria et al. (2022), which uses an adapted protocol from Tintor et al. (2020): *Fo*5176 was routinely maintained on PDA plates at room temperature. For each co-cultivation experiment, five 5 mm plugs were cut from the plate to inoculate a 50 ml yeast nitrogen base (YNB) medium supplemented with 1% sucrose, and this culture was incubated at room temperature for four days, shaking at 120 rpm. Spores were harvested by filtering through mirocloth and centrifugation at 3000 rcf for 10 min and were then resuspended in 25 ml autoclaved MilliQ water. The *A. thaliana* plants were grown on vertical plates with half-strength basal MS medium with vitamins for 11 days, and then transferred to a horizontal plate, in which the MS medium was removed, except for a roughly 2-3 cm strip at the top (Tintor *et al.*, 2020). The rest of the plate was filled up with quarter-strength liquid basal MS medium with vitamins. The seedlings were places onto the narrow solid medium strip, with the roots hanging into the liquid medium. Fungal spores were then added into the liquid medium. The plates were wrapped in aluminum foil, with only the above-ground organs exposed to light and placed back into the growth chamber.

Imaging

We observed earliest infection of *A. thaliana* by *Fo*5176 at three days post inoculation (dpi) with spores, so we imaged seedlings and fungi daily from 3 dpi to 11 dpi. Imaging was done on a Leica M205 FA stereomicroscope using ET CFP (ET436/20x ET480/40m), ET mCHER (ET560/40x ET630/75m) and F/AL488 (ET480/40x ET535/50m) filters for the detection of mTurquoise2, tdTomato and mNeonGreen, respectively. A magnification of 80× was used unless noted otherwise in the figure legend, and imaging and detection conditions, such as illumination power, exposure time, gain, etc. were kept constant across all measurements, to make images comparable and allow for at least a semi-quantitative assessment of expression strength of the different reporters. Images were taken with the Leica Application Suite software, then exported and processed using Fiji and the GNU image manipulation program (GIMP) (Schindelin *et al.*, 2012).

Acknowledgements:

This work was funded by the Australian Research Council (grant no. DE200101560) and was further supported by a seed grant from the Melbourne University Botany Foundation. LW was supported by the China Scholarship Council. H-WC was supported by a Graduate Research

478 Scholarship from the University of Melbourne. Imaging was done on instruments maintained by

479 the Biological Optical Microscopy Platform (BOMP) and the BioSciences Microscopy Unit at the

480 University of Melbourne. We would like to thank Dr. Imre E. Somssich for critical reading of the

481 manuscript and helpful comments and suggestions.

References

Berrocal-Lobo M, Molina A. 2004. Ethylene Response Factor 1 Mediates Arabidopsis
Resistance to the Soilborne Fungus Fusarium oxysporum. Molecular Plant-Microbe Interactions
17, 763–770.

Calabria J, Rast-Somssich MI, Wang L, Chen H-W, Watt M, Idnurm A, Persson S,
Somssich M. 2022. pGG-PIP: A GreenGate (GG) entry vector collection with Plant Immune
system Promoters (PIP). bioRxiv, 521163.

Chen YC, Wong CL, Muzzi F, Vlaardingerbroek I, Kidd BN, Schenk PM. 2014. Root
defense analysis against Fusarium oxysporum reveals new regulators to confer resistance.
Scientific reports **4**, 5584.

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. The Plant Journal **16**, 735–43.

Cole SJ, Yoon AJ, Faull KF, Diener AC. 2014. Host perception of jasmonates promotes
infection by Fusarium oxysporum formae speciales that produce isoleucine- and leucine-
conjugated jasmonates. Molecular Plant Pathology **15**, 589–600.

Czymmek KJ, Fogg M, Powell DH, Sweigard J, Park S-Y, Kang S. 2007. In vivo time-lapse
documentation using confocal and multi-photon microscopy reveals the mechanisms of invasion
into the Arabidopsis root vascular system by Fusarium oxysporum. Fungal genetics and biology
44, 1011–23.

Diener AC, Ausubel FM. 2005. RESISTANCE TO FUSARIUM OXYSPORUM 1, a Dominant
Arabidopsis Disease-Resistance Gene, Is Not Race Specific. Genetics **171**, 305–321.

Dita M, Barquero M, Heck D, Mizubuti ESG, Staver CP. 2018. Fusarium Wilt of Banana:
Current Knowledge on Epidemiology and Research Needs Toward Sustainable Disease
Management. Frontiers in Plant Science **9**, 1–21.

van der Does HC, Duyvesteijn RGE, Goltstein PM, van Schie CCN, Manders EMM,
Cornelissen BJC, Rep M. 2008. Expression of effector gene SIX1 of Fusarium oxysporum
requires living plant cells. Fungal genetics and biology **45**, 1257–64.

Edgar CI, McGrath KC, Dombrecht B, Manners JM, Maclean DC, Schenk PM, Kazan K.
2006. Salicylic acid mediates resistance to the vascular wilt pathogen Fusarium oxysporum in the
model host Arabidopsis thaliana. Australasian Plant Pathology **35**, 581.

Goedhart J, von Stetten D, Noirclerc-Savoye M, Lelimousin M, Joosen L, Hink MA, van
Weeren L, Gadella Jr. TWJ, Royant A. 2012. Structure-guided evolution of cyan fluorescent
proteins towards a quantum yield of 93%. Nature communications **3**, 751.

Gordon TR. 2017. Fusarium oxysporum and the Fusarium Wilt Syndrome. Annual Review of Phytopathology **55**, 23–39.

Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM. 2000. pGreen: a versatile and flexible binary Ti vector for Agrobacterium-mediated plant transformation. Plant molecular biology **42**, 819–32.

Holmes DR, Grubb LE, Monaghan J. 2018. The jasmonate receptor COI1 is required for AtPep1-induced immune responses in Arabidopsis thaliana. BMC Research Notes **11**, 555.

Holsters M, Silva B, van Vliet F, *et al.* 1980. The functional organization of the nopaline A. tumefaciens plasmid pTiC58. Plasmid **3**, 212–230.

Hou S, Tsuda K. 2022. Salicylic acid and jasmonic acid crosstalk in plant immunity. Essays in Biochemistry, 1–10.

Idnurm A, Urquhart AS, Vummadi DR, Chang S, Van de Wouw AP, López-Ruiz FJ. 2017. Spontaneous and CRISPR/Cas9-induced mutation of the osmosensor histidine kinase of the canola pathogen Leptosphaeria maculans. Fungal Biology and Biotechnology **4**, 12.

Journot-Catalino N, Somssich IE, Roby D, Kroj T. 2006. The transcription factors WRKY11 and WRKY17 act as negative regulators of basal resistance in Arabidopsis thaliana. The Plant cell **18**, 3289–3302.

Kidd BN, Kadoo NY, Dombrecht B, Tekeoglu M, Gardiner DM, Thatcher LF, Aitken EAB, Schenk PM, Manners JM, Kazan K. 2011. Auxin Signaling and Transport Promote Susceptibility to the Root-Infecting Fungal Pathogen Fusarium oxysporum in Arabidopsis. Molecular Plant-Microbe Interactions **24**, 733–748.

Koncz C, Schell J. 1986. The promoter of TL-DNA gene 5 controls the tissue-specific expression of chimaeric genes carried by a novel type of Agrobacterium binary vector. Molecular & General Genetics **204**, 383–396.

Lampropoulos A, Sutikovic Z, Wenzl C, Maegele I, Lohmann JU, Forner J. 2013. GreenGate - a novel, versatile, and efficient cloning system for plant transgenesis. PLOS ONE **8**, e83043.

Lee D-H, Lee H, Belkhadir Y. 2021. Coding of plant immune signals by surface receptors. Current Opinion in Plant Biology **62**, 102044.

Liu Z, Wu Y, Yang F, Zhang Y, Chen S, Xie Q, Tian X, Zhou J-M. 2013. BIK1 interacts with PEPRs to mediate ethylene-induced immunity. Proceedings of the National Academy of Sciences of the United States of America **110**, 6205–6210.

Lukan T, Coll A. 2022. Intertwined Roles of Reactive Oxygen Species and Salicylic Acid Signaling Are Crucial for the Plant Response to Biotic Stress. International Journal of Molecular Sciences **23**, 5568.

Lyons R, Stiller J, Powell J, Rusu AG, Manners JM, Kazan K. 2015. Fusarium oxysporum triggers tissue-specific transcriptional reprogramming in Arabidopsis thaliana. PLOS ONE **10**, e0121902.

Ma L-J, van der Does HC, Borkovich KA, *et al.* 2010. Comparative genomics reveals mobile pathogenicity chromosomes in Fusarium. Nature **464**, 367–73.

Martínez-Soto D, Yu H, Allen KS, Ma L-J. 2022. Differential colonization of the plant vasculature between endophytic versus pathogenic Fusarium oxysporum strains. Molecular Plant-Microbe Interactions, 1–27.

Masachis S, Segorbe D, Turrà D, *et al.* 2016. A fungal pathogen secretes plant alkalinizing peptides to increase infection. Nature Microbiology **1**, 16043.

McCombe CL, Greenwood JR, Solomon PS, Williams SJ. 2022. Molecular plant immunity against biotrophic, hemibiotrophic, and necrotrophic fungi. Essays in Biochemistry **0**, 1–13.

Miller JH. 1972. Formulas and recipes. Experiments in molecular genetics. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory, 433.

Murashige T, Skoog F. 1962. A Revised Medium for Rapid Growth and Bio Assays with Tobaoco Tissue Cultures. Physiologia Plantarum **15**, 473–497.

Pitsili E, Phukan UJ, Coll NS. 2020. Cell Death in Plant Immunity. Cold Spring Harbor Perspectives in Biology **12**, a036483.

Pogány M, von Rad U, Grün S, Dongó A, Pintye A, Simoneau P, Bahnweg G, Kiss L, Barna B, Durner J. 2009. Dual Roles of Reactive Oxygen Species and NADPH Oxidase RBOHD in an Arabidopsis- Alternaria Pathosystem. Plant Physiology **151**, 1459–1475.

Redkar A, Sabale M, Schudoma C, *et al.* 2022. Conserved secreted effectors contribute to endophytic growth and multi-host plant compatibility in a vascular wilt fungus. The Plant Cell.

Schindelin J, Arganda-Carreras I, Frise E, *et al.* 2012. Fiji: an open-source platform for biological-image analysis. Nature methods **9**, 676–82.

Shaner NC, Campbell RE, Steinbach PA, Giepmans BNG, Palmer AE, Tsien RY. 2004. Improved monomeric red, orange and yellow fluorescent proteins derived from Discosoma sp. red fluorescent protein. Nature biotechnology **22**, 1567–72.

Shaner NC, Lambert GG, Chammas A, *et al.* 2013. A bright monomeric green fluorescent protein derived from Branchiostoma lanceolatum. Nature methods **10**, 407–9.

Somssich M. 2018. A short history of Arabidopsis thaliana (L.) Heynh. Columbia-0. PeerJ Preprints **e26931v3**, 1–7.

Somssich M. 2019. A short history of plant transformation. PeerJ Preprints, 1–28.

Summerell BA. 2019. Resolving Fusarium: Current Status of the Genus. Annual review of phytopathology **57**, 323–339.

Takken FLW, Rep M. 2010. The arms race between tomato and Fusarium oxysporum. Molecular Plant Pathology **11**, 309–314.

Thatcher LF, Manners JM, Kazan K. 2009. Fusarium oxysporum hijacks COI1-mediated jasmonate signaling to promote disease development in Arabidopsis. The Plant Journal **58**, 927–39.

Tintor N, Paauw M, Rep M, Takken FLW. 2020. The root-invading pathogen Fusarium
oxysporum targets pattern-triggered immunity using both cytoplasmic and apoplastic effectors.
New Phytologist **227**, 1479–1492.

Tintor N, Ross A, Kanehara K, Yamada K, Fan L, Kemmerling B, Nürnberger T, Tsuda
K, Saijo Y. 2013. Layered pattern receptor signaling via ethylene and endogenous elicitor
peptides during Arabidopsis immunity to bacterial infection. Proceedings of the National
Academy of Sciences of the United States of America **110**, 6211–6216.

Torres MÁ, Jones JDG, Dangl JL. 2005. Pathogen-induced, NADPH oxidase–derived reactive
oxygen intermediates suppress spread of cell death in Arabidopsis thaliana. Nature Genetics **37**,
1130–1134.

Wang L, Calabria J, Chen H-W, Somssich M. 2022*a*. The Arabidopsis thaliana – Fusarium
oxysporum strain 5176 pathosystem: an overview (M Höfte, Ed.). Journal of Experimental
Botany, erac263.

Wang Y, Pruitt RN, Nürnberger T, Wang Y. 2022*b*. Evasion of plant immunity by microbial
pathogens. Nature Reviews Microbiology **20**, 449–464.

Zhou F, Emonet A, Dénervaud Tendon V, Marhavý P, Wu D, Lahaye T, Geldner N. 2020.
Co-incidence of damage and microbial patterns controls localized immune responses in roots.
Cell **180**, 440-453.e18.

 Supplementary Figures

Figure S1: Expression patterns of *EDS16* and *PR1* in uninfected control plants
A, B: Overlays of bright field and fluorescence images showing fluorescence of the *EDS16* reporter in a root tip (A) and in differentiated cells (B). **C, D**: Fluorescence (C) and bright field (D) image showing fluorescence of the *PR1* reporter in differentiated cells.

Figure S2: Expression patterns of *AOS*, *ERF1*, and *ACS2* in uninfected control plants
A: Overlay of bright field and fluorescence image showing expression of *AOS* at lateral root emergence sites. 12x magnification. **B:** Magnification of such a site. **C, D**: Overlay of bright field and fluorescence (C) and fluorescence-only (D) images showing onset of *ERF1* expression at the start of the differentiation zone. 32x magnification. **E, F**: Overlay of bright field and fluorescence (E) and fluorescence-only (F) images showing fluorescence from the *ERF1* reporter in differentiated cells. **G, H**: Overlay of bright field and fluorescence (G) and fluorescence-only (H) images showing expression of *ACS2* in the root tip and young differentiating cells. Note that these images are composites of three different images to account for a changing focal plane. **I, J**: Overlay of bright field and fluorescence (I) and fluorescence-only (J) images showing fluorescence from the *ACS2* reporter in differentiated cells.

Figure S3: Expression of *PDF1.2* in differentiated tissue following colonization of the root by *Fo*5176
A, B: Bright field (A), and fluorescence (B) images of the differentiation zone of a root expressing the *PDF1.2* reporter (yellow) in response to colonization by *Fo*5176 (magenta). Colonization via the tips of lateral roots has reached the main root at its differentiation zone. In this zone *PDF1.2* is responsive to the infection, showing upregulation in the vasculature.