1 **Relation between pH alterations in cellular models of amyotrophic lateral sclerosis and TDP-43**
2 **protein aggregation**

Charlotte Slek[1&], Yara Al Ojaimi[1&], Sylviane Marouillat[1], Patrick Vourc'h[1,2], Débora Lanznaster[1*#],
Hélène Blasco[1,2#]

1-UMR 1253, iBrain, Université de Tours, INSERM, Tours, France

2-Service de Biochimie et Biologie Moléculaire, CHRU de Tours, France.

[&,#] Equal contribution

*Corresponding author: Débora Lanznaster, PhD

E-mail: debora.lanznaster@univ-tours.fr

Abstract

Amyotrophic lateral sclerosis (ALS) is the most common degenerative disease of the motor neuron.
TDP-43 aggregates constitute a hallmark of ALS and play a major role in the motoneuron degeneration.
Several factors can influence the ability of TDP-43 to form aggregates. It was previously shown that, *in*
vitro, mild pH modification can induce a conformational change in TDP-43 structure thus increasing its
propensity to aggregate. Therefore, we first analyzed the intracellular pH alterations in a cellular model
of ALS as well as fibroblasts derived from ALS patients. HEK293T cells overexpressing wildtype TDP-
43 presented an increase in the fluorescence signal of the probe BCECF, which is sensitive to low pH.
We also observed an increase in BCECF signal (i.e., low pH) in cultures of fibroblasts obtained from
ALS patients as compared to controls. On the other hand, we tested the effect of pH alterations on the
localization and aggregation of TDP-43. HEK293T cells overexpressing wildtype TDP-43 were
incubated for 1h or 2h with solutions of different pH (6.4, 7.2 and 8.0). Incubation of cells for 1h in
solutions of pH 6.4 and 8.0 increased TDP-43 aggregates, an effect lost after a 2h incubation. In these
conditions, TDP-43 aggregation was accompanied by its mislocalization to the cytoplasm. Our results
suggest that a decrease in cellular pH is observed in a cellular model of TDP-43 proteinopathy and in
fibroblasts derived from ALS patients compared to controls. Alterations in cellular pH also seemed to
have an effect on the aggregation and cellular localization of TDP-43. Further studies should focus on
understanding how cellular pH alterations affect TDP-43, which might offer a new way to mitigate TDP-
43 pathology in ALS.

Keywords: amyotrophic lateral sclerosis, TDP-43, aggregates, pH

1. Introduction

Amyotrophic lateral sclerosis (ALS) is a degenerative disease characterized by the progressive death of motor neurons, leading to paralysis of skeletal muscles and eventually respiratory muscles, resulting in the death of patients within 3 to 5 years following the onset of the disease (Hardiman et al., 2017). Despite the progress made in understanding the pathophysiology of ALS, the mechanisms underlying motor neurodegeneration are still poorly understood. An abnormal aggregation of various proteins, particularly the TAR DNA binging protein 43 (TDP-43) has been reported in motor neurons. In fact, TDP-43 aggregates are considered as the hallmark of ALS (Hergesheimer et al., 2019) which has been identified as the major component of intracellular pathogenic aggregates in patients with ALS (Arai et al., 2006; Neumann et al., 2006).

Physiological TDP-43 shuttles between the nucleus and the cytoplasm but in pathological conditions, TDP-43 forms cytosolic aggregates, leading to the loss of protein function and cellular homeostasis. For instance, pathological mechanisms observed in ALS such as ischemia, mitochondrial alterations, and oxidative stress, which can affect intracellular pH have been associated with the cytoplasmic mislocalization and aggregation of TDP-43 (Gao et al., 2019; Wang et al., 2019; Wang et al., 2013). Concomitantly, several studies have shown that alterations in the pH can affect the structure of recombinant TDP-43 (Doke and Jha, 2022; Mompean et al., 2016a). However, no studies have assessed the effect of such alterations on TDP-43 in cells. We therefore hypothesize that there is a bidirectional interplay between pH-altering metabolic alterations in ALS and TDP-43 proteinopathy. The goal of our study was therefore to assess whether the intracellular pH was changed in a cellular model of ALS and patient fibroblasts, and investigate the impact of such changes on the localization and aggregation of cellular TDP-43.

2. Materials and Methods

2.1 Cells

HEK293T cells (ATCC, USA) were cultured in standard conditions (Lanznaster et al., 2019) and plated at a concentration of 0.3×10^6 cells/well the day before transfection.

Dermal fibroblasts from five ALS patients (cultured from the biopsy obtained at diagnosis, information in Supplementary material) and one healthy subject (man, 60 years-old; from Lonza Ltd, Basel, Switzerland) with roughly the same passage number, were cultivated in DMEM supplemented with 10% (v/v) FBS and 1% of Penicillin/Streptomycin.

2.2 pH solutions

pH solutions were prepared to induce changes in intracellular pH, as previously described (Swietach et al., 2012). The stock solution was prepared with 125 mM NaCl, 4.5 mM KCl, 20 mM Hepes, 20 mM MES, 1 mM $CaCl_2$ and 1 mM $MgCl_2$ (original pH of 4.4). Once this solution was obtained, the pH was adjusted with NaOH or HCl in order to obtain three solutions of pH 6.4, 7.2 and 8.0. Cells were incubated with these solutions for 1h or 2h, which did not modify the viability of the cells (Supplementary material).

2.3 Overexpression of TDP-43

84   Overexpression of TDP-43 was done as previously described (Lanznaster et al., 2019). Briefly,
85   HEK293T cells were transiently transfected with 2µg or 3µg of TDP-43-expressing plasmid. Control
86   plasmid (pcDNA vector) was transfected at the highest concentration (3µg). Assays were realized 48 h
87   after transfection.

2.4 Assessment of intracellular pH

To analyse the intracellular pH, cells were incubated with 0.5µM BCECF-AM (2',7'-Bis-(2-
Carboxyethyl)-5-(and-6)-Carboxyfluorescein, Acetoxymethyl Ester, ThermoScientific), a fluorescent
probe sensitive to variations in intracellular pH (Supplementary material). After 15 minutes of
incubation, the fluorescence was read at 535nm by confocal microscopy (Carre et al., 2015). The
fluorescence intensity was analysed using ImageJ software and is given as Relative Fluorescence Unit
(RFU).

2.5 Localization and aggregation of TDP-43

To study the localization of TDP-43 and its aggregation in different pH conditions, HEK293T cells were
plated on coverslips pre-treated with poly-D-lysine (10 ug/mL) and transfected with the plasmid GFP-
wtTDP-43 (3 µg). After 48 hours, cells were incubated for either one or two hours with the different pH
solutions (6.4, 7.2, 8.0) previously described. Ten minutes before reading the fluorescence, 1µL of
Hoechst 33342 (1:1000) was added. Then, cells were fixed with 4% paraformaldehyde solution (diluted
in PBS). Images were obtained by confocal microscopy (Leica SP8) and 100 cells per condition were
analysed for TDP-43 localization (nuclear, cytoplasmic, or both) and aggregation.

2.6 Statistics analysis

The statistical analyses were carried out with the GraphPad Prism 8 software. Data is reported as mean
± Standard Error of the Mean (SEM). Differences in pH regarding the fibroblasts of control and ALS
patients were analysed with the Mann-Whitney test. The localization of the TDP-43 protein was
analysed with the two-way ANOVA test. Differences in pH regarding increasing concentrations of TDP-
43 and the influence of pH on TDP-43 aggregation were analysed with the Kruskal-Wallis test. Results
were considered statistically significant at $p<0.05$.

3. Results

3.1   pH alteration in cellular models of ALS

3.1.1   pH evaluation in a TDP-43-overexpressing cellular model of ALS

After 48h of TDP-43 overexpression, HEK293T cells were incubated with the fluorescent probe
BCECF-AM. The images shown in **Figure 1 (upper panel)** pinpoint to two main observations: 1) for
the control cells, a basal level of fluorescence is visible, probably due to ion exchanges present to
maintain physiological pH; and 2) cells overexpressing TDP-43 fluoresce more intensely than mock-
transfected cells. Control cells (basal fluorescence) show a mean fluorescence of 18.45±3.2 RFU, that
is statistically different only from TDP-43-overexpressing cells transfected with 3 µg of plasmid (TDP-
43 2 µg: 37.85 ± 5.8 RFU, $p=0.0662$; and TDP-43 3 µg: 43.69 ± 4.5 RFU, $p=0.0127$). These results
reflect a tendency for the pH to be altered in cells overexpressing TDP-43, an observation that seems to
be dependent on the quantity transfected of TDP-43 plasmid.

3.1.2   pH assessment in ALS patients' fibroblasts

The intracellular pH of dermal fibroblasts obtained from ALS patients is different from that of fibroblasts obtained from a healthy control. The images shown in **Figure 1 (lower panel)** reflect a more pronounced fluorescence of the BCECF probe in the fibroblasts of ALS compared to control (mean ± S.E.M.: 46.2 ± 0.9 RFU and 25.2 ± 1.2 RFU; $p=0.007$, respectively).

3.2   Relationship between pH changes and TDP-43 protein mislocalization and aggregation

3.2.1   Effect of the pH on the localisation of TDP-43

First, we analysed the cellular TDP-43 localisation for each pH condition (**Figure 2A-H**). Our results indicate that the localization of overexpressed TDP-43 varies with the pH. Accordingly, after an incubation of 1h at physiological pH (7.2), TDP-43 was found to be evenly distributed between the cytoplasmic and nuclear compartments. On the other hand, there was an exacerbation of TDP-43 mislocalization following pH alterations. A significant increase of cytoplasmic TDP-43 was observed at pH 6.4 ($p=0.04$) and pH 8 ($p=0.05$) compared to pH 7.2. However, after 2 hours of incubation no statistical significance was noted.

3.2.2   Effect of pH on TDP-43 aggregation

The analysis of cells overexpressing TDP-43 aggregates showed that there is a significant increase in protein aggregation following a 1h incubation in solutions of pH 6.4 and 8.0 compared to the physiological pH condition (7.2) ($p=0.05$; **Figure 2I**). This increase was not observed following 2 hours of incubation (**Figure 2J**), which is therefore in correlation with the results obtained previously.

4.   Discussion

Intracellular pH is decreased in cellular models of ALS

We found a decrease in the intracellular pH of fibroblasts from ALS patients, suggesting cellular acidosis. Concomitantly, several studies have shown that TDP-43 is mislocalized and aggregated in fibroblasts from ALS patients (Riancho et al., 2020; Sabatelli et al., 2015). The decrease in intracellular pH was also observed in HEK293T cells overexpressing TDP-43, a cellular model of ALS (Lanznaster et al., 2019). Under prolonged or chronic stressful conditions, the cell's ability to autoregulate its intracellular environment decreases which can explain the eventual acidification of the intracellular environment.

Acidosis is commonly observed in different pathologies of the central nervous system such as Alzheimer's disease, Parkinson's disease, multiple sclerosis, and vascular dementia (Arias et al., 2008; Ding et al., 2000; Friese et al., 2007; Harguindey et al., 2007; Vergo et al., 2011). The metabolic alterations and disease features that have been noted in ALS such as microglial activation (Morihata et al., 2000), impaired oxidative phosphorylation (Erecinska et al., 1993), endoplasmic reticulum stress (Aoyama et al., 2005) can be potentially linked to altered pH regulation. Interestingly, motoneurons were found to be particularly sensitive to acidic environments (Behan et al., 2013). A significant increase in acid-sensing ion channels 2 (ASIC2) was also noted in the motoneurons of G93A-hSOD1 mice and sporadic ALS patients, supporting the involvement of acidosis in ALS (Behan et al., 2013). In one study, increasing the intracellular pH in a mouse model of ALS seemed to increase the lifespan by 13% without affecting the severity of the symptoms (Kuo et al., 2014).

Acute altered intracellular pH exacerbates the mislocalization and aggregation of TDP-43 in a cellular model of ALS

TDP-43 is particularly sensitive to changes in its environment which can shift the equilibrium towards favouring the formation of oligomeric forms (Doke and Jha, 2022). Under stressful cellular conditions, TDP-43 undergoes liquid-liquid phase separation (LLPS) and is recruited to stress granules (Colombrita et al., 2009; Khalfallah et al., 2018). For instance, stress granules can be formed in the context of metabolic dysfunction caused by starvation, hypoxia, or mitochondrial abnormalities (Farny et al., 2009; Jain et al., 2016; Jin et al., 2022; Riback et al., 2017; van der Laan et al., 2012), which can enhance TDP-43 LLPS and aggregation. These stressors are partly linked to intracellular acidosis which affects the protonation state of charged amino acid residues and therefore alters the structure and conformation of proteins (Jin et al., 2022; Yao and Haddad, 2004). Such modifications might allow proteins like TDP-43 to act as biosensors of changes in the intracellular environment (Rabouille and Alberti, 2017).

Changes in the extracellular pH have been shown to induce parallel but less pronounced changes in the intracellular pH (Bouyer et al., 2004; Buckler et al., 1991; Glunde et al., 2002; Putnam, 2001). Here we showed, for the first time, that pH alterations in the cellular environment can increase TDP-43 aggregation and cytoplasmic mislocalization. In cells overexpressing TDP-43, we observed a decrease in the nuclear localization of TDP-43 at pH 6.4 and pH 8 compared to the physiological pH. The extent of TDP-43 mislocalization to the cytoplasm was less pronounced after a 2 h incubation with the different solutions, which might be reflecting the ability of the cell to adjust to acute environmental alterations and restore homeostasis. The same pattern was observed for TDP-43 aggregation which significantly increased following 1h of incubation at pH 6.4 and pH 8 but not after 2 h. For instance, cells were shown to exhibit a quick and sharp fall in their intracellular pH within seconds of being put under acidic stress, but then started to restore their pH after 2 hours (Adam et al., 2021). These results might suggest an implication of TDP-43 in the response of cells to abnormal pH changes.

How intracellular pH induces TDP-43 mislocalization and aggregation has not been well investigated. Since TDP-43 is known for its ability to spontaneously aggregate, particularly through its intrinsically disordered C-terminal domain (CTD) (Johnson et al., 2009), it is complicated to assess the effect of microenvironmental alterations on the full length protein. Therefore, most studies that investigated the effect of pH on TDP-43 folding have used isolated protein domains rather than the full-length protein (Patni and Jha, 2021; Pillai and Jha, 2019, 2020). In addition, the results obtained seem to vary according to the recombinant TDP-43 forms used and the protocols applied (i.e. protein concentration, temperature, buffer pH, salt concentration).

For instance, under extreme acidic conditions (pH 2.5), the RNA-recognition motifs (RRM1 and RRM2) of TDP-43 were shown to undergo a conformational opening reaction linked to the protonation of buried ionizable residues, thus exposing hydrophobic patches and forming metastable oligomers that act as precursors to solid aggregates (Pillai and Jha, 2020). A subsequent study revealed that the protonation of the side chain of a histidine residue at position 166 is critical for the misfolding of the RRM domains (Patni and Jha, 2021). However, the acidic pH conditions used in these studies are extreme values that are not usually observed in the cytoplasmic environment of cells, even in cases of severe acid-base disturbances (Bouyer et al., 2004; Buckler et al., 1991; Glunde et al., 2002; Tolkovsky and Richards, 1987). Other groups have tried to decipher the biochemical alterations induced by pH changes and affecting the folding and aggregation of TDP-43 under more physiologically relevant conditions. The N-terminal domain (NTD) of TDP-43 was found to be stably folded over a wide range of pH conditions, in part due to the presence of residues with negatively charged carboxylate groups that mutually repel each other at pH values below 10 (Mompean et al., 2016b). Despite it not being itself unfolded, the NTD seems to mediate TDP-43 oligomerization and subsequent CTD misfolding (Tsoi et

al., 2017). While it was first suggested that this oligomerization is mediated by hydrophobic interactions
(Li et al., 2018), a more likely explanation is the pH-dependent decrease in electrostatic repulsion
(Babinchak et al., 2019; Mompean et al., 2016a). In fact, it seems like the balance between hydrophobic
interactions and electrostatic forces, both of which are affected by pH, is a key player in regulating the
folding and aggregation of TDP-43, particularly its CTD. Overall, it seems that TDP-43 aggregation is
mainly mediated by the CTD at the pH conditions between 6 and 8 that are used in our study.

Although the previously described studies show that TDP-43 domains can be structurally altered
upon pH changes, the impact observed on separate protein domains might not be reproducible in the
full-length protein given the presence of intramolecular interactions (Wei et al., 2016). In addition, all
these studies were performed on recombinant purified proteins, and not in cellular models of TDP-43
proteinopathy. Given the extensive post-translational modifications that TDP-43 can undergo inside
cells, such studies do not take into account these modifications that can highly affect the structure and
solubility of proteins (Hergesheimer et al., 2019). For these reasons, it is pivotal to study the effect of
pH changes on TDP-43 in more physiologically relevant settings, such as cellular models of ALS.

Conclusion

Given the reciprocal effect on one another of TDP-43 and pH alterations, it remains unclear
whether pathological TDP-43 drives pH changes or vice versa. This interplay between TDP-43 and pH
might take place through intermediary ALS-associated mechanisms (e.g. metabolic alterations,
mitochondrial dysfunction, hypoxia) and constitute a feedforward loop whereas TDP-43 proteinopathy
decreases the intracellular pH and intracellular pH alterations exacerbate TDP-43 proteinopathy. Our
results implicate intracellular pH changes in the pathophysiology of ALS and offer a new potential target
for therapeutic interventions. Future studies should look at the effect of pH alterations on TDP-43 in
neurons, which are particularly sensitive to changes in their environment.

Acknowledgements.

The authors would like to thank LabEx MAbImprove (ANR-10-LABX-53-01) and the Région Centre
Val de Loire for their financial support.

References

Adam, C., Paolini, L., Gueguen, N., Mabilleau, G., Preisser, L., Blanchard, S., Pignon, P., Manero, F., Le
Mao, M., Morel, A., et al. (2021). Acetoacetate protects macrophages from lactic acidosis-induced
mitochondrial dysfunction by metabolic reprograming. Nat Commun *12*, 7115.
Aoyama, K., Burns, D.M., Suh, S.W., Garnier, P., Matsumori, Y., Shiina, H., and Swanson, R.A. (2005).
Acidosis causes endoplasmic reticulum stress and caspase-12-mediated astrocyte death. J Cereb Blood
Flow Metab *25*, 358-370.
Arai, T., Hasegawa, M., Akiyama, H., Ikeda, K., Nonaka, T., Mori, H., Mann, D., Tsuchiya, K., Yoshida, M.,
Hashizume, Y., et al. (2006). TDP-43 is a component of ubiquitin-positive tau-negative inclusions in
frontotemporal lobar degeneration and amyotrophic lateral sclerosis. Biochem Biophys Res Commun
351, 602-611.
Arias, R.L., Sung, M.L., Vasylyev, D., Zhang, M.Y., Albinson, K., Kubek, K., Kagan, N., Beyer, C., Lin, Q.,
Dwyer, J.M., et al. (2008). Amiloride is neuroprotective in an MPTP model of Parkinson's disease.
Neurobiology of disease *31*, 334-341.
Babinchak, W.M., Haider, R., Dumm, B.K., Sarkar, P., Surewicz, K., Choi, J.K., and Surewicz, W.K. (2019).
The role of liquid-liquid phase separation in aggregation of the TDP-43 low-complexity domain. J Biol
Chem *294*, 6306-6317.

Behan, A.T., Breen, B., Hogg, M., Woods, I., Coughlan, K., Mitchem, M., and Prehn, J.H. (2013).
Acidotoxicity and acid-sensing ion channels contribute to motoneuron degeneration. Cell Death Differ
20, 589-598.
Bouyer, P., Bradley, S.R., Zhao, J., Wang, W., Richerson, G.B., and Boron, W.F. (2004). Effect of
extracellular acid-base disturbances on the intracellular pH of neurones cultured from rat medullary
raphe or hippocampus. J Physiol *559*, 85-101.
Buckler, K.J., Vaughan-Jones, R.D., Peers, C., Lagadic-Gossmann, D., and Nye, P.C. (1991). Effects of
extracellular pH, PCO2 and HCO3- on intracellular pH in isolated type-I cells of the neonatal rat carotid
body. J Physiol *444*, 703-721.
Carre, G., Ouedraogo, M., Magaud, C., Carreyre, H., Becq, F., Bois, P., Supuran, C.T., Thibaudeau, S.,
Vandebrouck, C., and Bescond, J. (2015). Vasorelaxation induced by dodoneine is mediated by calcium
channels blockade and carbonic anhydrase inhibition on vascular smooth muscle cells. J
Ethnopharmacol *169*, 8-17.
Colombrita, C., Zennaro, E., Fallini, C., Weber, M., Sommacal, A., Buratti, E., Silani, V., and Ratti, A.
(2009). TDP-43 is recruited to stress granules in conditions of oxidative insult. J Neurochem *111*, 1051-
1061.
Ding, D., Moskowitz, S.I., Li, R., Lee, S.B., Esteban, M., Tomaselli, K., Chan, J., and Bergold, P.J. (2000).
Acidosis induces necrosis and apoptosis of cultured hippocampal neurons. Exp Neurol *162*, 1-12.
Doke, A.A., and Jha, S.K. (2022). Effect of In Vitro Solvation Conditions on Inter- and Intramolecular
Assembly of Full-Length TDP-43. J Phys Chem B *126*, 4799-4813.
Erecinska, M., Nelson, D., Dagani, F., Deas, J., and Silver, I.A. (1993). Relations between intracellular
ions and energy metabolism under acidotic conditions: a study with nigericin in synaptosomes,
neurons, and C6 glioma cells. J Neurochem *61*, 1356-1368.
Farny, N.G., Kedersha, N.L., and Silver, P.A. (2009). Metazoan stress granule assembly is mediated by
P-eIF2alpha-dependent and -independent mechanisms. RNA *15*, 1814-1821.
Friese, M.A., Craner, M.J., Etzensperger, R., Vergo, S., Wemmie, J.A., Welsh, M.J., Vincent, A., and
Fugger, L. (2007). Acid-sensing ion channel-1 contributes to axonal degeneration in autoimmune
inflammation of the central nervous system. Nat Med *13*, 1483-1489.
Gao, J., Wang, L., Yan, T., Perry, G., and Wang, X. (2019). TDP-43 proteinopathy and mitochondrial
abnormalities in neurodegeneration. Mol Cell Neurosci *100*, 103396.
Glunde, K., Dussmann, H., Juretschke, H.P., and Leibfritz, D. (2002). Na(+)/H(+) exchange subtype 1
inhibition during extracellular acidification and hypoxia in glioma cells. J Neurochem *80*, 36-44.
Hardiman, O., Al-Chalabi, A., Chio, A., Corr, E.M., Logroscino, G., Robberecht, W., Shaw, P.J., Simmons,
Z., and van den Berg, L.H. (2017). Amyotrophic lateral sclerosis. Nat Rev Dis Primers *3*, 17085.
Harguindey, S., Reshkin, S.J., Orive, G., Arranz, J.L., and Anitua, E. (2007). Growth and trophic factors,
pH and the Na+/H+ exchanger in Alzheimer's disease, other neurodegenerative diseases and cancer:
new therapeutic possibilities and potential dangers. Curr Alzheimer Res *4*, 53-65.
Hergesheimer, R.C., Chami, A.A., de Assis, D.R., Vourc'h, P., Andres, C.R., Corcia, P., Lanznaster, D., and
Blasco, H. (2019). The debated toxic role of aggregated TDP-43 in amyotrophic lateral sclerosis: a
resolution in sight? Brain : a journal of neurology *142*, 1176-1194.
Jain, S., Wheeler, J.R., Walters, R.W., Agrawal, A., Barsic, A., and Parker, R. (2016). ATPase-Modulated
Stress Granules Contain a Diverse Proteome and Substructure. Cell *164*, 487-498.
Jin, X., Zhou, M., Chen, S., Li, D., Cao, X., and Liu, B. (2022). Effects of pH alterations on stress- and
aging-induced protein phase separation. Cell Mol Life Sci *79*, 380.
Johnson, B.S., Snead, D., Lee, J.J., McCaffery, J.M., Shorter, J., and Gitler, A.D. (2009). TDP-43 is
intrinsically aggregation-prone, and amyotrophic lateral sclerosis-linked mutations accelerate
aggregation and increase toxicity. J Biol Chem *284*, 20329-20339.
Khalfallah, Y., Kuta, R., Grasmuck, C., Prat, A., Durham, H.D., and Vande Velde, C. (2018). TDP-43
regulation of stress granule dynamics in neurodegenerative disease-relevant cell types. Sci Rep *8*, 7551.
Kuo, S.W., Jiang, M., and Heckman, C. (2014). Potential involvement of intracellular pH in a mouse
model of amyotrophic lateral sclerosis. Amyotrophic lateral sclerosis & frontotemporal degeneration
15, 151-153.

Lanznaster, D., Bourgeais, J., Bruno, C., Hergesheimer, R.C., Thepault, R.A., Vourc'h, P., Corcia, P.,
Andres, C.R., Herault, O., and Blasco, H. (2019). TDP-43-Mediated Toxicity in HEK293T Cells: A Fast and
Reproducible Protocol To Be Employed in the Search of New Therapeutic Options against Amyotrophic
Lateral Sclerosis. Cells *9*.

Li, H.R., Chen, T.C., Hsiao, C.L., Shi, L., Chou, C.Y., and Huang, J.R. (2018). The physical forces mediating
self-association and phase-separation in the C-terminal domain of TDP-43. Biochim Biophys Acta
Proteins Proteom *1866*, 214-223.

Mompean, M., Chakrabartty, A., Buratti, E., and Laurents, D.V. (2016a). Electrostatic Repulsion
Governs TDP-43 C-terminal Domain Aggregation. PLoS Biol *14*, e1002447.

Mompean, M., Romano, V., Pantoja-Uceda, D., Stuani, C., Baralle, F.E., Buratti, E., and Laurents, D.V.
(2016b). The TDP-43 N-terminal domain structure at high resolution. FEBS J *283*, 1242-1260.

Morihata, H., Nakamura, F., Tsutada, T., and Kuno, M. (2000). Potentiation of a voltage-gated proton
current in acidosis-induced swelling of rat microglia. The Journal of neuroscience : the official journal
of the Society for Neuroscience *20*, 7220-7227.

Neumann, M., Sampathu, D.M., Kwong, L.K., Truax, A.C., Micsenyi, M.C., Chou, T.T., Bruce, J., Schuck,
T., Grossman, M., Clark, C.M., et al. (2006). Ubiquitinated TDP-43 in frontotemporal lobar degeneration
and amyotrophic lateral sclerosis. Science *314*, 130-133.

Patni, D., and Jha, S.K. (2021). Protonation-Deprotonation Switch Controls the Amyloid-like Misfolding
of Nucleic-Acid-Binding Domains of TDP-43. J Phys Chem B *125*, 8383-8394.

Pillai, M., and Jha, S.K. (2019). The Folding and Aggregation Energy Landscapes of Tethered RRM
Domains of Human TDP-43 Are Coupled via a Metastable Molten Globule-like Oligomer. Biochemistry
58, 608-620.

Pillai, M., and Jha, S.K. (2020). Early Metastable Assembly during the Stress-Induced Formation of
Worm-like Amyloid Fibrils of Nucleic Acid Binding Domains of TDP-43. Biochemistry *59*, 315-328.

Putnam, R.W. (2001). Intracellular pH regulation of neurons in chemosensitive and nonchemosensitive
areas of brain slices. Respir Physiol *129*, 37-56.

Rabouille, C., and Alberti, S. (2017). Cell adaptation upon stress: the emerging role of membrane-less
compartments. Curr Opin Cell Biol *47*, 34-42.

Riancho, J., Castanedo-Vazquez, D., Gil-Bea, F., Tapia, O., Arozamena, J., Duran-Vian, C., Sedano, M.J.,
Berciano, M.T., Lopez de Munain, A., and Lafarga, M. (2020). ALS-derived fibroblasts exhibit reduced
proliferation rate, cytoplasmic TDP-43 aggregation and a higher susceptibility to DNA damage. Journal
of neurology *267*, 1291-1299.

Riback, J.A., Katanski, C.D., Kear-Scott, J.L., Pilipenko, E.V., Rojek, A.E., Sosnick, T.R., and Drummond,
D.A. (2017). Stress-Triggered Phase Separation Is an Adaptive, Evolutionarily Tuned Response. Cell *168*,
1028-1040 e1019.

Sabatelli, M., Zollino, M., Conte, A., Del Grande, A., Marangi, G., Lucchini, M., Mirabella, M., Romano,
A., Piacentini, R., Bisogni, G., et al. (2015). Primary fibroblasts cultures reveal TDP-43 abnormalities in
amyotrophic lateral sclerosis patients with and without SOD1 mutations. Neurobiol Aging *36*, 2005
e2005-2005 e2013.

Swietach, P., Hulikova, A., Patiar, S., Vaughan-Jones, R.D., and Harris, A.L. (2012). Importance of
intracellular pH in determining the uptake and efficacy of the weakly basic chemotherapeutic drug,
doxorubicin. PloS one *7*, e35949.

Tolkovsky, A.M., and Richards, C.D. (1987). Na+/H+ exchange is the major mechanism of pH regulation
in cultured sympathetic neurons: measurements in single cell bodies and neurites using a fluorescent
pH indicator. Neuroscience *22*, 1093-1102.

Tsoi, P.S., Choi, K.J., Leonard, P.G., Sizovs, A., Moosa, M.M., MacKenzie, K.R., Ferreon, J.C., and Ferreon,
A.C.M. (2017). The N-Terminal Domain of ALS-Linked TDP-43 Assembles without Misfolding. Angew
Chem Int Ed Engl *56*, 12590-12593.

van der Laan, A.M., van Gemert, A.M., Dirks, R.W., Noordermeer, J.N., Fradkin, L.G., Tanke, H.J., and
Jost, C.R. (2012). mRNA cycles through hypoxia-induced stress granules in live Drosophila embryonic
muscles. Int J Dev Biol *56*, 701-709.

Vergo, S., Craner, M.J., Etzensperger, R., Attfield, K., Friese, M.A., Newcombe, J., Esiri, M., and Fugger,
L. (2011). Acid-sensing ion channel 1 is involved in both axonal injury and demyelination in multiple
sclerosis and its animal model. Brain : a journal of neurology *134*, 571-584.

Wang, P., Deng, J., Dong, J., Liu, J., Bigio, E.H., Mesulam, M., Wang, T., Sun, L., Wang, L., Lee, A.Y., et al.
(2019). TDP-43 induces mitochondrial damage and activates the mitochondrial unfolded protein
response. PLoS Genet *15*, e1007947.

Wang, W., Li, L., Lin, W.L., Dickson, D.W., Petrucelli, L., Zhang, T., and Wang, X. (2013). The ALS disease-
associated mutant TDP-43 impairs mitochondrial dynamics and function in motor neurons. Hum Mol
Genet *22*, 4706-4719.

Wei, Y., Lim, L., Wang, L., and Song, J. (2016). Inter-domain interactions of TDP-43 as decoded by NMR.
Biochem Biophys Res Commun *473*, 614-619.

Yao, H., and Haddad, G.G. (2004). Calcium and pH homeostasis in neurons during hypoxia and
ischemia. Cell Calcium *36*, 247-255.

Figures.

Figure 1. Lower pH in cellular models of ALS. HEK293T cells (upper panel) overexpressing increasing
amounts of TDP-43 present more fluorescence due to BCECF than control (mock-transfected) cells.
Images represent five independent experiments. N= 5; *p=0,0127; Kruskal-Wallis test. Dermal
fibroblasts (lower panel) obtained from ALS patients also present increased BCECF fluorescence
compared to fibroblasts from a control subject. Images represent two independent experiments. N= 2;
**p=0,0007; Mann Withney test.

383
384 **Figure 2**. TDP43 localization in function of pH and incubation time : 1h (A,C,E,G,) or 2h (B,D,F,H). n
385 = 4 ; *P = 0,04 (two-way ANOVA); I: percentage of TDP43 aggregation after 1h incubation with
386 different pH solutions or (J) after 2h incubation. n = 4, *P = 0,05 (one-way ANOVA followed by a
387 Kruskal-Wallis test).