MetaPredict: A machine learning-based tool for predicting metabolic modules in incomplete

bacterial genomes

D. Geller-McGrath[1], Kishori M. Konwar[2], V.P. Edgcomb[3], M. Pachiadaki[1], J. W. Roddy[4],

T. J. Wheeler[4], J. E. McDermott[5,6]

[1]Biology Department, Woods Hole Oceanographic Institution, Woods Hole, MA 02543

[2]Luit Consulting, Revere, MA 02151

[3]Department of Marine Geology and Geophysics, Woods Hole Oceanographic Institution,

Woods Hole, MA 02543

[4]Department of Pharmacy Practice and Science, University of Arizona, Tucson, AZ 85721

[5]Computational Sciences Division, Pacific Northwest National Laboratory, Richland,

Washington, 99352

[6]Department of Molecular Microbiology and Immunology, Oregon Health & Science

University, Portland, Oregon 97239

Abstract

The reconstruction of complete microbial metabolic pathways using 'omics data from environmental samples remains challenging. Computational pipelines for pathway reconstruction that utilize machine learning methods to predict the presence or absence of KEGG modules in incomplete genomes are lacking. Here, we present MetaPredict, a software tool that incorporates machine learning models to predict the presence of complete KEGG modules within bacterial genomic datasets. Using gene annotation data and information from KEGG module databases, MetaPredict employs neural network and XGBoost stacked ensemble models to reconstruct and predict the presence of KEGG modules in a genome. MetaPredict can be used as a command line tool or as an R package, and both options are designed to be run locally or on a compute cluster. In our benchmarks, MetaPredict makes robust predictions of KEGG module presence within highly incomplete genomes.

Introduction

Microbial communities play a key role in all major biogeochemical cycles on Earth. Accurate and complete identification of microbial metabolic pathways within genomic data is crucial to understanding the function of microorganisms. This identification of pathways within genomic data, and assessment of their expression, provides important insight into their influence on the chemistry of their environment and their mediation of interactions with other organisms.

In recent decades, the scientific community has significantly advanced its capability to gather and sequence genomes from microorganisms. Key steps in the process of working with isolated genomes, single-amplified genomes (SAGs), or metagenome assembled genomes (MAGs), are identifying genes coding for enzymes that catalyze metabolic reactions and inferring the metabolic potential of the associated organism from these data. These analyses involve comparing protein-coding sequences with homologous sequences from reference metabolic pathway databases including KEGG (Kanehisa et al. 2000) and MetaCyc (Caspi et al. 2018). Environmental genomes that are recovered from high-throughput sequencing samples vary in their degree of completeness due to numerous factors including limited coverage of low-

abundance microbes, composition-based coverage biases, insertion-deletion errors, and
substitution errors (Browne et al. 2020). Enzymes encoded in genomes are also missed due to
limitations in protein annotation methods, that is, undiscovered protein families may be
undetected by traditional homology-based methods. This can limit the ability to determine the
extent to which these organisms (or communities) can catalyze metabolic reactions and form
pathways.

Sequencing biases, novel protein families, and incomplete gene and protein annotation
databases lead to missing, ambiguous, or inaccurate gene annotations that create incomplete
metabolic networks in recovered environmental genomes. Existing algorithms for identifying
incomplete reactions ("filling gaps"), in metabolic networks largely fall into two categories of
approaches: those based on network topology (such as the method utilized by Gapseq
(Zimmerman et al. 2021)), and those that utilize pre-defined pathway cutoffs (such as those used
by METABOLIC (Zhou et al. 2022)). Network topology and pathway gene presence/absence
cutoffs, however, can lead to underestimation of pathways that are present, particularly in highly
incomplete genomes. Parsimony-based algorithms such as MinPath detect gaps in a metabolic
network and identify the minimum number of modifications to the network that can be made to
activate those reactions (Ye et al. 2009); its conservative approach, however, can lead to
underestimation of the metabolic pathways present in a sample. Other modern tools, such as
DRAM (Shaffer et al. 2020), provide annotations for metagenomic sequences but do not closely
tie these to metabolic pathways. Flux-balance analysis (e.g. Escher-FBA [Rowe et al. 2018])
utilizes genome-scale metabolic models of organisms and requires experimental growth data for
model parameterization; it is not easily applied to incomplete genome data, and the additional
required experimental measurements may prohibit application in many use cases.

An emerging set of methods utilize machine learning models to a related problem of
classifying microbial organisms' niches based on their genomic features. One such example is a
tool called Traitar, which utilizes Support Vector Machines (SVMs) to predict lifestyle and
pathogenic traits in prokaryotes based on gene family abundance profiles (Weiman et al. 2016).
Other recent approaches have used machine learning approaches to train models using eukaryote
MAG and transcriptome data to classify trophic mode (autotroph, mixotroph, or heterotroph)
based on gene family abundance profiles (Lambert et al. 2021, Alexander et al. 2021). To our
knowledge, there are no existing tools that predict the presence/absence of KEGG metabolic

modules (with gap-filling options) via machine learning models trained on gene features of high-
quality genomes.
Here, we present "MetaPredict", an open-source tool for metabolic pathway prediction based on
a XGBoost (Chen and Guestrin, 2016) and neural network stacked ensemble classification
framework. MetaPredict addresses critical deficiencies in existing metabolic pathway
reconstruction tools that limit the utility and predictive power of 'omics data: it connects
manually curated, current knowledge of metabolic pathways from the KEGG database with

Figure 1. Overview of the MetaPredict pipeline. Input genome annotations are read into MetaPredict as a data object. The data are scanned for present KEGG modules and are formatted into a feature matrix. The feature matrix is then used to make predictions for all incomplete modules (or modules specified by the user). A summary and detailed reconstruction and prediction objects, along with gap-filling options are returned in a list as the final output.

machine learning methods to reconstruct and predict the presence or absence of KEGG

metabolic modules within genomic datasets including bacterial isolate genomes, MAGs, and

SAGs.

The models underlying MetaPredict contain metabolic reaction and pathway information

from taxonomically diverse bacterial isolate genomes and MAGs from the NCBI RefSeq

(O'Leary et al. 2016) and Genome Taxonomy Database (GTDB, Parks et al. 2021) databases.

The set of metabolic modules from the KEGG database is the basis of the tool's metabolic

module reconstruction and prediction. The KEGG database contains metabolic pathway

information for thousands of prokaryotic species and strains. KEGG modules are functional units

of metabolic pathways composed of sets of ordered reaction steps. Examples include carbon

fixation pathways, nitrification, biosynthesis of vitamins, and transporters or two-component

systems (see Supplementary Table 1 for a description of the distribution of modules covered by

MetaPredict). MetaPredict is designed to run on the command line locally or on a computing

cluster and is available as an R package from GitHub (https://github.com/d-

mcgrath/MetaPredict).

A detailed overview of the MetaPredict pipeline is provided in Figure 1. The tool accepts

as input gene annotations (of one or more genomes) with associated KEGG ortholog (KO) gene

identifiers and first reconstructs both complete and incomplete KEGG metabolic modules, then

predicts the presence or absence of all incomplete modules (or modules specified by the user).

Annotations can be used as input from tools such as KofamScan (Aramaki et al. 2020), DRAM,

blastKOALA (Kanehisa et al. 2016), ghostKOALA (Kanehisa et al. 2016), or a custom list of

KO identifier gene annotations. MetaPredict utilizes a stacked ensemble binary classification

approach to predict the presence or absence of complete KEGG modules in incomplete genomes

based on the KO gene identifiers of input genomes. MetaPredict classification models produced

accurate results on held-out test genome annotation datasets even when the data were highly

incomplete. A set of 293 neural network/XGBoost stacked ensemble models, one for each of 293

KEGG modules, made predictions with a high degree of precision on all test datasets and with

high recall on genomes with an estimated completeness as low as 35%. False positive predictions

were rare in all tests, while false negatives increased when predictions were made with highly

incomplete gene annotation information, as would be expected. An additional set of 183 neural

network models (not stacked ensembles) trained to classify the presence of KEGG modules

rarely present in the training data made predictions with robust precision and recall on genomes
with a completeness estimate of at least 60%, exhibiting very similar precision and recall
characteristics to the stacked ensembles. We believe that MetaPredict will be a valuable tool to
enable studies of metabolic potential in environmental microbiome studies as well as synthetic
biology effort.
137
138

Results and Discussion

MetaPredict's performance metrics on held-out test datasets suggest its models predict
with high fidelity when at least 40% of gene annotations are recovered from a reconstructed
genome (Figure 2). The efficacy of MetaPredict models was assessed using incomplete gene
annotation data simulated from whole genomes, as well as from genomes reconstructed from
reads that had been randomly downsampled. We further benchmarked MetaPredict against
custom presence/absence classification rules, and existing gap-filling tools METABOLIC and
Gapseq.

Benchmarking MetaPredict and alternative strategies on downsampled NCBI RefSeq and GTDB
data

We compared the performance of MetaPredict's stacked ensemble models to two classes
of competitor classifiers: naive rule-based classifiers and various other machine learning model
architectures. The evaluation was performed on test datasets comprised of isolate and
metagenome-assembled genomes from the GTDB and NCBI databases (see Methods). When
evaluating with the same sets of randomly downsampled gene annotations, we found that each
competing method performed worse than MetaPredict (Figure 2). We assessed two naïve
classification methods. First, we devised a classification rule based on the completeness of a
KEGG module relative to the number of genes retained after downsampling: if, in a
downsampled genome, the number of genes involved in a KEGG module are present in at least
the same proportion of all genes retained, the KEGG module is classified as 'present,' otherwise
it is labeled 'absent'. For example, if 50% of gene annotations were removed from a genome

during downsampling, then any KEGG module for which 50% of its associated genes are
retained would be reported as 'present'. The results of this naive approach (Figure 2) show that
the relative completeness of a KEGG module alone is not a robust classification strategy. The
second classification rule that we tested was: for all gene annotation sets in the dataset, if any
genes were present in an annotation set that were unique to a KEGG module (relative to other
modules) then the module was classified as 'present', otherwise it was labelled 'absent'. The
results of this naive approach (Figure 2) suggest that the presence of rare protein annotations or
genes unique to a certain KEGG module is not always a strong indicator of the presence of a
module in a genome. Ultimately, the performance of these naive classifiers indicate that
MetaPredict's models have the advantage of incorporating information from genes outside of
KEGG modules. We additionally compared the performance of various machine learning model
architectures. Of these, the XGBoost (no ensemble) and neural network (no ensemble)
architectures were the next-best performing models (Figure 2).

Figure 2. Comparison of performance metrics of MetaPredict's 293 stacked ensemble binary classification models to
next-best performing XBoost and neural network machine learning models as well as two naïve classification rules.
Downsampled gene annotations of high-quality genomes used in held-out test sets are from NCBI RefSeq and

GTDB databases. Each boxplot displays the distribution of model performance metrics for predictions on randomly
sampled versions of the gene annotation test sets in downsampling increments of 10% (90% down to 10%, from
right to left). The binary classifier performances are based on the classification of the presence or absence of KEGG
modules in the complete versions of the gene annotations that were downsampled for model testing.

MetaPredict's ensemble strategy produced the best observed performance. Mean F1 score of the
models was at least 0.90 when predicting on test datasets in which 30-90% of gene annotations
had been retained. MetaPredict rarely made false positive predictions based on data from highly
incomplete gene annotation sets; the average precision of the models was consistently above 0.94
for all held-out test sets. MetaPredict also did not misclassify most negative class observations;
the 293 models had an overall average specificity above 0.9 on all test datasets. The quality of
MetaPredict's negative class predictions was robust on test datasets containing at least 40% of
the complete gene annotation data. The mean recall decreased incrementally with decreased gene
annotation data and was below 0.86 on only on test sets containing 30% or less of the complete
gene annotation data. The drop in recall and negative predictive value on highly downsampled
test data was indicative of an increase in false negative predictions when MetaPredict models
were assessing KEGG module presence/absence based on highly incomplete information.

Benchmarking MetaPredict against GEM repository MAGs

MetaPredict was further tested on gene annotations from a set of 100 high-quality
metagenome-assembled genomes from the GEM genome repository (Figure 3.). The MAGs
selected from the repository had an estimated completeness of 100 and estimated contamination
of 0, and MIMAG quality score of "High Quality". The genomes belonged to 15 taxonomic
classes and were recovered from 11 different environments, primarily from human-associated
and built environment metagenomes (see Supplementary Fig. 4 for GEM genome taxonomic
distributions and environmental sources). We created a list of 9 GEM datasets by randomly
downsampling the data to retain 10% to 90% of gene annotations (in 10% increments) as in the
previous section. MetaPredict classified the presence/absence of KEGG modules in each MAG.
Overall results were comparable to MetaPredict's performance on the benchmark based on
GTDB and NCBI databases. The models excelled at predicting the presence or absence of

KEGG modules in genomes when at least 50% of gene annotations were randomly retained.

Predictions were less reliable when 40% or less gene annotation data was retained.

Figure 3. Boxplots of performance metrics of 293 MetaPredict models on high-quality bacterial GEM MAGs (*n* =

100). Model performance metrics are for predictions on downsampled versions of GEM genome gene annotations

in decreasing increments of 10% (retaining 10-90% of the annotations in each test set). MetaPredict's stacked

ensemble models were benchmarked against XGBoost and neural network model architectures.

 Benchmarking MetaPredict against existing tools on a deataset with downsampled reads

In addition to model assessments made through downsampling protein annotations, we

evaluated a second set of held-out test set genomes (*n* = 50) that we downsampled at the

sequence read level. We downsampled genome sequence reads to assemble and annotate

fragmented versions of these test set genomes from NCBI and GTDB. MetaPredict's

performance on this test set closely resembled protein annotation random sampling results

(Figure 4a, Figure 4b).

Figure 4. Performance metrics boxplots of 293 stacked ensemble classification models. Downsampled sequence

reads of high-quality genomes used as a second held-out test set are from NCBI RefSeq and GTDB databases. **Panel**

a: Boxplots display the distribution of model performance metrics for predictions of KEGG module

presence/absence on simulated incomplete genomes downsampled at the sequence read level by MetaPredict models

and METABOLIC. Downsampling increments were chosen based on average estimated completeness of the test set

genomes at each increment to reflect a range of estimated completeness thresholds. **Panel b:** Average estimated

genome completeness distributions of test set genomes that were downsampled at the sequence read level and then

assembled and annotated.

MetaPredict had an average F1 score for all 293 models of 0.92 on the second held-out test sets

that had an average estimated genome completeness of 35%. The similarity of these results to the

gene annotation downsampling approach validates the latter approach that was used more

broadly to assess MetaPredict. We compared MetaPredict's performance to module

presence/absence predictions made on the same data with METABOLIC, which is a command

line tool that performs gene annotations and estimates the completeness of KEGG Modules in

individual genomes and prokaryotic microbial communities (Figure 4a). We found that

MetaPredict's models made much more sensitive predictions on test set genomes. We also

benchmarked MetaPredict against Gapseq based on both tools' predictions of the presence or

absence of a particular KEGG pathway in test set genomes (Figure 5a). MetaPredict

outperformed predictions made by Gapseq, particularly on genomes that had had a high

proportion of reads removed prior to assembly.

Figure 5. Performance metrics boxplots of MetaPredict and Gapseq predictions for KEGG pathway map00290

(Valine, leucine, and isoleucine biosynthesis) which contains KEGG modules M00019, M00432, M00535, and

M00570. For MetaPredict predictions, the whole KEGG pathway was considered present if the aforementioned

KEGG modules were all present. Downsampled sequence reads of high-quality genomes used as a second held-out

test set are from NCBI RefSeq and GTDB databases. **Panel a:** Line segments display model performance metrics

for MetaPredict and Gapseq predictions of KEGG pathway map00290 presence/absence on simulated incomplete

genomes downsampled at the sequence read level. Downsampling increments were chosen based on average

estimated completeness of the test set genomes at each increment to reflect a range of estimated completeness

thresholds. **Panel b:** Average estimated genome completeness distributions of test set genomes that were

downsampled at the sequence read level and then assembled and annotated.

We fitted and tested an additional 183 neural network models for rare KEGG modules

present in less than 5% of the training data (Supplementary Figs. 2b, 5-6). The neural network

models were not incorporated into ensembles with XGBoost models due to the severe class

imbalance for these KEGG modules. These models performed optimally on genomes that were

estimated to be at least 60% complete.

Conclusion

MetaPredict is an open-source tool that can be used to characterize the functional

potential of one or more sample genomes by detecting complete KEGG modules and predicting

the presence or absence of those that are incomplete or missing. The tool accepts sets of gene

annotations of individual genomes in KO gene identifier format as input. This type of annotation

format can be acquired by annotating a genome of interest using KEGG-based annotation tools

such as KofamScan (Aramaki et al., 2020), blastKOALA (Kanehisa et al. 2016), or

ghostKOALA (Kanehisa et al. 2016). MetaPredict also provides gene gap-filling options by

suggesting putative KO gene annotations that could fill in missing gaps in predicted modules.

MetaPredict further validates the use of gene copy frequency within a genome as a

feature for bacterial metabolic function predictions. The performance metrics of MetaPredict on

NCBI/GTDB and GEM test datasets validated the use of stacked ensemble models to predict the

presence/absence of KEGG metabolic modules with high fidelity on sparse to near-complete

bacterial genomes. MetaPredict's binary classification models consistently made predictions with

high precision, recall and specificity on simulated and real genomes using gene annotation and

sequence read downsampling methods. The predictive power of the stacked ensemble models
was most limited when predicting on 10%-30% of protein annotations, and when the mean
estimated completeness of reconstructed genomes from downsampled reads was below 35%. We
suggest that optimal performance with MetaPredict can be achieved when at least 40% of genes
have been recovered in an input bacterial gene annotation dataset.

MetaPredict will facilitate more complete and accurate reconstruction of the metabolic
potential encoded within bacterial genomes from a diverse array of environments and will
enhance the ability to infer what metabolisms they are capable of, and/or how they may respond
to perturbations. MetaPredict connects the field of machine learning with the growing
community of environmental microbiologists using genomic sequencing techniques and will help
transform and improve the way they work with environmental genomic datasets.

Materials and Methods

Filtering genome database metadata, downloading high-quality genomes, and gene annotation

The NCBI RefSeq (Release 205) database metadata file was downloaded and filtered to
retain only the information for all bacterial genomes classified as "Complete genome", which are
defined on the NCBI assembly help webpage "all chromosomes are gapless and have no runs of
10 or more ambiguous bases (Ns), there are no unplaced or unlocalized scaffolds, and all the
expected chromosomes are present (i.e. the assembly is not noted as having partial genome
representation). Plasmids and organelles may or may not be included in the assembly but if
present, then the sequences are gapless." This resulted in 17,491 complete NCBI genomes.

The GTDB bacterial metadata file for release 95 was downloaded and filtered to keep the
information for all genomes with an estimated completeness greater than 99, an estimated
contamination of 0, and a MIMAG (Bowers et al., 2017) quality score of "High Quality". A total
of 30,760 non-redundant bacterial genomes from the two database metadata files were
downloaded using the ncbi-genome-download command line tool (Blin, K.). The RefSeq
genomes (17,491 total) were downloaded from the RefSeq FTP server, and the GTDB genomes
(13,269 total) were downloaded from the GenBank FTP server (Supplementary Figure 1). Genes

were called using Prodigal (Hyatt et al., 2010), and the KofamScan command line tool (Aramaki
et al. 2020) was used to generate gene annotations (in KO gene identifier format) for all of the
genomes using the KOfam set of Hidden Markov Models (HMMs) available for download from
the KEGG database (Kanehisa et al. 2002).

Formatting gene annotation data, fitting KEGG module classification models

The gene copy number data of the downloaded genomes was formatted in a matrix
containing KO gene identifier counts in columns and genomes in rows. The label of each model
was the presence/absence of a KEGG module, as was determined using the KEGG modules
downloaded from the KEGG database. For the module to be categorized as present, every step of
the module had to be present in a genome, otherwise it was designated as absent. 293 KEGG
modules were individually modelled using the gene annotation data of the genomes consolidated
from the NCBI and GTDB databases. All 293 models were constructed for KEGG modules for
which at least 5% of the training genomes contained the module genes. The constructed models
classify the presence or absence of complete KEGG modules based on the gene annotations of a
genome. Training and test sets contained both complete and incomplete gene annotations of
bacterial genomes from a diverse array of phyla (Supplementary Fig. 1). The incomplete
annotations used in training and testing of MetaPredict's models were constructed from complete
genome annotations that were randomly downsampled to retain 10-90% of the total gene content.
The training datasets had a size of 24,517 observations, for which one observation was a vector
in which each element was a count of the frequency that an individual KO identifier was
annotated in a genome. The test datasets each contained 6,129 observations. The percent of
observations with a positive class (a complete KEGG module 'present' in the gene annotations)
in the training and test datasets varied, with a mean of 28.24% (Supplementary Fig. 2a).
A neural network and XGBoost stacked ensemble classification approach was chosen to
model the relationship between whole genome KO gene identifier annotation data and the
presence of metabolic modules. Model stacking is an approach in which several independently-
trained models (often comprised of multiple machine learning architectures) are applied to an
input, and their output is passed to another machine learning model (a 'meta-learner'), which
produces a final classification. The stacking approach in MetaPredict employs an L1-regularized

linear model as a meta-learner to learn how to best combine predictions from the ensemble of
models. Training data was divided into a subset used to train ensemble members (24,517
observations) and the remaining entries (so called 'out-of-fold' predictions) held-out for meta-
learner training. The same training data was used to train each of the ensemble models, and the
meta-learner was trained on the out-of-fold predictions made by candidate ensemble members.
L1-regularization was utilized by the meta-learner to pick the final number of ensemble members
from the pool of candidates. For each model trained, the meta-learner then retained ensemble
members with the best performance on out-of-fold data and assigned weights to the predictions
made by each ensemble member for future predictions on unseen data. Features used in each
training dataset for classification were the gene copy number for all genes that were assigned KO
gene identifiers (for which one observation was a vector in which each element was a count of
the times that an individual KO gene identifier was annotated in a genome). Information gain,
which is a measure of the statistical dependence between two variables, was utilized for feature
selection to decrease model complexity and avoid over-fitting using the recipeselectors R
package (Steven Pawley, 2022). This method iteratively calculates the mutual information
between the dependent variable (KEGG module presence/absence) and each independent
variable (the KO gene identifiers) and measures the reduction in uncertainty for the dependent
variable, given the values of the independent variable. The mutual information, $I(X\,;Y)$, between
two random variables X and Y is: $I(X\,;Y) = H(X) - H(X|Y)$, where $H(X)$ is the entropy for X,
and $H(X|Y)$ is the conditional entropy for X given Y. The number of best-scoring features
retained in each model's train/test datasets were chosen for each model through cross validation.
XGBoost models and feed-forward neural networks with a single hidden layer were used as the
machine learning architectures in MetaPredict's stacked ensemble models.

The full set of available genome annotations was used to form each model's train and test
datasets. Stratified random sampling, in which the genome dataset observations were divided
into strata and sampled independently, was used to generate 80% train/20% test dataset splits
that each contained data observations with preserved proportions of positive ('KEGG module
present') and negative ('KEGG module absent') classes that were present in the genome dataset
(see boxplot of the distribution of module presence/absence classes in Supplementary Figure 2a).
The Tidymodels R package (Kuhn M., Wickham H., 2020) was used to fit 293 neural
network/XGBoost stacked ensemble models each corresponding to a unique KEGG module.

The 293 MetaPredict stacked ensemble models were fit using the Stacks R package
(https://github.com/tidymodels/stacks). First, potential XGBoost and neural network parameter
values were assessed using 10-fold nested cross validation to maximize the Receiver Operating
Characteristic (ROC). Grid search was utilized for the XGBoost parameters *mtry*, *min n*, *tree*
depth, *learn rate*, *loss reduction*, and *sample size*. After locking choices for these parameters, a
separate grid of values for neural etwork parameters *hidden units*, *penalty*, *dropout* and
activation were then assessed. The Stacks package functions were then utilized to fit Least
Absolute Shrinkage and Selection Operator (LASSO) meta-learner models to out-of-fold
predictions made by the XGBoost and neural network models during cross validation via grid
search. This was done over a range of λ values 1×10^{-6} to 1×10^{-1}, and the λ that maximized the
ROC was chosen for each ensemble member. The l_1-penalization of the LASSO model
facilitated the selection of final model ensemble members from the pool of candidate members
for each stacked model. For 183 KEGG modules rarely present in the training data (< 5%
genomes contained the full module), neural network models were chosen based on their
outperformance of XGBoost models on test datasets instead due to severe class imbalance and
inability to fit stacked ensemble models to these KEGG modules.

We assessed and benchmarked MetaPredict's models against two naïve classification
methods. First, we hypothesized that we could predict the presence of a KEGG module if, after
downsampling test sets of gene annotations, the proportion of module genes present in the
dataset was greater than or equal to the percentage of annotations retained in the dataset. If the
proportion of genes involved in a KEGG module were present in a dataset observation at least
equivalently to the proportion of gene annotations retained after downsampling, the module was
classified as 'present', otherwise it was classified as 'absent'. The second naïve classification
rule was: for all gene annotation sets in the dataset, if any genes were present in an annotation set
that were unique to a KEGG module (relative to all other KEGG modules) then the module was
classified as 'present', otherwise it was labelled 'absent'. We additionally benchmarked
MetaPredict's stacked ensemble models against several other machine learning model types
including standalone neural network and XGBoost models.

Evaluating models on test data, including test data randomly downsampled to simulate varying

degrees of genome incompleteness

Each of MetaPredict's models was validated on a held-out test set with 6,129 observations and

the performance metrics were extracted using the Tidymodels R package. The performance

metrics used in evaluating the models were precision, recall, specificity, F1 score, positive

predictive value, and negative predictive value (Table 1).

Metric	Definition
Precision	true positive/(true positive + false positive)
Recall	true positive/(true positive + false negative)
Specificity	true negative/(true negative + false positive)
F1 score	$2 \times ((\text{precision} \times \text{recall})/(\text{precision} + \text{recall}))$
Positive predictive value	$\text{recall} \times \text{prevalence} / ((\text{recall} \times \text{prevalence}) + ((1 - \text{specificity}) \times (1 - \text{prevalence})))$
Negative predictive value	$\text{specificity} \times (1 - \text{prevalence}) / ((1 - \text{recall}) \times \text{prevalence}) + (\text{specificity} \times (1 - \text{prevalence}))$
Prevalence*	(true positive + false negative) / (true positive + false positive + true negative + false negative)

Table 1. Definitions of machine learning model performance metrics used to assess MetaPredict models.
*Prevalence is defined due to its use in the definitions of negative and positive predictive value.

The genome annotations from each test set were then randomly downsampled to simulate

recovered gene annotations from incomplete genomes. 10% to 90% of genes from each

annotation set were randomly retained (in increments of 10%) and used as input for MetaPredict

predictions of KEGG module presence/absence. The model performance metrics in each

simulation were extracted using the Tidymodels package.

Testing models with a set of high-quality metagenome-assembled genomes from the Genomes

from Earth's Microbiomes online repository

MetaPredict was further validated on another test set of genome annotations extracted from the

Genomes from Earth's Microbiomes (GEM; Nayfach et al. 2021) repository of MAGs. The

GEM metadata file was downloaded from the repository (Nayfach et al. 2021) and filtered to

retain a random sample of 100 MAGs with a CheckM (Parks et al. 2015) estimated completeness

of 100, an estimated contamination of 0, and a MIMAG quality score of "High Quality". The

method for this assessment was the same as was described above for testing MetaPredict model

performances on the held-out test data.

Evaluating models on test data downsampled at the read level

A second held-out set of genomes ($n = 50$) was downloaded from NCBI/GTDB databases using

the SRA Toolkit (SRA Toolkit Development Team) and SRA explorer (Phil Ewels). The raw

sequencing reads were filtered using fastp (Chen et al. 2018), and the quality-filtered reads were

randomly downsampled using seqtk (Li, H. 2012). Downsampled reads were assembled into

genomes using the SPAdes assembler (Bankevich et al., 2012), genes were called with Prodigal

and then annotated using KofamScan. MetaPredict's stacked ensemble models were then used to

predict the presence or absence of all 293 KEGG modules in each genome and predictions were

then cross-referenced with their known presence/absence based on the unmodified test dataset. In

addition to simple approaches described above, the METABOLIC (Zhou et al. 2022) and Gapseq

(Zimmerman et al. 2021) tools were evaluated on the same benchmark dataset. Gapseq and

MetaPredict were compared by evaluating predictions for the presence or absence of the KEGG

pathway map00290 (Valine, leucine, and isoleucine biosynthesis) which contains KEGG

modules M00019, M00432, M00535, and M00570. For MetaPredict predictions, the whole

KEGG pathway was considered present if the aforementioned KEGG modules were all predicted

as present, otherwise it was classified as absent.

Gapfilling for incomplete modules predicted as present

MetaPredict provides enzyme gap-filling options for KEGG modules predicted as present by
suggesting putative KO gene annotations missing from an input genome's gene annotations that
could fill in missing gaps in predicted modules.
463
464

Acknowledgements

D. Geller-McGrath acknowledges funding from DOE SCGR Fellowship for the 2020 Solicitation
2 in Computational Biology and Bioinformatics. We would like to thank K. Konwar (Facebook,
MIT, Broad Institute) for his guidance with conceptualization of this bioinformatics tool, and A.
Solow (WHOI) for initial helpful initial discussion about statistical approaches. JEM, JWR and
TJW were supported by the Department of Energy (DOE) Office of Biological and
Environmental Research (BER) through the "Machine-Learning Approaches for Integrating
Multi-Omics Data to Expand Microbiome Annotation" project. PNNL is operated for the DOE
by Battelle Memorial Institute under Contract DE-AC05-76RL01830.

Corresponding author

Correspondence to David Geller-McGrath and Jason McDermott.

Ethics declarations
The authors declare no competing interests.
480
481

482   **References**
483

1. Kanehisa, M. *et al.* KEGG: kyoto encyclopedia of genes and genomes. *Nucleic acids*
research 28, 27–30 (2000). https://www.genome.jp/tools/kofamkoala/, Kanehisa et al.
2002
2. Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes. *Nucleic*
acids research 46, D633–D639 (2018).
3. Browne, Patrick Denis, et al. "GC bias affects genomic and metagenomic reconstructions,
underrepresenting GC-poor organisms." *GigaScience* 9.2 (2020): giaa008.

4. Zimmermann, Johannes, Christoph Kaleta, and Silvio Waschina. "gapseq: informed
prediction of bacterial metabolic pathways and reconstruction of accurate metabolic
models." *Genome biology* 22.1 (2021): 1-35.

5. Zhou, Zhichao, et al. "METABOLIC: high-throughput profiling of microbial genomes for
functional traits, metabolism, biogeochemistry, and community-scale functional
networks." *Microbiome* 10.1 (2022): 1-22.

6. Ye, Y. *et al.* A parsimony approach to biological pathway reconstruction/inference for
genomes and metagenomes. *PLoS Comput Biol* 5, e1000465 (2009).

7. Shaffer, M. *et al.* DRAM for distilling microbial metabolism to automate the curation of
microbiome function. *Nucleic Acids Research* 48, 8883 8900 (2020).

8. Rowe, Elliot, Bernhard O. Palsson, and Zachary A. King. "Escher-FBA: a web
application for interactive flux balance analysis." BMC systems biology 12.1 (2018): 1-7.

9. Weimann, Aaron, et al. "From genomes to phenotypes: Traitar, the microbial trait
analyzer." *MSystems* 1.6 (2016): e00101-16.

10. Lambert, Bennett S., et al. "The dynamic trophic architecture of open-ocean protist
communities revealed through machine-guided metatranscriptomics." *Proceedings of the*
National Academy of Sciences 119.7 (2022): e2100916119.

11. Alexander, Harriet, et al. "Eukaryotic genomes from a global metagenomic dataset
illuminate trophic modes and biogeography of ocean plankton." *bioRxiv* (2021).

12. Chen, Tianqi, and Carlos Guestrin. "Xgboost: A scalable tree boosting system."
Proceedings of the 22nd acm sigkdd international conference on knowledge discovery
and data mining. 2016.

13. O'Leary, Nuala A., et al. "Reference sequence (RefSeq) database at NCBI: current status,
taxonomic expansion, and functional annotation." *Nucleic acids research* 44.D1 (2016):
D733-D745.

14. Parks, Donovan H., et al. "GTDB: an ongoing census of bacterial and archaeal diversity
through a phylogenetically consistent, rank normalized and complete genome-based
taxonomy." *Nucleic Acids Research* (2021).

15. Aramaki, Takuya, et al. "KofamKOALA: KEGG ortholog assignment based on profile
HMM and adaptive score threshold." *Bioinformatics* 36.7 (2020): 2251-2252.

16. Kanehisa, Minoru, Yoko Sato, and Kanae Morishima. "BlastKOALA and GhostKOALA:
KEGG tools for functional characterization of genome and metagenome
sequences." *Journal of molecular biology* 428.4 (2016): 726-731.

17. Bowers, Robert M., et al. "Minimum information about a single amplified genome
(MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and
archaea." *Nature biotechnology* 35.8 (2017): 725-731.

18. Blin, K., https://github.com/kblin/ncbi-genome-download, version 0.2.10

19. Hyatt, Doug, et al. "Prodigal: prokaryotic gene recognition and translation initiation site
identification." *BMC bioinformatics* 11.1 (2010): 1-11.

20. Kanehisa, Minoru. "The KEGG database." *Novartis found symp*. Vol. 247. 2002.

21. Steven Pawley, 2022, https://github.com/stevenpawley/recipeselectors, version 0.0.1

22. Kuhn, Max, and Hadley Wickham. "Tidymodels: a collection of packages for modeling
and machine learning using tidyverse principles." *Boston, MA, USA.[(accessed on 10*
December 2020)] (2020).

23. Nayfach, Stephen, et al. "A genomic catalog of Earth's microbiomes." *Nature*
biotechnology 39.4 (2021): 499-509.
https://genome.jgi.doe.gov/portal/GEMs/GEMs.home.html

24. Parks, Donovan H., et al. "CheckM: assessing the quality of microbial genomes
recovered from isolates, single cells, and metagenomes." *Genome research* 25.7 (2015):
1043-1055.

25. SRA Toolkit Development Team,
https://trace.ncbi.nlm.nih.gov/Traces/sra/sra.cgi?view=software)

26. Phil Ewels, https://sra-explorer.info/

27. Chen, Shifu, et al. "fastp: an ultra-fast all-in-one FASTQ
preprocessor." *Bioinformatics* 34.17 (2018): i884-i890.

28. Li, Heng. "seqtk Toolkit for processing sequences in FASTA/Q formats." *GitHub* 767
(2012): 69.

29. Bankevich, Anton, et al. "SPAdes: a new genome assembly algorithm and its applications
to single-cell sequencing." *Journal of computational biology* 19.5 (2012): 455-477.

563 **Supplementary Figures**

567 Supplementary Figure 1. Distribution of phyla of bacterial genomes from which annotation data was used during

568 model training and testing. See Supplementary Table 2 for the full metadata table.

575
576
577 Supplementary Figure 2. Boxplot of the ratio of positive to negative classes of observations in all MetaPredict
578 stacked ensemble training and test datasets (**a**, *n* = 293) and neural network models (**b**, *n* = 183). Each train/test
579 dataset split contains the same distribution of observation classes.

Heatmap of KEGG module M00122 test dataset features

601
602
603 Supplementary Figure 3. Gene copy number data for the set of features (genes) used by MetaPredict's stacked
604 ensemble model for KEGG module M00122 (cobalamin biosynthesis). This gene annotation data is from the held-

605   out test set of downsampled protein annotation data for KEGG module M00122, and is only showing the gene copy

606   number of genes used by the stacked ensemble model for making predictions.

Supplementary Figure 4. **Panel a**: Bar chart of the taxonomic distribution of genomes from the GEM repository

used during model validation. **Panel b**: Bar chart of the environmental sources of metagenomes the MAGs from this

test set were recovered from.

Supplementary Figure 5. Boxplots of performance metrics of 183 neural network and XGBoost binary classification

models. Downsampled gene annotations of high-quality genomes used in held-out test sets are from NCBI RefSeq

and GTDB databases. Each boxplot displays the distribution of model performance metrics for predictions on

randomly sampled versions of the gene annotation test sets in downsampling increments of 10% (90% down to 10%,

from right to left). The binary classifier performances are based on the classification of the presence or absence of

KEGG modules in the complete versions of the gene annotations that were downsampled for model testing.

649 Supplementary Figure 6. Boxplots of performance metrics of 183 neural network models on high-quality bacterial

650 GEM MAGs ($n = 100$). Model performance metrics are for predictions on downsampled versions of GEM genome

651 gene annotations in decreasing increments of 10% (retaining 10-90% of the annotations in each test set).

652