# High-resolution glucose tracing and in situ imaging reveals regulated lactate production in human pancreatic β cells

Federica Cuozzo[1#], Katrina Viloria[1,2#], Ali H. Shilleh[2], Daniela Nasteska[1,2], Charlotte Frazer-Morris[2], Zicong Jiao[1,3$], Adam Boufersaoui[1], Bryan Marzullo[1], Hannah R. Smith[1], Caroline Bonner[4], Julie Kerr-Conte[4], Francois Pattou[4], Rita Nano[5,6], Lorenzo Piemonti[5,6], Paul R.V. Johnson[7], Rebecca Spiers[7], Jennie Roberts[1], Gareth G. Lavery[1,8], Anne Clark[2], Carlo D.L. Ceresa[2], Leanne Hodson[2], Amy P. Davies[9], Guy. A. Rutter[9,10,11], Masaya Oshima[12], Raphaël Scharfmann[12], Matthew J. Merrins[13], Ildem Akerman[1], Daniel A. Tennant[1*], Christian Ludwig[1*] and David J. Hodson[1,2*]

[1]Institute of Metabolism and Systems Research (IMSR), and Centre of Membrane Proteins and Receptors (COMPARE), University of Birmingham, Birmingham, UK.
[2]Oxford Centre for Diabetes, Endocrinology and Metabolism (OCDEM), NIHR Oxford Biomedical Research Centre, Churchill Hospital, Radcliffe Department of Medicine, University of Oxford, Oxford, UK.
[3]Geneplus-Beijing, Changping District, Beijing 102206, China.
[4]University of Lille, Institut National de la Santé et de la Recherche Médicale (INSERM), Centre Hospitalier Universitaire de Lille (CHU Lille), Institute Pasteur Lille, U1190 -European Genomic Institute for Diabetes (EGID), F59000, Lille, France.
[5]San Raffaele Diabetes Research Institute, IRCCS Ospedale San Raffaele, Milan, Italy.
[6]Vita-Salute San Raffaele University, Milan, Italy.
[7]Nuffield Department of Surgical Sciences, University of Oxford, John Radcliffe Hospital, Oxford, United Kingdom
[8]Centre for Systems Health and Integrated Metabolic Research (SHiMR), Department of Biosciences, School of Science and Technology, Nottingham Trent University, Nottingham, UK.
[9]Section of Cell Biology and Functional Genomics, Division of Diabetes, Endocrinology and Metabolism, Department of Metabolism, Digestion and Reproduction, Imperial College London, London, U.K.
[10]CHUM Research Centre and Faculty of Medicine, University of Montreal, Montreal, QC, Canada.
[11]Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore.
[12]Université Paris Cité, Institut Cochin, INSERM U1016, CNRS UMR 8104, 75014 Paris, France.
[13]Department of Medicine, Division of Endocrinology, Diabetes, and Metabolism, University of Wisconsin-Madison, Madison, WI 53705, USA; William S. Middleton Memorial Veterans Hospital, Madison, WI 53705, USA

[#]Joint first authors

*Correspondence should be addressed to:
david.hodson@ocdem.ox.ac.uk, c.ludwig@bham.ac.uk, d.tennant@bham.ac.uk

**Key words:** Glucose tracing, islet, metabolism, pyruvate, lactate, LDH, pyruvate dehydrogenase, imaging, GC-MS, NMR

**Short title:** Human β cells robustly produce lactate

**ABSTRACT**

Using $^{13}C_6$ glucose labeling coupled to GC-MS and 2D $^1H$-$^{13}C$ HSQC NMR spectroscopy, we have obtained a comparative high-resolution map of glucose fate underpinning β cell function. In both mouse and human islets, the contribution of glucose to the TCA cycle is similar. Pyruvate-fueling of the TCA cycle is primarily mediated by the activity of pyruvate dehydrogenase, with lower flux through pyruvate carboxylase. While conversion of pyruvate to lactate by lactate dehydrogenase (LDH) can be detected in islets of both species, lactate accumulation is six-fold higher in human islets. Human islets express LDH, with low-moderate *LDHA* expression and β cell-specific LDHB expression. LDHB inhibition increases glucose-dependent lactate generation in mouse and human β cells, and decreases $Ca^{2+}$-spiking frequency without affecting ATP/ADP levels. Thus, we show that LDHB limits glucose-stimulated lactate generation in β cells. Further studies are warranted to understand how lactate impacts β cell metabolism and/or function.

**HIGHLIGHTS**

- Human and rodent islets generate lactate following glucose stimulation.
- β cells specifically express LDHB, which acts to limit lactate generation.
- LDHB inhibition influences $Ca^{2+}$ spiking frequency without affecting ATP/ADP ratio.

**eTOC**

Cuozzo et al show that glucose-stimulated rodent and human islets generate lactate. Transcriptomic and imaging analyses reveal that LDHB is specifically expressed in β cells and unexpectedly restrains lactate production. LDHB expression and thus regulated lactate generation might reflect a key mechanism underlying β cell metabolism, function and survival.

## INTRODUCTION

β cells are highly-adapted as glucose sensors and need to balance glucose-dependent insulin release with housekeeping metabolic functions. The traditional view of β cell metabolism focuses on a tight relationship between blood glucose concentration, oxidative phosphorylation and stimulus-secretion coupling. Following a rise in glycemia, glucose enters the β cell through facilitated transport via low affinity glucose transporters (GLUT1 and GLUT2 in humans and rodents, respectively) [1,2]. Glucose is then phosphorylated by a low affinity hexokinase, glucokinase (GK), leading to closure of the ATP-sensitive potassium ($K_{ATP}$) channels (reviewed in [3,4]). The increase in membrane voltage then drives $Ca^{2+}$ flux through voltage-dependent $Ca^{2+}$ channels [3], which together with amplifying signals (amino acids, isocitrate, cAMP etc) [4-6], drives first and second phase insulin granule exocytosis. Direct conversion of pyruvate to lactate is thought to be suppressed in the β cell due to low levels of lactate dehydrogenase A (*LDHA*) [7-10], ensuring that the majority of pyruvate enters the TCA cycle.

Recent studies have challenged the canonical view of β cell metabolism by showing that ATP/ADP generation might be compartmentalized. $K_{ATP}$ channels are locally regulated by a membrane-associated glycolytic metabolon that is assisted by the mitochondrial phosphoenolpyruvate (PEP) cycle [11-13]. The glucose-dependent rise in the ATP/ADP ratio raises mitochondrial voltage to stall oxidative phosphorylation and the TCA cycle, activating anaplerotic flux through pyruvate carboxylase (PC) and the PEP cycle, which supports ATP production by pyruvate kinase (PK) until $K_{ATP}$ channels are closed. Following membrane depolarization, the rise in ADP supports a highly oxidative state that depends on high TCA cycle flux and pyruvate consumption by pyruvate dehydrogenase (PDH), which supports oxidative phosphorylation and sustained secretion [11,12,14]. These two complementary states might dictate the mitochondrial fate of pyruvate.

Alongside compartmentalized ATP/ADP generation, anaplerotic metabolism serves as an important source of coupling or amplifying factors for glucose-stimulated insulin secretion [15-17]. For example, PC directs pyruvate entry into the TCA cycle, which feeds isocitrate into isocitrate dehydrogenase 1 (IDH1) to support 2-ketoglutarate (2-KG)/NADPH generation and SENP1 activation [6,18]. This reaction is further supported by metabolism of glutamine by the reductive TCA cycle [19].

Despite the clear importance of metabolism for β cell insulin release and phenotype, we are still lacking a high-resolution, integrated view of β cell glucose fate. In particular, most data using glucose tracing and GC-MS/NMR spectroscopy has been derived from insulinoma cell lines, which provide the requisite cell mass for metabolite detection/annotation. However, insulinoma cell lines have to balance the need for insulin secretion with proliferation, an energy-consuming process [15,17,20-23], and fail to display normal cell heterogeneity known to influence metabolism [24-26]. In addition, species-differences in islet cell composition and β cell function have been described [27-29], yet their influence on energetics is still unclear. Lastly, bulk metabolomics has been informative for understanding β cell metabolism [30,31] and, while sensitive, lacks the resolution required to pinpoint glucose fate during glycolysis and the TCA cycle. Recent studies have shown the power of metabolic tracing to compare stem-derived β cell and human islet metabolism, highlighting key differences in glucose metabolism, metabolite trafficking and sensitivity to various fuels [32,33]. However, these studies did not (understandably) combine GC-MS/LC-MS with NMR for TCA metabolite

annotation, nor cross-compare human and mouse islets. Thus, our understanding of β cell
glucose metabolism in mouse and human islets remains incomplete.

In the present study, we combine GC-MS-based $^{13}C_6$ glucose tracing with the resolution of
2D $^1H$-$^{13}C$ HSQC NMR multiplet analysis to map glucose fate in islets with high sensitivity.
By applying this dual approach to human and mouse samples, we are able to provide a
detailed cross-species depiction of glucose metabolism. Detailed examination of $^{13}C$
labelling patterns confirms that PDH is the major contributor to the TCA cycle. We further
show that pyruvate is directly converted to lactate in both human and mouse islets.
However, lactate accumulation is much higher in human islets. Analysis of multiple
transcriptomic datasets shows that *LDHB* expression is restricted to human β cells,
confirmed using immunohistochemistry for LDHB protein. In situ metabolic imaging using
specific inhibitors shows that LDHB limits lactate production, but does not affect glucose-
stimulated ATP/ADP and $Ca^{2+}$ rises. We thus provide an integrated view of mouse and
human islet metabolism, show that β cell-specific LDHB expression limits lactate generation,
and suggest that the role of lactate in β cell metabolism, function and survival is worth
revisiting.

**RESULTS**

**Glucose contribution to TCA cycle in human and mouse islets**

To investigate glucose handling, mouse and human islets were incubated overnight with $^{13}C_6$
glucose prior to metabolite extraction and GC-MS and 2D $^1$H,$^{13}$C HSQC-NMR spectroscopy
(Figure 1A). To allow sufficient glucose flux for detection of $^{13}$C incorporation into TCA
metabolites, without inducing glucotoxicity, 10 mM $^{13}C_6$ glucose was used. The incorporation
of $^{13}$C from $^{13}C_6$ glucose into the TCA cycle metabolites was then established via mass
isotopologues distribution (MID) analysis (Figure 1B).

Suggesting a similar progression of glycolysis and the TCA cycle, glucose incorporation into
glycerol-3-phosphate (G-3-P) (Figure S1A) and the major metabolites malate, alanine, and
glutamate was not different between mouse and human islets (Figure 1C-E). However, a
slight but significant increase in m+2/m+3 aspartate and fumarate was detected in mouse
islets (Figure 1F and G), reflecting an increased contribution of glucose-derived pyruvate
into the TCA cycle via acetyl-CoA. Total aspartate and alanine levels did not differ between
the species (Figure 1H and I), whereas malate and fumarate levels were lower in mouse
(Figure 1J and K). Glutamate levels were ~3-fold higher in mouse versus human islets,
despite similar MIDs, implying that there is a larger contribution of non-labelled glutamate to
the total glutamate pool in this species e.g. through glutamine transport (Figure 1L).

**Lactate generation is higher in human compared to mouse islets**

To obtain a higher definition view of pyruvate management, its contribution to the production
of alanine and lactate was assessed. In both species, glucose incorporation could be
detected in m+2 and m+3 lactate, derived from the TCA cycle and direct pyruvate
conversion, respectively (Figure 2A-C). While the MID for alanine was similar in islets from
both species (Figure 1D), the accumulation of m+2 and m+3 lactate was significantly (~ 6-
fold) higher in humans (Figure 2A). The total amount of lactate was higher in human than in
mouse islets (Figure 2B-D). Supporting the notion of increased lactate production through
glycolytic input in human islets, the m+3 G-3-P /m+3 lactate labeling ratio was higher in
mouse versus human islets (Figure S1B). While accumulation of m+2 lactate was expected,
we were surprised to detect significant m+3 lactate accumulation in human islets, since *Ldha*
has been shown to be "disallowed" in rodent β cells, hence preventing alternative fates for
pyruvate [7-10].

Suggesting that m+3 lactate is derived mainly from pyruvate, rather than multiple rounds of
the TCA cycle, were the following findings: 1) m+4 malate and m+4 aspartate were
significantly lower compared to m+2 and m+3 isotopomers (Figure 1C and F); and 2) m+4
malate and m + 4 aspartate were similar in human and mouse, despite significantly higher
m+3 lactate in human (Figure 1C and F). In addition, previous tracing studies have shown
m+3 lactate accumulation in iPSC-derived islets and human islets after 1 hour of tracing [33].
Lastly, lack of lactate detection in previous studies with mouse islets and rodent beta cell
lines is unlikely due to the shorter timecourses used, since we were also unable to observe
meaningful m+3 lactate generation after 24 hours incubation with $^{13}C_6$-glucose.

**High-resolution annotation of $^{13}C_6$ glucose tracing data**

To identify isotopomer patterns with high-resolution, the MID analysis of $^{13}C_6$ glucose-traced

human and mouse islets was annotated with 2D $^1H$-$^{13}C$ HSQC NMR multiplet analysis. From

uniformly labeled glucose, $^{13}C$ atoms are incorporated into the metabolites of the TCA cycle

through the activity of PDH and PC (Figure 3A, B). This leads to the formation of labeling

patterns within the chemical structure of each metabolite that are specific to the pathway

from which they are produced (Figure 3A, B). Therefore, the positions of $^{13}C$ atoms within

each metabolite can be utilized to elucidate the relative activities of PDH and PC. To define

the different isotopomer patterns a numerical notation was used, where the numbers 0 and 1

indicate $^{12}C$ and $^{13}C$ atoms, respectively. Confirming the accuracy of the approach and the

robustness of our findings, the accumulation of $lactate_{111}$ (i.e. fully-labeled lactate) was

significantly higher in human compared to mouse islets, in line with the MID glucose-tracing

data (Figure 2A-C) (Figure 3C-E).

**TCA cycle fuelling is more dependent on flux through PDH than PC in human and**

**mouse islets**

In both human and mouse islets, $lactate_{110}$ made a greater contribution to the m+2

isotopologue pool than the other possible isotopomers (Figure 3C-E). This finding suggests

that TCA-derived lactate is produced primarily from the oxidative TCA cycle rather than the

reductive metabolism of PC-derived glutamate, from which $pyruvate_{011}$ and then $lactate_{011}$

would arise (Figure 3C-E). We also noticed that the majority of alanine was either 000 or

111, with only a very minor contribution to the other isotopomers (Figure 3F-H). As such, the

labeled portion of alanine is exclusively produced from pyruvate upstream of the TCA cycle,

meaning that the accumulation of $pyruvate_{110}$ from $malate_{1100}$ (i.e. TCA-derived) is almost

entirely employed to regenerate cytoplasmic $lactate_{110}$ (Figure 3F-H). $Alanine_{111}$

accumulation was slightly (~20%) higher in human than mouse islets, reflecting a greater

contribution of transamination toward amino acid production (Figure 3F-H). Supporting the

lactate isotopomer data, the contribution of $^{13}C_6$ glucose to the labeling patterns of glutamate

was found to be similar in humans and mice (Figure 3I, J). In both species, the most

abundant labeled isotopomer was $glutamate_{00011}$ (Figure 3I, J), which is derived from TCA

cycle flux through the activity of PDH (Figure 3A, J). Together, these findings provide further

evidence that pyruvate management in the pancreatic β cell occurs primarily through PDH at

the stimulatory glucose concentration used here [15,34].

**LDH is expressed at higher levels in human compared to mouse β cells**

<span style="color:red">Given that lactate generation was detected in human islets, and to a lesser extent mouse</span>

<span style="color:red">islets, we set out to investigate whether LDH protein is expressed in β cells. To this end</span>, we

performed immunohistochemistry using an antibody with cross-reactivity against LDHA,

LDHB and LDHC (i.e. total LDH). Since LDHC is undetectable in the pancreas [35] and is

largely confined to the testis [36], we assume that total LDH staining in pancreas represents

LDHA and LDHB isoforms. Punctate cytoplasmic LDH staining was detected in human

endocrine and exocrine pancreas, including in β cells (Figure4A). LDH expression was much

lower in mouse compared to human islets, with slightly higher levels in the exocrine versus

endocrine compartments (Figure 4A and C). However, following 8 weeks high-fat diet (HFD)

feeding to induce beta cell de-differentiation, LDH protein expression increased ~ 2-fold

versus age-matched standard diet controls (Figure 4B and C). Of note, LDH expression also

increased in the exocrine compartment, with HFD mouse pancreata resembling more their

human counterpart (Figure 4A-D). Occasional intra-islet cells were found to express very

high LDH levels in both human and mouse, likely representing endothelial cells known to be
enriched for LDHA [37] (correlation for enrichment = 0.502; Human Protein Atlas,
proteinatlas.org) (Figure 4A and B). β cell de-differentiation was confirmed in the same
samples using immunostaining for PDX1, which was significantly lower under HFD versus
standard chow conditions (Figure 4E and F).

Strong LDH staining could be detected in human liver sections derived from patients with
non-alcoholic steatohepatitis (NASH), in which hepatocytes express high LDH levels
(correlation for enrichment = 0.502; Human Protein Atlas, proteinatlas.org) (Figure 4G) [38,39].
As expected from studies of enzyme activity, LDH expression was higher in liver than in β
cells (Figure S2A-C) (LDH CTCF mean ± SD: $1.16 \times 10^6 \pm 2.06 \times 10^5$ versus $2.85 \times 10^5 \pm$
$4.90 \times 10^4$, liver versus β cell; P<0.0001, Mann-Whitney test) (n = 240 cells, n = 3 donors) [8].
All results were confirmed with multiple immunostaining runs, using both 40x and 63x
objectives (Figure S2A-D).

**_LDHB_ is expressed in human β cells within the endocrine compartment**

Next, we interrogated published RNA-sequencing studies to understand which LDH
isoenzymes might underlie lactate generation in human islets. Using data from FACS-
purified fractions, _LDHB_ was found to be specifically- and highly-expressed in β cells (shown
also in [40]) within the islet, whereas the opposite convention was observed for _LDHA_ (i.e.
specifically and highly expressed in α cells) (Figure 5A). Analysis of single cell RNA-seq data
confirmed these findings across multiple independent datasets (Figure 5B and C), as
visualized according to endocrine and non-endocrine cell clusters (Figure 5D and E).
Confirming accuracy of the clustering-based analysis (and inability of β cells to export
lactate), the lactate transporter MCT1, encoded by _SLC16A1_, could not be detected in any
islet endocrine cell type (Figure 5F), as previously reported [41,42].

~~While LDHB is generally thought to convert lactate -> pyruvate~~Recent studies in cancer cells
have shown that pyruvate -> lactate conversion is unaffected in single knockouts of LDHA
and LDHB, and only in a double LDHA/B knockout is pyruvate no longer converted to lactate
[43,44]. Thus, LDHB can theoretically compensate for LDHA activity when LDHA expression
levels are low. However, we cannot exclude the possibility that LDHA may also catalyze
pyruvate to lactate conversion in β cells as they do contain detectable LDHA mRNA, based
on both single cell RNA-sequencing and bulk RNA-sequencing of FACS purified α cells and
β cells (Figure 5A-D). This is consistent with the open chromatin conformation and
transcription factor binding to the LDHA promoter in the human islet (Figure 5G). By
contrast, _LDHB_ possessed two specific enhancers within the CTCF boundaries, suggestive
of β cell-specific regulation, although no transcription factor binding was observed at the
promoter (Figure 5H).

**LDHB protein localizes predominantly to human β cells within the endocrine**
**compartment**

To confirm transcriptomic findings, immunohistochemistry was performed using an antibody
against LDHB, validated by the human tissue atlas using TMT-MS, protein array and
subcellular localization (https://www.proteinatlas.org/ENSG00000111716-
LDHB/summary/antibody). To further validate the antibody for use in human β cells, western
blotting for LDHB was performed on extracts from EndoC-βH1 cells treated with either siCTL

or siLDHB. Confirming antibody specificity, a three-fold reduction in LDHB expression could
be seen in siLDHB versus siCTL samples, with knockdown further confirmed by qRT-PCR
for mRNA levels (Figure S3A and B).

Similarly to total LDH (i.e. LDHA/LDHB), LDHB was located throughout the cytoplasm as
punctate spots (Figure 6A). In line with the transcriptomic analysis and further confirming
antibody specificity, LDHB protein could be readily observed throughout the β cell
compartment, but was undetectable in the majority (81%) of α cells (Figure 6A-C). We did
however notice a small subpopulation (~19%) of α cells with high levels of LDHB,
comparable to those detected β cells (Figure 6B-D). Likewise, a proportion of β cells could
be differentiated by their absent or low expression of LDHB (26%) (Figure 6B-D). As for
LDH, all results were confirmed with multiple immunostaining runs, using both 40x and 63x
objectives (Figure S2A-D).

Since isolated islets were used for metabolic tracing, and pancreas digestion and culture
could conceivably influence LDH/LDHB expression levels, immunostaining was repeated in
isolated human islets. Almost identical results were obtained in isolated human islets and
pancreas sections, suggesting that culture time does not influence LDH/LDHB levels and
ergo lactate generation (Figure 6E-H).

**LDHB inhibition slightly increases lactate generation in human β cells without**
**affecting the cytosolic ATP/ADP setpoint**

While GC-MS- and NMR-based stable isotope tracing provide detailed information about
glycolysis and the TCA cycle, this method requires hundreds of islets and is not amenable to
purified β cell and α cell fractions.

To establish the contribution of the β cell compartment to lactate generation, we used Ad-
RIP-Laconic, a β cell-specific lactate FRET sensor [45,46]. Experiments in human islets showed
that high (17 mM) glucose was able to increase intracellular lactate levels after a lag of ~ 5
mins (Figure 7A-H). To investigate any role of LDHB, islets were pre-incubated in vehicle or
AXKO-0046 for 2 hours, a specific LDHB inhibitor with no detectable activity on LDHA [47].
AXKO-0046 was used at 10 µM, which corresponds to its maximal inhibitory concentration
[47]. While AXKO-0046 did not appreciably alter glucose-stimulated lactate dynamics, there
was a small (10-20%) but replicable increase in glucose-stimulated lactate generation
compared to control in islets from all donors examined (Figure 7A-I). Similar results were
observed in mouse β cells, which express LDHB albeit at lower levels compared to human β
cells (Figure 7J-L).

Given that LDHB reduces lactate generation, we next looked at whether LDHB inhibition
influences glucose-stimulated ATP/ADP and $Ca^{2+}$ rises in human β cells. Vehicle-treated
human islets responded to high glucose with increases in ATP/ADP ratios, which were not
significantly influenced by LDHB inhibition (Figure 7M). Similarly, glucose-stimulated $Ca^{2+}$
fluxes were identical between vehicle- and AXKO-0046-treated islets (Figure 7N and O). A
more detailed analysis of $Ca^{2+}$ oscillations, however, showed a reduction in $Ca^{2+}$ spiking
frequency in the presence of AXKO-0046 (Figure 7N and O), in keeping with previous
results showing that lactate directly opens $K_{ATP}$ channels [13].

Lastly, we attempted to replicate experiments in EndoC-βH5 spheroids, which also express
LDHB (Figure S3C). Unexpectedly, EndoC-βH5 spheroids did not tolerate long pre-

incubation times with higher concentrations of AXKO-0046, showing signs of blebbing. Thus, spheroids were instead pre-incubated with 100 nM AXKO-0046. At this concentration, AXKO-0046 suppressed both glucose-stimulated ATP/ADP ratios and $Ca^{2+}$ fluxes (Figure S3D-G). Analysis of transcriptomic data revealed that EndoC-βH5 spheroids express moderate LDHA levels, which might explain the exaggerated effects of LDHB inhibition versus primary human β cells (64.11 ± 2.60 versus 247.73 ± 5.00 TPM, *LDHA* versus *LDHB*; mean ± SD; taken from GSE224732 [48]).

**DISCUSSION**

Using $^{13}C_6$ glucose labeling coupled to GC-MS and 2D $^{1}H$-$^{13}C$ HSQC NMR spectroscopy, we have been able to obtain a high-resolution map of glucose fate within human and mouse islets. Unexpectedly, both human and mouse islets accumulate m+2 and m+3 lactate, meaning that lactate produced downstream of the TCA cycle (m+2) as well as via pyruvate -> lactate conversion (m+3) contribute equally to overall lactate generation. However, pyruvate -> lactate conversion was much higher (~ 6-fold) in human compared to mouse islets, in line with relative LDH/LDHB expression levels. Finally, we show that labeled lactate and glutamate accumulate as lactate$_{110}$ and glutamate$_{00011}$, confirming greater flux through PDH versus PC in both species, and the ultimate fate of glucose oxidation in the TCA cycle [11]. The major findings are schematically represented in Figure 8.

The observation that the islet lactate pool is derived from both TCA cycle- and pyruvate-derived sources suggests that mechanisms must be in place for direct pyruvate conversion. In many tissues, pyruvate would be converted to lactate by LDH, however, lactate generation is minimal in purified rat beta cells [49], and the *Ldha* isozyme of the enzyme has been shown to be disallowed or absent in the murine pancreatic β cell [8,10,50]. In keeping with these findings and further supporting a more limited role for LDH in rodents, lactate and LDH protein could only be detected at low levels in mouse islets, although glucose-stimulated lactate rises could still be detected by us and others in single β cells[46]. By contrast, LDH expression in human β cells was ~2-fold higher than in mouse β cells, albeit less than that in the human liver, known to strongly express LDH. Analysis of multiple published scRNA-seq datasets further showed that human β cells specifically and strongly express *LDHB* within the islet, which encodes the beta subunit of LDH, confirming previous single cell screening studies by van Gurp et al [40]. LDHA expression was also detected at low-moderate levels based upon bulk RNA-seq of purified human β cells.

Oxygen availability and ergo NADH:NAD$^{+}$ are both known to influence affinity of LDHA and LDHB for their substrates, with anaerobic conditions favoring pyruvate to lactate conversion and vice versa under normoxia. Notably, glucose-stimulation leads to large increases in beta cell $O_2$ consumption sufficient to activate HIF enzymes [51,52]. Supporting the findings here, recent studies using Seahorse assays have shown that glucose-stimulated oxygen consumption ratio is ~ 30% higher in mouse versus human islets due to higher leak (with caveats about inter-islet heterogeneity noted) [53]. As such, for a given $O_2$ supply, mouse islets consume more $O_2$, which should lead to increased lactate generation. However, lactate generation remains low compared to human islets, since mouse islets express less LDHA/LDHB.

It is possible that α-cells contribute to the accumulation of lactate. Human α cells account for ~35% of the entire islet and express *LDHA* at levels six times higher than β cells [54,55]. However, a major source of α cell lactate is via monocarboxylate transporters [9,56,57], which are unlikely to play a role here as lactate was absent from the tracing medium. In addition, while the total amount of lactate was only doubled in humans compared to mice, the m+3 (TCA-derived) and m+3 (glycolytically-derived) lactate accumulation was ~six-fold higher in human *versus* mouse islets, which cannot be accounted for solely by differences in α cell proportion. Total LDH levels were found to be similar in human α-cells and β cells, although a recent study suggests that plasma membrane-compartmentalized LDH activity differs between human α and β-cells, and plays a more significant role in α-cells [13]. Lastly, studies

with Laconic, a lactate FRET sensor, showed that human β cells are capable of generating
significant intracellular lactate levels in response to stimulation with high glucose
concentration. Mouse β-cells were also found to generate cytosolic lactate in response to
glucose (also [46]) using the same approach, but the species were not quantitatively
compared. Taken together, these data suggest that α cell lactate only makes a very minor
contribution to the whole m+3 and m+2 lactate increase detected here. We note that human
islet preps also typically comprise ~ 50% acinar, duct and other non-islet cells, where
LDHA/LDHB is enriched versus islet cells (see Figure 3D and E). While islets were hand-
picked, filtered and cultured for 2-3 days during which acinar cells are expected to die off, we
cannot exclude a contribution of the exocrine and other compartments to the findings here.

Human and mouse islets display a greater accumulation of lactate$_{110}$, rather than lactate$_{011}$.
While the accumulation of lactate$_{110}$ is indistinguishable in the PDH- and PC-mediated TCA
cycle, the 011 isotopomer would only derive from the reductive metabolism of PC-derived
glutamate. Corroborating this, in islets from both sschuiecies, the major glutamate
isotopomer derived from exogenous $^{13}C_6$ glucose was glutamate$_{00011}$. Although glutamate is
not a TCA cycle metabolite, it is in rapid exchange with α-KG and can be used as a read-out
of TCA cycle flux through PDH or PC. Consequently, the accumulation of glutamate$_{00011}$
provides further evidence for a higher reliance of the TCA cycle on the activity of PDH,
rather than PC. Although PC and PDH were thought to contribute equally to the TCA cycle in
β-cells [20-22], previous studies have shown that high glucose concentrations in vitro, more
reflective of those seen post-prandially in vivo, are associated with an increase toward PDH
activity [11,15,34]. Our studies thus show that the relative contribution of PC to the TCA cycle is
much lower than PDH (~20%), confirming findings from Alves et al in glucose-traced INS-1
cells [15] and Lewandowski et al in human islets [11]. This does not mean that PC flux is
unimportant, since PC is much more responsive to glucose stimulation than PDH [11].

While anaplerosis through PC is relatively limited in the β cell, we note that glucose carbons
can repeatedly transit the PEP cycle to increase ATP/ADP in the cytosol [11] (before pyruvate
is ultimately oxidized in the TCA cycle). As such, PC is able to make disproportionate
contributions to $K_{ATP}$ channel closure, and hence the triggering phase of insulin secretion, by
generating plasma-membrane localized increases in ATP/ADP [14]. Moreover, PC might make
relatively more important contributions to anaplerosis in stressed human beta cells in which
increased PC activity reduces NO production to counteract inflammation [58]. PC is also
important for funneling pyruvate into generation of the GSIS-coupling factors citrate and
isocitrate, which sustain 2-ketoglutrate and NADPH production [18]. Nonetheless, insulin
secretion is not the only energy sink on the β cell, and glucose flux through PDH is likely to
provide a source of acetyl-CoA to support other demands such as continued insulin release
and protein synthesis [49]. Indeed, insulin biosynthesis and ion pumping have both been
shown to be greater β cell energy sinks than exocytosis [59].

What might be the role of direct pyruvate to lactate conversion in pancreatic β cells? Since
the action of LDH leads to oxidation of NADH to NAD$^+$, lactate accumulation could provide a
source of reducing equivalents to support other NADH-producing metabolic pathways.
Providing evidence for an important contribution of pyruvate to lactate conversion in
NADH/NAD+ balance, the most abundant lactate isotopomers were lactate$_{111}$ and lactate$_{110}$,
which respectively represent lactate generation upstream and downstream of the TCA cycle.
Since the conversion of pyruvate -> lactate is associated with the generation of cytosolic
NAD$^+$, higher levels of total lactate$_{111}$ in humans might reflect an increase in the activity of

NADH-producing pathways relative to rodents. Notably, the alanine isotopomer distribution showed accumulation of alanine$_{000}$ and alanine$_{111}$, suggesting that pyruvate accumulated downstream of the TCA cycle is employed in the regeneration of lactate$_{110}$. Arguing against this possibility, β cells lack the capacity to extracellularly transport lactate, and are thought to already have a large capacity to produce reducing equivalents, e.g. via the glycerol phosphate and malate-aspartate shuttles [60]. Although membrane-associated LDH was found to form nanodomains with K$_{ATP}$ channels and support the local production of NAD$^+$ for GADPH, the contribution of LDH activity (at least in mouse β-cells) was small relative to other unidentified plasma membrane-associated NADH oxidases that support this function. Lastly, significant pools of cytosolic pyruvate and lactate, and sufficient LDH to equilibrate these pools, might provide a buffer to minimise wide fluctuations in NADH/NAD (or NAD(P)H/NADP), offering some protection against oxidative stress [61].

In summary, by combining MID and multiplet analyses, we show that glucose makes a similar contribution to glycolysis and the TCA cycle in human and mouse islets. Furthermore, the isotopomer distribution confirms that, in both species, the relative activity of PDH is higher than that of PC at elevated glucose concentration. However, the generation of fully-labeled lactate was found to be much higher in human versus mouse islets, with LDHB and acting to limit lactate levels. Together, these results demonstrate that lactate generation should be reconsidered in light of beta cell development, metabolism and function.

## Limitations

There are a number of limitations in the present studies. Firstly, glucose-tracing studies in purified α cells and β cells are warranted, although should be interpreted in light of loss of cell-cell interactions, changes in cell phenotype, as well as input material required. Nonetheless, our data provide a detailed and interrogable map of glucose metabolism pertaining to steady state insulin release in human and mouse islets, as well as other critical β cell housekeeping functions. Secondly, glucotoxicity might induce the upregulation of disallowed genes in β cells [62]. However, it is unlikely that the timings (12 hrs) and glucose concentration (10 mM) used here would overtly influence human β cell lactate generation, since *LDHA* or *LDHB* were not found to be differentially expressed in human islets exposed to 22.2 mM glucose for 4 days [63]. Thirdly, to increase resolution of glucose fate-mapping, tracing should be performed at different timepoints, similarly to recent studies [32,33]. Such studies are at the moment limited by 800 MHz NMR access and Helium availability. Fourthly, functional studies in primary human beta cells deleted for LDHA/LDHB are warranted, although there are concerns that the other LDH isoenzymes might be upregulated over chronic timecourses. Lastly, the effects of LDHB inhibition on glucose-stimulated lactate generation were relatively small. This could reflect the potency of AXKO-for LDHB in the tissue setting, existence of lactate microdomains, or the high glucose concentration used. Future studies looking at glucose concentration curves, as well as tracing-based analyses will be valuable.

 **STAR METHODS**

 **RESOURCE AVAILABILITY**

**Lead contact**

Further information and requests for resources and reagents should be directed to and will
be fulfilled by the lead contact, David J. Hodson (david.hodson@ocdem.ox.ac.uk).

**Materials availability**

This study did not generate any new unique reagents.

**Data and code availability**

This paper analyses existing, publicly available data. Accession numbers for the datasets
are listed in the key resources table.

All data reported in this paper will be shared by the lead contact upon request.

**EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS**

**Mice**

Male 8 to 12 week-old CD1 mice (Charles River stock no. 022) were used as tissue donors.
Mice were socially-housed in specific-pathogen free conditions under a 12 hour light-dark
cycle with ad libitum access to food and water, relative humidity 55 ± 10% and temperature
21 ± 2 °C.

Animal studies were regulated by the Animals (Scientific Procedures) Act 1986 of the U.K.
(Personal Project Licences P2ABC3A83 and PP1778740). Approval was granted by the
University of Birmingham and University of Oxford Animal Welfare and Ethical Review
Bodies (AWERB).

**Human**

Human islets (Lille): human pancreatic tissues were harvested from brain-dead adult donors
in accordance with the Lille clinical islet transplantation program's traceability requirements
(clinicaltrials.gov, NCT01123187, NCT00446264, NCT01148680), and were approved in
agreement with French regulations and the Ethical Committees of the University of Lille and
the Centre Hospitalier Régional Universitaire de Lille.

Human islets (Milan): the use of human islets for research was approved by the Ethics
Committee of San Raffaele Hospital in Milan (IPF002-2014).

Human islets (Oxford): human pancreata were retrieved from Donors after Brain-Stem Death
(DBD) with appropriate consent and ethical approval under 09/H0605/2 (REC: Oxfordshire
Rec B). Islets were isolated in the DRWF Oxford Human Islet Isolation Facility using
established isolation methods.

Human pancreas sections (Oxford): post-mortem pancreas samples were obtained from
donors, with appropriate permissions registered under CUREC R83564/RE001.

Studies with human islets and pancreata were approved by the University of Birmingham
Ethics Committee, the University of Oxford Ethics Committee, as well as the National
Research Ethics Committee (REC 16/NE/0107, Newcastle and North Tyneside, UK).

Liver sections (Oxford): Authorisation for research use of steatotic livers unsuitable for
transplantation was obtained by a specialist nurse in organ donation in accordance with
NHSBT guidelines, with appropriate permissions registered under REC 14/LO/0182. The
study was approved by the North East – Tyne and Weir South research ethics committee
(16/NE/0248) and by the NHSBT Research, Innovation and Novel Technologies Advisory
Group.

## METHOD DETAILS

### Study design

No data were excluded, and all individual data points are reported in the figures. The
measurement unit is animal or donor, with experiments replicated independently on different
days. Islet isolation is a nuisance variable and as such data are taken from independent islet
preparations. Samples and animals were allocated to treatment groups in a randomized
manner to ensure that all states were represented in the different experiment arms. MID
analysis was performed by a user blinded to sample identity. For metabolic tracing, nine
samples per investigated state are required to correctly reject the null hypothesis (two-tailed
test) for an effect size (d) = 1.5 (calculated in GPower 3.1).

### Mouse islets

Animals were culled using a schedule-1 method followed by injection of the common bile
duct with 1 mg/mL collagenase NB 8 (Serva) in RPMI 1640 (Gibco) and pancreas
dissection. After dissection, the pancreas was incubated in a water bath at 37°C for 12 min.
Subsequently, the tissues were shaken in 15 mL of RPMI 1640 and centrifuged for 1 min at
1500 rpm three times to induce mechanical digestion. Islets were separated using
Histopaque-1119 and 1083 (Sigma-Aldrich) gradients, before hand-picking and culture.
Unless otherwise stated, the islets obtained were kept in culture in RPMI 1640
supplemented with 10% fetal bovine serum (FBS, Gibco), 100 units/mL penicillin, and 100
µg/mL streptomycin (Sigma-Aldrich), at 37°C and 5% $CO_2$.

For immunohistochemistry, sections were cut from formalin-fixed, paraffin-embedded
(FFPE) pancreata obtained from wild-type male and female C57BL6 mice fed standard diet
or high fat diet (60% fat, Research Diets, catalog D12492) for 8-12 weeks. To reduce the
number of animals used in experiments in line with NC3Rs policy, blocks were re-used from
a previous study [64], new sections cut at different depth, and immunostained with different
markers (LDH, PDX1, see below).

### EndoC-βH1 and EndoC-βH5 cells

523 Total protein was extracted from provided EndoC-βH1 cells using RIPA lysis buffer. Lysis
524 buffer was supplemented with EDTA-free protease inhibitor and phosphatase inhibitor
525 cocktails. Lysates were resolved on SDS–PAGE, transferred to a nitrocellulose membrane
526 using the Trans-Blot Turbo Transfer Pack, blocked in 5% milk for 1 hr, followed by an
527 incubation with a polyclonal Anti-LDHB antibody (HPA019007) at 4 °C overnight. After

washing, membrane was then incubated for 2 hr at room temperature with secondary antibody conjugated to horseradish peroxidase (HRP). After incubation with Clarity Western ECL Substrate, bands were detected with Bio-Rad GelDoc Biosystems Western Blotting imager.

EndoC-βH1 cells were transfected using Lipofectamine RNAiMAX with ON-TARGETplus siRNA targeting *LDHB* (L-009779-00, Dharmacon, Horizon Discovery) at a final concentration of 80nM. Non-targeting control pool (D-001810-10, Dharmacon, Horizon Discovery) was used as a control. Briefly, siRNA and Lipofectamine RNAiMAX were combined in OptiMEM and applied to the cells. Medium was changed 2.5h later for fresh EndoC-βH1 culture medium. Cells were harvested in RIPA lysis buffer or in RLT lysis buffer for mRNA purification (Qiagen).

Lysis buffer was supplemented with EDTA-free protease inhibitor and phosphatase inhibitor cocktails. Lysates were resolved on SDS–PAGE, transferred to a nitrocellulose membrane using the Trans-Blot Turbo Transfer Pack, blocked in 5% milk for 1 hr, followed by an incubation with a polyclonal Anti-LDHB antibody (HPA019007) at 4 °C overnight. After washing, membranes were then incubated for 2 hr at room temperature with secondary antibody conjugated to horseradish peroxidase (HRP). After incubation with Clarity Western ECL Substrate, bands were detected with Bio-Rad GelDoc Biosystems Western Blotting imager.

Primers used for PCR were as follows: PPIA (housekeeping) Fw primer: ATGGCAAATGCTGGACCCAACA, PPIA (housekeeping) Rv primer: ACATGCTTGCCATCCAACCACT, LDHB Fw primer: TGATGGATCTGCAGCATGGG and LDHB Rv primer: CAGATTGAGCCGACTCTCCC.

To generate EndoC-βH5 (Human Cell Design) pseudoislets, 1 million cells were seeded in Aggrewell Multiwell plates in ULT1β1 culture media (Human Cell Design) at 37°C and 5% CO2. Pseudoislets aggregated in 3-4 days and media was changed every 2 days.

**Human islets**

Islets were cleared of possible debris via filtration with a 40 μm cut-off filter, hand-picked and cultured in CMRL medium (Corning) containing 5.5 mM glucose, 10% FBS, 100 units/mL penicillin, 100 μg/mL streptomycin and 0.1% amphotericin B (Sigma-Aldrich) at 37°C and 5% $CO_2$. Human donor characteristics are reported in Table S1.

### $^{13}C_6$ glucose tracing

For $^{13}C_6$ glucose tracing, 60 (for GC-MS) or 150-230 (for NMR) islets were used. Isolated islets were cultured in RPMI 1640, no glucose medium (Gibco), supplemented with 10% BSA, 10% FBS, 100 units/mL penicillin, and 100 μg/mL streptomycin plus 10 mM $^{13}C_6$ glucose (Sigma-Aldrich). After 24 h, the metabolites were extracted adding HPLC-grade methanol, HPLC-grade distilled $H_2O$ containing 1 μg/mL D6-glutaric acid and HPLC-grade chloroform (all from Sigma-Aldrich) in a 1:1:1 ratio, to the islets. Following centrifugation, the polar fractions were collected and vacuum dried before either GC-MS or NMR analyses. A 24 h tracing duration was used to allow steady state labelling of both glycolytic and TCA metabolites in the same sample, as well as sufficient $^{13}C$ incorporation for NMR-based labelling pattern annotation. While antibiotics have been shown to influence mitochondrial

570 function, research islet isolation is a non-sterile procedure. Antibiotic use is therefore
571 justified, since low grade infection is likely to exert a larger (and unnoticed) effect on
572 mitochondrial activity.

**GC-MS**

The dried polar extracts were prepared for GC-MS analysis through solubilization in 40 µL of
2% methoxyamine hydrochloric acid in pyridine (Fisher Scientific) at 60°C for 60 min and
derivatization with 60 µL of N-tertbutyldimethylsilyl-N-methyltrifluoroacetamide (MTBSTFA)
with 1% tertbutyldimethyl-chlorosilane (TBDMCS) (both from Sigma-Aldrich). The
suspension was further incubated at 60°C for 60 min, before being centrifuged at 13300 rpm
for 10 min at 4°C and transferred to chromatography vials with a glass insert (Restek) for
GC-MS analysis. The samples were analyzed on an Agilent 8890 GC and 5977B MSD
system. To do this, 1 µL of sample was injected in splitless mode with helium carrier gas at a
rate of 1.0 mL/min. The compound detection was carried out in scan mode and total ion
counts of each metabolite were normalized to the internal standard D6-glutaric acid and
corrected for natural $^{13}C$ abundance [65]. Retention times, ion counts and MID data are
provided in Table S2 (mouse) and Table S3 (human).

**NMR spectroscopy**

Following the $^{13}C_6$ glucose tracing, the dried polar metabolites were resuspended in 60 µL of
phosphate buffer: 57.8 mM disodium phosphate ($Na_2HPO_4$, Sigma-Aldrich), 42.2 mM
monosodium phosphate ($NaH_2PO_4$, Sigma-Aldrich), 0.5 mM 3-(trimethylsilyl) propionic-
2,2,3,3-d4 acid sodium salt (D4-TMSP, Sigma-Aldrich) in deuterium oxide ($D_2O$, Sigma-
Aldrich). Subsequently, the samples were centrifuged for 10 min at 14800 rpm and
sonicated in an ultrasonic bath for 5 min, before being loaded into NMR tubes (outer
diameter: 1.7 mm, Bruker) for acquisition. A Bruker Neo 800 MHz NMR spectrometer
equipped with a 1.7 mm z-PFG TCI Cryoprobe was used to acquire 2D $^{1}H$,$^{13}C$-HSQC NMR
spectra. The HSQC spectra were acquired with echo/anti-echo gradient coherence selection
with an additional pre-saturation for suppressing the residual water resonance. The spectral
widths were 15.6298 ppm and 189.7832 ppm in the $^{1}H$ and $^{13}C$ dimension, 512 complex
data points were acquired for the $^{1}H$ dimension and 25% (512) out of 2048 complex data
points were acquired for the $^{13}C$ indirect dimension using a non-uniform sampling scheme.
Apparent $^{13}C$,$^{13}C$ J-coupling was enhanced four-fold. The interscan relaxation delay was set
to 1.5 s. 2D $^{1}H$,$^{13}C$-HSQC spectra were reconstructed via the compressed sensing IRLS
algorithm using the MddNMR (version 2.5) [66] and NMRPipe (version 9.2) [67] software. All
NMR spectra were analyzed in the MATLAB based MetaboLab software package [68].

**Immunohistochemistry**

Formalin-fixed, paraffin-embedded pancreas, liver and isolated islets were cut at 5µm and
deparaffinised. Human donor characteristics are reported in Table S4. PBS with 2% BSA
and 0.2% Triton X was used to block sections for 1 hour at room temperature. Heat induced
antigen retrieval was performed with 10 mM citrate buffer (pH6) using a microwave for 10
minutes and cooled for 10 minutes in cold water. Sections were incubated with primary
antibodies overnight at room temperature, followed by washes with PBS- 2% BSA and 0.2%
Triton X. Secondary antibodies were incubated for 2 hours at room temperature. Sections
were mounted with DAPI (Vectashield, cat no. H-1500) and stored at 4°C. Primary

antibodies used were mouse anti-LDH (Santa Cruz Biotechnology, cat no. sc-133123, RRID:
AB_2134964), rabbit polyclonal anti-LDHB (Sigma-Aldrich, cat no. HPA019007,
RRID:AB_2670008), guinea-pig anti-PDX1 (Abcam, cat# ab47308, RRID:AB_777178),
guinea pig anti-insulin (Abcam, cat no. ab7842, RRID: AB_306130), and mouse monoclonal
anti-glucagon (Sigma-Aldrich, cat# G2654, RRID:AB_259852). Human and mouse LDHA,
LDHB and LDHC share 94.0%, 97.9% and 74.5% protein sequence identity, respectively.
We note however that LDHC is undetectable in the human pancreas [35] and is largely
confined to the testis [36]. For LDHA and LDHB detection, we used an amplification step
consisting of anti-mouse biotinylated (DAKO, cat no. E0413) or anti-rabbit biotinylated
(NOVEX, cat no. A16039, RRID: AB_2534713) antibodies, followed by incubation with
streptavidin-FITC (Sigma-Aldrich, Cat# S-3762). For all other markers, secondary antibodies
used were anti-guinea pig Alexa Flour 568 (Thermo Fisher, cat no. A-11075, RRID:
AB_141954), anti-guinea pig Alexa Flour 647 (Thermo Fisher, cat no. A-21450, RRID:
AB_141882) and anti-mouse Alexa Flour 568 (Thermo Fisher, cat no. A-11004, RRID:
AB_2534072).

Sections were imaged using an Olympus FV3000 confocal microscope equipped with high-
sensitivity spectral detectors and 40x, 1.25 NA and 60x, 1.30 NA silicone objectives.
Excitation and emission wavelengths for DAPI, Alexa Fluor 488, Alexa Flour 568 and Alexa
Flour 647 were $\lambda ex = 405$ nm / $\lambda em = 406 - 461$ nm, $\lambda ex = 488$ nm / $\lambda em = 499 - 520$ nm,
$\lambda ex = 561$ nm / $\lambda em = 579 - 603$ nm, $\lambda ex = 640$ nm / $\lambda em = 649 - 700$ nm respectively.

**Lactate, $Ca^{2+}$ and ATP/ADP imaging**

Isolated islets were transduced 24-48 hrs with the beta cell specific lactate FRET sensor,
Ad-RIP-Laconic [45,46], prior to widefield imaging using a Nikon Ti-E base equipped with 89
North LDI-7 Laser Diode Illuminator, 25x / 0.8 NA objective and Prime BSI Express sCMOS.
Excitation was delivered at $\lambda = 445$ nm and emission detected at $\lambda = 460–500$ and
$\lambda = 520–550$ nm for mTFP and Venus, respectively. All experiments were performed in
HEPES-bicarbonate buffer containing (in mmol/L) 120 NaCl, 4.8 KCl, 24 NaHCO3, 0.5
Na2HPO4, 5 HEPES, 2.5 CaCl2, 1.2 MgCl2, supplemented with 3–17 mmol/L d-glucose,
and bubbled with 95% $O_2$ / 5% $CO_2$. Laconic intensity was calculated as the ratio of
mTFP/Venus. AXKO-0046 (MedChemExpress, cat no. HY-147216) was used as a selective
nanomolar-affinity small molecule inhibitor of LDHB ($IC_{50} = 42$ nM, $IC_{MAX} = 10^{-5}$ nM).

$Ca^{2+}$ and ATP/ADP imaging was performed in human islets and EndoC- βH5 spheroids
loaded with Fluo8 or transduced with Ad-RIP-Perceval-HR, respectively [11,69]. Islets were
imaged as above, with excitation delivered at $\lambda = 470$ nm using a CrestOptics X-light V2
spinning disk head; and emission detected at $\lambda = 500-550$.

**Image analysis**

Images were deconvolved using Imaris Clearview (Oxford Instruments) and an AMD
Radeon Pro W5500 GPU with 8GB GDDR6. Image analysis was performed in ImageJ (NIH)
using corrected total cell fluorescence (CTCF), which is the integrated density corrected for
the product of cell area and mean background fluorescence [70,71].

**Transcriptomics analysis**

Quantification data for all published scRNA-seq datasets were kindly provided by Leon Van
Gurp [40]. In brief, pseudo-counts were normalized for each data set using Seurat [72], and the
cell identity was assigned based on the requirement for hormone gene expression to be in
the top 1% expressed genes in each cell using Aucell [73]. Quantification for published FACS
sorted alpha and β cells were obtained from GEO database repository under Arda et al [74].
The raw read files for each cell type were merged, trimmed and the transcripts were
quantified using Kallisto [75] or aligned and quantified as previously described [76], with similar
results.

For Uniform Manifold Approximation Projection (UMAP) plots, datasets were downloaded
from GSE150724 [40], GSE84133 [77] and GSE114297 [78]. GSE84133 [77] contained a single
donor with type 2 diabetes, excluded here. Seurat R package was used to filter, cluster, and
identify cell types. Cells were filtered if they had less than 500 unique genes and if more
than 20% of their UMI counts were of mitochondrial origin. Only moderate filtering of
mitochondrial DNA was conducted because of the nature of the cells. After filtering, raw UMI
counts were normalized by scaling to 10,000 followed by log2-transformation. Normalized
gene expression was used to find the top 2000 most variable genes. These genes were
ranked based on the number of samples they were deemed a top variable gene, scaled to
generate z-scores and used to run Principal component analysis (PCA). The first 50
principal components were used to define anchors between datasets using Reciprocal PCA
(RPCA). RPCA was run with Donor 1 from [40] as a reference. Subsequently, PCA was run on
the integrated data and a neighborhood graph built using 20 principal components.
Clustering was carried out using a resolution of 0.2 and visualized in 2D space (Uniform
Manifold Approximation Plot) for cluster identification. Cells clustered according to their cell
type, as marked by their expression of INS, GCG and SST. Most cells in the original clusters
had ambient levels of the non-identifying hormone, which might reflect cross-contamination
during FACs [79]. However, there were cells with low expression of the cluster-identifying
hormone, and cells with high expression of the non-identifying hormones. These cells were
excluded on the premise that they represent doublets, or misclustered cells. After cell type
identification, raw counts from each cell were aggregated (summed) according to cell type
and donor for 'pseudobulk' representation. Markers used for cell type identification are listed
in Table S5.

**QUANTIFICATION AND STATISTICAL ANALYSIS**

GraphPad Prism 9 (version 9.2.0) was used for all statistical analyses. Data distribution was
assessed using the D'Agostino-Pearson normality test. Pairwise comparisons were made
using Welch's test (assuming non-equal standard deviation between groups). Multiple
interactions were determined using one-way ANOVA or two-way ANOVA, with Sidak's post-
hoc test. For non-parametric data, pairwise comparisons were made using Mann-Whitney
test, and multiple interactions determined using Kruskal-Wallis test, with Dunn's post-hoc
test. Individual datapoints are shown in bar graphs. Unless otherwise stated in the figure
legend, all error bars represent mean ± S.E.M. and a p-value less than 0.05 was considered
significant: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$, ****$p < 0.0001$.

## AUTHOR CONTRIBUTIONS

F.C., K.V., A.S.H. performed experiments, analyzed data and wrote the manuscript. D.N., A.C., C.F-M. and H.R.S. performed experiments. Z.J. and I.A. performed bioinformatic analysis. R.N., L.P., C.B., F.P., J.K-C., P.R.V.J. and R.S. isolated and provided human islets. C.L. performed $^1$H,$^{13}$C-HSQC NMR experiments and analysis. G.G.L., D.T., A.B., B.M. and J.R. ran GC-MS on $^{13}$C$_6$ glucose labelled samples and provided analysis. L.H., A.C., C.D.L.C. provided human liver samples. A.P.D. and G.A.R. provided scRNA-seq analysis. R.S. and M.O. participated in antibody validation. M.J.M. provided Ad-RIP-Laconic and also advised on lactate imaging studies. D.J.H. supervised the studies, provided analysis and wrote the manuscript with input from all authors. All authors read and approved the studies.

## FUNDING

D.J.H. was supported by MRC (MR/S025618/1) and Diabetes UK (17/0005681 and 22/0006389) Project Grants, as well as a UKRI ERC Frontier Research Guarantee Grant (EP/X026833/1). This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Starting Grant 715884 to D.J.H.). A.S. was supported by a Novo Nordisk – Oxford Fellowship. G.G.L. was supported by a Wellcome Trust Senior Fellowship (104612/Z/14/Z). D.N. was supported by a Diabetes UK RD Lawrence Fellowship (23/0006509). L.H. was supported by British Heart Foundation Senior Basic Science Research Fellowships (FS/15/56/31645 and FS/SBSRF/21/31013). C.D.L.C. was supported by a Clinical Research Training Fellowship from the MRC. G.A.R was supported by a Wellcome Trust Investigator Award (212625/Z/18/Z), MRC Programme grant (MR/R022259/1), Diabetes UK Project grant (BDA16/0005485), CRCHUM start-up funds, an Innovation Canada John R Evans Leader Award (CFI 42649) an NIH-NIDDK (R01DK135268) project grant and a CIHR-JDRF team grant (CIHR-IRSC TDP-186358 and JDRF 4-SRA-2023-1182-S-N). R.S. was supported by the Dutch Diabetes Research Foundation and the DON Foundation. M.J.M was supported by the NIH/NIDDK (R01DK113013 and R01DK113103) and VA BLR&D (I01BX005113). I.A. was supported by RD Lawrence fellowship (Diabetes UK, 20/0006136), and Academy of Medical Sciences Springboard (SBF006\1140). D.T. was supported by a Cancer Research UK Programme Grant (C42109/A24747). The research was funded by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. The project involves an element of animal work not funded by the NIHR but by another funder, as well as an element focussed on patients and people appropriately funded by the NIHR. The DRWF Oxford Human Islet Isolation Facility was funded by the Diabetes Research and Wellness Foundation (DRWF) and Juvenile Diabetes Research Foundation (JDRF). Human pancreas sections were provided by the University of Alberta in Edmonton Alberta Diabetes Institute IsletCore (https://www.bcell.org/adi-isletcore.html) with the assistance of the Human Organ Procurement and Exchange (HOPE) program, Trillium Gift of Life Network (TGLN), and other Canadian organ procurement organizations.

## ACKNOWLEDGEMENTS

The authors thank Prof. Patrick E. MacDonald and Dr. Jocelyn Manning Fox (Alberta
Diabetes Institute IsletCore at the University of Alberta in Edmonton) for provision of human
pancreas sections. We would like to acknowledge the support and resources of the
Birmingham Metabolic Tracer Analysis Core.

**DISCLOSURE STATEMENT**

G.A.R. has received grant funding from, and is a consultant for Sun Pharmaceuticals
Industries Ltd. D.J.H. receives licensing revenue from Celtarys Research.

**REFERENCES**

1.	De Vos, A., Heimberg, H., Quartier, E., Huypens, P., Bouwens, L., Pipeleers, D., and Schuit, F. (1995). Human and rat beta cells differ in glucose transporter but not in glucokinase gene expression. Journal of Clinical Investigation *96*, 2489-2495. 10.1172/jci118308.

2.	Thorens, B., Sarkar, H.K., Kaback, H.R., and Lodish, H.F. (1988). Cloning and functional expression in bacteria of a novel glucose transporter present in liver, intestine, kidney, and beta-pancreatic islet cells. Cell *55*, 281-290. 10.1016/0092-8674(88)90051-7.

3.	Rorsman, P., and Ashcroft, F.M. (2018). Pancreatic beta-Cell Electrical Activity and Insulin Secretion: Of Mice and Men. Physiol Rev *98*, 117-214. 10.1152/physrev.00008.2017.

4.	Rutter, G.A., Pullen, T.J., Hodson, D.J., and Martinez-Sanchez, A. (2015). Pancreatic beta-cell identity, glucose sensing and the control of insulin secretion. Biochem J *466*, 203-218. 10.1042/BJ20141384.

5.	Henquin, J.C. (2000). Triggering and amplifying pathways of regulation of insulin secretion by glucose. Diabetes *49*, 1751-1760.

6.	Ferdaoussi, M., Dai, X., Jensen, M.V., Wang, R., Peterson, B.S., Huang, C., Ilkayeva, O., Smith, N., Miller, N., Hajmrle, C., et al. (2015). Isocitrate-to-SENP1 signaling amplifies insulin secretion and rescues dysfunctional beta cells. J Clin Invest *125*, 3847-3860. 10.1172/JCI82498.

7.	Ainscow, E.K., Zhao, C., and Rutter, G.A. (2000). Acute overexpression of lactate dehydrogenase-A perturbs beta-cell mitochondrial metabolism and insulin secretion. Diabetes *49*, 1149-1155. DOI 10.2337/diabetes.49.7.1149.

8.	Sekine, N., Cirulli, V., Regazzi, R., Brown, L.J., Gine, E., Tamarit-Rodriguez, J., Girotti, M., Marie, S., MacDonald, M.J., Wollheim, C.B., and et al. (1994). Low lactate dehydrogenase and high mitochondrial glycerol phosphate dehydrogenase in pancreatic beta-cells. Potential role in nutrient sensing. Journal of Biological Chemistry *269*, 4895-4902.

9.	Schuit, F., Van Lommel, L., Granvik, M., Goyvaerts, L., de Faudeur, G., Schraenen, A., and Lemaire, K. (2012). beta-cell-specific gene repression: a mechanism to protect against inappropriate or maladjusted insulin secretion? Diabetes *61*, 969-975. 10.2337/db11-1564.

10.	Pullen, T.J., Khan, A.M., Barton, G., Butcher, S.A., Sun, G., and Rutter, G.A. (2010). Identification of genes selectively disallowed in the pancreatic islet. Islets *2*, 89-95. 10.4161/isl.2.2.11025.

11.	Lewandowski, S.L., Cardone, R.L., Foster, H.R., Ho, T., Potapenko, E., Poudel, C., VanDeusen, H.R., Sdao, S.M., Alves, T.C., Zhao, X., et al. (2020). Pyruvate Kinase Controls Signal Strength in the Insulin Secretory Pathway. Cell Metabolism *32*, 736-750.e735. 10.1016/j.cmet.2020.10.007.

12.	Foster, H.R., Ho, T., Potapenko, E., Sdao, S.M., Huang, S.M., Lewandowski, S.L., VanDeusen, H.R., Davidson, S.M., Cardone, R.L., Prentki, M., et al. (2022). β-cell deletion of the PKm1 and PKm2 isoforms of pyruvate kinase in mice reveals their essential role as nutrient sensors for the KATP channel. eLife *11*. 10.7554/eLife.79422.

13.	Ho, T., Potapenko, E., Davis, D.B., and Merrins, M.J. (2023). A plasma membrane-associated glycolytic metabolon is functionally coupled to K(ATP) channels in pancreatic alpha and beta cells from humans and mice. Cell Rep *42*, 112394. 10.1016/j.celrep.2023.112394.

14.	Merrins, M.J., Corkey, B.E., Kibbey, R.G., and Prentki, M. (2022). Metabolic cycles and signals for insulin secretion. Cell Metabolism *34*, 947-968. 10.1016/j.cmet.2022.06.003.

15.	Alves, T.C., Pongratz, R.L., Zhao, X., Yarborough, O., Sereda, S., Shirihai, O., Cline, G.W., Mason, G., and Kibbey, R.G. (2015). Integrated, Step-Wise, Mass-

Isotopomeric Flux Analysis of the TCA Cycle. Cell Metab *22*, 936-947.
10.1016/j.cmet.2015.08.021.

16.    Farfari, S., Schulz, V., Corkey, B., and Prentki, M. (2000). Glucose-regulated
anaplerosis and cataplerosis in pancreatic beta-cells: possible implication of a
pyruvate/citrate shuttle in insulin secretion. Diabetes *49*, 718-726.

17.    Lu, D., Mulder, H., Zhao, P., Burgess, S.C., Jensen, M.V., Kamzolova, S., Newgard,
C.B., and Sherry, A.D. (2002). 13C NMR isotopomer analysis reveals a connection
between pyruvate cycling and glucose-stimulated insulin secretion (GSIS). Proc Natl
Acad Sci U S A *99*, 2708-2713. 10.1073/pnas.052005699.

18.    Joseph, J.W., Jensen, M.V., Ilkayeva, O., Palmieri, F., Alárcon, C., Rhodes, C.J., and
Newgard, C.B. (2006). The Mitochondrial Citrate/Isocitrate Carrier Plays a
Regulatory Role in Glucose-stimulated Insulin Secretion. Journal of Biological
Chemistry *281*, 35624-35632. 10.1074/jbc.M602606200.

19.    Zhang, G.-F., Jensen, M.V., Gray, S.M., El, K., Wang, Y., Lu, D., Becker, T.C.,
Campbell, J.E., and Newgard, C.B. (2021). Reductive TCA cycle metabolism fuels
glutamine- and glucose-stimulated insulin secretion. Cell Metabolism *33*, 804-
817.e805. 10.1016/j.cmet.2020.11.020.

20.    Cline, G.W., Lepine, R.L., Papas, K.K., Kibbey, R.G., and Shulman, G.I. (2004). 13C
NMR isotopomer analysis of anaplerotic pathways in INS-1 cells. J Biol Chem *279*,
44370-44375. 10.1074/jbc.M311842200.

21.    Cline, G.W., Pongratz, R.L., Zhao, X., and Papas, K.K. (2011). Rates of insulin
secretion in INS-1 cells are enhanced by coupling to anaplerosis and Kreb's cycle
flux independent of ATP synthesis. Biochem Biophys Res Commun *415*, 30-35.
10.1016/j.bbrc.2011.09.153.

22.    Simpson, N.E., Khokhlova, N., Oca-Cossio, J.A., and Constantinidis, I. (2006).
Insights into the role of anaplerosis in insulin secretion: A 13C NMR study.
Diabetologia *49*, 1338-1348. 10.1007/s00125-006-0216-5.

23.    Malinowski, R.M., Ghiasi, S.M., Mandrup-Poulsen, T., Meier, S., Lerche, M.H.,
Ardenkjær-Larsen, J.H., and Jensen, P.R. (2020). Pancreatic β-cells respond to fuel
pressure with an early metabolic switch. Scientific Reports *10*. 10.1038/s41598-020-
72348-1.

24.    Benninger, R.K.P., and Hodson, D.J. (2018). New Understanding of β-Cell
Heterogeneity and In Situ Islet Function. Diabetes *67*, 537-547.

25.    Benninger, R.K.P., and Kravets, V. (2021). The physiological role of β-cell
heterogeneity in pancreatic islet function. Nature Reviews Endocrinology.
10.1038/s41574-021-00568-0.

26.    Nasteska, D., Fine, N.H.F., Ashford, F.B., Cuozzo, F., Viloria, K., Smith, G., Dahir,
A., Dawson, P.W.J., Lai, Y.-C., Bastidas-Ponce, A., et al. (2021). PDX1LOW
MAFALOW β-cells contribute to islet function and insulin release. Nature
Communications *12*, 674. 10.1038/s41467-020-20632-z.

27.    Rodriguez-Diaz, R., Dando, R., Jacques-Silva, M.C., Fachado, A., Molina, J.,
Abdulreda, M.H., Ricordi, C., Roper, S.D., Berggren, P.O., and Caicedo, A. (2011).
Alpha cells secrete acetylcholine as a non-neuronal paracrine signal priming beta cell
function in humans. Nature Medicine *17*, 888-892. 10.1038/nm.2371.

28.    Cabrera, O., Berman, D.M., Kenyon, N.S., Ricordi, C., Berggren, P.O., and Caicedo,
A. (2006). The unique cytoarchitecture of human pancreatic islets has implications
for islet cell function. Proceedings of the National Academy of Sciences of the United
States of America *103*, 2334-2339. 0510790103 [pii]10.1073/pnas.0510790103.

29.    Hodson, D.J., Mitchell, R.K., Bellomo, E.A., Sun, G., Vinet, L., Meda, P., Li, D., Li,
W.H., Bugliani, M., Marchetti, P., et al. (2013). Lipotoxicity disrupts incretin-regulated
human beta cell connectivity. Journal of Clinical Investigation *123*, 4182-4194.
10.1172/JCI68459.

30.    Spegel, P., Sharoyko, V.V., Goehring, I., Danielsson, A.P., Malmgren, S., Nagorny,
C.L., Andersson, L.E., Koeck, T., Sharp, G.W., Straub, S.G., et al. (2013). Time-

resolved metabolomics analysis of beta-cells implicates the pentose phosphate
pathway in the control of insulin release. Biochem J *450*, 595-605.
10.1042/BJ20121349.
31. Wallace, M., Whelan, H., and Brennan, L. (2013). Metabolomic analysis of pancreatic
beta cells following exposure to high glucose. Biochimica et Biophysica Acta (BBA) -
General Subjects *1830*, 2583-2590. 10.1016/j.bbagen.2012.10.025.
32. Barsby, T., Vähäkangas, E., Ustinov, J., Montaser, H., Ibrahim, H., Lithovius, V.,
Kuuluvainen, E., Chandra, V., Saarimäki-Vire, J., Katajisto, P., et al. (2023). Aberrant
metabolite trafficking and fuel sensitivity in human pluripotent stem cell-derived islets.
Cell Reports *42*. 10.1016/j.celrep.2023.112970.
33. Balboa, D., Barsby, T., Lithovius, V., Saarimaki-Vire, J., Omar-Hmeadi, M., Dyachok,
O., Montaser, H., Lund, P.E., Yang, M., Ibrahim, H., et al. (2022). Functional,
metabolic and transcriptional maturation of human pancreatic islets derived from
stem cells. Nat Biotechnol *40*, 1042-1055. 10.1038/s41587-022-01219-z.
34. Lu, Mulder, H., Zhao, P., Burgess, S.C., Jensen, M.V., Kamzolova, S., Newgard,
C.B., and Sherry, A.D. (2002). 13C NMR isotopomer analysis reveals a connection
between pyruvate cycling and glucose-stimulated insulin secretion (GSIS). Proc Natl
Acad Sci U S A *99*, 2708-2713. 10.1073/pnas.052005699.
35. Alonso, L., Piron, A., Morán, I., Guindo-Martínez, M., Bonàs-Guarch, S., Atla, G.,
Miguel-Escalada, I., Royo, R., Puiggròs, M., Garcia-Hurtado, X., et al. (2021).
TIGER: The gene expression regulatory variation landscape of human pancreatic
islets. Cell Reports *37*. 10.1016/j.celrep.2021.109807.
36. Goldberg, E., Eddy, E.M., Duan, C., and Odet, F. (2009). LDHC: The Ultimate Testis-
Specific Gene. Journal of Andrology *31*, 86-94. 10.2164/jandrol.109.008367.
37. Segerstolpe, Å., Palasantza, A., Eliasson, P., Andersson, E.-M., Andréasson, A.-C.,
Sun, X., Picelli, S., Sabirsh, A., Clausen, M., Bjursell, M.K., et al. (2016). Single-Cell
Transcriptome Profiling of Human Pancreatic Islets in Health and Type 2 Diabetes.
Cell Metabolism *24*, 593-607. 10.1016/j.cmet.2016.08.020.
38. Uhlen, M., Fagerberg, L., Hallstrom, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A.,
Sivertsson, A., Kampf, C., Sjostedt, E., Asplund, A., et al. (2015). Proteomics.
Tissue-based map of the human proteome. Science *347*, 1260419.
10.1126/science.1260419.
39. Thul, P.J., Akesson, L., Wiking, M., Mahdessian, D., Geladaki, A., Ait Blal, H., Alm,
T., Asplund, A., Bjork, L., Breckels, L.M., et al. (2017). A subcellular map of the
human proteome. Science *356*. 10.1126/science.aal3321.
40. van Gurp, L., Fodoulian, L., Oropeza, D., Furuyama, K., Bru-Tari, E., Vu, A.N.,
Kaddis, J.S., Rodriguez, I., Thorel, F., and Herrera, P.L. (2022). Generation of
human islet cell type-specific identity genesets. Nat Commun *13*, 2020.
10.1038/s41467-022-29588-8.
41. Pullen, T.J., Sylow, L., Sun, G., Halestrap, A.P., Richter, E.A., and Rutter, G.A.
(2012). Overexpression of Monocarboxylate Transporter-1 (Slc16a1) in Mouse
Pancreatic beta-Cells Leads to Relative Hyperinsulinism During Exercise. Diabetes
*61*, 1719-1725. 10.2337/db11-1531.
42. Zhao, C., Wilson, M.C., Schuit, F., Halestrap, A.P., and Rutter, G.A. (2001).
Expression and distribution of lactate/monocarboxylate transporter isoforms in
pancreatic islets and the exocrine pancreas. Diabetes *50*, 361-366.
43. Deng, H., Gao, Y., Trappetti, V., Hertig, D., Karatkevich, D., Losmanova, T., Urzi, C.,
Ge, H., Geest, G.A., Bruggmann, R., et al. (2022). Targeting lactate dehydrogenase
B-dependent mitochondrial metabolism affects tumor initiating cells and inhibits
tumorigenesis of non-small cell lung cancer by inducing mtDNA damage. Cellular
and Molecular Life Sciences *79*. 10.1007/s00018-022-04453-5.
44. Ždralević, M., Brand, A., Di Ianni, L., Dettmer, K., Reinders, J., Singer, K., Peter, K.,
Schnell, A., Bruss, C., Decking, S.-M., et al. (2018). Double genetic disruption of
lactate dehydrogenases A and B is required to ablate the "Warburg effect" restricting

tumor growth to oxidative metabolism. Journal of Biological Chemistry *293*, 15947-
15961. 10.1074/jbc.RA118.004180.

45.   Jekabsons, M., San Martín, A., Ceballo, S., Ruminot, I., Lerchundi, R., Frommer, W.B., and Barros, L.F. (2013). A Genetically Encoded FRET Lactate Sensor and Its Use To Detect the Warburg Effect in Single Cancer Cells. PLoS ONE *8*. 10.1371/journal.pone.0057712.

46.   Sdao, S.M., Ho, T., Poudel, C., Foster, H.R., De Leon, E.R., Adams, M.T., Lee, J.H., Blum, B., Rane, S.G., and Merrins, M.J. (2021). CDK2 limits the highly energetic secretory program of mature beta cells by restricting PEP cycle-dependent K(ATP) channel closure. Cell Rep *34*, 108690. 10.1016/j.celrep.2021.108690.

47.   Shibata, S., Sogabe, S., Miwa, M., Fujimoto, T., Takakura, N., Naotsuka, A., Kitamura, S., Kawamoto, T., and Soga, T. (2021). Identification of the first highly selective inhibitor of human lactate dehydrogenase B. Scientific Reports *11*. 10.1038/s41598-021-00820-7.

48.   Blanchi, B., Taurand, M., Colace, C., Thomaidou, S., Audeoud, C., Fantuzzi, F., Sawatani, T., Gheibi, S., Sabadell-Basallote, J., Boot, F.W.J., et al. (2023). EndoC-βH5 cells are storable and ready-to-use human pancreatic beta cells with physiological insulin secretion. Molecular Metabolism *76*. 10.1016/j.molmet.2023.101772.

49.   Schuit, F., De Vos, A., Farfari, S., Moens, K., Pipeleers, D., Brun, T., and Prentki, M. (1997). Metabolic fate of glucose in purified islet cells. Glucose-regulated anaplerosis in beta cells. J Biol Chem *272*, 18572-18579. 10.1074/jbc.272.30.18572.

50.   Lemaire, K., Thorrez, L., and Schuit, F. (2016). Disallowed and Allowed Gene Expression: Two Faces of Mature Islet Beta Cells. Annu Rev Nutr *36*, 45-71. 10.1146/annurev-nutr-071715-050808.

51.   Sato, Y., Endo, H., Okuyama, H., Takeda, T., Iwahashi, H., Imagawa, A., Yamagata, K., Shimomura, I., and Inoue, M. (2011). Cellular hypoxia of pancreatic beta-cells due to high levels of oxygen consumption for insulin secretion in vitro. J Biol Chem *286*, 12524-12532. 10.1074/jbc.M110.194738.

52.   Bensellam, M., Duvillie, B., Rybachuk, G., Laybutt, D.R., Magnan, C., Guiot, Y., Pouyssegur, J., and Jonas, J.C. (2012). Glucose-induced O(2) consumption activates hypoxia inducible factors 1 and 2 in rat insulin-secreting pancreatic beta-cells. PLoS One *7*, e29807. 10.1371/journal.pone.0029807.

53.   Taddeo, E.P., Stiles, L., Sereda, S., Ritou, E., Wolf, D.M., Abdullah, M., Swanson, Z., Wilhelm, J., Bellin, M., McDonald, P., et al. (2018). Individual islet respirometry reveals functional diversity within the islet population of mice and human donors. Mol Metab *16*, 150-159. 10.1016/j.molmet.2018.07.003.

54.   Sanchez, P.K.M., Khazaei, M., Gatineau, E., Geravandi, S., Lupse, B., Liu, H., Dringen, R., Wojtusciszyn, A., Gilon, P., Maedler, K., and Ardestani, A. (2021). LDHA is enriched in human islet alpha cells and upregulated in type 2 diabetes. Biochem Biophys Res Commun *568*, 158-166. 10.1016/j.bbrc.2021.06.065.

55.   Moin, A.S.M., Cory, M., Gurlo, T., Saisho, Y., Rizza, R.A., Butler, P.C., and Butler, A.E. (2020). Pancreatic alpha-cell mass across adult human lifespan. Eur J Endocrinol *182*, 219-231. 10.1530/EJE-19-0844.

56.   Zaborska, K.E., Dadi, P.K., Dickerson, M.T., Nakhe, A.Y., Thorson, A.S., Schaub, C.M., Graff, S.M., Stanley, J.E., Kondapavuluru, R.S., Denton, J.S., and Jacobson, D.A. (2020). Lactate activation of alpha-cell KATP channels inhibits glucagon secretion by hyperpolarizing the membrane potential and reducing Ca(2+) entry. Mol Metab *42*, 101056. 10.1016/j.molmet.2020.101056.

57.   Pullen, T.J., and Rutter, G.A. (2013). When less is more: the forbidden fruits of gene repression in the adult beta-cell. Diabetes Obes Metab *15*, 503-512. 10.1111/dom.12029.

58.   Fu, A., Alvarez-Perez, J.C., Avizonis, D., Kin, T., Ficarro, S.B., Choi, D.W., Karakose, E., Badur, M.G., Evans, L., Rosselot, C., et al. (2020). Glucose-dependent

partitioning of arginine to the urea cycle protects β-cells from inflammation. Nature
Metabolism *2*, 432-446. 10.1038/s42255-020-0199-4.

59.　Ainscow, E.K., and Rutter, G.A. (2002). Glucose-Stimulated Oscillations in Free
Cytosolic ATP Concentration Imaged in Single Islet β-Cells. Diabetes *51*, S162-
S170. 10.2337/diabetes.51.2007.S162.

60.　Campbell, J.E., and Newgard, C.B. (2021). Mechanisms controlling pancreatic islet
cell function in insulin secretion. Nat Rev Mol Cell Biol *22*, 142-158. 10.1038/s41580-
020-00317-7.

61.　Gerber, P.A., and Rutter, G.A. (2017). The Role of Oxidative Stress and Hypoxia in
Pancreatic Beta-Cell Dysfunction in Diabetes Mellitus. Antioxid Redox Signal *26*,
501-518. 10.1089/ars.2016.6755.

62.　Bensellam, M., Jonas, J.C., and Laybutt, D.R. (2018). Mechanisms of beta-cell
dedifferentiation in diabetes: recent findings and future research directions. J
Endocrinol *236*, R109-R143. 10.1530/JOE-17-0516.

63.　Marselli, L., Piron, A., Suleiman, M., Colli, M.L., Yi, X., Khamis, A., Carrat, G.R.,
Rutter, G.A., Bugliani, M., Giusti, L., et al. (2020). Persistent or Transient Human β
Cell Dysfunction Induced by Metabolic Stress: Specific Signatures and Shared Gene
Expression with Type 2 Diabetes. Cell Reports *33*. 10.1016/j.celrep.2020.108466.

64.　Viloria, K., Nasteska, D., Ast, J., Hasib, A., Cuozzo, F., Heising, S., Briant, L.J.B.,
Hewison, M., and Hodson, D.J. (2023). GC-Globulin/Vitamin D–Binding Protein Is
Required for Pancreatic α-Cell Adaptation to Metabolic Stress. Diabetes *72*, 275-289.
10.2337/db22-0326.

65.　Fernandez, C.A., Des Rosiers, C., Previs, S.F., David, F., and Brunengraber, H.
(1996). Correction of 13C mass isotopomer distributions for natural stable isotope
abundance.　J　Mass　Spectrom　*31*,　255-262.　10.1002/(SICI)1096-
9888(199603)31:3<255::AID-JMS290>3.0.CO;2-3.

66.　Kazimierczuk, K., and Orekhov, V.Y. (2011). Accelerated NMR Spectroscopy by
Using Compressed Sensing. Angewandte Chemie International Edition *50*, 5556-
5559. 10.1002/anie.201100370.

67.　Delaglio, F., Grzesiek, S., Vuister, G., Zhu, G., Pfeifer, J., and Bax, A. (1995).
NMRPipe: A multidimensional spectral processing system based on UNIX pipes.
Journal of Biomolecular NMR *6*. 10.1007/bf00197809.

68.　Ludwig, C., and Günther, U.L. (2011). MetaboLab - advanced NMR data processing
and analysis for metabolomics. BMC Bioinformatics *12*. 10.1186/1471-2105-12-366.

69.　Tantama, M., Martinez-Francois, J.R., Mongeon, R., and Yellen, G. (2013). Imaging
energy status in live cells with a fluorescent biosensor of the intracellular ATP-to-
ADP ratio. Nat Commun *4*, 2550. 10.1038/ncomms3550.

70.　Gavet, O., and Pines, J. (2010). Progressive activation of CyclinB1-Cdk1 coordinates
entry to mitosis. Dev Cell *18*, 533-543. 10.1016/j.devcel.2010.02.013.

71.　Viloria, K., Nasteska, D., Briant, L.J.B., Heising, S., Larner, D.P., Fine, N.H.F.,
Ashford, F.B., da Silva Xavier, G., Ramos, M.J., Hasib, A., et al. (2020). Vitamin-D-
Binding Protein Contributes to the Maintenance of α Cell Function and Glucagon
Secretion. Cell Reports *31*. 10.1016/j.celrep.2020.107761.

72.　Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial
reconstruction of single-cell gene expression data. Nat Biotechnol *33*, 495-502.
10.1038/nbt.3192.

73.　Aibar, S., Gonzalez-Blas, C.B., Moerman, T., Huynh-Thu, V.A., Imrichova, H.,
Hulselmans, G., Rambow, F., Marine, J.C., Geurts, P., Aerts, J., et al. (2017).
SCENIC: single-cell regulatory network inference and clustering. Nat Methods *14*,
1083-1086. 10.1038/nmeth.4463.

74.　Arda, H.E., Li, L., Tsai, J., Torre, E.A., Rosli, Y., Peiris, H., Spitale, R.C., Dai, C., Gu,
X., Qu, K., et al. (2016). Age-Dependent Pancreatic Gene Regulation Reveals
Mechanisms Governing Human beta Cell Function. Cell Metab *23*, 909-920.
10.1016/j.cmet.2016.04.002.

75.  Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal
probabilistic RNA-seq quantification. Nat Biotechnol *34*, 525-527. 10.1038/nbt.3519.
76.  Akerman, I., Kasaai, B., Bazarova, A., Sang, P.B., Peiffer, I., Artufel, M., Derelle, R.,
Smith, G., Rodriguez-Martinez, M., Romano, M., et al. (2020). A predictable
conserved DNA base composition signature defines human core DNA replication
origins. Nat Commun *11*, 4826. 10.1038/s41467-020-18527-0.
77.  Baron, M., Veres, A., Wolock, S.L., Faust, A.L., Gaujoux, R., Vetere, A., Ryu, J.H.,
Wagner, B.K., Shen-Orr, S.S., Klein, A.M., et al. (2016). A Single-Cell Transcriptomic
Map of the Human and Mouse Pancreas Reveals Inter- and Intra-cell Population
Structure. Cell Syst *3*, 346-360 e344. 10.1016/j.cels.2016.08.011.
78.  Xin, Y., Kim, J., Okamoto, H., Ni, M., Wei, Y., Adler, C., Murphy, A.J., Yancopoulos,
G.D., Lin, C., and Gromada, J. (2016). RNA Sequencing of Single Human Islet Cells
Reveals Type 2 Diabetes Genes. Cell Metab *24*, 608-615.
10.1016/j.cmet.2016.08.018.
79.  Mawla, A.M., and Huising, M.O. (2019). Navigating the Depths and Avoiding the
Shallows of Pancreatic Islet Cell Transcriptomes. Diabetes *68*, 1380-1393.
10.2337/dbi18-0019.
80.  Lawlor, N., George, J., Bolisetty, M., Kursawe, R., Sun, L., Sivakamasundari, V.,
Kycia, I., Robson, P., and Stitzel, M.L. (2017). Single-cell transcriptomes identify
human islet cell signatures and reveal cell-type–specific expression changes in type
2 diabetes. Genome Research *27*, 208-222. 10.1101/gr.212720.116.
81.  Wang, Y.J., Schug, J., Won, K.J., Liu, C., Naji, A., Avrahami, D., Golson, M.L., and
Kaestner, K.H. (2016). Single-Cell Transcriptomics of the Human Endocrine
Pancreas. Diabetes *65*, 3028-3038. 10.2337/db16-0405.
82.  Xin, Y., Gutierrez, G.D., Okamoto, H., Kim, J., Lee, A.H., Adler, C., Ni, M.,
Yancopoulos, G.D., Murphy, A.J., and Gromada, J. (2018). Pseudotime Ordering of
Single Human beta-Cells Reveals States of Insulin Production and Unfolded Protein
Response. Diabetes. 10.2337/db18-0365.
83.  Enge, M., Arda, H.E., Mignardi, M., Beausang, J., Bottino, R., Kim, S.K., and Quake,
S.R. (2017). Single-Cell Analysis of Human Pancreas Reveals Transcriptional
Signatures of Aging and Somatic Mutation Patterns. Cell *171*, 321-330 e314.
10.1016/j.cell.2017.09.004.
84.  Akerman, I., Tu, Z., Beucher, A., Rolando, D.M.Y., Sauty-Colace, C., Benazra, M.,
Nakic, N., Yang, J., Wang, H., Pasquali, L., et al. (2017). Human Pancreatic beta Cell
lncRNAs Control Cell-Specific Regulatory Networks. Cell Metab *25*, 400-411.
10.1016/j.cmet.2016.11.016.

1054
1055

**FIGURES AND FIGURE LEGENDS**

1058
1059

**Figure 1: MID analysis of glucose fate in human and mouse islets. A)** Schematic
showing GC-MS and $^1$H-NMR-based $^{13}C_6$ glucose-tracing protocol in primary islets. **B)**
Schematic showing mass isotopomer distribution (MID) analysis of $^{13}C_6$ glucose-tracing
data. **C-E)** MID analysis showing similar incorporation of $^{13}C$ from $^{13}C_6$ glucose into malate
(C), alanine (D) and glutamate (E) in human and mouse islets. **F, G**) MID analysis showing
increased incorporation of $^{13}C$ from $^{13}C_6$ glucose into m+2 and m+3 aspartate (F), and m+2
fumarate (G) in mouse compared to human islets. **H, I)** Total amount of extracted aspartate
(H) and alanine (I) is similar in human and mouse islets. **(J-L)** Total amount of extracted
malate (J) and fumarate (K) is decreased in mouse relative to human islets, whereas
glutamate (L) is increased. For all data, n = 18 independent replicates from 9 human donors;

1070    n = 10 islet preparations from 15 animals. C-G were analyzed using two-way ANOVA and
1071    Sidak's post-hoc test. H-L were analyzed using Welch's test. Bar graphs show individual
1072    datapoints and mean ± SEM. AU = arbitrary unit.
1073
1074

1075

1076

**Figure 2: Human islets produce lactate. A-C)** MID analysis shows detectable glucose
incorporation into m+2 (TCA cycle) and m+3 (pyruvate conversion) lactate, with more
accumulation in human (A, B) versus mouse (A, C) islets (n = 18 independent replicates
from 9 human donors; n = 10 islet preparations from 15 animals) (two-way ANOVA and
Sidak's post-hoc test). **D)** Total lactate generation is higher in human compared to mouse
islets (n = 18 independent replicates from 9 human donors; n = 10 islet preparations from 15
animals) (Welch's test). Bar graphs show individual datapoints and mean ± SEM.

1084
1085
1086

1087

**Figure 3: Incorporation of $^{13}C$ from $^{13}C_6$ glucose into TCA cycle metabolites through**
**PDH and PC**. **A)** White and blue circles, respectively, show the incorporation of $^{12}C$ and $^{13}C$
into TCA cycle metabolites arising from metabolism of pyruvate by PC. **B)** White and red
circles, respectively, represent $^{12}C$ and $^{13}C$ atoms as incorporated from $^{13}C_6$ glucose into the
TCA cycle through the conversion of pyruvate to acetyl-CoA by PDH. **C-E)** Lactate$_{000}$,
lactate$_{111}$ and lactate$_{110}$ are the most abundant isotopomers (C) in both humans and mice
(D), although the incorporation of $^{13}C$ from $^{13}C_6$ glucose into lactate$_{111}$ is significantly higher
in human than mouse islets (D, E). **F-H)** $^{13}C$ incorporation into alanine isotopomers (F) is
similar in human and mouse islets (G), with alanine$_{111}$ being the most represented labeled
isotopomer (G, H). **I, J**) The distribution of labeling patterns for glutamate (I) are similar in
human and mouse islets (J), with glutamate$_{00011}$ being the most abundant labeled
isotopomer in both species (J). For all data, n = 16-17 islet preparations, 9 human donors
and n = 12-15 islet preparations, 7-8 animals. Data were analyzed using two-way ANOVA

and Sidak's post-hoc test.  Bar graphs show individual datapoints and mean ± SEM. AU =

arbitrary unit.

1103

**Figure 4: Human islets express LDHA, LDHB and LDHC protein. A)** LDH expression, assessed using an antibody with cross-reactivity against LDHA, LDHB and LDHC, is higher in human β cells compared to mouse β cells. **B)** LDH expression increases in mouse exocrine tissue and β cells following 8-12 weeks high fat diet (HFD) feeding versus age-matched standard diet (SD) controls. **C)** Corrected total cell fluorescence (CTCF) quantification of LDH protein expression in human β cells (n = 240 cells, 3 donors), SD mouse β-cells (n = 220 cells, 3 animals), and HFD mouse β cells (n = 250 cells, 3 animals) (Kruskal-Wallis test, Dunn's post-hoc test). **D)** As for G), but showing LDH immunoreactivity in the exocrine pancreas of mice fed either SD or HFD (n = 230-250 cells, 3 animals) (Mann-Whitney test). **E)** PDX1 expression is decreased in β cells of mice fed HFD for 8 weeks versus SD controls (n = 240 cells, n = 3 animals). **F)** CTCF) quantification of PDX1 protein expression in SD and HFD mouse β-cells (n = 240 cells, 3 animals) (Mann-Whitney test). **G)** Immunostaining showing strong LDH expression in human liver from donors with non-alcoholic steatohepatitis (n = 240 cells, n = 3 donors) (Mann-Whitney test). Representative images are shown. Scale bar = 30 μm. Bar graphs show individual datapoints and mean ± SEM. Violin plot shows median and interquartile range. AU = arbitrary unit.

1123
1124
**Figure 5: Human β-cells specifically express *LDHB*. A)** Normalized mRNA levels
(transcripts per million, TPM) for *INS, GCG, LDHA* and *LDHB* genes in fluorescent-activated
cell-sorted (FACS) α and β cell samples (re-analysis of data from [74]). **B, C)** Normalized
LDHA (B) and LDHB (C) expression in α, β, δ and γ cells from five independent human islet
single cell RNA-sequencing experiments [37,80-83]. Data from each study was subjected to re-
assignment of cell identity based upon strict criteria (see Methods). **D-F)** UMAP plots
showing *LDHA* (D), *LDHB* (E) and *SLC16A1* (F) raw counts clustered according to cell type
(data taken from n = 18 donors in [40,77,78]). **G, H)** Genome browser snapshot of transcription
factor binding, histone modification (ChIP-seq, targets as indicated) and RNA-sequencing

experiments performed on human islets for LDHA (G) and LDHB (H) [84]. Scales in A-C
represent RPKM.
1136
1137

1138
1139
**Figure 6: LDHB is predominantly localized to β cells within the islet. A)** LDHB protein
expression strongly co-localizes with insulin (INS) expression in human pancreas sections.
Zoom-in (middle panel) shows absence of LDHB in glucagon (GCG)+ cells. Zoom-in (bottom
panel) shows a small subpopulation of GCG+ cells with detectable LDHB staining. **B)**
Quantification of LDHB expression in GCG+ and INS+ cells in human pancreas sections,
including sub-analysis of GCG+ segregated by high (Hi) and low (Lo) LDHB levels (> 8 x $10^4$
CTCF) (n= 150 cells, 3 donors) (Mann-Whitney test). **C)** Pie-charts showing the proportions
of GCG+ and INS+ cells that express Hi or Lo LDHB in human pancreas sections (n= 150
cells, 3 donors). **D)** Representative images (zoom-in from A) showing classification of LDHB
Hi and Lo cells for GCG and INS. **E)** As for A), but showing that LDHB protein expression
remains strongly co-localized with INS in isolated human islets. **F)** Quantification of LDHB
expression in GCG+ and INS+ cells in isolated human islets, including sub-analysis of
GCG+ segregated by high (Hi) and low (Lo) LDHB levels (> 8 x $10^4$ CTCF) (n= 180 cells, 3
donors) (Mann-Whitney test). **G, H)** As for B and C, but isolated human islets (n= 180 cells,
3 donors). Scale bar = 30 μm. Violin plot shows median and interquartile range.
1155
1156

**Figure 7: Effects of LDHB inhibition on glucose-stimulate lactate, ATP/ADP and Ca²⁺ in human β cells. A)** 17 mM glucose stimulates rapid increases in cytosolic lactate in vehicle-treated islets from donor 1, measured using the β cell-specific FRET sensor RIP-Laconic. Glucose-stimulated lactate generation is amplified by an LDHB inhibitor (AXKO-0046) (n = 10-15 islets). **B)** Summary bar graph showing significant amplification and inhibition of lactate by AXKO-0046 (n = 10-15 islets) (one-way ANOVA and Sidak's post-hoc test). **C)** Representative images showing human β cell lactate levels in the presence of vehicle orAXKO-0046. **D-F)** As for A-C, except donor 2 (n = 10-14 islets) (unpaired t-test). **G-I)** As for A-C, except donor 3 (n = 10-12 islets) (unpaired t-test). **J-L)** AXKO-0046 amplifies glucose-stimulated lactate generation in mouse β cells. **M)** AXKO-0046 does not significantly influence glucose-stimulated ATP/ADP ratios in human β cells, measured using RIP-Perceval-HR (n = 3 donors, 19-23 islets) (unpaired t-test). **N, O)** Individual traces (N) and summary bar graph (O) showing that AXKO-0046 influences Ca²⁺ spiking frequency, but not Ca²⁺ levels in human β cells (n = 3 donors, 87-101 cells) (Mann-Whitney test). Unless other stated, traces shown mean ± SEM. Bar graphs show individual datapoints and mean ± SEM. Scale bar = 83 μm. Arrows and box show AUC calculation boundaries. G3, 3 mM glucose; G17, 17 mM glucose. Islets were pre-incubated with 10 μM AXKO-0046.

1178
1179
**Figure 8: Schematic showing pyruvate management in human islets. A)** In the current
view (top) of human β cell metabolism, glycolytically-derived pyruvate enters the TCA
through the actions of pyruvate dehydrogenase (PDH) and pyruvate carboxylase (PC). The
PEP cycle and extra-mitochondrial ADP might also contribute to $K_{ATP}$ channel regulation and
the triggering phase of insulin secretion. Alternative fates for pyruvate (i.e. generation of
lactate) are suppressed due to low levels of LDH/*LDHA* expression. **B)** The high-resolution
view of β cell metabolism reveals that LDHB contributes to regulation of lactate generation.
The TCA cycle intermediary malate is also converted to pyruvate, which is then further
converted to lactate leading to two inter-dependent pools of reducing equivalents to support
REDOX (schematic produced in Inkscape, Inkscape Project).
1190